


NORTHERN MANITOBA AND NORTHERN SASKATCHEWAN

Waterfowl Breeding Population Survey

2003



The data presented in this report are preliminary and subject to further editing. Final estimates will be available from the U.S. Fish and Wildlife Service, Office of Migratory Bird Management, Laurel, Maryland 20708

**Waterfowl Breeding Population Survey:
Northern Manitoba, Northern Saskatchewan**

STRATA SURVEYED: 21-24

DATES: May 17-June 4, 2003

DATA SUPPLIED BY: U. S. Fish and Wildlife Service

AERIAL CREW

Pilot/Observer	Fred Roetker, Flyway Biologist Division of Migratory Bird Management Lafayette, LA
Observer	Pat Stinson, Wildlife Biologist Migratory Bird Field Office Jackson, MS

ABSTRACT: Waterfowl numbers were mixed in northern Saskatchewan and northern Manitoba. Mallard numbers were 21% below 2002 values. Results from selected species are listed below:

	Estimates in thousands	% change from 2002	% change from 10 yr. mean	% change from 1955-02 mean
Mallard	610.4	-21.2%	-24.3%	-25.2%
Wigeon	188.0	7.3%	9.1%	-7.6%
GW-Teal	267.1	3.7%	13.4%	77.8%
Scaup	324.6	-8.4%	-9.6%	-28.4%
Ringneck	319.3	-39.5%	1.7%	28.8%

METHODS: The procedures followed in conducting this survey are contained in the Standard Operating Procedures for Aerial Breeding Population and Habitat Surveys in North America, Section III, revised 1987. Waterfowl data were collected using an aerial onboard computerized recording system designed to attribute each waterfowl observation with a respective latitude/longitude location. Observations were then logged along with sample details, i.e., strata transects, segments, time, and climatic conditions. This survey system began testing in Alaska during 1996 and has been gradually been implemented in each survey crew area since. During this transition, the Migratory Bird Management Office elected to redesign the sampling of stratum 25. This effort, effective during the 1998 and 1999 surveys, was intended to more appropriately sample the number and distribution of breeding waterfowl across the stratified habitat type. Historically, nine segments were concentrated in the Saskatchewan River delta and were not all oriented on latitude. Data from this stratum was included in the northern Manitoba report. In the new design, all thirty segments have been oriented on latitude with only five segments in the river delta area. There are now a total of five transects evenly distributed throughout the stratum, however, the sampling area is still identical to the historic survey. Direct comparisons of the 1998-2003 and historic data should be made only after evaluating this design change. Beginning with the May 2000 survey, the Migratory Bird Office elected to report Stratum 25 data in the Southern Manitoba Waterfowl Breeding Population Survey report since the southern Manitoba crew typically flies Stratum 25 and collects the data. Historic tables and graphs in this report have been adjusted to accommodate this change.

WEATHER AND HABITAT CONDITIONS: A much more normal spring break-up occurred across the region after the exceptionally late break-up of 2002. A wide range of habitat conditions prevailed across northern Saskatchewan in 2003. Dry basins continued to persist in the southwest portion of the survey area. Conditions were wetter to the north. Between Cree Lake and Lake Athabasca, conditions were very wet, and flooding occurred along the Otherside River during our survey. However, in the bulk of the country between these extremes, habitat conditions appeared ideal. Most smaller riverine habitat is stabilized by beavers, and ideal ponds were abundant throughout many drainages. In Manitoba, water levels tended to be lower. Much lower than average winter and spring precipitation reduced flows in small streams to the point that many beaver ponds were dry or quite recessional. The larger river and lake systems were also well below normal levels. Miles and miles of exposed shoreline habitat existed along the major flowages such as the Burntwood and the Nelson. Rocks were exposed everywhere, loafing sites were abundant, and associated upland meadows should offer stable nesting opportunities. The low water in major rivers and lakes should not adversely impact waterfowl nesting, however the critical conditions that persist along the small streams and beaver ponds will probably limit the production capacity of these typically ideal habitats.

BREEDING POPULATION ESTIMATES: Table 1 presents breeding waterfowl population data along with comparisons to 2002, the 1992 to 2002 average, and the 1955-2002 average. Appendix 1 presents earlier waterfowl breeding estimates.

CONCLUSIONS: Breeding waterfowl numbers were mixed in 2003. Habitat conditions ranged from mostly good in Saskatchewan to poor across much of northern Manitoba. The low water levels in many small streams and beaver ponds in northern Manitoba may limit production potential.

Table 1. Status of waterfowl breeding population estimates (thousands, adjusted for visibility bias) by species and stratum with comparisons against the previous year, the previous 10-year mean, and the long-term mean for Northern Saskatchewan and Northern Manitoba.

Species/Ponds	Stratum (2003)						% Change From				
	21	22	23	24	2003 Total	2002 Total	10-Year Mean	Long-Term Mean	2002	10-Year Mean	Long-Term Mean
Ducks											
Dabblers											
Mallard	89.5	92.4	193.5	235.0	610.4	774.5	806.6	816.1	-21.2%	-24.3%	-25.2%
Am. black duck	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.6	--	-100.0%	-100.0%
Gadwall	0.7	8.1	0.0	10.1	19.0	14.1	12.6	16.0	34.7%	51.2%	18.5%
Am. wigeon	31.1	22.9	26.3	107.7	188.0	175.1	172.3	203.3	7.3%	9.1%	-7.6%
Am. green-winged teal	45.4	27.0	73.7	121.0	267.1	257.7	235.5	150.2	3.7%	13.4%	77.8%
Blue-winged teal	21.2	60.8	15.3	51.1	148.4	201.2	125.2	177.2	-26.2%	18.5%	-16.2%
N. shoveler	0.8	1.7	0.0	1.2	3.7	19.1	25.9	23.3	-80.8%	-85.9%	-84.3%
N. pintail	0.6	1.9	0.0	2.7	5.2	9.0	10.9	25.0	-42.3%	-52.0%	-79.2%
Subtotal	189.3	215.0	308.8	528.8	1241.7	1450.7	1389.6	1412.8	-14.4%	-10.6%	-12.1%
Divers											
Redhead	0.7	14.4	0.0	0.0	15.1	6.7	6.4	11.7	124.0%	134.4%	28.8%
Canvasback	1.2	3.8	0.0	0.0	5.0	14.5	14.6	26.0	-65.7%	-66.1%	-80.9%
Scaups	64.5	71.1	76.1	113.0	324.6	354.5	359.0	453.3	-8.4%	-9.6%	-28.4%
Ring-necked duck	41.2	37.7	84.3	156.1	319.3	527.8	314.0	247.9	-39.5%	1.7%	28.8%
Goldeneyes	23.6	48.8	76.6	116.3	265.2	368.7	280.2	157.6	-28.1%	-5.4%	68.3%
Bufflehead	37.6	39.6	118.5	50.9	246.5	236.0	204.0	133.2	4.5%	20.8%	85.1%
Ruddy Duck	1.3	11.6	2.7	0.0	15.7	7.1	3.4	5.7	119.6%	366.2%	172.9%
Subtotal	170.1	226.9	358.1	436.2	1191.4	1515.3	1181.6	1035.4	-21.4%	0.8%	15.1%
Miscellaneous											
Long-tailed duck	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	--	--	-100.0%
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	--	-100.0%	-100.0%
Scoters	7.9	0.7	19.0	29.8	57.4	55.7	56.7	56.4	3.0%	1.2%	1.7%
Mergansers	39.5	30.5	78.5	55.0	203.5	308.5	223.1	175.8	-34.1%	-8.8%	15.8%
Subtotal	47.4	31.2	97.5	84.7	260.8	364.2	279.9	232.2	-28.4%	-6.8%	12.3%
Total Ducks	406.8	473.1	764.4	1049.7	2694.0	3330.2	2851.1	2680.4	-19.1%	-5.5%	0.5%
Canada Goose	23.9	22.9	21.7	101.7	170.2	213.6	165.4	94.9	-20.3%	2.8%	79.3%
Am. coot	0.5	4.0	0.0	1.6	6.1	101.5	38.8	59.5	-94.0%	-84.2%	-89.7%

Table 2. Northern Saskatchewan, Northern Manitoba stratum data sheet, May 2003.

	Strata				
	21	22	23	24	Total
Survey Square miles in stratum	44,922	20,278	68,320	82,966	216,486
Square miles in sample	418.5	166.5	301.5	513.0	1,399.5
Linear miles in sample	1,674	666	1,206	2,052	5,598
Number of transects in sample	11	6	9	8	34
Number of segments in sample	93	37	67	114	311
Expansion factor	107.341	121.790	226.600	161.727	

	Strata				
Current Year Design	21	22	23	24	Total
Square mile in stratum	44,922	20,278	68,320	82,966	216,486
Square miles in sample	396	166.5	297	513	1,372.5
Linear miles in sample	1,584	666	1188	2,052	5,490
Number of transects in sample	10	6	9	8	33
Number of segments in sample	88	37	66	114	305
Expansion factor	113.439	121.790	230.035	161.727	

Appendix 1. Long-term trend in adjusted waterfowl breeding population estimates (thousands).

Species/Ponds	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964
Ducks										
Dabblers										
Mallard	572.4	724.4	504.1	904.6	757.9	388.1	430.4	422.2	396.3	511.6
Am. black duck	0.0	0.0	0.0	0.0	2.7	3.6	2.4	10.1	1.6	13.4
Gadwall	1.0	1.0	5.6	0.0	25.7	1.0	34.3	9.9	9.4	8.5
Am. wigeon	105.4	22.9	45.8	23.6	86.6	123.2	104.0	112.4	103.8	147.2
Am. green-winged teal	53.0	35.6	12.3	0.0	76.2	10.7	7.8	27.2	26.5	75.2
Blue-winged teal	13.8	35.9	33.5	143.4	92.6	40.8	13.8	170.8	169.9	305.4
N. shoveler	4.2	1.9	0.0	2.6	19.7	0.0	0.0	9.0	5.8	55.4
N. pintail	21.0	48.0	17.8	9.4	44.6	11.4	111.7	12.8	32.3	33.3
Subtotal	770.7	869.8	619.2	1083.6	1105.9	578.8	704.2	774.4	745.6	1150.0
Divers										
Redhead	30.4	0.0	1.0	0.0	0.0	21.1	6.2	11.4	12.8	30.2
Canvasback	22.3	22.5	0.0	2.3	45.9	13.3	43.4	5.9	42.7	154.4
Scaups	449.2	228.2	779.6	359.9	449.5	131.5	215.2	239.2	277.3	312.5
Ring-necked duck	113.2	46.3	0.0	0.0	2.5	23.1	36.7	90.4	193.1	140.6
Goldeneyes	57.3	1.2	6.4	191.1	489.5	219.8	136.9	126.6	62.3	39.3
Bufflehead	45.9	15.7	18.8	27.4	26.9	124.8	28.0	54.6	39.8	30.0
Ruddy Duck	0.0	0.0	0.0	0.0	0.0	0.0	0.0	23.8	9.9	0.1
Subtotal	718.3	313.8	805.9	580.7	1014.3	533.6	466.5	551.9	638.0	707.0
Miscellaneous										
Oldsquaw	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scoters	12.1	6.8	71.8	26.7	100.5	5.5	44.4	33.9	13.9	39.5
Mergansers	160.2	51.5	35.7	89.3	77.2	75.0	40.6	99.7	125.7	73.7
Subtotal	172.3	58.3	107.6	116.1	177.7	80.5	85.0	133.7	139.5	113.2
Total Ducks	1661.4	1241.9	1532.6	1780.4	2297.9	1192.9	1255.7	1459.9	1523.2	1970.3
Canada Goose	71.3	65.5	48.7	53.1	147.2	26.5	94.8	17.4	15.7	42.6
Am. coot	0.0	0.0	0.0	0.0	22.5	7.9	0.0	5.5	17.3	6.3

Species/Ponds	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974
Ducks										
Dabblers										
Mallard	719.7	453.0	1363.3	1221.0	1368.1	1834.2	869.6	724.6	849.6	540.0
Am. black duck	2.6	1.5	7.8	5.8	0.0	1.1	0.0	0.0	0.0	0.0
Gadwall	26.4	22.6	51.3	48.6	46.8	28.9	7.7	18.3	22.5	9.7
Am. wigeon	185.9	178.3	143.4	321.0	345.3	532.7	379.9	240.0	252.1	168.4
Am. green-winged teal	67.1	58.3	80.6	68.1	138.1	102.6	134.2	111.7	99.7	119.8
Blue-winged teal	88.6	65.3	170.9	613.5	392.2	421.7	49.9	78.4	83.6	39.6
N. shoveler	38.1	26.8	10.3	27.0	40.2	50.4	19.6	14.9	10.0	7.3
N. pintail	18.1	22.7	27.9	48.5	58.3	72.7	42.7	14.9	23.7	14.1
Subtotal	1146.6	828.5	1855.6	2353.4	2389.0	3044.3	1503.7	1202.8	1341.2	898.9
Divers										
Redhead	29.6	11.6	27.4	49.4	12.7	49.6	11.3	5.2	7.5	0.8
Canvasback	33.1	25.0	42.1	55.2	18.3	99.4	5.0	11.7	40.0	3.1
Scaups	388.7	232.9	395.7	481.9	537.6	693.3	357.6	652.4	521.4	628.0
Ring-necked duck	172.1	239.5	184.2	277.0	242.1	393.0	338.7	221.6	139.1	110.7
Goldeneyes	62.6	66.6	13.8	39.3	141.5	213.4	107.5	89.4	155.6	103.8
Bufflehead	30.7	42.3	57.5	150.0	107.0	122.6	172.7	148.7	111.9	128.1
Ruddy Duck	7.9	5.0	12.4	37.1	1.3	10.0	0.0	3.9	1.0	0.0
Subtotal	724.8	622.8	733.1	1089.8	1060.5	1581.4	992.8	1133.0	976.5	974.4
Miscellaneous										
Oldsquaw	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scoters	17.2	39.6	33.2	81.0	126.7	35.5	46.3	92.3	60.0	37.2
Mergansers	138.9	68.1	159.3	118.6	153.5	144.4	100.6	118.9	125.8	128.8
Subtotal	156.0	107.7	192.5	199.6	280.2	179.9	146.9	211.2	185.7	166.0
Total Ducks	2027.4	1559.0	2781.2	3642.8	3729.6	4805.6	2643.5	2547.0	2503.4	2039.3
Canada Goose	96.3	44.6	77.8	69.7	119.7	48.2	57.9	48.7	29.6	29.1
Am. coot	15.5	8.2	29.3	193.7	20.5	47.9	17.3	13.6	41.7	7.8


Appendix 1, continued. Long-term trend in adjusted waterfowl breeding population estimates (thousands).

Species/Ponds	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
Ducks										
Dabblers										
Mallard	628.2	1074.0	891.0	1406.0	1282.0	968.7	1074.3	821.5	700.9	539.7
Am. black duck	0.0	0.6	6.3	2.6	0.7	1.1	1.0	0.0	0.5	0.0
Gadwall	4.5	8.9	27.0	9.4	4.4	17.8	50.9	13.0	15.0	8.1
Am. wigeon	172.1	247.8	246.7	337.0	272.3	308.4	313.2	232.9	278.9	242.8
Am. green-winged teal	104.4	172.5	165.8	383.7	296.3	393.3	166.7	137.5	174.7	127.0
Blue-winged teal	119.3	311.4	293.4	291.3	111.2	253.6	376.3	160.1	65.5	148.5
N. shoveler	25.3	10.3	56.5	17.9	14.5	27.9	40.2	16.6	18.1	17.8
N. pintail	40.5	35.2	38.9	58.3	25.7	21.6	41.6	16.3	15.2	14.7
Subtotal	1094.3	1860.7	1725.6	2506.2	2007.1	1992.4	2064.0	1398.0	1268.9	1098.7
Divers										
Redhead	3.1	17.0	8.8	7.0	7.6	10.0	27.0	8.0	22.9	7.7
Canvasback	16.1	22.9	28.8	12.4	8.6	24.3	44.9	19.8	13.5	16.5
Scaups	584.6	435.6	470.4	548.8	889.9	435.8	404.1	459.3	738.4	430.5
Ring-necked duck	207.2	279.6	187.2	266.6	167.1	332.8	445.2	283.3	284.1	352.2
Goldeneyes	81.0	242.5	182.5	85.6	66.1	48.8	331.8	110.0	163.6	105.5
Bufflehead	109.2	203.8	147.5	190.2	189.4	148.0	177.5	121.9	221.3	108.6
Ruddy Duck	0.0	4.3	0.7	4.9	4.0	3.4	27.1	1.5	17.1	0.0
Subtotal	1001.1	1205.7	1026.0	1115.6	1332.7	1003.1	1457.6	1003.8	1460.9	1021.0
Miscellaneous										
Oldsquaw	0.0	0.7	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scoters	52.6	78.6	89.2	88.3	78.9	108.9	86.4	119.2	198.0	64.0
Mergansers	262.1	114.3	181.3	296.3	152.3	282.0	263.8	147.5	302.6	273.0
Subtotal	314.7	193.6	271.6	384.6	231.2	390.9	350.2	266.7	500.6	337.0
Total Ducks	2410.1	3260.1	3023.2	4006.3	3571.0	3386.4	3871.8	2668.5	3230.4	2456.6
Canada Goose	31.1	66.3	50.5	168.1	116.4	80.3	96.8	52.6	93.6	64.9
Am. coot	21.0	50.2	228.5	55.9	12.6	324.3	526.1	33.3	24.5	94.8


Species/Ponds	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
Ducks										
Dabblers										
Mallard	509.6	413.9	673.5	913.1	839.1	853.9	947.9	1016.1	928.7	1005.7
Am. black duck	0.4	0.5	0.5	0.0	0.5	0.0	0.9	2.9	2.0	1.1
Gadwall	3.3	9.7	9.2	22.6	8.6	8.6	34.9	7.7	18.4	24.1
Am. wigeon	180.8	142.2	183.8	264.1	287.2	254.5	247.8	202.4	209.7	344.3
Am. green-winged teal	84.9	137.9	199.1	198.6	119.9	170.6	192.3	325.9	320.9	214.1
Blue-winged teal	236.1	65.9	127.0	485.6	281.1	170.4	357.0	374.4	338.0	132.2
N. shoveler	10.9	5.4	20.7	65.3	16.3	33.2	25.1	95.4	25.2	36.0
N. pintail	4.6	13.7	14.9	12.9	7.8	10.7	25.5	9.1	3.8	8.6
Subtotal	1030.5	789.3	1228.6	1962.1	1560.4	1502.0	1831.5	2033.8	1846.8	1766.1
Divers										
Redhead	8.1	12.0	10.0	14.1	2.1	5.7	1.4	7.7	1.1	19.6
Canvasback	14.1	4.6	7.5	31.2	14.6	23.3	16.1	94.7	16.1	9.5
Scaups	696.1	386.2	404.8	419.0	327.1	509.8	981.1	713.3	438.1	318.3
Ring-necked duck	341.5	288.5	445.0	291.2	181.0	278.3	332.6	834.5	373.0	323.1
Goldeneyes	66.4	236.8	145.7	39.6	40.9	57.1	104.4	328.7	178.4	112.6
Bufflehead	102.3	116.8	103.9	86.9	131.4	174.4	171.7	365.8	286.4	163.1
Ruddy Duck	0.7	5.2	5.1	2.8	5.7	28.8	10.7	7.2	0.0	0.0
Subtotal	1229.2	1049.9	1122.0	884.8	702.9	1077.4	1618.0	2351.9	1293.1	946.1
Miscellaneous										
Oldsquaw	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scoters	47.8	21.6	52.3	25.1	19.8	23.2	44.0	17.6	34.1	46.6
Mergansers	240.4	246.2	287.3	224.1	176.2	173.1	317.2	180.3	186.4	323.8
Subtotal	288.2	267.8	339.5	249.2	196.0	196.3	361.2	197.9	220.5	370.3
Total Ducks	2548.0	2107.1	2690.2	3096.2	2459.3	2775.7	3810.8	4583.7	3360.3	3082.6
Canada Goose	90.7	90.8	112.0	126.3	115.1	127.6	102.1	110.5	133.9	127.1
Am. coot	20.1	16.9	169.1	158.8	129.9	33.2	95.6	17.1	25.7	23.5

Species/Ponds	1995	1996	1997	1998	1999	2000	2001	2002	2003
Ducks									
Dabblers									
Mallard	1008.6	570.3	843.6	722.8	940.6	800.3	471.1	774.5	610.4
Am. black duck	0.0	0.0	0.0	0.0	3.7	0.0	0.0	0.0	0.0
Gadwall	24.6	4.3	15.1	4.7	7.4	6.0	6.9	14.1	19.0
Am. wigeon	156.0	144.1	163.3	157.7	173.7	114.0	84.8	175.1	188.0
Am. green-winged teal	275.7	193.6	284.4	228.6	255.8	158.7	165.2		257.7
Blue-winged teal	77.7	53.9	39.6	103.9	114.3	80.0	111.7	201.2	148.4
N. shoveler	53.5	25.0	32.7	19.6	18.1	18.1	11.9	19.1	3.7
N. pintail	25.9	2.8	22.0	7.8	7.5	14.8	6.3	9.0	5.2
Subtotal	1621.9	994.1	1400.7	1245.1	1521.1	1191.8	857.9	1450.7	1241.7
Divers									
Redhead	8.5	3.8	2.5	18.5	0.0	2.8	0.8	6.7	15.1
Canvasback	13.1	17.2	23.2	15.9	17.9	5.5	13.5	14.5	5.0
Scaups	391.3	390.6	488.9	204.9	494.9	269.7	239.0	354.5	324.6
Ring-necked duck	276.5	191.0	274.7	280.6	383.2	224.8	285.1	527.8	319.3
Goldeneyes	335.9	264.7	338.2	271.4	251.5	234.1	446.3	368.7	265.2
Bufflehead	156.9	135.5	200.0	211.7	234.6	224.7	191.1	236.0	246.5
Ruddy Duck	5.0	2.8	9.7	0.0	3.8	0.0	5.1	7.1	15.7
Subtotal	1187.1	1005.6	1337.3	1003.0	1386.0	961.4	1180.9	1515.3	1191.4
Miscellaneous									
Oldsquaw	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Eiders	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0
Scoters	63.4	40.6	118.4	55.5	68.4	55.3	29.0	55.7	57.4
Mergansers	257.3	169.6	241.0	179.1	192.9	200.0	172.3	308.5	203.5
Subtotal	320.7	210.2	359.4	234.6	261.3	255.3	202.2	364.2	260.8
Total Ducks	3129.7	2209.9	3097.4	2482.7	3168.4	2408.6	2241.0	3330.2	2694.0
Canada Goose	136.1	169.8	109.6	143.3	286.6	168.6	165.9	213.6	170.2
Am. coot	17.6	11.8	13.6	29.6	38.0	83.9	43.0	101.5	6.1


Strata 21-24 Mergansers




Strata 21-24 Mallard




Strata 21-24 American black duck




Strata 21-24 Gadwall




Strata 21-24 American wigeon




Strata 21-24 American green-winged teal




Strata 21-24 Blue-winged teal




Strata 21-24 Northern shoveler




Strata 21-24 Northern pintail




Strata 21-24 Redhead




Strata 21-24 Canvasback




Strata 21-24 Scaups




Strata 21-24 Ring-necked duck




Strata 21-24 Goldeneyes




Strata 21-24 Bufflehead




Strata 21-24 Long-tailed duck




Strata 21-24 Eiders




Strata 21-24 Scoters




Strata 21-24 Ruddy Duck




Strata 21-24 Canada Goose




Strata 21-24 American coot




Strata 21-24 Dabblers



Strata 21-24 Divers



Strata 21-24 Miscellaneous



Strata 21-24 Total Ducks

