

STATE AND FEDERAL THREATENED, ENDANGERED, AND OTHER SPECIES OF CONCERN
 LIKELY TO OCCUR IN HOLMES COUNTY FLORIDA
 Compiled by U.S. Fish and Wildlife Service April 2008

Common Name	Scientific Name	Status State	Status FWS	Natural Communities
FISH:				
Gulf sturgeon	<i>Acipenser oxyrinchus desotoi</i>	SSC	T CH	ESTUARINE: various MARINE: various habitats RIVERINE: alluvial and blackwater streams
Bluenose shiner	<i>Pteronotropis welaka</i>	SSC		RIVERINE: blackwater, alluvial, and spring-run streams
AMPHIBIANS & REPTILES:				
Flatwoods salamander	<i>Ambystoma cingulatum</i>	SSC	T CH(p)	PALUSTRINE: wet flatwoods, dome swamp, basin swamp, ruderal TERRESTRIAL: mesic flatwoods (reproduces in ephemeral wetlands within this community)
Eastern indigo snake	<i>Drymarchon corais couperi</i>	T	T	ESTUARINE: tidal swamp PALUSTRINE: hydric hammock, wet flatwoods TERRESTRIAL: mesic flatwoods, upland pine forest, sandhills, scrub, scrubby flatwoods, rockland hammock, ruderal
Coal skink	<i>Eumeces anthracinus</i>		ce	PALUSTRINE: seepage slope, baygall TERRESTRIAL: upland pine forest, upland hardwood forest, mesic flatwoods
Gopher tortoise	<i>Gopherus polyphemus</i>	SSC	ce	TERRESTRIAL: sandhills, scrub, scrubby flatwoods, xeric hammocks, coastal strand, ruderal
Pine barrens treefrog	<i>Hyla andersonii</i>	SSC		PALUSTRINE: seepage slope, baygall RIVERINE: seepage stream
Alligator snapping turtle	<i>Macroclmys temminckii</i>	SSC	ce	ESTUARINE: tidal marsh LACUSTRINE: river floodplain lake, swamp lake RIVERINE: alluvial stream, blackwater stream
Florida pine snake	<i>Pituophis melanoleucus mugitus</i>	SSC	ce	LACUSTRINE: ruderal, sandhill upland lake TERRESTRIAL: sandhill, scrubby flatwoods, xeric hammock, ruderal
Gopher frog	<i>Rana capito</i>	SSC	ce	TERRESTRIAL: sandhill, scrub, scrubby flatwoods, xeric hammock (reproduces in ephemeral wetlands within these communities)
BIRDS:				
Bachman's sparrow	<i>Aimophila aestivalis</i>		ce	TERRESTRIAL: various, ruderal
Little blue heron	<i>Egretta caerulea</i>	SSC		ESTUARINE: marshes, shoreline PALUSTRINE: floodplains, swamps RIVERINE: shoreline
Snowy egret	<i>Egretta thula</i>	SSC		ESTUARINE: marshes, tidal swamps, shoreline LACUSTRINE: lake edges PALUSTRINE: swamp, floodplain, ruderal RIVERINE: shoreline

E=endangered, T=threatened, P=proposed, C=candidate, s/a=similar appearance, SSC=species of special concern, ce=consideration encouraged, CH=Critical Habitat, BGEPA=Bald and Golden eagle protection act

This is not an exhaustive list of where species do occur, but a guide to indicate areas that might require surveys if appropriate habitat exists. Please contact Florida Natural Areas Inventory (850-224-8207) for additional species location information.

STATE AND FEDERAL THREATENED, ENDANGERED, AND OTHER SPECIES OF CONCERN
 LIKELY TO OCCUR IN HOLMES COUNTY FLORIDA
 Compiled by U.S. Fish and Wildlife Service April 2008

Common Name	Scientific Name	Status State	Status FWS	Natural Communities
Tricolored heron	<i>Egretta tricolor</i>	SSC		ESTUARINE: marshes, tidal swamps, shoreline LACUSTRINE: lake edges PALUSTRINE: swamp, floodplain, ruderal RIVERINE: shoreline
Arctic peregrine falcon	<i>Falco peregrinus tundrius</i>	E	ce	ESTUARINE: winters along coasts LACUSTRINE: various PALUSTRINE: various TERRESTRIAL: various, ruderal
Southeastern kestrel	<i>Falco sparverius paulus</i>	T	ce	ESTUARINE: various habitats PALUSTRINE: various habitats TERRESTRIAL: open pine forests, clearings, ruderal, various
Wood stork	<i>Mycteria americana</i>	E	E	ESTUARINE: marshes LACUSTRINE: floodplain lakes, marshes (feeding), various PALUSTRINE: marshes, swamps, various
Red-cockaded woodpecker	<i>Picoides borealis</i>	SSC	E	TERRESTRIAL: mature pine forests
Black Skimmer	<i>Rynchops niger</i>	SSC		ESTUARINE: various LACUSTRINE: various RIVERINE: various TERRESTRIAL: ocean beaches, beach dune, ruderal. Nests common on rooftops.
MAMMALS:				
Gray bat	<i>Myotis grisescens</i>	E	E	PALUSTRINE: caves, various TERRESTRIAL: caves, various
Southeastern big-eared bat	<i>Plecotus rafinesquii</i>		ce	PALUSTRINE: various, floodplains TERRESTRIAL: pine and hardwood forests, ruderal, various
Eastern chipmunk	<i>Tamias striatus</i>	SSC		TERRESTRIAL: slope forest, upland hardwood forest, upland pine forest
Florida black bear	<i>Ursus americanus floridanus</i>	T	ce	PALUSTRINE: titi swamps, floodplains TERRESTRIAL: pine and hardwood forests
INVERTEBRATES:				
Rayed creekshell (mussel)	<i>Anodontooides radiatus</i>		ce	RIVERINE: Small to medium sized creeks in substrates of mud, sandy mud, or sand and gravel (Panhandle watersheds: Apalachicola, Chipola, Escambia, Choctawhatchee)
Southern sandshell	<i>Hamiota australis</i>		C	Riverine: found in small to medium-sized creeks and rivers in with slow to moderate currents over sandy substrates sometimes with some silt. Endemic to the Escambia, Yellow, and Choctawhatchee River drainages of Alabama and Florida
Fuzzy pigtoe	<i>Pleurobema strodeanum</i>		C	Riverine: small to medium-sized creeks and rivers with slow to moderate currents in sand and sand with some silt. Endemic to the Escambia, Yellow, and Choctawhatchee River drainages of Alabama and Florida.

E=endangered, T=threatened, P=proposed, C=candidate, s/a=similar appearance, SSC=species of special concern, ce=consideration encouraged, CH=Critical Habitat, BGEPA=Bald and Golden eagle protection act

This is not an exhaustive list of where species do occur, but a guide to indicate areas that might require surveys if appropriate habitat exists. Please contact Florida Natural Areas Inventory (850-224-8207) for additional species location information.

STATE AND FEDERAL THREATENED, ENDANGERED, AND OTHER SPECIES OF CONCERN
 LIKELY TO OCCUR IN HOLMES COUNTY FLORIDA
 Compiled by U.S. Fish and Wildlife Service April 2008

Common Name	Scientific Name	Status State	Status FWS	Natural Communities
Tapered pigtoe	<i>Fusconaia burkei</i>		C	Riverine: Small to medium-sized creeks to large rivers in stable sand or sand and gravel substrata, occasionally occurring in silty sand in slow to moderate current. Endemic to the Choctawhatchee River drainage of Alabama and Florida.
Choctaw bean	<i>Villosa choctawensis</i>		C	Riverine: Small to large creeks and rivers with moderate current over sand to silty-sand substrates. Endemic to the Escambia, Yellow, and Choctawhatchee River drainages of Alabama and Florida.
Downy rainbow (mussel)	<i>Villosa villosa</i>		ce	RIVERINE: small streams to large rivers in sand or muddy sand substrates (Panhandle watersheds: Apalachicola, Chipola, Escambia, Choctawhatchee, Ochlockonee, Suwannee)
PLANTS:				
Apalachicola wild indigo	<i>Baptisia megacarpa</i>	E		PALUSTRINE: floodplain forest TERRESTRIAL: upland mixed forest, slope forest
Mountain laurel	<i>Kalmia latifolia</i>	T		RIVERINE: seepage stream bank TERRESTRIAL: slope forest, seepage stream banks
Cucumber magnolia	<i>Magnolia acuminata</i>	E		TERRESTRIAL: slope forest, upland mixed forest
Primrose-flower butterwort	<i>Pinguicula primulifolia</i>	E		PALUSTRINE: bogs, pond margins, margins of spring runs
Yellow fringed orchid	<i>Platanthera ciliaris</i>	T		PALUSTRINE: bogs, wet flatwoods TERRESTRIAL: Bluff
Orange azalea	<i>Rhododendron austrinum</i>	E		PALUSTRINE: bottomland forest RIVERINE: seepage stream bank TERRESTRIAL: slope forest, upland mixed forest
White-top pitcher plant	<i>Sarracenia leucophylla</i>	E	ce	PALUSTRINE: wet prairie, seepage slope, baygall edges, ditches
Parrot pitcher plant	<i>Sarracenia psittacina</i>	T		PALUSTRINE: wet flatwoods, wet prairie, seepage slope
Decumbant pitcher plant	<i>Sarracenia purpurea</i>	T		PALUSTRINE: Bogs

E=endangered, T=threatened, P=proposed, C=candidate, s/a=similar appearance, SSC=species of special concern, ce=consideration encouraged, CH=Critical Habitat, BGEPA=Bald and Golden eagle protection act

This is not an exhaustive list of where species do occur, but a guide to indicate areas that might require surveys if appropriate habitat exists. Please contact Florida Natural Areas Inventory (850-224-8207) for additional species location information.