

Maricopa County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Arizona cliffrose	<i>Purshia subintegra</i>	Endangered	Evergreen shrub of the rose family (Roseaceae). Bark pale gray and shreddy. Young twigs covered with dense hairs. Leaves have 1-5 lobes and edges curl downward (revolute). Flowers: 5 petals, white or yellow <0.5 inches long.	Graham, Maricopa, Mohave, Yavapai	< 4,000 ft	White limestone soils derived from tertiary lakebed deposits.	Occurs in central Arizona at Horseshoe Lake, in the Burro Creek drainage, and near Cottonwood in the Verde Valley.
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large, adults have white head and tail. Height 28-38 inches; wingspan 66-96 inches. Dark with varying degrees of mottled brown plumage. Feet bare of feathers.	Gila, Graham, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	Varies	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Some birds are nesting residents while a larger number winters along rivers and reservoirs. Once endangered (32 FR 4001, 03-11-1967; 43 FR 6233, 02-14-78) because of reproductive failures from pesticide poisoning and loss of habitat, this species was downlisted to threatened on August 11, 1995, and delisted August 8, 2007. Threatened status reinstated for Desert nesting bald eagles.
California Brown pelican	<i>Pelecanus occidentalis californicus</i>	Proposed delisted	Large, dark gray-brown water bird with webbed feet, pouch underneath its long bill, and wingspan of 7 ft. Adults have a white head and neck, brownish black breast, and silver gray upper parts.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	Varies	Coastal land and islands; species found occasionally around Arizona's lakes and rivers.	Subspecies endangered due to pesticides. Considered an uncommon transient in Arizona. Most observations recorded along the Colorado River and in the Gila Valley. Individuals known to wander up from Mexico in summer and fall. There are no breeding records for Arizona. Populations also exist in California and Mexico.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
California Least Tern	<i>Sterna antillarum browni</i>	Endangered	Least terns are smallest of the North American Terns. Body length is 21 to 24 cm (8 to 9 inches) with a wingspan of 45 to 51cm (18 to 20 inches). Characterized by a black crown and loreal stripe on their head, snowy white forehead and underside, and gray upperparts. Outer two primaries are black, bill is yellow or orange with black tip, and legs are orange. Males have a wider dark loreal stripe but sexes are mostly distinguished by behavior. Immatures have darker plumage, dark bill, and dark eye strips on white heads.	Maricopa, Mohave, Pima	< 2,000 ft	Open, bare or sparsely vegetated sand, sandbars, gravel pits, or exposed flats along shorelines of inland rivers, lakes, reservoirs, or drainage systems.	Breeding occasionally documented in Arizona; migrants may occur more frequently. Feeds primarily on fish in shallow waters and secondarily on invertebrates. Nests in a simple scrape on sandy or gravelly soil.
Desert pupfish	<i>Cyprinodon macularius</i>	Endangered	Small (2 inches) smoothly rounded body shape with narrow vertical bars on the sides. Breeding males blue on head and sides with yellow on tail. Females and juveniles tan to olive colored back and silvery sides.	Cochise, Graham, Maricopa, Pima, Pinal, Santa Cruz, Yavapai	< 4,000 ft	Shallow springs, small streams, and marshes. Tolerates saline and warm water.	Two subspecies are recognized: Desert Pupfish (<i>C.m. macularis</i>) and Quitobaquito Pupfish (<i>C.m. eremus</i>). Critical habitat includes Quitobaquito Springs, Pima County, portions of San Felipe Creek, Carrizo Wash, and Fish Creek Wash, Imperial County, California.
Gila topminnow	<i>Poeciliopsis occidentalis occidentalis</i>	Endangered	Small (2 inches), guppy-like, live bearing, lacks dark spots on its fins. Breeding males are jet black with yellow fins.	Cochise, Gila, Graham, Maricopa, Pima, Santa Cruz, Yavapai	< 4,500 ft	Small streams, springs, and cienegas vegetated shallows.	Species historically also occurred in backwaters of large rivers but is currently isolated to small streams and springs.
Lesser long-nosed bat	<i>Leptonycteris curasoae yerbabuena</i>	Endangered	Elongated muzzle, small leaf nose, and long tongue. Yellowish brown or gray above and cinnamon brown below. Tail minute and appears to be lacking. Easily disturbed.	Cochise, Gila, Graham, Greenlee, Maricopa, Pima, Pinal, Santa Cruz, Yuma	1,600-11,500 ft	Desert scrub habitat with agave and columnar cacti present as food plants.	Day roosts in caves and abandoned tunnels. Forages at night on nectar, pollen, and fruit of paniculate agaves and columnar cacti. This species is migratory and is present in Arizona usually from April to September and south of the border the remainder of the year.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Medium sized with dark eyes and no ear tufts. Brownish and heavily spotted with white or beige.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai	4,100-9,000 ft	Nests in canyons and dense forests with multi-layered foliage structure.	Generally nest in older forests of mixed conifer or ponderosa pine/gambel oak type, in canyons, and use variety of habitats for foraging. Sites with cool microclimates appear to be of importance or are preferred. Critical habitat was finalized on August 31, 2004 (69 FR 53182) in Arizona in Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Navajo, Pima, Pinal, Santa Cruz, and Yavapai counties.
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6,000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Big River fish also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam; Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizona/New Mexico border to Coolidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.
Sonoran pronghorn	<i>Antilocapra americana sonoriensis</i>	Endangered	Upperparts tan; underparts, rump, and two bands across the neck are white. Male has two black cheek pouches. Hoofed with slightly curved black horns having a single prong. Smallest and palest of the pronghorn subspecies.	Maricopa, Pima, Yuma	2,000-4,000 ft	Broad intermountain alluvial valleys with creosote-bursage and palo verde-mixed cacti associations.	Typically, bajadas are used as fawning areas and sandy dune areas provide food seasonally. Cacti (jumping cholla) appears to make up substantial part of diet. This subspecies also occurs in Mexico.
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 8,500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886). In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Woundfin	<i>Plagopterus argentissimus</i>	Endangered	Small (4 inches) silver minnow with fairly large fins and a sharp dorsal fin spine.	Mohave and Maricopa	< 4,500 ft	Inhabits shallow, warm, turbid, fast-flowing water. Tolerates high salinity.	Native population only in Virgin River. Designated critical habitat includes the Virgin River and its 100-year floodplain. Experimental non-essential populations (50 FR 30193, 07-24-1985) designated in portions of the Verde, Gila, San Francisco, and Hassayampa rivers and Tonto Creek. Species also occurs in Washington County, UT and Clark County, NV.
Yuma clapper rail	<i>Rallus longirostris yumanensis</i>	Endangered	Water bird with long legs and short tail. Long, slender decurved bill. Mottled brown or gray on its rump. Flanks and undersides are dark gray with narrow vertical stripes producing a barring effect.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	< 4,500 ft	Fresh water and brackish marshes.	Species is associated with dense emergent riparian vegetation. Requires wet substrate (mudflat, sandbar) with dense herbaceous or woody vegetation for nesting and foraging. Channelization and marsh destruction are primary sources of habitat loss.
Roundtail Chub	<i>Gila robusta</i>	Candidate	Member of the minnow family Cyprinidae and characterized by streamlined body shape. Color usually olive gray with silvery sides and a white belly. Breeding males develop red or orange coloration on the lower half of the cheeks and on the bases of paired fins. Individuals may reach 49.0 cm (19.3 in) but usually average 25-30 cm (9.8 - 11.8 in).	Apache, Coconino, Gila, Graham, Greenlee, LaPaz, Maricopa, Mohave, Navajo, Pinal, and Yavapai	1,000-7,500 ft.	Cool to warm waters of rivers and streams, often occupy the deepest pools and eddies of large streams.	Historical range of roundtail chub included both the upper and lower Colorado River basins. A 2009 status review determined that the lower Colorado River basin roundtail chub population segment (Arizona and New Mexico) qualifies as a distinct vertebrate population segment (DPS). Populations in the Little Colorado, Bill Williams, and Gila River basins are considered candidate species.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill that is blue-black with yellow on the lower half. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Neotropical migrant that winters primarily in South America and breeds primarily in the U.S. (but also in southern Canada and northern Mexico). As a migrant it is rarely detected; can occur outside of riparian areas. Cuckoos are found nesting statewide, mostly below 5,000 feet in central, western, and southeastern Arizona. Concern for cuckoos are primarily focused upon alterations to its nesting and foraging habitat. Nesting cuckoos are associated with relatively dense, wooded, streamside riparian habitat, with varying combinations of Fremont cottonwood, willow, velvet ash, Arizona walnut, mesquite, and tamarisk. Some cuckoos have also been detected nesting in velvet mesquite, netleaf hackberry, Arizona sycamore, Arizona alder, and some exotic neighborhood shade trees.
American peregrine falcon	<i>Falco peregrinus anatum</i>	Delisted	A crow-sized falcon with slate blue-gray on the back and wings, and white on the underside; a black head with vertical "bandit's mask" pattern over the eyes; long pointed wings; and a long wailing call made during breeding. Very adept flyers and hunters, reaching diving speeds of 200 mph.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	3,500-9,000 ft	Areas with rocky, steep cliffs, primarily near water, where prey (primarily shorebirds, songbirds, and waterfowl) concentrations are high. Nests are found on ledges of cliffs, and sometimes on man-made structures such as office towers and bridge abutments.	Species recovered with over 1,650 breeding birds in the US and Canada.
Arizona agave	<i>Agave arizonica</i>	Delisted	Member of the agave family. Has rosettes of bright green leaves, 17-24cm long and 2-4cm wide, broadest in the middle. Flowers are small, pale yellow, and jar shaped.	Gila, Maricopa, Yavapai	3,600-5,800 ft	Occurs on open slopes in chaparral or juniper grasslands. Prefers shallow, cobbled, and gravelly soils on steep slopes.	Arizona agave is a hybrid produced by a crossing of two other common agave species (<i>A. chrysantha</i> x <i>A. toumeyana</i> ssp. <i>toumeyana</i>).