

Annual Report 2011

Project No. 1982-013-03
Annual Stock Assessment - CWT (USFWS)

By
Stephen M. Pastor

U.S. Fish and Wildlife Service
Columbia River Fisheries Program Office
1211 SE Cardinal Court, Suite 100
Vancouver, WA 98683
February 2013

Funded By
Bonneville Power Administration

INTRODUCTION

In 1989 the Bonneville Power Administration (BPA) began funding the evaluation of production groups of juvenile anadromous fish not being coded-wire tagged for other programs. These groups were the "Missing Production Groups". Production fish released by the U.S. Fish and Wildlife Service (FWS) without representative coded-wire tags during the 1980s are indicated as blank spaces on the survival graphs in this report. This program is now referred to as "Annual Stock Assessment - CWT"

The objectives of the "Annual Stock Assessment" program are to:

- estimate the total survival of each production group,
- estimate the contribution of each production group to fisheries, and
- prepare an annual report for USFWS hatcheries in the Columbia River basin.

Coded-wire tag recovery information will be used to evaluate the relative success of individual hatchery brood stocks. This information can also be used by salmon harvest managers to develop plans to allow the harvest of excess hatchery fish while protecting threatened, endangered, or other stocks of concern.

All fish release information, including marked/unmarked ratios, is reported to the Pacific States Marine Fisheries Commission (PSMFC). Fish recovered in the various fisheries or at the hatcheries are sampled to recover coded-wire tags. This recovery information is also reported to PSMFC.

This report has been prepared annually starting with the report labeled "Annual Report 1994". Although the current report has the title "Annual Report 2010", it was written in fall of 2012 through early winter 2013 using data available from RMIS at that time; and submitted as final in March 2013. The main objective of the report is to evaluate survival of groups of hatchery fish which have been tagged under this ongoing project.

METHODS

The Annual Stock Assessment Report reports coded-wire tagging for the most recently completed contract period, and evaluates the survival and contribution of fish tagged, raised and released at National Fish Hatcheries in the Columbia River Basin.

Release information used in this report is collected with the U.S. Fish and Wildlife Service Columbia River (information) System (CRiS). Information prior to the Columbia River Basin wide implementation of CRiS, and from USFWS hatcheries in Idaho, is obtained from the interagency Regional Mark Information System.

Lists of coded-wire tags are obtained from the CRiS sr80s file by use of the CWTetc program. This currently involves the additional step of downloading Dworshak and Kooskia release information from the RMIS database, and adding it to the sr80s file. In past years numerous discrepancies have been noted between the original reports of releases, and those in RMIS.

Recovery information is obtained from the PSMFC Regional Mark Information System (RMIS) coded-wire tag database web site. This database is continuously being updated by the contributing agencies: California Fish and Game (CDFG), Canadian Department of Fisheries and Oceans (CDFO), Oregon Department of Fisheries and Wildlife (ODFW), Washington Department of Fisheries and Wildlife (WDFW), and US Fish and Wildlife Service (FWS); and the updates are reflected annually in this report.

Coded-wire tag recovery information is retrieved in the TS1 (Tag Summary) format. The TS1 report contains both the observed number of recoveries, and the estimated number of recoveries based on the sampling rate. The TS1 report is used precisely because it is a Summary. The number of individual recoveries, and the number of individual recovery sites preclude there use in this report. The TS1 report summarizes individual recoveries and recovery sites into broad categories based on combinations of geographic and other criteria, and provide a consistent means of reporting from year to year.

It should also be noted that fisheries in some tributaries such as the Clackamas River, and its tributary Eagle Creek, are poorly sampled (or not at all) for recovery of coded-wire tags. State agencies often rely on harvest estimates based on sport license returns in these tributaries, and this information is not available through RMIS and the TS1 format.

TS1 reports were downloaded from RMIS in fall of 2012. These TS1 reports are downloaded in ASCII format. A dBASE V program transforms these ASCII files into a single dbf file.

A Stock Assessment Reference Document is prepared for each hatchery, brood year, and species. Because many fish were released without representative coded-wire tags before the Annual Stock Assessment program began, a single Production Expansion Factor (PEF - the total number of fish released divided by the total number of tagged fish released) is calculated for each hatchery, brood year, species, and stage of fish released. This PEF is used to expand estimated recovery information for untagged fish released along with tagged fish, and to determine the overall contribution and survival rate for each facility.

Until the late 1990s, the adipose fin was removed from all fish receiving a coded-wire tag. The missing adipose fin facilitated the recovery of tags from returning or harvested jacks and adults.

In the late 1990s the goal of protecting weak, threatened, or endangered populations of fish, while maintaining fisheries on healthy populations, led to the removal of the adipose fin from most hatchery fish. The "mass marking" of hatchery steelhead, Coho, and Chinook makes it possible to harvest only hatchery fish. Wild fish with adipose fins are not harvested in "selective fisheries". In order to assess the success of selective fisheries, and to serve as an "index" for wild fish populations, some hatchery fish are coded-wire tagged, but not adipose fin clipped. These fish will not be harvested in any selective fisheries that may occur. A dbase program was written to expand recoveries for each coded-wire tag, rather than calculating a Production Expansion Factor. This program is used for all current brood years, and has now been used for many older brood years beginning with brood year 1990.

Stock Assessment Reference Summary printouts list the following information for each brood year, species, hatchery, and stage released at the hatchery: release information, the total number of observed recoveries, where recoveries occurred, the number of expanded recoveries from the PSMFC TS1 report, the number of recoveries expanded to include unmarked fish released, and a summary of where fish were recovered. The most recent brood years have been reprocessed, and all brood years for all hatcheries and species are included in this report. Text describing the major attributes of the contribution for each hatchery, species, and brood year is prepared and included in the report. Graphs of survival and relative proportion of recoveries in major areas are prepared and included in the report.

In November 2008 the program that summarizes data for the Fisheries graphs was amended to add harvest in the Estuary Sport fishery to the Columbia R. Sport numbers.

"Residualized" fish, or "mini-jacks" from yearling releases, are not included in estimates of survival.

In 2009 the SA program was refined to perform the second expansion, which "corrects" for the fact that not all released fish are coded-wire tagged, and expands total recovery for the untagged and/or unmarked fish for each coded-wire tag, recovery site, and year. The effect of this change was noticeable even at the Warm Springs NFH facility, where 100% of released fish are coded-wire tagged before being released.

The SAsumRY.prg creates a file imported into the Releases and Returns tables that follow the graphs of survival rates, and fisheries for each hatchery and species.

RESULTS

Four hundred and seven "Annual Stock Assessment" tags were recovered at National Fish Hatcheries in 2012, while the number of fish tagged under this program was 175,751.

A brief description of the estimated survival and contribution for each species raised and released at national fish hatcheries follows. These descriptions are followed by graphs for each hatchery and species.

Abernathy Fish Technology Center

Abernathy Fish Technology Center (FTC) is located 14 miles west of Longview, Washington, on Abernathy Creek, approximately three miles upstream from its confluence with the Columbia River at an altitude of 175 feet above sea level. Abernathy FTC began operations in 1959. From 1980 through the 1990s, lower river tule fall Chinook were the only species reared on a production basis at Abernathy as part of Mitchell Act (NOAA Fisheries) funding. Information of these tules can be found at the end of this report.

Columbia River Program Fisheries Office marking crews have been coded-wire tagging steelhead at Abernathy since January 2003. CRFPO personnel have also read wire from recovered adults. The RMIS database contains one recovery from these tagged fish.

Carson National Fish Hatchery

Carson National Fish Hatchery (NFH) is located 13 miles northwest of Carson, Washington, at the confluence of the Wind River and Tye Springs (River Mile 18), at 1,180 feet above sea level. Carson NFH began operation in 1938 by rearing and releasing fall Chinook salmon and trout.

In 1956, the hatchery was remodeled under the Mitchell Act. Construction of a fish ladder at Shipperd Falls (River Mile 2.1) in 1955 made it possible for spring Chinook to pass upstream to the hatchery. Trapping of spring Chinook at Bonneville Dam began in 1955, after Washington shore trapping facilities were completed, and continued through 1961. Those trapped fish were used to establish the Carson run of spring Chinook in the Wind River. Other species such as steelhead, brook trout, rainbow trout, kokanee, and coho have also been raised at Carson. Coho were raised as late as 1981. Current production involves adult collection, egg incubation and rearing of spring Chinook salmon.

Please note that sport and tribal harvest based on recovery of coded wire tags in the Columbia River as well as the tributaries (for example Wind River) are all reported as Columbia River harvest.

Brood year 1982 to 1985 spring Chinook from Carson were marked with coded-wire tags for a density study. The average total survival of these brood years is 0.22%. The majority of off station recoveries were in the freshwater sport fishery. There were an estimated 20 recoveries of brood year 1982 and 1983 fish in Canadian waters. An estimated 43 recoveries of brood year 1985 fish occurred in ocean fisheries in Alaska, Canada, and Washington.

Because of a WDFW fish marking study at Carson, WDFW sampled returning adults and originally reported recoveries for return years 1993 through 1996. During the change over from format 3.2 to 4.0, USFWS reported these recoveries.

The survival rate for Carson brood year 1988 is 0.4044% using the rd2 program which expands each coded-wire tag, rather than using a PEF, and including one additional expanded freshwater sport recovery reported by WDFW. The majority of contribution to fisheries was in the Columbia River freshwater sport, expanded to 5,339. Recoveries in the treaty ceremonial fisheries were expanded to a catch of 1,023.

Brood year 1989 was also part of the fish marking study conducted by WDFW. The estimated survival for this brood year is 0.1368%, less than a third of the previous brood year. The majority of fish taken off station (2,095) were taken in the freshwater sport fishery reported by WDFW. Two observed recoveries in the Washington ocean troll fishery were expanded to a total of 12 fish caught.

The survival rate for brood year 1990 is 0.0778%, a total of 1,801 fish from a release of over 2 million. A third of the fish entering the Columbia River were harvested before reaching Carson. Most of these recoveries, estimated at 125, were freshwater sport reported by WDFW. Treaty ceremonial fishery recoveries reported by ODFW show 32 fish taken, followed by 27 in the Columbia River sport fishery.

Brood year 1991 survival is now estimated to be 0.0239%. This is less than brood year 1990 and the lowest to date. The largest off station take of fish was the ODFW reported treaty ceremonial harvest of 69 fish. Four fish were recovered in WDFW spawning ground surveys, expanded to 12. This was the last brood year for the WDFW mark study.

Survival for brood year 1992 is 0.6145%, the highest since brood year 1988. An estimated 82 fish were harvested in the California ocean troll fishery, an unusual event. Since all of the coded-wire tags released with brood year 1992 were used for a density study, none of the tag codes represents the majority of fish released. Recoveries for this brood year at Carson NFH were originally reported by WDFW.

The 1993 brood year now has the fifth highest survival for this hatchery at 0.4371%. An estimated 42 fish were harvested in the California ocean troll fishery as jacks. The harvest of these fish off of California occurred the same year as the harvest of 82 fish from brood year 1992. WDFW freshwater sport fishery recoveries are expanded to a harvest of 4,494 fish, nearly as many as returned to the hatchery. Since all of the coded-wire tags released with brood year 1993 were used for a new density study, none of the tag codes represents the majority of the fish released.

Brood year 1994 survival is now estimated at 0.1243%. There are an expanded 882 fresh water sport recoveries, 1,082 expanded returns to hatcheries, and 165 expanded spawning ground recoveries. This equals a total of 2,141 expanded recoveries for the 1994 brood year.

There are an additional 4 PSC expanded WDFW freshwater sport recoveries this year, increasing the estimated survival for brood year 1995 to 0.3770%, in the upper third of successful brood years

for Carson NFH. There have been an estimated 997 expanded off station recoveries from these fish, all in the Columbia Basin.

The estimated survival for Carson **brood year 1996** is 1.1039% with an estimated 9,097 off station recoveries, and 10,047 returns to the hatchery. This brood year is one of the most successful yet for Carson NFH. Over seventy-five hundred fish were harvested in the WDFW reported freshwater sport fishery. The second greatest harvest was for treaty ceremonial (859).

Brood year 1997 has one of the highest survival rates yet for Carson NFH spring Chinook at 1.2826%. Over ten thousand fish were harvested in the Columbia River, including eighty two hundred in sport fisheries, and 1,353 in the Columbia River gill net fishery, and 259 for treaty ceremonial. Fourteen fish were also harvested in British Columbia.

Brood year 1998 total survival is estimated to be 1.3635%. Nearly thirteen thousand fish were harvested in the Columbia River Basin. The majority of the harvest was sport - over ten thousand - followed by a gill net harvest of 1,960.

The estimated survival for **brood year 1999** is 1.446%, which is nearly equal to that of the 1998 brood year. Nearly thirteen thousand four hundred fish were harvested in the Columbia River with a sport harvest of 9,450.

With additional Age5 FWS and WDFW recoveries **brood year 2000** Carson fish have an overall survival of 0.6967%. This brood year contributed seven thousand one hundred twenty-two to Columbia River fisheries: 1,200 in the gill net fishery, nearly six thousand to fisheries within the Columbia Basin. An additional 26 fish were taken in both the Treaty Ceremonial, and Ocean Troll fishery in Canada. Coded-wire tag recoveries expand to a return of 2,949 fish to Carson NFH, close to the Age Composition derived estimate of 3,044.

Brood year 2001 survival is about 1/3 of that of the previous brood year at 0.2551%. Approximately two thirds of this return was harvested within the Columbia Basin, mostly as sport catch: 1,184 reported by WDFW and 750 by ODFW. Gill nets took an estimated 483 fish, and 31 fish were caught in treaty fisheries.

Brood year 2002 now has an estimated total survival rate of 0.3615%, below the average to date of 0.4788%, but higher than that of the previous brood year. 2,451 fish were harvested in all freshwater sport fisheries with most (1,990) being taken in Freshwater Sport Fisheries.

A handful of new recoveries results in an estimated of survival 0.1922% for **brood year 2003**, less than half of the mean for all Carson brood years (0.4788%). Recoveries occurred primarily in the ODFW reported Columbia River and Freshwater Sport fisheries, a combined 1,011. Another 73 fish went to the gillnet fishery, and 14 to treaty ceremonial.

Brood year 2004's 0.5434% survival is above the "to date" average of 0.52%. While the freshwater sport fishery took nearly half the harvest (2,084), gillnets and Columbia River Sport had nearly equal shares of 1,444 and 1,316 fish respectively. One observed Treaty Ceremonial recovery was expanded to 20 fish.

Freshwater Sport fishing took the greatest share of **brood year 2005** off station recoveries, 1,696. Gillnet, Columbia River Sport, and net fisheries took 357, 970, and 192 fish each. Overall survival is estimated at 0.4446%.

Survival for brood year 2006 jumped to 1.3918%, essentially the same as the previous high of 1.398% of brood year 1998. Over twelve thousand fish were harvested in Columbia River fisheries: 5,770 in gillnets with ODFW reporting 2,290 sport recoveries and WDFW reporting 4,120. Another 6,102 by this estimate made it back to National fish hatcheries.

Dworshak National Fish Hatchery

Dworshak NFH is located at the confluence of the North Fork Clearwater River and the main stem Clearwater River about three miles west of Orofino, Idaho, at 1,000 feet above sea level. Dworshak NFH first began operations in 1969 rearing summer steelhead and resident trout to mitigate for fish lost by the construction and operation of Dworshak Dam. The facility was expanded in 1982 under the Lower Snake River Compensation Program (LSRCP) to rear spring Chinook salmon. The hatchery is now used to produce spring Chinook and summer steelhead.

Spring Chinook

Total recoveries for **brood year 1985** were 0.0201%, the first release of 100% Rapid River stock. The **brood year 1986** yearling release resulted in an estimated 2,595 fish, or 0.2306%.

Twelve different coded-wire tag groups were released with **brood year 1987**. Only three were in yearling fish released at the hatchery. Coded-wire tag release information was obtained from the StreamNet Distributed System. Total survival is estimated at 0.0041%. The ODFW reported freshwater sport harvest took an estimated 13 fish.

Overall survival for **brood year 1988** is now 0.0605%. This brood year contributed mainly to the treaty ceremonial fishery in the Columbia River.

Yearlings from **brood year 1989** survived only half as well as brood year 1988. A total of 70 fish were harvested in the Columbia basin, and 3 were recovered at WDFW hatcheries. Fifty-two fish were taken in the treaty ceremonial fishery. Overall survival is 0.0353%.

Coded-wire tag information for **brood year 1990** was obtained from the StreamNet Distributed System. Total estimated survival for this brood year was 0.0027%, or 26 fish from a release of almost 960,000. The treaty ceremonial fishery reported by ODFW took two fish from this brood year.

Brood year 1991 has an estimated overall survival rate of 0.0021%, the lowest of tagged brood years in this report. There was only one off station recovery from this brood year.

All **brood year 1992** spring Chinook released at Dworshak were represented by coded-wire tags, so the rd2 program was used to estimate the total survival of 0.0584%. An estimated 78 fish, or 10% of the total, were recovered at a variety of Columbia River Basin sites other than Dworshak.

Numerous recoveries of **brood year 1993** fish by FWS were added to the RMIS database in the year 2007. Location recovered is Dworshak NFH and Kooskia NFH. Total survival is now estimated to be 0.2675%, or 3,508 fish. The rd program was used since fish unrepresented by coded-wire tags were released. There were an estimated 721 recoveries off station, including 4 in Alaska and 4 in British Columbia ocean fisheries with an estimated 2,779 fish returning to the hatchery. This brood year also provided an estimated 177 fish for treaty fisheries.

Numerous recoveries of **brood year 1994** fish by FWS were added to the RMIS database in the year 2007. Location recovered is Dworshak NFH and Kooskia NFH. Although this more than doubles total estimated survival, it still stands at 0.0807%, less than half of the average to date as of 2008. Because of a very low adult return, there was also a low egg take, and a subsequent release of just 100,775 fish for brood year 1994. A mere 83 fish were recovered. The largest number of off station recoveries was in the IDFG reported freshwater sport category, but was only 14 fish. There were also 3 expanded recoveries in the Alaska ocean troll fishery. There was also 1 FWS reported recovery at Warm Springs NFH.

The addition of numerous recoveries of **brood year 1995** fish by FWS were added to the RMIS database in the year 2007, and the removal of an earlier, now found to be erroneous group of fresh water Fish Trap recoveries results in an estimated survival of 0.1696% for brood year 1995. Because the 1995 return to Dworshak was a total of 125 fish, only 53,078 yearlings were released. Only an estimated 90 fish were recovered from this brood year, eighty-six of them at hatcheries.

The estimation of survival for **brood year 1996** is 0.0979%. Columbia River sport (429), treaty ceremonial (228) and gill net (147) fisheries took more fish than are currently in the RMIS database for recoveries at FWS hatcheries.

Brood year 1997 has an estimated survival of 0.5363% even with only 5 estimated recoveries of coded-wire tagged fish at National Fish Hatcheries, a record. The contribution to freshwater fisheries was over fifty-five hundred fish. Fish from this brood year (29) were even caught by ocean trollers as reported by ODFW.

Brood year 1998 has the highest survival rate to date at 0.8582%, and a total of 8,711 fish.

Columbia River sport and gill net fisheries took over forty-seven hundred fish in 2004 and 2005. Fish (24) were also caught by Alaskan ocean troll fishers.

With the addition of FWS hatchery recoveries from 2004, **brood year 1999** survival is higher than average at 0.1093%. Gill net and sport recoveries in the Columbia River Basin account for all off station recoveries. Twenty-two of those expanded recoveries were reported by Idaho.

Survival increases substantially for **brood year 2000** at 0.5203% with the addition of FWS hatchery recoveries from Dworshak NFH. This survival is now the third highest on record for Dworshak spring Chinook. Columbia River contribution from this brood year nearly equaled the return to the hatchery (2,313 / 2,846). One observed ocean troll recovery by CDFO is expanded to a total of 47 fish.

As of this report the average survival is 0.2194%, and **brood year 2001** is about half of that average 0.1192%. Total survival was 1,232 fish from a release of 1,033,982 with 648 returning to FWS hatcheries. There were 448 sport fishery recoveries and 96 in gill nets.

Now in sixth place when ranked by survival is **brood year 2002** with survival of 0.2398% and a Columbia River contribution of 904. These fish were harvested more equally between gill nets and sport: 328 vs 432. IDFG reported most of the sport recoveries with 248 fish.

The survival for **brood year 2003** is 0.1128% about half that of the average to date. A harvest of eighty fish was reported by Idaho, while another 580 were reported by ODFW and WDFW. WDFW reporting includes 96 in spawning ground surveys.

With very few expanded recoveries for **brood year 2004** the estimated survival is 0.2668%, high enough to be above the current average. Two hundred fifty-three fish were caught in Idaho, but the vast majority of recoveries were reported by ODFW. Over two thousand fish were harvested from this brood year.

The number of **brood year 2005** spring chinook released from Dworshak was changed significantly after release information was downloaded in January of 2012. Two copies of the Stock Assessment report were produced to document the change. Using the latest numbers, survival is 0.0321%, or 309 fish from a release of nearly a million, the lowest survival rate since the early 1990s brood years.

Brood year 2006 survived at a somewhat better rate of 0.0617% and resulted in 367 harvested fish within the basin. Gillnet fisheries harvested 219 fish. An estimated 11 fish were found on spawning grounds in the Methow, Chewuch, and Entiat Rivers.

Summer Steelhead

Estimates of survival for **brood year 1980**, well before the "Missing Group" tagging began, were added to this report in the report titled with the year 2000. This Stock Assessment Summary printout is not available. Total survivals for brood years 1980, 1981, and 1982 were 2.9305%, 1.2033%, and 0.3813% respectively. Although the Stock Assessment printout shows 0 Escapement to NFH, this results from the fact that Idaho Department of Fish and Game (IDFG) reported hatchery recoveries at Dworshak NFH in the early 1980s.

Tagged fish from **brood year 1983** (release year 1984 with CWTs 051335, 102516, and 102517) are listed as "Off-site direct hatchery release" in the StreamNet database.

No coded-wire tagged fish were released on site in brood year 1983 or 1984.

Survival for brood year 1985 has stabilized at 1.1867%. Over 5,400 fish were taken in the Columbia River gill net fishery, and over 4,100 were taken in the freshwater sport fishery reported by Idaho.

Brood year 1986 has an overall recovery rate of 1.9046%. This brood year contributed mainly to the freshwater sport fishery in Idaho. Many of these steelhead were also recovered in the Columbia River gill net fishery.

Total survival rate for **brood year 1987** is 1.0603%. Six fish were taken in British Columbia ocean trolling, but the great majority of fish (8,324) were recovered in the Columbia River basin. The majority of "Columbia River" harvest was in the ODFW reported gill net fishery (4,124), and the IDFG reported freshwater sport fishery (3,816).

Brood year 1988 summer steelhead yielded 492 observed recoveries with an overall estimated survival rate of 0.5579%. The vast majority of fish recovered off station were taken in the Columbia River gill net fishery (2,696), followed by the Columbia River sport fishery (576). IDFG

reported freshwater sport recoveries expanded to 184 fish. There was 1 observed recovery of an age-2 fish in the squid gill net by-catch and foreign research vessel records reported by National Marine Fisheries Service (NMFS)

Overall survival for **brood year 1989** is 0.8264%. Most fish were taken in the Columbia River gill net fishery (3,714) out of a total of 4,907 fish taken in the various Columbia freshwater and estuary fisheries. Sixty-three fish were taken off the coast of British Columbia. IDFG recoveries are expanded to 244.

Brood year 1990 release data were obtained from the StreamNet database. Except for one different date, 90 additional fish, and a significant difference in the total weight of fish released, StreamNet records are similar to those originally reported for releases at the hatchery. The total survival rate is 0.4821% or 5,750 fish. Most of the harvested fish were taken in the Columbia River gill net fishery (2,575). Twenty-five fish were also taken in the Oregon ocean sport fishery.

Total survival for brood 1991 is 0.2388%, near the bottom of survival rates for Dworshak steelhead. A total of 2,923 fish survived from this brood year. Seventy-five fish were harvested in the Quinault reported coastal gill net fishery. ODFW reports Columbia River gill net and sport harvest of 984 and 149 fish, and another 328 fish as freshwater sport.

Survival for brood year 1992 is now estimated to be 0.2259%, using the rd2 program. Twenty-nine percent of the fish were recovered before returning to the hatchery. Off station recoveries occurred in the Columbia River gill net and freshwater sport fisheries of all three Northwestern states, with Washington reporting the largest share.

The current estimated survival for brood year 1993 is 0.218%, the third lowest to date. All reported off station recoveries were within the Columbia River basin, and 643 fish were harvested there, mostly in sport fisheries.

Observed recoveries for **brood year 1994** are expanded to a total of 1,065 estimated recoveries or 0.0878% by the rd2 program. More fish were recovered off station (769) than returned to the hatchery, according to coded-wire tag recoveries.

Survival for **brood year 1995** is about half that of the previous brood year, or 0.0441%.

Numerous recoveries of **brood year 1996** fish by FWS were added to the RMIS database in the year 2007. Location recovered is Dworshak NFH and Kooskia NFH. This brood year now has a total estimated survival of 0.1732% with 1,320 fish returning to the hatchery. While IDFG reported the equivalent of 4 fresh water sport recoveries, WDFW had 752, gill nets had 174, and another 107 were caught in other Columbia River fisheries.

When this report was first prepared, there were no age 5 recoveries in RMIS, and, more importantly, there were no reported recoveries at any hatchery of any age, including Dworshak. Age 3, 4, and 5 recoveries are now in database, and allow a more accurate estimation of total survival for brood year 1997, a total of 4,325 fish. WDFW reported fresh water sport fisheries recoveries are equivalent to 772 fish. Next in magnitude is the Columbia River gill net fishery with 438, all at age 3. Recoveries in other within basin sport fisheries equaled 268.

Due to continuing irreconcilable differences between original and RMIS reporting, original CRiS records for all subsequent brood years will be stored in the CRiS data file srDwor, and replaced with records from RMIS.

FWS recoveries for **brood year 1998** at Dworshak are now in RMIS giving an above average survival of 0.8582%. Columbia River gill net (2,282) and sport fisheries (2,440) harvested the great majority of fish from this brood year. Alaska ocean trollers also took 24 fish.

Survival for **brood year 1999** is 0.1009%, equal to 336 fish out of 125,452 released. Approximately one third of recoveries were in sport fisheries in the basin and the gill net fishery.

With 366 observed recoveries the estimated survival of **brood year 2000** is 0.2762%. With slightly more than half of the fish recovered at the hatchery, the Columbia River contribution included 773 in the gill net fishery, with the remainder taken in sport fishers in Idaho, Oregon and Washington.

The survival of **brood year 2001** can be described as poor at 0.1897%, with only 2 other brood years being lower out of a total 23. Off station recoveries were mostly IDFG sport and Columbia River gill net, with a few score reported by ODFW.

Brood year 2002 currently has a near average 0.3384% survival with about half of the fish recovered off station and a survival of nearly forty-one hundred fish. Close to twenty-two

hundred fish were caught in the Columbia basin, over half of total survival. One WDFW observed freshwater sport recovery disappeared from RMIS during 2010.

Survival for **brood year 2003** is 0.2309% with only 178 fish recovered at National Fish Hatcheries. Either no fish returned in 2007, or, more likely, they haven't been reported. Other releases of fish from this brood year, which may have influenced the total return to the hatchery, are not returned in queries of RMIS. We note, however, that 8 WDFW observed freshwater recoveries went missing in 2010.

Estimated survival for **brood year 2004** Dworshak summer steelhead is 0.2315%, a little less than the mean survival for this run; currently 0.5084%. Gillnet and sport fisheries within the basin split the harvest of these fish for a total of 1,451 while 1,147 escaped to the hatchery.

A doubling of the survival is now on record for **brood year 2005** at 0.5296%, yielding 6,390 fish. Most fish were taken in gillnets (2,192) while Idaho sport fishers caught 1,086. This survival is very close to average of those through 2006, 0.5266%.

Eagle Creek National Fish Hatchery

Eagle Creek National Fish Hatchery is located about seven miles from Estacada, Oregon, on Eagle Creek, approximately ten miles above its confluence with the Clackamas River at an elevation of 950 feet. The hatchery was authorized by the Mitchell Act of 1938 (52 Stat. 345), as amended in 1946.ⁱ The hatchery was constructed in 1956, and is currently operated as part of the Columbia River Fishery Development Program administered by NOAA - Fisheries. Two ladders enable fish to ascend Eagle Creek. The lower or Dywer Falls, 6 feet high and 5 miles above the mouth, was laddered by USFWS in 1958. An inadequate ladder on the 10 foot high middle falls, 9 miles above the mouth, was replaced with a new ladder in 1956.ⁱⁱ

Although Eagle Creek NFH has raised fall and spring Chinook in past years, production is now limited to coho and winter steelhead.

Because fisheries in Eagle Creek, the Clackamas River, and the Willamette River are sampled poorly, or not at all, for coded-wire tags, and 19 of 27 brood years were untagged, a long term survival trend is not available.

Coho

Coded-wire tags were used in a multiple year density study including brood years 1979, 1980 and 1981. **Brood year 1980** survival rate was 1.3546%, and brood year 1981 survival was 1.0413%. Oregon ocean fisheries took nearly four times more fish than Washington ocean fisheries in brood year 1980. Brood year 1981 had an almost even split between the two fisheries.

No coded-wire tags were released in **brood years 1982 through 1987**.

The estimated total recovery from **brood year 1988** was 42,345 fish, or 4.181% when using a PEF. Nearly three hundred thousand of the coho released were from the Sandy Hatchery, and these fish did not have a unique coded-wire tag. This is highest survival rate on record for Eagle Creek coho. This brood year contributed 11,340 fish to the Oregon ocean fisheries, and 8,930 were harvested in the Columbia River fisheries.

The **1989 brood year** coho survival rate was 0.9446%, about one quarter of the very high 4.181% survival rate of brood year 1988 coho. Over 3,800 of the coho from this brood year were caught off the Oregon coast in the sport and commercial fisheries. Eighteen hundred were harvested in the ocean off Washington, and 200 off California. Over 200 fish were taken in British Columbia. Eight hundred twenty were also harvested in the Columbia River.

Brood year 1990 coho has only 31 observed recoveries and an overall survival rate of 0.1228% making this the lowest survival rate since coded-wire tagging has been done consistently. Fish from Big Creek and Sandy were released along with Eagle Creek fish this year. Fewer than 600 fish were taken in the ocean, and 310 in the Columbia River. Five hundred of the ocean recoveries were reported by WDFW. The remaining 96 were split evenly between Oregon and California. Coded-wire tagged returns to the hatchery were estimated to be less than the ocean harvest of this brood year.

Total survival rate for **brood year 1991** coho is estimated to be 0.3402%. Fish from the ODFW Sandy hatchery were released this year, so even though all fish were assigned to a coded-wire tag, they are not truly representative. Only 137 fish were harvested in the ocean (91 by British Columbia and 46 by Washington), and 205 in the Columbia River. Escapement to the hatchery was about 3,300 fish. Although this brood year did better than the 1990 fish, it was much less successful than 1980, 1981, 1988 or 1989.

Brood year 1992 survival rate is estimated at 0.4758%, an improvement over the previous two broods, but still below average for broods in the 1980s. Escapement to the hatchery made up the largest segment of fish recovered, as was also the case for brood year 1991. Washington ocean fisheries took 750 fish and Oregon ocean fisheries harvested 277 fish. An additional 197 fish were taken in the Columbia River.

Brood year 1993 releases included both forced and volitional releases of Eagle Creek stock and Toutle stock coho. Expansion of individual coded-wire tags rather than using a PEF yields a total survival of 0.2375%, placing this brood year in the lower third of survival rates. There were an estimated 121 recoveries in the Columbia River, and 358 in the ocean off of British Columbia and Washington.

Total survival for **brood year 1994** was even lower at 0.1758%. WDFW reported ocean recoveries result in an ocean harvest of 263 fish, with 66 fish harvested in the Columbia River estuary sport fishery.

Brood year 1995 was the first brood year at Eagle Creek in which coho production was "mass

marked" with an adipose fin clip. A small number of fish were not adipose fin clipped, but received a coded-wire tag. Since externally marked fish may be harvested at a different rate than the unmarked fish, recoveries have been expanded for each coded-wire tag, rather than with a PEF. WDFW Columbia River estuary sport now has an additional 4 expanded recoveries. Survival is now estimated to be 2.1002%, in the upper third of brood years.

A doubling of ODFW reported Columbia River gill net fishery recoveries increases total survival for Eagle Creek brood year 1996 coho to 1.7191%, putting this brood year in the upper half when ranked by survival. Two thousand five hundred forty-nine fish were harvested in the ocean and 1,586 in the Columbia River.

A recalculated expansion for recoveries at the hatchery, plus other changes in reporting results in an estimated survival of 3.9776% for Eagle Creek brood year 1997 coho. Only brood years 1998 and 1988 are higher. Over fifty-two hundred fish were harvested in the ocean, almost equally divided between Oregon and Washington with an additional 95 recoveries estimated for California ocean sport fisheries.

An adjustment to the expansion rate of NFH recovered tags increases the estimate of survival for brood year 1998 to a record 4.9126%. This new hatchery return rate is closer to the rate determined with the Age Composition Analysis of the two return years. Over eighty-five hundred fish were taken in WDFW ocean fisheries, and over fourteen hundred in ODFW reported ocean fisheries. This gives a total of 9,965 fish harvested in the ocean. Additionally, ODFW had reported that an additional eighty-five hundred fish were taken in the Columbia River.

The survival rate for brood year 1999 is a very respectable 1.9272%. Ocean recoveries (883) were almost evenly split between Oregon and Washington, but most of the harvest occurred in the Columbia River (1,380). Expansion of CWT recoveries shows a return to the hatchery of 11,457 fish.

There were 511 observed recoveries of coded-wire tags from **brood year 2000**, which when expanded give a total survival rate of 1.5508%, a small reduction since last year. This is about average for Eagle Creek NFH coho. Most off hatchery recoveries were in the Columbia River gill net and estuary (472). Ocean fishery reports account for 524 expanded recoveries: 301 in Oregon, and 223 in Washington. Over seven thousand fish returned to the hatchery.

Eagle Creek had been releasing around one million coho per year since at least the early 1980s. Because of limited harvest opportunities, however, the production goal was reduced to 500,000. The 2000 brood year was the first year this new goal was implemented.

Brood year 2001 survival is estimated to be 1.2103, which is about average. Ocean harvest was split 374 to 256 for Washington and Oregon respectively.

An estimated survival of 1.5391% means that **brood year 2002** was slightly below average. This brood year contributed nearly 600 fish to ocean fisheries, and about five hundred to Columbia River harvest. Meanwhile the expanded recoveries at Eagle Creek were 7,480, while actual counts at the hatchery give a return of ninety-five hundred.

Survival for **brood year 2003** is similar to that of 2000 at 1.1521%, and slightly below average as of 2010. Nine hundred sixty-two fish were caught in the ocean, while 603 were taken in the estuary and fresh water. Meanwhile, using the CWT information, 4,477 fish returned to the hatchery.

Total survival for **brood year 2004** is estimated at 3.1056%, well above the average to date of 1.7843%, with a contribution of 1,734 fish to the ODFW Ocean Sport fishery. A mix of WDFW reported fisheries, ocean and in river, harvested another 1,370 fish.

Three hundred forty-one fish were harvested in the Columbia River Basin from **brood year 2005**: gill net and sport taking nearly equal shares, and another sixty-three fish caught in estuary sport fisheries. Four hundred forty-four fish were taken in WDFW and ODFW ocean fisheries, again nearly splitting the take. The low estimated survival for this brood year, 0.7916% is a result of CWT recoveries at the hatchery under estimating return of Age 3 fish by about half.

Survival for **brood year 2006** is now estimated at 3.5696%. ODFW reported Ocean Sport fisheries and WDFW private sport fisheries took almost equal numbers of fish: 2,327 and 2,025. Gillnetters harvested 1,071, and various ocean and Columbia River Estuary fisheries harvested about one thousand fish combined for a combined off station harvest of approximately seven thousand fish.

Initial survival estimate for **brood year 2007** is 0.4099%.

Entiat National Fish Hatchery

Entiat NFH is located on the Entiat River, west of Entiat, Washington. Elevation is 980 feet above sea level. Construction began in 1940 under the Grand Coulee Fish Maintenance Project and operations began a year later. This report addresses the long term spring Chinook salmon program through brood year 2006.

Yearling spring Chinook at Entiat were coded-wire tagged beginning with **brood year 1988**. Expanded off station recoveries include 57 in the treaty ceremonial fishery and 6 in the Columbia River sport fishery. Expanding CWT recoveries for brood year 1988 gave an estimated survival of 0.1023%, about average for this hatchery and species. Eggs from Winthrop NFH were raised and released at Entiat as part of this brood year.

There were 56 observed recoveries of the 111,207 tagged fish released with **brood year 1989**. Expansion of these recoveries yields an overall survival rate of 0.0519%. Seven fish were taken in the Columbia River sport fishery. Production for this brood year included fish raised from eggs transferred from Leavenworth NFH.

There were only four observed recoveries from the 95,682 tagged fish released with **brood year 1990**. Using the rd2 program yields an estimated total survival of 0.0047% for this brood year, the worst survival on record for Entiat spring Chinook. All recoveries were at the hatchery.

Brood year 1991 was a bit more successful than the previous brood year with a total survival rate of 0.0279%, using the rd2 program. Nine fish were taken in the Columbia River treaty ceremonial fishery. All observed (28) recoveries were of age-4 fish.

A total recovery rate of 0.0346% puts **brood year 1992** third from last when ranked by survival rate for yearling releases at Entiat. The freshwater fish trap category accounts for most of the off station recoveries, although there were also a few treaty ceremonial and WDFW hatchery recoveries.

Total survival for **brood year 1993** was 0.0383%, better than the previous two brood years, but still ranking as fourth worst. Five coded-wire tagged fish were recovered in fish traps. While most returned to the Entiat, an estimated 19 total fish were recovered at WDFW hatcheries.

The yearling release for **brood year 1994** included fish received as eggs from Leavenworth NFH. Total survival was estimated to be 0.0667% with an estimated 10 non-hatchery recoveries. This survival rate is in the middle of Entiat yearling spring Chinook survivals.

The estimated total survival for **brood year 1995** is 0.3282%. There were 3 ODFW reported ocean troll recoveries and 8 recoveries in the Columbia Basin. WDFW hatchery recoveries were at Wells. Brood year 1995 is in the top fourth of survivals

Brood year 1996 has provided a high estimated survival for Entiat NFH spring Chinook, now at 0.5855%. This was in spite of the use of a blank coded-wire tag for 16,053 of 124,536 yearling fish released. This blank wire could yield no off station recoveries. There were three expanded ocean recoveries, this time reported by British Columbia. Treaty ceremonial and sport recoveries were the next two highest off station recovery categories with an estimated 54 and 36 recoveries respectively. As usual the lion's share of recoveries (91%) was at Entiat NFH.

Entiat **brood year 1997** currently has yielded an estimated total survival of 0.7732%, the highest to date for Entiat. Off station recoveries were mostly in the ODFW reported Columbia River sport (211) and gill net (211) fisheries. There were also an estimated 11 treaty ceremonial recoveries.

The third highest number of recoveries now comes from brood year 1998: an estimated 2,067 recoveries and a survival rate of 0.5747%. Most non station recoveries were in the ODFW reported Columbia River gill net fishery (357), followed by the Columbia River sport fishery (94).

Brood year 1998's survival of 0.5747% puts it in third place for overall survival of Entiat brood years. Four hundred ninety-three fish were harvested in the Columbia River with 357 in the gill net fishery and 94 in sport fisheries.

Survival falls to less than one fifth of the previous year for brood year 1999 at 0.089%. An estimated 33 fish were recovered off station: 11 in spawning ground surveys, 13 in the Columbia River gill nets, and 19 in sport fisheries.

Although survival for **brood year 2000** rises to 0.1591, it is still below the average of 0.17%. Gill net and sport fisheries were responsible for recovery of 95 fish: 55 and 40 respectively. An estimated 12 fish were found in spawning ground surveys.

Fish from **brood year 2001** had a survival of 0.2206%, which is above average. One hundred nine

fish were taken in Columbia Basin sport fisheries, and another 54 in gill nets; while CWT recoveries at Entiat expand to an estimated 699 fish.

Estimated survival for **brood year 2002** is half that of the previous brood year, 0.1215%. Total number recovered from this brood year is estimated at 427 fish.

Based on the available coded-wire tag data, survival for brood year 2003 is one of the lowest on record at 0.0254%. However, no coded-wire tags were recovered at Entiat NFH in 2007 even though a total of 627 fish returned. Eight of the returning fish were listed as jacks by the hatchery with 69 listed as "unknown" sex. Any comparison with other returns should be undertaken with the incomplete sampling in mind.

Survival for **brood year 2004** is 0.1522%. Although this is lower than the average to date of 0.2306%, it must be noted that there was no sampling of CWTs at the hatchery in 2007, and half the heads recovered in 2008 were lost. Off station recoveries included 170 in Columbia River gillnets and 144 in Columbia River sport fisheries.

Brood year 2005 has an estimated survival of 0.2478% with 338 fish recovered within the basin rather than at the hatchery.

No brood year 2006 spring Chinook were released at the station.

Brood year 2009 summer Chinook will be reported on when returns from that new program are complete.

Kooskia National Fish Hatchery

Kooskia NFH is situated along Clear Creek, just upstream of the confluence with the Middle Fork Clearwater River, approximately 75 miles southeast of Lewiston, Idaho, at an altitude of 1,295 feet. The hatchery was authorized in 1961, first operated in 1969, and is currently used for adult collection and rearing of spring Chinook salmon.

Only two broods during the 1980s received tag codes at Kooskia. An age-0 release of brood year 1980 fish contained an IDFG tag code. Fish raised at Kooskia were released at Dworshak NFH with an IDFG tag code. Fish from Hagerman NFH were released at Kooskia in 1983.

Brood year 1988 fish at Kooskia were the first fish to be marked under the Missing Production Group program. Total survival for brood year 1988 was 1,083 fish, or 0.2683%. Treaty ceremonial fishing took most of the fish harvested off station (156). Nineteen fish were also taken in the ODFW test fishery net.

No tagged fish were released with brood year 1989.

An estimated 77 fish were recovered from brood year 1990, resulting in a survival rate of 0.0106%. Eleven of those fish were taken in the Columbia River sport fishery in 1994.

Total fish released for Kooskia NFH brood year 1991 was 343,437, including coded-wire tag 052925 with 60,585 fish. Current RMIS records show 50,585 fish released with this tag code, making a total release of 333,437. Total recoveries for this brood year consist of a jack and an adult recovered at a national fish hatchery for a survival of 0.0006%.

Fish traps and non-USFWS hatcheries reported the majority of off station recoveries for brood year 1992. An estimated 4 fish were taken in the treaty ceremonial fishery. With 86 recoveries reported by USFWS, total estimated survival rate is 0.0317% using the rd2 program, better than the previous two brood years.

Survival for brood year 1993 is estimated to be 0.0327%, right in the middle of survivals. Nine fish were taken in the California ocean sport fishery, an unusual occurrence. Thirty-two fish were harvested in the treaty ceremonial fishery. Fish without coded-wire tags, but with right ventral fin clips, were also released eight miles above the hatchery. There were an unusual number of recoveries of spring Chinook in California ocean sport and troll fisheries in 1996. Spring Chinook from Carson, Little White, and Warm Springs were also taken, as were tule fall Chinook from Spring Creek.

There were 8 observed recoveries from the release of 271,681 tagged fish in brood year 1994. The estimated number of recoveries in the ODFW fish trap is 521 from one observed recovery, a very high ratio. No recoveries have been reported from FWS hatcheries. A group of fish was released without a representative coded-wire tag. The PEF for this brood year is 1.23, and leads to a total survival of 0.0027%.

Fewer than 16,500 tagged fish were released in **brood year 1995** in a total of 16,598 fish. There is now 1 reported recovery by an IDFG hatchery for a survival of 0.0060%.

With 15 observed recoveries, **brood year 1996** total survival is estimated to be 0.0649%, or 28 fish. Ten fish were harvested in the gill net fishery, 8 for treaty ceremonial purposes, and 7 in Columbia River sport fisheries.

Brood year 1997 coded-wire tag release information was obtained from the RMIS database. It is substantially different than numbers reported through CRiS, as noted on the Stock Assessment Reference Summary. Previous numbers will be replaced with the numbers in RMIS next year. The survival rates for coded-wire tags in this release are expanded to a record 0.8767% survival for Kooskia NFH, without any FWS reported recoveries. Contributions were made in the following fisheries: Columbia River gill net and sport (1,000 and 865), freshwater sport (109), treaty ceremonial (66). There were 4 PSC expanded recoveries in the Oregon reported non-treaty ocean troll in 2001.

Release numbers in RMIS differ from those reported to CRiS for **brood year 1998**. Numbers in CRiS have been edited to conform to numbers reported to RMIS. Brood year 1998 has the third highest survival rate for Kooskia NFH spring Chinook at 0.76632% now that FWS recoveries are in RMIS. These FWS recoveries expand to a hatchery return of 852 fish, with a Columbia River contribution of 1,946. Nine hundred of those recoveries were by Columbia River gill nets, and sport fisheries took 917 fish. Spawning ground survey information shows an estimated 67 fish at Kooskia NFH from this brood year.

RMIS records for **brood year 1999** total 82,974 fish vs 80,430 previously recorded in the CRiS database. With the addition of 14 observed recoveries by FWS, estimated survival for this brood

year is 0.1410%. Columbia River gill nets took the largest number of fish at 22, and with the estimated take in the sport fishery at fourteen fish. There were a total of 5 expanded recoveries in spawning ground surveys.

Brood year 2000 RMIS release records for CWT 054516 appear to include fish released off station, but in Clear Creek (CRIS sr80s Dist-KK). Age4 and Age5 recoveries by FWS have come on line since the last report, and the current survival for this brood year is 0.3068%. Columbia River sport fisheries reported the highest number of recoveries for this brood year, estimated at a total of 432 fish, followed by gill net (359), then other freshwater sport with 135.

Survival for **brood year 2001** is about a fifth that for the previous return at 0.0511%, a total of 280 fish. Using the CWT recoveries, an estimated 185 fish returned to the hatchery. Although Columbia River contribution is only 85 fish, 10 were also taken in ocean troll fisheries reported by Oregon.

In the "middle of the pack" **brood year 2002** achieved a survival of 0.1289%. Over forty percent of surviving fish were retrieved off station. Most of those recoveries were in various sport fisheries in river with 48 also taken in gill nets. Twelve fish were caught in the BC Ocean non-treaty Troll fishery.

Brood year 2003 releases from Kooskia showed that 49,973 yearlings were released at "PUBLIC ROAD BRIDGE". That release appears to have been combined with the release at the hatchery before being submitted to RMIS, but with slightly different numbers. This original data has been archived and replaced with data downloaded from RMIS. The estimated survival of 0.0574% is about one quarter of the average to date. Harvest consisted of 210 fish taken within the Columbia River Basin.

Survival increased tenfold for **brood year 2004** at 0.5578%, and is nearly twice the average. Three thousand five hundred forty-two fish were recovered from this release. Most were caught in ODFW reported Columbia River Sport fishery (1,316) followed by 700 in gillnets. Return to the hatchery nearly matched ODFW sport at 1,211.

The next **brood year** survival, **2005**, returns to a level more like that of 2001 through 2003, or 0.1159%. Although all fish are coded as the same stock, and all have an adipose clip, a group of approximately forty-five hundred fish were released and not associated with a coded-wire tag. Given the low survival (a total of only 202 fish were recovered) from a release of around one hundred eighty thousand, that group would not change recovery information in any significant way.

Survival rebounds to 0.3908% for **brood year 2006**. With a release of 112,344 fish this results in a total of 439 fish, 127 of which escaped to the hatchery. Sport and gillnet fisheries shared the majority of the catch, with 17 fish going to Treaty Subsistence fishing.

Leavenworth National Fish Hatchery

Leavenworth NFH is located about four miles south of Leavenworth, Washington, along Icicle Creek, a tributary to the Wenatchee River. Elevation is 1,155 feet. The hatchery was originally authorized in 1937 by the Grand Coulee Fish Maintenance Project and was re-authorized in 1938 by the Mitchell Act. The facility began operations in 1942 and is currently used for producing spring Chinook salmon.

Fish from **brood year 1985** were released at Leavenworth with two different tag codes. Fish with those same tag codes were also inadvertently released in the Yakima River, and are therefore not included in this report.

Fish from **brood year 1986** were harvested mainly in the freshwater sport (757) and treaty ceremonial fisheries (402). One recovery from the ODFW treaty ceremonial fishery is no longer in the PSMFC database. Total estimated survival is estimated at 0.1174%.

Brood year 1987 spring Chinook salmon released as yearlings have an estimated total survival of 0.2785%. Off station recoveries were mainly in the freshwater sport fishery and treaty ceremonial fishery. Nearly 3,000 fish were taken in the Columbia River. One tag, which was expanded to 28 fish, was recovered in the British Columbia ocean troll fishery. The 1987 brood year spring Chinook released as sub-yearlings fingerlings survived at a rate of only 0.0030%. This release of 939,426 fish yielded a freshwater sport catch of 12 fish, and a hatchery escapement of 16 fish.

Brood year 1988 spring Chinook fared quite a bit better than brood year 1987 with an estimated survival rate of 0.4751%. Most fish were taken in the freshwater sport fishery (3,469). In addition, ODFW reported a treaty ceremonial harvest of 1,315. Twenty-five fish were harvested in the WDFW reported treaty troll fishery. Fingerling releases from this brood year survived at an overall rate of only 0.0034%. There were only seven observed recoveries from a release of nearly 300,000 tagged fish.

An agency 63 tag code was released with **brood year 1989** age-0 Leavenworth fish raised at Wells State Fish Hatchery (SFH) and returned to Leavenworth for further rearing. These fish survived at a rate of 0.0889%. Using the calculated PEF of 4.41, 13 fish were harvested in the British Columbia ocean troll fishery, 163 in the freshwater sport fishery, and 66 in the treaty ceremonial fishery. However, this PEF includes 400,000 fish released at one-third the size of the fish with the coded-wire tags.

Yearling fish from **brood year 1989** survived at a rate of 0.1946%, more than twice the rate for fish released at age-0. The great majority of off station recoveries occurred in the Columbia River sport fishery (1,585). An additional 360 fish were taken in the treaty ceremonial fishery.

Using the rd2 program, rather than the PEF method, the estimated number of fish surviving from **brood year 1990** yearlings was only 78 fish from a total release of over 2 million fish, a very low survival rate of 0.0034%. Treaty ceremonial harvest of this brood year is estimated to have been only 23 fish. Age-0 fish released from this brood year had no representative coded-wire tag.

Survival of **brood year 1991** is now estimated at 0.0213%, a total of 374 fish, using the rd2 program. This survival is over six times greater than brood year 1990. Fourteen age-4 fish were taken in the ODFW reported treaty ceremonial fishery. Age-0 fish released from this brood year had no representative coded-wire tag. Observed recoveries include age-2 fish recovered at the hatchery, 89. There are fewer PSC expanded recoveries, 82, which does not include these fish.

Survival was up again with a 0.0754% rate for **brood year 1992** using the rd2 program. Other than 987 hatchery recoveries, there were 124 fresh water sport recoveries, in addition to treaty ceremonial (32) and Columbia River gill net harvest (5).

7

Brood year 1993 survival is now at 0.2748% using the rd2 program, still increasing from the low point of brood year 1990. Only the last three brood years in the 80s had higher survivals. There was a freshwater sport harvest of 1,067, followed by treaty ceremonial (61), and freshwater net (60).

The estimated survival for **brood year 1994** is 0.0580%. An estimated 34 fish were recovered in treaty ceremonial fisheries, and 50 in freshwater sport fisheries.

Survival for **brood year 1995** is estimated to be 0.1232% with over three hundred fish being harvested off station. Fish were taken in gill net, fresh water sport, and treaty ceremonial fisheries.

Estimated survival for **brood year 1996** is 0.3747% in the upper third of survivals. There were 2 recoveries in Alaska, expanded to a total estimated 26. There were also a total of 388 estimated

recoveries in the WDFW reported fresh water sport recoveries, and over two hundred each in gill net, spawning ground, and treaty ceremonial fisheries.

Brood year 1997 shows an estimated total survival of 0.7675%, a record setting brood year. The greatest number of fish recovered off station was in the Columbia River sport fishery (2,140). The gill net fishery took an estimated 2,100 fish, followed by recoveries in spawning ground surveys (750), and treaty ceremonial (127).

Slightly lower than the previous brood year at 0.7190%, but still the second highest on file, brood year 1998 contributed about thirty-seven hundred fish to Columbia River fisheries. Thirty-nine fish were harvested in the ocean. A downward adjustment of a CDFO expansion factor reduced this survival in 2006.

Estimated survival for brood year 1999 is 0.0961%, an increase over last year due to the addition of 3 observed recoveries by WDFW. This is in the lower third of recorded survivals. Gill net and sport recoveries totaled 666 fish, and there were an estimated 87 recoveries in spawning ground surveys.

Additional recoveries from Leavenworth **brood year 2000** have appeared in data down loaded from RMIS. Therefore, the total number of fish that were caught or returned to the hatchery is 5,799, equivalent to a survival of 0.3731%. Nearly twenty-five hundred fish were taken in the Columbia Basin, in both sport and gill net fisheries. WDFW spawning ground surveys reported an estimated 292 fish. Alaska accounts for 2 observed troll recoveries, equivalent to 5 total fish.

The estimate of survival for **brood year 2001** is 0.2515%. Columbia River contribution is estimated at 750 fish. Most of those fish were taken in the ODFW reported "River sport" fishery. The gill net fishery harvested an estimated 144 fish.

Brood year 2002 estimated survival is 0.1025%, less than half of the through BY 2006 average of 0.24%. Freshwater sport fisheries harvested most of the fish not returning to the hatchery; a total of 467 fish. Spawning ground surveys account for 86 adults, and 24 were taken in Treaty Ceremonial fisheries.

A survival of 0.0745% for **brood year 2003** is also well below average. While fresh water sport fisheries account for 316 fish harvested, another 164 fish were caught by gill net fishers. A few fish each were taken in Ocean Troll (CDFO) and Treaty Ceremonial.

Brood year 2004 is currently rated as the third highest surviving brood year at Leavenworth at 0.4811% with additional expanded recoveries. While 1,387 fish escaped to the hatchery, over three thousand were harvested in the Columbia River. The four hundred thirty observed off station recoveries is a record for Leavenworth. The prime harvesters of the brood year were Columbia River gill net, and sport fishers who nearly split the catch: 1,260 verses 1,540.

Additional recoveries give **brood year 2005** a survival estimate of 0.1809%, well below average. ODFW's Columbia River Sport recoveries account for 404 fish with gill netters harvesting 168, while WDFW Freshwater Sport reports a take of 120 fish.

Survival for the next **brood year, 2006**, increases by more than three times to 0.7617%, even without recoveries of five year olds.

Summer steelhead at Leavenworth have never been marked with coded-wire tags.

Little White Salmon National Fish Hatchery

Little White Salmon NFH is located on the Little White Salmon River, 12 miles east of Stevenson, Washington, at an elevation of 90 feet. The hatchery began operations in 1898 and was remodeled and expanded in 1958 as part of the Columbia River Fishery Development Program (Mitchell Act). Current production consists of rearing upriver bright fall and spring Chinook salmon. Coho are raised, but no longer released at Willard NFH, which is administered by Little White Salmon. Willard is listed separately in this document. Recent production of tule fall Chinook will be reported on when results are available.

Spring Chinook

Both age-0 and yearling spring Chinook have been released from the hatchery. **Brood years 1982 through 1984** were coded-wire tagged to evaluate age-0 and yearling releases. Average survival for age-0 fish was 0.11%, compared with an average survival of 0.39% for yearling releases. Columbia River gill nets took a greater proportion of age-0 fish than yearlings (9.7% vs. 3.2%) for these three broods. Release of coded-wire tagged age-0 fish was resumed with brood year 1991.

Marking of spring Chinook resumed for **brood year 1988** under the Missing Production Group program. Brood year 1988 fish released at age-0 were not marked. The overall survival rate for yearling fish from this brood year was 1.0457%. Nearly 2,400 yearling release fish were harvested in the Columbia River, with over 1,500 in the WDFW reported freshwater sport fishery alone. Another 730 fish were taken in the treaty ceremonial fishery.

Brood year 1989 shows a total survival of 2,270 fish, or 0.2230%, down from last year with the elimination of an age-2 recovery. Nearly 1,200 fish were taken in the freshwater sport fishery, and 85 in the treaty ceremonial fishery. Age-0 fish were released, but without a coded-wire tag.

The survival rate for **brood year 1990** is 0.0155% with no reported off station recoveries. Coded-wire tagged fish from this brood year were released from the Willard NFH facility, (a sub-station located upstream from Little White Salmon NFH) along with a total of 869,952 fish. Another 807,742 untagged, unmarked fish were released at Little White Salmon NFH. There was no age-0 release from this brood year.

Only 232 fish survived out of 809,079 yearling fish released from **brood year 1991** (0.0287%), using the rd2 program. There were only 7 recoveries, identical to brood year 1990. There were no recoveries from 94,295 tagged fish released at age-0.

Brood year 1992 yearlings now have a survival rate of 0.5456% and an estimated total survival of 5,426 fish. Over one thousand fish were caught in the WDFW freshwater sport fishery, and another 790 in the freshwater net fishery.

Fingerlings from **brood year 1992** now have an estimated survival rate of 0.0121%, or only 61 fish from a release of 503,458.

The rd2 program estimates total survival as 0.2175% for **brood year 1993**. Recoveries from 1997 are expanded to 330 freshwater net, and 709 freshwater sport recoveries. There were an additional estimated 16 fish harvested in the treaty ceremonial fishery. This brood year has a below average survival rate.

Estimated survival for **brood year 1994** is 0.0385%, the third lowest percentage on record. Only 18 fish were estimated to have been taken in the treaty ceremonial fishery with just 352 returning to the hatchery.

Survival for **brood year 1995** is estimated to be 0.4042%, which places it in the mid-range of survivals. Recoveries occurred in freshwater sport (257), treaty ceremonial (34), gill net (26), and on spawning grounds (12) in addition to 2,381 returns to the hatchery.

The estimated total survival for brood year 1996 is 0.4352%, in the top half of survivals. Freshwater sport (723) and treaty ceremonial (285) fisheries were responsible for the majority of off station recoveries with an estimated 131 fish also being taken in the Columbia River gill net fishery.

Brood year 1997 is in the middle of survivals with 0.3685%. More than twenty-three hundred fish were harvested in Columbia River sport fisheries. Seven hundred twenty fish were taken in the Columbia River gill net fishery and 96 as treaty ceremonial.

A survival rate of 0.6306 puts **brood year 1998** third best of all Little White spring Chinook releases. This brood year contributed over twenty-five hundred fish to sport fisheries, eleven hundred to the gill net fishery, and 174 to treaty ceremonial harvest.

The estimated 0.1446% survival rate for **brood year 1999** is in the middle of completed brood years. Nearly one thousand fish were harvested in the Columbia River in sport (705) and gill net (225) fisheries.

The release of one million plus **brood year 2000** spring Chinook yearlings from Little White resulted in a total contribution of 2,655 and a calculated return to the hatchery of 1,138. Percent survival is 0.2559. Columbia River contribution is 1,474, higher than return to the hatchery (1,138).

Survival is halved for **brood year 2001** at 0.1037%, only 50 observed recoveries, and an estimated total of 1,025 recovered. Only one tagged Age3 fish was recovered at the hatchery, while age composition shows 12 jacks while the hatchery jack count was 17. ODFW estimated sport catch is 345, and the gill net catch is 165.

Brood year 2002 survival is 0.6414%, the third highest to date for Little White spring Chinook. Ocean fisheries north of Oregon: Washington, British Columbia, and Alaska, hit Age4 fish from this brood year, taking one hundred five fish. About twenty-four hundred fish were caught in the Columbia basin sport fisheries, and 441 in gill nets.

Survival for **brood year 2003** is an estimated 0.3201% with a total of 2,496 recovered. 1,434 fish were harvested off station while 1,062 made it back to the hatchery. Most fish (804) were harvested in Freshwater Sport reported by WDFW with gillnets taking less than half of that (318).

While further recoveries are expected, survival for **brood year 2004** is currently estimated at 0.5201%, near the top quarter in terms of success. Off station recovery of these fish was higher than the return to the hatchery, with 1,310 taken in the gill net fishery alone.

With no listed Age 5 recoveries, **brood year 2005** exceeds the survival rate for the previous brood year at 0.6711%. And, once again, more fish were harvested than escaped to the hatchery; a harvest of 4,632 and escapement of 2,120. The Columbia River Sport fishery was responsible for the great majority of the harvest with a take of 1,760 followed by WDFW's Freshwater Sport of 1,168 fish.

Currently, survival for **brood year 2006** is estimated to be 2.3158%, more than twice the previous highest survival rate.

Upriver Bright Fall Chinook

Brood year 1983 through 1985 upriver brights were marked for both a normal age-0 release and an extended rearing release. Average percent survival for the fingerling release from these three brood years was 1.1%, compared to the survival of 0.3882% for the extended rearing fish. The extended rearing program strategy is no longer being used. The survival rates for two of the three brood years with extended rearing are ranked fourth and fifth of eighteen releases for this program.

Brood year 1989 was the first upriver bright brood year to be marked under the Missing Production Group program. Fifty-eight observed recoveries were expanded to a total of 3,530 for an overall survival rate of 0.2454%. Most of the harvested fish were taken in the ocean (1,733), with 931 in Alaska, 577 in British Columbia and 225 in California. An additional 160 fish were harvested in the Columbia River gill net fishery, and 642 were recovered in spawning ground surveys.

Current recovery information and use of the new rd2 program shows a 0.3285% recovery rate for brood year 1990, higher than that for brood year 1989. Little White Salmon upriver bright fall Chinook appear to be the only group of fish in this report which had a higher rate of survival for brood year 1990 than brood year 1989. Unlike Spring Creek tule fall Chinook, where fish were released in March, April, and May, these fish were released in June. These upriver bright fish are also recovered farther north than Spring Creek tules. Alaska ocean troll and ocean sport took most of these brood year 1990 fish (2,161), followed by Canadian troll and ocean sport fishers (1,844), and the Columbia River gill net fishery (1,041). Spawning ground surveys account for an additional 1,047 fish.

Survival for brood year 1991 is 0.2039%, approximately 60% of brood year 1990. Alaskan troll (856) and British Columbian fishers (570) harvested fish in the ocean, and the Columbia River gill net fishery took 920 fish. Although brood years 1990 and 1991 had very low survival rates basin wide, these releases are in the middle, and fifth from the lowest.

The estimate of survival for brood year 1992 is now 0.3623% with a total survival of 6,763 fish. WDFW added 6 observed spawning ground recoveries in 1998. Alaska and British Columbia ocean fisheries took 787 and 118 fish respectively. Columbia River gill nets took 1,140 fish, and the

river sport fishery accounted for 275. Survival for this brood year is average.

Survival from brood year 1993 is now 13,540 fish or 0.7531%, more than double the survival for brood year 1992. This is the second best survival for this hatchery and species. Alaska and Canada took nearly equal shares in the ocean catch (1,545 and 1,222), and over thirty-five hundred fish were harvested in various Columbia River fisheries.

Brood year 1994 is now estimated to have survived at a rate of 0.0826%, only about one ninth the rate of the previous brood year, and the next to lowest for this hatchery and species. Alaskan fishers took 229 fish from a total of 309 estimated recoveries in the ocean. An estimated 299 fish were recovered in Columbia Basin spawning ground surveys.

The estimated survival for brood year 1995 is 0.2422%, with over nine hundred recoveries in Alaskan waters, from a total 5,079 estimated recoveries. This survival is in the lower half of survivals for this hatchery and species.

Brood year 1996 is in the lower third for Little White Salmon upriver bright fall Chinook at 0.1388%. Even so, fish were recovered in Alaska (764) and California (44) ocean fisheries.

Brood year 1997 is in the upper third of the survivals at 0.4139%. Nearly eleven hundred fish were harvested in Alaskan waters. The five thousand plus fish recovered in the Columbia River are particularly noteworthy. Over three thousand of those resulted from WDFW conducted spawning ground surveys. This high number of estimated spawning ground recoveries is undoubtedly due to the hatchery restricting the number of fish entering the adult holding pond during spawning season. This practice was begun in 2000. This brood year contributed fifteen hundred fish to the Columbia River gill net fishery.

Overall survival for brood year 1998 is now 0.3617%, a bit lower than the previous brood year. The expansion factor for eighteen 2003 recoveries observed in the Columbia River Gill net fishery by ODFW has been reduced, lowering the resultant expanded recoveries from 111 to 42. Over twelve hundred fish were harvested in Columbia River gill nets. Over twenty-six hundred fish from brood year 1998 are estimated to have been recovered on spawning grounds. Spawning ground recoveries were no doubt higher than usual due to the fact that all returning fish were not allowed to enter the Little White Salmon adult holding building.

Intermittent ladder operations occurred at Little White Salmon NFH during 2002-2004.ⁱⁱⁱ

Spawning ground recoveries expand to nearly seven thousand fish for brood year 1999. This is a result of restriction adult returns to the hatchery. The estimated total survival is 16,473 fish or 0.8359%, the second highest survival to date. Close to four thousand fish were taken in Alaska non-treaty troll fisheries, and about twenty-five hundred in Columbia River gill nets.

The current estimate of survival for **brood year 2000** is 0.3194% and a total survival of 6,189 fish. Alaskan fisheries took the lion's share of this brood year with 1,480 being caught mostly by trollers. Eighteen hundred fish found on spawning grounds are again the result of the hatchery preventing a large part of the returns from entering the hatchery.

Brood year 2001 is in the bottom third of successful years, and a changed expansion factor for ADFG Ocean Sport lowers that survival to 0.2859%. Harvest was split roughly equally between ocean and fresh water. Alaskan ocean harvest was the greatest number by far (1,140) followed by 310 by BC, and 30 in WA. Intermittent ladder operations sent an estimated 1,650 fish onto spawning grounds while another nine hundred fish were harvested in river with 1,210 taken in gill nets.

Continuing a short term downward trend in survival, **brood year 2002** survival is now 0.1362%. In spite of this low survival, nearly a third of these fish were harvested in the ocean: 440 off of Alaska, 396 off of British Columbia, and 88 off of Washington. Ladder closures sent 814 onto nearby spawning grounds. Estimates made with the coded-wire tag information, however, underestimates the return by 760 fish; a fact made clear by looking at actual counts of returning upriver brights at Little White Salmon NFH. Age Composition information shows a return of 1,101 fish.

A slight rise in survival is noted for **brood year 2003** at 0.1708%. Still, 980 fish were harvested along the West Coast from Oregon to Alaska, and 530 within the Columbia River Basin. Three hundred sixty fish were recovered on spawning grounds.

A just about the same as the previous brood year's survival, or 0.1765%, has been determined for **brood year 2004**. Nearly half of recovered fish were taken before returning to the Columbia River with 822 in Alaska, 346 in BC, and 24 in Washington waters. In basin harvest was 393 fish.

Brood year 2005 survival is higher than all but 3 other brood years at 0.7263%. Ocean and Columbia River recoveries were nearly equal at approximately three thousand each. Within the

basin the vast majority (2,189) was caught in gillnets. In the ocean Alaska harvested the greatest share with a take of 1,652.

To date the average estimated survival for Little White upriver bright fall Chinook is 0.3698%.

Spring Creek National Fish Hatchery

Spring Creek NFH is located on the Columbia River at Underwood, Washington, about 30 miles upstream of Bonneville Dam. Elevation at the hatchery is 93 feet above sea level. The hatchery was constructed in 1900 and began operations a year later. Fish were trapped in the Big White Salmon River from 1901 through the 1950s. Spring Creek NFH was remodeled in 1955 under the Mitchell Act and redone in 1970 under the John Day Mitigation Act. Spring Creek NFH currently produces tule fall Chinook.

The survival rate for **brood year 1984** was 0.0462%, and was 0.1294% for brood year 1985. Even with the low survival rate of these two brood years, thousands of fish were harvested in both the ocean and the Columbia River.

Survival for **brood year 1986** is estimated at 0.4328%, 46,050 fish from a release of 10,640,406. Over 22,000 fish were harvested in the ocean and 14,288 were taken in the Columbia River gill net fishery.

Brood year 1987 fish were recovered primarily in commercial fisheries off the coasts of Canada and Washington, as well as in the Columbia River gill net fishery. Total survival for the 1987 brood year is 0.3087%, or 27,326 fish from a release of 8,850,757.

Brood year 1988 fish were recovered primarily in the Columbia River gill net fishery (24,168). They also contributed to the ocean fisheries in Washington (17,417), Canada (11,671), and Oregon (6,319). Overall survival is estimated at 0.5168%.

Fish from **brood year 1989** fared slightly worse than brood year 1988 with an overall survival rate of 0.4855% or 49,683 fish. The greatest number of recoveries was in the ocean off Washington (14,894), and in the Columbia River (12,548). The Columbia River gill net fishery took the great majority of fish harvested in the Columbia River. British Columbia took 9,984 fish, and ocean harvest off Oregon totaled 3,551 fish.

The estimated total survival for **brood year 1990** drops to 0.1295%. This is one of the lower survival rates since brood year 1980. Most of the harvest of these fish occurred in the Columbia River, 5,017 fish, followed by the Washington ocean harvest of 4,811, and the Canadian harvest of 3,345.

Brood year 1991 survival rate, expanding for each coded-wire tag, is 0.1374%, only slightly better than brood year 1990. British Columbia ocean troll and ocean sport fisheries took the largest number of fish from this brood year at 7,787 fish. The Columbia River gill net fishery harvested 7,001 fish.

Survival for **brood year 1992** is 0.1620%, or 23,488 fish. The Columbia River gill net fishery harvested 10,508 fish, and 3,119 were taken in the ocean. The great majority of fish harvested in the ocean were taken by Canadian fishers. Brood year 1992 was the first year of a three year fin clip study at Spring Creek, and the PEF method was used for this calculation.

Brood year 1993 survival was greater than 1992 at 0.2197%. This survival rate was calculated by the rd program. Oregon reported slightly more ocean recoveries than Canada, but the Columbia River gill net fishery harvested greater than three times more fish than the combined ocean harvest. This brood year contributed more than twenty-four thousand fish to fisheries. The 1993 brood year was the second of three years of a three year fin clip study at Spring Creek.

In the third year of the fin clip study (**brood year 1994**) fish were released which were not represented by any recoverable coded-wire tags. Therefore, the PEF method was used (rd.prg). Survival is now estimated to be 0.1038%. Nevertheless, 4,430 fish were harvested in the ocean, mostly in British Columbia. An additional 6,259 fish were harvested in the Columbia River.

The current total survival estimate for **brood year 1995** is 0.0492%, the lowest since brood year 1984. Nevertheless, 3,586 fish were harvested in the Columbia River, and over eighteen hundred in the ocean fisheries, the majority of which were taken off the coast of Oregon.

The estimated survival for **brood year 1996** is a greatly improved 0.4249, nearly ten times better than the previous brood year, and the best survival since brood year 1990. Over twenty-six thousand fish were harvested in the Columbia River and Estuary, and an estimated 16,111 were caught in West Coast ocean fisheries. Seventy-two hundred fish were harvested in WDFW reported ocean fisheries, fifty-five hundred in Oregon, and three thousand in British Columbia.

There are now over one thousand observed recoveries from **brood year 1996**, and the estimated survival is 0.4250%. Ocean harvest was highest for Washington (7,247), Oregon (5,492), and British Columbia (3,093). Over twenty-six thousand fish were harvested in the Columbia River fisheries, with 19,213 fish making it back to the hatchery (when calculated by expansion).

Nevertheless, this return is near the median for Spring Creek tules.

Changed expansion factors and additional recoveries raise survival to 0.1145% for **brood year 1997**, now fourth from last for Spring Creek. Columbia River gill netters took the majority of fish off station (5,957). Canadian ocean sport and troll fishers harvested an additional 3,414. Washington and Oregon ocean fisheries took 744 and 1,129 fish respectively.

The addition of nearly six hundred observed recoveries increases the estimated survival for **brood year 1998** to 1.3003%, yielding 137,732 returning or harvested fish. ODFW report the greatest number of ocean recoveries (19,361). 44,427 fish were taken in the Columbia River. WDFW reported ocean tag code recoveries that are equivalent to 12,843 fish, and CDFO catch was 8,593.

Brood year 1999 is now the most successful in the series represented in this report with a survival rate of 1.4381% and total of 83,258 fish. Columbia River contribution was 80,592, and ocean recoveries totaled 83,341.

The estimated survival is 1.1122% for **brood year 2000**. BC fisheries harvested over twenty thousand fish, nearly seventeen thousand by ocean trollers, and about thirty-five hundred in sport fishing. Ocean trollers were responsible for about three quarters of the 15,007 fish harvest in Washington ocean fisheries, with charter and sport fisheries taking the remaining fifty-two hundred. Intermittent ladder operations led to an estimated 4,192 on spawning grounds.

iv

Survival is essentially the same for **brood year 2001** at 1.1246%. Another 4,479 fish were found on spawning grounds out of a total estimated survival of 181,285 fish. British Columbia fisheries harvested the greatest number of fish in the ocean (36,002), approximately the same number as was harvested by gill net in the Columbia River (39,671). Total Ocean take was 65,768 fish. Nearly forty thousand fish were harvested in the Columbia River gill net fishery, while about twenty-five hundred were caught in the in river sport fisheries.

This next **brood year, 2002** drops to below the average (when including brood year 2003) of 0.5413% at 0.2945%. Ocean and Columbia Basin fisheries come close to splitting the harvest evenly: 12,487 in the ocean and 13,554 in the basin. The number of fish estimated on spawning grounds dropped to 740.

Survival drops to 0.1631% for **brood year 2003**, placing it in the lower third of survivals. Even at this level of survival, 7,073 fish were harvested in the ocean, mainly from Washington to British Columbia, and another eighty-five hundred were caught in river. Canadian fisheries took the great majority of the catch. Columbia River gill nets harvested more fish than ocean fisheries (8,118 and 7,073 respectively). The 203 spawning ground recoveries are 2.5% of return to the hatchery.

Brood year 2004 is in the bottom 20% of survival rates at 0.0716% and well below the brood year 2006 inclusive average of 0.5208%. Still, over forty-nine hundred fish were harvested in the Columbia River Basin, and nearly twenty-six hundred were caught in ocean fisheries. Gill nets were responsible for the lion's share of the in basin harvest (4,452) while troll Fisheries in British Columbia caught the vast majority of fish taken in the ocean.

Survival rises to an average value of 0.5216% for the next **brood year, 2005**. Release of these fingerlings yielded a total survival of nearly ninety-nine thousand jacks and adults with nearly two thirds of them going to the harvest. Over thirty-eight thousand fish were taken in gillnets, the largest means of harvest for these fish in the basin. BC and Washington took most of the ocean harvest with 9,389 and 8,543 respectively.

Survival for **brood year 2006** is 0.1732%, as it was in last year's report. Even with this relatively weak showing, the release of fifteen and a half million fingerlings from this brood year provided an estimated 12,511 fish for Columbia River harvest, and 4,4415 in the ocean. Brood year 2006 is in the lower third by survival rate of those on record.

The initial survival estimate for brood year 2007 is 0.8805%, a rebound from brood year 2006 and the highest since brood year 2001.

Warm Springs National Fish Hatchery

Warm Springs NFH is located on the Warm Springs River, approximately 14 miles north of Warm Springs, Oregon at 1,525 feet above sea level. The hatchery was authorized in 1966 and began operations in 1978. The Confederated Tribes of the Warm Springs Indian Reservation of Oregon entered into an agreement with the USFWS to increase tribal fishing opportunities. Operations at the hatchery presently consist of adult collection, egg incubation and rearing of spring Chinook salmon.

Warm Springs has a unique fall release strategy that involves releasing the same coded-wire tags in the fall and in the spring. Therefore fall release and spring release fish are combined in this report.

The **1987 brood year** is the first group of fish coded-wire tagged since the late 1970s. Since then, every fish released from the hatchery has been marked. The 1987 brood year had a 0.1272% survival and contributed mainly to the freshwater sport fishery on the Deschutes River. The number of fish released has previously been reported as 609,409 but the total released for this brood year is 661,019. The number of fish with unshed coded-wire tags is 514,633. Groups of Left Ventral (LV) and Right Ventral (RV) fin clipped fish without CWTs were also released. Ventral clipped fish are represented by CWTd fish that were raised in the same raceway. The use of either the PEF or "by tag code" method of estimating survival is problematic. The "by tag code" method was used.

As with the previous brood year, there were LV and RV fish released along with the coded-wire tagged fish and expansions were done for each CWT. Brood year 1988 survived at a rate of 0.1699%, better than **brood year 1987**. These fish contributed primarily to the freshwater sport fishery in Oregon and the treaty fisheries in both the Columbia River and Deschutes River.

LV and RV clipped fish were released again with **brood year 1989**, and the "by tag code" method was used to estimate survival. Total survival for brood year 1989 now stands at 209 fish, or 0.0195%. Sixty-three fish were taken in the freshwater sport fishery, 10 in treaty subsistence, and 4 in treaty ceremonial.

Only an estimated 30 fish survived from a release of 659,507 from the **1990 brood year**, a survival rate of 0.0045%, using the rd2 program. There were 24 recoveries at the hatchery and 6 in the Oregon freshwater sport fishery.

The rd2 program for **brood year 1991** yields a survival of 0.0169%, or only 94 fish. Treaty ceremonial and subsistence took a total of nine fish.

There was a big jump in survival for **brood year 1992**. Over eight hundred fish were recovered to give a survival rate of 0.1569%, when calculated by rd2 program, near the median of survivals. Treaty fisheries harvested 20 fish, while the freshwater sport harvest was 111.

The estimated total survival rate of 0.2564% for **brood year 1993** is in the middle third of survivals. While treaty fisheries took only 3 fish, freshwater sport and fish traps recovered 57. An unusual harvest of 5 fish in California ocean fisheries was noted for this brood year. In all likelihood ODFW hatchery recoveries include the tags from 99 adults that were transferred to the ODFW Round Butte hatchery.

The **brood year 1994** release at Warm Springs included fish raised from eggs received from the ODFW Round Butte Hatchery. Fish were also received from Round Butte as yearlings. Round Butte fish were coded-wire tagged and fin clipped by removing the adipose and left ventral fins. A total of 321,363 Round Butte fish were released. The total survival for this brood year is 0.1366%, near the median for this hatchery and species. Treaty fisheries reported taking 13 fish before they returned to the hatchery.

Brood year 1995 achieved a new record high survival rate in this series with an estimated survival of 0.4163%. Over nineteen hundred fish were recovered from this release. More than 95% of the fish returned to the hatchery, with 55 additional fish reported as ODFW hatchery recoveries, which are most likely from 50 adults transferred to the ODFW Round Butte hatchery. There were again recoveries reported in the California troll fishery. The treaty ceremonial fishery reported and estimated 15 recoveries.

Total estimated survival for **brood year 1996** set a new record for Warm Springs spring Chinook at 0.6122%. Fish were available for both treaty fisheries (316) and freshwater sport fisheries (905) and, to a lesser extent the Columbia River gill net fishery (31).

Warm Springs NFH spring Chinook **brood year 1997** survival continues an upward trend at 0.6641%. There were 11 expanded recoveries in the ODFW reported non-treaty ocean troll fishery from this

brood year, the first time this has been observed. Most of the Columbia River Basin recoveries, which totaled 1,687, were in sport fisheries. There were an estimated 53 recoveries in tribal fisheries. This survival was adjusted due to a recalculation of the expansion rate for recoveries at Warm Springs NFH.

Brood year 1998 sets a record for **Warm Springs spring Chinook** with a survival of 1.4569%. There were recoveries again in the Oregon ocean troll fisheries (16). For the first time there were 3 recoveries in Alaska. Over two thousand fish were taken in freshwater sport fisheries, and nearly fifteen hundred were harvested by Columbia River gill netters. Treaty harvest was 77. This survival was adjusted due to a recalculation of the expansion rate for recoveries at Warm Spring NFH.

Although survival is only 63% of the previous brood year at 0.9837%, **brood year 1999** survival is the third highest on record. (A slight adjustment in the Columbia River gill net recoveries since last year added one expanded recovery.) British Columbia and Oregon reported ocean recoveries, albeit a total contribution of 3 fish. Total freshwater sport harvest was 871 and 302 were taken in gill nets. Treaty harvest was 21 fish.

Estimated survival for **brood year 2000** is 1.2396%, the second most successful on record. Twenty-six fish were harvested in the ocean, the majority of those in Canada. Tribal fishers took 9 fish in the ceremonial and subsistence fisheries. Sport fisheries reported harvest of 681 fish, and 480 were taken in gill nets.

Survival dropped to a sixth of the previous brood year at 0.2051% for **brood year 2001**. A correction of expansion factor for fish sampled at Warm Springs resulted in a slight drop since last year. Gill net recoveries totaled 53, and freshwater sport recoveries totaled 192. Treaty subsistence took another 2 fish, and 7 fish were caught from BC through Oregon.

With an estimated survival of 0.6205% (reduced by 1 recalculated expanded recovery), **brood year 2002** is ranked fourth best of brood years to date, although ninety-four percent of recoveries were at the hatchery. The Columbia River contribution of 145 fish had most recovered in sport fisheries, while 19 fish were harvested in the ocean, as reported by CDFO.

Brood year 2003 survival is 0.1475%, well below the average to date of 0.4164%. Brood year 2003 is ranked 14 of 21 by survival. Only 154 fish were harvested off station from the entire brood year.

Survival increases to the current average for **brood year 2004**, at 0.4502%. Nearly all off station recoveries were in either the gill net or sport fisheries in the Columbia River (307/385) with another 96 in "freshwater sport". One observed ODFW ocean troll recovery expands to a total of 3.

A nearly identical survival of 0.4266% has been recorded for **brood year 2005**. Sport fisheries took three times the number of fish as gillnets, 367 to 120. Over four times as many fish returned to the hatchery as were harvested off station.

Survival for **brood year 2006** rose from the previous brood year to 0.6373%, and this is now the second most successful release from Winthrop. Close to four hundred fish were harvested in the Columbia River, nearly equally divided between gillnet, and sport fisheries. Three fish were also taken in the British Columbia non-treaty ocean troll fishery.

The initial estimate for brood year 2007 is again higher than the previous one at 0.7168%.

Willard National Fish Hatchery

Willard NFH is situated four miles upstream of Little White Salmon NFH on the Little White Salmon River at an altitude of 900 feet. Willard is part of the Little White Salmon NFH complex. Willard NFH was constructed in 1952 under authorization of the Mitchell Act and was originally planned as a fall Chinook hatchery. The hatchery was switched to a coho facility because of cold water temperatures. Since the mid-1960s, the hatchery has been used primarily for coho production. Adult coho were trapped and spawned at Little White Salmon NFH. Coho are presently raised for the Yakama Nation's Wenatchee River program, but no longer released at Willard NFH.

Fish from **brood year 1988** contributed mostly to ocean fisheries (27,996 fish) from British Columbia (1,473) to California (5,189). An additional 8,776 fish were harvested in the Columbia River. Fifty-five hundred of those were taken in the gill net fishery, and 3,200 in the estuary sport fishery. Total survival rate was estimated to be 1.6976%.

Brood year 1989 coho survived at a rate of 0.3247%. Total survival was estimated to be 8,355 fish from a release of 2,573,323. They contributed mainly in ocean sport and non-treaty troll fisheries of Oregon (3,126) and Washington (1,779).

Using the rd2 program, total survival for **brood year 1990** was only 1,722 fish or 0.1097%. Fish were harvested primarily in sport fisheries off the coasts of Washington (795), Oregon (221) and California (63). One hundred twenty-six fish were taken in the Washington non-treaty troll fishery. One hundred twenty-nine fish were taken in the Columbia River sport fishery.

Survival of **brood year 1991** was even lower than 1990 with an overall survival of 0.0720%, or 2,207 fish from a release of over 3 million, using the rd2 program. The only fish harvested from this brood year were 550 fish taken in the Columbia River, 489 in the gill net fishery and 61 in the freshwater sport fishery.

There was some improvement in **brood year 1992** with a survival rate of 0.1007%. Washington ocean sport fisheries took 240 fish, and Oregon ocean sport 120. The Columbia River gill net fishery took an additional 161 fish for a total off station harvest of 521 fish from a release of nearly 2 million.

Brood year 1993 survival is in the same range as the previous three broods at 0.1219% when calculated by the rd2 program. Reporting by ODFW shows 85 fish taken in the Columbia River sport harvest, and 14 at an ODFW hatchery. All other recoveries were at Little White Salmon NFH.

The estimate of survival for **brood year 1994** is 0.2379%, a total of 5,602 fish, most of which were adults returning to the hatchery. Coho from the WDFW Kalama River hatchery were released as part of this brood year, and did not have a unique coded-wire tag. Since there was no representative coded-wire tag for these fish, the rd program was used to estimate survival.

The survival for **brood year 1995** is 0.5383%, now the third best during the 1990s. This higher survival rate is to a large extent due to the release at Little White (rather than Willard) of coho from the ODFW Klaskanine hatchery. CWT recoveries show that these fish survived at a rate over 3 times higher than the Willard fish. There were an estimated 928 recoveries in the ocean and 1,918 in the Columbia River. The Klaskanine fish were 100% adipose fin clipped, and, although there was a coded-wire tag only (no adipose fin clip) group released this brood year, Willard coho were not "mass marked" with an adipose fin clip.

Brood Year 1996 Bonneville coho, some raised at the ODFW Cascade hatchery, and eggs incubated at the Eagle Creek NFH, were released along with Willard fish to make up this brood year. Bonneville stock fish did not receive a unique coded-wire tag. A PEF was calculated for this brood year. The calculated total survival of 0.156% was higher than only three other brood years. Only 72 fish from this brood year were caught in the ocean, and just 324 in the Columbia River.

The estimation of total survival for **brood year 1997** is 0.8639%. This appears to be the first brood year that was mass marked by removing the adipose, except for one group with a coded-wire tag and no adipose fin clip. The brood year has the second highest rate of survival to date. Ocean fisheries reported by ODFW account for 1,286 adults harvested, compared to 724 reported by WDFW, and 1,450 in the Columbia River.

ODFW recoveries for **brood year 1998** are once again in the RMIS database, and total survival is now estimated at 1.0344%. Nearly five thousand fish were harvested in Washington ocean fisheries, 1,853 in ODFW reported ocean fisheries, and 3,931 fish were taken in the Columbia River.

Both Little White and Eagle Creek stock were released for **brood year 1999**, and the two stocks did not receive separate coded-wire tags. Releases for brood year 1999 included two coded-wire tags, one of them a "double index" tag where fish were not adipose clipped; and with an adipose clip. This is now standard operating procedure. However, it appears that the DIT tag was put into fish

from Eagle Creek which were raised and released at Willard. Estimated survival for brood year 1999 is 0.2821% with 385 ocean recoveries and a contribution of 818 in the Columbia River.

Changes in WDFW reporting give **brood year 2000** increases survival to 1.1865%, the second best survival rate to date. Ocean contribution is now estimated to be 4,522 with the harvest there split almost equally between Washington and Oregon reported fisheries. Approximately nine hundred fish were caught by Columbia River gill net and estuary sport fishers. Close to five hundred fish total were estimated to have been found in spawning ground surveys by both ODFW and WDFW.

The **brood year 2001** release did not fare as well, with an estimated survival of 0.8227%, but still had the fourth highest survival rate. Ocean contribution was 1,133 fish while Columbia River non hatchery recoveries were 453 with 197 of those in spawning ground surveys.

The last release coho at Willard NFH was of **brood year 2002** fish, and fish were released in January at 25/#, rather than in April at 15/#. The estimated survival for these fingerlings is 0.0757%. Only 63 fish were taken in various Columbia River fisheries with an ocean harvest of 84.

Winthrop National Fish Hatchery

Winthrop NFH is situated along the Methow River, near the town of Winthrop, Washington. Elevation is 1,760 feet above sea level. Nine dams separate Winthrop from the Pacific Ocean. The hatchery is authorized by both the Grand Coulee Fish Maintenance Project (1937), and the 1938 Mitchell Act.

The facility began operations in 1942 by trapping adult sockeye and steelhead at Rock Island Dam and transporting them to the hatchery. By 1951 the hatchery was rearing sockeye, spring Chinook, steelhead, kokanee, coho, and resident trout. Until 1996 the program had been simplified to spring Chinook only. Coho and steelhead now being raised at Winthrop have not been coded-wire tagged. Tagged summer Chinook were released in 1996.

Spring Chinook

Spring Chinook were coded-wire tagged beginning with brood year 1989. There were 16 observed recoveries from the 107,670 tagged fish released. Using the newer rd2 method of expanding for each tag code, these recoveries are expanded to a total of only 81 fish for an overall survival rate of 0.0077%. An estimated 27 fish were recovered in the freshwater sport fishery, and an estimated 10 fish were recovered by WDFW hatcheries.

All spring Chinook yearling fish released from Winthrop beginning with brood year 1990 have had representative coded-wire tags. Therefore, the rd2 program has been used for all subsequent brood years.

There is only one recovery from coded-wire tagged **brood year 1990** Winthrop spring Chinook. That recovery was at a WDFW hatchery. This recovery is expanded to a total of 8 to include unmarked fish that were probably also killed at that hatchery. The overall survival rate is therefore estimated to be 0.0013%.

There are no **brood year 1991** recoveries from the 189,187 tagged fish released. Brood year 1992 contributed an estimated 74 fish to a WDFW freshwater fish trap from the total estimated recovery of 94 fish, a survival rate of 0.0169%.

Survival for brood year 1993 has been reported as the highest on record to date for Winthrop at 0.0429%, up from last year. WDFW reported more hatchery recoveries than USFWS, and there were no other off station recoveries. It appears that 8 USFWS recoveries are missing from RMIS.

Brood year 1994 sets the new record for Winthrop Spring Chinook at 0.0543%. Fifty-eight of the hatchery recoveries were reported by WDFW. These adults were trapped at Wells Dam and taken to the WDFW Methow hatchery. There was also one spawning ground survey recovery at river mile 48 in the Methow River.

Brood year 1995 fish were raised and released at both Winthrop NFH and Methow SFH. Release records in the RMIS database (reported by WDFW) are somewhat different than records provided by Winthrop. Returning fish from this brood year were captured at Wells Dam and transported to Winthrop NFH. Thus, there were 117 WDFW reported hatchery recoveries vs. 3 FWS reported hatchery recoveries. Two fish were recovered in Canadian ocean fisheries, and the only other recoveries were in the WDFW spawning ground group. Total survival for this brood year is estimated at 0.8609%. Although this is by far the highest survival to date, the unusual circumstances must be noted.

The addition of four WDFW gill net fishery recoveries increases the estimated rate of survival to 0.3559% for **brood year 1996** Winthrop spring Chinook. WDFW recoveries yield an estimated 139 fish recovered on spawning grounds, all age-4 fish in the year 2000. Three fish were reported caught by Alaska.

Survival for **brood year 1997** is lower than for 1996 at 0.2836%. The half million release produced an estimated contribution of 1,546 jacks and adults. Gill net and sport fisheries intercepted the greatest number of fish: 507 and 378 respectively.

The **brood year 1998** release of 364,632 fish yielded a contribution of 1,440 for a survival rate of 0.3813%. Columbia River harvest was split almost equally between gill net and sport fisheries. Returns to the hatchery were also split nearly equally between FWS and WDFW.

Survival for **brood year 1999** is 0.0444%, in the middle of survivals for Winthrop, but still about one tenth of the previous brood year. Based on this accounting, only 48 fish returned to the hatchery, while another 30 were harvested or found elsewhere. However, it should be noted that the return to the hatchery by coded-wire tags is less than one tenth of the number by age composition.

Brood year 2000 is now in fourth place for best survivals with a total survival of 0.3041%, in **Winthrop NFH spring Chinook** history. About one fourth of fish recovered at hatcheries were recovered at the nearby Methow Hatchery. Another 132 fish, more than recovered at the Methow Hatchery, were found in spawning ground surveys.

An examination of the relationship between hatchery recoveries and spawning ground recoveries shows that the proportion of Winthrop fish recovered on spawning grounds has increased dramatically from the early 1990s through the early 2000s. A combination of high survival rates for recent brood years, ESA issues, and co-manager agreements in the Methow Basin required hatchery staff to block returning fish from entering the hatchery after the hatchery escapement goal had been reached.

The calculated survival for **brood year 2001**, 0.1111%, is below average for Winthrop spring Chinook to date. Spawning ground recoveries were again a large part of "recovered" fish, estimated at 141 from a total "survival" of 513. About 66 fish were taken in fisheries within the Columbia River Basin, and 134 fish were found on spawning grounds.

2002 brood year has essentially the same estimated survival at 0.1057%. An even greater proportion of recoveries were in spawning ground surveys, 220 out of 611 total estimated survival, or, more than one third of total survival. Sport and Treaty Ceremonial account for only 6 fish.

The survival estimate for **brood year 2003** is 0.0703%. While this is about 1/3 of the average to date, it is also in the middle of survivals by ranking. Unusual events with brood year 1995 fish, which have a survival rate twice that of the next highest brood year, account for this discrepancy. Spawning Ground recoveries account for close to forty percent of total survival.

An estimated survival of 0.2186% for **brood year 2004** is above the current average of 0.1996%. One hundred eighteen fish were harvested in gill nets. Expansion of WDFW spawning ground recoveries shows that 35% of fish were recovered in spawning ground surveys. Age 2 fish were recovered in a test fishery in Alaska, and in a FWS Spawning Ground Survey. Although these show up in the "PSC Expanded Recoveries" with a number of 0, they don't in the other sections of the Stock Assessment print out.

Recoveries at the hatchery in 2009 have not yet made their way into RMIS, complicating the processing for this **brood year, 2005**, as well as the next two brood years. The data from the CRIS database was, however, integrated with RMIS data to meet the needs of this report. A program (Bio2ts1) was written to accomplish this. The combining of datasets yields a survival rate of 0.1119%, which is in the lower tier of survivals. More fish were estimated to have been recovered in Spawning Ground surveys than returned to the hatchery.


Survival for **brood year 2006** is the second best on record at 0.6373%. While seventeen hundred fish escaped to the hatchery, 1,077 were found in Spawning Ground surveys. The gillnet fishery harvested 405 fish, the only reported harvest.

Initial recoveries for brood year 2007 suggest a survival rate of about half that of the previous year at 0.3334%.


Coho

Winthrop began releasing coded-wire tagged coho with brood year 2002 in production per the Yakima Nation. However, coded-wire tags recovered at Winthrop have never been reported to RMIS, making it impossible to accurately report on this program.


Carson NFH Spring Chinook yearlings




Carson NFH Spring Chinook yearlings




Dworshak NFH Spring Chinook yearlings




Dworshak NFH Spring Chinook fingerlings




Dworshak NFH Summer Steelhead yearlings




Dworshak NFH Summer Steelhead yearlings




Eagle Creek NFH Coho yearlings




Eagle Creek NFH Coho yearlings




Entiat NFH Spring Chinook yearlings




Entiat NFH Spring Chiook yearlings




Kooskia NFH Spring Chinook yearlings




Kooskia NFH Spring Chinook yearlings




Leavenworth NFH Spring Chinook yearlings




Leavenworth NFH Spring Chinook yearlings




Little White NFH Spring Chinook yearlings




Little White NFH Spring Chinook yearlings




Little White NFH Upriver Bright Fall Chinook fingerlings




Little White NFH Upriver Bright Fall Chinook fingerlings




Spring Creek NFH Tule Fall Chinook fingerlings Total Percent Survival




Spring Creek NFH Tule Fall Chinook fingerlings




Warm Springs NFH Spring Chinook yearlings




Warm Springs NFH Spring Chinook yearlings




Willard NFH Coho yearlings




Willard NFH Coho yearlings



Winthrop NFH Spring Chinook yearlings



Winthrop NFH Spring Chinook yearlings



Abernathy Fish Technology Center

Abernathy Fish Technology Center (FTC) is located 14 miles west of Longview, Washington, on Abernathy Creek, approximately three miles upstream from its confluence with the Columbia River at an altitude of 175 feet above sea level. Abernathy FTC began operations in 1959. From 1980 through the 1990s, lower river tule fall Chinook were the only species reared on a production basis at Abernathy as part of Mitchell Act (NOAA Fisheries) funding.

Coded-wire tags were released in fish from brood years 1980 and 1981. Current estimated survival rates are 1.1511% and 0.2084%. Changes in reporting by the Canadian Department of Fisheries and Oceans (CDFO) were noted in the 1997 report. Nevertheless, the CDFO ocean troll fishery took most of these fish.

There were no releases of marked fish for brood years 1982 to 1988.

Marking fish with coded-wire tags was resumed for brood year 1989 under the Missing Production Group program. Brood year 1989 contributed mainly to troll and sport fisheries in Washington (1,628) and British Columbia (977). Overall survival was 0.1836% or 3,296 fish.

Fish from brood year 1990 had a low survival rate of 0.0746%. Approximately sixty thousand fish were released a month earlier and at one tenth the weight of fish represented by coded-wire tags. Therefore, the rd2 program was used to expand the contribution of each group of coded-wire tagged fish. An estimated 99 fish were taken in the ocean off Washington, and 130 in the Columbia River estuary sport fishery. An additional 645 fish were estimated to have been in Washington Department of Fish and Wildlife (WDFW) spawning ground surveys, based on 25 expanded recoveries.

Total survival is now estimated to be 0.2195% for brood year 1991. An estimated 1,156 fish were recovered in spawning ground surveys, nearly twice the number returning to Abernathy. Ocean harvest was 913 fish and 1,470 fish were harvested in the Columbia River.

Total survival for brood year 1992 is now 0.2455%. Hatcheries other than Abernathy recovered an additional 444 fish from the Columbia River contribution of 1,316. Of the 437 fish harvested in the ocean, 348 were in British Columbia.


Total survival for brood year 1993 is 0.2169%. There were 165 recoveries in the ocean, 528 in the Columbia River and 285 at the hatchery.

Brood year 1994 survival fell to an estimated 548 fish, or 0.0381%. WDFW hatcheries reported more recoveries than FWS.


There were only 8 observed recoveries of CWTs released with brood year 1995 fish, but four more than for the previous brood year. These recoveries are expanded to a total estimated contribution of only 298 fish from a release of 1,037,507 for an estimated survival of 0.0224%. The low survival rates for these two brood years are consistent with the low survival for fish released from Spring Creek NFH.

Release of tule fall Chinook occurred through brood year 1998 without coded-wire tags. Brood year 1998 was the last year production tules were released.


Abernathy FTC Tule Fall Chinook fingerlings




Abernathy FTC Tule Fall Chinook fingerlings



Lower River spring Chinook Survival by Brood Year



Mid River spring Chinook Survival by Brood Year



Carson NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 2315382 | 1610 | 190 | 0 | 1 | 1801 | 0.0778 |
| 91 | 2321285 | 450 | 92 | 0 | 12 | 554 | 0.0239 |
| 92 | 2040568 | 8137 | 4289 | 82 | 31 | 12539 | 0.6145 |
| 93 | 2195192 | 4720 | 4825 | 42 | 8 | 9595 | 0.4371 |
| 94 | 1722621 | 1082 | 1045 | 0 | 14 | 2141 | 0.1243 |
| 95 | 907708 | 2425 | 997 | 0 | 0 | 3422 | 0.377 |
| 96 | 1734188 | 10047 | 8953 | 0 | 144 | 19144 | 1.1039 |
| 97 | 1415744 | 8086 | 9890 | 14 | 168 | 18158 | 1.2826 |
| 98 | 1430022 | 6876 | 12789 | 0 | 327 | 19992 | 1.398 |
| 99 | 1608684 | 7218 | 13067 | 0 | 327 | 20612 | 1.2813 |
| 00 | 1449361 | 2910 | 6962 | 25 | 0 | 9897 | 0.6829 |
| 01 | 1673255 | 1816 | 2426 | 20 | 0 | 4262 | 0.2547 |
| 02 | 1417986 | 2675 | 2451 | 0 | 0 | 5126 | 0.3615 |
| 03 | 1470134 | 1702 | 1128 | 0 | 0 | 2830 | 0.1925 |
| 04 | 1209384 | 1648 | 4924 | 0 | 0 | 6572 | 0.5434 |
| 05 | 1158425 | 1935 | 3215 | 0 | 0 | 5150 | 0.4446 |
| 06 | 1330836 | 6102 | 12420 | 0 | 0 | 18522 | 1.3918 |
| 07 | 1216198 | 2949 | 1917 | 27 | 0 | 4893 | 0.4023 |

Dworshak NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 959369 | 22 | 4 | 0 | 0 | 26 | 0.0027 |
| 91 | 467222 | 9 | 1 | 0 | 0 | 10 | 0.0021 |
| 92 | 1278273 | 688 | 56 | 0 | 3 | 747 | 0.0584 |
| 93 | 1311445 | 2890 | 587 | 8 | 23 | 3508 | 0.2675 |
| 94 | 102903 | 60 | 19 | 3 | 1 | 83 | 0.0807 |
| 95 | 53078 | 86 | 4 | 0 | 0 | 90 | 0.1696 |
| 96 | 973400 | 3410 | 901 | 0 | 2 | 4313 | 0.4431 |
| 97 | 1044511 | 896 | 2932 | 6 | 0 | 3834 | 0.5949 |
| 98 | 1017873 | 3338 | 5333 | 24 | 40 | 8735 | 0.8582 |
| 99 | 333120 | 253 | 111 | 0 | 0 | 364 | 0.1093 |
| 00 | 1000561 | 2864 | 2295 | 47 | 0 | 5206 | 0.5203 |
| 01 | 1033982 | 656 | 552 | 0 | 24 | 1232 | 0.1192 |
| 02 | 1074009 | 1696 | 776 | 0 | 112 | 2584 | 0.2406 |
| 03 | 1070826 | 792 | 552 | 0 | 96 | 1440 | 0.1345 |
| 04 | 1007738 | 1563 | 2212 | 0 | 0 | 3775 | 0.3746 |
| 05 | 963211 | 193 | 116 | 0 | 0 | 309 | 0.0321 |
| 06 | 939000 | 204 | 364 | 0 | 11 | 579 | 0.0617 |

Dworshak NFH summer Steelhead yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 1192593 | 1014 | 1567 | 28 | 0 | 2609 | 0.2188 |
| 91 | 1224101 | 486 | 1096 | 25 | 0 | 1607 | 0.1313 |
| 92 | 1217990 | 1946 | 806 | 0 | 0 | 2752 | 0.2259 |
| 93 | 1410283 | 2432 | 643 | 0 | 0 | 3075 | 0.218 |
| 94 | 1213577 | 2771 | 900 | 0 | 0 | 3671 | 0.3025 |
| 95 | 1377435 | 2007 | 615 | 0 | 0 | 2622 | 0.1904 |
| 96 | 1361034 | 1320 | 1037 | 0 | 0 | 2357 | 0.1732 |
| 97 | 1228944 | 2821 | 1451 | 0 | 0 | 4272 | 0.3476 |
| 98 | 1479940 | 4273 | 1311 | 0 | 0 | 5584 | 0.3773 |
| 99 | 1427083 | 4344 | 2712 | 0 | 0 | 7056 | 0.4944 |
| 00 | 1247550 | 1775 | 1671 | 0 | 0 | 3446 | 0.2762 |
| 01 | 1365623 | 1820 | 771 | 0 | 0 | 2591 | 0.1897 |
| 02 | 1210919 | 1936 | 2114 | 0 | 0 | 4050 | 0.3345 |
| 03 | 1202055 | 187 | 2589 | 0 | 0 | 2776 | 0.2309 |
| 04 | 1122064 | 1306 | 2130 | 0 | 0 | 3436 | 0.3062 |
| 05 | 1206565 | 3024 | 4456 | 0 | 0 | 7480 | 0.6199 |
| 06 | 154138 | 207 | 225 | 0 | 2 | 434 | 0.2816 |

Eagle Creek NFH coho yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 1087346 | 429 | 286 | 620 | 0 | 1335 | 0.1228 |
| 91 | 1060888 | 3267 | 205 | 137 | 0 | 3609 | 0.3402 |
| 92 | 980327 | 3696 | 215 | 1109 | 0 | 5020 | 0.5121 |
| 93 | 987877 | 1867 | 121 | 358 | 0 | 2346 | 0.2375 |
| 94 | 996618 | 1423 | 66 | 263 | 0 | 1752 | 0.1758 |
| 95 | 769509 | 14031 | 1052 | 1078 | 0 | 16161 | 2.1002 |
| 96 | 1010044 | 13249 | 1566 | 2549 | 0 | 17364 | 1.7191 |
| 97 | 1147711 | 39182 | 1214 | 5255 | 0 | 45651 | 3.9776 |
| 98 | 1006688 | 30964 | 8351 | 9966 | 174 | 49455 | 4.9126 |
| 99 | 711927 | 11485 | 1323 | 883 | 29 | 13720 | 1.9272 |
| 00 | 508321 | 6856 | 451 | 523 | 20 | 7850 | 1.5443 |
| 01 | 505400 | 5170 | 360 | 640 | 21 | 6191 | 1.225 |
| 02 | 557016 | 7322 | 386 | 572 | 0 | 8280 | 1.4865 |
| 03 | 524356 | 4477 | 580 | 1001 | 23 | 6081 | 1.1597 |
| 04 | 459486 | 10145 | 634 | 3418 | 73 | 14270 | 3.1056 |
| 05 | 528546 | 3399 | 339 | 444 | 2 | 4184 | 0.7916 |
| 06 | 511342 | 11425 | 1530 | 5243 | 55 | 18253 | 3.5696 |
| 07 | 324944 | 1005 | 87 | 204 | 24 | 1320 | 0.4062 |
| 08 | 355612 | 1418 | 123 | 263 | 26 | 1830 | 0.5146 |

Entiat NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 343150 | 16 | 0 | 0 | 0 | 16 | 0.0047 |
| 91 | 376462 | 96 | 9 | 0 | 0 | 105 | 0.0279 |
| 92 | 378729 | 110 | 21 | 0 | 0 | 131 | 0.0346 |
| 93 | 391211 | 144 | 6 | 0 | 0 | 150 | 0.0383 |
| 94 | 335593 | 213 | 11 | 0 | 0 | 224 | 0.0667 |
| 95 | 200486 | 647 | 8 | 3 | 0 | 658 | 0.3282 |
| 96 | 350784 | 1926 | 108 | 3 | 17 | 2054 | 0.5855 |
| 97 | 354238 | 2277 | 442 | 0 | 20 | 2739 | 0.7732 |
| 98 | 359667 | 3148 | 787 | 0 | 35 | 3970 | 1.1038 |
| 99 | 397855 | 311 | 32 | 0 | 28 | 371 | 0.0933 |
| 00 | 533720 | 740 | 204 | 0 | 16 | 960 | 0.1591 |
| 01 | 395689 | 750 | 177 | 0 | 8 | 935 | 0.2363 |
| 02 | 386833 | 368 | 92 | 0 | 10 | 470 | 0.1215 |
| 03 | 401240 | 26 | 60 | 0 | 16 | 102 | 0.0254 |
| 04 | 322516 | 112 | 319 | 9 | 60 | 500 | 0.155 |
| 05 | 362854 | 561 | 254 | 0 | 84 | 899 | 0.2478 |

Kooskia NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 727251 | 61 | 16 | 0 | 0 | 77 | 0.0106 |
| 91 | 333437 | 2 | 0 | 0 | 0 | 2 | 0.0006 |
| 92 | 305813 | 90 | 9 | 0 | 0 | 99 | 0.0324 |
| 93 | 722906 | 121 | 68 | 15 | 16 | 220 | 0.0327 |
| 94 | 333794 | 0 | 9 | 0 | 0 | 9 | 0.0027 |
| 95 | 16598 | 1 | 0 | 0 | 0 | 1 | 0.006 |
| 96 | 43165 | 381 | 26 | 0 | 1 | 408 | 0.9452 |
| 97 | 570290 | 656 | 2039 | 9 | 6 | 2710 | 1.1573 |
| 98 | 365150 | 852 | 1878 | 0 | 68 | 2798 | 0.7663 |
| 99 | 82974 | 75 | 38 | 0 | 4 | 117 | 0.141 |
| 00 | 549861 | 719 | 927 | 16 | 25 | 1687 | 0.3068 |
| 01 | 548097 | 185 | 85 | 10 | 0 | 280 | 0.0511 |
| 02 | 591380 | 450 | 294 | 12 | 6 | 762 | 0.1289 |
| 03 | 616385 | 150 | 204 | 0 | 0 | 354 | 0.0574 |
| 04 | 634947 | 1225 | 2226 | 0 | 91 | 3542 | 0.5578 |
| 05 | 174218 | 113 | 89 | 0 | 0 | 202 | 0.1159 |
| 06 | 112344 | 127 | 309 | 0 | 3 | 439 | 0.3908 |

Leavenworth NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 2286828 | 67 | 11 | 0 | 0 | 78 | 0.0034 |
| 91 | 1757931 | 360 | 14 | 0 | 0 | 374 | 0.0213 |
| 92 | 1522846 | 987 | 161 | 0 | 0 | 1148 | 0.0754 |
| 93 | 1712648 | 3477 | 1200 | 0 | 30 | 4707 | 0.2748 |
| 94 | 1706060 | 905 | 84 | 0 | 0 | 989 | 0.058 |
| 95 | 919025 | 811 | 248 | 0 | 73 | 1132 | 0.1232 |
| 96 | 1701753 | 5064 | 1000 | 26 | 286 | 6376 | 0.3747 |
| 97 | 1636402 | 5677 | 6120 | 0 | 762 | 12559 | 0.7675 |
| 98 | 1680904 | 7136 | 3630 | 39 | 1190 | 11995 | 0.7136 |
| 99 | 1630089 | 814 | 665 | 0 | 88 | 1567 | 0.0961 |
| 00 | 1554362 | 3298 | 2231 | 4 | 278 | 5811 | 0.3739 |
| 01 | 1288893 | 2481 | 665 | 15 | 81 | 3242 | 0.2515 |
| 02 | 1422100 | 771 | 587 | 9 | 90 | 1457 | 0.1025 |
| 03 | 1476046 | 596 | 486 | 6 | 12 | 1100 | 0.0745 |
| 04 | 1005505 | 1399 | 3140 | 63 | 235 | 4837 | 0.4811 |
| 05 | 1177568 | 1358 | 708 | 0 | 64 | 2130 | 0.1809 |
| 06 | 1539668 | 7560 | 5956 | 36 | 300 | 13852 | 0.8997 |
| 07 | 1685038 | 2205 | 1098 | 0 | 81 | 3384 | 0.2008 |

Little White Salmon NFH spring Chinook yearling Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 1677694 | 260 | 0 | 0 | 0 | 260 | 0.0155 |
| 91 | 809079 | 232 | 0 | 0 | 0 | 232 | 0.0287 |
| 92 | 994588 | 2895 | 1931 | 0 | 600 | 5426 | 0.5456 |
| 93 | 1057864 | 1246 | 1055 | 0 | 0 | 2301 | 0.2175 |
| 94 | 961515 | 322 | 18 | 0 | 0 | 340 | 0.0354 |
| 95 | 682623 | 2381 | 339 | 0 | 56 | 2776 | 0.4067 |
| 96 | 1066702 | 3459 | 1402 | 0 | 44 | 4905 | 0.4598 |
| 97 | 1074173 | 576 | 3358 | 0 | 24 | 3958 | 0.3685 |
| 98 | 1115384 | 3221 | 3813 | 0 | 0 | 7034 | 0.6306 |
| 99 | 1016574 | 480 | 990 | 0 | 0 | 1470 | 0.1446 |
| 00 | 1037382 | 1138 | 1475 | 42 | 0 | 2655 | 0.2559 |
| 01 | 1012339 | 540 | 510 | 0 | 0 | 1050 | 0.1037 |
| 02 | 1036733 | 3493 | 3017 | 105 | 35 | 6650 | 0.6414 |
| 03 | 779716 | 1062 | 1434 | 0 | 0 | 2496 | 0.3201 |
| 04 | 1014274 | 2530 | 2735 | 0 | 10 | 5275 | 0.5201 |
| 05 | 1006105 | 2120 | 4624 | 0 | 8 | 6752 | 0.6711 |
| 06 | 934438 | 4408 | 17672 | 0 | 240 | 22320 | 2.3886 |
| 07 | 1439901 | 952 | 1584 | 0 | 712 | 3248 | 0.2256 |

Little White Salmon NFH upriver bright fall Chinook Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 4029158 | 7142 | 1041 | 4005 | 1047 | 13235 | 0.3285 |
| 91 | 2862783 | 2461 | 1146 | 1426 | 804 | 5837 | 0.2039 |
| 92 | 1866901 | 3892 | 1459 | 905 | 507 | 6763 | 0.3623 |
| 93 | 1797922 | 6678 | 1576 | 3306 | 1980 | 13540 | 0.7531 |
| 94 | 1967894 | 708 | 308 | 309 | 300 | 1625 | 0.0826 |
| 95 | 2097206 | 2634 | 407 | 1213 | 825 | 5079 | 0.2422 |
| 96 | 2153118 | 1329 | 565 | 808 | 286 | 2988 | 0.1388 |
| 97 | 1999435 | 1918 | 2503 | 1334 | 2520 | 8275 | 0.4139 |
| 98 | 2149397 | 2394 | 1204 | 1525 | 2651 | 7774 | 0.3617 |
| 99 | 1970592 | 2207 | 2066 | 4892 | 7220 | 16385 | 0.8315 |
| 00 | 1937764 | 910 | 1279 | 2490 | 1730 | 6409 | 0.3307 |
| 01 | 2074295 | 1350 | 1440 | 1490 | 1650 | 5930 | 0.2859 |
| 02 | 2084184 | 352 | 748 | 924 | 814 | 2838 | 0.1362 |
| 03 | 2031737 | 1600 | 530 | 980 | 360 | 3470 | 0.1708 |
| 04 | 1902741 | 1774 | 323 | 1185 | 70 | 3352 | 0.1762 |
| 05 | 1802174 | 6912 | 2856 | 3234 | 174 | 13176 | 0.7311 |
| 06 | 2061924 | 6868 | 2205 | 1747 | 613 | 11433 | 0.5545 |
| 07 | 2001760 | 13300 | 4287 | 3953 | 606 | 22146 | 1.1063 |

Spring Creek NFH tule fall Chinook Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 14348604 | 5609 | 4408 | 8566 | 0 | 18583 | 0.1295 |
| 91 | 19072355 | 9417 | 7844 | 8938 | 9 | 26208 | 0.1374 |
| 92 | 14310020 | 8831 | 10305 | 3309 | 742 | 23187 | 0.162 |
| 93 | 15607856 | 10471 | 18212 | 5279 | 325 | 34287 | 0.2197 |
| 94 | 15990014 | 7676 | 4319 | 4430 | 171 | 16596 | 0.1038 |
| 95 | 16440491 | 3418 | 2403 | 1821 | 440 | 8082 | 0.0492 |
| 96 | 14547325 | 21201 | 23061 | 16388 | 1178 | 61828 | 0.425 |
| 97 | 15619429 | 5943 | 6308 | 4721 | 0 | 16972 | 0.1087 |
| 98 | 10592075 | 51756 | 41344 | 43276 | 1356 | 137732 | 1.3003 |
| 99 | 16065945 | 68983 | 70881 | 78415 | 7884 | 226163 | 1.4077 |
| 00 | 10569810 | 39000 | 32376 | 42600 | 4200 | 118176 | 1.1181 |
| 01 | 16117906 | 68208 | 42325 | 66246 | 4479 | 181258 | 1.1246 |
| 02 | 15079904 | 18703 | 12448 | 12516 | 740 | 44407 | 0.2945 |
| 03 | 14653529 | 8193 | 8429 | 7073 | 203 | 23898 | 0.1631 |
| 04 | 14533488 | 3037 | 4452 | 2593 | 329 | 10411 | 0.0716 |
| 05 | 15239054 | 37852 | 41500 | 19360 | 140 | 98852 | 0.6487 |
| 06 | 15472561 | 10374 | 12015 | 4415 | 0 | 26804 | 0.1732 |
| 07 | 14899270 | 20901 | 68468 | 41296 | 520 | 131185 | 0.8805 |

Warm Springs NFH spring Chinook Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 659507 | 24 | 6 | 0 | 0 | 30 | 0.0045 |
| 91 | 557014 | 85 | 9 | 0 | 0 | 94 | 0.0169 |
| 92 | 550664 | 731 | 133 | 0 | 0 | 864 | 0.1569 |
| 93 | 398142 | 951 | 65 | 5 | 0 | 1021 | 0.2564 |
| 94 | 420886 | 562 | 13 | 0 | 0 | 575 | 0.1366 |
| 95 | 467427 | 1915 | 28 | 3 | 0 | 1946 | 0.4163 |
| 96 | 790422 | 3562 | 1277 | 0 | 0 | 4839 | 0.6122 |
| 97 | 811570 | 3709 | 1670 | 11 | 0 | 5390 | 0.6641 |
| 98 | 770419 | 7637 | 3568 | 19 | 0 | 11224 | 1.4569 |
| 99 | 827665 | 6941 | 1194 | 7 | 0 | 8142 | 0.9837 |
| 00 | 618822 | 6462 | 1183 | 26 | 0 | 7671 | 1.2396 |
| 01 | 655740 | 1094 | 244 | 7 | 0 | 1345 | 0.2051 |
| 02 | 454312 | 2655 | 145 | 19 | 0 | 2819 | 0.6205 |
| 03 | 738162 | 934 | 153 | 2 | 0 | 1089 | 0.1475 |
| 04 | 676086 | 2243 | 798 | 3 | 0 | 3044 | 0.4502 |
| 05 | 620753 | 2142 | 503 | 3 | 0 | 2648 | 0.4266 |
| 06 | 346568 | 1558 | 398 | 3 | 0 | 1959 | 0.5653 |
| 07 | 580887 | 2903 | 1259 | 1 | 1 | 4164 | 0.7168 |

Winthrop NFH spring Chinook Releases and Returns

| Brood Year | Number Released | Columbia | | | Spawning Ground | Total Recoveries | Smolt to Adult Survival |
|------------|-----------------|----------|---------------|---------------|-----------------|------------------|-------------------------|
| | | Hatchery | River Harvest | Ocean Harvest | | | |
| 90 | 624771 | 8 | 0 | 0 | 0 | 8 | 0.0013 |
| 91 | 950624 | 0 | 0 | 0 | 0 | 0 | 0 |
| 92 | 556313 | 20 | 74 | 0 | 0 | 94 | 0.0169 |
| 93 | 770847 | 328 | 3 | 0 | 0 | 331 | 0.0429 |
| 94 | 112395 | 59 | 1 | 0 | 1 | 61 | 0.0543 |
| 95 | 14520 | 116 | 4 | 2 | 3 | 125 | 0.8609 |
| 96 | 324851 | 939 | 83 | 3 | 131 | 1156 | 0.3559 |
| 97 | 545062 | 583 | 916 | 3 | 44 | 1546 | 0.2836 |
| 98 | 377696 | 678 | 568 | 5 | 189 | 1440 | 0.3813 |
| 99 | 175869 | 53 | 11 | 6 | 8 | 78 | 0.0444 |
| 00 | 201604 | 452 | 27 | 0 | 134 | 613 | 0.3041 |
| 01 | 461678 | 333 | 46 | 0 | 134 | 513 | 0.1111 |
| 02 | 578307 | 384 | 6 | 0 | 221 | 611 | 0.1057 |
| 03 | 550214 | 225 | 9 | 2 | 151 | 387 | 0.0703 |
| 04 | 484090 | 567 | 121 | 0 | 370 | 1058 | 0.2186 |
| 05 | 589693 | 314 | 36 | 0 | 320 | 670 | 0.1136 |
| 06 | 509045 | 1762 | 405 | 0 | 1077 | 3244 | 0.6373 |
| 07 | 371959 | 897 | 59 | 0 | 284 | 1240 | 0.3334 |

ⁱⁱ Environmental survey report pertaining to salmon and steelhead in certain rivers of eastern Oregon and the Willamette River and its tributaries: Pt. II. Survey reports of the Willamette River and its tributaries. Fish Commission of Oregon, Research Division Willis, Raymond A, Melvin D Collins and Roy E Sams Oregon. Fish Commission. Research Division 1960

ⁱⁱⁱ Assessments to Determine the Effect of Current and Alternate Ladder Operations on Brood Stock Collection and Behavior of Hatchery Fall Chinook Salmon at Little White Salmon National Fish Hatchery During 2004-05

Rod Engle and Doug Olson

U.S. Fish and Wildlife Service

Columbia River Fisheries Program Office

1211 SE Cardinal Court, Suite 100

Vancouver, WA 98683

Speros Doulos and Jim Rockowski

Little White Salmon National Fish Hatchery

56961 State Route 14

Cook, WA 98605

August 14, 2006

^{iv} Assessments to Determine the Effect of Current and Alternate Ladder Operations on Brood Stock Collection and Behavior of Hatchery Fall Chinook Salmon at Spring Creek National Fish Hatchery during 2003-2005

U.S. Fish and Wildlife Service

Rod Engle and Doug Olson

Columbia River Fisheries Program Office

Larry Marchant and Mark Ahrens

Spring Creek National Fish Hatchery

2003-2005 ladder closures

Two closures were planned for the 2004 and 2005 hatchery return . These closures were planned to occur during weekends near the beginning and peak of the hatchery return. Closures began near 1:00 PM on Fridays and ended near 1:00 PM on Sundays.

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 04 Spring Chinook raised at Carson NFH
 released in Wind River

02/05/2013
 rd2
 04CaSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|---------------|--------|------------|
| 04/10/2006 | 448,453 | 17.2 | 22,527 | 052182 | Carson NFH |
| 04/10/2006 | 552,520 | 17.2 | 23,099 | 052183 | Carson NFH |
| 04/10/2006 | <u>208,411</u> | 17.6 | <u>24,546</u> | 052184 | Carson NFH |
| Totals | 1,209,384 | | 70,172 | | |

| Where Recovered | Total | 162 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|-----------|----------|----------|----|----------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 1,444 | | 90 | 2 | | 92 |
| Columbia River Sport | ODFW | 1,316 | | 71 | 4 | | 75 |
| Freshwater Net | WDFW | 60 | | 3 | | | 3 |
| Freshwater Sport | WDFW | 2,084 | 21 | 85 | | | 106 |
| Hatchery | FWS | 1,648 | 5 | 88 | 3 | | 96 |
| Treaty Ceremonial | ODFW | <u>20</u> | | <u>1</u> | | | <u>1</u> |
| Totals | | 6,572 | <u>26</u> | 338 | <u>9</u> | | 373 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 6,572 | 100 | 0.5434 |
| Total Survival | 6,572 | | 0.5434 |
| Escapement to NFH | 1,648 | 25 | 0.1363 |
| Columbia River contribution | 4,924 | 75 | 0.4071 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 4,924 | 1,648 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 05 Spring Chinook raised at Carson NFH
 released in Wind River

02/05/2013
 rd2
 05CaSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|---------------|--------|------------|
| 04/12/2007 | 528,799 | 17.2 | 24,772 | 052185 | Carson NFH |
| 04/12/2007 | 448,288 | 17.1 | 24,654 | 052186 | Carson NFH |
| 04/12/2007 | <u>181,338</u> | 17.2 | <u>24,701</u> | 052187 | Carson NFH |
| Totals | 1,158,425 | | 74,127 | | |

| Where Recovered | Total | 194 Observed Recoveries | | | | | Total |
|------------------------|--------------|---------------------------------|----|------------|----|----|------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 357 | | 5 | 25 | 4 | | 34 |
| Columbia River Sport | 970 | | 4 | 62 | | | 66 |
| Freshwater Net | 192 | | 1 | 11 | | | 12 |
| Freshwater Sport | 1,696 | | | 114 | | | 114 |
| Hatchery | <u>1,935</u> | | 7 | <u>120</u> | 5 | | <u>132</u> |
| Totals | 5,150 | | 17 | 332 | 9 | | 358 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 5,150 | 100 | 0.4446 |
| Total Survival | 5,150 | | 0.4446 |
| Escapement to NFH | 1,935 | 38 | 0.1670 |
| Columbia River contribution | 3,215 | 62 | 0.2775 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 3,215 | 1,935 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 06 Spring Chinook raised at Carson NFH
 released in Wind River

12/14/2012
 rd2
 06CaSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|---------------|--------|------------|
| 04/11/2008 | 160,272 | 17.7 | 25,012 | 052585 | Carson NFH |
| 04/11/2008 | 626,743 | 17.6 | 19,380 | 052590 | Carson NFH |
| 04/11/2008 | <u>543,821</u> | 17.1 | <u>25,040</u> | 054805 | Carson NFH |
| Totals | 1,330,836 | | 69,432 | | |

| Where Recovered | Total | 426 Observed Recoveries | | | | | Total |
|-----------------------------|--------|---------------------------------|-----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet ODFW | 5,770 | | | 256 | 2 | | 258 |
| Columbia River Sport ODFW | 2,290 | | | 86 | 8 | | 94 |
| Freshwater Net WDFW | 194 | | 1 | 7 | | | 8 |
| Freshwater Sport ODFW | 24 | | | 4 | | | 4 |
| Freshwater Sport WDFW | 4,120 | | 72 | 156 | 12 | | 240 |
| Hatchery FWS | 6,102 | | 44 | 261 | 10 | | 315 |
| Test Fishery Net ODFW | 22 | | | 1 | | | 1 |
| Totals | 18,522 | | 117 | 771 | 32 | | 920 |

| | | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 18,522 | 100 | 1.3918 |
| Total Survival | 18,522 | | 1.3918 |

| | | | |
|-----------------------------|--------|----|--------|
| Escapement to NFH | 6,102 | 33 | 0.4585 |
| Columbia River contribution | 12,420 | 67 | 0.9332 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 12,420 | 6,102 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 07 Spring Chinook raised at Carson NFH
 released in Wind River

12/14/2012
 rd2
 07CaSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|---------------|--------|------------|
| 04/13/2009 | 617,520 | 16.5 | 24,804 | 053876 | Carson NFH |
| 04/13/2009 | 381,783 | 17.3 | 22,128 | 053877 | Carson NFH |
| 04/13/2009 | <u>216,895</u> | 17.3 | <u>24,681</u> | 054181 | Carson NFH |
| Totals | 1,216,198 | | 71,613 | | |

| Where Recovered | Total | 192 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|-----------|------------|----|----|------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 352 | | 16 | | | 16 |
| Columbia River Sport | ODFW | 693 | | 37 | | | 37 |
| Freshwater Net | WDFW | 34 | | 2 | | | 2 |
| Freshwater Sport | ODFW | 17 | | 1 | | | 1 |
| Freshwater Sport | WDFW | 796 | | 60 | | | 60 |
| Hatchery | FWS | 2,949 | 9 | 156 | | | 165 |
| Sport (private) | WDFW | 27 | 3 | | | | 3 |
| Test Fishery Net | ODFW | 25 | | 1 | | | 1 |
| Totals | | <u>4,893</u> | <u>12</u> | <u>273</u> | | | <u>285</u> |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 27 | 1 | 0.0022 |
| Escapement to Columbia River | 4,866 | 99 | 0.4001 |
| Total Survival | 4,893 | | 0.4023 |
| Escapement to NFH | 2,949 | 60 | 0.2425 |
| Columbia River contribution | 1,917 | 39 | 0.1576 |

| Ocean | Contribution by Area | |
|----------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 27 WA | Columbia River | Escapement to NFH |
| 0 OR | 1,917 | 2,949 |
| 0 CA | | |
| 27 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 04 Spring Chinook raised at Dworshak NFH
 released in Clearwater River

02/08/2013
 rd2
 04DwSCS

1,007,738 released with 134,384 tagged fish and 2 CWTs

| Where Recovered | Total | 322 Observed Recoveries | | | | | Total |
|-----------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet ODFW | 847 | | | 110 | 2 | | 112 |
| Columbia River Sport ODFW | 1,012 | | | 120 | 12 | | 132 |
| Freshwater Sport IDFG | 253 | | | 33 | | | 33 |
| Freshwater Sport ODFW | 22 | | | 3 | | | 3 |
| Freshwater Sport WDFW | 56 | | 7 | | | | 7 |
| Hatchery FWS | 1,539 | | 74 | 125 | 6 | | 205 |
| Hatchery IDFG | 16 | | | 2 | | | 2 |
| Hatchery ODFW | 8 | | | 1 | | | 1 |
| Test Fishery Net ODFW | 15 | | | 2 | | | 2 |
| Treaty Ceremonial ODFW | 7 | | | 1 | | | 1 |
| Totals | 3,775 | | 81 | 397 | 20 | | 498 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 3,775 | 100 | 0.3746 |
| Total Survival | 3,775 | | 0.3746 |
| Escapement to NFH | 1,539 | 41 | 0.1527 |
| Columbia River contribution | 2,236 | 59 | 0.2219 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 2,236 | 1,539 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 02/06/2013
 Broodyear 05 Spring Chinook raised at Dworshak NFH rd2
 released in Clearwater River 05DwSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 03/29/2007 | 90,032 | 0.0 | 61,328 | 050797 | Dworshak NFH |
| 03/28/2007 | 89,441 | 0.0 | 63,772 | 051187 | Dworshak NFH |
| Totals | 179,473 | | 125,100 | | |

Handwritten notes:
 64,307
 63,845
 128,152
 260
 410,083
 424,976
 835,059

Observed Recoveries
 PSC Expanded Recoveries

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|------------------------|------|-------|----|----|-----|----|----|-------|
| Columbia River Gillnet | ODFW | 11 | | | 11 | | | 11 |
| Columbia River Sport | ODFW | 76 | | | 76 | | | 76 |
| Freshwater Sport | IDFG | 27 | | 6 | 21 | | | 27 |
| Freshwater Sport | ODFW | 1 | | | 1 | | | 1 |
| Hatchery | FWS | 192 | | 38 | 154 | | | 192 |
| Hatchery | IDFG | 1 | | | 1 | | | 1 |
| Test Fishery Net | ODFW | 1 | | | 1 | | | 1 |
| Totals | | 309 | | 44 | 265 | | | 309 |

| | | % of total | % of release |
|------------------------------|-----|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 309 | 100 | 0.1722 |
| Total Survival | 309 | | 0.1722 |
| Escapement to NFH | 192 | 62 | 0.1070 |
| Columbia River contribution | 117 | 38 | 0.0652 |

NUMBER is different than last year and wont be replaced!
 Please investigate!
 TAGGED is different than last year and wont be replaced!
 Please investigate!

Ocean Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 117 | 192 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 05 Spring Chinook raised at Dworshak NFH
 released in Clearwater River

02/07/2013
 rd2
 05DwSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 03/29/2007 | 474,390 | 0.0 | 64,307 | 050797 | Dworshak NFH |
| 03/28/2007 | 488,821 | 0.0 | 63,845 | 051187 | Dworshak NFH |
| Totals | 963,211 | | 128,152 | | |

| Where Recovered | Total | 260 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 11 | | 11 | | | 11 |
| Columbia River Sport | ODFW | 76 | | 76 | | | 76 |
| Freshwater Sport | IDFG | 27 | 6 | 21 | | | 27 |
| Freshwater Sport | ODFW | 1 | | 1 | | | 1 |
| Hatchery | FWS | 192 | 38 | 154 | | | 192 |
| Hatchery | IDFG | 1 | | 1 | | | 1 |
| Test Fishery Net | ODFW | 1 | | 1 | | | 1 |
| Totals | | 309 | 44 | 265 | | | 309 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 309 | 100 | 0.0321 |
| Total Survival | 309 | | 0.0321 |
| Escapement to NFH | 192 | 62 | 0.0199 |
| Columbia River contribution | 117 | 38 | 0.0121 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 117 | 192 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 02/06/2013
 Broodyear 06 Spring Chinook raised at Dworshak NFH rd2
 58664 417,858 released in Clearwater River 06DwSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 04/02/2008 | 58,771 | 0.0 | 58,664 | 054132 | Dworshak NFH |
| 04/03/2008 | 59,799 | 0.0 | 59,250 | 054133 | Dworshak NFH |
| Totals | 118,570 | | 117,914 | | |

| Where Recovered | Total | 326 Observed Recoveries | | | | | Total |
|------------------------|----------|---------------------------------|-----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW 219 | | | 211 | 8 | | 219 |
| Columbia River Sport | ODFW 89 | | 13 | 68 | 8 | | 89 |
| Fish Trap (freshwater) | ODFW 3 | | 3 | | | | 3 |
| Freshwater Sport | IDFG 35 | | 17 | 17 | 1 | | 35 |
| Freshwater Sport | ODFW 3 | | | 3 | | | 3 |
| Freshwater Sport | WDFW 14 | | | 10 | 4 | | 14 |
| Hatchery | FWS 197 | | 74 | 111 | 12 | | 197 |
| Hatchery | IDFG 7 | | 3 | 3 | 1 | | 7 |
| Spawning Ground | FWS 6 | | | 6 | | | 6 |
| Spawning Ground | WDFW 5 | | 3 | 2 | | | 5 |
| Test Fishery Net | ODFW 1 | | | 1 | | | 1 |
| Totals | 579 | | 113 | 432 | 34 | | 579 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 579 | 100 | 0.4883 |
| Total Survival | 579 | | 0.4883 |
| Escapement to NFH | 197 | 34 | 0.1661 |
| Columbia River contribution | 382 | 66 | 0.3222 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 382 | 197 |
| 0 CA | | |
| 0 Total | | |

939,000 117,914
 382
 - 3
 12
 367

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 06 Spring Chinook raised at Dworshak NFH
 released in Clearwater River

02/07/2013
 rd2
 06DwSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 04/02/2008 | 476,522 | 0.0 | 58,664 | 054132 | Dworshak NFH |
| 04/03/2008 | 462,478 | 0.0 | 59,250 | 054133 | Dworshak NFH |
| Totals | 939,000 | | 117,914 | | |

| Where Recovered | Total | 326 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|-----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 219 | | 211 | 8 | | 219 |
| Columbia River Sport | ODFW | 89 | 13 | 68 | 8 | | 89 |
| Fish Trap (freshwater) | ODFW | 3 | 3 | | | | 3 |
| Freshwater Sport | IDFG | 35 | 17 | 17 | 1 | | 35 |
| Freshwater Sport | ODFW | 3 | | 3 | | | 3 |
| Freshwater Sport | WDFW | 14 | | 10 | 4 | | 14 |
| Hatchery | FWS | 197 | 74 | 111 | 12 | | 197 |
| Hatchery | IDFG | 7 | 3 | 3 | 1 | | 7 |
| Spawning Ground | FWS | 6 | | 6 | | | 6 |
| Spawning Ground | WDFW | 5 | 3 | 2 | | | 5 |
| Test Fishery Net | ODFW | 1 | | 1 | | | 1 |
| Totals | | 579 | 113 | 432 | 34 | | 579 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 579 | 100 | 0.0617 |
| Total Survival | 579 | | 0.0617 |
| Escapement to NFH | 197 | 34 | 0.0210 |
| Columbia River contribution | 382 | 66 | 0.0407 |

Ocean

Contribution by Area

- 0 AK
- 0 BC
- 0 WA
- 0 OR
- 0 CA
- 0 Total

Columbia River
 382

Escapement to NFH
 197

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 07 Spring Chinook raised at Dworshak NFH
 released in Clearwater River

02/06/2013
 rd2
 07DwSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 03/26/2009 | 67,690 | 0.0 | 67,690 | 053572 | Dworshak NFH |
| 03/25/2009 | 65,226 | 0.0 | 65,226 | 053573 | Dworshak NFH |
| Totals | 132,916 | | 132,916 | | |

| Where Recovered | Total | 154 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 25 | | 2 | 23 | | | 25 |
| Columbia River Sport | 31 | | | 16 | 15 | | 31 |
| Fish Trap (freshwater) | 1 | | | 1 | | | 1 |
| Freshwater Sport | 23 | | | 22 | 1 | | 23 |
| Hatchery | 110 | | 14 | 96 | | | 110 |
| Hatchery | 3 | | | 3 | | | 3 |
| Hatchery | 1 | | | 1 | | | 1 |
| Spawning Ground | 4 | | | 4 | | | 4 |
| Totals | 198 | | 16 | 166 | 16 | | 198 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 198 | 100 | 0.1490 |
| Total Survival | 198 | | 0.1490 |

| | | | |
|-----------------------------|-----|----|--------|
| Escapement to NFH | 110 | 56 | 0.0828 |
| Columbia River contribution | 88 | 44 | 0.0662 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 88 | 110 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 07 Coho raised at Eagle Creek NFH
 released in Eagle Creek

12/14/2012
 rd2
 07ECCOS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------------|------|---------------|--------|-----------------|
| 04/10/2009 | 290,882 | 20.2 | 25,069 | 054182 | Eagle Creek NFH |
| 04/10/2009 | <u>34,062</u> | 20.7 | <u>24,925</u> | 054183 | Eagle Creek NFH |
| Totals | 324,944 | | <u>49,994</u> | | |

| Where Recovered | Total | 187 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|-----------|------------|----|----|------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 51 | 7 | | | | 7 |
| Columbia River Sport | ODFW | 36 | 3 | | | | 3 |
| Hatchery | FWS | 1,005 | 12 | 179 | | | 191 |
| Ocean Sport | ODFW | 108 | | 9 | | | 9 |
| Spawning Ground | ODFW | 24 | | 2 | | | 2 |
| Sport (charter) | WDFW | 12 | | 1 | | | 1 |
| Sport (private) | WDFW | <u>84</u> | | <u>7</u> | | | <u>7</u> |
| Totals | | <u>1,320</u> | <u>12</u> | <u>208</u> | | | <u>220</u> |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 204 | 15 | 0.0628 |
| Escapement to Columbia River | 1,116 | 85 | 0.3434 |
| Total Survival | 1,320 | | 0.4062 |
| Escapement to NFH | 1,005 | 76 | 0.3093 |
| Columbia River contribution | 111 | 8 | 0.0342 |

| Ocean | Contribution by Area | |
|-----------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 96 WA | Columbia River | Escapement to NFH |
| 108 OR | 111 | 1,005 |
| 0 CA | | |
| 204 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/14/2012
 Broodyear 08 Coho raised at Eagle Creek NFH rd2
 released in Eagle Creek 08ECCOS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------------|------|---------------|--------|-----------------|
| 04/12/2010 | 330,600 | 13.8 | 24,815 | 054371 | Eagle Creek NFH |
| 04/12/2010 | <u>25,012</u> | 14.1 | <u>24,825</u> | 054372 | Eagle Creek NFH |
| Totals | 355,612 | | <u>49,640</u> | | |

| Where Recovered | Total | 299 Observed Recoveries | | | | | Total |
|------------------------|--------------|---------------------------------|------------|----|----|----|------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 6 | | 6 | | | | 6 |
| Columbia River Sport | 78 | | 6 | | | | 6 |
| Estuary Sport | 39 | | 3 | | | | 3 |
| Hatchery | 1,418 | 3 | 287 | | | | 290 |
| Ocean Sport | 211 | | 19 | | | | 19 |
| Spawning Ground | 26 | | 2 | | | | 2 |
| Sport (charter) | 52 | | 4 | | | | 4 |
| Totals | <u>1,830</u> | <u>3</u> | <u>327</u> | | | | <u>330</u> |

| | | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 263 | 14 | 0.0740 |
| Escapement to Columbia River | 1,567 | 86 | 0.4406 |
| Total Survival | 1,830 | | 0.5146 |
| Escapement to NFH | 1,418 | 77 | 0.3987 |
| Columbia River contribution | 149 | 8 | 0.0419 |

| Ocean | Contribution by Area | |
|-----------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 52 WA | Columbia River | Escapement to NFH |
| 211 OR | 149 | 1,418 |
| 0 CA | | |
| 263 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 02 Spring Chinook raised at Kooskia NFH
 released in Clear Creek

02/07/2013
 rd2
 02KoSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|--------|--------|-------------|
| 03/30/2004 | 161,535 | 0.0 | 26,643 | 053735 | Kooskia NFH |
| 03/30/2004 | 429,845 | 0.0 | 70,897 | 054137 | Kooskia NFH |
| Totals | 591,380 | | 97,540 | | |

| Where Recovered | Total | 98 Observed Recoveries | | | | | Total |
|--------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 48 | | | | 8 | | 8 |
| Columbia River Sport | 174 | | | 25 | 4 | | 29 |
| Fish Trap (freshwater) | 12 | | | 2 | | | 2 |
| Freshwater Sport | 30 | | | 5 | | | 5 |
| Freshwater Sport | 18 | | | 3 | | | 3 |
| Hatchery | 444 | | 5 | 69 | | | 74 |
| Hatchery | 6 | | | 1 | | | 1 |
| Ocean Troll (non-treaty) | 12 | | | 2 | | | 2 |
| Spawning Ground | 6 | | | 1 | | | 1 |
| Treaty Ceremonial | 12 | | | 2 | | | 2 |
| Totals | 762 | | 5 | 110 | 12 | | 127 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 12 | 2 | 0.0020 |
| Escapement to Columbia River | 750 | 98 | 0.1268 |
| Total Survival | 762 | | 0.1289 |
| Escapement to NFH | 444 | 58 | 0.0751 |
| Columbia River contribution | 306 | 40 | 0.0517 |

Ocean

Contribution by Area

| | | |
|----------|----------------|-------------------|
| 0 AK | | |
| 12 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 306 | 444 |
| 0 CA | | |
| 12 Total | | |

Stock Assessment Reference Summary from CRIS and PSMFC TS1
 Broodyear 03 Spring Chinook raised at Kooskia NFH
 released in Clear Creek

02/07/2013
 rd2
 03KoSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|----------------|--------|-------------|
| 04/01/2005 | <u>616,385</u> | 0.0 | <u>100,895</u> | 051786 | Kooskia NFH |
| Totals | <u>616,385</u> | | <u>100,895</u> | | |

| Where Recovered | Total | 39 Observed Recoveries | | | | | Total |
|------------------------|----------|---------------------------------|----------|-----------|-----------|----|-----------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 102 | | | 2 | 15 | | 17 |
| Columbia River Sport | 84 | | | 11 | 3 | | 14 |
| Freshwater Sport | 18 | | | 2 | 1 | | 3 |
| Hatchery | 144 | | 1 | 19 | 4 | | 24 |
| Hatchery | <u>6</u> | | | <u>1</u> | | | <u>1</u> |
| Totals | 354 | | <u>1</u> | <u>35</u> | <u>23</u> | | <u>59</u> |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 354 | 100 | 0.0574 |
| Total Survival | 354 | | 0.0574 |
| Escapement to NFH | 144 | 41 | 0.0234 |
| Columbia River contribution | 210 | 59 | 0.0341 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 210 | 144 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 04 Spring Chinook raised at Kooskia NFH
 released in Clear Creek

02/07/2013
 rd2
 04KoSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|--------|--------|-------------|
| 04/03/2006 | 634,947 | 0.0 | 97,082 | 051784 | Kooskia NFH |
| Totals | 634,947 | | 97,082 | | |

| Where Recovered | Total | 296 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 700 | | | 100 | | | 100 |
| Columbia River Sport | 1,316 | | | 177 | 11 | | 188 |
| Freshwater Sport | 119 | | 1 | 16 | | | 17 |
| Freshwater Sport | 28 | | | 4 | | | 4 |
| Freshwater Sport | 63 | | 6 | 3 | | | 9 |
| Hatchery | 1,211 | | 47 | 124 | 2 | | 173 |
| Hatchery | 14 | | 2 | | | | 2 |
| Spawning Ground | 84 | | 1 | 11 | | | 12 |
| Treaty Ceremonial | 7 | | | 1 | | | 1 |
| Totals | 3,542 | | 57 | 436 | 13 | | 506 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 3,542 | 100 | 0.5578 |
| Total Survival | 3,542 | | 0.5578 |

| | Total | % of total | % of release |
|-----------------------------|-------|------------|--------------|
| Escapement to NFH | 1,211 | 34 | 0.1907 |
| Columbia River contribution | 2,331 | 66 | 0.3671 |

Ocean

Contribution by Area

0 AK
 0 BC
 0 WA
 0 OR
 0 CA
 0 Total

Columbia River
 2,331

Escapement to NFH
 1,211

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 05 Spring Chinook raised at Kooskia NFH
 released in Clear Creek

02/07/2013
 rd2
 05KoSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|-------------|
| 04/02/2007 | 4,525 | 0.0 | 0 | - | Kooskia NFH |
| 04/02/2007 | 110,967 | 0.0 | 107,452 | 051785 | Kooskia NFH |
| 04/02/2007 | 42,474 | 0.0 | 40,420 | 102480 | Kooskia NFH |
| 04/02/2007 | 20,777 | 0.0 | 19,787 | 107571 | Kooskia NFH |
| Totals | 178,743 | | 167,659 | | |

155 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|------------------------|------|-------|----|----|-----|----|----|-------|
| Columbia River Gillnet | ODFW | 13 | | | 13 | | | 13 |
| Columbia River Sport | ODFW | 62 | | | 62 | | | 62 |
| Freshwater Sport | IDFG | 12 | | 2 | 10 | | | 12 |
| Hatchery | FWS | 112 | | 18 | 94 | | | 112 |
| Hatchery | IDFG | 1 | | | | 1 | | 1 |
| Treaty Subsistence | FWS | 2 | | | | 2 | | 2 |
| Totals | | 202 | | 20 | 179 | 3 | | 202 |

| | | % of total | % of release |
|------------------------------|-----|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 202 | 100 | 0.1130 |
| Total Survival | 202 | | 0.1130 |
| Escapement to NFH | 112 | 55 | 0.0627 |
| Columbia River contribution | 90 | 45 | 0.0504 |

Ocean

Contribution by Area

- 0 AK
- 0 BC
- 0 WA
- 0 OR
- 0 CA
- 0 Total

Columbia River
 90

Escapement to NFH
 112

Stock Assessment Reference Summary from CRIS and PSMFC TS1
 Broodyear 06 Spring Chinook raised at Kooskia NFH
 released in Clear Creek

02/07/2013
 rd2
 06KoSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|-------------|
| 04/07/2008 | 112,344 | 0.0 | 107,945 | 053575 | Kooskia NFH |
| Totals | 112,344 | | 107,945 | | |

| Where Recovered | Total | 247 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 137 | | | 137 | | | 137 |
| Columbia River Sport | 119 | | | 119 | | | 119 |
| Freshwater Sport | 26 | | 3 | 23 | | | 26 |
| Freshwater Sport | 2 | | | 2 | | | 2 |
| Freshwater Sport | 6 | | | 6 | | | 6 |
| Hatchery | 127 | | 16 | 111 | | | 127 |
| Spawning Ground | 3 | | 3 | | | | 3 |
| Test Fishery Net | 2 | | | 2 | | | 2 |
| Treaty Subsistence | 17 | | | 17 | | | 17 |
| Totals | 439 | | 22 | 417 | | | 439 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 439 | 100 | 0.3908 |
| Total Survival | 439 | | 0.3908 |
| Escapement to NFH | 127 | 29 | 0.1130 |
| Columbia River contribution | 312 | 71 | 0.2777 |

Ocean

Contribution by Area

- 0 AK
- 0 BC
- 0 WA
- 0 OR
- 0 CA
- 0 Total

Columbia River
 312

Escapement to NFH
 127

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/28/2012
 Broodyear 07 Spring Chinook raised at Leavenworth NFH rd2
 released in Icicle Creek 07LeSCS

| | | | | | |
|------------|-----------|------|---------|--------|-----------------|
| Date | Number | #/lb | Tagged | CWT | AT |
| 04/28/2009 | 1,685,038 | 18.3 | 196,529 | 054195 | Leavenworth NFH |
| Totals | 1,685,038 | | 196,529 | | |

| | | | | | | | | | |
|------------------------|------|--|-------|---------------------------------|----|-----|----|----|-------|
| | | | | 261 Observed Recoveries | | | | | |
| | | | | === PSC Expanded Recoveries === | | | | | |
| Where Recovered | | | Total | 2s | 3s | 4s | 5s | 6s | Total |
| Columbia River Gillnet | ODFW | | 234 | | | 26 | | | 26 |
| Columbia River Sport | ODFW | | 153 | | | 17 | | | 17 |
| Freshwater Sport | ODFW | | 9 | | | 1 | | | 1 |
| Freshwater Sport | WDFW | | 711 | | | 79 | | | 79 |
| Hatchery | FWS | | 2,196 | | 12 | 232 | | | 244 |
| Hatchery | WDFW | | 0 | | | | | | 0 |
| Spawning Ground | WDFW | | 81 | | | 9 | | | 9 |
| Totals | | | 3,384 | | 12 | 364 | | | 376 |

| | | | | | |
|------------------------------|--|--|-------|-------|---------|
| | | | | % of | % of |
| | | | | total | release |
| Ocean contribution | | | 0 | 0 | 0.0000 |
| Escapement to Columbia River | | | 3,384 | 100 | 0.2008 |
| Total Survival | | | 3,384 | | 0.2008 |

| | | | | | |
|-----------------------------|--|--|-------|----|--------|
| Escapement to NFH | | | 2,196 | 65 | 0.1303 |
| Columbia River contribution | | | 1,188 | 35 | 0.0705 |

| | | |
|---------|----------------------|-------------------|
| Ocean | Contribution by Area | |
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 1,188 | 2,196 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 raised at Leavenworth NFH released in Icicle Creek
 Broodyear 06 Spring Chinook
 12/28/2012 rd2 06LeSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|-----------|------|---------|--------|-----------------|
| 04/28/2008 | 741,443 | 18.3 | 199,442 | 052573 | Leavenworth NFH |
| 04/28/2008 | 798,225 | 18.5 | 198,658 | 053784 | Leavenworth NFH |
| Totals | 1,539,668 | | 398,100 | | |

2,205 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|--------------------------|------|--------|----|-----|-------|-----|----|-------|
| Columbia River Gillnet | ODFW | 2,184 | | 2 | 526 | 18 | | 546 |
| Columbia River Sport | ODFW | 2,552 | | 11 | 470 | 157 | | 638 |
| Freshwater Sport | ODFW | 60 | | | 10 | 5 | | 15 |
| Freshwater Sport | WDFW | 1,144 | | 64 | 165 | 57 | | 286 |
| Hatchery | FWS | 7,556 | | 253 | 1,368 | 268 | | 1,889 |
| Hatchery | WDFW | 4 | | | 1 | | | 1 |
| Ocean Sport | CDFO | 12 | | | 3 | | | 3 |
| Ocean Troll (non-treaty) | ADFG | 24 | | 6 | | | | 6 |
| Spawning Ground | WDFW | 300 | | 17 | 32 | 26 | | 75 |
| Test Fishery Net | ODFW | 16 | | 1 | 3 | | | 4 |
| Totals | | 13,852 | | 354 | 2,578 | 531 | | 3,463 |

| | | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 36 | 0 | 0.0023 |
| Escapement to Columbia River | 13,816 | 100 | 0.8973 |
| Total Survival | 13,852 | | 0.8997 |
| Escapement to NFH | 7,556 | 55 | 0.4908 |
| Columbia River contribution | 6,260 | 45 | 0.4066 |

| Ocean | Contribution by Area | |
|----------|----------------------|-------------------|
| 24 AK | | |
| 12 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 6,260 | 7,556 |
| 0 CA | | |
| 36 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 06 Spring Chinook raised at L White Salmon NFH rd2
 released in L White Salmon River 06LWSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|---------------|--------|--------------------|
| 04/17/2008 | 373,607 | 14.8 | 44,178 | 052965 | L White Salmon NFH |
| 04/17/2008 | <u>560,831</u> | 14.8 | <u>69,586</u> | 052975 | L White Salmon NFH |
| Totals | 934,438 | | 113,764 | | |

1,135 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|------------------------|------|---------------|----|------------|--------------|-----------|----|--------------|
| Columbia River Gillnet | ODFW | 6,480 | | 28 | 762 | 20 | | 810 |
| Columbia River Sport | ODFW | 2,584 | | 11 | 289 | 23 | | 323 |
| Freshwater Net | WDFW | 1,352 | | 60 | 104 | 5 | | 169 |
| Freshwater Sport | WDFW | 7,208 | | 321 | 580 | | | 901 |
| Hatchery | FWS | 4,408 | | 50 | 484 | 17 | | 551 |
| Spawning Ground | WDFW | 240 | | 6 | 4 | 20 | | 30 |
| Test Fishery Net | ODFW | 48 | | 1 | 5 | | | 6 |
| Totals | | <u>22,320</u> | | <u>477</u> | <u>2,228</u> | <u>85</u> | | <u>2,790</u> |

| | Total | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 22,320 | 100 | 2.3886 |
| Total Survival | 22,320 | | 2.3886 |
| Escapement to NFH | 4,408 | 20 | 0.4717 |
| Columbia River contribution | 17,912 | 80 | 1.9169 |

Ocean

Contribution by Area

| | | |
|---------|----------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 17,912 | 4,408 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 07 Spring Chinook raised at L White Salmon NFH rd2
 released in L White Salmon River 07LWSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|------|----------|--------|--------------------|
| 04/16/2009 | 590,763 | 15.0 | 71,221 | 053769 | L White Salmon NFH |
| 04/16/2009 | 424,569 | 16.5 | 75,394 | 054180 | Willard NFH |
| 04/16/2009 | <u>424,569</u> | 16.5 | <u>0</u> | 054180 | Willard NFH |
| Totals | 1,439,901 | | 146,615 | | |

| Where Recovered | Total | 217 Observed Recoveries | | | | | Total |
|------------------------|-------|---------------------------------|----------|-----------|----|----|-----------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | ODFW | 288 | | 36 | | | 36 |
| Columbia River Sport | ODFW | 72 | | 9 | | | 9 |
| Freshwater Net | WDFW | 344 | | 43 | | | 43 |
| Freshwater Sport | ODFW | 16 | | 2 | | | 2 |
| Freshwater Sport | WDFW | 872 | | 109 | | | 109 |
| Hatchery | FWS | 944 | 1 | 117 | | | 118 |
| Hatchery | WDFW | 0 | | | | | 0 |
| Spawning Ground | WDFW | <u>712</u> | | <u>89</u> | | | <u>89</u> |
| Totals | | 3,248 | <u>1</u> | 405 | | | 406 |

| | | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 3,248 | 100 | 0.2256 |
| Total Survival | 3,248 | | 0.2256 |
| Escapement to NFH | 944 | 29 | 0.0656 |
| Columbia River contribution | 2,304 | 71 | 0.1600 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 2,304 | 944 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 04 Upriver Bright FCS raised at L White Salmon NFH rd2
 released in L White Salmon River 04LWURB

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|----------------|-------|----------------|------------|--------------------|
| 06/29/2005 | 442,868 | 122.0 | 0 | - | Willard NFH |
| 06/23/2005 | 1,259,477 | 81.3 | 179,130 | 0501040509 | L White Salmon NFH |
| 06/23/2005 | <u>200,396</u> | 68.6 | <u>193,212</u> | 052581 | L White Salmon NFH |
| Totals | 1,902,741 | | 372,342 | | |

| Where Recovered | Total | 552 Observed Recoveries | | | | | Total |
|--------------------------|--------------|---------------------------------|-----------|------------|------------|----------|------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 280 | | 7 | 29 | 4 | | 40 |
| Commercial Seine | 70 | | | | 10 | | 10 |
| Estuary Sport | 14 | | | 2 | | | 2 |
| Freshwater Net | 29 | | | 4 | 1 | | 5 |
| Hatchery | 1,774 | 20 | 77 | 268 | 78 | 5 | 448 |
| Multiple Gear | 7 | | | 1 | | | 1 |
| Ocean Sport | 21 | | | | 3 | | 3 |
| Ocean Sport | 147 | | | 14 | 7 | | 21 |
| Ocean Troll (non-treaty) | 717 | | | 59 | 46 | | 105 |
| Ocean Troll (non-treaty) | 199 | | 8 | 32 | 21 | | 61 |
| Ocean Troll (non-treaty) | 2 | | 2 | | | | 2 |
| Spawning Ground | 70 | 3 | | 5 | 2 | | 10 |
| Sport (private) | 3 | | | 3 | | | 3 |
| Treaty Troll | 19 | | | 2 | | | 7 |
| Totals | <u>3,352</u> | <u>23</u> | <u>99</u> | <u>419</u> | <u>172</u> | <u>5</u> | <u>718</u> |

| | | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 1,185 | 35 | 0.0623 |
| Escapement to Columbia River | 2,167 | 65 | 0.1139 |
| Total Survival | 3,352 | | 0.1762 |
| Escapement to NFH | 1,774 | 53 | 0.0932 |
| Columbia River contribution | 393 | 12 | 0.0207 |

| Ocean | Contribution by Area | |
|-------------|----------------------|-------------------|
| 815 AK | | |
| 346 BC | | |
| 24 WA | Columbia River | Escapement to NFH |
| 0 OR | 393 | 1,774 |
| 0 CA | | |
| 1,185 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 05 Upriver Bright FCS raised at L White Salmon NFH rd2
 released in L White Salmon River 05LWURB

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|-----------|------|---------|--------|--------------------|
| 06/22/2006 | 219,610 | 82.2 | 219,169 | 052875 | L White Salmon NFH |
| 06/22/2006 | 1,582,564 | 82.2 | 228,976 | 052876 | L White Salmon NFH |
| Totals | 1,802,174 | | 448,145 | | |

1,305 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|--------------------------|------|--------|----|-----|-----|-------|----|-------|
| Columbia River Gillnet | ODFW | 2,210 | | 36 | 194 | 87 | 3 | 320 |
| Columbia River Sport | ODFW | 322 | | 13 | 13 | 20 | | 46 |
| Commercial Seine | ADFG | 42 | | | | 6 | | 6 |
| Estuary Sport | ODFW | 70 | | 3 | | 7 | | 10 |
| Freshwater Net | WDFW | 247 | | 14 | 23 | | | 37 |
| Freshwater Sport | WDFW | 7 | | | | 1 | | 1 |
| Groundfish Observer (Ber | NMFS | 14 | | | | 2 | | 2 |
| Hatchery | FWS | 6,903 | 29 | 282 | 369 | 1,197 | 10 | 1,887 |
| Hatchery | ODFW | 9 | | | 3 | | | 3 |
| Mixed Net and Seine | CDFO | 1 | 1 | | | | | 1 |
| Mixed Net and Seine | WDFW | 1 | | 1 | | | | 1 |
| Multiple Gear | ADFG | 7 | | 1 | | | | 1 |
| Ocean Sport | ADFG | 112 | | | 8 | 7 | 1 | 16 |
| Ocean Sport | CDFO | 602 | | 15 | 35 | 35 | 1 | 86 |
| Ocean Sport | ODFW | 28 | | | | 4 | | 4 |
| Ocean Troll (non-treaty) | ADFG | 1,414 | | 8 | 116 | 78 | | 202 |
| Ocean Troll (non-treaty) | CDFO | 800 | | 30 | 145 | 30 | 1 | 206 |
| Ocean Troll (non-treaty) | WDFW | 43 | | | 6 | 1 | | 7 |
| Ocean Troll - Day Boat | ADFG | 77 | | | 11 | | | 11 |
| Spawning Ground | WDFW | 174 | 2 | 11 | 13 | 4 | | 30 |
| Sport (private) | WDFW | 67 | | 7 | 6 | | | 13 |
| Treaty Troll | WDFW | 26 | | 7 | 1 | | | 8 |
| Totals | | 13,176 | 32 | 428 | 943 | 1,479 | 16 | 2,898 |

| | | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 3,234 | 25 | 0.1794 |
| Escapement to Columbia River | 9,942 | 75 | 0.5517 |
| Total Survival | 13,176 | | 0.7311 |
| Escapement to NFH | 6,903 | 52 | 0.3830 |
| Columbia River contribution | 3,039 | 23 | 0.1686 |

| Ocean | Contribution by Area | |
|-------------|----------------------|-------------------|
| 1,652 AK | | |
| 1,403 BC | | |
| 137 WA | Columbia River | Escapement to NFH |
| 28 OR | 3,039 | 6,903 |
| 0 CA | | |
| 3,220 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 06 Upriver Bright FCS raised at L White Salmon NFH rd2
 released in L White Salmon River 06LWURB

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|-----------|------|---------|--------|--------------------|
| 06/28/2007 | 931,722 | 77.1 | 97,716 | 053886 | L White Salmon NFH |
| 06/28/2007 | 931,722 | 77.1 | 98,815 | 053887 | L White Salmon NFH |
| 06/28/2007 | 98,654 | 77.1 | 97,667 | 053888 | L White Salmon NFH |
| 06/28/2007 | 99,826 | 77.1 | 98,828 | 053889 | L White Salmon NFH |
| Totals | 2,061,924 | | 393,026 | | |

| Where Recovered | Total | 708 Observed Recoveries | | | | | Total |
|-------------------------------|--------|---------------------------------|-----|-------|-----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet ODFW | 1,881 | | 26 | 120 | 55 | | 201 |
| Columbia River Sport ODFW | 150 | | 6 | 4 | 6 | | 16 |
| Commercial Seine ADFG | 10 | | 1 | | | | 1 |
| Estuary Sport ODFW | 36 | | 4 | | | | 4 |
| Freshwater Net WDFW | 75 | 1 | 7 | | | | 8 |
| Freshwater Sport WDFW | 63 | | 7 | | | | 7 |
| Hake Trawl Fishery, Shor NMFS | 9 | 1 | | | | | 1 |
| Hatchery FWS | 6,793 | 131 | 117 | 984 | 129 | | 1,361 |
| Hatchery ODFW | 75 | | 2 | 3 | 5 | | 10 |
| Ocean Sport ADFG | 104 | | | 6 | 5 | | 11 |
| Ocean Sport CDFO | 284 | | 16 | 11 | 3 | | 30 |
| Ocean Troll (non-treaty) ADFG | 630 | | 3 | 39 | 24 | | 66 |
| Ocean Troll (non-treaty) CDFO | 584 | | 15 | 88 | 20 | | 123 |
| Ocean Troll (non-treaty) WDFW | 29 | | | 3 | | | 3 |
| Spawning Ground WDFW | 613 | 3 | 6 | 37 | 72 | | 118 |
| Sport (charter) WDFW | 3 | | | | 3 | | 3 |
| Sport (private) WDFW | 55 | | 6 | 1 | | | 7 |
| Treaty Troll WDFW | 39 | | 3 | 2 | | | 5 |
| Totals | 11,433 | 136 | 219 | 1,298 | 322 | | 1,975 |

| | | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 1,747 | 15 | 0.0847 |
| Escapement to Columbia River | 9,686 | 85 | 0.4698 |
| Total Survival | 11,433 | | 0.5545 |
| Escapement to NFH | 6,793 | 59 | 0.3294 |
| Columbia River contribution | 2,893 | 25 | 0.1403 |

| Ocean | Contribution by Area | |
|-------------|----------------------|-------------------|
| 744 AK | | |
| 868 BC | | |
| 126 WA | Columbia River | Escapement to NFH |
| 0 OR | 2,893 | 6,793 |
| 0 CA | | |
| 1,738 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 12/31/2012
 Broodyear 07 Upriver Bright FCS raised at L White Salmon NFH rd2
 released in L White Salmon River 07LWURB

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|-----------|------|---------|--------|--------------------|
| 07/03/2008 | 901,117 | 96.0 | 99,711 | 054191 | L White Salmon NFH |
| 07/03/2008 | 900,903 | 96.0 | 99,954 | 054192 | L White Salmon NFH |
| 07/03/2008 | 99,763 | 96.0 | 99,763 | 054193 | L White Salmon NFH |
| 07/03/2008 | 99,977 | 96.0 | 99,977 | 054194 | L White Salmon NFH |
| Totals | 2,001,760 | | 399,405 | | |

1,334 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|--------------------------|------|--------|-----|-------|-------|----|----|-------|
| Columbia River Gillnet | ODFW | 3,771 | 9 | 76 | 334 | | | 419 |
| Columbia River Sport | ODFW | 468 | | 12 | 40 | | | 52 |
| Commercial Seine | ADFG | 9 | 1 | | | | | 1 |
| Estuary Sport | ODFW | 30 | | | 6 | | | 6 |
| Freshwater Net | WDFW | 18 | 2 | | | | | 2 |
| Groundfish Observer (Gul | NMFS | 54 | | 6 | | | | 6 |
| Hatchery | FWS | 13,188 | 149 | 1,352 | 1,087 | | | 2,588 |
| Hatchery | ODFW | 112 | 2 | 6 | 8 | | | 16 |
| Ocean Sport | ADFG | 207 | | 3 | 17 | 3 | | 23 |
| Ocean Sport | CDFO | 333 | | 12 | 25 | | | 37 |
| Ocean Sport | ODFW | 21 | | 2 | 3 | | | 5 |
| Ocean Troll (non-treaty) | ADFG | 2,264 | | 10 | 169 | 77 | | 256 |
| Ocean Troll (non-treaty) | CDFG | 18 | | | 2 | | | 2 |
| Ocean Troll (non-treaty) | CDFO | 614 | | 11 | 115 | | | 126 |
| Ocean Troll (non-treaty) | ODFW | 8 | | | 8 | | | 8 |
| Ocean Troll (non-treaty) | WDFW | 10 | | | 2 | | | 2 |
| Ocean Troll - Day Boat | ADFG | 81 | | | 6 | 3 | | 9 |
| Spawning Ground | WDFW | 606 | 3 | 18 | 121 | | | 142 |
| Sport (private) | WDFW | 78 | | 6 | 8 | | | 14 |
| Treaty Troll | WDFW | 256 | | 26 | 14 | | | 40 |
| Totals | | 22,146 | 166 | 1,540 | 1,965 | 83 | | 3,754 |

| | | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 3,953 | 18 | 0.1975 |
| Escapement to Columbia River | 18,193 | 82 | 0.9089 |
| Total Survival | 22,146 | | 1.1063 |
| Escapement to NFH | 13,188 | 60 | 0.6588 |
| Columbia River contribution | 5,005 | 23 | 0.2500 |

| Ocean | Contribution by Area | |
|-------------|----------------------|-------------------|
| 2,561 AK | | |
| 947 BC | | |
| 344 WA | Columbia River | Escapement to NFH |
| 29 OR | 5,005 | 13,188 |
| 18 CA | | |
| 3,899 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 05 Tule Fall Chinook raised at Spring Creek NFH
 released in Columbia River

02/01/2013
 rd2
 05SCFCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|------------|-------|---------|--------|------------------|
| 05/05/2006 | 149,582 | 149.5 | 148,553 | 052871 | Spring Creek NFH |
| 04/17/2006 | 149,243 | 131.6 | 148,657 | 052872 | Spring Creek NFH |
| 05/05/2006 | 3,271,426 | 133.6 | 147,338 | 052873 | Spring Creek NFH |
| 04/17/2006 | 4,077,775 | 127.8 | 147,870 | 052874 | Spring Creek NFH |
| 03/02/2006 | 74,653 | 171.2 | 74,504 | 052969 | Spring Creek NFH |
| 03/02/2006 | 74,848 | 171.2 | 74,702 | 052970 | Spring Creek NFH |
| 03/02/2006 | 3,587,014 | 171.2 | 73,383 | 052971 | Spring Creek NFH |
| 03/02/2006 | 3,854,513 | 171.2 | 74,317 | 052972 | Spring Creek NFH |
| Totals | 15,239,054 | | 889,324 | | |

1,953 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|--------------------------|------|--------|-----|-------|-----|----|----|-------|
| Columbia River Gillnet | ODFW | 38,229 | 225 | 891 | 171 | | | 1,287 |
| Columbia River Sport | ODFW | 364 | | 7 | | | | 7 |
| Estuary Sport | ODFW | 1,587 | 11 | 34 | | | | 45 |
| Estuary Sport | WDFW | 1,320 | 12 | 30 | 3 | | | 45 |
| Hake Trawl Fishery, At-S | NMFS | 65 | 65 | | | | | 65 |
| Hatchery | FWS | 36,278 | 508 | 1,378 | 387 | | | 2,273 |
| Hatchery | ODFW | 1,574 | 35 | 22 | 8 | | | 65 |
| Ocean Sport | CDFO | 5,352 | 10 | 150 | 11 | | | 171 |
| Ocean Sport | ODFW | 272 | 11 | | | | | 11 |
| Ocean Trawl By-Catch | ODFW | 44 | 2 | | | | | 2 |
| Ocean Trawl By-Catch | WDFW | 197 | 11 | | | | | 11 |
| Ocean Troll (non-treaty) | CDFO | 4,037 | 6 | 266 | 10 | 5 | | 287 |
| Ocean Troll (non-treaty) | ODFW | 1,047 | | 32 | 1 | | | 33 |
| Ocean Troll (non-treaty) | WDFW | 2,589 | 2 | 148 | 6 | | | 156 |
| Spawning Ground | WDFW | 140 | | | 5 | | | 5 |
| Sport (charter) | WDFW | 1,091 | 16 | 54 | 4 | | | 74 |
| Sport (private) | WDFW | 3,229 | 46 | 110 | 13 | | | 169 |
| Treaty Troll | WDFW | 1,437 | 17 | 60 | 4 | | | 81 |
| Totals | | 98,852 | 977 | 3,182 | 623 | 5 | | 4,787 |

% of total % of release

| | | | |
|------------------------------|--------|----|--------|
| Ocean contribution | 19,360 | 20 | 0.1270 |
| Escapement to Columbia River | 79,492 | 80 | 0.5216 |
| Total Survival | 98,852 | | 0.6487 |

| | | | |
|-----------------------------|--------|----|--------|
| Escapement to NFH | 36,278 | 37 | 0.2381 |
| Columbia River contribution | 43,214 | 44 | 0.2836 |

Ocean Contribution by Area

| | | |
|--------------|----------------|-------------------|
| 0 AK | | |
| 9,389 BC | | |
| 8,543 WA | | |
| 1,363 OR | Columbia River | Escapement to NFH |
| 0 CA | 43,214 | 36,278 |
| 19,295 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 06 Tule Fall Chinook raised at Spring Creek NFH
 released in Columbia River

02/01/2013
 rd2
 06SCFCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|------------|-------|---------|--------|------------------|
| 03/09/2007 | 368,263 | 128.1 | 25,097 | 052570 | Spring Creek NFH |
| 04/12/2007 | 4,110,491 | 88.6 | 99,904 | 052577 | Spring Creek NFH |
| 03/09/2007 | 25,062 | 128.1 | 24,963 | 052584 | Spring Creek NFH |
| 03/05/2007 | 361,272 | 149.2 | 25,080 | 052588 | Spring Creek NFH |
| 03/05/2007 | 25,109 | 149.2 | 24,961 | 052589 | Spring Creek NFH |
| 03/09/2007 | 331,936 | 128.1 | 49,701 | 052895 | Spring Creek NFH |
| 03/09/2007 | 50,694 | 128.1 | 49,813 | 052896 | Spring Creek NFH |
| 03/05/2007 | 5,771,649 | 149.2 | 49,657 | 052897 | Spring Creek NFH |
| 03/05/2007 | 50,206 | 149.2 | 49,622 | 052898 | Spring Creek NFH |
| 04/12/2007 | 100,082 | 88.6 | 99,889 | 053484 | Spring Creek NFH |
| 05/01/2007 | 98,478 | 74.2 | 97,493 | 053485 | Spring Creek NFH |
| 05/01/2007 | 3,396,257 | 74.2 | 97,493 | 053592 | Spring Creek NFH |
| 03/09/2007 | 342,545 | 128.1 | 50,089 | 054318 | Spring Creek NFH |
| 03/09/2007 | 50,310 | 128.1 | 49,838 | 054334 | Spring Creek NFH |
| 03/05/2007 | 50,263 | 149.2 | 49,662 | 054335 | Spring Creek NFH |
| 03/05/2007 | 339,944 | 149.3 | 49,356 | 054336 | Spring Creek NFH |
| Totals | 15,472,561 | | 892,618 | | |

367 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|--------------------------|------|--------|----|-----|----|----|----|-------|
| Columbia River Gillnet | ODFW | 11,114 | 16 | 198 | 40 | | | 254 |
| Columbia River Sport | ODFW | 626 | | 16 | | | | 16 |
| Estuary Sport | WDFW | 275 | | 15 | | | | 15 |
| Hatchery | FWS | 9,878 | 21 | 518 | 5 | | | 544 |
| Hatchery | ODFW | 496 | | 24 | 3 | | | 27 |
| Mixed Net and Seine | WDFW | 42 | 2 | | | | | 2 |
| Ocean Sport | CDFO | 1,445 | | 24 | 9 | | | 33 |
| Ocean Troll (non-treaty) | CDFO | 1,491 | 18 | 56 | 12 | | | 86 |
| Ocean Troll (non-treaty) | ODFW | 348 | | | 3 | | | 3 |
| Ocean Troll (non-treaty) | WDFW | 635 | | 31 | 1 | | | 32 |
| Sport (charter) | WDFW | 2 | 2 | | | | | 2 |
| Sport (private) | WDFW | 435 | 4 | 17 | 4 | | | 25 |
| Treaty Troll | WDFW | 17 | 9 | 8 | | | | 17 |
| Totals | | 26,804 | 72 | 907 | 77 | | | 1,056 |

| | Total | % of total | % of release |
|------------------------------|--------|------------|--------------|
| Ocean contribution | 4,415 | 16 | 0.0285 |
| Escapement to Columbia River | 22,389 | 84 | 0.1447 |
| Total Survival | 26,804 | | 0.1732 |

| | | | |
|-----------------------------|--------|----|--------|
| Escapement to NFH | 9,878 | 37 | 0.0638 |
| Columbia River contribution | 12,511 | 47 | 0.0809 |

Ocean Contribution by Area

| | | |
|-------------|----------------|-------------------|
| 0 AK | | |
| 2,936 BC | | |
| 1,131 WA | Columbia River | Escapement to NFH |
| 348 OR | 12,511 | 9,878 |
| 0 CA | | |
| 4,415 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 07 Tule Fall Chinook raised at Spring Creek NFH
 released in Columbia River

02/04/2013
 rd2
 07SCFCS

14,899,270 released with 891,550 tagged fish and 20 CWTs

| Where Recovered | Total | 1,713 Observed Recoveries | | | | | Total |
|--------------------------|-------|---------------------------------|-------|-------|-----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Coastal Gillnet | WDFW | 330 | | 5 | 5 | | 10 |
| Columbia River Gillnet | ODFW | 64,708 | 537 | 1,334 | 97 | | 1,968 |
| Columbia River Sport | ODFW | 565 | 5 | 12 | | | 17 |
| Estuary Sport | ODFW | 1,245 | 10 | 28 | 3 | | 41 |
| Estuary Sport | WDFW | 1,950 | 61 | 27 | 3 | | 91 |
| Hake Trawl Fishery, Shor | NMFS | 1 | | 1 | | | 1 |
| Hatchery | FWS | 19,455 | 1,186 | | 92 | | 1,278 |
| Hatchery | ODFW | 1,446 | 53 | 43 | | | 96 |
| Mixed Net and Seine | WDFW | 72 | | 3 | | | 3 |
| Ocean Sport | CDFG | 112 | | 4 | | | 4 |
| Ocean Sport | CDFO | 4,139 | 38 | 134 | 17 | | 189 |
| Ocean Sport | ODFW | 627 | 6 | 18 | | | 24 |
| Ocean Sport | WDFW | 56 | 1 | | | | 1 |
| Ocean Troll (non-treaty) | CDFO | 7,481 | | 385 | 32 | | 417 |
| Ocean Troll (non-treaty) | ODFW | 8,008 | | 248 | 4 | | 252 |
| Ocean Troll (non-treaty) | WDFW | 10,258 | | 567 | 3 | | 570 |
| Spawning Ground | WDFW | 520 | 8 | 12 | | | 20 |
| Sport (charter) | WDFW | 3,679 | 36 | 129 | 3 | | 168 |
| Sport (private) | WDFW | 5,226 | 118 | 184 | 2 | | 304 |
| Treaty Troll | WDFW | 1,307 | 12 | 70 | | | 82 |
| Totals | | 131,185 | 2,071 | 3,204 | 261 | | 5,536 |

| | | % of total | % of release |
|------------------------------|---------|------------|--------------|
| Ocean contribution | 41,296 | 31 | 0.2772 |
| Escapement to Columbia River | 89,889 | 69 | 0.6033 |
| Total Survival | 131,185 | | 0.8805 |
| Escapement to NFH | 19,455 | 15 | 0.1306 |
| Columbia River contribution | 70,434 | 54 | 0.4727 |

| Ocean | Contribution by Area | |
|--------------|----------------------|-------------------|
| 0 AK | | |
| 11,620 BC | | |
| 20,928 WA | Columbia River | Escapement to NFH |
| 8,635 OR | 70,434 | 19,455 |
| 112 CA | | |
| 41,295 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 02/01/2013
 Broodyear 05 Spring Chinook raised at Warm Springs NFH rd2
 released in Warm Springs River 05WSSCS

620,753 released with 582,771 tagged fish and 27 CWTs

| Where Recovered | Total | 1,140 Observed Recoveries | | | | | Total |
|-------------------------------|--------------|---------------------------------|------------|--------------|-----------|----|--------------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet ODFW | 120 | | | 120 | | | 120 |
| Columbia River Sport ODFW | 367 | | 10 | 353 | 4 | | 367 |
| Fish Trap (freshwater) ODFW | 2 | | | 2 | | | 2 |
| Freshwater Sport ODFW | 5 | | 4 | | 1 | | 5 |
| Hatchery FWS | 2,138 | | 162 | 1,958 | 18 | | 2,138 |
| Hatchery ODFW | 4 | | 3 | 1 | | | 4 |
| Ocean Troll (non-treaty) ADFG | 3 | | | 3 | | | 3 |
| Test Fishery Net ODFW | 9 | | | 9 | | | 9 |
| Totals | 2,648 | | 179 | 2,446 | 23 | | 2,648 |

| | | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 3 | 0 | 0.0005 |
| Escapement to Columbia River | 2,645 | 100 | 0.4261 |
| Total Survival | 2,648 | | 0.4266 |
| Escapement to NFH | 2,138 | 81 | 0.3444 |
| Columbia River contribution | 507 | 19 | 0.0817 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 3 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 507 | 2,138 |
| 0 CA | | |
| 3 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1 02/01/2013
 Broodyear 07 Spring Chinook raised at Warm Springs NFH rd2
 released in Warm Springs River 07WSSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|------------------|
| 04/14/2009 | 125,296 | 33.7 | 123,896 | 053479 | Warm Springs NFH |
| 04/20/2009 | 59,797 | 27.3 | 59,797 | 053770 | Warm Springs NFH |
| 04/20/2009 | 60,027 | 30.8 | 59,912 | 053771 | Warm Springs NFH |
| 04/20/2009 | 78,049 | 29.8 | 77,582 | 053772 | Warm Springs NFH |
| 04/20/2009 | 88,460 | 30.1 | 87,944 | 053869 | Warm Springs NFH |
| 04/20/2009 | 83,034 | 30.2 | 82,262 | 053870 | Warm Springs NFH |
| 04/14/2009 | 86,224 | 27.6 | 85,896 | 053885 | Warm Springs NFH |
| Totals | 580,887 | | 577,289 | | |

| Where Recovered | Total | 1,544 Observed Recoveries | | | | | Total |
|--------------------------|-------|---------------------------------|-----|-------|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet | 256 | | 9 | 238 | 9 | | 256 |
| Columbia River Sport | 90 | | 5 | 82 | 3 | | 90 |
| Fish Trap (freshwater) | 2 | | | 2 | | | 2 |
| Freshwater Sport | 891 | | 12 | 878 | 1 | | 891 |
| Freshwater Sport | 16 | | | 16 | | | 16 |
| Hatchery | 2,902 | | 167 | 2,735 | | | 2,902 |
| Hatchery | 1 | | | 1 | | | 1 |
| Ocean Troll (non-treaty) | 1 | | | 1 | | | 1 |
| Spawning Ground | 1 | | | 1 | | | 1 |
| Test Fishery Net | 4 | | | 4 | | | 4 |
| Totals | 4,164 | | 193 | 3,958 | 13 | | 4,164 |

| | | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 1 | 0 | 0.0002 |
| Escapement to Columbia River | 4,163 | 100 | 0.7167 |
| Total Survival | 4,164 | | 0.7168 |
| Escapement to NFH | 2,902 | 70 | 0.4996 |
| Columbia River contribution | 1,261 | 30 | 0.2171 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 1 WA | Columbia River | Escapement to NFH |
| 0 OR | 1,261 | 2,902 |
| 0 CA | | |
| 1 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 05 Spring Chinook raised at Winthrop NFH
 released in Methow River

02/01/2013
 rd2
 05WtSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 04/11/2007 | 218,411 | 17.9 | 217,974 | 051791 | Winthrop NFH |
| 04/11/2007 | 44,419 | 18.7 | 44,331 | 052177 | Winthrop NFH |
| 04/11/2007 | 36,757 | 19.6 | 36,757 | 052178 | Winthrop NFH |
| 04/11/2007 | 150,506 | 18.3 | 150,210 | 052678 | Winthrop NFH |
| 04/11/2007 | 109,933 | 18.3 | 109,715 | 052877 | Winthrop NFH |
| 04/11/2007 | 29,667 | 18.7 | 29,667 | 054863 | Winthrop NFH |
| Totals | 589,693 | | 588,654 | | |

| Where Recovered | Total | 451 Observed Recoveries | | | | | Total |
|-----------------------------|-------|---------------------------------|-----|-----|----|----|-------|
| | | === PSC Expanded Recoveries === | | | | | |
| | | 2s | 3s | 4s | 5s | 6s | |
| Columbia River Gillnet ODFW | 8 | | | 8 | | | 8 |
| Columbia River Sport ODFW | 26 | | | 26 | | | 26 |
| Freshwater Sport ODFW | 2 | | | 2 | | | 2 |
| Hatchery FWS | 294 | | 33 | 254 | 7 | | 294 |
| Hatchery WDFW | 20 | | 4 | 16 | | | 20 |
| Spawning Ground WDFW | 320 | | 64 | 256 | | | 320 |
| Totals | 670 | | 101 | 562 | 7 | | 670 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 670 | 100 | 0.1136 |
| Total Survival | 670 | | 0.1136 |
| Escapement to NFH | 294 | 44 | 0.0499 |
| Columbia River contribution | 376 | 56 | 0.0638 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 376 | 294 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 06 Spring Chinook raised at Winthrop NFH
 released in Methow River

02/01/2013
 rd2
 06WtSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 04/14/2008 | 27,650 | 19.8 | 27,318 | 052179 | Winthrop NFH |
| 04/14/2008 | 198,171 | 15.2 | 195,408 | 052574 | Winthrop NFH |
| 04/14/2008 | 102,205 | 14.7 | 100,979 | 053179 | Winthrop NFH |
| 04/14/2008 | 85,873 | 14.6 | 84,678 | 053180 | Winthrop NFH |
| 04/14/2008 | 95,146 | 15.7 | 87,684 | 053181 | Winthrop NFH |
| Totals | 509,045 | | 496,067 | | |

2,314 Observed Recoveries
 === PSC Expanded Recoveries ===

| Where Recovered | | Total | 2s | 3s | 4s | 5s | 6s | Total |
|------------------------|------|-------|----|-----|-------|-----|----|-------|
| Columbia River Gillnet | ODFW | 405 | | 2 | 384 | 19 | | 405 |
| Hatchery | FWS | 1,736 | | 68 | 1,568 | 100 | | 1,736 |
| Hatchery | WDFW | 26 | | 5 | 19 | 2 | | 26 |
| Spawning Ground | FWS | 11 | | | 11 | | | 11 |
| Spawning Ground | WDFW | 1,066 | | 464 | 538 | 64 | | 1,066 |
| Totals | | 3,244 | | 539 | 2,520 | 185 | | 3,244 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 3,244 | 100 | 0.6373 |
| Total Survival | 3,244 | | 0.6373 |
| Escapement to NFH | 1,736 | 54 | 0.3410 |
| Columbia River contribution | 1,508 | 46 | 0.2962 |

| Ocean | Contribution by Area | |
|---------|----------------------|-------------------|
| 0 AK | | |
| 0 BC | | |
| 0 WA | Columbia River | Escapement to NFH |
| 0 OR | 1,508 | 1,736 |
| 0 CA | | |
| 0 Total | | |

Stock Assessment Reference Summary from CRiS and PSMFC TS1
 Broodyear 07 Spring Chinook raised at Winthrop NFH
 released in Methow River

02/01/2013
 rd2
 07WtSCS

| Date | Number | #/lb | Tagged | CWT | AT |
|------------|---------|------|---------|--------|--------------|
| 04/16/2009 | 74,877 | 16.1 | 73,841 | 053576 | Winthrop NFH |
| 04/16/2009 | 144,970 | 17.2 | 135,451 | 054299 | Winthrop NFH |
| 04/16/2009 | 152,112 | 16.2 | 139,436 | 054364 | Winthrop NFH |
| Totals | 371,959 | | 348,728 | | |

994 Observed Recoveries

=== PSC Expanded Recoveries ===

| Where Recovered | Total | 2s | 3s | 4s | 5s | 6s | Total |
|-----------------------------|-------|----|-----|-------|----|----|-------|
| Columbia River Gillnet ODFW | 36 | | | 33 | 3 | | 36 |
| Columbia River Sport ODFW | 13 | | 4 | 9 | | | 13 |
| Freshwater Sport WDFW | 9 | | | 9 | | | 9 |
| Hatchery FWS | 860 | | 181 | 679 | | | 860 |
| Hatchery WDFW | 37 | | 8 | 29 | | | 37 |
| Spawning Ground WDFW | 284 | | 11 | 273 | | | 284 |
| Test Fishery Net ODFW | 1 | | | 1 | | | 1 |
| Totals | 1,240 | | 204 | 1,033 | 3 | | 1,240 |

| | Total | % of total | % of release |
|------------------------------|-------|------------|--------------|
| Ocean contribution | 0 | 0 | 0.0000 |
| Escapement to Columbia River | 1,240 | 100 | 0.3334 |
| Total Survival | 1,240 | | 0.3334 |
| Escapement to NFH | 860 | 69 | 0.2312 |
| Columbia River contribution | 380 | 31 | 0.1022 |

Ocean

Contribution by Area

- 0 AK
- 0 BC
- 0 WA
- 0 OR
- 0 CA
- 0 Total

Columbia River
 380

Escapement to NFH
 860