



OFFICE OF THE SECRETARY
**U.S. Department
of the Interior**

www.doi.gov

News Release

Date: June 17, 2014

Contact: Jessica Kershaw, Interior_Press@ios.doi.gov
Kirby-Lynn Shedlowski, kirbylynn_shedlowski@nps.gov

Lisa Iams, liams@usbr.gov

Tom Buckley, tom_buckley@fws.gov

Once Thought Locally Extinct, Endangered Razorback Suckers Discovered Spawning in Grand Canyon National Park

GRAND CANYON, Ariz. – The Department of the Interior announced today that researchers recently discovered Razorback suckers (*Xyrauchen texanus*), an endangered fish species, spawning in the lower Colorado River within Grand Canyon National Park. The detection of larval Razorback suckers, believed to have been missing from the Grand Canyon since the 1990s, provides evidence that these fish may be naturally reproducing in an area where the species has not been seen in more than 20 years.

“The discovery that the Razorback suckers are spawning in the national park far upstream from Lake Mead is good news for this endangered species,” said Assistant Secretary for Water and Science Anne Castle. “It is also a demonstration of the power of the partnership among Interior’s Bureau of Reclamation, National Park Service, U.S. Fish and Wildlife Service and U.S. Geological Survey, which have worked for recovery of this species side-by-side with Arizona state officials, Indian tribes, private landowners and researchers.”

The larval fish were first detected on April 14, 2014 and again in multiple samples from both April and May, confirming the occurrence of spawning and indicating that suitable habitat is available to support larger populations of this species.

“I’m proud to see that Grand Canyon provides habitat once again for this unique species, as well as for the endangered Humpback chub, and I’m committed to working with our cooperators to continue native fish conservation efforts within the park,” said Dave Uberuaga, the Superintendent of Grand Canyon National Park.”

Unique to the Colorado River Basin, Razorback suckers were once widespread and abundant throughout the Colorado River and its tributaries from the Green River in Wyoming to the Gulf of California. Because of basin-wide alterations in habitat and the introduction of nonnative

species, however, spawning and survival to adulthood were known to occur only in Lake Mead National Recreation Area.

Researchers typically determine the age of captured Razorback suckers and monitor their habitat use, reproduction, and movements, using specialized equipment designed to detect signals emitted from sonic tags implanted within the fish.

While the monitoring of the Razorback sucker has been conducted in the Lake Mead National Recreation Area for several years, the study was extended, beginning in the spring of 2014, from the Lake Mead Inflow area near Pearce Ferry upstream to Lava Falls Rapid within Grand Canyon National Park.

On March 16, 2014, in cooperation with the Nevada Department of Wildlife, BIO-WEST, Inc., and the Arizona Game and Fish Department, the National Park Service and Bureau of Reclamation successfully released nine adult endangered razorback suckers in the Colorado River within Grand Canyon National Park, downstream of Lava Falls (River Mile 180).

During a subsequent April monitoring trip in Grand Canyon conducted by BIO-WEST, Inc., American Southwest Ichthyological Researchers LLC, Bureau of Reclamation, and the National Park Service, biologists located several of the newly released sonic-tagged fish as well as previously tagged fish that had migrated upstream from Lake Mead.

“The most surprising result was finding larval Razorback suckers at 9 of the 47 locations in the park – far upstream from Lake Mead,” said Mark McKinstry, a biologist with the Bureau of Reclamation.

These results illustrate the resiliency of this fish to adapt to changing conditions according to Brandon Albrecht and Ron Kegerries of BIO-WEST, Inc., who have been monitoring the Lake Mead population for 18 years and are lead biologists for the Grand Canyon Razorback research.

Brian Healy, fisheries program manager for the national park, said “Razorback suckers continue to surprise us in Grand Canyon, first with the discovery of adults after 20 years of absence, and now with spawning within the park. We’re all hoping to see evidence that these larval fish survive to adulthood in the coming years.”

“This exciting news suggests that Grand Canyon is becoming a significant basin-wide haven for the endangered fishes in the Colorado River,” said Lesley Fitzpatrick, biologist for the U.S. Fish and Wildlife Service.

Razorback suckers are named for the bony keel on their backs. They are the largest species of suckers that live in the Colorado River and reach a maximum length of 36 inches. They may live 40 years or more, feeding on a variety of insects and crustaceans.

This study is part of a cooperative effort funded by the Bureau of Reclamation, with the National Park Service, U.S. Fish and Wildlife Service, Lower Colorado River Multi-Species Conservation Program, Nevada Department of Wildlife, Arizona Game and Fish Department, BIO-WEST, Inc., American Southwest Ichthyological Researchers LLC., and Hualapai Tribe and is a component of the recently released Comprehensive Fisheries Management Plan for Grand Canyon National Park.

The U.S. Geological Survey has conducted extensive previous research on how the aquatic ecosystem of the Colorado River in the Grand Canyon has been heavily influenced by Glen Canyon Dam and the decades of controlled release of water for power generation.

Additional information about Grand Canyon National Park's fisheries program is available online at <http://www.nps.gov/grca/naturescience/fish.htm>.

###

