

Bartonella spp. Exposure in Northern and Southern Sea Otters in Alaska and California

Sebastian E. Carrasco,¹ Bruno B. Chomel,^{1,2} Verena A. Gill,³ Angela M. Doroff,⁴ Melissa A. Miller,^{1,5} Kathleen A. Burek-Huntington,⁶ Rickie W. Kasten,² Barbara A. Byrne,¹ Tracey Goldstein,¹ and Jonna A.K. Mazet¹

Abstract

Since 2002, an increased number of northern sea otters (*Enhydra lutris kenyoni*) from southcentral Alaska have been reported to be dying due to endocarditis and/or septicemia with infection by *Streptococcus infantarius* subsp. *coli*. *Bartonella* spp. DNA was also detected in northern sea otters as part of mortality investigations during this unusual mortality event (UME) in Kachemak Bay, Alaska. To evaluate the extent of exposure to *Bartonella* spp. in sea otters, sera collected from necropsied and live-captured northern sea otters, as well as necropsied southern sea otters (*Enhydra lutris nereis*) unaffected by the UME, were analyzed using an immunofluorescent antibody assay. Antibodies against *Bartonella* spp. were detected in sera from 50% of necropsied and 34% of presumed healthy, live-captured northern sea otters and in 16% of necropsied southern sea otters. The majority of sea otters with reactive sera were seropositive for *B. washoensis*, with antibody titers ranging from 1:64 to 1:256. *Bartonella* spp. antibodies were especially common in adult northern sea otters, both free-living (49%) and necropsied (62%). Adult stranded northern sea otters that died from infectious causes, such as opportunistic bacterial infections, were 27 times more likely to be *Bartonella* seropositive than adult stranded northern sea otters that died from noninfectious causes ($p < 0.001$; 95% confidence interval 2.62–269.4). Because *Bartonella* spp. antibodies were detected in necropsied northern sea otters from southcentral (44%) and southwestern (86%) stocks of Alaska, as well as in necropsied southern sea otters (16%) in southcentral California, we concluded that *Bartonella* spp. exposure is widely distributed among sea otter populations in the Eastern Pacific, providing context for investigating future disease outbreaks and monitoring of *Bartonella* infections for sea otter management and conservation.

Key Words: Northern sea otter—Southern sea otter—*Bartonella* spp.—Antibodies—*Bartonella washoensis*—Alaska.

Introduction

NORTHERN AND SOUTHERN SEA OTTERS (*Enhydra lutris kenyoni* and *E. l. nereis*) are keystone predators that inhabit the near-shore coastal waters of Alaska and California, respectively (Riedman and Estes 1990). Because of a decline in the southwestern stock of northern sea otters in Alaska and slow population recovery of southern sea otters in California, these populations are currently listed as “threatened” under the United States Endangered Species Act (US

Fish and Wildlife Department 2008; US Geological Survey 2010). Causes of the decline and slow recovery are multifactorial and mediated through complex interactions among sea otter populations and numerous biotic and environmental factors, including infectious diseases (Kreuder et al. 2003, Goldstein et al. 2009, Miller et al. 2010b, Counihan-Edgar et al. 2012). Epidemiological studies have demonstrated that southern sea otters are at high risk for exposure to gastrointestinal protozoal pathogens and opportunistic bacteria in the coastal waters of central California (Miller et al. 2008,

¹Wildlife Health Center, One Health Institute, School of Veterinary Medicine, University of California, Davis, California.

²Department of Population Health and Reproduction, School of Veterinary Medicine, University of California, Davis, California.

³US Fish and Wildlife Service, Marine Mammals Management, Anchorage, Alaska.

⁴Kachemak Bay Research Reserve, Homer, Alaska.

⁵California Department of Fish and Wildlife, Marine Wildlife Veterinary Care and Research Center, Santa Cruz, California.

⁶Alaska Veterinary Pathology Services, Eagle River, Alaska.

Miller et al. 2010a). Recent data also suggest that northern sea otters in Alaska have suffered increased exposure to and infection with pathogenic microorganisms that could make them more susceptible to predation and disease-associated mortality (Goldstein et al. 2009, Goldstein et al. 2011, Brownstein et al. 2011).

Because a large number of northern sea otters were found dead in southcentral Alaska beginning in 2006, the US Working Group on Marine Mammal Unusual Mortality Events declared an unusual mortality event (UME; Gill 2006). Since that time, diagnostic efforts have been directed toward understanding the causes of the sea otter UME. *Streptococcus infantarius* subsp. *coli* (a member of *Streptococcus bovis/equinus* complex, SB/E) infection was identified to be highly associated with the mortality in Kachemak Bay and surrounding areas. This bacterium was isolated from heart valves of northern sea otters with vegetative valvular endocarditis (VVE), with or without septicemia (Gill 2006). Recently, we reported that up to 45% of examined northern sea otters had detectable *Bartonella* spp. DNA, and 33% of the heart valves from animals that died with VVE were co-infected with *Bartonella* spp. and *S. infantarius* subsp. *coli* (Carrasco et al. 2014).

Bartonella spp. are fastidious Gram-negative bacteria that infect erythrocytes and vascular endothelial cells of hosts and are usually transmitted by blood-sucking arthropods (Chomel et al. 2009). This facultative intracellular bacterium causes persistent asymptomatic bacteremia and has been associated with debilitating and possibly life-threatening illnesses, including encephalopathies, bacillary angiomatosis, myocarditis, and valvular endocarditis in domestic animals and humans (Breitschwerdt et al. 2010). Clinical manifestations vary on the basis of the immune status of the host, the infecting *Bartonella* strain, and the co-evolutionary history of a *Bartonella* spp. with its animal host (Chomel et al. 2009). In their mammalian reservoir hosts, *Bartonella* spp. often cause asymptomatic chronic bacteremia, whereas in the nonreservoir hosts this pathogen is typically associated with multiple disease processes (Chomel et al. 2009). *Bartonella* spp. have been increasingly isolated or detected in blood or lesions from live-captured or moribund marine animals (Maggi et al. 2008, Morick et al. 2009); however, little is known about the epidemiological significance, as hosts and transmission dynamics of this organism in the marine environment are relatively unstudied.

Because *Bartonella* DNA was recently detected in northern and southern sea otter carcasses from Alaska and California (Carrasco et al. 2014), our main objective was to assess exposure in northern and southern sea otters to this usually vector-borne pathogen. The goals of this study were two-fold: (1) To determine the frequency and distribution of serum antibodies against selected *Bartonella* species in northern and southern sea otters and (2) to evaluate risk factors associated with *Bartonella* exposure in both populations. Our findings provided baseline serological data and insight into the epidemiology of bartonellosis in sea otters to contribute to conservation efforts.

Materials and Methods

Sampling

Serum samples were collected from 44 apparently healthy, live-captured northern sea otters sampled in Kachemak Bay, Alaska, in 2007 by the United States Fish and Wildlife Ser-

vice (USFWS), Anchorage, Alaska. Blood was collected from anesthetized sea otters using well-established protocols for immobilization, sampling, and release (Doroff and Badajos 2010). Samples were also obtained from 48 northern sea otters that stranded and were necropsied between 2004 and 2009. Postmortem examinations of northern sea otters were performed at the Anchorage USFWS office by a veterinary pathologist or by veterinarians with marine mammal necropsy experience.

At necropsy, northern sea otters were categorized into either the infectious or noninfectious group on the basis of the diagnosed cause of death, such as opportunistic bacterial infections. Thirty-three individuals were categorized into an “infectious” group, whereas those animals dying due to trauma and/or other causes and with no evidence of concurrent infectious disease were included in a “noninfectious” group ($n=15$). Bacterial pathogens isolated or detected in lesions of necropsied northern sea otters included *S. infantarius* subsp. *coli* or SB/E organisms, *Bartonella* spp., and *Streptococcus phocae*, as previously described (Carrasco et al. 2014). Stranded sea otters were found dead or moribund in south-central Alaska, Kodiak Archipelago, and the western side of Cook Inlet, including Ninilchik, Homer and Kachemak Bay, Whittier, Kodiak Island, Hallo Bay, and Kamishak Bay (Fig. 1). Sea otters from southcentral stock that had stranded in the Ninilchik, Whittier, Homer, and Kachemak Bay regions were further subcategorized into a “Kachemak Bay area” group, whereas animals from southwest stock that had stranded in Kamishak Bay, Hallo Bay, and Kodiak Island were pooled into a “southwest Alaska” group. These categories were selected because mortalities caused by SB/E infection were common in northern sea otters from southcentral stock in Kachemak Bay and surrounding areas (adjacent to the threatened southwestern stock).

In addition, 148 serum samples were obtained from necropsied southern sea otters that had stranded during 2001–2009 in California. Postmortem examinations of southern sea otters were performed at the Marine Wildlife Veterinary Care and Research Center, California Department of Fish and Wildlife (CDFW), Santa Cruz, California, by a veterinary pathologist to establish the cause of death. As with the northern sea otters, southern sea otters that were classified into the “infectious” group generally were diagnosed with a cause of death due to either bacterial or parasitic pathogens or had bacterial or parasitic infections as a contributing cause of death ($n=72$ animals), whereas animals that were diagnosed with other noninfectious causes of death or had contributing factors that were not associated with infectious disease were included in a “noninfectious” group ($n=76$). Acknowledged pathogens of southern sea otters included a range of bacterial, protozoal, and acanthocephalan parasites, as previously described (Kreuder et al. 2003, Miller et al. 2010a). These sea otters were found dead or moribund on beaches in various geographic areas along the central California coast, including Monterey Bay, Spanish Bay, Carmel Bay, Julia Pfeiffer State Beach, San Simeon Bay and surroundings, Morro Bay and surroundings, Diablo Canyon, and San Luis Obispo Bay and Pismo Beach (Fig. 1.).

The age of each sea otter was estimated at the time of necropsy based on body mass, weight, dentition, pelage characteristics, and reproductive status (Riedman and Estes 1990, Von Biela et al. 2011). Pups, juveniles, and subadults




FIG. 1. Maps of Alaska and California showing stranding areas of northern and southern sea otters that were necropsied and tested by indirect immunofluorescent antibody (IFA) assay for *Bartonella* spp. (A) Alaska stranding locations 2004–2009. Eighteen of the seropositive northern sea otter carcasses stranded on beaches from Kachemak Bay area (44%, 18/41), whereas six positive otter carcasses (86%, 6/7) were found in southwest Alaska group, including Kamishak Bay area ($n = 4$) and Kodiak Island ($n = 2$). (B) California stranding locations 2001–2009. Nineteen (22%, 19/85) seropositive samples came from otters stranding near Monterey Bay. Five additional seropositive samples (11%, 5/46) were obtained from the sea otter carcasses stranding in the lower portion of the southern sea otter range near Morro Bay ($n = 1$), San Luis Obispo Bay ($n = 2$), and Pismo Beach ($n = 2$).

were categorized as “immature” otters, and adults and aged adults were categorized as “adult” otters. Nutritional condition was assessed by the amount of subcutaneous fat present and by palpation of the ribs, backbone, and other bony protuberances (Doroff and Mulcahy 1997). Otters were categorized as being in “poor” nutritional condition if they were thin or emaciated with scant subcutaneous fat and evidence of muscle loss, or in “good” nutritional condition if they had moderate to abundant subcutaneous fat.

Serology

Antibodies against *Bartonella henselae* type II (also known as Marseille; isolate U4, University of California–Davis), *Baronella clarridgeiae* (ATCC 51734), and *Bartonella washoensis* (isolate UCD-dog2, University of California–Davis) in serum and/or whole blood samples from northern and southern sea otters were detected using an indirect immunofluorescent antibody (IFA) assay. The IFA procedure was similar to the procedure described by Henn et al. (2007) with a slight modification. Fluorescein isothiocyanate (FITC)-conjugated goat anti-ferret immunoglob-

ulin G (IgG) (Bethyl Laboratories, Montgomery, TX) was used for detection of primary antibodies. *B. henselae*, *B. clarridgeiae*, and *B. washoensis* strains were used as surrogate bacteria for this assay due to the lack of success obtaining a *Bartonella* isolate from sea otter tissues. Samples were examined at dilutions of 1:64 and 1:128 using a fluorescence microscope at 400× magnification. The intensity of bacillus-specific fluorescence was scored subjectively from 1 to 4, as previously described (Chang et al. 2000). Samples with a fluorescence score of ≥ 2 at a dilution of 1:64 were considered positive because this antibody titer has proven to indicate prior exposure and current infection by *Bartonella* spp. in dogs and cats (Chomel et al. 1995, Pappalardo et al. 1997). Samples testing positive were titrated in serial two-fold dilutions until the fluorescence score was below the limit of detection (< 2). Each slide was examined in a blinded fashion by three readers, and fluorescence scores were averaged.

Statistical analysis

Data collected on each necropsied northern and southern sea otter included animal identification number, age class, sex,

stranding date and location, date of sampling, weight, nutritional condition, and cause of death from necropsy findings, including pathological lesions and reported laboratory results for any postmortem bacterial culture or parasitology result. Additional SB/E or *S. infantarius* subsp. *coli* and *S. phocae* culture results were provided by the USFWS from microbiological reports from the Microbiology Laboratory (Veterinary Medical Teaching Hospital, University of California–Davis). Chi-squared analyses were used to evaluate associations between *Bartonella* IFA results and age, sex, nutritional condition, stranding area, and animal grouping as “infectious” or “non-infectious” cause of death using JMPv8.0 software (SAS Institute Inc, www.jmp.com). Associations between *Bartonella* IFA results and live-captured otters and necropsied northern sea otters were also analyzed using chi-squared analyses. Stratified analyses were considered where appropriate after univariate analyses were completed. Odds ratios (ORs) with 95% confidence intervals (CIs) were used to evaluate the strength of association of *Bartonella* IFA results with each of the variables that were significant in the stratified analysis. A Fisher exact test was performed to analyze variables with limited sample sizes (Fisher 1935). For all tests, *p* values ≤ 0.05 were considered statistically significant. Sampling sites were mapped using ArcGIS 10.2 software (ESRI, Redlands, CA) for all necropsied northern and southern sea otters tested by serology.

Results

Serological findings

Antibodies reactive to *Bartonella* spp. were detected in 34% (15/44) and 50% (24/48) of serum samples from live-captured and necropsied northern sea otters, respectively, with antibody titers ranging from 1:64 to 1:256 (Tables 1 and 2). By comparison, *Bartonella* spp. seroprevalence was lower (16%, 24/148) for southern sea otters (Table 1). The majority of positive sea otter sera were reactive for *B. washoensis*, as antibodies reactive to *B. washoensis* ($\geq 1:64$ IFA titers) were detected in 27% (12/44) of sera from live-captured northern sea otters (Table 1). Two (4.5%, 2/44)

TABLE 1. PREVALENCE OF ANTIBODIES AGAINST *B. HENSELAE* (BH), *B. CLARRIDGEIAE* (BC), AND *B. WASHOENSIS* (BW) IN APPARENTLY HEALTHY, LIVE-CAPTURED NORTHERN SEA OTTERS AND NECROPSIED NORTHERN AND SOUTHERN SEA OTTERS USING AN INDIRECT IMMUNOFLOUORESCENT ANTIBODY (IFA) TEST

	Live-captured northern sea otters (n=44)	Necropsied northern sea otters (n=48)	Necropsied southern sea otters (n=148)
<i>Bartonella</i> spp.			
<i>B. washoensis</i>	27% (12/44)	14% (7/48)	4.7% (7/148)
<i>B. clarridgeiae</i>	2.2% (1/44)	ND	1.3% (2/148)
<i>B. henselae</i>	ND	ND	2.0% (3/148)
Bh and Bw	ND	25% (12/48)	5.4% (8/148)
Bh and Bc	ND	2% (1/48)	ND
Bc and Bw	4.5% (2/44)	2% (1/48)	1.3% (2/148)
Bh, Bc, and Bw	ND	6.2% (3/48)	1.3% (2/148)
Total <i>Bartonella</i> positives	34% (15/44)	50% (24/48)	16% (24/148)

ND, Not detected.

TABLE 2. THE DISTRIBUTION OF *BARTONELLA* INDIRECT IMMUNOFLOUORESCENT ANTIBODY TITERS IN LIVE-CAPTURED NORTHERN SEA OTTERS AND NECROPSIED NORTHERN AND SOUTHERN SEA OTTERS

<i>Bartonella</i> IFA titer	Live- captured northern sea otters (n=44)	Necropsied northern sea otters (n=48)	Necropsied southern sea otters (n=148)	Total number of samples by titer level (n=240)
$\geq 1:256^a$	1	4	1	6
$\geq 1:128^a$	7	11	7	25
$\geq 1:64^a$	7	9	16	32
$< 1:64$	29	24	124	177

^aAntibody titers $\geq 1:64$ were classified as seropositive. IFA, indirect immunofluorescent assay.

were seropositive for *B. clarridgeiae* and *B. washoensis*, and one sample was seropositive only to *B. clarridgeiae* ($\geq 1:128$). Of 48 necropsied northern sea otters sampled, seven (14%, 7/48) reacted only to *B. washoensis* and 12 (25%, 12/48) were reactive to *B. henselae* and *B. washoensis* (Table 1). Antibodies reactive to *B. washoensis* ($\geq 1:64$) were detected in 13% (19/148) of sera from necropsied southern sea otters (Table 1). Seven samples (4.7%, 7/148) were seropositive only to *B. washoensis*, and eight samples (5.4%, 8/148) tested positive for *B. henselae* and *B. washoensis*.

Association of *Bartonella* seroprevalence with demographic and spatial factors

Significant associations were identified between *Bartonella* seropositivity in northern sea otter carcasses and adult age class, poor nutritional condition, and death due to infectious causes (Table 3). *Bartonella* seropositivity was not more common in necropsied northern sea otters when compared to apparently healthy live animals. A much higher proportion of *Bartonella*-seropositive stranded northern sea otters died with infectious disease as a primary or major contributing cause of death, when compared to stranded northern sea otters that died from noninfectious causes (Table 3). Adult and aged adult stranded northern sea otters that died with infectious disease were 27 times more likely (Fisher exact test, $p < 0.001$; 95% CI 2.62–269.4) to be seropositive to *Bartonella* spp. than northern sea otters that died from noninfectious causes. A similar age association was observed for *Bartonella*-seropositive, live-captured northern sea otters ($p = 0.04$). Of 28 adult, live-captured northern sea otters tested, 12 (49%) were seropositive, compared to two out of 16 immature sea otters (13%; $p = 0.04$).

In contrast to observed patterns for northern sea otters, no significant associations were observed between *Bartonella* seropositivity and age class, sex, nutritional condition, or cause of death for southern sea otter carcasses. Southern and northern sea otter carcasses that tested positive for *Bartonella* spp. were found on beaches in various geographic areas of California and Alaska (Fig. 1).

Discussion

The high proportion of *B. washoensis*-seropositive sea otters was consistent with our recent detection of *Bartonella* DNA in sea otter heart valves from Alaska and California

TABLE 3. DEMOGRAPHIC RISK FACTOR CATEGORIES EVALUATED FOR THEIR ASSOCIATION WITH *BARTONELLA* SEROPOSITIVITY IN NECROPSIED NORTHERN AND SOUTHERN SEA OTTERS

Risk factor	Categories	Northern sea otters (n = 48) ^a	Southern sea otters (n = 148) ^b	p values
		% Seropositive	% Seropositive	
Sex	Male	45.7 (16/35)	12.3 (10/81)	a = 0.33 / b = 0.16
	Female	61.5 (8/13)	20.8 (14/67)	
Age	Immature	25 (4/16)	13 (6/46)	a = 0.01 / b = 0.48
	Adult	62.5 (20/32)	17.6 (18/102)	
Nutritional condition	Poor	62 (18/29)	17.8 (17/95)	a = 0.02 / b = 0.46
	Good	27.7 (5/18)	13.2 (7/53)	
Cause of death	Noninfectious	13.3 (2/15)	10.5 (8/76)	a ≤ 0.001 / b = 0.08
	Infectious	66.6 (22/33)	22.2 (16/72)	

(Carrasco et al. 2014). The *Bartonella* DNA detected was closely related to *B. henselae*, *B. washoensis*, and *Candidatus Bartonella volans*, suggesting that northern and southern sea otters are infected with more than one *Bartonella* species (Carrasco et al. 2014). Consistent with that finding, seroreactivity to greater than one *Bartonella* spp. was detected in 67% (16/24) and 50% (12/24) of *B. washoensis*-positive samples from northern and southern sea otter carcasses, respectively. This result was not surprising given that serological cross-reactivity of these three *Bartonella* spp. and others has been previously reported in sera from *Bartonella*-infected humans and animals (Maurin et al. 2002, Chomel et al. 2003, Tsuneoka et al. 2004, Boulouis et al. 2005). In contrast to samples from sea otter carcasses, the majority of the sera from presumably healthy, live-captured northern sea otters showed exclusive reactivity against *B. washoensis* (86%; 12/14 *B. washoensis*-positive samples). The combined PCR (Carrasco et al. 2014) and serology results from this study indicated that at least *B. washoensis*-like organisms were circulating among northern sea otters, consistent with recent findings for closely related American river otters (*Lontra canadensis*), that were also infected with either *B. washoensis*-like or *B. volans*-like organisms (Chinnadurai et al. 2010).

Bartonella exposure was detected among all northern sea otter age groups, but it was most common in adults. This finding is consistent with serosurveys of *B. henselae* in cats (Guptill et al. 2004) and *Bartonella vinsonii* subsp. *berkhoffii* in coyotes (Chang et al. 1999, Chang 2000) that showed that *Bartonella* seroprevalence increased with age, whereas *Bartonella* bacteremia decreased as the age of these other species increased. Because seropositivity was common and the risk of exposure increased with age, it is likely that *Bartonella* spp. exposure is endemic in northern sea otters in Alaska. *Bartonella* spp. are usually transmitted among terrestrial reservoirs by blood-sucking arthropods such as sand flies, fleas, or lice (Chomel et al. 2009); however, little is known regarding potential vectors for *Bartonella* spp. transmission in aquatic animals. Interestingly, *Bartonella* DNA identical to that from *Bartonella*-infected harbor seals was detected in hematophagous lice that are common pin-niped ectoparasites (Thompson et al. 1998, Morick et al. 2009). This finding suggested that marine host-adapted vectors may be able to transmit *Bartonella* spp.

Reports of co-infections of *Bartonella* spp. and other opportunistic pathogens are common in domestic animals and

humans (Breitschwerdt and Kordick 2000, Boulouis et al. 2005). Our investigation revealed a similar relationship between *Bartonella* seropositivity and death due to infectious causes for stranded northern sea otters, as 16 of the 22 seropositive animals from the infectious group died with SB/E-associated endocarditis or septicemia. Furthermore six of these 22 seropositive northern sea otters were co-infected with *Bartonella* DNA and SB/E organisms, as was recently described (Carrasco et al. 2014). Interestingly, five of these six seropositive animals had antibodies reactive to *B. washoensis* and detectable *Bartonella* DNA that was closely related to *B. washoensis* and *Candidatus B. volans* (Carrasco et al. 2014). Some *B. washoensis*-seropositive northern sea otters that had died due to VVE yielded titers ≥ 1:256, as commonly reported in *Bartonella*-associated endocarditis in dogs and humans (Maurin et al. 2002, MacDonald et al. 2004). However, the clinical impact of disseminated *Bartonella* infection in sea otters was not clear. The observed high mortality due to VVE for northern sea otters from this region and confirmation of co-infection by *Bartonella* and other pathogenic bacteria in prior studies (Brownstein et al. 2011, Carrasco et al. 2014) suggests that *Bartonella* co-infections may be a contributor to the development of debilitating disease in northern sea otters.

Previous studies in dogs and humans with infective endocarditis have shown evidence of serological cross-reactivity between *Bartonella* spp. and intracellular bacteria (LaScola and Raoult 1996, Maurin et al. 2002, Macdonald et al. 2004). Although it is reasonable to speculate that SB/E-infected northern sea otters could have cross-reactive antibodies to *Bartonella* antigens using our IFA-based assay, data from our laboratory and others have shown that cross-reactivity between *Bartonella* spp. and *Streptococcus* spp. is rare when samples are evaluated with this serological assay (Sykes et al. 2006, Vermeulen et al. 2010). However, a potential limitation of our findings is that we cannot rule out the possibility of cross-reactive antibodies to other intracellular pathogens that have shown to interfere in the diagnosis of bartonellosis in dogs and humans (Maurin et al. 2002, MacDonald et al. 2004).

Exposure to *Bartonella* spp. was widely distributed among sea otters from the southcentral stock of Alaska, because positive samples were detected in otters from Ninilchik, Whittier, and Kachemak Bay. In addition, *Bartonella* spp. antibodies were detected in the southwestern stock of northern sea otters that inhabit areas adjacent to Kachemak

Bay, including Kamishak Bay and Kodiak Island. This broad distribution of seropositivity suggested that this bacterium is endemic in northern sea otters. Due to shared haul-out sites and rafting behavior of sea otters in southcentral Alaska (Doroff and Badajos 2010) and the wide range and diversity of seroprevalence to *Bartonella* spp. documented here, it appears that *Bartonella* may be transmitted horizontally between sea otters, possibly by yet unidentified arthropod vector/vectors. It is also possible that a common vector may be involved in the transmission of *Bartonella* between northern sea otters and river otters since fecal analyses in haul-out sites in Kachemak Bay showed that otters likely used the same haul-out sites as coastal river otters at different times of day (Doroff et al. 2012).

In conclusion, this study together with our recently published molecular findings (Carrasco et al. 2014) confirms that *Bartonella* organisms have been present in sea otters over the past decade and provides evidence for widespread exposure to *Bartonella* spp. among northern and southern sea otters. This bacterium seems to be regionally endemic for northern sea otter subspecies in Kachemak Bay and surrounding areas. Our findings provide baseline data on the frequency and distribution of *Bartonella* exposure among northern and southern sea otters in Alaska and California, providing context for investigating future disease outbreaks and monitoring of *Bartonella* infections for sea otter management and conservation.

Acknowledgments

The authors thank the personnel from the US Fish and Wildlife Service (USFWS), Anchorage, Alaska; Alaska SeaLife Center, Seward, Alaska; and the Marine Wildlife Veterinary Care and Research Center, California Department of Fish and Wildlife (CDFW), Santa Cruz, California; as well as the Homer stranding network for assistance with animal captures, necropsies, and sample collection. We also thank Kate Thomas for helping to illustrate sampling areas and Dana Jensi (USFWS) and Erin Dodd and Eva Berberich (CDFW) for their assistance on demographic and spatial information of sea otters and logistics for collection and shipping samples. Field sample collections in Alaska were conducted under authority of a Marine Mammal Protection Act permit issued by the Division of Management Authority, USFWS (MA041309-4 permit number) and under an Institutional Animal Care and Use Committee (IACUC) review by the University of Alaska. This research was supported by grant no. D08ZO-046 from the Morris Animal Foundation; the Karen C. Drayer Wildlife Health Center graduate student fellowship, University of California, Davis; and an Oceans and Human Health training grant no. NA06OAR4310119 (Training Tomorrow's Ecosystem and Public Health Leaders Using Marine Mammals as Sentinels of Oceanic Change) from the National Oceanic and Atmospheric Administration (NOAA). S.E.C. was also supported by grant no. NA06OAR4310119 and National Institutes of Health National Research Service Award T32 AI 060519.

Author Disclosure Statement

No competing financial interests exist.

The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the US Fish and Wildlife Service.

References

- Boulouis HJ, Chang CC, Henn JB, Kasten RW, et al. Factors associated with the rapid emergence of zoonotic *Bartonella* infections. *Vet Res* 2005; 36:383–410.
- Breitschwerdt EB, Kordick DL. *Bartonella* infection in animals: Carriership, reservoir potential, pathogenicity, and zoonotic potential for human infection. *Clin Microbiol Rev* 2000; 13:428–38.
- Breitschwerdt EB, Maggi RG, Chomel BB, Lappin MR. Bartonellosis: An emerging infectious disease of zoonotic importance to animals and human beings. *J Vet Emerg Crit Care (San Antonio)* 2010; 20:8–30.
- Brownstein D, Miller MA, Oates SC, Byrne BA, et al. Antimicrobial susceptibility of bacterial isolates from sea otters (*Enhydra lutris*). *J Wildl Dis* 2011; 47:278–292.
- Carrasco SE, Chomel BB, Gill VA, Kasten RW, et al. Novel *Bartonella* infection in northern and southern sea otters (*Enhydra lutris kenyoni* and *Enhydra lutris nereis*). *Vet Micro* 2014; 170:325–334.
- Chang C, Yamamoto K, Chomel BB, Kasten RW, et al. Seroepidemiology of *Bartonella vinsonii* subsp. *berkhoffii* infection in California coyotes, 1994–1998. *Emerg Infect Dis* 1999; 5:711–715.
- Chang CC, Kasten RW, Chomel BB, Simpson DC, et al. Coyotes (*Canis latrans*) as the reservoir for a human pathogenic *Bartonella* sp.: Molecular epidemiology of *Bartonella vinsonii* subsp. *berkhoffii* infection in coyotes from central coastal California. *J Clin Microbiol* 2000; 38:4193–4200.
- Chinnadurai SK, Birkenheuer AJ, Blanton HL, Maggi RG, et al. Prevalence of selected vector-borne organisms and identification of *Bartonella* species DNA in North American river otters (*Lontra canadensis*). *J Wildl Dis* 2010; 46:947–950.
- Chomel BB, Abbott RC, Kasten RW, Floyd-Hawkins KA, et al. *Bartonella henselae* prevalence in domestic cats in California: Risk factors and association between bacteremia and antibody titers. *J Clin Microbiol* 1995; 33:2445–2450.
- Chomel BB, Wey AC, Kasten RW. Isolation of *Bartonella washoensis* from a dog with mitral valve endocarditis. *J Clin Microbiol* 2003; 41:5327–5332.
- Chomel BB, Boulouis HJ, Breitschwerdt EB, Kasten RW, et al. Ecological fitness and strategies of adaptation of *Bartonella* species to their hosts and vectors. *Vet Res* 2009; 40:29.
- Counihan-Edgar KL, Gill VA, Doroff AM, Burek KA, et al. Genotypic characterization of *Streptococcus infantarius* subsp. *coli* isolates from sea otters with infective endocarditis and/or septicemia and from environmental mussel samples. *J Clin Microbiol* 2012; 50:4131–4133.
- Doroff AM, Badajos O. Monitoring survival and movement patterns of sea otters (*Enhydra lutris kenyoni*) in Kachemak Bay, Alaska, August 2007–April 2010. Final Report. Kachemak Bay Research Reserve, Homer, Alaska, 2010:1–18.
- Doroff AM, Mulcahy D. A field guide to general necropsy and tissue collection for sea otters (*Enhydra lutris*) in Alaska. Marine Mammals Management Technical Report MMM. U.S. Fish and Wildlife Service, 1997:97–93.
- Doroff A, Badajos O, Corbell K, Jensi D, et al. Assessment of sea otter (*Enhydra lutris kenyoni*) diet in Kachemak Bay, Alaska 2008–2010. *IUCN Otter Spec Group Bull* 2012; 29:1:15–23.
- Fisher RA. The logic of inductive inference. *J R Stat Soc Ser A* 1935; 98:39–82.
- Gill VA. Marine mammal unusual mortality event initiation protocol for northern sea otters. US Fish and Wildlife Service Report, Anchorage Alaska, 2006:1–11.

- Goldstein T, Mazet JA, Gill VA, Doroff AM, et al. Phocine distemper virus in northern sea otters in the Pacific Ocean, Alaska, USA. *Emerg Infect Dis* 2009; 15:925–927.
- Goldstein T, Gill VA, Tuomi P, Monson D, et al. Assessment of clinical pathology and pathogen exposure in sea otters (*Enhydra lutris*) bordering the threatened population in Alaska. *J Wildl Dis* 2011; 47:579–592.
- Guptill L, Wu CC, HogenEsch H, Slater LN, et al. Prevalence, risk factors, and genetic diversity of *Bartonella henselae* infections in pet cats in four regions of the United States. *J Clin Microbiol* 2004; 42:652–659.
- Henn JB, Gabriel MW, Kasten RW, Brown RN, et al. Gray foxes (*Urocyon cinereoargenteus*) as a potential reservoir of a *Bartonella clarridgeiae*-like bacterium and domestic dogs as part of a sentinel system for surveillance of zoonotic arthropod-borne pathogens in northern California. *J Clin Microbiol* 2007; 45:2411–2418.
- Kreuder C, Miller MA, Jessup DA, Lowenstine LJ, et al. Patterns of mortality in southern sea otters (*Enhydra lutris nereis*) from 1998–2001. *J Wildl Dis* 2003; 39:495–509.
- La Scola B, Raoult D. Serological cross-reactions between *Bartonella quintana*, *Bartonella henselae*, and *Coxiella burnetii*. *J Clin Microbiol* 1996; 34:2270–2274.
- Maggi RG, Raverty SA, Lester SJ, Huff DG, et al. *Bartonella henselae* in captive and hunter-harvested beluga (*Delphinapterus leucas*). *J Wildl Dis* 2008; 44:871–877.
- MacDonald KA, Chomel BB, Kittleson MD, Kasten RW, et al. A prospective study of canine infective endocarditis in northern California (1999–2001): Emergence of *Bartonella* as a prevalent etiologic agent. *J Vet Intern Med* 2004; 18:56–64.
- Maurin M, Rolain JM, Raoult D. Comparison of in-house and commercial slides for detection by immunofluorescence of immunoglobulins G and M against *Bartonella henselae* and *Bartonella quintana*. *Clin Diagn Lab Immunol* 2002; 9:1004–1009.
- Miller MA, Miller WA, Conrad PA, James ER, et al. Type X *Toxoplasma gondii* in a wild mussel and terrestrial carnivores from coastal California: New linkages between terrestrial mammals, runoff and toxoplasmosis of sea otters. *Int J Parasitol* 2008; 38:1319–1328.
- Miller MA, Byrne BA, Jang SS, Dodd EM, et al. Enteric bacterial pathogen detection in southern sea otters (*Enhydra lutris nereis*) is associated with coastal urbanization and freshwater runoff. *Vet Res* 2010a; 41:1.
- Miller MA, Conrad PA, Harris M, Hatfield B, et al. A protozoal-associated epizootic impacting marine wildlife: Mass-mortality of southern sea otters (*Enhydra lutris nereis*) due to *Sarcocystis neurona* infection. *Vet Parasitol* 2010b; 172: 183–194.
- Morick D, Osinga N, Gruys E, Harrus S. Identification of a *Bartonella* species in the harbor seal (*Phoca vitulina*) and in seal lice (*Echinophthirius horridus*). *Vector Borne Zoonotic Dis* 2009; 9:751–753.
- Pappalardo BL, Correa MT, York CC, Peat CY, et al. Epidemiologic evaluation of the risk factors associated with exposure and seroreactivity to *Bartonella vinsonii* in dogs. *Am J Vet Res* 1997; 58:467–471.
- Riedman ML, Estes JA. The sea otter (*Enhydra lutris*): Behavior, ecology, and natural history. US Fish and Wildlife Service, Biological Report; 1990:126.
- Sykes JE, Kittleson MD, Pesavento PA, Byrne BA, et al. Evaluation of the relationship between causative organisms and clinical characteristics of infective endocarditis in dogs: 71 cases (1992–2005). *J Am Vet Med Assoc* 2006; 228:1723–1734.
- Thompson PM, Corpe HM, Reid RJ. Prevalence and intensity of the ectoparasite *Echinophthirius horridus* on harbour seals (*Phoca vitulina*): Effects of host age and inter-annual variability in host food availability. *Parasitology* 1998; 117:393–403.
- Tsuneoka H, Umeda A, Tsukahara M, Sasaki K. Evaluation of indirect fluorescence antibody assay for detection of *Bartonella clarridgeiae* and Seroprevalence of *B. clarridgeiae* among patients with suspected cat scratch disease. *J Clin Microbiol* 2004; 42:3346–3349.
- US Fish and Wildlife Service (USFWS). Northern sea otters (*Enhydra lutris kenyoni*): Southwest Alaska Stock 2008. Available at <http://alaska.fws.gov/fisheries/mmm/seaotters/reports.htm> Accessed October, 2013.
- United States Geological Survey (USGS). Spring 2010 Mainland California Sea Otter Survey Results. USGS Western Ecological Research Center, Unpublished USGS report. Available at <http://www.werc.usgs.gov/Project.aspx?ProjectID=91>
- Vermeulen MJ, Verbakel H, Notermans DW, Reimerink JH, et al. Evaluation of sensitivity, specificity and cross-reactivity in *Bartonella henselae* serology. *J Med Microbiol* 2010; 59:743–745.
- Von Biela VR, Testa JW, Gill VA, Burns JM. Evaluating cementum to determine past reproduction in northern sea otters. *J Wildl Management* 2008; 72:618–624.

Address correspondence to:

Sebastian E. Carrasco
Wildlife Health Center, One Health Institute
School of Veterinary Medicine
University of California
Davis, CA 95616

E-mail: secarrasco@ucdavis.edu, scarrasc@iupui.edu