

NORTHERN SEA OTTER (*Enhydra lutris kenyoni*): Southeast Alaska Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Sea otters occur in nearshore coastal waters of the U.S. along the North Pacific Rim from the Aleutian Islands to California. The species is most commonly observed within the 40-meter depth contour since animals require frequent access to benthic foraging habitat in subtidal and intertidal zones (Reidman and Estes 1990). Sea otters in Alaska are not migratory and generally do not disperse over long distances, although movements of tens of kilometers are normal (Garshelis and Garshelis 1984). Individuals are capable of long distance movements of over 100 km (Garshelis et al. 1984); however, movements of sea otters are likely limited by geographic barriers, high energy requirements of the animals, and social behavior.

Applying the phylogeographic approach of Dizon et al. (1992), Gorbics and Bodkin (2001) identified three sea otter stocks in Alaska: southeast, southcentral, and southwest. The ranges of these stocks are defined as follows: (1) Southeast Alaska stock extends from Dixon Entrance to Cape Yakataga; (2) Southcentral Alaska stock extends from Cape Yakataga to Cook Inlet including Prince William Sound, the Kenai Peninsula coast, and Kachemak Bay; and (3) Southwest Alaska stock includes the Alaska Peninsula and Bristol Bay coasts, the Aleutian, Barren, Kodiak, and Pribilof Islands (Figure 1).




Figure 1. Approximate distribution of northern sea otters in Alaska waters (shaded area)

POPULATION SIZE

Historically, sea otters occurred across the North Pacific Rim, ranging from Hokkaido, Japan through the Kuril Islands, the Kamchatka Peninsula, the Commander Islands, the Aleutian Islands, peninsular and south coastal Alaska and south to Baja California, Mexico (Kenyon 1969). In the early 1700s, the worldwide population was estimated to be between 150,000 (Kenyon 1969) and 300,000 individuals (Johnson 1982). Prior to large-scale commercial exploitation, indigenous people of the North Pacific hunted sea otters. Although it appears that harvests periodically led to local reductions of sea otters (Simenstad et al. 1978), the species remained abundant throughout its range until the mid-1700s. Following the arrival in Alaska of Russian explorers in 1741, extensive commercial harvest of sea otters over the next 150 years resulted in the near extirpation of the species. When sea otters were afforded protection by the International Fur Seal Treaty in 1911, probably fewer than 2,000 animals remained in 13 remnant colonies (Kenyon 1969).

Although population regrowth began following legal protection, no remnant colonies of sea otters existed in southeast Alaska. As part of efforts to re-establish sea otters in portions of their historical range, otters from Amchitka Island and Prince William Sound were translocated to other areas (Jameson et al. 1982). These translocation efforts met with varying degrees of success. From 1965 to 1969, 412 otters (89 percent from Amchitka Island in southwest Alaska, and 11 percent from Prince William Sound in southcentral Alaska) were translocated to 6 sites in southeast Alaska (Jameson et al. 1982). In the first 20 years following translocation, these populations grew in numbers and expanded their range (Pitcher 1989).

Nearly all of the current population estimates for the southeast Alaska stock were developed using aerial survey methods of Bodkin and Udevitz (1999). The lone exception was a survey of the outer coastline from the western boundary of the stock at Cape Yakataga to Cape Spencer conducted by U.S. Geological Survey (USGS) in 2000 (N=32, CV=0.378). In 2002, USGS also surveyed Glacier Bay (N=1,266, CV=0.15) and the northern half of the southeast Alaska (N=1,838, CV=0.17; Bodkin and Esslinger 2006). The southern half was surveyed by USGS in 2003 (N=5,845; CV=0.14). In 2005, the U.S. Fish and Wildlife Service (Service) surveyed Yakutat Bay using the same methods (N=1,582; CV=0.33; Gill and Burn 2007). The most recent population estimates for the southeast Alaska stock are presented in Table 1.

Table 1. Population estimates for the southeast Alaska stock of northern sea otters. Previous stock assessment report (SAR) total is from August 2002.

Survey Area	Year	Unadjusted Estimate	Adjusted Estimate	CV	N _{MIN}	Reference
North Gulf of Alaska	2000	15	32	0.378	24	USGS unpublished data
Glacier Bay	2002		1,266	0.15	1,117	Bodkin and Esslinger (2006)
Northern Southeast Alaska	2002		1,838	0.17	1,594	Bodkin and Esslinger (2006)
Southern Southeast Alaska	2003		5,845	0.14	5,198	Bodkin and Esslinger (2006)
Yakutat Bay	2005		1,582	0.334	1,203	Gill and Burn (2007)
Current Total			10,563		9,136	
Previous SAR Total			12,632		9,266	

Glacier Bay was also surveyed as recently as 2006, with a resulting estimate of 2,785 sea otters (Bodkin and Esslinger 2006). The increase in sea otter abundance in Glacier Bay cannot be explained by reproduction alone, indicating that there has been substantial redistribution of sea otters in the past several years (Bodkin and Esslinger 2006). Therefore, to avoid double-counting of animals in both the Glacier Bay and northern southeast Alaska survey areas, we used the 2002 estimate for Glacier Bay, combined with adjusted estimates for the remainder of the stock, which results in a total estimate of 10,563 sea otters for the southeast Alaska stock.

Minimum Population Estimate

The minimum population estimate (N_{MIN}) for this stock is calculated using Equation 1 from the PBR Guidelines (Wade and Angliss 1997): $N_{MIN} = N/\exp(0.842 * [\ln(1+[CV(N)]^2)]^{1/2})$. The N_{MIN} for each survey area is presented in Table 1. The estimated N_{MIN} for the southeast Alaska stock is 9,136 sea otters.

Current Population Trend

Prior to the most recent aerial surveys, the trend for this stock of sea otters had been one of growth (Pitcher 1989, Agler et al. 1995). Comparing the current population estimate with that of the previous stock assessment report suggests that the southeast Alaska stock may not have continued to increase in abundance (USGS unpublished data). The comparison of abundance estimates is complicated by substantial differences in methods between the 1994 skiff survey of Agler et al. (1995) and the USGS aerial surveys; however, GIS analysis of the most recent surveys compared with original data from Pitcher (1989) indicates that range expansion from the outer coast to inner, protected waters has not occurred. The distribution of sea otters has changed; however, with substantial immigration into Glacier Bay in the past decade. In addition, residents of southeast Alaska also report changes in sea otter distribution, and consider the population to be healthy in their local areas.

Sea otter abundance in Yakutat Bay has also increased over the last decade, likely through reproduction, although some amount of immigration cannot be ruled out (Gill and Burn 2007). During this process, otters appear to have expanded their range to include the western shores of Yakutat Bay.

Although the estimated population size of this stock is lower than in the previous stock assessment report, due to improved precision in some of the estimates, the value for N_{MIN} is comparable. Therefore, the current population trend for the southeast Alaska stock is believed to be stable.

MAXIMUM NET PRODUCTIVITY RATE

Estes (1990) estimated a population growth rate of 17 to 20% per year for four northern sea otter populations expanding into unoccupied habitat. Although maximum productivity rates have not been measured through much of the sea otter's range in Alaska, in the absence of more detailed information, the rate of 20% calculated by Estes (1990)

is considered the best available estimate of R_{MAX} . There is insufficient information available to estimate the current net productivity rate for this population stock.

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act of 1972 (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $PBR = N_{MIN} \times 0.5 R_{MAX} \times F_R$. The recovery factor (F_R) for this stock is 1.0 (Wade and Angliss 1997) as population levels have remained stable with a known human take. Thus, for the southeast stock of sea otters, $PBR = 914$ animals ($9,136 \times 0.5(0.2) \times 1.0$).

ANNUAL HUMAN CAUSED MORTALITY

Fisheries Information

A complete list of fisheries and marine mammal interactions is published annually by NOAA-Fisheries, the most recent of which was published on November 27, 2007 (72 FR 66048). Although numerous fisheries exist within the range of the southeast Alaska stock of northern sea otters, none have been identified as interacting with this stock.

The estimated level of incidental mortality and serious injury of this stock can be estimated from fishery observer programs that monitor a portion of commercial fisheries in Alaska and report injury and mortality of marine mammals incidental to those operations. No incidents of sea otter incidental take have been observed in trawl, longline, or pot groundfish fisheries in southeast Alaska from 1989-2006 (Perez 2003; Perez 2006; Perez 2007).

An additional source of information on the number of sea otters killed or injured incidental to commercial fishery operations in Alaska are fisher self-reports required of vessel-owners by NOAA-Fisheries. From 1990 to 1993, self-reported fisheries data reflected no sea otter kills or injuries in southeast Alaska. Self-reports were incomplete for 1994 and not available for 1995 or 1996. Between 1997 and 2005, there were no records of incidental take of sea otters by commercial fisheries in this region. Credle et al. (1994) considered fisher self-reports to be a minimum estimate of incidental take as these data are most likely negatively biased.

The total fishery mortality and serious injury for the southeast Alaska stock of the northern sea otter has achieved a zero mortality and serious injury rate.

Oil and Gas Development

Exploration, development, and transport of oil and gas resources can adversely impact sea otters and nearshore coastal ecosystems in Alaska. Sea otters rely on air trapped in their fur for warmth and buoyancy. Contamination with oil drastically reduces the insulative value of the pelage, and consequently, sea otters are among the marine mammals most likely to be detrimentally affected by contact with oil. It is believed that sea otters can survive low levels of oil contamination (< 10% of body surface), but that greater levels (>25%) will lead to death (Costa and Kooyman 1981, Siniff et al. 1982). Vulnerability of sea otters to oiling was demonstrated by the 1989 Exxon Valdez oil spill in Prince William Sound. Total estimates of mortality for the Prince William Sound area vary from 750 (range 600 - 1,000) (Garshelis 1997) to 2,650 (range 500 - 5,000) (Garrot et al. 1993) otters. Statewide, it is estimated that 3,905 sea otters (range 1,904 - 11,257) died in Alaska as a result of the spill (DeGange et al. 1994). At present, abundance of sea otters in some oiled areas of Prince William Sound remains below pre-spill estimates, and evidence from ongoing studies suggests that sea otters and the nearshore ecosystem have not yet fully recovered from the spill (Bodkin et al. 2002, Stephensen et al. 2001).

There is currently no oil and gas development in southeast Alaska. Tankers carrying oil south from the Trans-Alaska Pipeline typically travel offshore and, therefore, pose a minimal risk to sea otters in southeast Alaska. As a result, no mortalities due to oil and gas development have been documented within the range of the southeast Alaska sea otter stock.

Subsistence/Native Harvest Information

The MMPA exempted Native Alaskans from the prohibition on hunting marine mammals, provided such taking was not wasteful. Alaska Natives are legally permitted to take sea otters for subsistence use or for creating and selling authentic handicrafts or clothing. Data for subsistence harvest of sea otters in southeast Alaska are collected by a mandatory Marking, Tagging and Reporting Program administered by the Service since 1988. Figure 2 provides a summary of harvest information for the southeast stock from 1989-2006. The mean reported annual subsistence take

during the past five complete calendar years (2002-2006) was 322 animals. Reported age composition during this period was 84% adults, 12% subadults, and 4% pups. Sex composition during the past five years was 70% males, 28% females, and 2% of unknown sex.

Research and Public Display

In the past five years, no sea otters were removed from the southeast Alaska stock for public display, nor were any sea otters captured and released for scientific research.

Other Factors

In August 2006, the Working Group on Marine Mammal Unusual Mortality Events reviewed information provided by the Service, and declared that a dramatic increase in sea otter strandings since 2002 constitutes an Unusual Mortality Event (UME) in accordance with Section 404 of the MMPA. The disease that typifies this UME is caused by a *Streptococcus infantarius* infection and has been observed over a broad geographic range in Alaska, including a few cases from southeast Alaska; however, the majority of cases have come from Kachemak Bay in the southcentral Alaska stock. It is not clear if the observed stranding pattern is representative of overall sea otter mortality, or an artifact of having a well-developed stranding network in the Kachemak Bay area. The Service will continue to work with the NOAA-Fisheries and Alaska SeaLife Center to develop the infrastructure for a statewide marine mammal stranding network in Alaska.




Figure 2. Reported subsistence harvest of northern sea otters from the southeast Alaska stock, 1989-2006.

STATUS OF STOCK

The level of direct human-caused mortality within the southeast Alaska stock does not exceed the PBR level, and the stock is neither listed as “depleted” under the MMPA or listed as “threatened” or “endangered” under the U. S. Endangered Species Act of 1973, as amended, nor is it likely to be listed as such in the foreseeable future. In addition, the level of human-caused mortality and serious injury is not likely to cause the stock to be reduced below its optimum sustainable population level. Therefore, the southeast Alaska stock of the northern sea otter is classified as non-strategic.

CITATIONS

- Agler, B. A., S. J. Kendall, P. E. Seiser, and J. R. Lindell. 1995. Estimates of marine bird and sea otter abundance in southeast Alaska during summer 1994. Migratory Bird Management, U.S. Fish and Wildlife Service, Anchorage, Alaska. 90pp.
- Bodkin, J. L., and M. S. Udevitz. 1999. An aerial survey method to estimate sea otter abundance. Pages 13-26 in G.W. Garner et al., editors. Marine Mammal Survey and Assessment Methods. Balekema, Rotterdam, Netherlands.
- Bodkin, J. L., B. E. Ballachey, T. A. Dean, A. K. Fukuyama, S. C. Jewett, L. M. McDonald, D. H. Monson, C. E. O’Clair, and G. R. VanBlaricom. 2002. Sea otter population status and the process of recovery from the Exxon Valdez spill. Marine Ecology Progress Series. 241:237-253.
- Bodkin, J.L. and G.E. Esslinger. 2006. Sea Otter Population Briefing, Southeast Alaska, 1969-2003; Data summary and initial interpretation. U.S. Geological Survey, Alaska Science Center report. 24pp.
- Costa, D. P., and G. L. Kooyman. 1981. Effects of oil contamination in the sea otter *Enhydra lutris*. Outer Continental Shelf Environmental Assessment Program. NOAA Final Report. La Jolla, California.
- Credle, V. A., D. P. DeMaster, M. M. Merlein, M. B. Hanson, W. A. Karp, and S. M. Fitzgerald (eds.). 1994. NMFS observer programs: minutes and recommendations from a workshop held in Galveston, Texas, November 10-11, 1993. U.S. Department of Commerce, NOAA Tech. Memo. NMFS-OPR-94-1. 96 pp.
- DeGange, A. R., A. M. Doroff, and D. H. Monson. 1994. Experimental recovery of sea otter carcasses at Kodiak Island, Alaska, following the Exxon Valdez oil spill. Marine Mammal Science 10:492-496.

- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conservation Biology* 6(1):24-36.
- Estes, J. A. 1990. Growth and equilibrium in sea otter populations. *Journal of Animal Ecology* 59:385-401.
- Garrott, R. A., L. L. Eberhard, and D. M. Burn. 1993. Mortality of sea otters in Prince William Sound following the Exxon Valdez oil spill. *Marine Mammal Science* 9:343-359.
- Garshelis, D. L., and J. A. Garshelis. 1984. Movements and management of sea otters in Alaska. *Journal of Wildlife Management* 48(3):665-678.
- Garshelis, D. L., A. M. Johnson, and J. A. Garshelis. 1984. Social organization of sea otters in Prince William Sound, Alaska. *Canadian Journal of Zoology* 62:2648-2658.
- Garshelis, D. L. 1997. Sea otter mortality estimated from carcasses collected after the Exxon Valdez oil spill. *Conservation Biology* 11(4): 905-916.
- Gill, V.A., and D.M. Burn. 2007. Aerial surveys of sea otters in Yakutat Bay, Alaska, 2005. U.S. Fish and Wildlife Service, Marine Mammals Management Office. Technical Report MMM 2007-01. 18pp.
- Gorbics, C. S., and J. L. Bodkin. 2001. Stock structure of sea otters (*Enhydra lutris kenyoni*) in Alaska. *Marine Mammal Science* 17(3): 632-647.
- Jameson, R.J., K.W. Kenyon, A.M. Johnson, and H.M. Wight. 1982. History and status of translocated sea otter populations in North America. *Wildlife Society Bulletin*. 10:100-107.
- Johnson, A. M. 1982. Status of Alaska sea otter populations and developing conflicts with fisheries. Pages 293-299 in: *Transactions of the 47th North American Wildlife and Natural Resources Conference*, Washington D.C.
- Kenyon, K. W. 1969. The sea otter in the eastern Pacific Ocean. *North American Fauna* 68. U.S. Department of the Interior, Washington D.C.
- Perez, M. A. 2003. Compilation of marine mammal incidental catch data for domestic and joint venture groundfish fisheries in the U.S. EEZ of the North Pacific, 1989-2001. NOAA Technical Memorandum NMFS-AFSC-138. 145 pp.
- Perez, M. A. 2006. Analysis of marine mammal bycatch data from the trawl, longline, and pot groundfish fisheries of Alaska, 1998-2004, defined by geographic area, gear type, and target groundfish catch species. NOAA Technical Memorandum NMFS-AFSC-167. 194 pp.
- Perez, M. A. 2007. Bycatch of marine mammals in the groundfish fisheries of Alaska, 2006. Alaska Fisheries Science Center Processed Draft Report. 67pp.
- Pitcher, K. W. 1989. Studies of southeastern Alaska sea otter populations: distribution, abundance, structure, range expansion and potential conflicts with shellfisheries. Anchorage, Alaska. Alaska Department of Fish and Game, Cooperative Agreement 14-16-0009-954 with U.S. Fish and Wildlife Service. 24 pp.
- Riedman, M. L., and J. A. Estes. 1990. The sea otter *Enhydra lutris*: behavior, ecology, and natural history. *Biological Report*; 90 (14). U.S. Fish and Wildlife Service.
- Simenstad, C. A., J. A. Estes, and K. W. Kenyon. 1978. Aleuts, sea otters, and alternate stable-state communities. *Science* 200:403-411. 127 pp.
- Siniff, D. B., T. D. Williams, A. M. Johnson, and D. L. Garshelis. 1982. Experiments on the response of sea otters *Enhydra lutris* to oil contamination. *Biological Conservation* 23: 261-272.
- Stephensen, S. W., D. B. Irons, S. J. Kendall, B. K. Lance, and L. L. MacDonald. 2001. Marine bird and sea otter population abundance of Prince William Sound, Alaska: trends following the T/V Exxon Valdez oil spill, 1989-2000. Restoration Project 00159 Annual Report. USFWS Migratory Bird Management, Anchorage, Alaska. 114 pp.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Department of Commerce, NOAA Technical Memo. NMFS-OPR-12. 93 pp.