

---

# **Annual Operating Plan**

**for  
Lower Snake River Fish and Wildlife Compensation Programs  
Grande Ronde and Imnaha Basins, Oregon**

**Spring Chinook Salmon  
Fall Chinook Salmon  
Coho Salmon  
Pacific Lamprey**

**For the Period of  
January 1 – December 31, 2025**

**Prepared by:  
Oregon Department of Fish and Wildlife  
Confederated Tribes of the Umatilla Indian Reservation  
Nez Perce Tribe**

**For  
Lower Snake River Compensation Plan  
USFWS  
and  
Bonneville Power Administration**

**Updated  
02-26-25**

## List of Figures, Tables, Sections

Contacts for Spring Chinook, Fall Chinook, and Coho Programs.....	3
Figure 1. Grande Ronde and Imnaha basin Chinook salmon hatchery facilities and release locations .....	4
Table 1. Grande Ronde and Imnaha basin Chinook salmon smolt release goals, including marking strategy.....	5
Figure 2. Grande Ronde and Imnaha basin Chinook salmon production timeline .....	6
Table 2. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2023 inventory, current as of 11/1/2024. Marks (CWT, PIT Tag, and AD) represent percentage or number of fish marked but does not account for tag loss and/or mortality. ....	7
Table 3. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2023 transport and release schedule in 2025. Inventory current as of 11/1/2024.....	9
Table 4. Brood year (BY) 2024 fry inventory with release goals and marking plan, current as of 11/1/2024. ....	10
Table 5. Pre-season return estimates for adult (age 4-5) Chinook salmon returning to Oregon Snake Basin tributaries in 2025. ....	11
Table 6. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2025 brood stock collection goals .....	12
Table 7. BY 2025 Upper Grande Ronde River weekly brood stock collection strategy, by sex and origin. (natural-origin, NOR, and hatchery-origin, HOR) Collection of 50% of the natural return and as many hatchery adults needed for brood number, up to 178 adults total .....	13
Table 8. BY 2025 Catherine Creek weekly brood stock collection strategy, by sex, and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon based on 36% natural in brood. Returns that are lower than anticipated in 2025 may necessitate a shift to collect and pass fish based on the maximum allowable under the sliding scale (40% of total natural and hatchery returns .....	14
Table 9. BY 2025 Lookingglass Creek weekly brood stock collection strategy, by sex and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon.....	15
Table 10. BY 2025 Lostine River weekly adult brood stock collection strategy, by sex and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon.....	16
Figure 3. BY 2025 broodstock collection schedule for Chinook salmon with mean PIT tag arrival at IR-4, and the range of trapping dates from 2016 to 2022. Total collections will include 282 adult spring Chinook salmon and 14 jacks. A detailed collection schedule will be developed when in-season projections are available.....	17
Table 11. Grande Ronde Fall Chinook salmon brood year 2024 (BY 24) inventory from Irrigon Fish Hatchery (IFH) current as of 11/1/24.....	18
Table 12. Grande Ronde basin Coho salmon brood year (BY) 2023 inventory, current as of 11/1/2024. ....	19
Table 13. Tentative release numbers and locations of lamprey in the Grande Ronde River basin in 2025. ....	20
<b>Standard Operating Procedures.....</b>	<b>21</b>
Table 14. Mark coordination for stocks at Lookingglass Hatchery. ....	26
<b>Grande Ronde and Snake River Fall Chinook Program .....</b>	<b>35</b>
<b>Grande Ronde Coho Salmon Program .....</b>	<b>36</b>

<b>Grande Ronde Lamprey / Freshwater Mussels.....</b>	<b>39</b>
<b>Freshwater Mussel Project Overview .....</b>	<b>40</b>
Table 1. Upper Grande Ronde River weir management guidelines for broodstock collection and passage .....	41
Table 2. Catherine Creek Spring Chinook broodstock/upstream passage management guidelines. ....	41
Table 3. Lookingglass Creek weir management guidelines for broodstock collection and passage. Current weir management strategy is consistent with Lookingglass Management Plan. Table is a placeholder until permanent agreement is in place.....	42
Table 4. Lostine River weir management guidelines for broodstock collection and passage. Assumes program goal of 250,000 smolts (166 adults for broodstock). ....	43
Table 5. Sliding scale management tool for Imnaha Chinook hatchery program utilized for managing disposition of Chinook salmon adults for broodstock and escapement to natural spawning areas (Table 1 excerpted from Imnaha spring Chinook HGMP May 2011). ....	44
Appendix A. Disinfections and Sanitation Guidelines for all LSRCP Hatcheries.....	45
Appendix B. Juvenile Chinook fish health monitoring plan and disease treatments.....	47
Appendix C. Adult Chinook fish health monitoring plan and disease treatments at Lookingglass Hatchery.....	48
Appendix D. Disease treatments and other drugs for adult Chinook Broodstock. ....	49
Appendix E. Standard Operating Procedures for Imnaha River Weir .....	51

## Contacts for Spring Chinook, Fall Chinook, and Coho Programs

Name	Agency	Phone	E-mail
<b>Operation and Maintenance</b>			
Mike McLean	CTUIR	541-429-7960	MikeMcLean@ctuir.org
Jen Krajcik	CTUIR	541-429-7278	JenniferKrajcik@ctuir.org
Becky Johnson	NPT	208-843-7320	Beckyj@nezperce.org
Rick Zollman	NPT	541-886-5364	Rickz@nezperce.org
Andrew Gibbs	ODFW	541-786-3114	Andrew.j.Gibbs@odfw.oregon.gov
Diane Deal	ODFW	541-437-9723	Diane.e.Deal@odfw.oregon.gov
Ron Harrod	ODFW	541-426-4467	Ron.l.Harrod@odfw.oregon.gov
Ace Trump	WDFW	509-382-1004	Ace.Trump@dfw.wa.gov
Alex Clark (Cascade Hatchery)	ODFW	541-374-8381	Alex.Clark@odfw.oregon.gov
Owen Davis	ODFW	541-963-2138	Owen.c.Davis@odfw.oregon.gov
Keenan Schmidt	ODFW	541-922-5732	Keenan.Schmidt@odfw.oregon.gov
Justin Herold	ODFW	541-426-4467	Justin.j.Herold@odfw.oregon.gov
Erika Brigante (Fish Health)	ODFW	541-962-3823	Erika.n.Brigante@odfw.oregon.gov
Sally Gee (Fish Health)	ODFW	541-962-3011	Sally.Gee@odfw.oregon.gov
<b>Monitoring &amp; Evaluation</b>			
Gene Shippentower	CTUIR	541-429-7287	GeneShippentower@ctuir.org
Les Naylor	CTUIR	541-429-7942	LesNaylor@ctuir.org
Carrie Crump	CTUIR	541-429-7945	CarrieCrump@ctuir.org
Joseph Feldhaus	ODFW	541-962-3724	Joseph.Feldhaus@odfw.oregon.gov
Pat Keniry	ODFW	541-962-3026	Patrick.j.Keniry@odfw.oregon.gov
Joseph Dittmer	ODFW	541-975-0805	Joseph.h.Dittmer@odfw.oregon.gov
Joe Lemanski	ODFW	541-975-4736	Joseph.r.Lemanski@odfw.oregon.gov
Jason Vogel	NPT	208-621-3602	Jasonv@nezperce.org
Bill Young	NPT	208-834-5290	Billy@nezperce.org
Brian Simmons	NPT	541-432-2515	brians@nezperce.org
Neal Espinosa	NPT	541-432-2502	neale@nezperce.org
Ryan Rumelhart	NPT	541-432-2512	ryanr@nezperce.org
Clark Watry	NPT	208-621-3549	clarkw@nezperce.org
Stuart Rosenberger	IPC	208-388-6121	Srosenberger@idahopower.com
Scott Putnam	IDFG	208-750-4243	Scott.Putnam@idfg.idaho.gov
<b>Fisheries Management</b>			
Preston Bronson	CTUIR	541-429-7277	PrestonBronson@ctuir.org
Jack Yearout	NPT	208-621-4638	Jacky@nezperce.org
Jeff Yanke	ODFW	541-318-7927	Jeff.Yanke@odfw.oregon.gov
Ethan Brandt	ODFW	541-805-0856	ethan.j.brandt@odfw.oregon.gov
Tyler Hoyt	ODFW	541-975-8606	tyler.j.hoyt@odfw.oregon.gov
Kyle Bratcher	ODFW	541-318-7928	Kyle.w.Bratcher@odfw.oregon.gov
Michael Lance	ODFW	541-318-7930	Michael.j.Lance@odfw.oregon.gov
<b>Federal</b>			
Nate Wiese	LSRCP	208-296-8759	Nathan_Wiese@fws.gov
Chris Starr	LSRCP	208-378-5329	Chris_Starr@fws.gov
Rod Engle	LSRCP	208-278-5298	Rod_Engle@fws.gov
Greg Burak	LSRCP	208-986-1120	Greg_Burak@fws.gov
Brett Farman	NOAA	503-231-6222	Brett.Farman@noaa.gov




Figure 1. Grande Ronde and Imnaha basin Chinook salmon hatchery facilities and release locations.

Table 1. Grande Ronde and Imnaha basin Chinook salmon smolt release goals, including marking strategy.

Stock	Smolt release goal	Acclimation Period		Direct Release	Ad clipped (%)	CWT (N)	CWT (%)	PIT Tags	
		First	Second					N	Source <sup>a</sup>
Upper Grande Ronde R.	250,000	125,000		125,000	50	250,000	100	2,600	LSRCP
Catherine Creek	150,000	150,000			100	100,000	67	21,000	CSS
Lookingglass Creek	250,000	250,000			100	120,000	48	5,000	LSRCP
Lostine River	250,000	125,000	125,000		100	126,000	50	6,000	LSRCP
Grande Ronde Basin	900,000					596,000		35,000	
Imnaha River	490,000	280,000		210,000	100	245,000	50	21,000	CSS
<b>Oregon Snake Basin</b>	<b>1,390,000</b>	<b>930,000</b>	<b>250,000</b>	<b>335,000</b>		<b>841,000</b>		<b>56,000</b>	

<sup>a</sup> Source = funding source for allocation of PIT tags: LSRCP (Lower Snake River Compensation Plan), CSS (Comparative Survival Study – Fish Passage Center).




Figure 2. Grande Ronde and Imnaha basin Chinook salmon production timeline.

Table 2. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2023 inventory, current as of 11/1/2024. Marks (CWT, PIT Tag, and AD) represent percentage or number of fish marked but does not account for tag loss and/or mortality.

Release Site	Pre-Split	Current Inventory		% CWT	CWT Codes	PIT Tag (N)	AD (%)
	Container	Container	N				
Upper Grande Ronde R. (GQ)	RW15	RW15	54,904	100	09-25-96	700	100
	RW16	RW16	56,606	100	09-25-95	700	100
	RW17	RW17	53,299	100	09-24-94	600	0
	RW18	RW18	55,516	100	09-24-93	600	0
<b>Upper Grande Ronde Total</b>			<b>220,325</b>			<b>2,600</b>	
Catherine Cr. (CQ)	RW13	RW13	79,413	66	09-00-94	10,500	100
	RW14	RW14	77,522	68	09-00-95	10,500	100
<b>Catherine Cr. Total</b>			<b>156,935</b>			<b>21,000</b>	
Lookingglass Cr. (LG)	RW1	RW1	67,190	47	09-25-65	1,600	100
		AHPA	51,675	46	09-25-64	900	100
		AHPB	51,622	46	09-25-64	800	100
	RW6	AHPC	51,575	46	09-25-64	900	100
		AHPD	51,630	47	09-25-65	800	100
<b>Lookingglass Cr. Total</b>			<b>273,692</b>			<b>5,000</b>	
Lostine River (LQ)	RW2	RW2	66,085	52	09-25-66	1,500	100
	RW3	RW3	73,463	46	09-25-67	1,500	100
	RW4	RW4	68,510	50	09-25-68	1,500	100
	RW5	RW5	65,453	51	09-25-69	1,500	100
<b>Lostine River Total</b>			<b>273,511</b>			<b>6,000</b>	

<sup>a</sup> Discussions ongoing to discontinue CWT only groups and Ad clip those groups in 2024.



Table 2 (cont.). Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2023 inventory, current as of 11/1/2024. Marks (CWT, PIT Tag, and AD) represent percentage or number of fish marked but does not account for tag loss and/or mortality.

Release Site	Pre-Split	Current Inventory		% CWT	CWT Codes	PIT Tag (N)	AD (%)
	Container	Container	N				
Imnaha River (IM)	RW 7& RW8	RW6	71,531	0	—	3,000	100
	RW7	RW7	71,487	0	—	3,000	100
	RW8	RW8	71,492	0	—	3,000	100
	RW9	RW9	69,943	92	09-24-97	3,000	100
	RW10	RW10	71,006	91	09-24-98	3,000	100
	RW11	RW11	71,978	90	09-24-99	3,000	100
	RW12	RW12	70,048	95	09-25-00	3,000	100
<b>Imnaha River Total</b>			<b>497,485</b>			<b>21,000</b>	

Table 3. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2023 transport and release schedule in 2025. Inventory current as of 11/1/2024.

<b>Stock</b>	<b>Transfer Date</b>	<b>Release Type</b>	<b>Container</b>	<b>To</b>	<b>N</b>	<b>Estimated Pounds</b>	<b>Release Date</b>
Grande Ronde R.	7-Apr	1 <sup>st</sup> Direct	RW 15, 16	UGR Acc.	108,813	4,946	9-Apr
Grande Ronde R.	11-April	2 <sup>nd</sup> Acclimation	RW 17, 18	UGR Acc.	111,509	5,069	24-Apr
<b>Grande Ronde Total</b>					<b>220,322</b>		
Catherine Creek	7-April	Acclimated	RW 13, 14	Catherine Acc.	156,932	7,133	21-Apr
<b>Catherine Total</b>					<b>156,932</b>		
Lookingglass Creek		Acclimated	RW1	Lookingglass Cr.	X	X	11-Apr
Lookingglass Creek		Force Out	A, B, C, D	Lookingglass Cr.	X	X	17-Mar
<b>Lookingglass Total</b>					<b>273,674</b>	13,684	
Lostine River	19-Mar	1 <sup>st</sup> Acclimation	RW 2, 3	Lostine R. Acc.	139,524	5,581	10-Apr
Lostine River	11-Apr	2 <sup>nd</sup> Acclimation	RW 4, 5	Lostine R. Acc.	133,928	5,357	23-Apr
<b>Lostine Total</b>					<b>273,452</b>		
Imnaha River	31-Mar	1 <sup>st</sup> Acclimation	RW 6, 7, 11, 12	Imnaha R. Acc.	285,044	14,252	4-Apr
Imnaha River	10-Apr	2 <sup>nd</sup> Acclimation	RW 8, 9, 10	Imnaha R. Acc	212,441	10,621	15-Apr
<b>Imnaha Total</b>					<b>497,485</b>		

Table 4. Brood year (BY) 2024 fry inventory with release goals and marking plan, current as of 11/1/2024.

<b>Stock</b>	<b>Stock-BY ID</b>	<b>Estimated Fry</b>	<b>Release Goal</b>	<b>% Ad Clip</b>	<b>% CWT</b>	<b>PIT Tag (N)</b>
Upper Grande Ronde	GQ 80-23	242,938	250,000	50	100	3,000
Catherine Creek	CQ 201-23	156,166	150,000	100	68	21,000
Lookingglass Creek	LG 81-23	224,940	250,000	100	48	5,000
Lostine River	LQ 200-23	288,305	250,000	100	45	6,000
Imnaha River	IM 29-23	461,420	490,000	100	61	21,000
<b>Total Fry to Pond</b>		<b>1,373,769</b>	<b>1,390,000</b>			<b>56,000</b>

Post Shock/Pick numbers

Estimated eggs/female pre-shock – Upper Grande Ronde, 3,347 – Catherine Creek, 3,494 – Lookingglass Creek, 3,612 – Lostine River, 4,219 – Imnaha River, 4,084

Table 5. Pre-season return estimates for adult (age 4-5) Chinook salmon returning to Oregon Snake Basin tributaries in 2025.

<b>Stock</b>	<b>Origin</b>	<b>BON</b>	<b>Conversion to LGD</b>	<b>LGD</b>	<b>Conversion to Tributary</b>	<b>Tributary Mouth</b>	<b>Conversion from BON to Tributary</b>
Upper Grande Ronde	Hatchery	698	77%	537	90%	<b>484</b>	69%
	Natural	121	86%	104	90%	<b>94</b>	77%
Catherine Creek	Hatchery	896	77%	690	90%	<b>621</b>	69%
	Natural	267	86%	230	90%	<b>207</b>	77%
Lookingglass Creek	Hatchery	721	77%	555	90%	<b>499</b>	69%
	Natural	119	86%	103	90%	<b>93</b>	77%
Lostine River	Hatchery	1,145	75%	859	90%	<b>773</b>	68%
	Natural	583	80%	466	90%	<b>420</b>	72%
<b>Grande Ronde Total</b>	Hatchery	3,460		2,641		<b>2,377</b>	
	Natural	1,090		903		<b>814</b>	
Imnaha River	Hatchery	1,630	75%	1,223	90%	<b>1,102</b>	68%
	Natural	669	80%	535	90%	<b>482</b>	72%
<b>Oregon Snake Basin</b>	Hatchery	5,090		3,864		<b>3,479</b>	
	Natural	1,759		1,438		<b>1,296</b>	

Table 6. Grande Ronde and Imnaha basin Chinook salmon brood year (BY) 2025 brood stock collection goals.

Stock	Goals <sup>a</sup>			Fecundity Estimate (eggs per female) <sup>b</sup>	% Planned Survival (green egg to smolt) <sup>b</sup>
	Female Collection	Females Spawned	Green Eggs Collected <sup>b</sup>		
Upper Grande Ronde (GQ)	100	81	249,215	3,091	100
Catherine Creek (CQ)	50	49	171,656	3,512	87
Lookingglass Creek (LG)	89	79	264,974	3,367	94
Lostine River (LQ)	82	72	287,277	3,966	87
<b>Grande Ronde Total</b>	<b>310</b>	<b>281</b>	<b>1,023,248</b>		
Imnaha (IM)	141	134	576,471	4,060	85
<b>Oregon Snake Basin Tributaries</b>	<b>451</b>	<b>415</b>	<b>1,599,719</b>		

<sup>a</sup> Subject to change based on in-season run projections.

<sup>b</sup> Fecundity, planned survival, and green egg collection goals reflect most recent 3-year estimates, but will not match females spawned. ODFW may collect more females if broodstock collection indicates smaller females.

Table 7. BY 2025 Upper Grande Ronde River weekly brood stock collection strategy, by sex and origin. (natural-origin, NOR, and hatchery-origin, HOR). Collection of 50% of the natural return and as many hatchery adults needed for brood number, up to 178 adults total.

Week	Est. NOR Timing (%)	Females		Males		Jack	
		NOR	HOR	NOR	HOR	NOR	HOR
4-May	0.0	0	0	0	0	0	0
11-May	0.0	0	0	0	0	0	0
18-May	0.1	0	0	0	0	0	0
25-May	8.9	2	7	2	5	0	0
1-Jun	38.1	11	28	11	20	2	2
8-Jun	14.3	4	10	4	7	1	1
15-Jun	11.2	3	8	3	6	1	0
22-Jun	23.1	6	17	6	12	1	1
29-Jun	4.3	1	3	1	2	0	0
6-Jul	0.0	0	0	0	0	0	0
13-Jul	0.0	0	0	0	0	0	0
20-Jul	0.0	0	0	0	0	0	0
27-Jul	0.0	0	0	0	0	0	0
3-Aug	0.0	0	0	0	0	0	0
<b>Total Goal</b>		27	73	27	51	5	4

Table 8. BY 2025 Catherine Creek weekly brood stock collection strategy, by sex, and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon based on 36% natural in brood. Returns that are lower than anticipated in 2025 may necessitate a shift to collect and pass fish based on the maximum allowable under the sliding scale (40% of total natural and hatchery returns).

Week	Est. NOR Timing (%)	Females		Males		Jack	
		NOR	HOR	NOR	HOR	NOR	HOR
4-May	0.1	0	0	0	0	0	0
11-May	0.0	0	0	0	0	0	0
18-May	0.1	0	0	0	0	0	0
25-May	8.1	1	3	1	2	0	0
1-Jun	24.7	4	8	4	7	2	1
8-Jun	20.3	4	7	4	5	1	1
15-Jun	14.5	3	5	3	4	0	0
22-Jun	24.7	5	8	5	7	0	0
29-Jun	5.6	1	1	1	2	0	0
6-Jul	1.5	0	0	0	0	0	0
13-Jul	0.4	0	0	0	0	0	0
20-Jul	0.1	0	0	0	0	0	0
27-Jul	0.0	0	0	0	0	0	0
3-Aug	0.0	0	0	0	0	0	0
<b>Total Goal</b>		18	32	18	27	3	2

Table 9. BY 2025 Lookingglass Creek weekly brood stock collection strategy, by sex and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon.

Week	Est. NOR Timing (%)	Females		Males		Jack	
		NOR	HOR	NOR	HOR	NOR	HOR
4-May	0.1	0	0	0	0	0	0
11-May	0.6	0	1	0	0	0	0
18-May	1.9	0	2	0	1	0	0
25-May	4.3	0	4	0	3	0	0
1-Jun	11.3	1	9	1	8	0	1
8-Jun	16.4	1	14	1	13	0	2
15-Jun	19.0	1	16	1	15	0	2
22-Jun	16.1	1	13	1	13	0	2
29-Jun	15.8	1	13	1	12	0	1
6-Jul	6.9	0	6	0	5	0	1
13-Jul	3.3	0	3	0	2	0	0
20-Jul	1.5	0	1	0	1	0	0
27-Jul	1.4	0	1	0	1	0	0
3-Aug	0.5	0	0	0	0	0	0
10-Aug	0.9	0	1	0	1	0	0
Total Goal		5	84	5	75	0	9



Table 10. BY 2025 Lostine River weekly adult brood stock collection strategy, by sex and origin (natural-origin, NOR, and hatchery-origin, HOR) for Chinook salmon.

Week	Est. NOR Timing (%)	Females		Males		Jacks	
		NOR	HOR	NOR	HOR	NOR	HOR
<1-Jun	0.0%	0	0	0	0	0	0
8-Jun	0.4%	0	0	0	0	0	0
15-Jun	3.9%	1	2	1	2	0	0
22-Jun	6.2%	2	3	1	3	0	0
29-Jun	6.3%	2	4	1	3	0	0
6-Jul	23.8%	6	14	6	13	0	1
13-Jul	27.2%	7	15	7	15	0	2
20-Jul	9.2%	2	5	2	5	0	2
27-Jul	3.3%	1	2	1	2	0	0
3-Aug	1.3%	0	1	0	1	0	0
10-Aug	1.0%	0	1	0	1	0	0
17-Aug	1.3%	0	1	0	1	0	0
24-Aug	3.4%	1	2	1	2	0	0
31-Aug	8.5%	2	5	2	5	0	1
7-Sep	4.0%	1	2	1	2	0	0
<b>Total Goal</b>		<b>25</b>	<b>57</b>	<b>23</b>	<b>55</b>	<b>0</b>	<b>6</b>

Figure 3. BY 2025 broodstock collection schedule for Chinook salmon with mean PIT tag arrival at IR-4, and the range of trapping dates from 2016 to 2022. Total collections will include 282 adult spring Chinook salmon and 14 jacks. A detailed collection schedule will be developed when in-season projections are available.




Table 11. Grande Ronde Fall Chinook salmon brood year 2024 (BY 24) inventory from Irrigon Fish Hatchery (IFH) current as of 11/1/24

<b>Release Site</b>	<b>Transfer Date</b>	<b>Scheduled Release</b>	<b>Release Type</b>	<b>Stock</b>	<b>Expected at Release</b>	<b>Current Inventory</b>	<b>CWT (%)</b>	<b>PIT (N)</b>	<b>AD (%)</b>
Big Canyon Acc. Release	End April	Mid-May	Acclimated	IFH (97H)	500,000	500,000	20	4,500	100

Table 12. Grande Ronde basin Coho salmon brood year (BY) 2023 inventory, current as of 11/1/2024.

Release Site	Cascade Raceway	Release Type	N	Fish Size (fpp)	Marks			Release Date
					CWT (%)	PIT (N)	AD (%)	
Wallowa River	17	Direct	76,549	20.0	0.0	4000	100	24-25 March
Wallowa River	18	Direct	76,761	20.0	0.0	–	100	24-25 March
Wallowa River	19	Direct	76,516	20.0	0.0	–	100	24-25 March
Wallowa River	20	Direct	76,534	20.0	0.0	–	100	24-25 March
Wallowa River	21	Direct	76,582	20.0	51.0	–	100	24-25 March
Wallowa River & Hurricane Cr.	22	Direct	76,546	20.0	84.4	–	100	24-25 March
Hurricane Cr.	23	Direct	76,540	20.0	0.0	1000	100	24-25 March
<b>Lostine River Total</b>			<b>536,028</b>					

Table 13. Tentative release numbers and locations of lamprey in the Grande Ronde River basin in 2025.

Program / Location	Life Stage	Estimated Release	Release Timing
<b>Confederated Tribes of the Umatilla Indian Reservation (CTUIR)</b>			
Upper Grande Ronde River	Adult	300	Fall 2025
Catherine Creek	Adult	350	Fall 2025
Lookingglass Creek	Adult	600	Fall 2025
Lower Grande Ronde	Adult	300	April/May 2025
Limber Jim Creek	Adult	50	Fall 2025
Indian Creek	Adult	100	Fall 2025
Sheep and McCoy Creeks	Adult	0	--
Wenaha River	Adult	800	Summer/Fall 2025
Five Points Creek	Adult	0	--
<b>CTUIR Total</b>		<b>2500</b>	
<b>Nez Perce Tribe (NPT)</b>			
Wallowa River	Adult	25	April-May 2025
Minam River	Adult	25	April-May 2025
Joseph Creek <sup>a</sup>	Adult	50	Sept-Oct 2025
<b>NPT Total</b>		<b>100</b>	

<sup>a</sup> Lamprey are released at three locations- at the confluence of Chesnimnus Creek and Joseph Creek, in the Wallowa River (at Minam boat ramp), and in the Minam River (at the old ODFW Dingle-Johnson site).

**Standard Operating Procedures**  
**for**  
**Lower Snake River Fish and Wildlife Compensation Programs**  
**Grande Ronde and Imnaha Basins, Oregon**

**Spring Chinook Salmon**  
**Fall Chinook Salmon**  
**Coho Salmon**  
**Pacific Lamprey / Freshwater Mussels**

**For the Period of**  
**2023-2028**

**Prepared by:**  
**Oregon Department of Fish and Wildlife**  
**Confederated Tribes of the Umatilla Indian Reservation**  
**Nez Perce Tribe**

**For**  
**Lower Snake River Compensation Plan**  
**USFWS Administration**  
**and**  
**Bonneville Power Administration**

**Updated**  
**11-16-24**

## Grande Ronde and Imnaha Spring/Summer Chinook

### I. Production Goals

- A. Consistent with production tables specified in the U.S. vs. Oregon fish management agreement, production objectives for each tributary is described below. All production is conventional. Release target size is 20-25 fpp.
1. Upper Grande Ronde = 250,000
  2. Catherine Creek = 150,000
  3. Lookingglass Creek = 250,000
  4. Lostine River = 250,000
  5. Imnaha River = 490,000

### II. Smolt Releases

#### A. Key Contacts

1. Grande Ronde - Hatcheries: CTUIR (McLean, Krajcik), ODFW (Deal, Gibson, Brandt, Davis), NPT (Zollman). Fish Research: CTUIR (Naylor, Crump) ODFW (Keniry, Feldhaus, Gibson, Lemanski, Brandt), NPT (Vogel, Simmons, Young), IDFG (Putnam), LSRCF (Starr, Engle, Burak), and NOAA (Farman).
2. Imnaha - NPT (B. Johnson, Rumelhart, Simmons, Young, Vogel), ODFW (Feldhaus, Keniry, Gibson, Bratcher), CTUIR (McLean, Krajcik), LSRCF (Starr, Engle, Burak), NOAA (Farman), IDFG (Putnam).

#### B. Acclimation facilities - All facilities should be set-up and operational at least 2 days prior to scheduled delivery of smolts.

1. Release numbers = Last physical inventory – mortalities.
2. Operators report final numbers to the ODFW LGH staff or Shari Beals.
3. Scan all acclimation mortalities for PIT tags. PIT tag data provided to NPT (Simmons) and ODFW (Keniry). Mortalities should be provided to fish health for examination.

#### C. When fish are transported, liberation staff will record the water temperature of the contributing facility or waterbody, the receiving facility or waterbody, and the transportation tank when fish are loaded and again when fish are released.

Please record this data on the fish liberation reports.

4. The difference in temperatures between the liberation tank and target water body should not exceed 10 °F (5.6 °C). If temperature range is >10 °F (5.6 °C), water from planting site should be pumped into the top of the tank while drafting water out of the bottom. After water in the tank has reached the correct temperature, wait for at least 30 minutes to allow fish to acclimate themselves to the temperature change before releasing them.

#### D. Contingencies – Under extreme conditions, smolts may be released earlier than scheduled. In those cases, notify downstream rotary trap operators immediately.

1. Grande Ronde Traps –Pat Keniry, Polly Gibson, Mike McLean, Les Naylor, Carrie Crump, Scott Putnam.

2. Imnaha Traps –Brian Simmons, Jason Vogel, Ryan Rumelhart, and Scott Putnam.
3. Notify co-managers, LSRCP, and NOAA within 24 hours.

**E. Site-specific scheduling**

1. Upper Grande Ronde River Acclimation
  - a. Fish are split equally into 4 containers.
  - b. One acclimated release – all CWT only.
  - c. One direct release - all Ad-CWT.
2. Catherine Creek Acclimation
  - a. Fish are split equally into 4 containers.
  - b. One acclimation phase.
3. Lookingglass Creek Fish Hatchery
  - a. Fish will start to be volitionally released on April 1 and forced out on April 15.
4. Lostine River Acclimation
  - a. Smolts are allowed to volitionally release after 7-10 days of acclimation before forced release.
  - b. Fish are split evenly into 4 containers.
  - c. Two releases – early and late (half of production each).
5. Imnaha River Acclimation
  - a. Three raceways (210,000 smolts) will be acclimated and released during the first acclimation and four raceways (280,000 smolts) will be acclimated and released during the second acclimation period.
  - b. Two releases –Fish will be acclimated in two acclimations, each one being 5-6 days then forced out with the second group coming in for acclimation immediately after. The dates are subject to change but are generated through barging start dates and travel time to Lower Granite Dam.

### **III. Adult Trapping**

**A. Key Contacts – Adult Trapping**

1. Transportation - Facility operators (NPT and CTUIR) coordinates all hauling and notifies LGH (Deal) of the stock, number being hauled, and estimated arrival time.
2. Communications – Operators should distribute summaries of collections and operations to co-manager regularly (weekly or bi-monthly). Wallowa Hatchery provides a summary of fish provided for distribution/food bank (Lostine and Imnaha stocks).
  - a. All non-native fish encountered in a trap will be dispatched, kept, reported, and given to the district fish biologists and fish health for record keeping.



- b. On Lookingglass Creek scan non-target species for PIT tags and record external marks and length. Report data to district fish biologists.

**B. Grande Ronde Basin trapping guidelines**

1. While in operation, trapping facilities are checked daily.
2. Captured fish will be collected for transport or released at the time they are first handled. On Catherine Creek and the Upper Grande Ronde, if more than 15 adults are in the trap then fish will be collected and transported or passed upstream that day. If less than 15 fish are in the trap, fish may be held up to 72 hours.
3. Monitor water temperatures, and adjust schedule to best coincide their work with the coolest water temperatures.
4. Operators may conduct walking surveys, making visual observations of fish below weir sites to anticipate trapping numbers. Surveys may include snorkeling.
5. Bull Trout and Lamprey may be encountered at trapping facilities. If captured; enumerate, measure (Bull Trout only), Lookingglass & Lostine only: take a genetic sample (Bull Trout only), and pass in the intended direction it was trapped. For Lamprey, notify contacts of any observations (*see* Section 4.A. Lamprey Overview).

**C. Imnaha trapping guidelines**

1. The Imnaha satellite facility is staffed 24 hours per day, 7 days per week, while operational.
2. After the first Chinook is captured, trap is processed daily through the last week of July.
3. After July, trap is processed during weekdays. Exception is if 10 or more natural Chinook salmon, or 30 or more total fish or one bull trout are estimated in the trap on Saturday or Sunday, the trap will be processed on that day.
4. Broodstock is collected on Monday and Thursday of each week, or as needed. Extra broodstock can be collected if there is a deficit in brood numbers the prior week.
5. On all other days collected fish are passed above, held for broodstock in pen, recycled, or transported to Wallowa Hatchery.
6. Recaptured recycle fish can be recycled multiple times with an additional punch and a preferred release location above the sport fishery.
7. Refer to Appendix E; Standard Operating Procedures for Imnaha River Weir, for further details.

**D. Trap installation and scheduling**

1. Upper Grande Ronde and Catherine Creek
  - a. Lead agency CTUIR.
  - b. Weir will be installed when flows are lowest (March), but the trap will not become operational until chinook are present through late-July if river conditions allow. Weir installed early March on Catherine Creek. The traps are typically staffed overnight after mid-April.

- c. The Upper Grande Ronde weir is removed when the daily maximum water temperature exceeds 68° F (20° C).
- d. UGR brood: 1 ROP Hat, 2ROP Wild + PIT tag
- e. CC brood: 2 LOP + PIT tag
- f. Outplants to LGC and Indian Creek will get 3 LOP.

## 2. Lookingglass Creek

- a. Lead agency ODFW.
- b. The intake trap at Lookingglass Hatchery is operated from early March (or as conditions allow) until the last spawning survey has been conducted (typically mid-September), or no later than September 30. From September 30 to March 1, the ladder will be operational with the fyke removed to allow for year-round volitional passage.
- c. Lower ladder will be operated to collect broodstock but will not be functional until subsistence fisheries have ended.
- d. Operators should note when pickets are removed and if/how many adults/jacks are flushed downstream. When pickets are pulled, operators will notify CTUIR personnel.
- e. LGC brood handling:
- f. Brood = 1LOP + PIT tag
- g. Fish passed upstream 1 ROP save punch for genetic sample, CTUIR will PIT tag all fish passed upstream and radio tag.
- h. Fishery recycled downstream: 2 ROP save punch for genetic sample.
- i. Recaps in general do not need to be marked multiple times. Just record data, record as recap, and pass in the direction they need to go.
- j. Recaps from fishery recycle (2 ROP):
- k. Passed upstream: After fishery, add 1ROP save for genetics
- l. Passed downstream: During fishery, no extra punch needed.
- m. Recaps from above (1 ROP): Always pass upstream, no extra mark needed.
- n. Recaps from CC outplants (3 LOP):
- o. A differential mark for ladder of capture will no longer be given. If above weir 1 ROP, below weir 2ROP.
- p. After fishery, pass all fish upstream except hatchery jacks.
- q. UGR strays captured will get 3 ROP and a zip tie.

Table 14. Mark coordination for stocks at Lookingglass Hatchery.

Disposition	Lookingglass Cr.	Catherine Cr	U. Grande Ronde	Imnaha
Passed upstream	1 ROP	1 ROP	1 ROP	1 ROP
Passed downstream	2 ROP	3 LOP	--	2 LOP (add 1 LOP if recycled again)
Broodstock	1 LOP + PIT tag	2 LOP + PIT tag	1 ROP h, 2 ROP w + PIT tag	No Punch
Strays to other stocks	3 ROP zip + PIT tag	--	--	--
<b>LGC Recaptures</b>	<b>Pass Upstream</b>	<b>Pass Downstream</b>		
1 ROP	No other mark	Don't do it.		
2 ROP	After fishery, add 1 ROP (for genetics) Makes 3 ROP	During fishery, no other mark		
3 LOP	After fishery add 1 LOP (for genetics) Makes 4 LOP	During fishery, no other mark		

### 3. Lostine River

- a. Lead agency NPT.
- b. Trapping begins mid-May for Chinook monitoring. The trap will operate until no Chinook are captured for 10 days after 1-Sep. During the pilot phase of the Grande Ronde Coho Salmon program, trapping will continue until mid-December or when icing prevents operation.

### 4. Imnaha River

- a. Lead agency ODFW.
- b. Trap is installed in early June, or as soon as river conditions allow, and operated until after the last spawning ground survey which occurs mid-September.
- c. Staffing guidelines:
  - i. Weekdays - Two (2) ODFW staff each day. NPT provides one (1) staff that will travel on site each day.
  - ii. Weekends - ODFW provide two (2) staff daily (one Lookingglass Hatchery staff stays on-site, another travels to the site on the weekend if needed). NPT provides one (1) staff that will travel on site each day. NPT hauls recycles and/or outplants. Any surplus fish beyond the capacity of the NPT tanker can be placed in a holding pen. Alternatively, one hatchery staff may haul fish.

E. Site-specific weir management guidelines

5. Upper Grande Ronde River - The Grande Ronde conventional program calls for collection of up to 50% of natural fish and up to 100% of conventional return to reach the broodstock goal (SOP Table 1).
  - a. Fish are handled without anesthesia.
  - b. Mark all hatchery and natural adults released upstream of the weir with a right opercle punch (1-ROP).
  - c. Collect scale samples from all natural-origin fish passed upstream.
  - d. Inject broodstock with erythromycin (or tulathromycin), oxytetracycline, and then transport to the adult holding ponds (see vii. Adult holding Ponds). PIT tag broodstock.
6. Catherine Creek – Pre-season run projections and the sliding scale (AOP Table 5 SOP Table 2) are used to determine brood collection and natural spawning objectives and may be reassessed with PIT tag data during the run.
  - a. Surplus - If surplus hatchery fish are available, the first 50 should be outplanted to Indian Creek (3 LOP), then 100 to Lookingglass Creek below the weir (3 LOP), then 50 more to Indian Creek (3 LOP), and remainder to Lookingglass below the weir (3 LOP). All fish over 800 mm will be used for brood or upstream passage only.
  - b. Fish are handled using electro narcosis.
  - c. Mark all hatchery and natural adults released upstream of the weir with a right opercle punch (1 ROP).
  - d. Collect scale samples from all natural origin fish passed upstream.
  - e. Inject broodstock with erythromycin (or tulathromycin) and oxytetracycline and transport to the adult holding ponds (see vii. Adult holding Ponds). PIT tag broodstock.
7. Lookingglass Creek – Contingent on annual approvals from NOAA Fisheries, disposition of Lookingglass Creek adults trapped at either the LGH intake weir or lower ladder will occur as follows:
  - a. At run size projections <400 adults (to the mouth of Lookingglass Creek): Balance broodstock collections with upstream escapement. Manage a 50:50 escapement to brood pass:keep ratio on a weekly basis.
  - b. At run size projections >400 adults: Broodstock collections follow the schedule outlined in Table 9. All adults not taken for brood are passed upstream.
  - c. At run size projections >750: Recycling of hatchery-origin adults back into the fishery will also be considered, with details specified in the AOP.
  - d. Disposition of trapped fish before and after 4-Jul follows guidelines provided in the Lookingglass Creek management plan.
  - e. Fish are handled without anesthesia.
  - f. All hatchery and natural adults released upstream of the weir captured at the upper or lower ladder will be marked with a right opercle punch (1-ROP).

- g. Genetic samples are collected from all adults captured. CTUIR will PIT tag all fish passed upstream and radio tag some. Additionally, scale samples are collected on all wild fish passed upstream.
  - h. Inject broodstock with erythromycin (or tulathromycin) and oxytetracycline, PIT tag, and transport to the adult holding ponds (*see vii. Adult holding Ponds*). All fish that are not injected with erythromycin, such as those that are received after August 1<sup>st</sup>, will be given 3 caudal fin punches.
  - i. Release all natural origin jacks upstream.
  - j. Do not release hatchery jacks upstream. Sacrifice all hatchery origin jacks for tag recoveries and provide carcasses for tribal distribution, foodbank, or bury. Hatchery jacks without CWT can be provided for distribution, foodbank, buried, or recycled below the weir on Lookingglass Creek.
  - k. Do not release identifiable out-of-basin (within or outside Grande Ronde basin) stray salmon upstream of the weir.
    - iii. Strays from the Upper Grande Ronde River trapped at Lookingglass receive a 3 ROP, PIT tag, and bright colored dorsal ziptie and added to the Grande Ronde broodstock or held for return to the river.
    - iv. Catherine Creek strays can be recycled below the weir on Lookingglass Creek, used for Catherine Creek brood or, if needed Lookingglass brood.
  - l. Additional guidance can be found in the Lookingglass Creek Spring Chinook Management Plan.
8. Lostine River - Pre-season run projections and the sliding scale (AOP Table 5 SOP Table 4) are used to determine brood collection and natural spawning objectives, and may be reassessed with PIT tag data during the run. The basic collection guidelines are as follows:
- a. Fish are handled using electro narcosis, with MS-222 as a backup. If fish are exposed to MS-222, a 21-day period is required before they are used for consumption, are released as carcass outplants, or are released alive. The MS-222 label must be adhered to. Fish held for distribution are sampled according to Appendix A.
  - b. Release all natural origin jacks upstream. Hatchery jacks should represent a maximum of ten percent (10%) of the total male hatchery fish released upstream. Natural- and hatchery-origin fish may be radio tagged and released at the weir.
    - i. Radio-tagged fish are included in estimates of the hatchery/natural composition of fish passed above the weir.
    - ii. Inject broodstock with oxytetracycline and transport to circular tanks in endemic building (*see vii. Adult holding Ponds*).
  - c. Hatchery-origin fish may be transferred to Wallowa Hatchery for distribution or recycled for harvest.
  - d. Surplus hatchery may be outplanted according to these guidelines:
    - i. Hatchery fish can be outplanted to the Wallowa River and tributaries (Bear Creek, Hurricane Creek, and Prairie Creek).

- ii. If outplants occur in new reaches, subsequent spawning success will be monitored in target reaches by NPT and/or ODFW staff.
- e. Marking - Fish captured at the Lostine weir, and not retained for broodstock, will receive an opercle punch that indicates their intended disposition:
  - i. Pass above weir = 1 LOP
  - ii. Trapped and hauled to acclimation = 2 LOP
  - iii. Recycled to fishery = 1 ROP
  - iv. Outplanted for natural spawning = 2 ROP
- 9. Imnaha River -In-season run projections and the sliding scale (SOP Table 5 AOP Table 4) are used to determine brood collection and natural spawning objectives, and may be reassessed with PIT tag data during the run.
  - a. See Appendix E. Standard Operating Procedures for the Imnaha River weir for handling guidelines.
  - b. ODFW staff will determine fish disposition on-site. Priorities for hatchery fish will be: 1) broodstock, 2) natural spawning above the weir, 3) recycle to fishery, 4) transport to Wallowa Hatchery and Lookingglass Hatcheries for CWT recovery, subsistence and/or foodbank use, and 5) outplants. Priorities for natural fish will be: 1) broodstock, and 2) natural spawning above the weir.
  - c. *Chinook salmon* - Inject broodstock females with erythromycin (or tulathromycin) and oxytetracycline. Males are injected with oxytetracycline and erythromycin (or tulathromycin). All fish that are not injected with erythromycin, such as those that are received after August 1<sup>st</sup>, will be given 3 caudal fin punches. Transport to the adult holding ponds (*see* vii. Adult holding Ponds). After broodstock and upstream passage, surplus hatchery jacks and adults can be utilized as described above.
    - i. If fish are exposed to MS-222, a 21-day period is required before they are used for consumption or are released as carcass outplants, or are released alive. The MS-222 label must be adhered to.
    - ii. A total of 300 live hatchery adults can be outplanted to Big Sheep Cr. and Lick Cr. If more adult out plants are anticipated, notification should be provided to NOAA Fisheries and ODFW Wallowa District.
    - iii. Surplus carcass jacks can be disposed of in Big Sheep Cr. and Lick Cr. Surplus live hatchery jacks can be released in Big Sheep Cr. And Lick Creek after the last redd count survey.
    - iv. Collect scales on ~50% of natural origin adults passed above the weir.
  - d. *Steelhead* – See Imnaha weir SOP (Appendix E).
  - e. *Bull Trout* – See Imnaha weir SOP (Appendix E).
  - f. Marking - Fish captured at the Imnaha weir, and not retained for broodstock, will receive an opercle punch that indicates their intended disposition:
    - i. Pass above weir = 1 ROP

- ii. Outplant = 1 LOP
- iii. Recycle = 2 LOP
- iv. Double recycle = 3 LOP (previously recycled fish will have another LOP applied).
- g. Mortalities and carcasses
  - i. Before redd surveys: Imnaha staff collects the first 10 weir mortalities, seals them intact in a labelled plastic bag, freezes them, and notifies Fish Health that they are ready for pathology sampling. Note: avoid collecting fallbacks for Fish Health samples. Dead fish found on the weir that are not collected for fish health should be sampled for bio data that includes: fork length, sex, pre-spawn status, record fin clip, the opercle punch (for recapture status), PIT tag, and collect snouts on carcasses with a CWT.
  - ii. After redd surveys: Fish research will sample mortalities as described above, then carcasses should be clearly identified as sampled (tails removed) and returned to the river below the weir.
  - iii. Biological data will be sent to ODFW Fish Research (Dittmer).

#### **F. Broodstock transportation**

1. CTUIR and NPT will transport fish from Upper Grande Ronde, Catherine Creek, and Lostine River sites. ODFW or NPT will provide transportation of fish from the Imnaha weir to Lookingglass Hatchery.
2. Broodstock should be hauled daily, and not held more than 72 hours. Imnaha broodstock will be hauled on Monday and Thursday, or as needed to make broodstock.
3. Drivers should complete a transfer data sheet, provide to Lookingglass Hatchery staff for data entry in the HMIS system.
4. Minimize shock by hauling during morning hours to take advantage of cooler stream temperatures. Temperature differences between transport container and facility water should not exceed 10° F (5.6° C). Temper if needed.
5. Handling - Fish will be netted from the transport tank and placed in holding tanks at Lookingglass Hatchery. Lookingglass Hatchery personnel will record all observations on data sheets and report to Fish Health at the end of the season.

#### **G. Adult holding at Lookingglass Hatchery**

1. Upper Grande Ronde – One (1) adult holding pond
2. Catherine Creek – One (1) adult holding pond
3. Lookingglass Creek – One (1) adult holding pond, food bank fish held in circular tank 19
4. Lostine River – Endemic building, circular tank 20 and 21. Food fish held at Wallowa Hatchery.
5. Imnaha River – One (1) adult holding pond, food fish held at Wallowa Hatchery.

### **IV. Spawning**

- A. Key Contacts – Crump, Greiner, McLean, Zollman, Brigante, Deal.

- B. Schedule** - A pre-sort to determine sex ratio and allow adjustments to brood stock collections, will occur mid to late July. The first sort will occur in early to mid-August, with an expected first spawn during mid-August. These days can be adjusted by workload and water temperatures. Fish are spawned weekly as described below:
1. Tuesday – Imnaha and Lostine
  2. Wednesday – Upper Grande Ronde, Catherine Creek, and Lookingglass Creek.
  3. Thursday – a catch up day if spawning was not completed Tuesday and/or Wednesday
- C. Anesthetic** – Electro-anesthesia used for all stocks.
- D. Spawning** – Broodstock are spawned at Lookingglass Hatchery. Sorting and spawning for each stock takes place on the same day.
1. Hatchery and co-manager staffs will determine fertilization matrices.
- E. Pairing**
2. Fertilize a maximum of 10% of the eggs fish with fish <601mm (CC, UGR, LGC) and <631mm (IM, LR)
  3. Maximize the amount of the eggs fertilized with five year old males, when available (UGR, LGC, and CC males >800mm, and LR and IM males >850mm).
  4. Large males may be spawned up to 3 times.
  5. Conduct jack spawning with 1 female to 1 jack matrix. Most adult spawning matrices will be 2 females x 2 males, but matrices of 1 x 1, 1 x 2, 2 x 1, or 3 x 2 can be used if necessary.
  6. Incubate fertilized eggs at Lookingglass hatchery.
  7. Determine fecundity at eye-up.
- F. General fertilization procedures**
1. Sort and euthanize ripe females.
  2. Collect eggs preventing addition of outside containments (other body parts).
  3. Store eggs separately for each individual female.
  4. Drain ovarian fluid from eggs.
  5. Sort males, spawn in dry cup.
  6. Mix sperm with eggs, activate with pathogen free water (~100 ml).
  7. Wait 60 seconds, rinse eggs.
  8. Treat fertilized and rinsed eggs in 100 ppm Iodophor solution for minimum of 45 minutes.
  9. Tray eggs, 1 tray per individual female.
- G. Surplus** - Fish that are surplus to broodstock needs may be returned to stream (send data to ODFW Fish Research). Fish injected with antibiotics will not be used for human consumption or released where legal harvest is possible. Fish



exposed to MS-222 will not be released or used for human consumption if the withdrawal period has not been met. The MS-222 label must be adhered to.

- H. Staffing support** - CTUIR will provide fish culture support for Upper Grande Ronde, Catherine, and Lookingglass stocks. Nez Perce Tribe will provide fish culture support for spawning of the Imnaha and Lostine River stocks.

## **V. Incubation**

- I.** All eggs will be incubated to eyed stage at Lookingglass Hatchery. Until eye-up, segregate individual females. After eye-up, eggs will be enumerated and segregated by disease profile. No eggs will be culled after egg enumeration, unless segregated by individual females after enumeration. If possible, only lower risk BKD eggs will be reared (<0.200 OD units). Eggs will be trayed down at 5,000-5,500 eggs/tray.
- A.** Eggs are incubated using chilled and un-chilled well water and UV treated Lookingglass Creek water incubators. Untreated creek water may have to be used in the event of a power outage or other emergency.
- B.** If excess production is expected (i.e., greater than 110% of smolt release levels outlined in Table 1 of the Annual Operating Plan), co-managers have the following dispositions available for each program:
  1. Upper Grande Ronde: up to 35,000 eggs or fry can be released into Meadow and Sheep Creeks.
  2. Catherine Creek: up to 21,000 eggs or fry can be released into Indian Creek
  3. Lookingglass Creek: up to 35,000 eggs or fry can be released into Lookingglass Creek
  4. Lostine River: up to 48,016 eyed eggs, fry, or parr can be released into Bear Creek, Hays Fork Prairie Creek, the Wallowa River, and Lostine Rivers.
  5. Imnaha River: up to 95,000 eggs or fry can be released into Lick Creek
  6. Specific locations, timing, and marking plans for excess will be coordinated on a case-by-case basis amongst co-managers.

## **VI. Smolt Production**

- A.** Size at release - Smolts are produced at a target size of 20-25 fpp at release.
- B.** Early Rearing – Fry are reared in double deep troughs at Lookingglass Hatchery on UV-treated Lookingglass Creek water. Fish are transferred outside to Lookingglass Creek water in April or May. Trough loading will be range from 30,000-50,000 per trough.
- C.** Final Rearing – In May, fish are moved to outdoor raceways. After marking and spawning, fish will be placed in final rearing containers. Refer to AOP for final ponding plan.

**D. Lookingglass Hatchery raceway allocation** - Eleven raceways are allocated for Grande Ronde tributary production, seven raceways for Imnaha production. Lookingglass Creek smolts are reared in the adult holding ponds (AHP). Current allocation by stock is:

1. Upper Grande Ronde River – 4 raceways.
2. Catherine Creek – 2 raceways.
3. Lookingglass Creek – 1 raceway and 4 adult holding ponds.
4. Lostine River – 4 raceways.
5. Imnaha River – 7 raceways.

## **VII. Marking**

J. Key Contacts- ODFW (Kevin Kennedy- Clackamas, Feldhaus, LaPoint, and Keniry), NPT (Vogel, Simmons)

K. Mark Type

1. Adipose clipping –occurs during the last two weeks of August.
2. Coded wire tagging (CWT) - occurs during the last two weeks of August.
3. PIT tagging –occurs during late-October.
4. See AOP for marking, CWT, and PIT tag number information.

## **VIII. Monitoring and Evaluation**

L. Key Contacts: ODFW (Keniry, Feldhaus), CTUIR (Naylor, Crump), NPT (Vogel, Simmons)

M. Data is collected on behalf of several ongoing monitoring efforts, including: program evaluations, monitoring differences between natural and hatchery production, growth and survival of natural-origin fish (in collaboration with ODFW Early Life History Project).

N. PIT tags are used to detect fish at main stem dams, in-stream arrays, and weirs to assess downstream and upstream survival and timing.

5. Catherine Creek and Imnaha stocks are part of the Comparative Survival Study (CSS), and therefore receives PIT tags from the Fish Passage Center.

O. Data collection efforts

1. During pre-liberation sampling the following data is collected per raceway:
  - a. Collect 100 weights
  - b. Collect 100 lengths
  - c. Collect 50 genetic samples for Genetic Stock Index (for each stock, not raceway)
  - d. Check 550 fish per tag code for tag retention
  - e. Check 350 fish per raceway for fin clip quality
2. While processing adult fish (live or mortalities) at trapping facilities, operators should collect the following data:
  - f. Data collection – count, length, marks/tags, sex, percent spawned, scan for and record PIT tag codes.

- g. Tissue collection – snout/scales, genetic samples.
- 3. During broodstock spawning and early incubation, collect the following:
  - a. Length (all fish), sex, weight (females), marks/tags, PIT tags, eyed egg weights, individual fecundity
  - b. Tissue collection – snout/scales/fin rays, kidney sample, genetic sample
- 4. During spawning ground surveys, the following data is collected:
  - a. Redds (count, GPS)
  - b. Live fish (count, age class [jack or adult])
  - c. Carcasses (count, GPS, length, marks/tags, snout/scales/fin rays, kidney sample, genetic sample)
- 5. Genetic tissue collection for monitoring and pedigree analysis
  - a. Un-punched Chinook carcasses recovered above weirs on Catherine Creek, Lookingglass Creek, and the Lostine River.

## **Grande Ronde and Snake River Fall Chinook Program**

### **IX. Smolt Release**

- A.** Grande Ronde River - Priority 10 in the 2018-2027 U.S. vs. Oregon production table B4 targets a total production of 500,000 sub-yearlings scheduled for release in the Wallowa River around mid-May at 50 fpp. Marks for this release are as follows:
  - 1. 100,000 AdCWT and 400,000 Ad Only
- B.** Couse Creek- Priority 12 in the 2018-2027 U.S. vs. Oregon production table B4 targets a total production of 700,000 scheduled for release into Couse Creek in mid-May at 50 fpp. Marks for this release are as follows:
  - 1. 100,000 AdCWT and 500,000 unmarked

### **X. Adult collections and spawning – refer to Lyons Ferry AOP.**

- A.** Incubation and Rearing
  - 1. Incubation at Lyons Ferry - After eye-up, inventory, and disease profiles, Lyons Ferry staff will combine eggs and ship to Irrigon Hatchery in December. Only eggs from females with ELISA OD values below 0.200 OD units are transferred. Fish are reared and tagged at Irrigon Hatchery prior to release.
  - 2. Couse Creek eggs will be shipped as green eggs.
- B.** Key Contacts
  - 1. Lyons Ferry Hatchery (A. Trump, Herr), ODFW (Schmidt, Keniry, Bratcher), CTUIR (McLean, Krajcik), NPT (Johnson, Simmons, Young), IDFG (Putnam), IPC (Rosenberger)

## **Grande Ronde Coho Salmon Program**

### **XI. Overview**

- B. Consistent with production tables specified in the U.S. vs. Oregon fish management agreement, the production objective for this program is 500,000 smolts released in the Wallowa River Basin (Eggleston Corner Rd, 100k into Hurricane Creek and 400k at Fish Hatchery Lane bridge).
- C. Consistent with production tables specified in the U.S. vs. Oregon fish management agreement, the production objective for this program is 500,000.
- D. Management objective – To reintroduce Coho salmon to Northeast Oregon, continue to provide measurable harvest benefits that these fish provide for treaty and non-treaty fisheries in the mainstem Columbia, and develop fisheries in the Snake and Grande Ronde rivers. Concurrently this program could reestablish natural production of Coho salmon in the Wallowa/Lostine River and provide the opportunity for natural recolonization of Coho in Grande Ronde tributaries (e.g., Minam and Wenaha rivers) that historically produced Coho salmon.
- E. The coho program described in this document is the first step of a phased approach to reintroduce coho salmon to the Grande Ronde basin. This first ‘pilot’ phase will direct stream release smolts during 2017-2024 and evaluate subsequent adult returns to determine if this Grande Ronde program can be self-sustaining.

### **XII. Smolt release**

- A. Final rearing - Fish are reared to smolt at Cascade Hatchery.
- B. Transportation – Coho smolts will be hauled from Cascade Hatchery to the Wallowa River Basin for acclimation (late-March) and release in April. Transportation is coordinated between NPT and ODFW.
- C. Release
- D. Wallowa River & tributaries – Fish will be released in the Wallowa River and tributaries outside of the Lostine River and will be coordinated between NPT and ODFW.
- E. Rotary screw trap operators in the Lostine River should be notified prior to release. Key contacts are Gibson and Keniry (ODFW), and Simmons and Watry (NPT). After that, NPT will provide support, if needed, to the Lostine River trap crew until the trap resumes normal operations.

### **XIII. Adult collections and spawning**

- A. Trap installation and scheduling

1. The first returns of adult coho salmon began in 2017.
2. The Lostine River weir will remain in operation following trapping for Chinook salmon, and all applicable guidelines described above for spring Chinook salmon will be followed (see Grande Ronde and Imnaha Spring Chinook Salmon, Adult Trapping).
3. Broodstock is not collected during this pilot phase of the program. Coho salmon are enumerated and sampled for marks (see monitoring and evaluation below).
4. During the pilot phase, disposition of coho will be coordinated annually with NPT and ODFW.

**B. Spawning**

5. Tanner Creek stock from the Bonneville Complex or other Tanner Creek sources will be used for the program until a local broodstock can be developed. Starting in 2024, broodstock from the Clearwater River may be used. However, collection of Tanner Creek broodstock may be necessary if returns to the Clearwater River are inadequate to supply eggs to the Grande Ronde Program.

**C. Incubation, and rearing**

6. Incubation and rearing –Rearing to the eyed egg stage may occur at Cascade Hatchery or at hatcheries in the Clearwater Basin. Eggs that are eyed within the Clearwater Basin will be transferred to Cascade Hatchery where they will be reared from eyed egg to smolt.

**D. Marking**

1. Smolts are marked consistent with the following guidelines, and revisions to marked schemes may be coordinated in the U.S. vs. Oregon forum:
  - a. Adipose clipping – 100%
  - b. Coded Wire Tagging (CWT) - Appropriate level of CWT to evaluate harvest contribution (~90,000 AdCWT).
  - c. PIT Tags - A portion of juveniles will be implanted with PIT tags to assess downstream survival (initial goal of 5,000 AdPIT, depending on funding).

**E. Monitoring and evaluation**

1. Objectives - During the pilot phase of this program, monitoring objectives will be limited to the following:
  - a. Enumerate adult returns to weir.
  - b. Estimate smolt-to-adult survival of release, contribution to fisheries, and straying using Coded Wire Tags (CWT).
  - c. Determine if natural spawning is occurring in the Lostine River.
    - iv. Other Grande Ronde tributaries may be surveyed (Minam, Wenaha, Wallowa Rivers).
2. Data Collection – Data will be limited to adult returns at the Lostine River weir, but may be expanded if resources allow. At the Lostine weir, operators collect the following data:
  - a. Enumerate coho catch.

- b. Disposition, as decided by co-managers annually (will be marked and released upstream).
- c. Examine for external marks.
- d. Estimate sex and measure length of individual fish.
- e. Scan for CWT and PIT Tags, collect genetic sample (opercle punch).

**F. Key Contacts**

- 3. Key Contacts – B. Johnson, Zollman, Vogel, Watry, Simmons (NPT); Gibbs, Harrod, Bratcher, (ODFW); McLean, Krajcik (CTUIR); Alex Clark (ODFW Cascade Hatchery).

## Grande Ronde Lamprey / Freshwater Mussels

### XIV. Lamprey Overview

- A. Objectives - The purpose of this stop gap effort is to avoid local extirpation in the Snake River Basin and maintain a population of ammocoetes that serve as a source of pheromone attractants drawing adults upstream to spawn in the abundant habitat in this region, thereby continuing a presence in the Snake River Basin until upstream adult and downstream juvenile passage problems are identified and corrected, and healthy, harvestable populations are restored.
- B. CTUIR program
  - 1. Adult collection – Adult lamprey are trapped and collected by CTUIR at main stem dams on the Columbia River.
  - 2. Adult holding – Once trapped, lamprey are held at Minthorn Springs facility in the Umatilla basin.
  - 3. Adult releases – Release locations and numbers will be determined on an annual basis. See Table 14 in AOP section.
  - 4. Key contacts – Aaron Jackson (CTUIR – aaronjackson@ctuir.org), Brandt (ODFW)
- C. NPT Program
  - 1. Adult collection – Adult lamprey are trapped and collected by NPT Fisheries staff at Bonneville, The Dalles, and John Day dams.
  - 2. Adult Holding – Once trapped, lamprey are held at Nez Perce Tribal hatchery on the Clearwater River through winter months.
  - 3. Adult releases – Adult Lamprey are released to spawn naturally in tributaries of the Clearwater and Grande Ronde Rivers, including Asotin Creek in Washington. In Oregon, lamprey have been released in the Wallowa and Minam Rivers, and Joseph Creek. Release locations will be determined on an annual basis, and as availability of lamprey allows.
  - 4. Key contacts – Tod Sween (NPT, tods@nezperce.org); Bratcher, Brandt (ODFW)
- D. Fish Health
  - 1. Fish Health recommends an examination (up to 5 grab-sampled) be conducted prior to lamprey being transferred to Oregon waters. At a minimum, all moribund and dead lamprey should be examined during rearing in Oregon and Idaho to develop a pathogen history. If unable to lethally sample due to tribal policy, then develop a pathogen history as best as possible with moribund and dead lamprey. For lamprey releases in Oregon, Fish Health recommends source lamprey for holding in Idaho come from direct transfer from the dam collection site or the CTUIR holding site near Pendleton, OR rather than Yakima Indian Nation site near Prosser.



## Freshwater Mussel Project Overview

### XV. Mussel Overview

- E. Objectives - The purpose is to evaluate the status and distribution of freshwater mussel populations in the Grande Ronde sub-basin. This information will be used to inform conservation efforts of remaining freshwater mussel populations and to identify and characterize broodstock populations for future restoration efforts.
- F. Planned activities - The CTUIR Freshwater Mussel Project will:
  - 1. Conduct surveys to identify and monitor freshwater mussel populations in the Grande Ronde and tributaries. Freshwater mussel surveys are visual, non-destructive surveys performed by snorkeling or wading in the channel during low water time periods.
  - 2. Collect mussel tissue samples for species identification or for population genetic characterization. Mussels are collected by hand, with minimal disturbance to substrates.
  - 3. Collect gravid adult females for broodstock. Individuals used for broodstock are collected by hand with minimal disturbance to substrates, non-brooding animals are returned to sediments in the same location.
  - 4. Conduct mussel surveys, salvages, translocations, and follow-up monitoring at restoration project sites with in-stream activities that could negatively impact freshwater mussels.
  - 5. The Xerces Society and the CTUIR Freshwater Mussel Project collaboratively maintain a database of western freshwater mussel records. Please help maintain this critical conservation tool by reporting mussel observations to the CTUIR Mussel project or the Xerces Society ([mussels@xerces.org](mailto:mussels@xerces.org)).
- G. Key Contacts – Alexa Maine (CTUIR, [AlexaMaine@ctuir.org](mailto:AlexaMaine@ctuir.org)).

Table 1. Upper Grande Ronde River weir management guidelines for broodstock collection and passage.

<b>Total adult escapement to the mouth (HOR+NOR)<sup>a</sup></b>	<b>HOR at mouth (%)</b>	<b>Maximum NOR in Broodstock (%)</b>	<b>HOR to retain for Broodstock (%)<sup>b</sup></b>	<b>HOR adults released above weir (%)</b>	<b>Minimum NOR in broodstock (%)</b>	<b>Strays allowed above weir (%)<sup>c</sup></b>
Any	Any	Up to 50	Up to 100	Up to 100	— <sup>d</sup>	≤5
<sup>a</sup> Pre-season estimate of total escapement <sup>b</sup> Conventional hatchery adults only, all captive brood adults released to spawn naturally or outplanted <sup>c</sup> For hatchery adults originating from different gene conservation groups (Rapid River stock or strays from outside the Grande Ronde basin) <sup>d</sup> Not decision factor at this level of escapement, percentage determined by other criteria						

Table 2. Catherine Creek Spring Chinook broodstock/upstream passage management guidelines.

<b>Total adult escapement to the mouth (HOR+NOR)<sup>a</sup></b>	<b>HOR at mouth (%)</b>	<b>Max of NOR Run in Broodstock (%)</b>	<b>HOR to retain for Broodstock (%)<sup>b</sup></b>	<b>HOR adults released above weir (%)</b>	<b>Minimum NOR in broodstock (%)</b>	<b>Strays allowed above weir (%)<sup>c</sup></b>
< 250	Any	40	40	— <sup>d</sup>	— <sup>d</sup>	≤5
251-500	Any	20 <sup>d</sup>	20	≤70	≥20	≤5
>500	Any	≤20	— <sup>e</sup>	≤50	≥30	≤5
<sup>a</sup> Pre-season estimate of total escapement <sup>b</sup> Conventional hatchery adults only, all captive brood adults released to spawn naturally or outplanted <sup>c</sup> For hatchery adults originating from different gene conservation groups (Rapid River stock or strays from outside the Grande Ronde basin) <sup>d</sup> Not to exceed 150,000 smolt production <sup>e</sup> Not decision factor at this level of escapement, percentage determined by other criteria						

Table 3. Lookingglass Creek weir management guidelines for broodstock collection and passage. Current weir management strategy is consistent with Lookingglass Management Plan. Table is a placeholder until permanent agreement is in place.

<b>Escapement Level</b>	<b>Passed Above (%)</b>	<b>Retain for Broodstock (%)</b>
150	67	33
200	60	40
250	55	45
300	50	50
>300 – adjustments will be made based on brood needs. If brood need has been met remainder to be released upstream		

Table 4. Lostine River weir management guidelines for broodstock collection and passage. Assumes program goal of 250,000 smolts (166 adults for broodstock).

<b>Estimated Natural Run of ADULTS to River Mouth (proportion of minimum abundance threshold recommended by ICTRT<sup>1</sup>)</b>	<b>Number of ADULT Natural Fish to River Mouth</b>	<b>Max % Natural ADULTS for broodstock</b>	<b>Number of ADULT Natural Fish Retained for Broodstock (Proportion of Natural Brood)</b>	<b>Max Proportion of ADULT hatchery fish released above weir</b>	<b>Minimum % natural ADULTS in Broodstock</b>
<b>&lt; 0.05 of Critical</b>	< 8	0	0	NA	NA
<b>0.05 – 0.5 of Critical</b>	8 – 74	50%	4 – 37	NA	NA
<b>0.5 of Critical - Critical</b>	75 – 149	40%	30 – 60	70%	20%
<b>Critical – 0.5 of Viable</b>	150 – 249	40%	60 – 100	60%	25%
<b>0.5 Viable – Viable</b>	250 – 499	30%	75 – 150	50%	30%
<b>Viable – 1.5 Viable</b>	500 – 749	30%	150 – 225	40%	40%
<b>1.5 – 2.0 Viable</b>	750 – 999	25%	188 – 250	25%	50%
<b>&gt;2.0 Times Viable</b>	> 1,000	25%	> 250	<10%	100%
<sup>1</sup> Lostine River contributes about 50% of production for Wallowa/Lostine Population - Viable level is 50% of TRT recommended minimum abundance threshold for Wallowa/Lostine population (1,000) after broodstock collection and fishery.					

Table 5. Sliding scale management tool for Imnaha Chinook hatchery program utilized for managing disposition of Chinook salmon adults for broodstock and escapement to natural spawning areas (Table 1 excerpted from Imnaha spring Chinook HGMP May 2011).

<b>Estimated natural run of ADULTS to river mouth as a proportion of minimum interior TRT minimum abundance threshold (MAT)</b>	<b>Number of ADULT natural fish to river mouth</b>	<b>Expected handle rate at weir of ADULT natural fish (50%)</b>	<b>Max % natural ADULTS for broodstock<sup>2</sup></b>	<b>Number of ADULT natural fish retained for broodstock</b>	<b>Max proportion of ADULT hatchery fish released above weir</b>	<b>% Natural ADULTS in Broodstock<sup>3</sup></b>
< .05 of Critical <sup>4</sup>	< 15 <sup>4</sup>	< 8 <sup>4</sup>	0	0	NA	NA
.05 - .5 of Critical	15 – 149	8 - 74	50%	04 - 37	NA	NA
.5 – Critical	150 -299	75 -149	40%	30 - 60	70%	20%
Critical - .5 of MAT	300 – 499	150 -249	40%	60 - 100	60%	25%
.5 Viable – MAT	500 – 999	250 - 499	30%	75 - 150	50%	30%
			<b>40%<sup>1</sup></b>			
Viable - 1.5 MAT	1000 – 1499	500 - 749	30%	150 – 225	40%	40%
			<b>40%<sup>1</sup></b>		30%	
1.5 - 2 x MAT	1500 – 1999	750 - 999	25%	188 – 250	25%	50%
> 2 x MAT	> 2000	> 1000	25%	> 250	<10%	100%
<sup>1</sup> Percentage highlighted will be implemented after three consecutive years (or 3 year mean) at viable or greater <sup>2</sup> Maximum percent of wild adults handled at weir collected for Broodstock <sup>3</sup> Target percent natural adults in Broodstock <sup>4</sup> In the submitted HGMP, the less-than (<) symbols shown above are displayed in error as greater-than (>) symbol						

## Appendix A. Disinfections and Sanitation Guidelines for all LSRCP Hatcheries.

### Specific Operational Recommendations

For background on the importance of these recommendations see page 1 of Appendix C (2013 AOP)

Applies to Who?	Prevention Control Measure or Sanitary Practice	Guideline Comment
All	Disinfect all gear/equipment prior to entering or leaving hatchery grounds	-As per attached iodophor protocol -Hatchery crew responsible for providing tub of 100 ppm iodophor
Hatchery Crew	Do not go from adult handling operations to juvenile operations activities unless all bib gear is thoroughly disinfected.	-As per attached iodophor protocol -it would be preferable to have bib gear designated for either adult or juvenile use.
Hatchery Crew	Pick mortality on a daily basis	-This is consistent with ODFW's statewide Fish Hatchery and Fish Health Management Policy.
All	Disinfect equipment when moving from raceway to raceway or tank to tank for <u>any</u> fish handling or pond cleaning activities	-As per attached iodophor protocol -Includes CWT, fin clipping and PIT tag operations. See footnote for marking.
All	Use footbaths upon entering or leaving the work area for a given program	-Use larger tub of disinfectant if involved in a spawning
All	Use a new disposable apron or disinfected personal rain gear while working with fish	
CTUIR Personnel operations at Lookingglass Hatchery	Disinfect all gear/equipment prior to entering or leaving hatchery grounds, Lookingglass Creek, or the intake building and when done with operations at intake	-CTUIR personnel responsible to maintain and use a tub of 100 ppm iodophor at intake building workstation
Hatchery Crew	Assure that individual raceway and tank mortality "picker equipment" is in place at each raceway and tank	-All use these for the specifically designated raceway
Hatchery Crew	Sanitize each raceway prior to use for the next brood year. (see below for recommendation)	-dry for a minimum of three days
Hatchery Crew	Keep footbaths located at strategic locations refreshed with disinfectant	-As per iodophor label, refreshed as needed
People at Spawns	Disinfect the spawning table and spawning work area between stocks and at the end of the day	-As per attached iodophor protocol
Research, Hatchery Crew & Fish Health Personnel	Handle and necropsy dead fish only in designated areas	-Adult morts: use concrete pad outside spawn area or concrete pad in endemic building at LGH -store snouts only in adult mortality freezer -Juvenile morts: store in freezer in designated area for this purpose.
PIT taggers	-PIT tagging supervisors maintain and keep footbaths by each door of PIT tagging trailer for use during operations -Assure that PIT tagging needles are new or clean and sharp -Disinfect in 70% Isopropyl alcohol -No re-use of PIT tag needles until air dried	-if PIT tag needles are re-used disinfect as per isopropyl protocol attached
Lib Truck Operators	Assure proper disinfection of tank and equipment prior to collection or transfer	-As per attached disinfectant application Summary

Appendix A (cont.): Disinfection and Sanitation Guidelines for all LSRCP hatcheries  
Summary of Recommended Disinfectants (Concentration and time) and for what Application.

Disinfectant*	Application	Concentration	Time	Comment
<b>Iodophor</b>	Nets, gear and equipment, clipping & tagging van, PIT tag stations, large tub disinfectant containers, spawning colanders and buckets, lib truck, footbaths, floors  Note: For raceway sanitization** – thoroughly clean the unit to remove dirt, spray or brush on 75-100 ppm iodophor and let this remain for a minimum of 10 minutes. Leave it to dry for a minimum of 3 days. Allow iodophor to dry and break down with exposure to light.  **If the above recommendation cannot be done then sanitize raceways by thoroughly cleaning them and leaving to dry for a minimum of 3 days.	100 ppm  Note: to make 100 ppm solution mix 6.7 oz of jug strength iodophor to 5 gallons H <sub>2</sub> O or 6.7 oz.=189ml	10 min.	-Equipment should be pre-rinsed to remove dirt, mucus or other organic material which reduces the efficacy of disinfection and sanitization  -Rinse equipment to remove harmful residue if equipment is going into standing water containing fish or fish are being placed into the equipment (tank or bucket). Remember that iodine at 1:20,000 is harmful to fish.  -Argentyne or other buffered iodophors such as Western Chemicals “PVP iodine” would be acceptable. <u>Note:</u> if DRAW 476 is used remember this product is 1.75% active iodine and unbuffered so should not be used for water- hardening eggs
	Water hardening eggs	100 ppm	Minimum 15 minutes	This is the statewide general practice
	Egg transfers - disinfection at receiving station	100 ppm	10 minutes	
<b>Virkon Aquatic</b>	Footbaths, nets, boots & gear			As per label
<b>Chlorine or Aqueous solution as sodium hypochlorite (Household Bleach)</b>	Lib truck tanks	10 ppm	10 min.	Organic matter binds and neutralizes
	Raceway disinfection	100 ppm		Left to dry and breakdown in sun. Need to assure that no bleach goes to effluent.

<sup>a</sup>Within a stock, operations will start with groups determined to be of lowest disease risk proceeding to raceways of higher disease risk. The latest fish health information should be used to determine the least risky raceway sequence.

<sup>b</sup> All chemical use will be done in accordance with label use and reporting requirements. Disinfecting and disinfected water must be disposed of in an approved manner.

Appendix B. Juvenile Chinook fish health monitoring plan and disease treatments.

Location	Stock	Examination Category	Protocol	Comment/Disease Treatment
Lookingglass Hatchery	200 (LR) 201(CC) 80 (GR) 29(IM) 81(LG)	Routine Monthly	<p>-Gill &amp; skin wet mounts from a combination of moribund and healthy fish.</p> <p>-A subset of mort/moribund per stock, kidney smears on TYE-S agar. Gill culture if suspect gill disease. <i>R. salmoninarum</i> assays (D-FAT or ELISA) if bacterial kidney disease is suspect.</p> <p>-tissues (gill/ kidney/spleen) will be assayed for cultivable viruses from a sub-sample of fish as is consistent with clinical signs (priority in raceways with increased daily loss).</p> <p>-Grab sampled and moribund fish will be assayed for EIBS (blood smears and hematocrits) during exams as needed (i.e., persistent anemia).</p>	<p>A 21-day Aquamycin feed will be implemented for all raceways for each stock, except for all the Lostine River and Imnaha River raceways. Disease outbreaks - treated on a case-by-case basis. Therapies and remedial measures are based on conventional and available treatments, new information, and innovation and other treatments to be implemented if recommended by Fish Health Services in order to maintain health of fish. Formalin treatments may be implemented for Ichthyobodo (costia) infestations. Fungus - Formalin flushes (1 hour) are prescribed after fin clipping, PIT tagging, VIE tagging and coded wire tagging for two consecutive days. Formalin requires a veterinarian prescription when used off-label. Treat CWD with medicated feed if necessary, using a Veterinary Feed Directive (VFD) through Fish Health Services.</p>
Lookingglass Hatchery	200 (LR) 201(CC) 80(GR) 29(IM) 81(LG)	Pre-transfer & Annual Myxobolus cerebralis testing	<p>-n = 60 grab-sampled smolts per stock</p> <p>-<i>R. salmoninarum</i> by ELISA</p> <p>-tissues (gill/kidney/spleen) from 5 fish pools for culturable viruses</p> <p>-wet mounts of skin &amp; gill tissue from a minimum of 5 fish</p> <p>-<i>O.mykiss</i> on water supply for 6 months (60 fish) for <i>Myxobolus cerebralis</i></p>	<p>Pre-transfer grab-sample numbers may vary depending on disease history and number of fish for a given brood year.</p> <p>A small group of <i>O. mykiss</i> should be reared at Lookingglass Hatchery for annual <i>M. cerebralis</i> testing 181 days after ponding. The timing of the 181 days is at the hatchery's discretion and should be chosen to coordinate with hatchery operations.</p>
Chinook acclimation IM, LR, CC & UGR		Pre-liberation	<p>-Smolt groups held at acclimation sites longer than 3 weeks will be evaluated with a lesser number of "grab-sampled" fish as in pre-transfer protocol above.</p> <p>-Mortalities will be examined.</p> <p>-If available, some fresh, non-frozen moribunds and mortalities will be examined.</p>	<p>Pre-liberation grab-sample numbers at acclimation sites may vary depending on disease history and number of fish for a given brood year.</p>



Appendix C. Adult Chinook fish health monitoring plan and disease treatments at Lookingglass Hatchery.

Stock	Examination Category	Protocol	Comment
200 (LR) 201 (CC) 80 (GR) 29 (IM) 81 (LGC)	Adult Spawners (Broodstock)	-A minimum of 60 females will be sampled for virus from each stock (or all females if <60). Samples will be individual or pooled (3 per pool) ovarian fluid and caeca/kidney/spleen sample pools not to exceed 5 fish. - All females for BKD by ELISA - If available, 10g egg samples from thirty females from each stock will be collected and flash-frozen at time of spawn for thiamine analysis.	ELISA results will be used to implement BKD prevention control through culling of eggs known to be of higher risk.
200 (LR) 201(CC) 80(GR) 29(IM) 81(LG)	Prespawning Mortality	- Up to 10/stock (before Sep 1 <sup>st</sup> ): -Kidney sampled for BKD by ELISA -Examined and worked up as needed per Fish Pathologist	<ul style="list-style-type: none"> <li>Note: additional mortality may be sampled</li> <li>Lookingglass Creek mortalities will be worked up with CTUIR staff to assure data collection covers all the needed information</li> </ul>
81 or 201 LG-CK	Spawning Ground Survey	-Collect a sub-sample of 30 kidney samples from adult Chinook above the weir (hatchery intake)	<ul style="list-style-type: none"> <li>Fish Health Request</li> </ul>

Appendix D. Disease treatments and other drugs for adult Chinook Broodstock.

Location	Stock	Treatment for	Chemical/ Drug	Protocol	Comment
Lookingglass	200 (LR) 201(CC) 80(GR)  29(IM) 81(LG)	Fungus Control	Formalin	Formalin administered 3 days per week at 167 ppm for 1 hr. If there is an indication, such as heavy headburn or saprolegnia, the hatchery will work with Fish Health to increase administration up to 7 days per week.  Hydrogen peroxide 3 days per week at 100 ppm	If formalin cannot be used then use hydrogen peroxide (second choice) at the recommendation of fish health services  Continue treatments throughout the entire spawning season.
Lostine, Catherine Creek, Upper Grande Ronde, Imnaha and Lookingglass	200 (LR) 201(CC) 80(GR) 29(IM) 81(LG)	BKD       Furunculosis-Enteric Redmouth & other gram negative bacterial infections	Erythromycin or tulathromycin (all stock except 200 LR)   Oxytetracycline (all stocks)	Injection 20 mg/Kg (erythromycin INAD) or 2.5 mg/kg (tulathromycin veterinary prescription)      Injection 10 mg/kg (Veterinary Prescription)	Inject fish kept for broodstock. Tulathromycin will be used instead of erythromycin when erythromycin is unavailable. Note that fish that are injected with tulathromycin cannot enter the river either as live fish or carcasses as tulathromycin has an infinite withdrawal period.

## **Carcass Stream Nutrient Enrichment Health Guidelines**

### **A. DEQ Fish Carcass Placement Guidelines 12/16/2022**

1. Salmon and steelhead carcass placements are not to exceed 2,500 lbs./mile/year. Distributions should occur August through May, and they should not occur during times of extreme water flow or temperature events. Carcasses are placed in streams only when or where they will not adversely impact water quality. Streams should be flowing when carcasses are distributed. Carcasses should be placed only within the wetted portion of the channel and in streams historically used by anadromous salmonids for spawning. Disease Control: Adhere to all applicable and current ODFW guidelines (see below).

### **A. ODFW Guidelines**

1. For carcass stream enrichment programs, medication exposure and disease will be considered to avoid amplifying pathogens in the environment. The withdrawal period of medications must be met prior to carcass distribution. For a withdrawal period to be met, the fish must be kept alive for the entire duration of withdrawal days post exposure.
  - a. Withdrawal period after MS-222 use: 21 days
  - b. Withdrawal period after erythromycin injection: 60 days
  - c. Withdrawal period after tulathromycin (Draxxin) injection: infinite (i.e. neither live fish nor carcasses can be placed in rivers)
  - d. Withdrawal period after oxytetracycline injection: 21 days
2. Only fish that survive to spawning will be used, not mortalities.
3. Because *Myxobolus cerebralis* is endemic in wild fish and returning hatchery adults in the Imnaha and Grande Ronde watersheds, fish carcasses from these watersheds will not be transferred to other watersheds.
4. Before using carcasses for stream enrichment, freeze carcasses for 7 days at -20°C. If possible, carcasses will be frozen and saved until test results are obtained for virus (and *M. cerebralis*, if being tested). Freezing carcasses is beneficial as it reduces parasitic, viral, and bacterial loads overall, thus minimizing the risk of transmission to the native populations. If freezing is not possible, see guidelines below:
  - a. If carcasses are to be distributed in the same stream and live adults of the same stock have already been distributed in that stream or exist there naturally, carcasses may be distributed as soon as they are available.
  - b. If carcasses are to be moved to reaches of the originating watershed where adults are not normally present or to a different watershed:
    1. If there is no previous history of virus from the last 5 years, carcasses can be used unless virus is detected and exceeds 30%. If virus exceeds 30%, enrichment will stop.
    2. If there is a history of virus in the last 5 years, carcasses will not be distributed outside of the watershed in which the hatchery is located until tests are completed and <30% virus is detected (carcasses will have to be frozen and saved until tests are completed).

## Appendix E. Standard Operating Procedures for Imnaha River Weir

### **Standard Operating Procedures (SOP) for processing fish at the Imnaha River Weir**

**Purpose:** The Imnaha satellite facility is an adult collection and juvenile acclimation and release facility for the Imnaha River Spring/Summer Chinook Salmon Program. The adult weir and trap collect returning Chinook salmon for program broodstock and allow for selective management of natural spawners upstream of the facility. During annual operations, non-target ESA-listed species are also encountered at the weir site. Bull trout are collected and marked for ongoing population monitoring, and post-spawn steelhead kelts are migrating downstream past the weir. This document is intended to provide guidance to operational staff for handling and processing ESA-listed species at the Imnaha facility, with the goal of minimizing stress and risk of latent injury to each individual.

**Annual Coordination:** Imnaha River weir operations are coordinated with funding agencies and co-managers during the Lower Snake River Compensation Plan (LSRCP) annual operations plan (AOP) process. Co-managers agree that the annual planning process will be consistent with this SOP, and meet the intent of this guidance. Significant deviations in annual operations may require broader coordination and an update of the SOP. Prior to each trapping season, staff involved in operating the weir facility should review this document, the AOP, ESA-permits, and any associated Section 7 consultation documents, and understand any constraints regarding changes and/or modifications to procedures pursuant to these other documents.

### **Equipment List by Species**

<u>Chinook Salmon</u>	<u>Bull Trout</u>	<u>Steelhead</u>
<ul style="list-style-type: none"><li>• Large dip net(s)</li><li>• Measuring board</li><li>• PIT tag reader</li><li>• Coded wire tag (CWT) reader</li><li>• Extra batteries (AA)</li><li>• Operculum punches</li><li>• Syringes</li><li>• Needles</li><li>• Antibiotics</li><li>• Sharps container</li><li>• Electronic thermometer</li><li>• Datasheets</li><li>• Scale Cards</li><li>• Snout identification number (SNID)</li></ul>	<ul style="list-style-type: none"><li>• Dip net(s)</li><li>• MS-222</li><li>• Buffer (Baking soda)</li><li>• Measuring board</li><li>• PIT tag reader</li><li>• Electronic thermometer</li><li>• PIT tag injectors</li><li>• PIT tag needles and tag (single use)</li><li>• Recovery tank</li></ul>	<ul style="list-style-type: none"><li>• Long-handled dip nets</li><li>• Anesthesia vessel</li><li>• MS-222</li><li>• Buffer (Baking soda)</li><li>• Scale Cards</li><li>• Genetic samples</li><li>• Measuring board</li><li>• PIT tag reader</li><li>• PIT tagging equipment</li></ul>

### **Trapping operations**

1. The following operational criteria will be in place during ladder operation and brood stock collection at the Imnaha Weir.

- a. A one foot (1.0) difference of water elevation between water in the ladder and in the river. This should correspond to 8.0 ft/second water velocity through the ladder aperture for fish passage and attractant flow into the river. The apparatus gate will be open during the trapping season unless the water elevation of one foot (1.0) cannot be satisfied.
  - b. An RM&E plan with additional operational information will be developed and coordinated through the Imnaha technical working group for implementation prior to 2020 operations (May/June).
2. Based on annual flows, and historical PIT tag detections and travel time of adult Chinook salmon in the Lower Imnaha River from (IR1/IR2 arrays) to the weir, the flexible start date of the weir occurs sometime in early June. This can be moved back or pushed forward based on agreement of the co-managers. The exact installation date will depend on environmental conditions. The weir will be operated until the second Friday in September, or until the last scheduled spawning ground survey for Chinook salmon.
3. The adult weir and trap will be staffed 24 hours per day, 7 days per week, while operational, beginning with the capture of the first Chinook or bull trout.
4. From initial operations through the last week of July, fish will be removed and processed daily.
5. After the last week of July and through the end of operations, fish will be removed and processed daily from Monday-Friday. During this latter period, if 10 or more natural Chinook salmon, or 30 or more fish total, are estimated in the trap on Friday and/or Saturday fish will be removed and processed during the weekend. If a bull trout is observed in the trap on Friday and/or Saturday, all fish will be removed and processed during the weekend.
6. Trap modifications
  - a. Modifications were made to the trap/ladder in 2018-2020 included:
    - i. A new screen was attached to the existing V entrance to help reduce escapement out of the trap.
    - ii. Shade cover over ladder.
    - iii. Improved maintenance of finger weir with varying flows.
    - iv. Another moveable bar was placed on the entrance to the trap to reduce escapement of smaller bull trout.
    - v. A flush gate will be installed in the third step of the ladder to be able to draw down the trap level if necessary.

**Morning trap and weir check** – Each morning, on-site staff at the Imnaha satellite facility will check trap conditions, recover weir and/or trap mortalities, and communicate staffing needs to Lookingglass Hatchery.

1. Measure and record facility temperatures in the raceway using an electronic thermometer. If water temperatures are expected to exceed 16°C (61°F) by 10:00 AM, notify staff of expedited bull trout procedures (see below).
  - a. If temperatures in excess of 16°C (61°F) are anticipated during subsequent processing days, notify Lookingglass Hatchery manager. Processing times may be adjusted to start earlier to avoid higher temperatures for all species.
2. Estimate the numbers of Chinook (by-origin) and bull trout in the trap.
  - a. If more than 30 Chinook salmon are estimated in the trap, assume that at least 10 are natural-origin.

- b. Bull trout can be differentiated from Chinook salmon by the presence of a white leading edge on their fins.
  - c. On Fridays and Saturdays, if trap abundance thresholds are met or a bull trout is encountered notify Lookingglass Hatchery manager to ensure staff is scheduled for weekend processing (*see above*; Trapping Operations, item 4).
- 3. Recover trap and weir mortalities. Ensure passage chutes within weir are unblocked.
  - a. On-site staff will check for Chinook salmon, steelhead, and bull trout mortalities and;
    - i. Inspect the intake and inspect the trap and surrounding walkways (for jump-outs). Remove any mortalities from the trap.
    - ii. Inspect the upstream face of the weir for mortalities, recover if flows safely allow.
    - iii. Inspect the passage chute opening for debris (if flows safely allow). Release/clean panels if debris/bedload may be lodged in chutes.
  - b. Mortality Recovery Procedures
    - 1. Bull Trout: Scan for existing PIT tag, measure fork and total length, and inspect for hooking wounds. Record location of recovery.
      - a. Retain all recovered mortalities in the freezer at the Imnaha satellite facility for further investigation. Immediately notify and provide recovery information to Kyle Bratcher, Wallowa District Fisheries Biologist, Rod Engle LSRCF, and Rick Wilkison IPC. Care must be taken in handling sick or injured fish to ensure effective treatment and care, and in handling dead specimens to preserve biological material in the best possible state. In conjunction with the care of sick or injured fish, or the preservation of biological materials from a dead animal, LSRCF has the responsibility to ensure that information relative to the date, time, and location of the fish when found, and possible cause of injury or death be recorded and provided to the Service.
    - 2. Steelhead: Steelhead monitoring will be discontinued in 2025
    - 3. Chinook salmon: All weir mortalities recovered before the first regularly scheduled spawning ground survey will be processed by on-site staff.
      - a. After the first annual spawning ground survey, which typically occurs the last week in August , weir carcasses should be processed from Friday to Tuesday.
      - b. For each carcass, weir staff will document the fork length (mm), sex (M, F), fin mark (e.g., AD = adipose clip; none = Unclipped), type of opercle mark present on the carcass (e.g., 1LOP, 1ROP), Percent spawn for female carcasses (0% = full of eggs, 100% = completely spawned), and presence/absence of a CWT. If there is a CWT, staff will collect a snout and each snout will receive a unique SNID. Weir staff does not need to collect scales or kidney samples. After the carcass is processed, the tail will be removed, and the carcass tossed back into the river below the weir.
      - c. Before redd surveys: Imnaha staff will collect the first 10 weir mortalities, seal them intact in a labelled plastic bag, freeze them, and notify Fish Health that they are ready for pathology sampling. Note: avoid collecting fallbacks or degraded carcasses for Fish Health samples.

**Processing shed setup** – Prior to staff arrival for daily processing, on-site staff at the weir will setup the processing shed and test equipment.

1. Test the tower door and braille to ensure operation before fish are crowded.
2. Inspect hoses, valves, and fittings used to supply water to the processing shed and liberation trucks.
3. Inspect and test oxygen supplies in the holding tank.
4. Setup processing tables and tanks. Place rubber mats around fish processing stations.
5. Ensure that all items listed above (*see*: equipment list by species) are present and operational.
  - a. Test PIT tag reader with a dummy tag.
6. Fill liberation trucks upon arrival. Install electronic thermometers and oxygen sensors, and monitor throughout fish processing.

### **Crowding / Sorting**

1. Begin filling the holding tank. Turn on the oxygen supply and monitor levels with a DO monitor.
2. Herd fish in the trap towards the tower door using long handled dip nets.
3. Lower the mechanical crowder on the west end of the trap.
4. Slowly move the mechanical crowder toward the tower door, monitoring for fish that may get impinged on the trap walls or floor.
5. Depending on trap densities, leave enough space for fish to move around freely in-between loading groups of fish into the tower.
6. While loading fish in the tower, move the mechanical crowder as far as possible towards the east end of the trap. Use a long-handled crowder to herd fish along the trap walls towards the tower door.
  - a. When loading a group into the tower, attempt to target no more than 20 Chinook salmon.
7. Once the trap is full and processing equipment is ready, raise the braille slowly to move fish into the holding tank.
  - a. Attempt to transfer no more than 20 Chinook salmon into the holding tank at one time.
8. The holding tank will have a divider, with a cover, that allows separation between Chinook salmon and bull trout and prevents fish from jumping out of the holding tank.
9. To the best degree possible, bull trout should be isolated from Chinook salmon as soon as possible to minimize interactions. Staff can opportunistically net bull trout as they spill from the tower into the holding tank, or net bull trout directly from the larger portion of the holding tank into the portion reserved for bull trout.
  - a. It is not necessary to separate *every* bull trout before processing begins. Additional stress will be caused by attempting to net fish, so staff should isolate what is immediately possible and separate the rest as Chinook salmon are processed.

### **Bull Trout Processing**

1. **General guidelines**
  - a. Bull trout will normally be processed before Chinook salmon. On high volume days, both species can be processed concurrently if staff resources allow.

- b. Bull trout will be processed in a manner that minimizes time out of water and risk of dropping (i.e., use nets at all times).
- c. Bull trout are thermally-intolerant; therefore, staff will note water temperatures prior to processing trap catch each day. Expedited processing procedures will be followed when water temperatures are expected to exceed 16°C (61°F) before processing is completed.

Normal procedures (≤61 F)	Expedited procedures (>61 F)
<ol style="list-style-type: none"> <li>1. Anesthetize with MS-222</li> <li>2. Buffer anesthetic bath</li> <li>3. Scan for existing PIT tags</li> <li>4. Measure for fork and total length (mm)</li> <li>5. Inspect for previous injuries, including hooking scars</li> <li>6. Mark new captures with PIT tag</li> <li>7. Transfer to release pipe.</li> </ol>	<ol style="list-style-type: none"> <li>1. Scan for existing PIT tags</li> <li>2. Estimate fork length (within 2-inch size class)</li> <li>3. Inspect for previous injuries, including hooking scars</li> <li>4. Transfer to release pipe.</li> </ol>

## 2. Anesthetization

- a. Tricaine methanesulfonate (MS-222) will be used as a primary anesthesia.
- b. Bull trout will be processed in a separate MS-222 tank.
  - i. For MS-222 as an Anesthetic
    1. Bull trout will be placed in separate MS-222 tank for anesthesia. The tank will be monitored for temperature to ensure dosage guidelines are met and/or changed with cool water regularly.
      - a. Dosage = 50-60 mg/L at 13-16C (moderate dosage)
      - b. Buffer with an equivalent dose of baking soda.
      - c. Duration = lack of equilibrium before 5 minutes, up to 5 minutes in solution.
      - d. Recovery of anesthetized bull trout will take place in the recovery channel after being released down the release pipe. (see Section 5 - Release).
      - e. Fish will be observed in the recovery channel several times during processing of fish. Staff is instructed to regularly inspect recovery channel for any impacts/concerns of recovery.

## 3. Marking / PIT-tagging

- a. PIT tagging equipment will be supplied by Idaho Power (IPC) and procedure/guidance provided by IPC staff will be followed for the PIT tagging process.
- b. All bull trout will be scanned for an existing PIT tag using a PIT tag reader. While scanning with a handheld device, emphasize the area of the fishes' body around the dorsal fin (see P4 instructions below).
- c. Record existing PIT tag codes as 'recaptures'.
- d. Measure



- e. Record both fork and total length (mm) for new and recaptured bull trout.
- f. Inspect for injuries or record external appearance and identify wounds
  - i. Wound location: Mouth/maxillary, head, body
  - ii. Size: 1) = less than 1 inches; 2) = 1 to 3 inches; 3) = over 3 inches
  - iii. Record whether wound is healed or unhealed?
- g. For previously unmarked (new) fish, apply a 12mm PIT tag to the dorsal sinus using a preloaded syringe.
- h. Properly dispose used PIT tag needles in a designated sharps container.

#### 4. Release

- a. Release Protocol
  - i. Bull trout will be released into the release channel and allowed to volitionally leave the facility. Fish can be hand netted from the anesthetic tank and released directly into the recovery channel.
    - a. Undesirable impacts/concerns would be observation of fish against the weir, dead in the recovery channel, or any weir mortalities following releases in the channel.
  - ii. Should undesirable impacts/concerns be observed during initial annual operations, handled bull trout will be placed into a recovery tank on a truck and then driven up to intake for release (by hand/net) identical to past operations (2016).

### **Chinook salmon processing**

#### 1. General guidelines

- a. Natural-origin fish will be differentiated from hatchery-origin fish by the presence of an adipose fin.
- b. ODFW staff will be responsible for determining fish disposition on-site. Weekly and/or daily guidance will be provided by the Lookingglass Hatchery Manager and/or Wallowa District staff.
- c. Co-managers have agreed to the following dispositions, in order of priority:

##### Natural-origin

- 1. Broodstock
- 2. Pass above weir for natural spawning

##### Hatchery-origin

- 1. Broodstock
- 2. Pass above weir for natural spawning
- 3. Recycle downstream for fishery contribution
- 4. Tribal distribution and/or foodbank
- 5. Outplant

#### 2. Processing / Marking

- a. Identify the origin of fish (natural or hatchery), estimate sex (male or female), and record the fork length (mm).
  - i. Sex identification will be based on phenotypic traits (snout shape/size and anal vent characteristics).
- b. Scan for a coded wire tag (CWT) near the snout. Note presence or absence of a CWT.

- c. Place fish on measuring/pit tag board and identify if the fish has a pit tag present. If present, scan fish with a handheld PIT reader and record the PIT tag code with other biological information.
- d. Designated staff will select a disposition for each fish based on broodstock and natural spawning objectives for that week.
- e. Salmon selected to spawn upstream will be released in the existing upstream release tube. All other dispositions will be transferred to the respective liberation truck for transport.
  - i. Fish designated for tribal distribution / food bank, or outplanting, may be held on-site until transportation is available. These fish will be held in a net pen in the rearing pond.
  - ii. Fish designated for tribal distribution/food bank, or any mortality (i.e., trap mort) or weir fallback is checked for CWT. For tribal distribution, no more than 25% of hatchery fish will have a snout removed. For Oregon foodbanks, a minimum of 50% of hatchery fish will have the snout removed. When a snout is collected, the sample is placed into a snout bag with a SND and all biological information (length, sex, origin, marks, etc.) is recorded on a separate datasheet along with the SNID.

### 3. Release


- a. After processing, Chinook salmon will be transferred via hand net to a liberation truck.
  - b. The tank will be monitored for oxygen saturation and water temperature while in transfer.
1. **Steelhead Processing** – During the early portion of the trapping season, upstream-migrating steelhead may be encountered in the Imnaha weir trap. Adult steelhead observed in the Imnaha trap box will be handled, processed and passed upstream.
  2. **Procedures**
    - a. Steelhead removed from the trap will be placed in an anesthetic vessel containing a buffered solution of tricaine methanesulfonate (MS-222) at a concentration of 80 mg/L (buffer dose = 80 mg/L baking soda).
    - b. After anesthetization, each fish will be examined for fin-clips, marks and/or tags, measured for fork length (cm), categorized by gender using secondary sexual characteristics, and origin determined (hatchery or natural).
    - c. Scales will be collected from the preferred area of the fish and a 1ROP opercle punch given to each steelhead. The resulting tissue will be retained for future genetic analysis.
    - d. Steelhead without a preexisting PIT tag will receive a tag for future detections on PIT tag arrays and downstream dams (kelts). These fish will be tagged in the cartilage of the pelvic girdle to facilitate tag retention during spawning.
    - e. Steelhead will be recovered in fresh water. After recovery, steelhead will be released in the upstream release tube.

**Downstream migrants** – Chinook salmon, bull trout, and steelhead (kelts) may be observed on the upstream side of the weir. On-site staff will make efforts (within reason) to safely pass steelhead kelts and bull trout downstream, while restricting the downstream movement of Chinook salmon.

1. If bull trout and/or steelhead are observed on the upstream side of the weir, on-site staff will assess feasibility of passing fish downstream.
  - a. Criteria should depend on flows and presence of Chinook salmon. Chinook salmon that are passed above the weir are uniquely marked and abundance is managed. Therefore, all steps should be taken to avoid allowing Chinook salmon downstream.
  - b. On-site staff will monitor the relative presence of fish above the weir, and select the best time of day to pass fish without allowing salmon downstream.
2. Staff will use long-handled nets to herd steelhead and bull trout toward a selected weir panel.
3. The selected weir panel will be temporarily raised to allow bull trout and steelhead downstream.
4. ODFW will document Imnaha weir kelt passage data.

**Dead Kelts** - Steelhead kelts that have died on the weir or floated down as carcasses will be retained for processing (if flows allow for recovery). After examination carcasses will be passed downstream.


CATHERINE CREEK STOCK SPRING CHINOOK  
BROOD YEAR 2023




**GRANDE RONDE STOCK SPRING CHINOOK  
BROOD YEAR 2023**

**Lookingglass Hatchery**


**Upper Grande Ronde Acclimation Facility**



**IMNAHA STOCK SPRING CHINOOK  
BROOD YEAR 2023**



# LOOKINGGLASS STOCK SPRING CHINOOK BROOD YEAR 2023



**LOSTINE STOCK SPRING CHINOOK  
BROOD YEAR 2023**

**Lookingglass Hatchery**

**Lostine Acclimation Facility**

August - September 2023

February 2024

April - May 2023

August - October 2024

April 2025

*Spawning*

*Incubation*

*Rearing*

*Release*

**Spawning Females**


**Egg Trays**

**Canadian Troughs**

**Pre-mark Raceways**

**Post-mark Raceways**

**Release**



- = CWT tagged raceways
- [Solid Blue Box] = 1<sup>st</sup> Acclimated Release
- [Solid Green Box] = 2<sup>nd</sup> Acclimated Release
- [Yellow Box] = Low BKD
- [Red Box] = Medium BKD

\* Calculations do not take mortality or program excess into account, as final release numbers have yet to be determined.

\* Total smolt numbers are not finalized, current as of 11/15/2024