

Biological Opinion

Kentucky Field Office's Revised Conservation Strategy for Forest Dwelling Bats in the Commonwealth of Kentucky 2025



Prepared by:

U.S. Fish and Wildlife Service
Kentucky Ecological Services Field Office
Frankfort, Kentucky

Field Supervisor

Date

Table of Contents

1. Introduction.....	1
2. Proposed Action	2
<i>Action Area</i>	3
<i>Conservation Measures.....</i>	3
<i>Conservation Benefits</i>	4
3. Indiana Bat.....	5
<i>Status of the Species</i>	5
<i>Description of the Indiana Bat</i>	5
<i>Life History of the Indiana Bat</i>	5
<i>Habitat Characteristics of the Indiana Bat.....</i>	7
<i>Numbers, Reproduction, and Distribution of the Indiana Bat.....</i>	9
<i>Conservation Needs of and Threats to the Indiana Bat</i>	11
4. Northern Long-eared Bat.....	13
<i>Status of the Species</i>	13
<i>Description of the Northern Long-eared Bat.....</i>	14
<i>Life History of the Northern Long-eared Bat.....</i>	14
<i>Habitat Characteristics of the Northern Long-eared Bat</i>	16
<i>Numbers, Reproduction, and Distribution of the Northern Long-eared Bat</i>	18
<i>Conservation Needs of and Threats to the Northern Long-eared Bat.....</i>	19
5. Environmental Baseline.....	20
<i>Indiana Bat.....</i>	20
<i>Northern Long-eared Bat</i>	22
6. Effects of the Action.....	23
<i>Effects of Noise and Vibration on the Indiana Bat and NLEB Applicable Science</i>	23
<i>Effects of Aquatic Resource Degradation on the Indiana Bat and Northern</i>	26
<i>Effects of Tree Removal on the Indiana Bat and Northern Long-eared Bat.....</i>	28
<i>Applicable Science – Removal of Summer Habitat (Summer Occupancy Timeframe)</i>	29
<i>Applicable Science – Removal of Summer Habitat (Hibernation Timeframe)</i>	30
<i>Applicable Science – Loss and Fragmentation of Forested Habitats</i>	34
<i>Applicable Science- Removal of Spring Staging/Fall Swarming Habitat</i>	36
<i>Summary of Effects of the Action on the Indiana Bat and Northern Long-eared Bat.....</i>	39
7. Cumulative Effects on the Covered Species	40

8. Conclusion of Effects on the Covered Species	41
9. Incidental Take Statement	44
<i>Indiana Bat</i>	44
<i>Northern Long-eared Bat</i>	45
<i>Reasonable and Prudent Measures</i>	46
<i>Terms and Conditions</i>	47
<i>Monitoring and Reporting Requirements</i>	47
10. Conservation Recommendations	47
11. Reinitiation Notice	48
13. LITERATURE CITED	49

1. Introduction

A biological opinion (BO) is the document that states the opinion of the U.S. Fish and Wildlife Service (Service) under the Endangered Species Act (ESA) of 1973, as amended, as to whether a Federal action is likely to:

- a) jeopardize the continued existence of species listed as endangered or threatened; or
- b) result in the destruction or adverse modification of designated critical habitat.

A BO evaluates the effects of a Federal Action, which include all consequences of the Action and from non-Federal actions unrelated to the proposed Action (cumulative effects), relative to the status of listed species and critical habitat. A Service opinion that concludes a proposed Federal action is not likely to jeopardize species and is not likely to destroy or adversely modify critical habitat fulfills the Federal agency's responsibilities under §7(a)(2) of the ESA.

“Jeopardize the continued existence” means to engage in an action that reasonably would be expected, directly or indirectly, to reduce appreciably the likelihood of both the survival and recovery of a listed species in the wild by reducing the reproduction, numbers, or distribution of that species (50 CFR §402.02).

This BO addresses the effects of the Service's implementation of the Conservation Strategy for Forest-Dwelling Bats in the Commonwealth of Kentucky (Strategy) on the federally endangered Indiana bat (*Myotis sodalis*) and northern long-eared bat (*Myotis septentrionalis*) (NLEB) (the covered species). The Strategy becomes effective as of the signature date of this BO. This Strategy will supersede the Revised 2016 Conservation Strategy for Forest-Dwelling Bats in the Commonwealth of Kentucky. For this intra-Service consultation, the Service's Kentucky Field Office (KFO) is the federal action agency, and the Service's Southeastern Regional Office (RO) is the consulting office.

Currently, the Indiana bat is the only species covered by this BO for which critical habitat has been designated within the Action Area. Impacts to designated critical habitat for the Indiana bat are not part of the Action and have not been considered in this BO. Any project impact to critical habitat for the Indiana bat will require a separate, project specific evaluation.

Section 9 of the Act and regulations issued under section 4(d) of the Act prohibit the taking of endangered and threatened species without special exemption. Federal agencies may obtain such exemption through the “Incidental Take Statement” (ITS) of a BO that supports a non-jeopardy finding for their proposed action. Incidental take results from a federal action but is not the purpose of the action. Incidental take may be allowed when the Service approves such take through an ITS. The ITS includes the amount or extent of anticipated take due to the federal action, reasonable and prudent measures (RPMs) to minimize the take, and terms and conditions (T&Cs) that must be observed when implementing those RPMs.

The Service has determined that the proposed Action may result in the incidental take of the Indiana bat and northern long-eared bat. Incidental take for this Action would be exceeded when **1,440 Indiana bats or 157 NLEBs are incidentally taken due to the removal of 20,000 acres of suitable forested habitat**. Any incidental take occurring as a result of this Action is exempted from the prohibitions of Section 9 of the ESA, provided that the non-discretionary

measures and conservation measures detailed in the ITS (Section 11 of this BO) become binding conditions of any permit, contract, or grant issued for implementing the Action.

2. Proposed Action

The Action is the KFO's implementation of the Strategy for an indefinite number of projects with Federal and non-Federal entities that may affect one or both covered species. The Strategy offers voluntary ESA compliance options that allow entities to ensure their actions: (1) do not jeopardize the continued existence of a covered species; and (2) obtain an exemption for take prohibitions, as provided in Section 10 of this BO, for a covered species that is incidental to their actions. The KFO participates in the Strategy and is responsible for determining if a specific project meets the requirements for voluntary participation, as well as implementing certain, predetermined conservation measures associated with the Strategy. Implementation of the Strategy provides recovery-focused conservation benefits for the covered species as mitigation for the removal of up to 20,000 acres of suitable forested habitat over a 10-year period.

The types of impacts to the covered species that are addressed in this BO and associated with the Strategy are limited to those adverse effects caused by the removal or alteration of suitable forested habitats that are reasonably certain to occur based on information provided during the technical assistance/informal consultation process. Forested habitat loss or alteration associated with the Strategy may be permanent or temporary and may occur while the covered bat species are likely to be present (occupied) or during the hibernation period when covered bat species are not expected to be present (unoccupied). Impacts may occur in areas where a covered species has been documented (known habitat) or where the presence of one or both covered species is presumed present (potential habitat) (i.e., reasonable certainty of occurrence has been determined using modeled occupancy data). Further, known habitats for either species may include known summer habitat and fall swarming/spring staging habitat around known hibernacula. All suitable forested habitats are considered to have reasonable certainty of species presence for both summer and fall swarming/spring staging use unless probable absence of the species for that habitat type has been demonstrated (e.g., through approved surveys). The KFO will limit impacts to no more than a cumulative total of 2,000 acres of suitable forested habitat removal per year for all covered species.

Additional coordination with the KFO will be required for any action expected to result in impacts to a covered species that were not evaluated in this BO, including any action expected to result in impacts to designated critical habitat of a covered species.

The KFO has chosen to exclude projects that could impact known or potential hibernacula due to their importance for conservation of the covered species and the difficulty in analyzing effects related to potential hibernaculum impacts. Additionally, project-specific review by the KFO will be required for the categories of projects listed below to determine if use of the Strategy is appropriate or if another ESA compliance option should be used:

- Individual projects resulting in the loss of more than 100 acres of forested habitat for any of the covered species.

- Projects occurring within one mile of Priority 1 (P1) or Priority 2 (P2) Indiana bat hibernacula.¹
- Projects occurring within ½ mile of Priority 3 (P3) or Priority 4 (P4) Indiana bat hibernacula or any northern long-eared bat hibernacula.¹
- Clearing of greater than 10 acres of forested habitat during the pup season.
- Projects that may result in adverse effects to the covered species from post-construction and/or operational activities other than loss of forested habitat (e.g., permanent lighting, noise and vibrations above pre-construction levels, prescribed fire).

Action Area

For purposes of consultation under ESA §7, the “Action Area” is defined as “all areas to be affected directly or indirectly by the Federal action and not merely the immediate area involved in the action” (50 CFR §402.02). The Action Area for this consultation includes all lands within the geo-political boundaries of the Commonwealth of Kentucky and those portions of Missouri, Illinois, Indiana, Ohio, West Virginia, Virginia, and Tennessee that occur within 20 miles of the Kentucky state line. Execution of multi-state projects utilizing the Strategy will require advance, written approval of the Service Field Office(s) involved.

This Action Area corresponds with the scope of the Strategy and recognizes that projects associated with the Strategy: (a) are likely to occur at scattered and undeterminable locations across Kentucky; (b) may cross into adjacent states; and (c) will vary in size and distribution on the landscape.

Conservation Measures

Conservation measures are proposed actions that will be undertaken by project proponents as part of the Action to benefit, promote the recovery of, and/or minimize effects to species affected by the Action.

Avoidance and Minimization Measures

Project proponents are encouraged to implement the following conservation measures, where applicable and appropriate, to minimize impacts to covered species. While implementation of these conservation measures would not necessarily remove the adverse effects expected from the Action, incorporation will minimize the effect of take on the covered species and reduce the required compensatory mitigation, saving the proponent time and money.

- Reduce the amount of suitable forested habitat removed by the project.
- Remove suitable trees between November 15 and March 31 (or October 16 and March 31 if outside known swarming habitat) when forested habitats are unoccupied by the covered species.
- Avoid removing suitable forested habitat during the pup season (May 15 – July 31).

¹Priority 1 (P1) hibernacula have a current or historical winter population of ≥10,000 Indiana bats; Priority 2 (P2) have 1,000-9,999 bats; Priority 3 (P3) have 50-999 bats; and Priority 4 (P4) have <50 bats (USFWS 2007).

- Avoid removing suitable forested habitat within one mile of a known Indiana bat P1 or P2 hibernaculum and ½ mile of a known Indiana bat P3 or P4 hibernaculum.
- Avoid removing suitable forested habitat within ½-mile of a known northern long-eared bat hibernaculum.
- Use Best Management Practices (BMPs) for sediment and erosion control in accordance with state permits, program requirements, and regulations during tree removal and construction.

Compensation Measures

The Strategy identifies compensatory mitigation options that project proponents may implement or fund to assist in the conservation and/or recovery of the covered species within the Action Area. Use of the Strategy becomes part of the project proponent's proposed action and requires that at least one of the compensatory mitigation options listed below is implemented as part of their proposed action:

- (a) protection of unprotected known habitat with a demonstrated significance to the covered species;
- (b) contribution of funds to Kentucky's Imperiled Bat Conservation Fund (IBCF) sufficient to offset the identified impacts when other measures are impractical or have limited value to the covered species' conservation and/or recovery;
- (c) purchase credits from an approved species conservation bank;
- (d) other activities that provide a tangible conservation benefit to the covered bat species proposed to the KFO for case-by-case evaluation.

The KFO proposes to participate in and implement the Strategy with the following limitations:

Projects implemented in association with the Strategy will impact no more than a cumulative total of 20,000 acres (not to exceed 2,000 acres annually) of known and/or potential Indiana bat habitat and/or northern long-eared bat habitat.

Conservation Benefits

As described in the Strategy, the KFO has identified conservation goals for the covered species based on recovery plans, published guidance, literature, and/or the best available scientific information. Proponents incorporating the Strategy will provide mitigation for impacts from their project to the covered species being impacted. This mitigation is anticipated to exceed (at a programmatic scale) what is needed to compensate for impacts to the species, yielding a net conservation benefit or gain. Conservation benefits, as well as impacts, are tracked by acres and the habitat type (e.g., staging/swarming, maternity), as it is the loss of suitable forested habitat that drives the adverse effects evaluated in this BO.

Conservation benefits may be achieved directly through a project proponent's specific actions or indirectly through the IBCF or a conservation bank. Regardless of the mechanism used, all conservation benefits will be aligned with the goals identified in the Strategy. These goals have been established to maximize the benefits to the species by targeting actions that will help protect and manage important habitat components for the covered species, especially during the most sensitive life-stages.

3. Indiana Bat

This section summarizes the best available data about the biology and current condition of the Indiana bat throughout its range that are relevant to formulating an opinion about the Action.

Status of the Species

The Service published its decision to list the Indiana bat as endangered on March 11, 1967 (Federal Register 32[48]:4001) under the Endangered Species Preservation Act of October 15, 1966 (80 Stat. 926; 16 U.S.C. 668aa[c]). The ESA of 1973 (87 Stat. 884, as amended; 16 U.S.C. 1531 et seq.) subsequently extended full legal protection from unauthorized take to the species. Critical habitat was designated for the species on September 24, 1976 (41 FR 14914). Thirteen hibernacula, including 11 caves and two mines in six states, were listed as critical habitat.

Description of the Indiana Bat

The Indiana bat is a temperate, insectivorous, migratory bat that hibernates in caves and mines in the winter and summers in forested areas. It is a medium-sized bat, having a wingspan of nine to 11 inches and weighing only one-quarter of an ounce. It has brown to dark-brown fur, and the facial area often has a pinkish appearance. The Indiana bat closely resembles the little brown bat (*Myotis lucifugus*) and the northern long-eared bat. It is distinguished from these species by its foot structure and fur color. The Indiana Bat Draft Recovery Plan (USFWS 2007) provides a comprehensive summary of the description of the species and is incorporated by reference.

Life History of the Indiana Bat

The Indiana bat hibernates in caves and mines in the winter (typically October through April) and migrates to forested summer habitat. When arriving at their traditional hibernacula from August to October, Indiana bats “swarm” for several weeks prior to hibernation. Some male bats may begin to arrive at hibernacula as early as July, but females typically arrive later. The time of greatest swarming activity in Indiana and Kentucky has been documented as early as September (Cope and Humphrey 1977). Swarming is a critical part of the life cycle when Indiana bats converge at hibernacula, mate, and forage until sufficient fat reserves have been deposited to sustain them through the winter (USFWS 1983). Swarming behavior typically involves large numbers of bats flying in and out of cave entrances throughout the night, while most of the bats continue to roost in trees during the day (Cope and Humphrey 1977). Body weight may increase by two grams within a short time, mostly in the form of fat. Copulation occurs on cave ceilings near the cave entrance during the latter part of the swarming period (USFWS 2007). Females may mate their first autumn, whereas males may not mature until the second year (USFWS 2007). By late September, many females have entered hibernation, but males may continue swarming well into October in what is believed to be an attempt to breed with late arriving females.

The initiation of hibernation may vary by latitude and annual weather conditions; however, most bats are hibernating by the end of November (USFWS 2007). Hibernation facilitates survival during winter when insect prey is unavailable. Hibernating Indiana bats cluster on cave ceilings

in densities of approximately 300-484 bats/ft² from approximately October through April. Like other cave bats, the Indiana bat naturally arouses during hibernation (Sealander and Heidt 1990). Arousal s are more frequent and longer at the beginning and end of the hibernation period (Sealander and Heidt 1990). Limited mating occurs throughout the winter and in early April as bats emerge (USFWS 2007).

Spring emergence occurs when outside temperatures have increased, and insects (prey) are more abundant (Richter et al. 1993). Most Indiana bats emerge in late March or early April; however, the timing of annual emergence may vary across the range, depending on latitude and annual weather conditions. Females emerge before males. Shortly after emerging from hibernation, the females become pregnant via delayed fertilization from the sperm that has been stored in their reproductive tracts through the winter (USFWS 2007). During the “staging” period, the bats forage for a few days or weeks near their hibernacula before migrating to their traditional summer roosting areas. Most populations leave their hibernacula to migrate to summer habitat by late April.

Initially, published literature indicated that Indiana bats migrate north for the summer maternity season (USFWS 2007, Gardner and Cook 2002). However, recent migration studies also document lateral and southward migrations (Copperhead 2017, Copperhead 2018, Copperhead 2019, Roby et al. 2019). Some reproductive females have been documented to migrate up to 418 miles (Butchkoski and Bearer 2016) to form maternity colonies, while others form maternity colonies within only a few miles of their hibernacula (U.S. Army Garrison Fort Drum 2011). Males are commonly found roosting near the hibernacula but have also been documented to migrate long distances to their summer habitat (Kurta and Rice 2002). Migration is stressful for the Indiana bat, particularly in the spring when their fat reserves and food supplies are low. As a result, adult mortality may be the highest in late March and April.

Female Indiana bats, like most temperate members of the family Vespertilionidae, give birth to one pup each year (Mumford and Calvert 1960, Humphrey et al. 1977, Thomson 1982). The proportion of female Indiana bats that produce offspring is not well documented. At a colony in Indiana, 23 of 25 female Indiana bats produced volant juveniles during the first year of the study, and 23 of 28 females the following year (Humphrey et al. 1977). Based on cumulative mist-netting captures over multiple years, Kurta and Rice (2002) estimated that 89% of adult females in Michigan maternity colonies were in reproductive condition (e.g., pregnant, lactating, or post-lactating). Mist-net surveys over multiple years at three post-WNS Kentucky maternity colonies indicated that 92% of adult females exhibited signs of reproduction. Although reproductive proportion trended lower in post-WNS captures (92% compared to 99% pre-WNS), no significant difference was found between pre- and post-WNS reproductive proportions (Rogers 2020).

Racey (1982) notes that a particular ratio of fat to lean mass is normally necessary for puberty and the maintenance of female reproductive activity in mammals. He suggests further that the variation in the age of puberty in bats is due to nutritional factors, possibly resulting from the late birth of young and their failure to achieve threshold body weight in their first autumn. Once puberty is achieved, reproductive rates frequently reach 100% among healthy bats of the family Vespertilionidae, and young, healthy female bats can mate in their first autumn as long as their prey base is sufficient to allow them to reach a particular fat to lean mass ratio.

Studies by Belwood (2002) show asynchronous births among members of a colony. This results in variation in size of juveniles (newborn to almost adult size young) in the same colony. Young Indiana bats are capable of flight within a month of birth. Young born in early June may be flying as early as the first week of July (Clark et al. 1987), with others flying from mid- to late-July. Mortality between birth and weaning was found to be about 8% (Humphrey et al. 1977).

Indiana bats feed on aquatic and terrestrial invertebrates. Diet varies seasonally and among different ages, sexes, and reproductive status (USFWS 1999). Numerous foraging habitat studies have found that Indiana bats forage in closed to semi-open forested habitats and forest edges located in floodplains, riparian areas, lowlands, and uplands. Old fields and agricultural fields are also used (USFWS 2007; Sparks et al. 2005). Indiana bats frequently forage along riparian corridors and obtain water from streams, ponds, and water-filled road ruts in forest uplands.

The average life span of the Indiana bat is five to 10 years, but banded individuals have been documented living as long as 14 and 15 years (Humphrey and Cope 1977). Using winter sampling of unknown-age bats over a 23-year period, Humphrey and Cope (1977) estimated annual survival. Female survivorship in an Indiana population was 76% for ages one to six years and 66% for ages six to 10 years. Male survivorship was 70% for ages one to six years and 36% for ages six to 10 years. Following 10 years, the survival rate for females dropped to only 4% (Humphrey and Cope 1977).

Habitat Characteristics of the Indiana Bat

Winter Habitat

Indiana bats roost in caves or mines with configurations that provide a suitable temperature and humidity microclimate (Brack et al. 2003, USFWS 2007). Requirements for hibernacula are discussed in the draft Recovery Plan for the species (USFWS 2007).

Summer Habitat

During summer, Indiana bats (males and females) use forested habitat for roosting, foraging, and commuting. Indiana bats are often associated with floodplain or riparian forests with large trees, scattered canopy gaps, and open understories (USFWS 2007). Research has showed adaptability in habitats used, including upland forests, forests altered by grazing, swine feedlots, row-crops, hay fields, residences, clear-cut harvests, and shelterwood cuts (Garner and Gardner 1992, USFWS 1999).

Suitability of a roost tree is determined by its condition (dead or alive), suitability of loose bark, solar exposure, spatial relationship to other trees, and its spatial relationship to water sources and foraging areas. Potentially suitable roost trees can be trees of any species with bark separating from the tree after the tree dies, senesces, or is injured. Live trees that exhibit peeling or shaggy bark, such as hickories (*Carya* spp.) and large white oaks (*Quercus alba*), may also be suitable roost trees. Many maternity colonies have been associated with oak-hickory and elm-ash-cottonwood forest types. Tree cavities, hollow portions of tree boles or limbs, and crevices and splits from broken tops occasionally have been used as roosts, usually by individual bats. Roost longevity is variable due to many factors, such as the rate at which bark sloughs off or the tree falls down. Some roosts may only be habitable for one to two years while species with good bark retention, such as slippery elm (*Ulmus rubra*), cottonwood (*Populus deltoides*), green ash (*Fraxinus pennsylvanica*), and various oaks (*Quercus* spp.) and hickories (*Carya* spp.) may provide habitat for four to eight years (USFWS 1999).

Trees over 40 cm (15.7 in) in diameter at breast height (dbh) are optimal for maternity colonies, while trees in excess of 22 cm (8.6 in) in dbh serve as alternate roosts (USFWS 2002). Females have been documented using roost trees as small as 14 cm (5.5 in) in dbh (Kurta 2005). The average size of a male roost is typically smaller than that of female maternity colonies; for example, one male was observed in a roost tree measuring only 6.4 cm (2.5 in) in dbh (Gumbert et al. 2002).

Maternity colonies have been documented to use eight to 34 roost trees per season (Callahan et al. 1997, Kurta et al. 2002; Roby pers comm 2024). The extent and configuration of the roosting area is probably determined by availability of suitable roost trees in a given area. Distances between roosts can be a few meters to a few kilometers (Kurta et al. 1996, 2002). Primary roosts are generally larger in diameter and located in openings or at the edge of forest stands, while alternate roosts can either be in openings or the interior of the forest stand. Maternity colony movements among multiple roosts seem to depend on micro-climate changes, particularly solar radiation (Humphrey et al. 1977). Cool temperatures can delay fetal development and growth of juveniles; therefore, selection of maternity roost sites may be critical to reproductive success. Kurta et al. (1993) suggest movement between roosts may be the way that bats deal with the ephemeral nature of roost trees. It is not known how many alternate roosts must be available to ensure retention of a colony within a particular area, but large, nearby forest tracts would improve the potential for an area to provide adequate roosting habitat (Callahan 1993, Callahan et al. 1997).

Information to date indicates that Indiana bats predominately forage, roost, and travel within wooded habitats or along their edges (Cope et al. 1974, Humphrey et. al., 1977, LaVal et al., 1977, LaVal and LaVal 1980, Gardner et. al., 1991a and b, Hobson and Holland 1995, Kiser and Elliot 1996, Butchkosi and Hassigner 2002, Rommé et al. 2002, Murray and Kurta 2004, Menzel et al. 2005, Sparks et al., 2005). Use of other habitat types appear to be infrequent relative to availability (Garner and Gardner 1992, Menzel et al., 2005, Sparks et al. 2005). The observations of Murray and Kurta (2004) indicate that Indiana bats will avoid traveling in open areas; of the 34 transmitter nights (= a single bat monitored through one night), no bats were detected crossing open areas but rather predictably, over 5 years, used a single tree-lined corridor to move from their roosting to foraging areas. Avoiding these open areas increased the distance bats needed to fly by up to 55 percent (=0.2 to 3.4 km extra distance flown) more than if they had taken a straight-line flight from their day roosts to their foraging areas. Similarly, investigators in Missouri (Ecology and Environment, Inc. 2009) found that the areas of activity for five radio-tagged bats were in heavily forested areas and along riparian corridors and forest edges. No bats were recorded in the open areas interspersed throughout the research area.

Conversely, research has shown that Indiana bats will cross open areas to travel between roosting and foraging habitat (Brack and Sparks, pers. comm. 2011). Adult females and volant juveniles from multiple maternity colonies in Kentucky have been documented flying back and forth across the Ohio River to forage and roost (Seiter et al 2022). The Ohio River's average width in these areas is approximately 1,500 feet. Whitaker and Brack (2006) documented a maternity roost in an isolated 0.7 ha woodlot where the closest woody habitat was a brushy fencerow of small trees 160 m (525 ft) away. Similarly, three years of radiotelemetry study on a maternity colony of Indiana bats in an agricultural landscape of Ohio documented Indiana bats often crossing open areas greater than 1 km in length (Kniowski 2011).

Given this information, it is reasonable to assume that Indiana bats may use edge habitat and seemingly isolated tracts in close proximity to occupied habitat but will rarely fly over large open areas. The distance from the forest edge (i.e., the area of non-wooded habitat) that Indiana bats are likely to travel through is unknown. We are aware of few studies that provide specific data on “capture distances from the forest edge.” Data garnered thus far by Stantec et al. (2010) show that of the 1124 foraging telemetry points from 21 radio-tagged Indiana bats, the vast majority (75 percent) of points were within 400 feet of a forest edge and 97 percent were within 1000 ft of forest edge. Drawing from all existing data, it is reasonable to conclude that Indiana bats are unlikely to occur within project areas located more than 1000 feet from wooded areas.

Home range size estimates for Indiana bat vary greatly among studies. Home range size may vary between seasons, sexes, and reproductive status of the females (Lacki et al. 2007).

MacGregor et al. (1999) found that males that choose to remain near their hibernacula ranged from 10 to 568 ha (25-1,404 ac). Other studies using radio telemetry tagging and various analysis methods (i.e., mean convex polygons, 95% adaptive kernel, 95% fixed kernel) have estimated average individual Indiana bat summer home range sizes of 205–828 ac (Menzel et al. 2005; Sparks et al. 2005; Watrous et al. 2006; Jachowski et al. 2014; Kniowski and Gehrt 2014).

Jachowski et al. (2014) found home range size averaged 101 ha (249 ac) for four males and 145 ha (358 ac) for eight females. Menzel et al. (2005) found no significant difference in male and female home range size in Illinois. Womack et al. (2013) found Missouri home ranges averaged 1,137 ha (2,809 ac) and lactating and pregnant females did not significantly differ in home range. Bergeson et al. (2013) analyzed a mean of 375 ha (927 ac) and 285 ha (704 ac) foraging range for females in Illinois using two different analysis techniques. Watrous et al. (2006) calculated a mean home range of 83 hectares (205 acres) for 14 female Indiana bats in Vermont. Without site-specific data, the Service generally considers the potential home range for an Indiana bat to include all suitable habitat within 4 km (2.5 mi) of documented roost(s) (USFWS 2011), recognizing the area of actual use may be just a portion of that area.

Indiana bats show a high degree of fidelity to roost trees, roosting areas, and foraging areas (Gardner et al. 1991; Humphrey et al. 1977; Kurta et al. 1996, 2002; Kurta and Murray 2002; Gumbert et al. 2002). Bats using familiar foraging and roosting areas are thought to benefit from decreased susceptibility to predators, increased foraging efficiency, and the ability to switch roosts in case of emergencies or alterations surrounding the original roost (Gumbert et al. 2002).

Spring and Fall Habitat

In the spring and fall, Indiana bats usually roost, forage, and commute in habitats similar to those selected during the summer. These areas are most typically within 10 miles of a P1/P2 hibernaculum² and five miles of a P3/P4 hibernaculum; however, use of habitat areas that are farther from hibernacula have been documented (Kiser and Elliot 1996; MacGregor et al. 1999; Rommé et al. 2002; Hawkins et al. 2005).

Numbers, Reproduction, and Distribution of the Indiana Bat

Indiana bats are found over most of the eastern half of the United States. Winter surveys in 2024 found hibernating Indiana bats dispersed across 15 states. However, over 95% of the estimated

² Priority 1 (P1) hibernacula have a current or historical winter population of $\geq 10,000$ Indiana bats; Priority 2 (P2) have 1,000-9,999 bats; Priority 3 (P3) have 50-999 bats; and Priority 4 (P4) have <50 bats (USFWS 2007).

range-wide population hibernated in four states – Missouri (37.6%), Indiana (37.1%), Illinois (13.2%), and Kentucky (7.9%) (USFWS 2024). Summer distribution of the Indiana bat occurs throughout a wider geographic area than its winter distribution. Most summer occurrences are from the upper Midwest including southern Iowa, northern Missouri, much of Illinois and Indiana, southern Michigan, Wisconsin, western Ohio, and Kentucky. However, many summer maternity colonies have been found in the northeastern states of Pennsylvania, Vermont, New Jersey, New York, West Virginia, and Maryland. Maternity colonies have also been found in the south, including southern Arkansas, Georgia, Alabama, Mississippi (Copperhead 2017, Copperhead pers. comm. 2014), southwestern North Carolina (Britzke et al. 2003, USFWS 2007). Non-reproductive summer records for the Indiana bat have also been documented in eastern Oklahoma, northern Mississippi, Alabama, and Georgia.

The data regarding Indiana bat abundance prior to federal listing are limited, but the available information, summarized in the draft Recovery Plan (USFWS 2007), suggests that Indiana bats were once far more abundant than they were in the 1960s. When the Indiana bat was originally listed as endangered in 1967, there were an estimated 883,300 bats, and most of these hibernated in a small number of hibernacula (Clawson 2002). Since the species was listed, its population numbers continued to decline through approximately 2001, with large population declines observed at hibernacula in Kentucky and Missouri. The range wide population estimate dropped approximately 57% from 1965 to 2001 (USFWS 2007). The range-wide, biennial population estimates increased from 2001 to 2007, indicating that the species' long-term decline had been arrested and likely reversed (USFWS 2024). However, the arrival of white-nose syndrome (or "WNS"; see discussion below) is the probable cause of the decline observed since 2007. The Service estimated the 2024 range-wide population at 631,786 bats (Figure 1) occurring in 194 hibernacula in 15 states, with the three most populous states being Missouri (237,733), Indiana (234,657), and Illinois (83,304). This represents a 2.8% decrease from the range-wide population estimate of 664,637 in 2007 (when WNS began). However, there has been an 8.3% increase since the 2022 survey.




Figure 1. Indiana bat range-wide population estimates from 1981 to 2022.

Conservation Needs of and Threats to the Indiana Bat

Destruction/Degradation of Hibernacula

Well-documented examples of modifications to Indiana bat hibernacula that affected the thermal regime of each cave and, thus the ability of the caves to support hibernating Indiana bats, is summarized in the draft revised Recovery Plan (USFWS 2007). Generally, threats to the integrity of hibernacula have decreased since the time that Indiana bats were listed as endangered under the ESA. Increasing awareness of the importance of cave microclimates to hibernating bats and regulatory authorities under the ESA have reduced, but not eliminated, threats to hibernacula. In addition to purposeful modifications, there are threats from stochastic events (e.g., collapse in mines, flooding).

Loss/Degradation of Forested Habitat

Loss of forest cover and degradation of forested habitats have been cited as contributing to the decline of Indiana bats (USFWS 1983, Garner and Gardner 1992, Drobney and Clawson 1995, Whitaker and Brack 2002). Throughout the range of the Indiana bat, there is less forest now than before European settlement (Smith et al. 2003), particularly within the core of the species' range in the Midwest. Conversion to agriculture has been the largest single cause of forest loss. The conversion of floodplain and bottomland forests, recognized as high quality habitats for Indiana bats, has been a particular cause of concern (Humphrey 1978). Since the 1950s, some marginal farmlands have been abandoned and allowed to revert to forest, contributing to a net increase in forest within the range of the Indiana bat, particularly in the Northeast (Smith et al. 2003). Forest cover has also increased within the Midwest Recovery Unit (Smith et al. 2003). Not only has the amount of forest cover increased since the 1950s, but also the average diameter of trees has increased (Smith et al. 2003), which may equate to an increased supply of suitable roost trees for Indiana bats.

Urbanization and development are currently the greatest contributors to forested habitat loss within the range of the Indiana bat (Wear and Greis 2002; U.S. Forest Service (USFS) 2005, 2006), which results in permanent conversion to land uses generally unsuitable for Indiana bats. At a study site in central Indiana, Indiana bats avoided foraging in a high-density residential area (Sparks et al. 2005), although maternity roosts have been found in low-density residential areas (Belwood 2002). Duchamp (2006) found that urban land use was inversely related to bat species diversity in north-central Indiana; several bat species, including the Indiana bat, were less likely to occur in landscapes with greater amounts of urban and suburban development. Development directly destroys habitat and fragments remaining habitat.

Forest cover is not a completely reliable predictor of where Indiana bat maternity colonies will be found on the landscape (Farmer et al. 2002). Indiana bat maternity colonies occupy habitats ranging from completely forested to areas of highly fragmented forest. Nonetheless, trends in forest cover are of interest relative to Indiana bats, with increasing forest cover suggesting at least the potential for improved habitat conditions. Conversely, in areas where almost all forest land has been lost, the absence of woodlands on the landscape certainly equates to less habitat than in prehistoric and early historic periods.

Forest conversion is expected to increase throughout the Indiana bats' range due to commercial and urban development, energy production and transmission, and natural changes. The 2010 Resources Planning Act Assessment projects forest losses of 6.5-13.8 million hectares (16–34 million acres) (or 4–8% of 2007 forest area) across the conterminous United States. Forest loss is

expected to be concentrated in the southern United States, with losses of 3.6-8.5 million hectares (9–21 million acres) (USFS 2012). Forest conversion causes loss of potential habitat, fragmentation of remaining habitat, and if occupied at the time of the conversion, direct injury or mortality to individuals.

Disturbance of Hibernating Bats

The original recovery plan for the species stated that human disturbance of hibernating Indiana bats was one of the primary threats to the species (USFWS 1983). The primary forms of human disturbance to hibernating bats result from cave commercialization (cave tours and other commercial uses of caves), recreational caving, vandalism, and research-related activities. Progress has been made in reducing the number of caves in which disturbance threatens hibernating Indiana bats, but the threat has not been eliminated. Biologists throughout the range of the Indiana bat were asked to identify the primary threat at specific hibernacula, and “human disturbance” was identified as the primary threat at 41% of P1, P2 and P3 hibernacula combined.

White-nose Syndrome

WNS is an infectious wildlife disease caused by a fungus of European origin, *Pseudogymnoascus destructans* (Pd), which poses a considerable threat to hibernating bat species throughout North America, including the Indiana bat. White-nose syndrome is responsible for unprecedented mortality of insectivorous bats in eastern North America (Blehert et al. 2009; Turner et al. 2011). Since the disease was first observed in New York in 2007 (later biologists found evidence from 2006 photographs), WNS has spread rapidly in bat populations from the East to the Midwest and the South.

WNS may cause behavioral changes in infected individuals. For example, at some WNS-affected sites, a shift of hibernating bats from traditional winter roosts to roosts unusually close to hibernacula entrances has been observed. Bats have also been observed flying outside of hibernacula during winter (often during the day) at some affected sites. At some sites, bat carcasses (particularly of the little brown bat) have been found outside affected hibernacula. Many infected bats do not survive the winter. The exact processes by which the fungal skin infection leads to death are not known; but depleted fat reserves (i.e., starvation) contribute to mortality (Reeder et al. 2012, Warnecke et al. 2012), and dehydration may also have a role (Willis et al. 2011, Cryan et al. 2013, Ehlman et al. 2013). It is also suspected that some of the affected bats that survive hibernation emerge in such poor condition that they die soon after emergence or during the summer. Among those bats that do survive, it appears that productivity of female survivors may be negatively affected (Franci et al. 2012; Pettit and O’Keefe 2017).

The Northeast Recovery Unit, where WNS was first observed in the winter of 2006-2007, lost over 70% of its Indiana bats between 2007 and 2015. At the time dead bats were first observed in the winter of 2006-2007, it is not known how long the (previously unidentified) fungus, Pd, had been present in affected sites. Based on subsequent observations as WNS spread, it appears that the arrival of the fungus in an area may precede large-scale fatality of bats by several years. Between 2011 and 2015, the Appalachian Recovery Unit, where WNS was confirmed in the winter of 2008-2009, declined by 84% and continues to decline. The Midwest Recovery Unit, where WNS was confirmed in the winter of 2010-2011, declined by 16% between 2011 and 2015. The Ozark-Central Recovery Unit, where WNS was confirmed in the winter of 2011-2012, declined by less than 1% between 2013 and 2015. However, populations in the Midwest and Ozark-Central Recovery Units have increased by 10.2 and 16.9%, respectively, since 2015.

As of 2016, WNS or Pd was confirmed in all states within the species' range. Further declines in Indiana bat populations from the disease may occur in the future.

Environmental Contaminants

With the restrictions on the use of organochlorine pesticides in the 1970s, this significant threat to Indiana bats was reduced. However, cholinesterase-inhibiting insecticides, organophosphates, and carbamates have now become the most widely used insecticides (Grue et al. 1997), and the impact of these chemicals on Indiana bats is not currently known. Because of the unique physiology of bats in relation to reproduction, high energy demands and sophisticated thermoregulatory abilities, much more research needs to be done with these pesticides and their effects on bats. These and other contaminants likely remain a significant and poorly understood threat to Indiana bats. USFWS (2007) summarizes known and suspected contaminant threats to bats.

Climate Change

The capacity of climate change to result in changes in the range and distribution of wildlife species is recognized, but detailed assessments of how climate change may affect specific species, including Indiana bats, are limited. During winter, only a small proportion of caves provide the right conditions for hibernating Indiana bats because of the species' very specific temperature requirements. Surface temperature is directly related to cave temperature, so climate change that involves increased surface temperatures will inevitably affect the suitability of hibernacula. Impacts on the availability or timing of emergence of insect prey are also likely. Loeb and Winters (2013) modeled potential changes in Indiana bat summer maternity range within the United States; in their model, the area suitable for summer maternity colonies of Indiana bats were forecasted to decline significantly.

Wind Turbines

There is growing concern that Indiana bats (and other bat species) may be threatened by the recent surge in construction and operation of wind turbines across the species' range. Not all facilities conduct fatality monitoring and, even when monitoring is conducted, only a small proportion of dead bats are likely to be found. Based on this information, it is likely that additional Indiana bat mortality has occurred at these facilities and at other wind facilities throughout the range of the species.

4. Northern Long-eared Bat

This section summarizes the best available data about the biology and current condition of the northern long-eared bat throughout its range that are relevant to formulating an opinion about the Action.

Status of the Species

The Service published its decision to list the northern long-eared bat as endangered on November 29, 2022 (87 FR 73488). No critical habitat has been designated for this species.

Description of the Northern Long-eared Bat

The northern long-eared bat is a temperate, insectivorous, migratory bat that hibernates in caves and mines in the winter and summers in forested areas. Adult body weight averages five to eight grams (0.2 to 0.3 ounces), with females tending to be slightly larger than males (Caceres and Pybus 1997, p. 3). Average body length ranges from 77 to 95 mm (3.0 to 3.7 inches). (Barbour and Davis 1969, p. 76; Caceres and Barclay 2000, p. 1). The fur is medium to dark brown on the dorsal side and tawny to pale brown on the ventral side, with dark brown (but not black) ears and wing membranes (Nagorsen and Brigham 1993, p. 87; Whitaker and Mumford 2009, p. 207). As indicated by its common name, the northern long-eared bat is distinguished from other *Myotis* species by its relatively long ears (average 17 mm (0.7 in); Whitaker and Mumford 2009, p. 207) that, when laid forward, extend beyond the nose up to five mm (0.2 in; Caceres and Barclay 2000, p. 1). The tragus (i.e., projection of skin in front of the external ear) is long (average 9 mm [0.4 in]; Whitaker and Mumford 2009, p. 207), pointed, and symmetrical (Nagorsen and Brigham 1993, p. 87; Whitaker and Mumford 2009, p. 207). Within its range, the species can be confused with the little brown bat or the western long-eared myotis (*Myotis evotis*). The northern long-eared bat can be distinguished from the little brown bat by its longer ears, tapered and symmetrical tragus, slightly longer tail, and less glossy pelage (Caceres and Barclay 2000, p. 1; Kurta 2013, in litt.). The northern long-eared bat can be distinguished from the western long-eared myotis by its darker pelage and paler membranes (Caceres and Barclay 2000, p. 1).

Life History of the Northern Long-eared Bat

The northern long-eared bat hibernates in caves and mines in the winter (typically October through April) and migrates to forested summer habitat. The swarming season occurs between the summer and winter seasons (Lowe 2012, p. 50), and the purpose of swarming behavior may include: introduction of juveniles to potential hibernacula, copulation, and stopping over sites on migratory pathways between summer and winter regions (Kurta et al. 1997, p. 479; Parsons et al. 2003, p. 64; Lowe 2012, p. 51; Randall and Broders 2014, pp. 109–110). During this period, heightened activity and congregation of transient bats around caves and mines is observed, followed later by increased sexual activity and bouts of torpor prior to winter hibernation (Davis and Hitchcock 1965, pp. 304–306; Fenton 1969, p. 601; Parsons et al. 2003, pp. 63–64).

The swarming period may occur between July and early October, depending on latitude within the species' range (Hall and Brenner 1968, p. 780; Fenton 1969, p. 598; Caire et al. 1979, p. 405; Kurta et al. 1997, p. 479; Lowe 2012, p. 86;). Individuals may investigate several cave or mine openings during the transient portion of the swarming period, and some individuals may use these areas as temporary daytime roosts or may roost in forest habitat adjacent to these sites (Kurta et al. 1997, pp. 479, 483; Lowe 2012, p. 51). Many of the caves and mines associated with swarming are also used as hibernacula for several other species of bats (Fenton 1969, p. 599; Whitaker and Rissler 1992, p. 132; Kurta et al. 1997, p. 484; Glover and Altringham 2008, p. 1498; Randall and Broders 2014, p. 109).

While northern long-eared bats are thought to predominantly overwinter in caves and abandoned mines, the species has also been observed overwintering in other types of habitats that offer similar conditions (e.g., temperature, humidity levels, air flow). The species may use these alternate hibernacula in areas where caves or mines are not present (Griffin 1945, p. 22). Further, Girder et al. (2016, p. 11) found northern long-eared bats to be present and active year-round on the coastal plain of North Carolina, where there is no known non-cavernicolous (cave-

like) hibernacula; therefore, it is likely this population was not (traditionally) hibernating. Also, in coastal North Carolina, northern long-eared bats were observed to be active most of the winter. Although torpor was observed, time spent in torpor was very short, with the longest torpor bout (i.e., hibernation period) for each bat averaging 6.8 days (Jordan 2020, p. 672). Similarly, the species has been recently documented as active during the hibernation season in other southern states (e.g., Alabama, Mississippi, and Louisiana). In those areas, it appears the species enters temporary periods of torpor and will roost in trees and culverts while in torpor. However, Kentucky is within the hibernating portion of the species' range.

Spring staging is the time between winter hibernation and spring migration to summer habitat (Whitaker and Hamilton 1998, p. 80). During this time, bats begin to gradually emerge from hibernation, exit the hibernacula to feed, but re-enter the same or alternative hibernacula to resume daily bouts of torpor (Whitaker and Hamilton 1998, p. 80). The staging period is likely short in duration (Caire et al. 1979, p. 405; Whitaker and Hamilton 1998, p. 80). In Missouri, Caire et al. (1979, p. 405) found that northern long-eared bats moved into the staging period in mid-March through early May. Sasse et al. (2014, p. 172) found pregnant females using a mine in late April and May in Arkansas. In Michigan, Kurta et al. (1997, p. 478) determined that by early May, two-thirds of the *Myotis* species, including the northern long-eared bat, had dispersed to summer habitat. Variation in timing (onset and duration) of staging for Indiana bats (*Myotis sodalis*) was based on latitude and weather (USFWS 2007, pp. 39–40, 42); similarly, timing of staging for northern long-eared bats is likely based on these same factors.

Migratory movements between seasonal habitats (summer roosts and winter hibernacula) of 56 kilometers (35 miles) to 89 kilometers (55 miles) have been documented (Griffin 1940b, pp. 235, 236; Caire et al. 1979, p. 404; Nagorsen and Brigham 1993 p. 88). The spring migration period typically runs from mid-March to mid-May (Easterla 1968, p. 770; Caire et al. 1979, p. 404; Whitaker and Mumford 2009, p. 207); fall migration typically occurs between mid-August and mid-October.

In the summer, reproductive females form maternity colonies and give birth to pups. Maternity colonies are generally small, numbering from about 30 (Whitaker and Mumford 2009, p. 212) to 60 individuals (Caceres and Barclay 2000, p. 3); however, larger colonies of up to 100 adult females have been observed (Whitaker and Mumford 2009, p. 212). Most studies have found that the number of individuals roosting together in a given roost typically decreases from pregnancy to post-lactation (Foster and Kurta 1999, p. 667; Lacki and Schwierjohann 2001, p. 485; Garroway and Broders 2007, p. 962; Perry and Thill 2007, p. 224; Johnson et al. 2012, p. 227). Northern long-eared bats exhibit fission-fusion behavior (Garroway and Broders 2007, p. 961), where members frequently coalesce to form a group (fusion), but composition of the group is in flux, with individuals frequently departing to be solitary or to form smaller groups (fission) before returning to the main spatially discrete unit or network (Barclay and Kurta 2007, p. 44). As part of this behavior, northern long-eared bats switch tree roosts often (Sasse and Pekins 1996, p. 95), typically every two to three days (Foster and Kurta 1999, p. 665; Owen et al. 2002, p. 2; Carter and Feldhamer 2005, p. 261; Timpone et al. 2010, p. 119). Patriquin et al. (2016, p. 55) found that roost switching and use varies regionally in response to differences in ambient conditions (e.g., precipitation, temperature).

Adult females give birth to a single pup (Barbour and Davis 1969, p. 104). Birthing within the colony tends to be synchronous, with most births occurring around the same time (Krochmal and Sparks 2007, p. 654). Parturition (birth) may occur as early as late May or early June (Easterla

1968, p. 770; Caire et al. 1979, p. 406; Whitaker and Mumford 2009, p. 213) and may occur as late as mid-July (Whitaker and Mumford 2009, p. 213). Juvenile volancy often occurs by 21 days after birth (Kunz 1971, p. 480; Krochmal and Sparks 2007, p. 651) and has been documented as early as 18 days after birth (Krochmal and Sparks 2007, p. 651).

Northern long-eared bats are nocturnal foragers and use hawking (i.e., catching insects in flight) and gleaning (i.e., picking invertebrates from surfaces) behaviors in conjunction with passive acoustic cues (Nagorsen and Brigham 1993, p. 88; Ratcliffe and Dawson 2003, p. 851). The species has a diverse diet including moths, flies, leafhoppers, caddisflies, and beetles (Griffith and Gates 1985, p. 452; Nagorsen and Brigham 1993, p. 88; Brack and Whitaker 2001, p. 207), with diet composition differing geographically and seasonally (Brack and Whitaker 2001, p. 208). The most common invertebrates found in the diets of northern long-eared bats are lepidopterans (moths) and coleopterans (beetles) (Brack and Whitaker 2001, p. 207; Lee and McCracken 2004, pp. 595–596; Feldhamer et al. 2009, p. 45; Dodd et al. 2012, p. 1122), with arachnids also being a common prey item (Feldhamer et al. 2009, p. 45).

Foraging patterns indicate a peak activity period within five hours after sunset followed by a secondary peak within eight hours after sunset (Kunz 1973, pp. 18–19). Brack and Whitaker (2001, p. 207) did not find significant differences in the overall diet of northern long-eared bats between morning (3 a.m. to dawn) and evening (dusk to midnight) feedings; however, there were some differences in the consumption of particular prey orders between morning and evening feedings. Additionally, no significant differences existed in dietary diversity values between age classes or sex groups (Brack and Whitaker 2001, p. 208).

Habitat Characteristics of the Northern Long-eared Bat

Winter Habitat

Northern long-eared bats are thought to predominantly overwinter in hibernacula that include caves and abandoned mines. These hibernacula have relatively constant, cooler temperatures (0 to 9 degrees Celsius or 32 to 48 degrees Fahrenheit) (Raesly and Gates 1987, p. 18; Caceres and Pybus 1997, p. 2; Brack 2007, p. 744), with high humidity and no strong currents (Fitch and Shump 1979, p. 2; van Zyll de Jong 1985, p. 94; Raesly and Gates 1987, p. 118; Caceres and Pybus 1997, p. 2). Bats are typically found roosting singly or in small numbers in cave or mine walls or ceilings, often in small crevices or cracks, sometimes with only the nose and ears visible, and thus are easily overlooked during surveys (Griffin 1940a, pp. 181–182; Barbour and Davis 1969, p. 77; Caire et al. 1979, p. 405; van Zyll de Jong 1985, p. 9; Caceres and Pybus 1997, p. 2; Whitaker and Mumford 2009, pp. 209–210).

This species has also been observed overwintering in other types of habitats that have similar conditions (e.g., temperature, humidity levels, air flow) to cave or mine hibernacula. The species may use these alternate hibernacula in areas where caves or mines are not present (Griffin 1945, p. 22). Individuals have been found using the following alternative hibernacula: abandoned railroad tunnels (USFWS 2015, p. 17977), the entrance of a storm sewer in central Minnesota (Goehring 1954, p. 435), a hydroelectric dam facility in Michigan (Kurta et al. 1997, p. 478), an aqueduct in Massachusetts (Massachusetts Department of Fish and Game 2012, unpublished data), and a dry well in Massachusetts (Griffin 1945, p. 22). More recently, northern long-eared bats were found in a crawl space within a dwelling in Massachusetts (Dowling and O'Dell 2018, p. 376) and a rock crevice in Nebraska (White et al. 2020, p. 114).

Summer Habitat

Northern long-eared bats typically roost singly or in maternity colonies underneath bark or, more often, in cavities or crevices of both live trees and snags (Sasse and Pekins 1996, p. 95; Foster and Kurta 1999, p. 662; Owen et al. 2002, p. 2; Carter and Feldhamer 2005, p. 262; Perry and Thill 2007, p. 222; Timpone et al. 2010, p. 119). The species is flexible in tree species selection and, while they may select for certain tree species regionally, are likely not dependent on certain species of trees for roosts throughout their range; rather, many tree species that form suitable cavities or retain bark will be used by the bats opportunistically (Foster and Kurta 1999, p. 668; Silvis et al. 2016, p. 12; Hyzy 2020, p. 62). Carter and Feldhamer (2005, p. 265) hypothesized that structural complexity of habitat or available roosting resources are more important factors than the actual tree species. Further, Silvis et al. (2012, p. 7) found forest successional patterns, stand, and tree structure to be more crucial than tree species in creating and maintaining suitable long-term roosting opportunities.

Males and non-reproductive females may also roost in cooler locations in the summer, such as caves and mines (Barbour and Davis 1969, p. 77; Amelon and Burhans 2006, p. 72). To a lesser extent, individuals have also been observed roosting in colonies in human-made structures, such as in buildings, in barns, on utility poles, behind window shutters, in bridges, and in bat houses (Mumford and Cope 1964, p. 72; Barbour and Davis 1969, p. 77; Cope and Humphrey 1972, p. 9; Burke 1999, pp. 77–78; Sparks et al. 2004, p. 94; Amelon and Burhans 2006, p. 72; Whitaker and Mumford 2009, p. 209; Timpone et al. 2010, p. 119; Bohrman and Fecske 2013, pp. 37, 74; Feldhamer et al. 2003, p. 109; Sasse et al. 2014, p. 172; USFWS 2015, p. 17984; Dowling and O'Dell 2018, p. 376). It has been hypothesized that use of human-made structures may occur in areas with fewer suitable roost trees (Henderson and Broders 2008, p. 960; Dowling and O'Dell 2018, p. 376). In north-central West Virginia, individuals were found to use artificial roosts more readily as distance from large forests [greater than 200 hectares (494 acres)] increased, suggesting that artificial roosts are less likely to be selected when there is greater availability of suitable roost trees (De La Cruz et al. 2018, p. 496).

Studies of northern long-eared bat home range are highly variable, and it is uncertain if male northern long-eared bat summer home range size are larger or smaller than females. Foster and Kurta (1999) studied Indiana bats at a site also occupied by northern long-eared bats and found that northern long-eared bats moved greater distances (anywhere from 6 to 2,000 m) between roosts. Timpone et al. (2010) reported the mean distance between capture sites and roosts was 1,700 m. Fill et al. (2021) recorded a male traveled 2,700 m and a female traveled 300 m between capture site and roost. Another study found foraging areas can be larger for females than males (Broders et al. 2006) with males traveling approximately 500 m for foraging and roosting and females traveling 2000 m; these distances were centered around roosting and foraging areas.

From data reported by Broders et al. (2006), we estimate home ranges for males as approximately 14 ha (35 ac) and 54 ha (134 ac) for females. Henderson and Broders (2008) estimated the distance moved for foraging females was 1,100 m, whereas the total roost area covered was 31 ha (77 ac). Owen et al. (2003) determined a mean maternity home range size of 65 ha (161 ac), and Gorman et al. (2022) estimated a minimum roosting area for a maternity colony at 88.4 ha (218 ac). The Programmatic Biological Opinion on Final 4(d) Rule for the Northern Long-Eared Bat and Activities from Take Prohibitions (Service 2016) indicates 1,000 ac as the area a northern long-eared bat colony utilizes. The Service assumed a 131 ha (325 ac) home range size for northern long-eared bat based on the average from reported studies (Service 2024b); however, the Service did not provide sources for its information.

Most foraging occurs above the understory one to three meters (three to 10 feet) above the ground but under the canopy (Nagorsen and Brigham 1993, p. 88) on forested hillsides and ridges, rather than along riparian areas (LaVal et al. 1977, p. 594; Brack and Whitaker 2001, p. 207). This coincides with data indicating that mature forests are an important habitat type for foraging (Caceres and Pybus 1997, p. 2; White et al. 2017, p. 8). Foraging also takes place over small forest clearings, water, and along roads (van Zyll de Jong 1985, p. 94). Northern long-eared bats seem to prefer intact mixed-type forests with small gaps (i.e., forest trails, small roads, or forest-covered creeks) in forest with sparse or medium vegetation for forage and travel rather than fragmented habitat or areas that have been clear cut (USFWS 2015, p. 17992).

Numbers, Reproduction, and Distribution of the Northern Long-eared Bat

The northern long-eared bat's range includes much of the eastern and north-central U.S. and all Canadian provinces west to the southern Yukon Territory and eastern British Columbia (Nagorsen and Brigham 1993, p. 89; Caceres and Pybus 1997, p. 1; Environment Yukon 2011, p. 10). In the U.S., the species' range reaches from Maine west to Montana, south to eastern Kansas, eastern Oklahoma, Arkansas, and east to South Carolina (Whitaker and Hamilton 1998, p. 99; Caceres and Barclay 2000, p. 2; Simmons 2005, p. 516; Amelon and Burhans 2006, pp. 71–72). The range includes all or portions of the following 37 states and the District of Columbia: Alabama, Arkansas, Connecticut, Delaware, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Vermont, Virginia, West Virginia, Wisconsin, and Wyoming. Historically, the species has been most frequently observed in the northeastern U.S., Quebec, and Ontario, with sightings increasing during swarming and hibernation (Caceres and Barclay 2000). The species is patchily distributed throughout most of its range, and historically was less common in the southern and western portions of the range than in the northern portion of the range (Amelon and Burhans 2006).

More than 780 hibernacula have been identified throughout the species' range in the U.S., although many hibernacula contain only a few (one to three) individuals (Whitaker and Hamilton 1998). Northern long-eared bats are documented in hibernacula in 29 of the 37 states in the species' range, including: Alabama (2), Arkansas (71), Connecticut (8), Delaware (2), Georgia (3), Illinois (21), Indiana (25), Kentucky (119), Maine (3), Maryland (8), Massachusetts (7), Michigan (103), Minnesota (11), Missouri (more than 269), Nebraska (2), New Hampshire (11), New Jersey (7), New York (90), North Carolina (22), Oklahoma (16), Ohio (7), Pennsylvania (112), South Carolina (2), South Dakota (21), Tennessee (58), Vermont (16), Virginia (8), West Virginia (104), and Wisconsin (67). Other States within the species' range have no known hibernacula due to no suitable hibernacula present, lack of survey effort, or existence of unknown winter habitat.

The current range and distribution of northern long-eared bats must be described and understood within the context of the impacts of WNS. Before the onset of WNS, the best available information on the species came primarily from summer surveys (primarily focused on the Indiana bat or other bat species) and some targeted research projects. In those efforts, the northern long-eared bat was frequently encountered and was considered the most common

myotis bat in many areas. Overall, the species was considered to be widespread and plentiful throughout its historic range (Caceres and Barclay 2000).

Conservation Needs of and Threats to the Northern Long-eared Bat

White-nose Syndrome

For over a decade, WNS has been the primary stressor on the northern long-eared bat. WNS is a disease of bats that is caused by the fungal pathogen *Pseudogymnoascus destructans* (Pd) (Blehert et al. 2009, entire; Turner and Reeder 2009, entire; Lorch et al. 2011, entire; Coleman and Reichard 2014, entire; Frick et al. 2017, entire; Bernard et al. 2020, entire; Hoyt et al. 2021, entire). The effects of WNS have been severe, with most summer and winter colonies experiencing significant declines following its arrival. Just four years after the discovery of WNS, Turner et al. (2011, pp. 18–19) estimated a 98% decline in winter counts across 42 sites in Vermont, New York, and Pennsylvania. Similarly, Frick et al. (2015, p. 5) estimated the arrival of WNS led to a tenfold decrease in northern long-eared bat colony size. Most recently, Cheng et al. (2021, entire) used data from 27 states and two provinces to conclude that WNS caused estimated population declines of 97–100% across 79% of the species’ range. Although variation exists among sites, an overwhelming majority of hibernating colonies have developed WNS and experienced serious impacts within two to three years after its arrival (Cheng et al. 2021, entire; Wiens et al. 2022, pp. 231–247). To date, there are no proven measures to reduce the severity of impacts from WNS.

Wind Related Mortality

Wind related mortality, while overshadowed by the disproportionate impacts to tree bats and by the enormity of WNS, is also a significant stressor for northern long-eared bats. There is notable spatial overlap between northern long-eared bat occurrences and wind facilities, and mortality has been documented at wind turbines. At the 2020 installed mega-watt capacity, it was estimated that 122 northern long-eared bats were killed annually at wind facilities. Analyses using data from Wiens et al. (2022, pp. 236–247) and Whitby et al. (2022, entire) suggest that the impact of wind related mortality is discernible in the ongoing decline of the species. “Feathering” (i.e., pitch turbine blades parallel with the prevailing wind direction to slow rotation speeds) has been used to reduce bat fatalities; however, the effectiveness of this method in reducing fatality rates for the northern long-eared bat has not been documented.

Climate Change

While the risk of exposure to climatic changes exists range-wide for the northern long-eared bat, the magnitude, direction, and seasonality of those changes are expected to vary across its range. Although there may be some benefits to the species from a changing climate, overall, negative impacts are anticipated. Species-specific observations for the northern long-eared bat are lacking; however, observed impacts on the little brown bat include reduced reproduction during drought conditions (Adams 2010, pp. 2440–2442) and decreased adult survival during dry years in the Northeast (Frick et al. 2010, pp. 131–133). While sufficient moisture is important, excessive precipitation during the spring can negatively affect insectivorous bats. Heavier precipitation events may lead to decreased insect availability and reduced echolocation ability (Geipel et al. 2019, p. 4), resulting in lower foraging success. Precipitation also wets bat fur, reducing its insulating value (Webb and King 1984, p. 190; Burles et al. 2009, p. 132) and increasing a bat’s metabolic rate (Voigt et al. 2011, pp. 794–795). Bats are likely to reduce their foraging bouts during heavy rain events, and reduced reproduction has been observed during cooler, wetter springs in the Northwest (Grindal et al. 1992, pp. 342–343; Burles et al. 2009, p.

136). Responses to climate change by the northern long-eared bat will vary throughout its range based on the extent of annual temperature rise in the future.

Habitat Loss

As previously discussed, northern long-eared bats require suitable roosting, foraging, and commuting habitat during the spring, summer, and fall. Any loss of these habitats is likely to influence the survival and reproductive success of the species. Data from the National Land Cover Database shows that deciduous forest landcover decreased across all northern long-eared bat representation units (RPUs) (e.g., Southeast, Eastern Hardwoods, Subarctic, Midwest, and East Coast) by 1.4 million acres from 2006 to 2016, for an average a loss of 140,000 acres per year. Other cover types that provide foraging opportunities, such as emergent wetland cover types, also decreased by an additional 1.4 million acres across all RPUs. Changes in suitable habitat availability may be linked to losses of roosting or foraging habitat, longer flights between suitable roosting and foraging areas due to habitat fragmentation, fragmentation of maternity colony networks, and direct injury or mortality. Impacts from forest habitat removal may range from minor (e.g., removal of a small portion of foraging habitat in unfragmented forested area with a robust population) to significant (e.g., removal of roosting habitat in highly fragmented landscapes with a small, disconnected population). Adverse impacts are more likely in areas with little forest or highly fragmented forests (e.g., western U.S. and central Midwestern states), as there is a higher probability of removing roosts or losing connectivity between roosting and foraging habitats.

The complete loss of or modification of winter roosts (rendering the site unsuitable) can impact both individual bats and populations. Additionally, disturbances within hibernacula can make a site unsuitable or harm individuals using it. Modifications to bat hibernacula can alter the ability of bats to access the site (Spanjer and Fenton 2005, p. 1110) and affect the airflow and microclimate of the subterranean habitat, affecting the ability of the cave or mine to support hibernating bats. Furthermore, bats present during any excavation or filling could be crushed or suffocated. Human entry or other disturbances to hibernating bats results in additional arousals from hibernation, increasing total energy expenditure at a time when bats rely on fat reserves. This is of particular importance for sites impacted by WNS, as more frequent arousals from torpor increases the probability of mortality in bats with limited fat stores (Willis and Boyles 2012, p. 96).

5. Environmental Baseline

This section is an analysis of the effects of past and ongoing human and natural factors leading to the current status of each covered species, its habitat, and ecosystem within the Action Area. The environmental baseline is a “snapshot” of the species’ health in the Action Area at the time of the consultation and does not include the effects of the Action under review.

Indiana Bat

Action Area Numbers, Reproduction, and Distribution

The Action Area lies near the center of the species’ range, and numerous records of the species occupying summer and winter habitat exist throughout the Action Area. Occurrences of the species are clearly tied to the availability of suitable summer and winter habitat. Potential winter habitat is static (assuming no anthropogenic alterations occur) in the landscape because the caves

and other underground features the species relies on for winter habitats do not change locations. However, the species will move from one hibernaculum to another to take advantage of better conditions or to abandon habitat that humans or other factors have altered or disturbed.

The occupied range of the species (i.e., the collective home ranges of all individuals) within both the total range and the Action Area is unknown but is likely considerably smaller than the total range and Action Area, respectively, due to the presence of unsuitable habitats and unoccupied suitable habitats within both. According to data maintained by the KFO, the Indiana bat is known from numerous locations distributed across the Action Area.

Kentucky's 108 known hibernacula, including five P1 hibernacula, support significant winter populations, with an estimated 49,498 bats hibernating in 2022. Additionally, 43 hibernacula occur within 20 miles of Kentucky's border, including five P1 hibernacula. P1 hibernacula within Kentucky and the 20-mile buffer collectively housed 214,620 bats in 2022, representing 37% of the species' estimated range-wide population. Summer records for the species are closely aligned with areas containing sufficient forest cover to support summer roosting behavior. Despite the arrival of white-nose syndrome (WNS) in 2011, Kentucky has not documented a significant winter population decline, and it appears that most, if not all, of the Indiana bat summer colonies that existed pre-WNS remain occupied by the species.

Known Habitat Features for Indiana bats in Kentucky:

This summary outlines known habitat features utilized by the Indiana bat in Kentucky, based on current species occurrence data:

- **Designated Critical Habitat:** Two cave hibernacula were designated as critical habitat for the Indiana bat in Kentucky on September 24, 1976 (41 FR 41914), including Bat Cave in Carter County and Coach Cave in Edmonson County.
- **Summer Colonies:** 471 individual summer habitat occurrences have been documented. This includes 360 maternity records compiled from mist net captures, acoustic detections, and verified maternity roosts and 111 non-maternity records of solitary males and non-reproductive females.
- **Hibernacula:** 108 documented hibernacula have been identified among the caves, mines, or cave-like features where bats could hibernate during the winter months. This includes 21 P1 and P2 hibernacula and their associated Swarming 1 habitat and 87 P3 and P4 hibernacula and their associated Swarming 2 habitat.
- **Bridges/Culverts:** Four man-made structures have been identified that provide roosting sites for Indiana bats during the non-hibernation season.

Action Area Conservation Needs and Threats

It is difficult to identify specific factors affecting the Indiana bat's environment within the Action Area, because the Action Area has been defined as the Commonwealth of Kentucky and all portions of adjoining states that occur within 20 miles of the Kentucky border. This BO is based on analysis at a programmatic level rather than at an individual project scale. However, we can determine that there are a number of current and long-term land uses, environmental conditions, and demographic trends which could affect Indiana bats within the Action Area, which were discussed in the species status section of the BO.

Northern Long-eared Bat

Action Area Numbers, Reproduction, and Distribution

The Action Area lies near the center of the species' range, and numerous records of the species occupying summer and winter habitat exist throughout the Action Area. Occurrences of the species are clearly tied to the availability of suitable summer and winter habitat.

Potential winter habitat is static (assuming no anthropogenic alterations occur) in the landscape because the caves and other underground features the species relies on for winter habitats do not change locations. However, the species will move from one hibernaculum to another to take advantage of better conditions or to abandon habitat that humans or other factors have altered or disturbed.

The occupied range of the species (i.e., the collective home ranges of all individuals) within both the total range and the Action Area is unknown but is likely considerably smaller than the total range and Action Area, respectively, due to the presence of unsuitable habitats and unoccupied suitable habitats within both. According to our records, the NLEB is known from numerous locations distributed across the Action Area.

Estimating populations of northern long-eared bats in Kentucky from winter counts has been challenging due to the bats' tendency to hibernate in cracks and crevices. In the summer occupancy season, northern long-eared bats roost under bark, in tree cavities, or in crevices of both live and dead trees. They will also roost in man-made structures, such as bridges and abandoned buildings, and in natural rock shelters and crevices in cliff lines. Prior to the arrival of WNS in 2011, the northern long-eared bat was one of the most common bat species encountered during the summer in Kentucky. In 2013, the first noticeable WNS effects appeared and led to significant declines in Kentucky's northern long-eared bat populations. Spring staging and fall swarming are significant time periods for mating and migration to hibernacula. As such, forested habitat is crucial for roosting, foraging, and commuting behavior.

Known Habitat Features for Northern Long-eared Bats in Kentucky

This summary outlines known habitat features utilized by the northern long-eared bat in Kentucky, based on current species occurrence data:

- **Designated Critical Habitat:** Critical habitat has not been designated for the northern long-eared bat.
- **Pre-WNS Summer Colonies:** 1,541 individual occurrences were documented prior to 2013 during the summer occupancy timeframe from mist net captures, acoustic detections, and verified maternity roosts.
- **Post-WNS Summer Colonies:** 195 individual occurrences have been documented since 2013 during the summer occupancy timeframe, including mist net captures, acoustic detections, and verified maternity roosts.
- **Hibernacula:** 172 caves, mines, or cave-like features have been documented as hibernacula during the winter months.
- **Bridges/Culverts:** One man-made structure has been documented to provide roosting sites for bats during the non-hibernation season.

Action Area Conservation Needs and Threats

It is difficult to identify specific factors affecting the NLEB's environment within the Action Area, because the Action Area has been defined as the Commonwealth of Kentucky and all portions of adjoining states that occur within 20 miles of the Kentucky border. This BO is based on analysis at a programmatic level rather than at an individual project scale. However, we can determine that there are several current and long-term land uses, environmental conditions, and demographic trends which could affect NLEBs within the Action Area, which were discussed in the species status section of the BO.

6. Effects of the Action

This section addresses the direct and indirect effects of the Action on the covered species, including the effects of consequences of the proposed action. Direct effects are caused by the action and occur at the same time and place. Indirect effects are caused by the action but are later in time and reasonably certain to occur. The Service believes the covered species are likely to experience similar impacts from removal of forested habitat due to similarities in species biology and life history and has decided to combine our analysis into one analysis for both species. When appropriate, we identify where applicable science may apply to a particular species; however, we expect the effects on the species from the stressors to be similar and have reflected this in our analysis.

The types of impacts to covered species that are addressed in this BO and associated with the Strategy are limited to those adverse effects caused by the removal or alteration of suitable forested habitats. Because the Action is programmatic and involves a variety of project types, we did not identify specific project components of the Action to be analyzed. Rather, we have identified three general stressors (i.e., the alteration of the environment that is relevant to the species) to the covered species that are reasonably certain to result from the Action: noise and vibration, aquatic resource degradation, and tree removal. All these stressors are associated with the removal or alteration of suitable forested habitats that likely occur during the implementation or construction timeframe of a project that is incorporated as a likely consequence of the proposed Action.

Below we discuss the best available science relevant to each stressor, then describe the Stressor-Exposure-Response pathways that identify the circumstances for an individual bat's exposure to the stressor (i.e., the overlap in time and space between the stressor and the covered species). Finally, we identify and consider how proposed conservation measures may reduce the severity of the stressor or the probability of an individual bat's exposure for each pathway.

Effects of Noise and Vibration on the Indiana Bat and NLEB Applicable Science

Bats exposed to noise and vibration may flush from their roosts. Bats that flush during the daytime are at greater risk of harm due to predation (Mikula et al. 2016). Additionally, bats that flush from roosts and/or avoid travel and foraging areas in response to this stressor may be harmed due to an increase in energy expenditure. Increased energy demands could have a significant effect on bats due to their low body mass. Because females require increased energy reserves during lactation (Kurta et al. 1989), an increased demand for energy in response to noise and vibrations could be especially detrimental to lactating females and, subsequently, their pups.

Information is limited regarding the effects of noise and vibration on the covered species; however, studies show that the Indiana bat can tolerate some level of noise and vibration. For example, several construction projects, prior to documentation of WNS, have occurred on Fort Drum adjacent to multiple known Indiana bat roosts (Johnson et al. 2011). Construction around these project sites occurred for multiple years during the active season. The last known capture and roosting locations of Indiana bats near these projects were within approximately 800 and 400 meters (0.5 and 0.25 mi) of the construction activities, respectively. Indiana bats also occupy another military installation, Fort Knox, suggesting that noise from machinery and training activities may disturb colonies of roosting bats, but such disturbances would have to be severe to cause roost abandonment (Hawkins et al. 2008). Gardner et al. (1991) had evidence that Indiana bats continued to roost and forage in an area with active timber harvest. This suggested that noise and exhaust emissions from machinery could possibly disturb colonies of roosting bats, but such disturbances would have to be severe to cause roost abandonment. Callahan (1993) noted the likely cause of the bats in his study area abandoning a primary roost tree was disturbance from a bulldozer clearing brush adjacent to the tree. In another study near I-70 and the Indianapolis Airport, a primary maternity roost was located 1,970 ft. (0.6 km) south of I-70 (3D/International, Inc. 1996). This primary maternity roost was not abandoned, despite constant noise from the interstate and airport runways. However, the roost's proximity to I-70 may be related to a general lack of suitable roosting habitat in the vicinity, and since the noise levels from the airport were not novel to the bats (i.e., the bats had apparently habituated to the noise) (USFWS 2002). Noise and vibration could cause an Indiana bat to flush from its roost, expending extra energy and making it more vulnerable to predation (Mikula et al. 2016).

Novel noises would be expected to result in some changes to bat behaviors, but research suggests that bats can become habituated to this stressor under certain circumstances.

Effects Pathway #1 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Noise and Vibration

<i>Exposure (time)</i>	The covered species will be exposed to this stressor between April 1 and October 15, which corresponds with the occupancy timeframe of summer habitat or through November 15 if in known swarming habitat.
<i>Exposure (space)</i>	The covered species may be exposed to this stressor wherever they occur within the Action Area.
<i>Resource affected</i>	The stressor is expected to affect individual bats of the covered species, including adults and/or juveniles of both sexes.
<i>Individual response</i>	The stressor is expected to cause a variety of potential responses by affected individuals, especially if noise and vibration exceed normal, ambient levels for the Action Area where forested habitat removal is occurring. These include responses that are unlikely to result in significant effects to bats while roosting, such as arousing during daylight hours, shifting within the roost, and increasing vocalizations, as well as minor shifts in use of foraging and commuting habitats while active at night. The covered species are also likely to experience the following potentially significant effects if the stressor causes bats to flush from their roosts:

Effects Pathway #1 – Indiana Bat and Northern Long-eared Bat

	<ul style="list-style-type: none"> Extra energy expenditure that may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females and juveniles. Increased chance of predation of individuals, especially if individuals flush during daylight hours when noise and vibration are most likely to occur during forest habitat removal operations.
<i>Conservation Measures</i>	<p>The conservation measure that would most directly apply and minimize effects related to this stressor is:</p> <ul style="list-style-type: none"> Remove suitable trees between November 15 and March 31 (or October 16 and March 31 if outside known swarming habitat) when forested habitats are unoccupied by the covered species.
<i>Interpretation</i>	<p>The effects of increased noise and vibrations will be greatest during removal or alteration of the forested habitat when bats may be roosting in trees immediately adjacent to noise- and/or vibration-producing activities and are more likely to flush from their roosts or alter their behavior. Forested habitat removal activities that require heavy equipment and associated vehicles, personnel, and tools will be used at this time and are likely to produce noise and vibrations in the Action Area above ambient levels. The noise and vibrations associated with these activities may affect the covered species by causing individuals to alter their behaviors, which may be temporary or permanent. Significant changes in noise levels or significant increases in vibrations above ambient levels are more likely to result in altered behaviors, such as flushing from roosts and avoidance of habitat areas in close proximity to the source of noise or vibrations. The novelty of the noise or vibrations and the relative level of those disturbances will also likely dictate the range of responses from individuals or colonies of a covered species. Flushing from roosts is expected to cause: (a) extra energy expenditure that may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females and juveniles; (b) an increased chance of predation of individuals, especially if individuals flush during daylight hours when noise and vibration are most likely to occur during the construction of a project; and (c) increase in probability that adult females may abandon roosts and/or non-volant young if the event occurs during the pup season. These are adverse effects that are likely to result in harm to the covered species, including injury or mortality of individuals.</p> <p>Individuals roosting, foraging, and/or commuting in other portions of the project vicinity outside of the project's disturbance footprint will be exposed to this stressor at variable levels that would decrease with distance from the disturbance footprint. Specifically, noise and vibration levels within portions of the project vicinity outside the disturbance footprint are expected to be highest at locations closer to the point of origin and diminish with increasing distance from the point of origin, due to the diminished effects of noise and vibrations with distance from the source (i.e., the noise or vibrations will be absorbed</p>

Effects Pathway #1 – Indiana Bat and Northern Long-eared Bat

	<p>and typically become less loud or noticeable). Therefore, individuals of a covered species within the project vicinity and outside of the disturbance footprint are more likely to be affected if they are closer to the noise and vibrations point of origin and if the noise or vibrations are significantly different than ambient levels (i.e., loud, repetitive, novel, etc.). Conversely, individuals of a covered species would be less likely to be affected by noise and vibrations the farther they are from the disturbance footprint and the noise and vibrations' point of origin. However, the likelihood that adverse effects will not occur cannot be discounted, because flushing from a roost may still occur and result in the same adverse effects noted in the previous paragraph.</p> <p>Noise and vibration disturbances from personnel and vehicles within the project vicinity during certain aspects of the forested habitat removal that do not involve heavy equipment (e.g., land surveying) are expected to be similar to ambient levels within the project area and covered species present are likely habituated to such noise and vibration. Based on these factors, effects to the covered species from noise and vibrations associated with removal or alteration of suitable forested habitat that do not involve heavy equipment are unlikely to occur or result in effects outside of the disturbance footprint and are considered discountable.</p>
<i>Effect</i>	The effect of this stressor is Harm to affected covered species, which can include physical injury to individuals and/or mortality of individuals.

Because of the difficulty in determining the number of individuals that will be adversely affected as a result of this specific stressor, the Service has determined that it is appropriate to consider the total amount of covered species adversely affected by using the analysis for removal of summer and spring staging/fall swarming habitat in this section. We believe that this reduces the potential to double count the number of individuals impacted by the entire Action. A small, but indeterminable, portion of the covered species are expected to be injured or killed due to this stressor.

Effects of Aquatic Resource Degradation on the Indiana Bat and Northern Long-eared Bat

Aquatic resource degradation may occur indirectly as a result of tree removal associated with a project. In addition, the placement of culverts in streams and drainage ditches during construction of access roads and construction entrances could disturb sediment and negatively affect water quality. Trenching at stream crossings will also result in sediment disturbance in the streams, and soil that is exposed during excavation and vegetation removal could enter streams through stormwater runoff. Spills and leaks of petroleum-based products and other contaminants from vehicles and heavy equipment could also enter streams and degrade water quality. Activities that reduce the quantity or alter the quality of aquatic resources could affect a covered species, even if conducted while individuals are not present. However, project proponents are required to use BMPs in accordance with applicable state permits, program requirements, and

regulations to minimize the potential for aquatic resource degradation. Implementation of these practices is anticipated to avoid some potential water quality impacts and minimize others.

Applicable Science

Drinking water is essential, especially when bats are actively foraging. The covered species will utilize a variety of water sources including streams, ponds, and water-filled road ruts in upland forest. Numerous foraging habitat studies have found that Indiana bats often forage in closed to semi-open forested habitats and forest edges located in floodplains, riparian areas, lowlands, and uplands; old fields and agricultural fields are also used (USFWS 2007). Northern long-eared bats typically forage under the canopy on forested hillsides and ridges rather than along riparian areas (LaVal et al. 1977, p. 594; Brack and Whitaker 2001, p. 207). However, forest-covered streams may also be used by the northern long-eared bats during foraging and travel (USFWS 2015, p. 17992).

The covered species collectively feed on a variety of aquatic and terrestrial insects. Indiana bat diets vary seasonally and among different ages, sexes, and reproductive status (USFWS 1999). Four orders of insects contribute most to the diet of the species: Coleoptera, Diptera, Lepidoptera, and Trichoptera (Belwood 1979; Lee 1993; Kiser and Elliot 1996; Murray and Kurta 2002). Various reports differ considerably in which of these orders is most important. Consistent use of moths, flies, beetles, and caddisflies throughout the year at various colonies suggests that Indiana bats are selective predators to a certain degree, but incorporation of other insects into the diet also indicate these bats can be opportunistic (Murray and Kurta 2002). Brack and LaVal (1985) and Murray and Kurta (2002) suggested that the Indiana bat may best be described as a “selective opportunist.” The northern long-eared bat has a diverse diet that includes aquatic insects such as caddisflies (Griffith and Gates 1985, p. 452; Nagorsen and Brigham 1993, p. 88; Brack and Whitaker 2001, p. 207).

The negative impacts of sedimentation on aquatic insect larvae are well-documented. In a literature review, Henley et al. (2000) summarized how stream sedimentation impacts aquatic insect communities. Sediment suspended in the water column affects aquatic insect food sources by physically removing periphyton from substrate and reducing light available for primary production of phytoplankton. Sediment that settles out of the water column onto the substrate fills interstitial spaces occupied by certain aquatic insect larvae. Increases in sedimentation can change the composition of the insect community in a stream. In a three-year study measuring sedimentation and macroinvertebrate communities before, after, and during disturbance from a highway construction site, Hendrick (2008) found increased turbidity and total suspended solids downstream from the construction that correlated with a shift in macroinvertebrate communities. The change, however, was not great; the Hilsenhoff Biotic Index decreased from “excellent” before construction to “good” after construction. The use of BMPs likely minimized the effects of the construction on the macroinvertebrate communities.

Effects Pathway #2 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Aquatic Resource Degradation (sedimentation)

<i>Exposure (time)</i>	The covered species may be exposed to this stressor between April 1 and October 15, which corresponds with the occupancy timeframe of summer habitat.
------------------------	---

Effects Pathway #2 – Indiana Bat and Northern Long-eared Bat	
<i>Exposure (space)</i>	Foraging habitat in the Action Area consisting of streams and adjacent areas where aquatic insect prey may be located.
<i>Resource affected</i>	The stressor is expected to affect foraging habitat, prey abundance (aquatic insects), and individual bats, including adults and/or juveniles of both sexes.
<i>Individual response</i>	The stressor is expected to cause a variety of potential responses by affected covered species, especially if water quality is degraded in a project area where tree removal is occurring. These include responses that are unlikely to result in significant effects to bats, such as minor shifts in use of drinking water sources and foraging habitat while active at night. The Service has no data that would suggest that these minor behavioral shifts would require extra energy expenditures or reduce foraging efficiency that would reduce fitness and/or result in reduced survival and/or reproductive success of individuals.
<i>Conservation Measures</i>	The conservation measure that would most directly apply and minimize effects related to this stressor is: <ul style="list-style-type: none"> • Use BMPs for sediment and erosion control in accordance with state permits, program requirements, and regulations during tree removal and construction.
<i>Interpretation</i>	The Service has no information that would clearly indicate that potential aquatic resource degradation during tree removal is likely to result in significant adverse effects on the covered species. The effects of sedimentation on aquatic resources are expected to be minimal due to the temporary nature of activities and implementation of the conservation measures. Drinking water sources and aquatic insect prey are not expected to be eliminated, and the covered species have shown they can use a variety of drinking water and prey sources and do not forage exclusively on aquatic insect prey.
<i>Effect</i>	This stressor is expected to have an insignificant effect on the covered species. No physical injury to individuals and/or mortality of individuals is expected to result from this stressor.

Effects of Tree Removal on the Indiana Bat and Northern Long-eared Bat

The Action would result in the removal and loss of up to 2,000 cumulative acres of forested habitat per year and no more than 100 acres of habitat per project. The majority of this removal of forested habitats would occur during implementation of projects that use the Strategy for ESA compliance. Trees removed during the April 1– October 15 timeframe may be occupied by the covered species when they are removed. We do not know which trees would be removed during the active timeframe or exactly which trees the covered species would be occupying. The resulting forested habitat loss would be permanent and may create gaps in habitat between larger blocks of forested habitat or potential foraging corridors within the Action Area. In addition to the applicable science discussed below, for “Removal of Summer Habitat (active and inactive timeframes)” we also consider the science for “Loss and Fragmentation of Forested Habitat” and “Removal of Spring Staging/Fall Swarming Habitat (active and inactive timeframes)” for our analysis of this specific stressor.

Applicable Science – Removal of Summer Habitat (Summer Occupancy Timeframe)

Injury and death of northern long-eared bats during tree felling has not been reported; however, Belwood (2002) reported on the felling of a dead maple in a residential lawn in Ohio that resulted in the displacement of multiple Indiana bats. Thirty-four individuals were found on the ground after the tree was felled, including a dead adult female and 33 non-volant young. Five of the young died on impact or a short time later. Adult females that were not displaced remained in the tree until nighttime. The remaining young were assumed to have been retrieved later by the adult females. The deaths of 11 adult female Indiana bats were also reported from Indiana during the felling of a shagbark hickory (John O. Whitaker, personal communication, 1986). Approximately 50 Indiana bats were observed flying from the loose bark of a dead American elm tree that was bulldozed. Eight of the bats were captured, including two adult females, two immature males, and four immature females, indicating that the tree was being used by a maternity colony (Cope et al. 1973). Risk of injury or death from being crushed when a tree is felled is most likely to impact non-volant pups, but adults may also be injured or killed. This risk is greater for adults during cooler weather when bats periodically enter torpor and would be unable to arouse quickly enough to respond (i.e., flush and potentially avoid being in the roost when it is felled).

In addition to the expenditure of additional energy to find new roost trees, the removal of primary or alternate maternity roosts can lead to the fragmentation or break up of a maternity colony (Sparks et al. 2003; Silvis et al. 2014a). The effect of colony fragmentation on Indiana bats is unknown; however, Indiana bats congregate in large maternity colonies due to the benefits it provides the species. Barclay and Kurta (2007) stated that Indiana bats benefit from the formation of maternity colonies through: (1) information sharing about roosting and foraging habitats, (2) reduced predation risk, and (3) thermoregulatory advantages. However, this colonial behavior also comes with potential risks, such as increased parasite transmission and competition for resources.

Effects Pathway #3 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Tree Removal, Removal of Summer Habitat (summer occupancy)

<i>Exposure (time)</i>	The covered species may be exposed to this stressor between April 1 and October 15, which corresponds with the occupancy timeframe of summer habitat.
<i>Exposure (space)</i>	Forested habitat that is proposed for removal at various locations throughout the Action Area.
<i>Resource affected</i>	The stressor is expected to affect summer habitat (i.e., roost trees, foraging, and commuting habitat) and individual bats, including adults and/or juveniles of both sexes.
<i>Individual response</i>	The stressor is expected to cause a variety of potential responses by affected individuals of the covered species. These include responses that are unlikely to result in significant effects to bats, such as arousing during daylight hours, shifting within the roost, increasing vocalizations, and minor shifts in use of foraging and commuting habitats while active at night. However, the covered species are also likely to experience the following potentially significant effects from the stressor:

Effects Pathway #3 – Indiana Bat and Northern Long-eared Bat

	<ul style="list-style-type: none"> • Bats struck by equipment or crushed by a felled tree will be injured or killed. • Colony fragmentation could decrease thermoregulation and foraging efficiency that may reduce fitness and result in reduced survival/reproductive success. • Extra energy expenditure that may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females and juveniles. • Increased chance of predation of individuals, especially if individuals flush during daylight hours when roost tree removal and removal of summer habitat are most likely to occur.
<i>Conservation Measures</i>	The conservation measure associated with the Strategy that would most directly apply and minimize effects related to this stressor is: <ul style="list-style-type: none"> • Remove suitable trees between November 15 and March 31 (or October 16 and March 31 if outside known swarming habitat) when forested habitats are unoccupied by the covered species.
<i>Interpretation</i>	Tree removal during the summer occupancy timeframe could result in death or injury to roosting individuals that are crushed by a felled tree, especially non-volant juveniles. Those bats that survive or flush from felled trees will be exposed to increased levels of predation and expend extra energy to find another suitable roost. The use of additional energy in response to habitat loss, especially when combined with the energy needs associated with normal life cycle processes during the summer timeframe (e.g., migration, pregnancy, lactation) or other stressors (e.g., WNS), is likely to reduce fitness and subsequently reduce survival and reproductive success for individuals of the covered species. These effects would be avoided (e.g., no individual mortality would occur) or reduced to some extent if all proposed tree removal does not occur during the summer occupancy timeframe.
<i>Effect</i>	The effect of this stressor is Harm to affected covered species, which can include physical injury to individuals and/or mortality of individuals.

Applicable Science – Removal of Summer Habitat (Hibernation Timeframe)

The potential for adverse effects to Indiana bats from tree removal during the hibernation timeframe is rooted in the well-documented knowledge that Indiana bats exhibit strong fidelity to their summer roosting areas and foraging habitat (Kurta et al. 2002; Garner and Gardner 1992; USFWS 2007). Adverse effects to Indiana bats associated with the removal of forested habitats occur through several pathways that lead to a reduction in individual fitness as a result of increased energy expenditure. This evaluation is supported by numerous bat researchers, including Kurta and Rice (2002), who commented:

“The U.S. Fish and Wildlife Service often allows potential roost trees to be cut after Indiana bats leave for hibernation in order to make way for developments such as new bridges, highways, and housing projects. This policy understandably is intended to allow human developments to

proceed while preventing direct "take" of Indiana bats. This practice, however, should be limited, because it destroys potential roost trees without establishing whether they actually are used by Indiana bats, which may leave the bats with no shelter when they return in spring in an energetically stressed condition. Upon returning, the bats have just completed 6-7 months of hibernation and an extensive migration, and they arrive already pregnant and at a time when air temperatures are low and food (flying insects) is scarce. Excessive precipitation and/or colder-than-average temperatures drastically reduce reproductive success of temperate bats (Grindal et al. 1992; Lewis 1993), and such negative effects likely would occur even during normal weather if Indiana bats do not have adequate shelter."

Northern long-eared bat colonies retain their identity and exhibit high site fidelity between years (Silvis et al. 2015). A colony's use of the same general roosting area from one year to the next may occur due to the return of at least some individuals from the prior year – either juveniles (e.g., Silvis et al. 2015, p. 11) or adults. Northern long-eared bat females have been shown to roost together for multiple summers in the same location, and individual females have been captured returning to the same small area for at least five consecutive summers (Foster and Kurta 1999, p. 665, Patriquin et al. 2010, Perry 2011).

To evaluate the effects of roost removal on the northern long-eared bat, Silvis et al. (2015, p. 5) removed a primary roost and five secondary roosts, respectively, from the roosting area of two colonies in a heavily forested area of Kentucky. No roosts were removed from the roosting area of a third colony. In the year after roost removal, individuals persisted in each area and did not appear to change their colony roosting areas (Silvis et al. 2015, p. 10). Despite the 'consistent patterns of space use between years' by the colonies, few individuals were recaptured in the second year. "Colony identity" remained intact, although turnover among the individuals that comprised each colony was high (Olivera-Hyde et al. 2019, p. 724). The return of juveniles from the first year may have been key in retention of the colonies' identities despite the high colony turnover (Silvis et al. 2015, p. 10-11). Females "exhibit fidelity to a general geographic area", but they may not settle into the same areas as in previous years (Olivera-Hyde et al. 2019, p. 724). Although the identity of each colony persisted, Silvis et al. (2015b, p. 12) detected signs of a "segmented roost network" in the colony from which five secondary roosts were removed. Those five roosts constituted 24% of the roosts identified during radio-tracking of colony members. This was consistent with a previous simulation in which removal of about 20% of roosts resulted in a 50% chance of colony fragmentation (Silvis et al. 2014b, p. 287).

Forested habitat loss or alteration during the hibernation timeframe (i.e., while the bats are not present) harms all covered species by requiring them to increase energy use to respond to the habitat loss or alteration when they return to summer habitats. This is likely to impair essential behavior patterns associated with sheltering (roosting), breeding, and/or feeding (foraging). This impairment, in turn, results in reduced survival and/or reproduction of the affected individuals. These effects are compounded because most of the returning bats are coming from hibernacula infected with WNS. Individuals surviving WNS have additional energetic demands. For example, when emerging from hibernation, WNS-affected bats have lower levels of fat reserves than non-WNS-affected bats (Reeder et al. 2012; Warnecke et al. 2012). Many may also have wing damage (Reichard and Kunz 2009; Meteyer et al. 2009) that makes migration and foraging more challenging. Females that survive migrating to their summer habitat must partition energy

resources between foraging, keeping warm, maintaining a successful pregnancy, rearing pups, and healing their own bodies.

Effects Pathway #4 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Tree Removal, Removal of Summer Habitat (hibernation)

<i>Exposure (time)</i>	Tree removal will occur between November 16 and March 31 (or October 16 and March 31 outside of known swarming areas) during the hibernation timeframe. The covered species may be exposed to this stressor between April 1 and October 15 (summer occupancy timeframe) the first summer bats return after tree removal.
<i>Exposure (space)</i>	Forested habitat that is proposed for removal at various locations throughout the Action Area.
<i>Resource affected</i>	The stressor is expected to affect summer habitat (i.e., roost trees, foraging, and commuting habitat) and individual bats of a covered species, including adults and/or juveniles of both sexes.
<i>Individual response</i>	<p>The stressor is expected to cause a variety of potential responses by individuals from the loss of summer habitat in areas where removal or alteration of forested habitat previously occurred as part of a project. These include responses that are unlikely to result in significant effects to bats, such as minor shifts in use of foraging and commuting habitats while active at night. However, the covered species are also likely to experience the following potentially significant effects from the stressor:</p> <ul style="list-style-type: none">• Extra energy expenditure to find new suitable roosts may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females and juveniles.• Colony fragmentation could decrease thermoregulation and foraging efficiency that may reduce fitness and result in reduced survival/reproductive success.• Colony fragmentation will increase the chance of predation for individuals.
<i>Interpretation</i>	Adults from the covered species are expected to experience adverse effects after they arrive at summer habitat in the first year after tree removal occurs. The extra energy to find new roosting habitat is in addition to what is already necessary for foraging, pup rearing, social interactions, or other activities. The use of additional energy in response to habitat loss, especially when combined with the energy needs associated with normal life cycle processes during the summer timeframe (e.g., migration, pregnancy, lactation) or other stressors (e.g., WNS), is likely to result in adverse effects. However, the covered species are expected to adapt to this stressor in subsequent years after new suitable roosting habitat is found.
<i>Effect</i>	The effect of this stressor is Harm to affected individuals of a covered species, which can include physical injury and/or mortality of individuals.

Amount or Extent of Adverse Effects - Summer Habitats – Indiana Bats

The Service finds it unlikely that all maternity colonies within the Action Area are known and is reasonably certain that all suitable forested habitats without a known maternity colony have the potential to contain a maternity colony, unless survey data indicates otherwise. Summer presence/probable absence surveys for the Indiana bat within Kentucky have found an average occupancy rate of 2.0 percent for survey sites in potential maternity habitat (2005-2022) (unpublished data, Service 2024). Applying this occupancy rate to the 20,000 acres of potential maternity habitat to be removed within the 10-year timeframe of the Action, we estimate that 400 of these acres are occupied by Indiana bat maternity colonies. The Service assumes that maternity colonies can be as small as 25 acres and as large as 2,809 acres, with an average of 739 acres of habitat per colony based on our literature review, that colonies do not overlap, and that each maternity colony represents 360 Indiana bats (120 adult females, 120 adult males, and 120 pups) in Kentucky (Roby et. al *in press*; Unpublished data, Service 2024). Based on these assumptions, the Action’s effects on summer maternity habitat for the Indiana bat would affect up to 360 Indiana bats:

- 20,000 acres potential maternity habitat X 0.020 occupancy rate = 400 acres.
- 400 acres ÷ 739 acres per maternity colony = 0.54 potential maternity colonies
- 1 potential maternity colony (0.54 rounded up) X 360 Indiana bats per colony = 360 total Indiana bats.

Amount or Extent of Adverse Effects - Summer Habitats – Northern Long-eared Bats

The Service finds it unlikely that all maternity colonies within the Action Area are known, and we are reasonably certain that all suitable forested habitats without a known maternity colony have the potential to contain a maternity colony, unless survey data indicates otherwise. Summer presence/probable absence surveys for the northern long-eared bat within Kentucky have found an average occupancy rate of 3.75 percent (KY post-WNS rates based on mist-netting data from 2014-2022). Applying this occupancy rate to the 20,000 acres of potential maternity habitat to be removed within the 10-year timeframe of the Action, we estimate that 750 of these acres are occupied by northern long-eared bat maternity colonies. The Service assumes that northern long-eared bat males are mostly sympatric with females and that male and female summer home range sizes are not significantly different. We also assume summer home ranges can be as small as 35 acres and as large as 5,659 acres (2,700 m radius), with an average of 279 acres and that colonies do not overlap. The Service also assumes maternity colonies range widely in size (reported range of 7 to 100; Owen et al. 2002;), although about 30-60 may have been most common (Service 2016) based on pre-WNS data. In 2023, the Service provided an updated maternity colony size based on colony counts published in several studies in 2022 (Service 2023a). Data for Kentucky estimates the average maternity colony size is 13 adult females per colony, and that each produces 1 juvenile (13 adult females + 13 adult males + 13 pups = 39 bats) (USFWS 2014b). Based on these assumptions, the Action’s effects on summer maternity habitat for the northern long-eared bat would affect up to 117 northern long-eared bats:

- 20,000 acres of potential maternity habitat X 0.0375 occupancy rate = 750 acres.
- 750 acres ÷ 279 acres per maternity colony = 2.69 potential maternity colonies
- 3 potential maternity colonies (2.69 rounded up) X 39 bats per colony = 117 northern long-eared bats.

Applicable Science – Loss and Fragmentation of Forested Habitats

In addition to removal of roosting habitat, tree removal often results in the loss and fragmentation of forested habitats, resulting in degradation of the covered species foraging and commuting habitat. Patterson et al. (2003) noted that the mobility of bats allows them to exploit fragments of habitat. However, they cautioned that reliance on already diffused resources (e.g., roost trees) leaves bats highly vulnerable, and that energetics may preclude the use of overly patchy habitats.

In a fragmented landscape, Indiana bats may have to fly across less suitable or unsuitable habitat, which could pose a greater risk from predators (e.g., raptors) (Mikula et al. 2016). As a result, Indiana bats consistently follow tree-lined paths rather than crossing large open areas (Gardner et al. 1991; Murray and Kurta 2004). Murray and Kurta (2004) found that Indiana bats increased their commuting distances by 55% to follow these paths rather than flying over large agricultural fields. However, if these corridors are not available, Indiana bats may be forced over open areas. For example, Kniowski and Gehrt (2014) observed Indiana bats flying across open expanses of cropland >1 km (0.6 mile) to reach remote, isolated woodlots or riparian corridors.

Indiana and NLEB maternity colonies in Illinois, Indiana, Michigan, and Kentucky have been shown to use the same roosting and foraging areas during subsequent years (Gardner et al. 1991; Humphrey et al. 1977; Kurta and Murray 2002; Kurta et al. 1996; Kurta et al. 2002). Bats using familiar roosting and foraging areas are thought to benefit from decreased susceptibility to predators, increased foraging efficiency, and the ability to switch roosts in case of emergencies or alterations surrounding the original roost (Gumbert et al. 2002). Conversely, bats that must use new or inferior habitats after a loss or alteration of their normal forested habitat would not have these same benefits. In addition, movement distances, foraging areas, and roosting areas used by female northern long-eared bats may be smaller in fragmented forest landscapes than in landscapes with larger amounts of suitable forest cover (Henderson and Broders 2008, p. 959). In these areas, the extent of available forest patches may constrict northern long-eared bat foraging areas and could even increase use of alternative roosts (e.g., buildings, Henderson and Broders 2008 p. 959-960).

Racey and Entwistle (2003) discussed the difficulties of categorizing space requirements in bats, as they are highly mobile and show relatively patchy use of habitat (and use of linear landscape features), although connectivity of habitats has some clear advantages (e.g., aid orientation, attract insects, provide shelter from wind and/or predators). In their southern Illinois study, Carter et al. (2002) found Indiana bat roosts in a highly fragmented landscape, although both the number of patches and mean patch size were higher in the area surrounding roosts than around randomly selected points. Kniowski and Gehrt (2014) suggest, to obtain similar resources, longer or more frequent commuting flights will be required by Indiana bats in highly fragmented landscapes with smaller, more distant suitable habitat patches when compared to landscapes with larger, more abundant habitat patches.

Effects Pathway #5 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Tree Removal, Loss and Fragmentation of Forested Habitats

<i>Exposure (time)</i>	Tree removal will be a one-time occurrence but may occur any time of year. The covered species may be exposed to this stressor while
------------------------	--

Effects Pathway #5 – Indiana Bat and Northern Long-eared Bat

	present between April 1 and October 15 or during the first summer bats return if tree removal occurs in the unoccupied time.
<i>Exposure (space)</i>	Forested habitat that is proposed for removal at various locations throughout the Action Area.
<i>Resource affected</i>	The stressor is expected to affect summer habitat (i.e., foraging and commuting habitat) and individual bats, including adults and juveniles of both sexes.
<i>Individual response</i>	<p>The stressor is expected to cause a variety of potential responses by affected individuals from the loss and fragmentation of summer habitat in the Action Area where loss and fragmentation of forested habitats has occurred as the result of a project. These include responses that are unlikely to result in significant effects to bats, such as minor shifts in use of foraging and commuting habitats while active at night. However, the covered species are also likely to experience the following potentially significant effects depending on the timing, location, and size of forest loss and fragmentation:</p> <ul style="list-style-type: none"> • Extra energy expenditure that may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females and juveniles.
<i>Conservation Measure</i>	<p>The conservation measures that would most directly apply and minimize effects related to this stressor are:</p> <ul style="list-style-type: none"> • Remove suitable trees between November 15 and March 31 (or October 16 and March 31 if outside known swarming habitat) when forested habitats are unoccupied by the covered species.
<i>Interpretation</i>	<p>Tree removal will create gaps within larger forested blocks and along linear forested corridors or expand gaps along the edges of larger forest blocks. Projects that remove small amounts of forested habitat within an already diverse landscape of forest blocks, tree lines, and open areas are anticipated to have an insignificant effect on the covered species. These types of gaps are not expected to make access to other forested habitat more difficult, require additional energy expenditure, or limit access to habitat because the covered species currently foraging and commuting in these types of areas are presumably unaffected by these gaps. Additionally, the gap may provide new or additional forest edge habitat that the covered species could use for foraging and commuting habitat.</p> <p>However, projects that remove large areas of forested habitat and/or fragment other areas of habitat to the extent it significantly alters a bats' behavior (i.e., foraging and commuting) are likely to adversely affect the covered species. These adverse effects may cause extra energy expenditure, reducing the fitness of individuals, especially for females and juveniles.</p>
<i>Effect</i>	The effect of this stressor is Harm to affected covered species, which can include physical injury to individuals and/or mortality of individuals.

Because of the difficulty in determining the number of individuals from a covered species that will be adversely affected as a result of this specific stressor, the Service has determined that it is appropriate to consider the total amount of individuals from each covered species adversely affected by using the analysis for removal of summer and spring staging/fall swarming habitat. We believe that this reduces the potential to double count the number of individual bats impacted by the entire Action. A small, but indeterminable, number of covered species are expected to be injured or killed due to this stressor.

Applicable Science- Removal of Spring Staging/Fall Swarming Habitat

The active spring staging and fall swarming periods are April 1 – May 14 and August 16 – November 15 respectively and are sensitive periods for the covered species. This is when mating occurs, bats are preparing for migration to summer habitats, and when bats are busy foraging to store sufficient fat reserves to survive winter hibernation. Suitable spring staging and fall swarming habitat is comprised of forested habitats where they roost, forage, and travel, which is most typically within 5 to 10 miles of a hibernaculum, depending on the species, available forested habitat, and number of individuals using a particular hibernaculum. This includes forested patches as well as linear features such as fencerows, riparian forests, and other wooded corridors. These wooded areas may be dense or loose aggregates of trees with variable amounts of canopy closure.

In general, covered species use roosting, foraging, and commuting habitat(s) in the spring and fall that are similar to those selected during the summer. Therefore, we are considering the applicable science discussed above for “Removal of Summer Habitat (active and inactive timeframes)”, and “Loss and Fragmentation of Forested Habitat” for our analysis of this specific sub-stressor.

Effects Pathway #6 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Tree Removal, Removal of Spring Staging /Fall Swarming Habitat
(staging/swarming occupancy)

<i>Exposure (time)</i>	The covered species may be exposed to this stressor between April 1 and May 14 and August 16 and November 15, which corresponds with the occupancy timeframe of spring staging and fall swarming habitat, respectively.
<i>Exposure (space)</i>	Forested habitat associated with hibernacula that is proposed for removal at various locations throughout the Action Area.
<i>Resource affected</i>	The stressor is expected to affect staging/swarming habitat (i.e., roosting, foraging, and commuting habitat) and individual bats, including adults and juveniles of both sexes.
<i>Individual response</i>	The stressor is expected to cause a variety of potential responses by affected individuals of a covered species from the loss of staging/swarming habitat where tree removal is occurring. These include responses that are unlikely to result in significant effects to bats, such as minor shifts in use of foraging and commuting habitats while active at night. However, the covered species are also likely to experience the following potentially significant effects from the stressor:

	<ul style="list-style-type: none"> • Bats struck by equipment or crushed by a felled tree will be injured or killed. • Extra energy expenditure may reduce the fitness of individuals and result in reduced survival and/or reproductive success, especially for females. • Increased chance of predation of individuals, especially if individuals flush during daylight hours when roost tree removal and removal of staging/swarming habitat are most likely to occur during the implementation component of a project.
<i>Conservation Measures</i>	<p>The conservation measures that would most directly apply and minimize effects related to this stressor are:</p> <ul style="list-style-type: none"> • Remove suitable trees between November 15 and March 31 (or October 16 and March 31 if outside known swarming habitat) when forested habitats are unoccupied by the covered species. • Avoid removing suitable forested habitat within ½-mile of a known Indiana bat P3 or P4 hibernaculum and one-mile of a known Indiana bat P1 or P2 hibernaculum. Avoid removing suitable forested habitat within ½-mile of a known northern long-eared bat hibernaculum.
<i>Interpretation</i>	<p>Tree removal during the staging/swarming occupancy timeframes could result in death or injury to roosting individuals that are crushed by a felled tree. Those bats that survive or flush from felled trees will be exposed to increased levels of predation and expend extra energy to find another suitable roost. This energy expenditure is in addition to what is likely necessary for foraging, social interactions, and other activities. The use of additional energy in response to habitat loss, especially when combined with the energy needs associated with normal life cycle processes during the staging/swarming timeframes (e.g., migration and mating) or other stressors (e.g., WNS), is likely to reduce fitness and subsequently reduce survival and reproductive success. These effects would be avoided (e.g., no individual mortality would occur) or reduced to some extent if all proposed tree removal within the staging/swarming habitat does not occur during the staging/swarming occupancy timeframe.</p>
<i>Effect</i>	<p>The effect of this stressor is Harm to affected covered species, which can include physical injury to individuals and/or mortality of individuals.</p>

Effects Pathway #7 – Indiana Bat and Northern Long-eared Bat

Activity: Removal or Alteration of Suitable Forested Habitat

Stressor: Tree Removal, Removal of Spring Staging/ Fall Swarming Habitat (hibernation)

<i>Exposure (time)</i>	Inactive timeframe (November 16 – August 15) removal will expose the covered species to adverse effects from April 1 – May 14 and August 16 – November 15, for one season after removal.
<i>Exposure (space)</i>	Forested habitat associated with hibernacula that is proposed for removal at various locations throughout the Action Area.

<i>Resource affected</i>	The stressor is expected to affect staging/swarming habitat (i.e., roosting, foraging, and commuting habitat) and individual bats, including adults and juveniles of both sexes.
<i>Individual response</i>	<p>The stressor is expected to cause a variety of potential responses by affected individuals from the loss of staging/swarming habitat in areas where tree removal previously occurred. These include responses that are unlikely to result in significant effects to bats, such as minor shifts in use of foraging and commuting habitats while active at night. However, the covered species are also likely to experience the following potentially significant effects from the stressor:</p> <ul style="list-style-type: none"> Increased effort to find new suitable roosting habitat requires extra energy expenditure that can reduce fitness and result in reduced survival / reproductive success. Increased effort to access sufficient foraging resources requires extra energy expenditure that can reduce fitness and result in reduced survival / reproductive success. Reduced foraging efficiency can reduce fitness and result in reduced survival / reproductive success.
<i>Conservation Measures</i>	<p>The conservation measures that would most directly apply and minimize effects related to this stressor are:</p> <ul style="list-style-type: none"> Avoid removing suitable forested habitat within ½-mile of a known Indiana bat P3 or P4 hibernaculum and one-mile of a known Indiana bat P1 or P2 hibernaculum. Avoid removing suitable forested habitat within ½-mile of a known northern long-eared bat hibernaculum.
<i>Interpretation</i>	<p>The covered species will experience adverse effects after they arrive at their spring staging/fall swarming habitat the first year after tree removal. The extra energy to find new habitat is in addition to what is necessary for foraging, social interactions, or other activities. The use of additional energy in response to habitat loss, especially when combined with the energy needs associated with normal life cycle processes (e.g., migration and mating) or other stressors (e.g., WNS), is likely to result in adverse effects. The covered species are expected to adapt to this stressor in subsequent years after they have found new suitable habitat.</p>
<i>Effect</i>	<p>The effect of this stressor is Harm to affected individuals of a covered species, which can include physical injury and/or mortality of individuals.</p>

Amount or Extent of Adverse Effects – Staging/Swarming Habitats – Indiana Bats

According to winter counts conducted at known Indiana bat hibernacula (P1, P2, P3, and P4), approximately 214,620 Indiana bats hibernate within the Action Area (USFWS unpublished 2022 data). To calculate the extent of adverse effects on Indiana bats due to removal of staging/swarming habitat, we use a worst-case scenario that assumes all 20,000 acres of habitat impacts throughout the 10-year timeframe of the Action occur within known staging/swarming habitat. We use the most recent winter count data to estimate the density of Indiana bats using the Action Area staging/swarming habitats that are within 10 miles of P1 and P2 hibernacula and

within 5 miles of P3 and P4 hibernacula. We assume an even distribution of bats and 50 percent forest cover within these circles:

- 214,620 bats in P1, P2, P3, and P4 hibernacula / 5.90 million acres of associated staging/swarming habitat = 0.036 bats per acre

To estimate how many Indiana bats the Action will affect in staging/swarming habitats, the bat density is applied to the acreage of staging/swarming habitat that we are assuming the Action may affect (20,000 acres of impact and 10 years of the Action). Based on these assumptions, the Action's effects on known staging/swarming habitat would affect up to 720 Indiana bats:

- 20,000 acres of staging/swarming habitat impact x 0.036 bats per acre = 720 Indiana bats

Very few, if any, of these 720 Indiana bats are expected to be injured or killed by the Action. Disruption of normal behavior as a result of physical disturbance and/or habitat modification or degradation will account for the majority of potential impacts.

Amount or Extent of Adverse Effects – Staging/Swarming Habitats – Northern Long-eared Bats

As discussed in the Status of the Species and Environmental Baseline, northern long-eared bats are difficult to detect during winter surveys, and the Service expects that many hibernacula are unknown. To be conservative and consider a larger effect on the species than is likely to occur, the Service has chosen to assume that all the projects implemented under the Strategy will impact a single northern long-eared bat staging/swarming population. The Service is also assuming that each 5-mile staging/swarming buffer contains 100 northern long-eared bats. This is likely a high estimate considering that most winter counts within the Action Area detect fewer than 10 northern long-eared bats per hibernaculum. However, we believe using this high estimate accounts for additional impacts that may occur to currently undocumented staging/swarming populations of the species. Using these assumptions to calculate a density for staging/swarming habitat and applying that density to the 20,000 acres impacted over the 10-year timeframe of the Action, the Service anticipates that up to 40 northern long-eared bats will be adversely affected by the Action's impacts to staging/swarming habitat:

- 5-mile radius around a hibernaculum encompasses 50,266 acres
- 100 bats ÷ 50,266 acres = 0.002 bats per acre
- 20,000 acres of staging/swarming habitat impact x 0.002 bats per acre = 40 bats

Very few, if any, of these 40 northern long-eared bats are expected to be injured or killed by the Action. Disruption of normal behavior as a result of physical disturbance and/or habitat modification or degradation will account for the majority of potential impacts to the species.

Summary of Effects of the Action on the Indiana Bat and Northern Long-eared Bat

The Action may occur within suitable staging/swarming habitat and/or suitable summer maternity roosting, foraging, and commuting habitat for the covered species. Impacts to the covered species will occur due to the removal or alteration of suitable forested habitat. Stressors to the covered species as a consequence of tree removal activities include noise and vibration, aquatic resource degradation, and tree removal (Table 1). Impacts to habitat could occur at any

time of year, including May 15 to July 31 when non-volant pups may be present, although this has been restricted under the Action to only 10 ac/project where it cannot be avoided.

It is difficult to determine the number of individuals from the covered species that will be adversely affected by these identified stressors during specific activities and timeframes. The Service has determined that the total number of covered species adversely affected should be based on the scenario that all 20,000 acres of habitat impacts would occur in summer maternity habitat or all 20,000 acres of habitat impacts will occur in staging/swarming habitat, but in reality, it will be some unknown quantity of both. Although unlikely to occur, this approach assumes the worst-case scenario while evaluating the extent of adverse effects on the species. We estimate that 360 Indiana bats and 117 NLEBs (females/males/juveniles, proportionally) utilizing summer maternity habitats, and 720 Indiana bats and 40 NLEBs (females/males, proportionally) utilizing staging/swarming habitats could be adversely affected by impacts associated with the Action.

Table 1. A summary of the effects of the Action on the Covered Species

Stressors: Activity	Adverse	Insignificant/Discountable
Noise and vibration: <i>removal or alteration of suitable forested habitat</i>	harm	
Aquatic resource degradation, sedimentation: <i>removal or alteration of suitable forested habitat</i>		insignificant
Tree removal, summer habitat (summer occupancy): <i>removal or alteration of suitable forested habitat</i>	harm	
Tree removal, summer habitat (hibernation): <i>removal or alteration of suitable forested habitat</i>	harm	
Tree removal, forest loss and fragmentation: <i>removal or alteration of suitable forested habitat</i>	harm	
Tree removal, staging/swarming habitat (staging/swarming occupancy): <i>removal or alteration of suitable forested habitat</i>	harm	
Tree removal, staging/swarming habitat (hibernation): <i>removal or alteration of suitable forested habitat</i>	harm	

7. Cumulative Effects on the Covered Species

For purposes of consultation under ESA §7, cumulative effects are the effects of future state, tribal, local, or private actions that are reasonably certain to occur in the Action Area. Future federal actions that are unrelated to the proposed action are not considered, because they require separate consultation under §7 of the ESA.

Land use activities that may affect the covered species and that are likely to occur within the Action Area include: timber harvest, all-terrain vehicle recreational use, recreational use of caves, and development associated with road, residential, industrial, and agricultural development, and other related activities that are not a direct consequence of the Action. These private actions are likely to occur within the Action Area, but the Service is unaware of any quantifiable information about the extent of private timber harvests within the Action Area, the amount of use of off-highway vehicles within the Action Area, or the amount of recreational use

of caves within the Action Area. Similarly, the Service does not have any information on the amount or types of residential, industrial, or agricultural development that will occur within the Action Area that may require removal of forested habitat without the mandate to consult with the Service. Therefore, the Service is unable to make any determinations or conduct any meaningful analysis of how these actions may or may not adversely and/or beneficially affect the covered species. It is possible that these activities may will have cumulative effects on the covered species and their habitat in certain situations (e.g., a private timber harvest during summer months within an unknown maternity colony may cause adverse effects to that maternity colony). We can only speculate as to the extent or severity of those effects, but these actions have been occurring for decades and are considered part of the environmental baseline for the current status of the species.

8. Conclusion of Effects on the Covered Species

In this section, we summarize and interpret the findings of the previous sections for the covered species (status, baseline, effects, and cumulative effects) relative to the purpose of a BO under §7(a)(2) of the ESA, which is to determine whether a Federal action is likely to jeopardize the continued existence of the species. We have considered the status of each species across its range, the status of each species within the Action Area, and the effects of the Action to each of the covered bat species. In our effects analysis, we identified how the covered species would be adversely affected by the Action. We anticipate that individuals from each of the covered species who utilize summer and staging/swarming habitat within the Action Area are likely to be adversely affected by the removal of forested habitat and will experience harm as a result of the Action. However, we do not anticipate that all individuals from the covered species that are adversely affected will experience harm based on the following reasons:

- Injury and/or mortality of adult and juvenile bats will be reduced if some or all of the habitat removal occurs during the hibernation timeframe (November 16 – March 31) when all of the covered bat species are not present in the Action Area.
- Injury and/or mortality of non-volant pups will be reduced if some or all of the habitat removal occurs outside the pup season (May 15 – July 31).
- The minimal amount of summer habitat removed each year from the Action Area versus the total availability of habitat within the Action Area.
- The removal of habitat throughout the Action Area will be random and spread across the variety of habitat types (e.g., known summer, known staging/swarming, suitable summer), and additional habitat will periodically become available independent of the Action.
- Limiting the size of each area where habitat can be removed by a single project will limit the scale of impacts and reduce the potential for loss of large, forested blocks or significant fragmentation of remaining forested habitat.

After reviewing the status of each covered species, the environmental baseline for the Action Area, the effects of the Action, and the inclusion of any cumulative effects in the environmental baseline, it is the Service’s biological opinion that the Action is not likely to jeopardize the continued existence of the Indiana bat or the northern long-eared bat. We reached this determination using the best available commercial and scientific information as described in the effects analysis of this BO while considering how those effects relate to the survival and

recovery potential (i.e., resiliency, redundancy, and representation) of the Indiana bat and northern long-eared bat, as described below:

Resiliency

Resiliency describes the ability of a species to withstand stochastic disturbance (arising from random factors). Resiliency is positively related to population size, growth rate, and fecundity and may be influenced by connectivity among populations. Generally, populations need enough individuals within habitats of adequate area and quality to maintain survival and reproduction in spite of disturbance. Resilient populations are better able to withstand disturbances such as random fluctuations in birth rates (demographic stochasticity), variations in rainfall (environmental stochasticity), or the effects of anthropogenic activities.

Indiana Bat

The recovery goals for the species include obtaining a minimum overall population estimate of 457,000 and demonstrating a positive population growth rate. The number of bats adversely affected by the Action would be 1,080, and we do not expect mortality of all of these Indiana bats. Additionally, 1,080 individuals represent only 0.18% of the 2022 range-wide estimate of Indiana bats (583,263). Therefore, the Action would adversely affect only a small proportion of the range-wide species' population. For these reasons, the Action will not reduce the resiliency of the Indiana bat.

Northern Long-eared Bat

The range of the northern long-eared bat includes 37 states and the District of Columbia in the eastern and north-central United States and portions of eight Canadian provinces. We anticipate the number of NLEB adversely affected by the Action would be 157, and we do not expect mortality of all these individuals. The Action would only adversely affect a small proportion of the range-wide species' population. In addition, the habitat losses associated with the Action are not expected to cause significant or meaningful reductions in habitat connectivity or habitat quality that would lead to negative population-level effects. For these reasons, the Action will not reduce the resiliency of the northern long-eared bat.

Redundancy

Redundancy describes the ability of a species to withstand catastrophic events (a rare destructive natural event or episode involving many populations). It "guards against irreplaceable loss of representation" (Redford et al. 2011, p. 42) and minimizes the effect of localized extirpation on the range-wide persistence of a species. Generally speaking, redundancy is best achieved by having multiple, resilient (connected) populations widely distributed across the species' range. Having multiple populations reduces the likelihood that all populations are affected simultaneously, while having widely distributed populations reduces the likelihood of populations possessing similar vulnerabilities to a catastrophic event. Given sufficient redundancy, single or multiple catastrophic events are unlikely to cause the extinction of a species. Therefore, as redundancy increases, species viability also increases.

Indiana Bat and Northern Long-eared Bat

If habitat removal occurs from May 15 to July 31, non-volant pups from the covered species may be impacted, and mortality would be higher than at other times of the year. However, implementation of the Strategy limits clearing of habitat during this time of year to only 10 acres per project, thus minimizing the number of maternity colonies affected by the proposed Action. Additionally, impacts are expected to be widely distributed across the Action Area and are unlikely to cause a reduction in any of the covered species' redundancy, as the likelihood of a catastrophic event (or large effect) would be reduced. This represents a very small proportion of the range-wide population for each of the covered species, and thus, the widely distributed impacts are unlikely to cause a reduction in redundancy of the Indiana bat and northern long-eared bat as the likelihood of a catastrophic event (or large effect) would be reduced.

Representation

Representation describes the ability of a species to adapt to changing environmental conditions over time and is characterized by the breadth of genetic and environmental diversity within and among populations. The more representation, or diversity, a species has, the more it is capable of adapting to changes (natural or human caused) in its environment. In the absence of species-specific genetic and ecological diversity information, we evaluate representation based on the extent and variability of habitat characteristics across the geographical range and other factors as appropriate.

Indiana Bat and Northern Long-eared Bat

The overall acreage of forested habitat within the Action Area that is suitable habitat for the covered species represents only a fraction of the total suitable habitat within the range of the covered species. The 20,000 acres of suitable habitat that will be removed for the Action represents only 0.001% of the total habitat available in the Action Area.

Additionally, no reduction in the distribution of Indiana bats or northern long-eared bats is expected because the Action Area will continue to support suitable habitat, and both species are expected to continue to occupy the Action Area. For these reasons, we do not expect the representation of the Indiana bat or northern long-eared bat to be reduced by the Action.

Further, implementation of the Strategy is expected to promote the survival and recovery of the covered species through protection and management of:

- 1) existing forested habitat that supports known maternity populations, particularly those that would expand existing conservation ownerships;
- 2) known priority hibernacula;
- 3) additional conservation lands that contain potential habitat for the species, particularly those that would expand existing conservation ownerships; and
- 4) additional conservation benefits from the avoidance and minimization that the Strategy inherently encourages (e.g., protection of known habitats, protection of non-volant pups, protection of staging/swarming bats).

Based on this analysis of resiliency, redundancy, and representation, we conclude that the effects of the Action will not appreciably reduce the likelihood of both the survival and recovery of the Indiana bat or northern long-eared bat.

9. Incidental Take Statement

ESA §9(a)(1) and regulations issued under §4(d) prohibit the take of endangered and threatened fish and wildlife species without special exemption. The term “take” in the ESA means “to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” (ESA §3(19)). In regulations, the Service further defines:

- “harass” is defined as an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying to such an extent as to significantly disrupt normal behavior patterns which include, but are not limited to, breeding, feeding or sheltering.” (50 C.F.R. § 17.3); .
- “harm” as “an act which actually kills or injures wildlife. Such act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding or sheltering;” (50 CFR §17.3); and
- “incidental take” as “takings that result from, but are not the purpose of, carrying out an otherwise lawful activity conducted by the Federal agency or applicant” (50 CFR §402.02).

Under the terms of ESA §7(b)(4) and §7(o)(2), taking that is incidental to a Federal agency action that would not violate ESA §7(a)(2) is not considered prohibited, provided that such taking is in compliance with the terms and conditions of an incidental take statement (ITS).

For the exemption in ESA §7(o)(2) to apply to the Action considered in this BO, the Service must undertake the non-discretionary measures described in this ITS and the conservation measures detailed in this BO, and these measures must become binding conditions of any permit, contract, or grant issued for implementing the Action. The Service has a continuing duty to monitor the activity covered by this ITS. The protective coverage of §7(o)(2) may lapse if the Service fails to (a) assume and implement the terms and conditions; or (b) require a permittee, contractor, or grantee to adhere to the terms and conditions of the ITS through enforceable terms that are added to the permit, contract, or grant document. To monitor the impact of incidental take, the project proponent must report the progress of the specific action and its impact on the covered species to the Service as specified in the respective sections of this ITS.

Amount or Extent of Take

This section specifies the amount or extent of take of the Indiana bat and northern long-eared bat, that the Action is reasonably certain to cause.

Indiana Bat

Based on our evaluation, the proposed Action is reasonably certain to cause the incidental take of 1,080 individual Indiana bats. This taking is expected primarily in the form of harm, although

some harassment may also occur. The mechanisms of this taking and the basis for our estimation of its extent are described in Section 6 (Effects of the Action) of this BO.

Table 4. Amount of incidental take of the Indiana bat caused by the Action.

Species	# of Individuals	Form of Take
Indiana bat	360	Harm (Tree Removal, Summer Habitat)
	720	Harm (Tree Removal, Staging/Swarming Habitat)

Surrogate Measures for Monitoring

The Service anticipates the incidental taking of Indiana bats associated with the Action will be difficult to detect for the following reasons:

- The individuals are small, mostly nocturnal, and when not hibernating, occupy forested habitats where they are difficult to find, capture, or observe;
- The Indiana bat forms maternity colonies under loose bark or in the cavities of trees, and males and non-reproductive females may roost individually, which makes finding roost trees difficult;
- Finding dead or injured specimens during or following project implementation is unlikely; and
- Most incidental take is in the form of non-lethal harm and not directly observable.

When it is not practical to monitor take in terms of individuals of the listed species, the regulations at 50 CFR §402.14(i)(1)(i) indicate that an ITS may express the amount or extent of take using a surrogate (e.g., a similarly affected species, habitat, or ecological conditions), provided that the Service also:

- describes the causal link between the surrogate and take of the listed species; and
- sets a clear standard for determining when the level of anticipated take has been exceeded.

Due to the difficulty of detecting take of Indiana bats caused by the proposed Action, the Service has decided to monitor the extent of taking using the acreage of suitable habitat removed by the Action. This surrogate measure is appropriate because the majority of the anticipated taking will result from habitat removal/alteration and activities associated with that alteration, and because it sets a clear standard for determining when the extent of taking is exceeded. The level of take anticipated in this BO is 2,000 acres per year over a 10-year period, totaling 20,000 acres of habitat. Instructions for monitoring and reporting take are provided below.

Northern Long-eared Bat

Based on our evaluation, the proposed Action is reasonably certain to cause the incidental take of 157 individual northern long-eared bats. This taking is expected in the form of harm. The mechanisms of this taking and the basis for our estimation of its extent are described in Section 11 (Effects of the Action) of this BO.

Table 5. Amount of incidental take of the northern-long eared bat caused by the Action.

Species	# of Individuals	Form of Take
Northern long-eared bat	117	Harm (Tree Removal, Summer Habitat)
	40	Harm (Tree Removal, Staging/Swarming Habitat)

Surrogate Measures for Monitoring

The Service anticipates the incidental taking of northern long-eared bats associated with this project will be difficult to detect for the following reasons:

- The individuals are small, mostly nocturnal, and when not hibernating, occupy forested habitats where they are difficult to find, capture, or observe;
- The northern long-eared bat forms small maternity colonies under loose bark or in the cavities of trees, and males and non-reproductive females may roost individually, which makes finding roost trees difficult;
- Finding dead or injured specimens during or following project implementation is unlikely; and
- Most incidental take is in the form of non-lethal harm and not directly observable.

When it is not practical to monitor take in terms of individuals of the listed species, the regulations at 50 CFR §402.14(i)(1)(i) indicate that an ITS may express the amount or extent of take using a surrogate (e.g., a similarly affected species, habitat, or ecological conditions), provided that the Service also:

- describes the causal link between the surrogate and take of the listed species; and
- sets a clear standard for determining when the level of anticipated take has been exceeded.

Due to the difficulty of detecting take of northern long-eared bats caused by the proposed Action, the Service has decided to monitor the extent of take using the acreage of suitable habitat removed by the Action. This surrogate measure is appropriate because the majority of the anticipated take will result from habitat removal/alteration and activities associated with that alteration, and because it sets a clear standard for determining when the extent of taking is exceeded. The level of take anticipated in this BO is 2,000 acres per year over a 10-year period, totaling 20,000 acres of habitat. Instructions for monitoring and reporting take are provided below.

Reasonable and Prudent Measures

The Action includes conservation measures, discussed previously, to minimize impacts to the covered species. Based on appropriate implementation of these measures, the Service believes that no additional “reasonable and prudent measures” will be necessary to minimize incidental take of the covered species caused by the Action.

Terms and Conditions

No reasonable and prudent measures to minimize incidental take caused by the Action are provided in this BO, so no terms and conditions for carrying out such measures are necessary.

Monitoring and Reporting Requirements

This section provides the specific instructions for such monitoring and reporting (M&R), including procedures for handling and disposing of any individuals of a species actually killed or injured. These M&R requirements are mandatory. Below we identify the M&R responsibilities of the KFO.

- 1.) The KFO shall keep records of the levels of incidental take exempted under this BO that are applied to projects that have agreed to voluntarily implement the Strategy. These records shall track the acres and the seasonality of habitat affected under each project and specify whether the affected habitat is known summer or staging/swarming habitat or potential summer or staging/swarming habitat.
- 2.) The KFO shall periodically audit projects implemented by cooperators to verify compliance with the Strategy. The selection of projects for these audits is at the KFO's discretion but shall occur at least once each year for the duration of this BO.
- 3.) The KFO, its cooperators, and any of their contractors must take care when handling dead or injured species covered by this BO or any other federally listed species that are found at project sites in order to preserve biological material in the best possible state and to protect the handler from exposure to diseases, such as rabies. Project cooperators are responsible for ensuring that evidence for determining the cause of death or injury is not unnecessarily disturbed. Reporting the discovery of dead or injured listed species is required in all cases to enable the Service to determine whether the level of incidental take exempted by this BO is exceeded and to ensure that the terms and conditions are appropriate and effective. Parties finding a dead, injured, or sick specimen of any endangered or threatened species, must promptly notify the Service's Division of Law Enforcement at 1875 Century Blvd., Suite 380, Atlanta, Georgia 30345 (Telephone: 404/679-7057) and the KFO at 330 West Broadway, Room 265, Frankfort, Kentucky 40601 (Telephone: 502/695-0468). The KFO is then responsible for notifying the RO Ecological Services program office at 1875 Century Boulevard, Suite 200, Atlanta, Georgia 30345 (Telephone 404/679-7085).

10. Conservation Recommendations

Section 7(a)(1) of the ESA directs Federal agencies to use their authorities to further the purposes of the Act by conducting conservation programs for the benefit of endangered and threatened species. Conservation recommendations are discretionary activities that an action agency may undertake to minimize or avoid the adverse effects of a proposed action, implement recovery plans, or develop information useful for the conservation of listed species. The Service offers the following conservation recommendations:

- 1.) The KFO will keep records of the amount of habitat purchased, managed, and protected and the amount of funding contributed to the Imperiled Bat Conservation Fund. The KFO

should use these records, and other information about conservation benefits to bats resulting from implementation of the Strategy, to inform an analysis of its overall effect in determining whether to renew and modify the program as the duration of the current program draws to a close.

- 2.) The KFO will create and maintain a geographic data base and query tool that allows cooperators to identify when proposed projects are located within known habitats for the covered species. The KFO will also maintain current pdf maps of the covered species known habitats for any cooperators who may not have access the geographic database.

11. Reinitiation Notice

This concludes formal consultation on the KFO's participation in and approval of cooperators' voluntary use of the Strategy and the effects on the covered species. Reinitiation of formal consultation is required where discretionary KFO involvement or control over the action has been retained (or is authorized by law) and if: (a) the amount or extent of incidental take is exceeded, (b) new information reveals effects of the KFO's action that may affect listed species or critical habitat in a manner or to an extent not considered in this consultation, (c) the KFO's action is later modified in a manner that causes an effect to the listed species or critical habitat not considered in this consultation, or (d) a new species is listed or critical habitat is designated that may be affected by the action. In instances where the amount or extent of incidental take is exceeded, any operations causing such take must cease pending reinitiation.

- 1) For this BO, the exempted incidental take would be exceeded when the take surpasses 2,000 acres of Indiana bat habitat removal or 2,000 acres of northern long-eared bat habitat removal in any year for a ten-year period. The total amount of incidental take, as measured by the habitat surrogate, covered for this period is 20,000 acres for each species. These are the amounts of habitat removal that are exempted from the prohibitions of section 9 of the Act by this BO.

This consultation was assigned FWS ID #2025-0127392. Please refer to this number in any correspondence concerning this consultation.

13. LITERATURE CITED

3D/International, Inc. 1996. 1996 field studies for interim mitigation for impacts to Indiana bats at the Indianapolis International Airport in Marion County, Indiana. 125pp.

Adams, R.A. 2010. Bat reproduction declines when conditions mimic climate change projections for western North America. *Ecology* 91(8):2437–2445.

Amelon, S., and D. Burhans. 2006. Conservation assessment: *Myotis septentrionalis* (northern long-eared bat) in the eastern United States. Pages 69–82 in Conservation assessments for five forest bat species in the eastern United States, Thompson, F. R., III, editor. U.S. Department of Agriculture, Forest Service, North Central Research Station, General Technical Report NC-260. St. Paul, Minnesota. 82pp.

Barbour, R.W., and W.H. Davis. 1969. *Bats of America*. The University of Kentucky Press, Lexington, Kentucky. 311pp.

Barclay, R.M.R., and A. Kurta. 2007. Ecology and Behavior of Bats Roosting in Tree Cavities and Under Bark. Chapter 2: pp. 17–60 in *Bats in Forests: Conservation and Management*, M.J. Lacki, J.P. Hayes, and A. Kurta, editors. The Johns Hopkins University Press, Baltimore, Maryland, 352pp.

Belwood, J.J. 1979. Feeding ecology of an Indiana bat community with emphasis on the endangered Indiana bat, *Myotis sodalis*. M.S. Thesis, University of Florida, Gainesville, FL. 103 pp.

Belwood, J.J. 2002. Endangered bats in suburbia: observations and concerns for the future. Pp. 193-198 in A. Kurta and J. Kennedy (eds.), *The Indiana bat: biology and management of an endangered species*. Bat Conservation International, Austin, TX.

Bergeson, S.M., T.C. Carter, and M.D. Whitby. 2013. Partitioning of foraging resources between sympatric Indiana and little brown bats. *J. of Mammalogy* 94(6):1311-1320.

Blehert, D.S., A.C. Hicks, M.J. Behr, C.U. Meteyer, B.M. Berlowski-Zier, E.L. Buckles, J. Coleman T.H., S.R. Darling, A. Gargas, R. Niver, J.C. Okoniewski, R.J. Rudd, and W.B. Stone. 2009. Bat White-Nose Syndrome: an emerging fungal pathogen? *Science*, 323:227.

Bohrman, J., and D. Fecske. 2013. White-nose syndrome surveillance and summer monitoring of bats at Great Swamp National Wildlife Refuge, Morris County, New Jersey. A final report prepared for United States Fish and Wildlife Service, 115pp.

Brack V. Jr. 2007. Temperatures and locations used by hibernating bats, including *Myotis sodalis* (Indiana bat), in a limestone mine: implications for conservation and management. *Journal of Environmental Management*, 40:739–746.

Brack, V., Jr. and R.K. LaVal. 1985. Food habits of the Indiana bat in Missouri. *Journal of Mammalogy* 66:308-315.

Brack V. Jr., and J.O. Whitaker. 2001. Foods of the northern myotis, *Myotis septentrionalis*, from Missouri and Indiana, with notes on foraging. *Acta Chiropterologica*, 3(2):203–210.

Brack V. Jr., S.A. Johnson, and R.K. Dunlap. 2003. Wintering populations of bats in Indiana, with emphasis on the endangered Indiana myotis, *Myotis sodalis*. *Proceedings of the Indiana Academy of Science* 112(1):61–74.

Broders, H.G., G.J. Forbes, S. Woodley, and I.D. Thompson. 2006. Range extent and stand selection for roosting and foraging in tree roosting northern long-eared bats and little brown bats in the Greater Fundy ecosystem, New Brunswick. *J. Wildlife Management* 70(5):1174–1184.

Burke, H.S. Jr. 1999. Maternity colony formation in *Myotis septentrionalis* using artificial roosts: the rocket box, a habitat enhancement for woodland bats? *Bat Research News*, 40:77–78.

Burles, D.W., R.M Brigham, R.A. Ring, and T.E. Reimchen. 2009. Influence of weather on two insectivorous bats in a temperate Pacific Northwest rainforest. *Canadian Journal of Zoology* 87:132–138.

Butchkoski, C. M., and J. M. Hassinger. 2002. Ecology of a maternity colony roosting in a building. Pp. 130–142 in A. Kurta and J. Kennedy, Eds. *The Indiana bat: biology and management of an endangered species*. Bat Conservation International, Austin, Texas, 253 pp.

Butchkoski, C.M., and S. Bearer. 2016. Summer bat-netting trends in Pennsylvania. Pp. 137–151, In C.M. Butchkoski, D.M. Reeder, G.G. Turner, and H.P. Whidden (Eds.). *Conservation and Ecology of Pennsylvania's Bats*. The Pennsylvania Academy of Science, East Stroudsburg, PA. 284 pp.

Caceres, M.C., and M.J. Pybus. 1997. Status of the northern long-eared bat (*Myotis septentrionalis*) in Alberta. Alberta Environmental Protection, Wildlife Management Division, Wildlife Status Report No. 3, Edmonton, AB, 19pp.

Caceres, M.C., and R.M.R. Barclay. 2000. *Myotis septentrionalis*. *Mammalian Species*, 634:1–4.

Caire, W., R.K. LaVal, M.L. LaVal, and R. Clawson. 1979. Notes on the ecology of *Myotis keenii* (Chiroptera, Vespertilionidae) in Eastern Missouri. *American Midland Naturalist*, 102(2): 404–407.

Callahan, E.V., III. 1993. Indiana bat summer habitat requirements. M.S. Thesis, University of Missouri, Columbia, MO.

Callahan, E.V., R.D. Drobney, and R.L. Clawson. 1997. Selection of summer roosting sites by Indiana bats (*Myotis sodalis*) in Missouri. *Journal of Mammalogy* 78:818–825.

Carter, T.C., and G. Feldhamer. 2005. Roost tree use by maternity colonies of Indiana bats and northern long-eared bats in southern Illinois. *Forest Ecology and Management*, 219:259–268.

Cheng, T.L., J.D. Reichard, J.T.H. Coleman, T.J. Weller, W.E. Thogmartin, B.E. Reichert, A.B. Bennett, H.G. Broders, J. Campbell, K. Etchison, D.J. Feller, R. Geboy, T. Hemberger, C. Herzog, A.C. Hicks, S. Houghton, J. Humber, J.A. Kath, R.A. King, S.C. Loeb, A. Massé, K.M. Morris, H. Niederriter, G. Nordquist, R.W. Perry, R.J. Reynolds, D.B. Sasse, M.R. Scafani, R.C. Stark, C.W. Stihler, S.C. Thomas, S.C. Thomas, G.G. Turner, S. Webb, B. Westrich, and W.F. Frick. 2021. The scope and severity of white-nose syndrome on hibernating bats in North America. *Conservation Biology* 2021:1–12.

Clark, B.K., J.B. Bowles, and B.S. Clark. 1987. Summer status of the endangered Indiana bat. *The American Midland Naturalist* 118(1): 32-39.

Clawson, R.L. 2002. Trends in population size and current status. Pp. 2-8 in A. Kurta and J. Kennedy (eds.), *The Indiana Bat: Biology and Management of an Endangered Species*. Bat Conservation International, Austin, TX.

Coleman, J.T.H., and J.D. Reichard. 2014. Bat white-nose syndrome in 2014: A brief assessment seven years after discovery of a virulent fungal pathogen in North America. *Outlooks on Pest Management* 25(6):374-377.

Cope, J.B., and S.R. Humphrey. 1972. Reproduction of the bats *Myotis keenii* and *Pipistrellus subflavus* in Indiana. *Bat Research News* 13(9).

Cope, J.B., and S.R. Humphrey. 1977. Spring and autumn swarming behavior in the Indiana bat, *Myotis sodalis*. *Journal of Mammalogy* 58:93-95.

Cope, J.B., A.R. Richter, and R.S. Mills. 1973. A summer concentration of the Indiana bat, *Myotis sodalis*, in Wayne County, Indiana. Earlham College. Richmond, IN.

Cope, J.B., A.R. Richter, and R.S. Mills. 1974. Concentrations of the Indiana bat, *Myotis sodalis*, in Wayne County, Indiana. *Proceedings of the Indiana Acadamy of Science* 83:482-484.

Copperhead Environmental Consulting, Inc. 2017. Spring Migration of Female Indiana Bats (*Myotis sodalis*) from Signature and Tobaccoport Saltpeter Caves in Tennessee. Report submitted to the U.S. Fish and Wildlife Service Tennessee Field Office.

Copperhead Environmental Consulting, Inc. 2018. Final Report: Statewide Mitigation and Monitoring for Indiana Bats: Arkansas DOT Job 001799. Report submitted to the Arkansas Department of Transportation and U.S. Fish and Wildlife Service Arkansas Field Office.

Copperhead Environmental Consulting, Inc. 2019. Final Report: Statewide Mitigation and Monitoring for Indiana Bats: Arkansas DOT Job 001799. Report submitted to the Arkansas Department of Transportation and U.S. Fish and Wildlife Service Arkansas Field Office.

Cryan P.M., C.U. Meteyer, D.S. Blehert, J.M. Lorch, D.M. Reeder, G.G. Turner, J. Webb, M. Behr, M. Verant, R.E. Russell, and K.T. Castle. 2013. Electrolyte Depletion in white-nose syndrome bats. *Journal of Wildlife Diseases* 49:398-402.

Davis, W.H., and H.B. Hitchcock. 1965. Biology and Migration of the Bat, *Myotis lucifugus*, in New England. *Journal of Mammalogy*, 46(2):296–313.

De la Cruz, J.L., R.L. Ward, and E.S. Schroder. 2018. Landscape characteristics related to use of artificial roosts by northern long-eared bats in north-central West Virginia. *Northeastern Naturalist*, 25(3):487–501.

Dodd, L.E., E.G. Chapman, J.D. Harwood, M.J. Lacki, and L.K. Rieske. 2012. Identification of prey of *Myotis septentrionalis* using DNA-based techniques. *Journal of Mammalogy*, 93(4):1119–1128.

Dowling, Z.R., and D.I. O'Dell. 2018. Bat use of an island off the coast of Massachusetts. *Northeastern Naturalist*, 25(3):362–382.

Drobney, R.D. and R.L. Clawson. 1995. Indiana bats. Pp. 97-98 in LaRoe, E.T., G.S. Farris, C.E. Puckett, P.D. Doran, and M.J. Mac (eds.), *Our living resources: a report to the nation on the distribution, abundance, and health of U.S. plants, animals, and ecosystems*. U.S. Department of the Interior, National Biological Service, Washington, DC. 530 pp. Available at: <http://biology.usgs.gov/s+t/noframe/c164.htm>.

Duchamp, J.E. 2006. Modeling bat community structure and species distribution across fragmented landscapes within the Upper Wabash River Basin. PhD dissertation. Purdue University, West Lafayette, Indiana.

Easterla, D.A. 1968. Parturition of Keen's myotis in southwestern Missouri. *Journal of Mammalogy*, 49(4):770.

Ehlman, S.M., J.J. Cox, and P.H. Crowley. 2013. Evaporative water loss, spatial distributions, and survival in white-nose syndrome affected little brown myotis: a model. *Journal of Mammalogy* 94(3):572–583.

Ecology and Environment, Inc. 2009. Draft Report: Indiana Bat (*Myotis sodalis*) Field Studies at Shuteye Creek Wind Farm Adair and Sullivan Counties, Missouri.

Environment Yukon. 2011. Yukon Bats. Government of Yukon, Environment Yukon, Whitehorse, Yukon. 22pp.

Farmer, A.H., B.S. Cade, and D.F. Stauffer. 2002. Evaluation of a habitat suitability index model. Pp. 172-179 in A. Kurta and J. Kennedy (eds.), *The Indiana bat: biology and management of an endangered species*. Bat Conservation International, Austin, TX.

Feldhamer, G.A., T.C. Carter, A.T. Morzillo, and E.H. Nicholson. 2003. Use of bridges as day roosts by bats in southern Illinois. *Transactions of the Illinois State Academy of Science* 96(2):107–112.

Feldhamer, G.A., T.C. Carter, and J.O. Whitaker Jr. 2009. Prey consumed by eight species of insectivorous bats from southern Illinois. *The American Midland Naturalist*, 162(1):43–51.

Fenton, M.B. 1969. Summer activity of *Myotis lucifugus* (Chiroptera:Vespertilionidae) at hibernacula in Ontario and Quebec. Canadian Journal of Zoology, 47(4):597–602.

Fill, C.T., C.R. Allen, J.F. Benson, and D. Twidwell. 2021. Roost use and movements of northern long-eared bats in a southeast Nebraska agricultural landscape. The American Midland Naturalist 185(2):241-248.

Fitch, J.H., and K.A. Shump, Jr. 1979. *Myotis keenii*. Mammalian Species, 121:1–3. Foden, W.B., B.E. Young, H.R. Akcakaya, R.A. Garcia, A.A. Hoffmann, B.A. Stein, C.D. Thomas, C.J. Wheatley, D. Bickford, J.A. Carr, D.G. Hole, T. G. Martin, M. Pacifici, J.W. Pearce-Higgins, P.J. Platts, P. Cisconti, J.E.M. Watson, and B. Huntley. 2018. Climate change vulnerability assessment of species. WIREs Climate Change 10(e551):1–36. <https://doi.org/10.1002/wcc.551>.

Foster, R.W., and A. Kurta. 1999. Roosting ecology of the Northern bat (*Myotis septentrionalis*) and comparisons with the endangered Indiana bat (*Myotis sodalis*). Journal of Mammalogy, 80(2):659–672.

Franci, K.E., W.M. Ford, D.W. Sparks, and V. Brack. 2012. Capture and reproductive trends in summer bat communities in West Virginia: Assessing the impact of white-nose syndrome. Journal of Fish and Wildlife Management 3(1): 33–42.

Frick, W.F., D.S. Reynolds, and T.H. Kunz. 2010. Influence of climate and reproductive timing on demography of little brown myotis *Myotis lucifugus*. Journal of Animal Ecology, 79:128– 136.

Frick, W.F., S.J. Puechmaille, J.R. Hoyt, B.A. Nickel, K.E. Langwig, J.T. Foster, K.E. Barlow, T. Bartonička, D. Feller, A. Haarsma, C. Herzog, I. Horaček, J. Kooij, B. Petrov, R. Reynolds, L. Rodrigues, C.W. Stihler, G.G. Turner, and A.M. Kilpatrick. 2015. Disease alters macroecological patterns of North American bats. Global Ecology and Biogeography, 24:741– 749.

Frick, W.F., T.L. Cheng, K.E. Langwig, J.R. Hoyt, A.F. Janicki, K.L. Parise, J.T. Foster, and A.M. Kilpatrick. 2017. Pathogen dynamics during invasion and establishment of white-nose syndrome explain mechanisms of host persistence. Ecology 98: 624–631.

Gardner, J.E., and E.A. Cook. 2002. Seasonal and geographic distribution and quantification of potential summer habitat. Pp. 9-20 in A. Kurta and J. Kennedy (eds.), The Indiana bat: biology and management of an endangered species. Bat Conservation International, Austin, TX.

Gardner, J.E., J.D. Garner, and J.E. Hofmann. 1991. Summer roost selection and roosting behavior of *Myotis sodalis* (Indiana bat) in Illinois. Unpublished report to Region-3 U.S. Fish and Wildlife Service, Fort Snelling, MN. 56 pp.

Garner, J.D., and J.E. Gardner. 1992. Determination of summer distribution and habitat utilization of the Indiana bat (*Myotis sodalis*) in Illinois. Unpublished Report. Endangered Species Coordinator, Region 3, Service, Twin Cities, MN.

Garroway, C.J., and H.G. Broders. 2007. Nonrandom association patterns at northern long-eared bat maternity roosts. *Canadian Journal of Zoology*, 85:956–964.

Geipel, I., M.J. Smeekes, W. Halfwerk, and R.A. Page. 2019. Noise as an informational cue for decision-making: the sound of rain delays bat emergence. *Journal of Experimental Biology*:222, jeb192005.

Glover, A.M., and J.D. Altringham. 2008. Cave selection and use by swarming bat species. *Biological Conservation*, 141:1493–1504.

Gorman, K. M., S.M. Deeley, E.L. Barr, S.R. Freeze, N. Kalen, M.S. Muthersbaugh, and W.M. Ford. 2022. Broad-scale geographic and temporal assessment of northern long-eared bat (*Myotis septentrionalis*) maternity colony-landscape association. *Endangered Species Research*, 47, 119-130. <https://doi.org/10.3354/esr01170> (Accessed October 8, 2024).

Grider, J.F., A.L. Larsen, J.A. Homyack, and M.C. Kalcounis-Rueppell. 2016. Winter activity of coastal plain populations of bat species affected by white-nose syndrome and wind energy facilities. *PLoS one*, 11(11):e0166512.

Griffin, D.R. 1940a. Reviewed notes on the life histories of New England cave bats. *Journal of Mammalogy*, 21(2):181–187.

Griffin, D.R. 1945. Travels of banded cave bats. *Journal of Mammalogy*, 26(1):15–23.

Griffith, L.A., and J.E. Gates. 1985. Food habits of cave-dwelling bats in the central Appalachians. *Journal of Mammalogy* 66(3):451–460.

Grindal, S., T. Collard, R. Brigham, and R.M.R. Barclay. 1992. The influence of precipitation on reproduction by *Myotis* bats in British Columbia. *The American Midland Naturalist* 128(2):339–344. doi:10.2307/2426468.

Grue, C.E., P.L. Gibert, and M.E. Seeley. 1997. Neurophysiological and behavioral changes in non-target wildlife exposed to organophosphate and carbamate pesticides: thermoregulation, food consumption and reproduction. *American Zoologist* 37:369-388.

Gumbert, M.W., J.M. O'Keefe, and J.R. MacGregor. 2002. Roost fidelity in Kentucky Pp. 143-152 in A. Kurta and J. Kennedy (eds.), *The Indiana Bat: Biology and Management of an Endangered Species*. Bat Conservation International, Austin, TX.

Hall, J.S., and F.J. Brenner. 1968. Summer netting of bats at a cave in Pennsylvania. *Journal of Mammalogy*, 49(4):779–781.

Hawkins, J.A., J. Jaskula, A. Mann, and V. Brack, Jr. 2005. Habitat Conservation Plan: 2004 telemetry study of autumn swarming behavior of the Indiana bat (*Myotis sodalis*). Report prepared for the Indiana Department of Natural Resources, Indianapolis, IN. 25 pp. plus appendices.

Hawkins, J.A., P.L. Sewell, and M.W. Gumbert. 2008. Final Report: Indiana bat survey and anthropogenic stimuli study conducted at US Army Garrison Fort Knox and Brashears creek study sites during summer 2007. Final Report submitted to ICI Services, LLC.

Henderson, L.E., and H.G. Broders. 2008. Movements and resource selection of the northern long-eared myotis (*Myotis septentrionalis*) in a forest-agriculture landscape. *Journal of Mammalogy*, 89(4):952–963.

Hendrick, L.B. 2008. Evaluation of the impacts of highway construction on sediment and benthic macroinvertebrates in Appalachian streams. PhD dissertation. West Virginia University, Morgantown, West Virginia.

Henley, W.F., M.A. Patterson, R.J. Neves, and A.D. Lemly. 2000. Effects of Sedimentation and turbidity on lotic food webs: a concise review for natural resource managers. *Reviews in Fisheries Science*. 8(2): 125-139.

Hobson, C.S., and Holland, J.N. 1995. Post-hibernation Movement and Foraging Habitat of a Male Indiana Bat, *Myotis sodalis* (Chiroptera: Vespertilionidae), in West Virginia. 101pp.

Humphrey, S.R. 1978. Status, winter habitat, and management of the endangered Indiana bat, *Myotis sodalis*. *Florida Scientist* 41:65-76.

Humphrey, S.R., and J.B. Cope. 1977. Survival rates of the endangered Indiana bat, *Myotis sodalis*. *Journal of Mammalogy* 58:32-36.

Humphrey, S.R., A.R. Richter, and J.B. Cope. 1977. Summer habitat and ecology of the endangered Indiana bat, *Myotis sodalis*. *Journal of Mammalogy*. 58:334-346.

Hyzy, B.A., R.E. Russel, A. Silvis, W.M. Ford, J. Riddle, and K. Russel. 2020. Investigating maternity roost selection by northern long-eared bats at three sites in Wisconsin. *Endangered Species Research*, 41:55–65, 2020.

Jachowski, D.S., J.B. Johnson, C.A. Dobony, J.W. Edwards, and W.M. Ford. 2014. Space use and resource selection by foraging Indiana bats at the northern edge of their distribution. *Endangered Species Research* 24:149-157.

Johnson J.B., J.W Edwards, W.M. Ford, K. Cunningham, and J.L. Rodrigue. 2011. Fort Drum Military Installation cantonment area Indiana myotis survey. Rutland, Vermont: U.S. Forest Service Green Mountain National Forest. Final Report 09-PA-11092000-016.

Johnson, J.B., W.M. Ford, and J.W. Edwards. 2012. Roost networks of northern myotis (*Myotis septentrionalis*) in a managed landscape. *Forest Ecology and Management*, 266:223-231.

Jordan, G. W. 2020. Status of an anomalous population of northern long-eared bats in coastal North Carolina. *Journal of Fish and Wildlife Management* 11(2):665–678.

Kiser, J.D. and C.L. Elliott. 1996. Foraging habitat, food habits, and roost tree characteristics of the Indiana bat (*Myotis sodalis*) during autumn in Jackson County, Kentucky. Report

prepared for Kentucky Department of Fish and Wildlife Resources, Nongame Program, Frankfort, KY. 65 pp.

Kniowski, A.B., and S.D. Gehrt. 2014. Home range and habitat selection of the Indiana bat in an agricultural landscape. *Journal of Wildlife Management* 78(3):503-512.

Krochmal, A.R., and D.W. Sparks. 2007. Timing of birth and estimation of age of juvenile *Myotis septentrionalis* and *Myotis lucifugus* in west-central Indiana. *Journal of Mammalogy*, 88(3):649–656.

Kniowski, Andrew B. 2011. Summer Ecology of the Indiana bat (*Myotis sodalis*) in an Agricultural Landscape. Ohio State University. PhD Dissertation. 184 pages.

Kunz, T.H. 1971. Reproduction of some Vespertilionid bats in central Iowa. *American Midland Naturalist*, 86(2):477–486.

Kunz, T.H. 1973. Temporal and spatial components of bat activity in central Iowa. *Journal of Mammalogy*, 54(1):14–32.

Kurta, A. 2013. Peer review of the 12-month finding on the eastern small-footed bat and northern long-eared bat.

Kurta, A. and H. Rice. 2002. Ecology and management of the Indiana bat in Michigan. *Michigan Academician* 33:361-376.

Kurta A., G.P. Bell, K.A. Nagy, and T.H. Kunz. 1989. Energetics of pregnancy and lactation in free-ranging little brown bat (*Myotis lucifugus*). *Physiological Zoology* 62: 804–818.

Kurta, A., D. King, J.A. Teramoto, J.M. Sibley, and K.J. Williams. 1993. Summer roosts of the endangered Indiana bat (*Myotis sodalis*) on the northern edge of its range. *American Midland Naturalist* 129:132-138.

Kurta, A., K.J. Williams, and R. Mies. 1996. Ecological, behavioral, and thermal observations of a peripheral population of Indiana bats (*Myotis sodalis*). Pages 102-117 in *Bats and Forests Symposium* (R. M. R. Barclay and R. M. Brigham, eds.). Research Branch, British Columbia Ministry of Forests, Victoria, British Columbia, Canada, Working Paper 23:1-292.

Kurta, A., J. Caryl, and T. Tipps. 1997. Bats and Tippy Dam: species composition, seasonal use, and environmental parameters. *Michigan Academician*, 29(4):473–490.

Kurta, A., S.W. Murray, and D.H. Miller. 2002. Roost selection and movements across the summer landscape. Pp. 118-129 in A. Kurta and J. Kennedy (eds.), *The Indiana bat: biology and management of an endangered species*. Bat Conservation International, Austin, TX.

Lacki, M.J., and J.H. Schwierjohann. 2001. Day-roost characteristics of northern bats in mixed mesophytic forest. *The Journal of Wildlife Management*, 65(3):482–488.

Lacki, M. J., S. K. Amelon, and M.D. Baker. 2007. Foraging ecology of bats in forests *in* Lacki, M.J., J.P. Hayes, and A. Kurta (eds), Bats in forests: conservation and management. Johns Hopkins University Press, Baltimore, MD.

LaVal, R.K., and M.L. LaVal. 1980. Ecological studies and management of Missouri bats, with emphasis on cave-dwelling species. Missouri Department of Conservation, Conservation Commission of the State of Missouri, Terrestrial Series No. 8.

LaVal, R.K., R.L. Clawson, M.L. LaVal, and W. Caire. 1977. Foraging behavior and nocturnal activity patterns of Missouri bats, with emphasis on the endangered species *Myotis griseescens* and *Myotis sodalis*. *Journal of Mammalogy*, 58(4):592–599.

Lee, Ya-Fu. 1993. Feeding Ecology of the Indiana bat, *Myotis sodalis*, and Resource Partitioning with *Myotis keenii* and *Myotis lucifugus*. M.S. Thesis, University of Tennessee, Knoxville. 77 pp.

Lee, Y.F., and G.F. McCracken. 2004. Flight activity and food habits of three species of Myotis bats (Chiroptera: Vespertilionidae) in sympatry. *Zool. Studies*. 43:589–597.

Loeb, S.C. and E.A. Winters. 2013. Indiana bat summer maternity distribution: effects of current and future climates. *Ecology and Evolution* 3(1): 103-114.

Lorch, J.M., C.U. Meteyer, M.J. Behr, J.G. Boyles, P.M. Cryan, A.C. Hicks, A.E. Ballmann, J.T.H. Coleman, D.N. Redell, D.M. Reeder, and D.S. Blehert. 2011. Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* 480:376–379.

Lowe, A.J. 2012. Swarming behaviour and fall roost-use of little brown (*Myotis lucifugus*) and northern long-eared bats (*Myotis septentrionalis*) in Nova Scotia, Canada. Master's Thesis. St. Mary's University, Halifax, Nova Scotia, Canada. 88pp.

MacGregor J., J. Kiser, M. Gumbert, and T. Reed. 1999. Autumn roosting habitat of male Indiana bats (*Myotis sodalis*) in a managed forest setting in Kentucky. Pp. 169-170 in J. Stringer and D. Loftis (eds), Proceedings 12th central hardwood forest conference, 1999 February 28-March 1-2, Lexington, KY. General Technical Report SRS-24. U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, NC.

Massachusetts Department of Fish and Game. 2012. Unpublished data

Menzel, J.M., W.M. Ford, M.A. Menzel, T.C. Carter, J.E. Gardner, J.D. Garner, and J.E. Hofmann. 2005. Summer habitat use and home-range analysis of the endangered Indiana bat. *Journal of Wildlife Management* 69(1):430-436.

Meteyer, C.U., E.L. Buckles, D.S. Blehert, A.C. Hicks, D.E. Green, V. Shearn-Bochsler, N.J. Thomas, A. Gargas, and M.J. Behr. 2009. Histopathologic criteria to confirm white-nose syndrome in bats (cited as: Pathology criteria for confirming white-nose syndrome in bats). *Journal of Veterinary Diagnostic Investigation* 21:411–414.

Mikula, P., F. Morelli, R.K. Lucan, D.N. Jones, and P. Tryjanowski. 2016. Bats as prey of diurnal birds: a global perspective. *Mammal Review* 46: 160-174.

Mumford, R.E., and L.L. Calvert. 1960. *Myotis sodalis* evidently breeding in Indiana. *Journal of Mammalogy* 41:512.

Mumford R.E., and J.B. Cope. 1964. Distribution and status of the chiroptera of Indiana. *American Midland Naturalist*, 72(2):473–489.

Murray, S.W., and A. Kurta. 2002. Spatial and temporal variation in diet. Pp. 182-192 in A. Kurta and J. Kennedy (eds.), *The Indiana bat: biology and management of an endangered species* Bat Conservation International, Austin, TX.

Murray, S.W., and A. Kurta. 2004. Nocturnal activity of the endangered Indiana bat (*Myotis sodalis*). *Journal of Zoology* 262:197–206.

Nagorsen, D.W., and R.M. Brigham. 1993. *Bats of British Columbia*. Royal British Columbia Museum, Victoria, and the University of British Columbia Press, Vancouver. 164 pp.

Olivera-Hyde, M., A. Silvis, E.M. Hallerman, W.M. Ford, and E.R. Britzke. 2019 Relatedness within and among northern long-eared bat (*Myotis septentrionalis*) colonies at a local scale. *Canada Journal of Zoology*. 97: 724–735.

Owen, S.F., M.A. Menzel, W.M. Ford, J.W. Edwards, B.R. Chapman, K.V. Miller, and P.B. Wood. 2002. Roost tree selection by maternal colonies of Northern long-eared Myotis in an intensively managed forest. USDA Forest Service. Newtown Square, Pennsylvania. 10 pp.

Parsons, S., K.L. Lewis, and J.M. Psyllakis. 2003. Relationships between roosting habitat of bats and decay of aspen in the sub-boreal forests of British Columbia. *Forest Ecology and Management* 177:559–570.

Patriquin, K.J., M.L. Leonard, H.G. Broders, and C.J. Garroway. 2010. Do social networks of female northern long-eared bats vary with reproductive period and age? *Behavioral Ecology and Sociobiology* 84:899–913.

Patriquin, K.J., M.L. Leonard, H.G. Broders, W.M. Ford, E.R. Britzke, and A. Silvis. 2016. Weather as a proximate explanation for fission–fusion dynamics in female northern long-eared bats. *Animal Behaviour* 122:47–57.

Patterson, B.D., M.R. Willig, and R.D. Stevens. 2003. Trophic strategies, niche partitioning, and patterns of ecological organization. In T.H. Kunz and M.B. Fenton (Eds), *Bat Ecology*. The University of Chicago Press.

Perry, R.W. 2011. Fidelity of bats to forest sites revealed from mist-netting recaptures. *Journal of Fish and Wildlife Management* 2(1):112–116; e1944-687X. doi: 10.3996/082010-JFWM-030

Perry, R. W., and R. E. Thill. 2007. Roost selection by male and female northern long-eared bats in a pine-dominated landscape. *Forest Ecology and Management* 247:220-226.

Pettit, J.L. and J.M. O'Keefe. 2017. Impacts of white-nose syndrome observed during long-term monitoring of a Midwestern bat community. *Journal of Fish and Wildlife Management* 8(1): 69-78.

Racey, P. 1982. Ecology of bat reproduction. pp 57–104 in *Ecology of Bats* (ed Kunz, T.). Boston University, Boston, MA.

Raesly, R.L., and J.E. Gates. 1987. Winter habitat selection by north temperate cave bats. *American Midland Naturalist* 1987:15–31.

Racey, P.A., and A.C. Entwistle. 2003. Conservation ecology of bats. In T.H. Kunz and M.B. Fenton (Eds), *Bat Ecology*. The University of Chicago Press.

Randall, J., and H.G. Broders. 2014. Identification and characterization of swarming sites used by bats in Nova Scotia, Canada. *Acta Chiropterologica*, 16:109–116.

Ratcliffe, J.M., and J.W. Dawson. 2003. Behavioral flexibility: the little brown bat, *Myotis lucifugus*, and the northern long-eared bat, *M. septentrionalis*, both glean and hawk prey. *Animal Behaviour*, 66:847–856.

Redford, K.H., G. Amato, J. Baillie, P. Beldomenico, E.L. Bennett, N. Clum, R. Cook, G. Fonseca, S. Hedges, F. Launay, S. Lieberman, G.M. Mace, A. Murayama, A. Putnam, J.G. Robinson, H. Rosenbaum, E.W. Sanderson, S.N. Stuart, P. Thomas, and J. Thorbjarnarson. 2011. What does it mean to conserve a (vertebrate) species? *BioScience* 61:39–48.

Reeder, D.M., C.L. Frank, G.G. Turner, C.U. Meteyer, A. Kurta, E.R. Britzke, M.E. Vodzak, S.R. Darling, C.W. Stihler, A.C. Hicks, R. Jacob, L.E. Grieneisen, S.A. Brownlee, L.K. Muller, and D.S. Blehert. 2012. Frequent arousal from hibernation linked to severity of infection and mortality in bats with white-nose syndrome. *Plos One* 7:e38920. <http://dx.doi.org/10.1371%2Fjournal.pone.0038920>

Reichard, J.D. and T.H. Kunz. 2009. White-nose syndrome inflicts lasting injuries to the wings of little brown myotis (*Myotis lucifugus*). *Acta Chiropterologica* 11:457–464. <https://doi.org/10.3161/150811009X485684>

Richter, A. R., S.R. Humphrey, J.B. Cope, V. Brack. 1993. Modified cave entrances: thermal effect on body mass and resulting decline of endangered Indiana bats (*Myotis sodalis*). *Conservation Biology*, 7(2):407-415.

Roby, P.L., Seiter, W.R., Brandenburg, M.G., Armstrong, M.P., and M.W. Gumbert. In Press. Long-term monitoring of Indiana bats on Fort Knox Military Base, Kentucky. *Journal of Fish and Wildlife Management*.

Roby, P.L., M.W. Gumbert, and M.J. Lacki. 2019. Nine years of Indiana bat (*Myotis sodalis*) spring migration behavior. *Journal of Mammalogy* XX(X):1-11.

Rogers, M.L. 2020. Master of Science. Eastern Kentucky University. Assessment of Indiana Bat Reproductive Condition, Recapture Trends, and Temperatures of Artificial Roosts in Kentucky. 118pp

Romme, R.C., A.B. Henry, R.A. King, T. Glueck, and K.Tyrell. 2002. Home Range Near Hibernacula in Spring and Autumn. 158pp.

Sasse, D.B., and P.J. Pekins. 1996. Summer roosting ecology of northern long-eared bats (*Myotis septentrionalis*) in the white mountain national forest. Bats and Forests Symposium October 1995, Victoria, British Columbia, Canada, pp. 91–101.

Sasse, D.B., M.L. Caviness, M.J. Harvey, J.L. Jackson, and P.N. Jordan. 2014. Journal of the Arkansas Academy of Science 68(32):170–173.

Sealander, J.A., and G.A. Heidt. 1990. Arkansas Mammals. The University of Arkansas Press, Fayetteville, Arkansas. 308 pp.

Seiter, W., P. Roby, and C. Birdsall. 2022. Indiana Bat Monitoring and Foraging Analysis in Hunting Area 6, Fort Knox, 2021. Final report submitted to Lee Andrews and Mike Armstrong, USFWS, Frankfort, KY. 152pp

Silvis, A., W.M. Ford, E.R. Britzke, N.R. Beane, and J.B. Johnson. 2012. Forest succession and maternity day-roost selection by *Myotis septentrionalis* in a mesophytic hardwood forest. International Journal of Forestry Research 2012:1–8.

Silvis, A., A.B. Kniowski, S.D. Gehrt, and W.M. Ford. 2014a. Roosting and foraging social structure of the endangered Indiana bat (*Myotis sodalis*). PLoS ONE 9(5):1-12.

Silvis, A., W.M. Ford, E.R. Britzke, and J.B. Johnson. 2014b. Association, roost use and simulated disruption of *Myotis septentrionalis* maternity colonies. Behavioural Processes, 103:283–290.

Silvis, A., W.M. Ford, and E.R. Britzke. 2015. Effects of hierarchical roost removal on northern long-eared bat (*Myotis septentrionalis*) maternity colonies. PLoS ONE, 10(1):1–17.

Silvis, A., R.W. Perry, and W.M. Ford. 2016. Relationships of three species of bats impacted by White-Nose Syndrome to forest condition and management. Gen. Tech. Rep. SRS-214. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 48 p.

Simmons, N.B. 2005. Order Chiroptera: Subfamily Myotinae. p. 516 in Mammal species of the world: a taxonomic and geographic reference, D.E. Wilson and D.M. Reeder, editors. The John Hopkins University Press, Baltimore, Maryland. 2000 pp.

Smith, W.B., P.D. Miles, J.S. Vissage, and S.A. Pugh. 2003. Forest resources of the United States, 2002. U.S. Department of Agriculture, Forest Service, North Central Research Station, St. Paul, MN. General Technical Report NC-241. 137 pp. Available at: <http://www.ncrs.fs.fed.us/pubs/viewpub.asp?key=1987>. (Accessed June 5, 2017).

Spanjer, G.R., and M.B. Fenton. 2005. Behavioral responses of bats to gates at caves and mines. Wildlife Society Bulletin 33:1101–1112

Sparks D.W., M.T. Simmons, C.L. Gummer, and J.E. Duchamp. 2003. Disturbance of roosting bats by woodpeckers and raccoons. *Northeastern Naturalist* 10:105-8.

Sparks, D.W., J.O. Whitaker, Jr., and C.M. Ritz. 2005. Foraging ecology of the endangered Indiana bat. Pp. 15-27 in K.C. Vories and A. Harrington (eds.), *The Proceedings of the Indiana bat and coal mining: a technical interactive forum*. Office of Surface Mining, U.S. Department of the Interior, Alton, IL.

Stantec 2010. Unpublished data from ongoing research along the Bellefontaine ridge in west - central Ohio submitted to U.S. Fish and Wildlife Service.

Thomson, C.E. 1982. *Myotis sodalis*. *Mammalian Species*. 163:1-5.

Timpone, J.C., J.G. Boyles, K.L. Murray, D.P. Aubrey, and L.W. Robbins. 2010. Overlap in roosting habits of Indiana bats (*Myotis sodalis*) and northern bats (*Myotis septentrionalis*). *American Midland Naturalist* 163:115–123.

Turner, G.G., D.M. Reeder, and J.T.H. Coleman. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North American bats and a look to the future. *Bat Research News* 52(2):13–27.

U.S. Army Garrison Fort Drum. 2011. Biological Assessment on the proposed activities on the Fort Drum Military Installation, Fort Drum, New York (2012-2014) for the Federally-endangered Indiana bat (*Myotis sodalis*).

U.S. Fish and Wildlife Service (USFWS). 1983. Recovery Plan for the Indiana Bat. Twin Cities, MN.

U.S. Fish and Wildlife Service (USFWS). 1999. Final biological opinion for the proposed streambank stabilization and the Yano Range and upgrade of the Wilcox Tank Range at Fort Knox, Kentucky. USFWS Cookeville Field Office, Cookeville, TN. 18 pp.

U.S. Fish and Wildlife Service (USFWS). 2002. Final biological opinion on the application for an incidental take permit for the federally endangered Indiana bat (*Myotis sodalis*) for the Six Points Road interchange and associated development. USFWS Bloomington Field Office, Bloomington, IN. 36 pp.

U.S. Fish and Wildlife Service (USFWS). 2007. Indiana bat (*Myotis sodalis*) Draft Recovery Plan: First Revision. Fort Snelling, MN. 258 pp.

U.S. Fish and Wildlife Service (USFWS). 2011. Indiana Bat Section 7 and Section 10 Guidance for Wind Energy Projects.
<http://www.fws.gov/midwest/endangered/mammals/inba/WindEnergyGuidance.html>.

U.S. Fish and Wildlife Service (USFWS). 2014b. Designation of Critical Habitat for *Leavenworthia exigua* var. *laciiniata* (Kentucky Glade Cress). May 6, 2014, *Federal Register* 79(87): 25689-25707.

U.S. Fish and Wildlife Service (USFWS). 2015. Endangered and Threatened Wildlife and Plants;

Threatened Species Status for the Northern Long-Eared Bat With 4(d) Rule; Final Rule and Interim Rule. Federal Register 80(63): 17974–18033.

U.S. Fish and Wildlife Service (Service). 2016. Programmatic Biological Opinion on Final 4(d) Rule for the Northern Long-Eared Bat and Activities from Take Prohibitions. Midwest Region. 103 pp.

U.S. Fish and Wildlife Service (Service). 2022. Kentucky Ecological Field Office, unpublished data.

U.S. Fish and Wildlife Service (Service). 2023. Biological Opinion: Effects to the Northern Long-Eared Bat from Planned and Ongoing Activities Being Implemented in the Eastern and Southern Regions of the U.S. Forest Service. 55 pp.

U.S. Fish and Wildlife Service (USFWS). 2024. Range-wide Indiana Bat and Northern Long-eared Bat Survey Guidelines. U.S. Fish and Wildlife Service, Region 3, Bloomington, MN. 95 pp.

U.S. Fish and Wildlife Service (Service). 2024b. Range-Wide Indiana Bat & Northern Long-Eared Bat Survey Guidelines. 15 pp and appendices. Available at: <https://www.fws.gov/media/range-wide-indiana-bat-and-northern-long-eared-bat-survey-guidelines> (Accessed July 3, 2024).

U.S. Fish and Wildlife Service (Service). 2024c. Kentucky Ecological Field Office, unpublished data

U.S. Forest Service (USFS). 2005. A snapshot of the northeastern forests. U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry. Newton Square, PA. NA-IN-01-06. 23 pp. Available at: http://na.fs.fed.us/pubs/misc/snap_shot/ss.pdf. (Accessed: December 13, 2007).

U.S. Forest Service (USFS). 2006. NCRS—the changing Midwest assessment. U.S. Department of Agriculture, Forest Service, North Central Research Station, St. Paul, MN. Available at: <http://www.ncrs.fs.fed.us/4153/deltawest/landcover>. (Accessed: January 25, 2006).

U.S. Forest Service (USFS). 2012. 2010 Resources Planning Act Assessment. Gen. Tech. Rep. WO-87. Washington, D.C., 198pp.

Van zyll de Jong, C.G. 1985. Handbook of Canadian mammals. National Museums of Canada, Ottawa, Canada. 212pp.

Voigt, C.C., K. Schneeberger, S.L. Voigt-Heucke, and D. Lewanzik. 2011. Rain increases the energy cost of bat flight. Biological Letters 7:793–795. doi:10.1098/rsbl.2011.0313

Warnecke, L., J.M. Turner, T.K. Bollinger, J.M. Lorch, V. Misra, P.M. Cryan, G. Wibbelt, D.S. Blehert, and C.K.R. Willis. 2012. Inoculation of bats with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. Proceedings of the National Academy of Sciences 109:6999–7003.

Watrous, K.S., T.M. Donovan, R.M. Mickey, S.R. Darling, A.C. Hicks, and S.L. von Oettingen. 2006. Predicting minimum habitat characteristics for the Indiana bat in the Champlain Valley. *Journal of Wildlife Management* 70(5):1228-1237.

Wear, D.N., and J.G. Greis (eds.). 2002. Southern forest resource assessment. U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, NC. General Technical Report SRS-53. 635 pp.

Webb, D.R., and J.R. King. 1984. Effects of wetting on insulation of bird and mammal coats. *Journal of Thermal Biology* 9(3):189–191.

Whitaker, J.O., and L.J. Rissler. 1992. Seasonal activity of bats at Copperhead Cave. *Proceedings of the Indiana Academy of Science*, 101:127–134.

Whitaker, J.O., and W.J. Hamilton. 1998. Order Chiroptera: Bats. Chapter 3: pp.89–102 in *Mammals of the eastern United States*, Third Edition, Comstock Publishing Associates, a Division of Cornell University Press, Ithaca, New York, 608pp.

Whitaker, J.O., Jr. and V. Brack, Jr. 2002. Distribution and summer ecology in Indiana. Pp. 48–54 in A. Kurta and J. Kennedy (eds.), *The Indiana bat: biology and management of an endangered species*. Bat Conservation International, Austin, TX.

Whitaker, J. O., D. W. Sparks, and V. Brack. 2006. Use of artificial roost structures by bats at the Indianapolis International Airport. *Environmental Management* 38: 28–36.

Whitaker, J.O., and R.E. Mumford. 2009. Northern Myotis. pp. 207–214. In *Mammals of Indiana*. Indiana University Press, Bloomington, Indiana. 688pp.

Whitby, M., B.J. Udell, A.M. Wiens, T. Cheng, W. Frick, B.E. Reichert, and J.D. Reichard. 2022. Summer mobile acoustic transect analysis for little brown, northern long-eared, and tricolored bat species status assessment. Chapter C in Straw, B.R., J. A. Martin, J.D. Reichard, and B.E. Reichert, editors. *Analytical Assessments in Support of the U.S. Fish and Wildlife Service 3-Bat Species Status Assessment*. Cooperator report prepared in cooperation with the U.S. Geological Survey, United States Fish and Wildlife Service and Bat Conservation International. <https://doi.org/10.7944/P9B4RWEU>.

White, J.W., P. Freeman, and C.A. Lemen. 2017. Habitat selection by the northern long-eared myotis (*Myotis septentrionalis*) in the Midwestern United States: Life in a shredded farmscape. *Transactions of the Nebraska Academy of Sciences and Affiliated Societies* 37: 1–10.

White, J.A., P.W. Freeman, H.W. Otto, and C.A. Lemen. 2020. Winter use of a rock crevice by northern long-eared myotis (*Myotis septentrionalis*) in Nebraska. *Western North American Naturalist*: Vol. 80: No. 1, Article 14.

Wiens, A.M., J. Szymanski, B.J. Udell, and W.E. Thogmartin. 2022. Winter colony count data assessment and future scenarios for the little brown, northern long-eared, and tricolored

bat species status assessment. Chapter E in Straw, B.R, J. A. Martin, J.D. Reichard, and B.E. Reichert, editors. Analytical Assessments in Support of the U.S. Fish and Wildlife Service 3-Bat Species Status Assessment. Cooperator report prepared in cooperation with the U.S. Geological Survey, United States Fish and Wildlife Service and Bat Conservation International. <https://doi.org/10.7944/P9B4RWEU>.

Willis, C.K.R, A.K. Menzies, J.G. Boyles, and M.S. Wojciechowski. 2012. Evaporative water loss is a plausible explanation for mortality of bats from white-nose syndrome. *Integrated and Comparative Biology* 51:364–373.

Womack, K. M., S. K. Amelon, and F. R. Thompson. 2013. Summer home range size of female Indiana bats (*Myotis sodalis*) in Missouri, USA. *Acta Chiropterologica* 15(2): 423–429

Personal Communications:

Brack and Sparks. 2011. Personal Communication.

Copperhead Environmental Consulting, Inc. 2014. Migration and Summer Maternity Colony Range Map.

Roby, Piper. 2024. Copperhead Consulting, Inc. Personal Communication.

Whitaker, John O. 1986. Personal Communication.