



Trinity River Mainstem Redd Survey Update 11/6/2015

The [Arcata Fish and Wildlife Office](#) along with the [California Department of Fish and Wildlife](#), [Yurok Tribal Fisheries Program](#), [Hoopa Valley Tribal Fisheries Department](#), and [Shasta-Trinity National Forest](#) survey the mainstem Trinity River to recover salmonid carcasses and count and map Chinook Salmon (*Oncorhynchus tshawytscha*) redds. The distribution of Chinook Salmon spawning is expected to change in response to rehabilitation efforts of the [Trinity River Restoration Program](#) and to the relative run size of natural vs. [Trinity River Hatchery](#) origin fish. Spawning surveys are attempted weekly between [Lewiston Dam](#) and the North Fork Trinity River (reaches 1 through 7) and every other week on the survey reaches downstream of the North Fork Trinity River (Figure 1, Table 1).

This update provides preliminary data on 2015 redd observations encountered and entered into our database as of 11/06/2015 (Table 1). Figure 2 compares preliminary 2015 cumulative redd counts for Reaches 1-10 with the previous 12 years (2002-2014). At this point, redds constructed by Chinook and Coho salmon are not differentiated. After the end of the season, redds are apportioned by association with hatchery or natural origin Chinook Salmon using carcass data.




Figure 1. Mainstem Trinity River salmon spawn survey area for this week's update.





Table 1. Preliminary count of redds by reach for the fall 2015 mainstem Trinity River redd survey. Reach 8 (Pigeon Point Run) and Reach 11 (Burnt Ranch Gorge) are not surveyed due to abundant whitewater.

Week start	Reach														Total	
	01	02	03	04	05	06	07	08	09	10	11	12	13	14		
08/31	0															0
09/07	0	0	0	NS	NS	0	0									0
09/14	0	0	0	0	0	0	0		1	0						1
09/21	12	24	9	9	25	19	15									113
09/28	13	34	13	30	28	37	39		24	13						231
10/05	19	42	67	34	49	30	39					5	0	0		285
10/12	10	45	19	15	32	43	30		115	64						373
10/19	26	24	31	11	14	44	28					24	7	1		210
10/26	43	27	6	5	28	24	25		74	61						293
11/02	43	17	2	0	10	6	4					52	37	15		186
11/09																0
11/16																0
11/23																0
11/30																0
12/07																0
12/14																0
Total	166	213	147	104	186	203	180		214	138		81	44	16		1,692

These data are preliminary and subject to revision.

“NS” = no survey

* = partial count due to technical difficulties




Figure 2. Fall 2015 cumulative Trinity River mainstem redd counts, Reaches 1 to 10 (Reach 8 excluded), as of 11/06/2015 compared to 2014 (dashed line) and the range of counts from 2002-2014 (blue polygon).

To access the latest version of this report on-line, see <http://www.fws.gov/arcata/fisheries/projectUpdates.html> and click on “Trinity River Redd survey update”

We annually track spawning success rates (i.e., pre-spawn mortality versus partially or fully spawned) of female salmon throughout the Trinity River spawning survey season. Figure 3 shows the cumulative percent of female salmon carcasses found pre-spawn compared to recent years. Pre-spawn mortality data is reported only when 10 female salmon carcasses are recovered. As of 11/06 there has been 3 female salmon that died before spawning, out of a total of 356 female carcasses.




Figure 3. Comparison of fall 2015 cumulative observed pre-spawn mortality as of 11/06/2015 compared to that observed recent years 2009-2014. Plots begin for each year when at least 10 female salmon carcasses have been encountered.

A Keyhole Markup Language file has been created to view redd distribution data with Google Earth. This data is created directly from the data collected by our field crews. Each point is attributed with the date that the redd was initially marked (i.e. “10/2” = October, 2nd).

Trinity Redds 2015 11 06

If Google Earth fails to launch automatically, the file can be used by first saving it to your computer then launching the Google Earth application. Select the menu option **File, Open**, locate the downloaded file (“Trinity_Redds_2015_11_06.kml”); and click the **Open** button.

Note: Background imagery in Google Earth is sometimes not projected with perfect accuracy. Redd locations may appear to be mapped on dry ground in those cases when viewed with Google Earth, but they are indeed in the water! Likewise, the river has changed since the most recently available Google Earth imagery.




Image of the Week: Fisheries technician, Spike (Hoopa Valley Tribe), holds a spawned female Chinook Salmon. Note the tablet computer and GPS unit. We use these electronics to record the location and characteristics of salmon carcasses and redds you see in this report.

For more information regarding mainstem Trinity River redd surveys, please contact Steve Gough (707-825-5197; steve_gough@fws.gov) or Derek Rupert (570-419-2823; derek_rupert@fws.gov).

To access the latest version of this report on-line, see <http://www.fws.gov/arcata/fisheries/projectUpdates.html> and click on “Trinity River Redd survey update”