

U.S. Fish & Wildlife Service

Arcata Fisheries Data Series Report DS 2006-01

Mainstem Klamath River Fall Chinook Salmon Spawning Survey 2001

Schyuler Grove, Patrick Mcneil, and Mark Magneson



U.S. Fish and Wildlife Service
Arcata Fish and Wildlife Office
1655 Heindon Road
Arcata, CA 95521
(707) 882-7201



June 2006



Funding for this study was provided by the Klamath River Basin Conservation Area Restoration Program under Project Number 05-FP-13 and the Klamath River Habitat Assessment Study administered by the Arcata Fish and Wildlife Office.

Disclaimer: The mention of trade names or commercial products in this report does not constitute endorsement or recommendation for use by the Federal government.

The Arcata Fish and Wildlife Office Fisheries Program reports its study findings through two publication series. The **Arcata Fisheries Data Series** was established to provide timely dissemination of data to local managers and for inclusion in agency databases. The **Arcata Fisheries Technical Reports** publishes scientific findings from single and multi-year studies that have undergone more extensive peer review and statistical testing. Additionally, some study results are published in a variety of professional fisheries journals.

Key words: Chinook, Klamath, redd, spawning survey.

The correct citation for this report is:

Grove S., P. McNeil, and M. Magneson 2006. Mainstem Klamath River fall Chinook salmon spawning survey 2001. U. S. Fish and Wildlife Service, Arcata Fish and Wildlife Office, Arcata Fisheries Data Series Report Number 2006-01, Arcata, California.

Table of Contents

	page
List of Tables	iv
List of Figures	iv
Introduction.....	1
Methods.....	3
Survey Procedures	3
Rafting Equipment.....	5
Individual Reaches.....	5
Reach 1: IGD river access (rkm 309.8) to Deliverance Camp river access (rkm 287.5).....	5
Reach 2: Ash Creek river access (rkm 285.7) to Beaver Creek riffle river access (rkm 261.9).....	5
Reach 3: Beaver Creek riffle river access to Blue Heron river access (rkm 234.3).....	5
Reach 4: Blue Heron river access to Seiad Bar river access (rkm 213.6).....	5
Reach 5: Seiad Bar river access to China Point river access (rkm 192.4).....	5
Reach 6: China Point river access to the Indian Creek confluence (rkm 173.8).....	6
Redd Densities	6
Collection of Redd Data.....	6
Water Temperature.....	6
Discharge	7
Water Clarity.....	7
Adult and Grilse Expansion.....	7
Results and Discussion	7
Individual Reaches.....	7
Reach 1: IGD to Ash Creek river access.....	7
Reach 2: Ash Creek river access to Beaver Creek riffle river access.....	7
Reach 3: Beaver Creek riffle river access to Blue Heron river access	9
Reach 4: Blue Heron river access to Seiad Bar river access.....	9
Reach 5: Seiad Bar river access to China Point river access.....	9
Reach 6: China Point river access to Indian Creek.....	9
All Reaches	15
Redd Densities	15
Water Temperature	15
Discharge	15
Water Clarity.....	15
Suction Dredge Mining.....	15
Adult Grilse Expansion.....	16
Summary	17
Acknowledgements.....	18
Literature Cited.....	18

List of Tables

	page
Table 1. Weekly summary and percent frequency of mainstem Klamath River fall Chinook salmon redd counts for Reaches 1 to 6, 1993 to 2001 (NS = No Survey, R1 = Iron Gate Dam to Ash Creek river access, R2 = R1 to Beaver Creek riffle river access, R3 = R2 to Blue Heron river access, R4 = R3 to Seiad Bar, R5 = R4 to China Point river access, R6 = R5 to Indian Creek river access).	8
Table 2. Fall Chinook salmon redds per river kilometer (rkm) for 10 river kilometer (approximate) sections on the mainstem Klamath River, 1993 to 2001 (“Ns” = No survey).	16
Table 3. Natural fall Chinook salmon spawning escapement adult and grilse expansion, Klamath River, 2001 (CDFG 2002).	17

List of Figures

	page
Figure 1. Overview map of the Klamath River Basin accessible to fall Chinook salmon.	2
Figure 2. Mainstem Klamath River Chinook salmon spawning survey project location including individual study reaches (1 to 6).	4
Figure 3. Mainstem Klamath River fall Chinook salmon redd density (redds/rkm) by reach, 2001.	10
Figure 4. Yearly USFWS Chinook salmon redd counts by reach, 1993 to 2001.	10
Figure 5. Redd distribution map for 2001, mainstem Klamath River, Iron Gate Dam to Lumgrey Creek.	11
Figure 6. Redd distribution map for 2001, mainstem Klamath River, Vesa Creek to Scott River.	12
Figure 7. Redd distribution map for 2001, mainstem Klamath River, Tom Martin Creek to Portuguese Creek.	13
Figure 8. Redd distribution map for 2001, mainstem Klamath River, Fort Goff Creek to Indian Creek.	14
Figure 9. Water temperatures (°C) at rkm 300.4 and discharge (ft ³ /s) from Iron Gate Dam (October 15 to December 14, 2001).	17

Mainstem Klamath River Fall Chinook Salmon Spawning Survey 2001

Schyuler Grove, Patrick McNeil, and Mark Magnuson

U. S. Fish and Wildlife Service, Arcata Fish and Wildlife Office
1655 Heindon Road; Arcata, California
Mark_Magnuson@fws.gov

Abstract. Results of the ninth annual US Fish and Wildlife Service fall Chinook salmon *Oncorhynchus tshawytscha* spawning survey conducted on the mainstem Klamath River are summarized. The survey was conducted for an eight week period (October 15 to December 14, 2001), covering 135.9 river kilometers (rkm; 84.4 river miles) between Iron Gate Dam and the confluence of Indian Creek at Happy Camp. A total of 2,172 redds were observed during the 2001 survey, which was 1.4 times higher than the annual average from the preceding eight years ($\bar{x} = 1,539$ redds). Thirty-eight percent ($n = 830$) of redds observed during the 2001 survey were located between Iron Gate Dam and the Shasta River. The highest redd density within 10 km river sections was between Iron Gate Dam and Cape Horn Creek (61 redds/rkm). The lowest redd densities in 2001 were observed between Shinar and China Creek (4.3 redds/rkm). Water clarity between Iron Gate Dam and Indian Creek ranged from 1.2 to 3.0 m, representing some of the highest visibility ever recorded during these surveys. During 2001, one redd was observed on a suction dredge tailing between Iron Gate Dam and Indian Creek.

Introduction

The Klamath River drains approximately 14,000 km² in Oregon and 26,000 km² in California. The majority of the watershed in California is within the boundaries of the Six Rivers, Klamath and Shasta-Trinity National Forests. The Yurok Indian Reservation, comprising about 139 km² in Humboldt and Del Norte counties, borders the lower 68 km of the Klamath River (Figure 1). The most important anadromous salmonid spawning tributaries in the basin include the Trinity River (the largest tributary in the basin) draining approximately 7,690 km², and the Shasta, Scott and Salmon rivers, each draining about 2,070 km². Iron Gate Dam (IGD), located at river kilometer (rkm) 310.3 on the Klamath River and Lewiston Dam located at rkm 249 on the Trinity River, are barriers to upstream passage of anadromous salmonids. Iron Gate Hatchery (IGH) and Trinity River Hatchery, located near the base of each dam, were constructed to mitigate for losses in natural fish production that resulted from dam construction (USFWS 1991).




Figure 1. Overview map of the Klamath River Basin accessible to fall Chinook salmon.

The Klamath River Basin historically supported large runs of Chinook *Oncorhynchus tshawytscha* and coho salmon *O. kisutch*, and steelhead trout *O. mykiss*, which contribute to economically and culturally important subsistence, sport and commercial fisheries (Leidy and Leidy 1984). Generations of Indians have fished in the drainage, with catches of salmon, steelhead, lamprey, and sturgeon historically providing the mainstay of the Indian economy in the area. Sport fishing for salmon and steelhead in the drainage may exceed 200,000 angler days annually. During the 1980's, the Klamath River stocks accounted for up to 30% of commercial Chinook salmon landings in northern California and Southern Oregon, averaging about 450,000 Chinook salmon per year (PFMC 1988).

Concern about the depletion of anadromous salmonid resources and associated habitat in the basin emerged around the turn of the century and has accelerated in recent decades, coincident with expanded logging and fishing operations, construction of dams, road construction, agricultural use, and other development. As in other river systems of the Pacific Northwest, Chinook salmon of the Klamath River Basin have been adversely impacted by habitat degradation and over-exploitation, as reflected by declining runs in recent decades (USFWS 1991).

On October 27, 1986 the US Congress enacted PL 99-552, the Klamath River Fish and Wildlife Restoration Act. This Act authorized the Secretary of the Interior to restore anadromous fish populations to optimum levels in the Klamath River Basin through the creation of the Klamath River Basin Conservation Area Restoration Program, functioning under the guidance of the Klamath River Fishery Management Council (USFWS 1991). The US Fish and Wildlife Service (USFWS) was funded through the Klamath River Basin Conservation Area Restoration Program to identify fall Chinook salmon spawning areas and collect information necessary to estimate the natural fall Chinook salmon spawning escapement on the mainstem Klamath River between IGD and the confluence of Indian Creek (rkm 173.8). This information is critical for monitoring long-term trends in escapement and for developing harvest management alternatives.

Methods

Survey Procedures

The 2001 Arcata Fish and Wildlife Office (AFWO) mainstem Klamath River redd survey protocol consists of six mainstem reaches (Figure 2) covering 135.9 rkm (84.4 river miles). The six reaches were surveyed weekly from IGD to the confluence of Indian Creek, unless adverse weather limited the visibility of the water to less than 1 m in depth. Reaches were surveyed between October 15 and December 14. Two individual crews were organized for conducting the survey. Each crew covered the same survey reaches each week to remain familiar with each survey reach and prior redd locations.




Figure 2. Mainstem Klamath River Chinook salmon spawning survey project location including individual study reaches (1 to 6).

Rafting Equipment.

Two 4.3 m (14 foot) inflatable Wing Catarafts were used for direct observation of fall Chinook salmon redds. These rafts are commonly used for use in white water rescue due to their stability and maneuverability in fast water. Rafts were stacked on a flatbed trailer and deployed at selected access sites along the study area. Each raft was equipped with a rowing frame, a modified observation platform, and anchoring system. Two personnel operated each raft (rower and observer).

Individual Reaches

Reach 1: IGD river access (rkm 309.8) to Deliverance Camp river access (rkm 287.5). Reach 1 is approximately 22.3 rkm (13.8 miles) in length (Figure 2) and was surveyed over a period of two days (Monday and Tuesday). Due to the high percentage of redds in Reach 1, two cataraft crews surveyed the entire reach from IGD to Deliverance Camp river access. One crew surveyed the south bank to mid-channel while the second crew surveyed from the north bank to mid-channel. A section of Reach 1 from the Interstate 5 (I-5) bridge (rkm 292.7) to the Deliverance Camp river access was completed in approximately two hours on the first day (Monday) of the survey week. On the second survey day of each week (Tuesday), the survey resumed from the IGD river access to the I-5 Bridge. This reach was completed in 8 to 10 hrs; depending on flows. The section from Deliverance Camp river access to Ash Creek river access (rkm 285.7) was not surveyed because past surveys have revealed lack of available spawning habitat.

Reach 2: Ash Creek river access (rkm 285.7) to Beaver Creek riffle river access (rkm 261.9). Reach 2 is approximately 23.8 rkm (14.7 miles) in length (Figure 2). The Ash Creek river access is located on the south bank of the Klamath River just downstream of Ash Creek and accessed from Klamath River road. The Beaver Creek riffle river access is located along the north bank on a large gravel bar downstream from Beaver Creek Bridge, just off Highway 96. One crew surveyed this reach, on the third survey day (Wednesday) of each week in about 7 hrs.

Reach 3: Beaver Creek riffle river access to Blue Heron river access (rkm 234.3). Reach 3 is approximately 27.6 rkm (17.1 miles) in length (Figure 2). The Blue Heron river access is on the south bank approximately 2 rkm upstream from the Scott River confluence, just off Highway 96. This reach was surveyed by one crew on the third survey day (Wednesday) of each week, taking about 8 hrs to complete.

Reach 4: Blue Heron river access to Seiad Bar river access (rkm 213.6). Reach 4 is approximately 20.8 rkm (12.9 miles) in length (Figure 2). The Seiad Bar river access is located along the north bank of the Klamath River on Walley Johnson's property. Access to the river is reached by taking Diamond J road from Highway 96 and then turning right onto an unmarked road located near the California Department of Transportation compound. Reach boundaries remain the same as in previous years. This reach was surveyed on the fourth day (Thursday) of each survey week by one crew, taking about 8 hrs to complete.

Reach 5: Seiad Bar river access to China Point river access (rkm 192.4). Reach 5 is approximately 21.2 rkm (13.2 miles) in length (Figure 2). China Point river access is

located along the north bank of the river, at the US Forest Service (USFS) river access just off Highway 96. One crew surveyed this reach, on the fourth day (Thursday) of each survey week, taking about 7 hrs to complete.

Reach 6: China Point river access to the Indian Creek confluence (rkm 173.8). Reach 6 is approximately 18.6 rkm (11.6 miles) in length (Figure 2). Due to the high concentration of redds in this area, this reach was split at Gordons Ferry river access (rkm 185). China Point river access to Gordons Ferry river access is approximately 7.4 rkm (4.6 miles). Gordons Ferry river access to Indian Creek confluence is approximately 11.3 rkm (7 miles). The Gordons Ferry river access is located on the north bank of the river just off Highway 96. This reach was split into two sections and covered by two crews on the fifth day (Friday) of each survey week, with each section taking about 5 hrs to complete.

Redd Densities

To describe the frequency of redd occurrence by location, the mainstem Klamath River between IGD and the confluence of Indian Creek was segregated into 14, approximately 10-rkm long river sections.

Collection of Redd Data.

A length of plastic flagging was attached to vegetation on the riverbank nearest to the position of each redd observation. Different colored flagging was used each week to ensure that redds were not double counted during the subsequent surveys. Date, number of redds per site, location in channel, and redd site number were recorded on flags. Redd sites were also recorded on river maps and on data forms. River reach, GPS mark number, tally number, location in channel (left or right bank, middle, side channel, split channel and pool tail-out), distance from bank, defended area (distance between redds in a cluster), number of adults and grilse, flag location, water temperature, weather conditions, river reach, and age of redd. Estimated redd ages were recorded as Redd Age Code 1, 2, or 3 depending on appearance. Fresh redds with bright substrate, little or no periphyton, and well-developed mounds were classified as Redd Age Code 1. Redds two to four weeks old with slightly flattened mounds and dulled substrate due to periphyton growth were classified as Redd Age Code 2. Redds older than four weeks, which were identifiable only by pit and/or mound presence and typically could not be distinguished from surrounding substrate by brightness differences, were classified as Redd Age Code 3. Only completed redds (which included both a pit and mound) were included in the daily counts. Test redds and small (<1.0 m²) redds were omitted.

Water Temperature.

Water temperature was recorded using an Optic Stowaway Tidbit on an hourly basis throughout the survey period at the downstream end of R-Ranch (5 rkm downstream of IGD). Hourly data were used to calculate mean daily water temperatures.

Discharge

Mean daily river flow was obtained from the US Geological Survey gauging station (Number 11516530; Latitude 41° 55' 41", Longitude 122°26'35" NAD 27; <http://waterdata.usgs.gov/ca/nwis/current/?type=flow>), located in the Klamath River just downstream of IGD. Mean daily river flow was recorded in ft³/s.

Water Clarity

Water clarity was measured each survey week by lowering a 20-cm diameter Secchi disc vertically into the water column. The disc was lowered until the black and white pattern was not discernable. The disc was then raised until the pattern became visible, and this depth was recorded in meters.

Adult and Grilse Expansion

The total number of redds counted in this survey was used by California Department of Fish and Game (CDFG) to estimate adult and grilse (two year old) spawning abundance (CDFG 2002). Adult numbers were calculated by multiplying the total redd count by two. This estimate assumes one male and one female salmon per redd. The adult and grilse age breaks for the mainstem Klamath River from IGD to Indian Creek were derived from scales and fork length data collected during carcass surveys in Reach 1 (KRTAT 2002).

Results and Discussion

A total of 2,172 Chinook salmon redds were counted between IGD and the confluence of Indian Creek in 2001, which was 1.4 times higher than the annual average (1,539 redds) from the preceding eight years (Table 1). The highest number of Chinook salmon redds/rkm was in Reach 1 (Figure 3), consistent with the findings of previous survey years. Redd counts from 2001 were the second highest recorded since 1993 (Figure 4). Based on field maps and notes from the survey, locations of redds for all reaches surveyed were mapped, as shown in Figures 5 to 8.

Individual Reaches

Reach 1: IGD to Ash Creek river access. A total of 830 redds were observed in this reach during the 2001 survey (Table 1; Figure 5). The 830 redds observed represent 38% of the total redd count for 2001. Redd density in this reach was 37.3 redds/rkm (Figure 3). Based on weekly redd counts, peak spawning (n = 323) occurred during the third week (October 29 to November 2) of this survey (Table 1).

Reach 2: Ash Creek river access to Beaver Creek riffle river access A total of 269 redds were observed in this reach during the 2001 survey (Table 1; Figures 5 and 6). The 269 redds observed represents 12% of the total redd count for 2001. Redd density in this reach was 11.3 redds/rkm (Figure 3). Peak spawning (n = 102) occurred during the second (October 22 to October 26) week of this survey (Table 1). The 269 redds counted in this reach represent the highest count since the project started in 1993 (Table 1).

Table 1. Weekly summary and percent frequency of mainstem Klamath River fall Chinook salmon redd counts for Reaches 1 to 6, 1993 to 2001 (NS = No Survey, R1 = Iron Gate Dam to Ash Creek river access, R2 = R1 to Beaver Creek riffle river access, R3 = R2 to Blue Heron river access, R4 = R3 to Seiad Bar, R5 = R4 to China Point river access, R6 = R5 to Indian Creek river access).

Year	Week	Reach						Total
		R1	R2	R3	R4	R5	R6	
1993	Oct 25 to 29	15	13	30	18	16	81	173
	Nov 1 to 5	67	24	4	1	15	5	116
	Nov 8 to 12	5	1	18	7	0	1	32
	Nov 15 to 18	0	0	4	5	0	0	9
	Total	87	38	56	31	31	87	330
	% Frequency	26	12	17	9	9	26	
1994	Oct 17 to 21	89	28	48	Ns	Ns	98	263
	Oct 24 to 28	278	59	77	113	98	124	749
	Oct 31 to Nov 4	375	20	46	42	16	33	532
	Nov 7 to 11	86	Ns	Ns	Ns	Ns	Ns	86
	Nov 14 to 18	3	2	7	4	5	5	26
	Total	831	109	178	159	119	260	1,656
% Frequency	50	7	11	10	7	16		
1995	Oct 16 to 20	138	12	70	26	30	139	415
	Oct 23 to 27	598	82	199	94	91	169	1,233
	Oct 30 to Nov 3	727	58	78	35	57	112	1,067
	Nov 6 to 10	277	26	49	13	25	50	440
	Nov 27 to Dec 1	39	9	14	4	12	3	81
	Total	1,779	187	410	172	215	473	3,236
% Frequency	55	6	13	5	7	15		
1996	Oct 21 to 25	290	31	96	10	118	39	584
	Oct 28 to Nov 1	291	29	25	22	42	92	501
	Nov 4 to 8	83	4	24	8	33	59	211
	Nov 11 to 15	40	0	6	0	7	23	76
	Total	704	64	151	40	200	213	1,372
	% Frequency	51	5	11	3	15	16	
1997	Oct 16	272	Ns	Ns	Ns	Ns	Ns	272
	Oct 20 to 24	252	37	69	89	29	136	612
	Oct 27 to 31	424	18	76	52	22	76	668
	Nov 3 to 7	70	7	13	16	8	27	141
	Nov 10 to 14	2	14	4	5	3	18	46
	Total	1,020	76	162	162	62	257	1,739
% Frequency	59	4	9	9	4	15		
1998	Oct 14 to 15	89	Ns	Ns	Ns	Ns	Ns	89
	Oct 19 to 23	180	45	67	15	20	45	372
	Oct 26 to 30	368	11	12	14	7	39	451
	Nov 2 to 6	226	22	33	10	9	28	328
	Nov 9 to 12	135	3	11	3	2	2	156
	Nov 15 to 19	12	1	3	0	1	2	19
Total	1,010	82	126	42	39	116	1,415	
% Frequency	71	6	9	3	3	8		
1999	Oct 13 to 15	98	3	Ns	Ns	Ns	Ns	101
	Oct 18 to 22	200	27	31	17	23	39	337
	Oct 25 to 27	304	23	20	Ns	Ns	Ns	347
	Nov 1 to 5	83	12	9	8	8	19	139
	Nov 8 to 12	37	2	2	1	5	11	58
	Nov 15 to 19	1	2	0	2	2	0	7
Total	723	69	62	28	38	69	989	
% Frequency	73	7	6	3	4	7		

Table 1. (Continued). Weekly summary and percent frequency of mainstem Klamath River fall Chinook salmon redd counts for Reaches 1 to 6, 1993 to 2001 (NS = No Survey, R1 = Iron Gate Dam to Ash Creek river access, R2 = R1 to Beaver Creek riffle river access, R3 = R2 to Blue Heron river access, R4 = R3 to Seiad Bar, R5 = R4 to China Point river access, R6 = R5 to Indian Creek river access).

Year	Week	Reach						Total
		R1	R2	R3	R4	R5	R6	
2000	Oct 16 to 20	327	92	69	25	10	19	542
	Oct 23 to 27	146	62	34	52	10	53	357
	Oct 30 to Nov 3	254	42	69	54	20	86	525
	Nov 6 to 10	57	12	15	21	2	16	123
	Nov 13 to 17	4	0	9	12	0	6	30
	Nov 20 to 22	1	Ns	Ns	Ns	Ns	Ns	1
	Total	788	208	196	164	42	180	1,578
% Frequency	50	13	12	10	3	11		
2001	Oct 15 to 19	92	24	28	21	2	23	190
	Oct 22 to 26	168	102	128	59	40	82	579
	Oct 29 to Nov 2	323	97	170	102	55	139	886
	Nov 5 to 9	155	10	40	12	31	29	277
	Nov 12 to 16	75	31	49	22	9	Ns	186
	Nov 26 to 30	17	Ns	Ns	Ns	Ns	Ns	17
	Dec 3 to 7	Ns	Ns	12	Ns	Ns	5	17
	Dec 10 to 14	Ns	5	8	4	3	Ns	20
Total	830	269	435	220	140	278	2,172	
% Frequency	38	12	20	10	6	13		

Reach 3: Beaver Creek riffle river access to Blue Heron river access. A total of 435 redds were counted in this reach during the 2001 survey (Table 1; Figure 6); comprising 20 % of the total redd count for the season. Redd density was 15.8 redds/rkm (Figure 3). Peak spawning (n = 170) in Reach 3 occurred during the third week (October 29 to November 2) of this survey (Table 1). The 435 redds observed during the survey is the highest count for this reach since the project started in 1993.

Reach 4: Blue Heron river access to Seiad Bar river access. A total of 220 redds were counted in this reach during the 2001 survey (Table 1; Figures 6 and 7); representing 10 % of the total redd count for the season. Redd density was 10.6 redds/rkm (Figure 3). Peak spawning (n = 102) in Reach 4 occurred during the third week (October 29 to November 2) of this survey (Table 1). The 220 redds observed during the survey is the highest count for this reach since the project started in 1993.

Reach 5: Seiad Bar river access to China Point river access. A total of 140 redds were counted in this reach during the 2001 survey (Table 1; Figures 7 and 8), accounting for 6 % of the season total. Redd density was 6.6 redds/rkm (Figure 3). Peak spawning (n = 55) occurred during the third week (October 29 to November 2) of this survey (Table 1). Reach 5 had the lowest redd count (n = 140) of any reach sampled in 2001 (Table 1).

Reach 6: China Point river access to Indian Creek. A total of 278 redds were counted in Reach 6 (Table 1; Figure 8), representing 13 % of the total redd count for 2001. The estimated redd density was 14.9 redds/rkm (Figure 3), the second highest for this reach




Figure 3. Mainstem Klamath River fall Chinook salmon redd density (redds/rkm) by reach, 2001.




Figure 4. Yearly USFWS Chinook salmon redd counts by reach, 1993 to 2001.




Figure 6. Redd distribution map for 2001, mainstem Klamath River, Vesa Creek to Scott River.




Figure 7. Redd distribution map for 2001, mainstem Klamath River, Tom Martin Creek to Portuguese Creek.




Figure 8. Redd distribution map for 2001, mainstem Klamath River, Fort Goff Creek to Indian Creek.

since 1993. Peak spawning (n = 139) in Reach 6 occurred during the third week (October 29 to November 2) of this survey (Table 1).

All Reaches

Spawning was observed throughout the mainstem river from IGD to Indian Creek and was consistent with previous survey data in that spatial distribution of redds was relatively lowest in Reach 5 (6.6 redds/rkm), highest in Reach 1 (37.2 redds/rkm), and intermediate for reaches 2, 3, 4 and 6. The highest weekly redd count also occurred during late October, similar to other survey years. Overall, the 2001 redd survey yielded the second highest count during the period of record (1993 to 2001; Figure 4).

Redd Densities

Redd densities in 2001 were highest (61.0 redds/rkm) between IGD and Cape Horn Creek. Redd densities were higher in 2001 than in 2000 for all but two 10-km Sections (Section 1 and 3). In 2001, the lowest redd density (4.3 redds/rkm) was between Shinar and China Creek (Table 2; Figure 9).

Water Temperature

Mean daily water temperatures decreased from 15.5 to 6.0 °C during this survey (October 15 and December 14, 2001). Water temperatures continued to decrease during the surveys, even though IGD flows remained fairly stable 1,300 to 1,410 cfs (Figure 10).

Discharge

Discharges for the 2001 survey ranged from 1,300 to 1,410 cfs (Figure 10). Some of the highest mean daily discharges (1,360 to 1,820 cfs) were during the 1999 survey. Some of the lowest occurred during 1994 (906 to 962 cfs).

Water Clarity

Vertical Secchi disc readings ranged from 1.2 to 3.0 m during this 2001 survey, compared to 1.2 to 2.0 m in 2000. The 1.2 m reading was observed on December 7, 2001 in Reach 6. The 3.0 m reading was observed on October 16 in Reach 1. Water clarity during the 2001 survey increased slightly from that experienced in 2001. Visibility generally decreased with higher river discharge, cloud cover, and precipitation.

Suction Dredge Mining

Recreational suction dredge mining was present throughout the survey from the Highway I-5 Bridge to Happy Camp. There was only one redd observed in 2001 on suction dredge tailings. Studies have indicated that redds constructed on dredge tailings are more unstable in high flows than those on naturally deposited substrate (Harvey and Lisle 1999).

Adult Grilse Expansion

The CDFG estimated the natural fall Chinook salmon spawner escapement for the mainstem Klamath River for 2001 at 9,832 adults and 1,016 grilse (Table 3). The adult/grilse estimates are based on male to female ratio and jack percentages observed at upper Klamath River tributary weirs (CDFG 2005). Based on spawning data from mainstem and tributary spawning surveys conducted by AFWO, USFS, CDFG, and Hoopa and Yurok tribes, the CDFG estimated that 132,945 adults spawned in-river or at hatcheries within the Klamath River Basin in 2001.

Table 2. Fall Chinook salmon redds per river kilometer (rkm) for 10 river kilometer (approximate) sections on the mainstem Klamath River, 1993 to 2001 (“Ns” = No survey).

Tributary Reach	Reach Length (rkm)	Year								
		1993	1994	1995	1996	1997	1998	1999	2000	2001
Iron Gate Dam to Cape Horn Creek	9.2	8.7	69.9	137.8	61.0	85.0	84.7	65.4	64.2	61.0
Cape Horn Creek to Shasta River	12.2	0.6	14.9	41.3	12.0	20.2	18.7	9.5	16.1	22.0
Shasta River to Humbug Creek	8.8	1.0	1.8	7.2	1.3	2.2	4.8	3.8	10.5	8.9
Humbug Creek to Vesa Creek	11.4	1.6	3.0	3.2	1.3	2.8	1.9	2.5	4.8	5.6
Vesa Creek to Little Humbug Creek	10.9	1.7	10.5	15.4	6.1	5.3	3.9	2.7	10.9	20.5
Little Humbug Creek to Kohl Creek	9.5	2.7	6.1	16.8	4.7	7.9	4.5	0.9	8.4	16.4
Kohl Creek to Kinsman Creek	11.0	2.4	4.0	14.5	5.3	3.6	5.5	3.0	5.2	13.2
Kinsman Creek to Kuntz Creek	9.8	0.8	6.8	3.9	3.4	1.6	0.6	1.2	3.4	9.3
Kuntz Creek to Walker Creek	10.3	2.0	8.8	12.0	2.3	14.6	3.5	1.8	10.5	15.0
Walker Creek to Portuguese Creek	9.4	2.7	9.0	13.4	8.0	1.9	1.4	2.2	4.1	8.1
Portuguese Creek to Shinar Creek	8.6	0.5	4.1	8.1	5.0	2.9	2.3	1.9	2.4	5.8
Shinar Creek to China Creek	7.2	3.1	10.6	19.9	11.4	4.3	0.8	0.4	0.8	4.3
China Creek to Ottley Gulch	8.3	4.9	14.0	23.4	17.7	13.6	8.1	4.3	14.3	25.9
Ottley Gulch to Indian Creek	9.9	2.4	9.1	18.9	6.7	13.4	4.9	3.1	6.2	6.4




Figure 9. Water temperatures (°C) at rkm 300.4 and discharge (ft³/s) from Iron Gate Dam (October 15 to December 14, 2001).

Table 3. Natural fall Chinook salmon spawning escapement adult and grilse expansion, Klamath River, 2001 (CDFG 2002).

Natural Spawning Area	Natural Spawners		Totals
	Grilse	Adults	
Mainstem Klamath River Iron Gate Dam to Indian Creek	1,016	9,832	10,848
Bogus Creek Basin	648	11,927	12,575
Shasta River Basin	2,641	8,452	11,093
Scott River Basin	744	5,398	6,142
Salmon River Basin	743	2,607	3,350
Misc. Klamath Tributaries upstream of Yurok Reservation	538	2,240	2,778
Yurok Reservation Tributaries	48	488	536
Total Natural Spawners	6,378	40,944	47,322

Summary

The 2001 fall Chinook salmon redd count of 2,172 was the second highest number observed since the initiation of these surveys in 1993. The highest redd count was 3,236

redds observed in 1995. Peak mainstem fall Chinook salmon spawning (41 %) occurred during the third survey week (October 29 to November 2, 2001) in all reaches except for Reach 2. In Reach 2, peak spawning occurred during the second survey week (October 22 to October 26).

Typical water clarity experienced during the 2001 survey was higher (1.2 to 3.0 m) than observed during past survey years.

Acknowledgements

The Arcata Fish and Wildlife Office wishes to acknowledge the Karuk Tribe of California, Department of Natural Resources, for providing support and the following personnel: Ron Reed, Mike Polmateer, and Dave Arwood II. In addition, we are grateful to AFWO employees Tom Shaw, George Guillen, and Nicholas Hetrick for editorial comments. We would also like to thank Wally Johnson for allowing us river access through his property at Seiad.

Literature Cited

- CDFG (California Department of Fish and Game). 2005. Klamath River Basin fall Chinook salmon spawner escapement, in-river harvest and run-size estimates 1978-2005a. CDFG. Available from W. Sinnen, CDFG, 5341 Ericson Way, Arcata, CA 95521
- Harvey, B.C. and T.E. Lisle. 1999. Scour of Chinook salmon Redds on Suction Dredge Tailings. *North American Journal of Fisheries Management* 19:613-617.
- KRTAT (Klamath River Technical Advisory Team). 2002. Klamath River fall Chinook age-specific escapement, river harvest, and run size estimates, 2001 run. Klamath Fishery Management Council, Yreka Fish and Wildlife Office, Yreka, California.
- Leidy, R.A. and G.R. Leidy. 1984. Life stage periodicities of anadromous salmonids in the Klamath River Basin, northwestern California. U.S. Fish and Wildlife Service, Division of Ecological Services, Sacramento, CA. 21 pp.
- PFMC (Pacific Fishery Management Council). 1988. Review of 1988 ocean salmon Fisheries. Portland, Oregon.
- USFWS 1991. Annual Report: Klamath River Fisheries Assessment Program, 1989. Coastal California Fishery Resource Office, Arcata, CA.