

# Warm Springs National Fish Hatchery - Spring Chinook Salmon Program FY 2022 Annual Report

Brook Silver, Todd Gilmore, David Hand, Jesse Rivera, and Jeremy Voeltz

Columbia River Fish and Wildlife Conservation Office 1211 SE Cardinal Court, Suite 100  
Vancouver, WA 98683

May 18, 2023

## Abstract

*In 1966, congress authorized the Warm Springs National Fish Hatchery to stock salmon and trout within the Confederated Tribes of the Warm Springs Reservation of Oregon reservation to increase tribal harvest opportunities. The current focus of the Warm Springs National Fish Hatchery is to produce spring Chinook Salmon for tribal harvest in the Deschutes and Columbia River and for on-reservation distribution to tribal members. The facility is managed as an integrated hatchery program to minimize genetic divergence between Warm Springs River hatchery and wild stocks. The Columbia River Fish and Wildlife Conservation Office conducts monitoring and evaluation of this hatchery program. This report summarizes broodstock need, juvenile production levels, and marking and tagging information for the past ten years. After juvenile release, the detection rates at Bonneville Dam, juvenile survival, adult returns, smolt-to-adult survival rates inferred from coded-wire tag recoveries, and adult age structures are reported. Special studies and recommendations for future studies supported by U.S. Fish and Wildlife Service funds are also discussed.*

For previous Columbia River Fish and Wildlife Program Office reports, please see:  
<https://www.fws.gov/office/columbia-river-fish-and-wildlife-conservation/reports>

**Disclaimer:**

*The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the U.S. Fish and Wildlife Service. The mention of trade names or commercial products in this report does not constitute endorsement or recommendation for use by the federal government.*

**The correct citation for this report is:**

Silver B.P., T. Gilmore, D. Hand, J. Rivera, and J. Voeltz. 2023. Warm Springs National Fish Hatchery - Spring Chinook Salmon Program, 2022 Annual Report. U.S. Fish and Wildlife Service, Columbia River Fish and Wildlife Conservation Office, Vancouver, WA. 57 p.

# Contents

<b>Abstract</b> .....	<b>1</b>
<b>Introduction</b> .....	<b>4</b>
<i>Program Description</i> .....	5
<i>Past Objectives</i> .....	6
<i>Present Objectives</i> .....	7
<b>Hatchery Operations Summary</b> .....	<b>10</b>
<i>On-Station Juvenile Production</i> .....	10
a) Egg-to-Smolt Survival.....	10
b) Juvenile Marking, Tagging, and Release Data.....	10
<i>Off-Station Juvenile Production</i> .....	13
a) PIT Tagging Program.....	13
b) Juvenile Survival.....	14
<i>Adult Returns: Smolt-to-Adult Survival, Detections, Age Structure, and Harvest Data</i> .....	16
a) Adult Returns .....	16
b) Bonneville Dam and Ladder Detections .....	19
c) Age Structure.....	21
d) Hatchery Returns by Stock.....	22
e) External Mark Accuracy by Stock .....	25
f) Adult Harvest.....	27
<b>2022 Run Reconstruction</b> .....	<b>29</b>
<b>2022 Wild Spring Chinook Management</b> .....	<b>31</b>
<i>Wild Adult Return</i> .....	31
<i>Wild Juvenile Release</i> .....	34
<b>2023 Run Forecast</b> .....	<b>36</b>
<i>Warm Springs River Wild stock</i> .....	36
<i>Warm Springs NFH program</i> .....	36
<b>Transfers</b> .....	<b>37</b>
<b>Other Fish counted and passed above Warm Springs NFH</b> .....	<b>45</b>
<b>Past M&amp;E Studies</b> .....	<b>47</b>
<b>Summary and Future Studies</b> .....	<b>49</b>
<i>Future M&amp;E Studies</i> .....	50
<b>Appendix A</b> .....	<b>52</b>
<b>References</b> .....	<b>54</b>



*Figure 1. Aerial photograph of Warm Springs NFH located along the Warm Springs River, within the Warm Springs Reservation of OR. U.S. Fish and Wildlife Service stock photograph.*

## Introduction

Salmon are an integral part of the spiritual and cultural identity of the Confederated Tribes of the Warm Springs Reservation of Oregon (CTWSRO) and are an essential component of their traditional and contemporary diet. Each year, returning salmon allow the transfer of traditional values from generation to generation. It is a tribal priority to meet current and future needs of the resource as well as those of the Tribe. Because the CTWSRO tribal population is growing, the need for salmon is more important than ever.

The tribes, states, and federal government share the responsibility to protect fish habitat and enhance fish runs in all waters. The Treaty of 1855 recognizes tribal sovereignty as the right of the CTWSRO to govern their members and manage their territories and resources. Furthermore, the federal government and its implementing agencies owe an affirmative duty to use their expertise and authority in meaningful consultation with CTWSRO and safeguard natural resources of crucial importance to self-government and prosperity. In 1959, the CTWSRO requested the U.S. Fish and Wildlife Service (Service) investigate the possibilities of salmon and steelhead enhancement on the Warm Springs Reservation. It was determined that operation of a national fish hatchery on the Warm Springs Reservation was pivotal for the enhancement of the anadromous fish runs. On May 31, 1966, Warm Springs National Fish Hatchery (WSNFH) was authorized by Federal Statute 184 to stock salmon and trout within the Warm Springs Reservation to increase tribal harvest opportunities. Since 1978, WSNFH has supplemented fish for harvest in the waters of the Warm Springs Reservation. Production from the hatchery is considered essential for the enhancement of spring Chinook Salmon (*Oncorhynchus tshawytscha*) populations and meeting tribal trust responsibilities.

The CTWSRO has the principal management responsibility for fishery resources on the Warm Springs Reservation. Since 1977 the Service and CTWSRO have worked together to draft hatchery operations and management plans to assure the operation of the hatchery is compatible with and compliments the Tribe's fishery management goals. This cooperative management of the hatchery provides tribal and sport harvest opportunities, enhances anadromous fish runs in Warm Springs Reservation waters, and meets the future needs of the resource and those of the CTWSRO while protecting wild fish populations.

The current focus of the WSNFH is to produce spring Chinook Salmon for tribal harvest in the Deschutes and Columbia River and for on-reservation distribution to tribal members. The facility is managed as an integrated hatchery program. The Service and Tribes have taken this integrated approach to managing the hatchery to not only produce fish, but also minimize genetic divergence between Warm Springs River hatchery and wild stocks, as well as determine what effects hatchery fish have on the ecosystem into which they are released (Olson et al. 2004). The Warm Springs River is one of two rivers in the Deschutes River subbasin that supports natural production of spring Chinook Salmon. Although spring Chinook Salmon are not listed under the Endangered Species Act (ESA), the WSNFH program does cause interactions with listed Mid-Columbia River summer Steelhead (Olson and Spateholts 2001). The safe passage of all wild fish populations, both downstream and upstream of WSNFH, is also an important goal. The hatchery is operated in compliance with the ESA (National Marine Fisheries Service (NMFS) 2007) and consistent with the 2018 - 2027 *United States v. Oregon* Management Agreement (NMFS 2018). The purpose of this report is to summarize programs conducted at the facility over the past ten years and describe additional studies conducted and supported by Service funds.

## Program Description

Warm Springs NFH is located at river kilometer (rkm) 16 of the Warm Springs River, within the Warm Springs Reservation of Oregon, approximately 23 kilometers (km) north of the town of Warm Springs (Fig. 1). The Warm Springs River enters the Deschutes River at rkm 135, which enters the Columbia River 329 kms from the Pacific Ocean. It is upstream of two main-stem dams on the Columbia River, Bonneville (rkm 235) and The Dalles (rkm 308), and downstream of the Pelton/Round Butte (rkm 161) dams on the Deschutes River. The facility is part of the Columbia River Gorge Complex and operated by the Service on land and water leased from the CTWSRO until February 1, 2043 (BIA 1967). The water intake structure and pumps are located at the hatchery site just upstream of a barrier dam across the Warm Springs River, adjacent to the hatchery facility. Prior water intake, water passes through a trash rack and traveling screen. The primary prevention of fish entrainment are the drum screens located in the intake structure behind the trash racks. In addition, a redundant fish bypass located in front of the traveling screens may deposit small fish below the barrier dam.

The hatchery currently has a staff of five full-time Service employees; the hatchery manager, three animal caretakers, and a maintenance mechanic. The Pacific Region Fish Health Program (PRFHP) manages fish health and disease prevention in accordance with Service Fish Health Policy and Implementation Guidelines and IHOT policies (Integrated Hatchery Operations Team (IHOT) 1995; USFWS 1995, 2004) and with protocols of Oregon Department of Fish and Wildlife (ODFW). Fish health personnel promptly manage any health problems to limit mortality and reduce disease transmission.




Figure 2. The Warm Springs NFH is located within the Warm Springs Reservation of Oregon and uses funds from the Service to support its rearing program.

## Past Objectives

Fish production began in 1978 with eggs from wild spring Chinook Salmon and steelhead (*O. mykiss*) captured from the existing natural runs passing the hatchery site. The steelhead program was terminated in 1981 because of disease, growth problems and physical limitations of the facility. To protect wild steelhead, only wild steelhead are passed above WSNFH and all known hatchery origin steelhead are sacrificed and distributed to the CTWSRO.

In 1984, the CTWSRO asserted that separating the hatchery and natural producing fish would best serve the fish and the needs of the tribal people. The CTWSRO proposed a two-stock concept, whereby only wild (unmarked) fish are passed above the hatchery. To this end, 100 percent of fish released from the hatchery are marked with a coded-wire tag (CWT) and an adipose fin-mark (AD) to distinguish them from wild fish. The differential marking of hatchery and wild fish provides consistent long-term data on the life-history patterns and possible changes that may occur within stocks. It also allows for maintenance of the genetic integrity of the naturally producing stock. To maintain the genetic and life-history characteristics of the wild population in the hatchery environment, the hatchery incorporates wild fish into its broodstock when wild returns are greater than 1,000 adults. The minimum escapement goal for naturally produced spring Chinook salmon above the hatchery is 1,000 adults, with a long-term goal of a run of 2,800, similar to runs before the hatchery was constructed (CTWSRO and USFWS 2007).

In 1996, WSNFH installed an automated fish passage system to minimize handling of natural fish and reduce pre-spawn mortality by separating out returning hatchery spring Chinook salmon with CWTs. During the spring Chinook migration period, generally from April 15th to September 30th, the barrier dam directed fish into the adult ladder. Fish swam through a tube in the adult ladder, which triggered a pneumatic gate if a CWT was detected. The goal was to have all CWT hatchery fish shunted to a holding pond and pass non-tagged fish to another catch pond where an underwater video camera monitored them as they swam out through the ladder and upstream of the hatchery. The minimum operating standard for the system was the removal of 95 percent of the fish with CWTs and 95 percent accuracy in counting upstream bound fish. However, the passage system failed to meet the efficiency standards and cost for upgrade was more than \$75,000 (Archibald 2013). In 2014, the system was decommissioned, and hatchery personnel manually sort all fish trapped in the holding ponds.

The release goal for juvenile spring Chinook released from the hatchery has ranged from 400,000 to 1.2 million during 1978 through 1991. From 1992 to the present, the juvenile release goal has been consistently set at 750,000. From brood years (BY) 1979 to 2007, there have been two release strategies, spring and fall. For brood years 1979 to 1992, a graded fall release strategy was employed. Fish were sorted into graded raceways by size and the larger fish were released into the Warm Springs River during the fall. Between brood years 1993 and 2007, a fall volitional release strategy was used at WSNFH. The release strategy was a partial volitional release, where raceways were opened for approximately 4 weeks, between October and November, and fish could volitionally exit the hatchery and enter the Warm Springs River. Based on Passive Integrated Transponder (PIT) tag monitoring of the fall release, anywhere from 10% to 60% of the fish in a raceway would exit during the fall period. Once the fall volitional release period ended, the raceways were closed and the remaining fish would be released the following spring, generally during a spring volitional release period of late-March through April. Fish remaining at the end of April were forced out to make room for the next year's brood. Studies from the 1980s indicated that most smolts released in the spring reached the estuary within three to four weeks, however the behavior of fish released in the fall was not clear (Cates 1992). Scale analysis of adult returns indicated that most fall-released fish that survived to adulthood over-wintered in fresh water before migrating to the ocean the following spring. Follow-up studies from 2000 to 2003 indicated that some fish released in the fall (5% to 36% of the total release each year) quickly migrated downstream and exited the Deschutes River within days of release; however, most of the fish released in the fall overwintered in the Deschutes River (Reagan et al. 2005). The size at release of fish at the hatchery was reduced during the early 2000s, from a size at spring release of 10-15 fish per pound to 20-30 fish per pound. It was thought that the smaller size of fish reared at WSNFH may have contributed to the overwintering behavior of the fall released fish (Reagan et al. 2005). Subsequent studies (brood years 2005-2007) using PIT tag detections of fish leaving the hatchery found that very few fish that left the hatchery during the fall survived to migrate downstream to Bonneville Dam or survived to adult return. The fall volitional release strategy ended with brood year 2007. A spring only release has been used at the hatchery since brood year 2008.

## Present Objectives

Operations at the hatchery presently consist of adult collection, egg incubation and rearing of spring Chinook salmon. The current hatchery broodstock objective is to spawn 650 - 693



Chinook Salmon adults with a 60:40 female:male spawning ratio with jacks (< 60 cm in length) making up 5% of the broodstock (USFWS 2019). To account for 10% mortality between collection and spawning, 726 - 770 adults are collected for broodstock proportionately through the run based on wild stock timing and may be adjusted if temperatures exceed 16 °C. To maintain the stock integrity and genetic diversity of hatchery and wild spring Chinook salmon, approximately 10 percent natural origin fish have been incorporated into broodstock collection based on pre-season forecasts and in-season run size updates. However, if the wild run is less than 1,000 fish, no wild fish will be collected for broodstock. In a USFWS review of the WSNFH spring Chinook salmon program, the Hatchery Review Team (USFWS 2006) recommended that the program maintain the current goal of a minimum of 10 percent natural-origin spring Chinook Salmon in the broodstock and continue to limit hatchery-origin spring Chinook salmon on the spawning grounds to less than 10 percent. Small numbers of wild adult returns over the past decade have resulted in no wild fish being incorporated into the hatchery broodstock in most years. In recent years the CTWSRO developed a supplementation plan with the Service to hold and spawn a portion of the wild return at the hatchery. Excess milt from wild males has been spawned with hatchery females to integrate wild fish into the hatchery population. Remaining surplus hatchery origin spring Chinook salmon are dispatched and provided to the CTWSRO for tribal needs. After spawning, spring Chinook salmon are either placed in a landfill or are used for nutrient enhancement after they have been screened for disease and treated (eviscerated and heat-baked) to prevent disease transmission.

During years of low returns to the hatchery or unexpected losses to production, consideration has been given to augmenting the hatchery production with eggs or juveniles from other hatchery programs. The primary source of eggs during years of shortfall is from ODFW's Round Butte Hatchery (RB), located within the Deschutes River Basin. Eggs and juveniles from Parkdale Hatchery, located within the Hood River Basin, have also been used to augment the WSNFH production in recent years. To maintain the WSNFH genetic stock, any releases from non-WSNFH stocks are differentially marked (e.g., left or right ventral clip) and coded-wire tagged to distinguish them from WSNFH fish upon return. These stocks are excluded from the broodstock and distributed to the CTWSRO or to RB if needed. If returns to WSNFH are projected to be below broodstock needs, RB fish returning to WSNFH may be spawned and their progeny reared and marked separately from WSNFH stock.

When considering whether to rear and release non-WSNFH stock fish, managers must consider the risks and benefits of such actions. The main benefit of importing non-Warm Springs stock fish during years of shortfalls is an increase in the likelihood of meeting the annual juvenile release goal, which in turn leads to increased adult returns, thereby increasing harvest opportunities. Risks include the possibility that mis-clipped/unclipped fish returning to WSNFH as adults could be spawned with Warm Springs hatchery and even wild fish. Fish that stray into the Warm Springs population have a lower ability for their progeny to be successful. If they are frequently brought into the population, it could reduce chances for the long-term survival of the population. The performance of Parkdale and Round Butte stocks reared and released at Warm Springs NFH is currently being assessed.

The current production goal is on-station spring release of 750,000 externally marked smolts into the Warm Springs River (NMFS 2018). All juvenile fish released from the hatchery are marked (CWT and AD) to differentiate them from naturally produced fish upon return. Approximately 15,000 juveniles have been PIT tagged annually since brood year 2005 (BY05) (release year



2007). In 2022, the *United States v. Oregon* Policy Committee approved the change to the production table to incorporate the wild supplementation program into the existing hatchery program, and to footnote the transfer of ~560,000 juveniles to Little White Salmon NFH during the summer due to high water temperature concerns at Warm Springs NFH.

#### Hatchery Management Goals (USFWS 2019)

1. Produce Spring Chinook Salmon consistent with *United States v. Oregon* production goals for annual tribal harvest opportunity in Deschutes River and Columbia River fisheries
2. Provide for distribution to tribal members and the community freezer at CTWSRO
3. Provide safe passage for wild fish consistent with CTWSRO management of the Warm Springs River

#### CRFWCO Monitoring and Evaluation Objectives:

1. Monitor and evaluate on-station rearing strategies
  2. Monitor and evaluate juvenile releases, off-station juvenile survival, and migration
  3. Monitor release to adult return survival
  4. Develop run-reconstruction of adult returns, including contribution to harvest and returns to the hatchery
  5. Produce run forecasts for wild and hatchery returns
  6. Track passage of wild fish
  7. Conduct special studies as needed in consultation with the Warm Springs Hatchery Evaluation Team
-

## Hatchery Operations Summary

### On-Station Juvenile Production

#### a) Egg-to-Smolt Survival

Survival metrics during the early life stages include:

1. 95% or higher survival from the egg to eye up stage
2. 90% survival from the egg to fry stage; and
3. 97% survival from fry to smolt stage

Mortality can occur during each of these life stages due to disease, injury, predation, starvation, deformities, and genetic anomalies. Throughout the rearing cycle, the hatchery has a maximum Flow Index < 1.0 and Density Index <0.2 to minimize disease risk (USFWS 2019). Hatchery staff report these metrics to describe their production levels and determine whether alternative rearing and release practices are needed to improve on-station survival when warranted. This data is collected by hatchery staff and is not part of this report. A bulleted summary of infrastructure modifications or on-station rearing notes unique to 2022 can be found in Appendix A.

#### b) Juvenile Marking, Tagging, and Release Data

Funds distributed by the Service are used to meet annual juvenile release goals, process adult returns, for costs associated with PIT tagging, and equipment maintenance. The facility has an annual release goal of 750,000 spring Chinook salmon into the Warm Springs River (release year is two years after brood year). Fish released contribute to sport, commercial, and tribal fisheries while also providing for adequate escapement for hatchery production. The actual number of juveniles produced at WSNFH has varied by release year (Table 1) with a total annual mean of 517,072 juveniles released since 2013.

Since 2013, the facility has achieved a mean juvenile size of 24.8 fish/lb. at the time of release. While all juveniles are given an AD mark and CWT, the actual number of fish with marks and tags at release is estimated based on mark quality and tag retention sampling. Approximately 97% of the total number of juveniles released are AD and CWTed, with the remaining 3% released as AD only due to coded-wire tag loss, and very small percentage with no mark and no CWT (due to missing marks during the marking process). The actual number of juveniles that are mass-marked annually are presented below (Table 1). CWT codes are stored in the Service's Columbia River Information System (CRiS) database at the Columbia River Fish and Wildlife Conservation Office (CRFWCO) and reported annually to the Regional Mark Processing Center, Regional Mark Information System (RMIS) database.

In fall 2017, WSNFH requested surplus eggs from Round Butte state hatchery due to concerns about another year of high egg loss. Fortunately, the brood year 2017 (BY17) production did not experience elevated losses, however this led them to have extra juveniles on station above their 750,000 +/- 10% target (~65,000 above the high end 825,000 allowable). A one-time release of the extra 65,000 spring Chinook juveniles on-station in 2019 was determined not to have a

substantial effect on ESA-listed species above and beyond what was considered in the BiOp covering the WSNFH program (Rich Turner, 3/18/2019).

Beginning with BY20, wild origin adult returns have been held and spawned at the WSNFH. These wild fish are generally spawned into three groups (wild x wild [treated as wild], wild x hatchery [treated as wild], and wild x hatchery [treated as hatchery]). Wild and wild x hatchery crosses are kept separate during incubation and ponding until a decision is made on their final disposition. The BY20 wild x hatchery cross juveniles were split 50:50 between hatchery production and left unmarked for the wild fish population. The hatchery portion of the BY20 wild x hatchery fish resulted in 11,745 (4.5 %) of the total 257,953 fish released in 2022 as part of the hatchery program.

In 2022, the planned release of BY20 fish was scheduled to occur around April 8, 2022 to coincide with spill in the Columbia River system beginning around April 10<sup>th</sup>. However, in February 2022, Fish Health staff documented Bacterial Kidney Disease (BKD) causing increased mortality. The Service and the CTWSRO made the decision to release the fish in early March due to the status of the disease outbreak and the theory that releasing fish early would lower fish density and reduce BKD impacts. Given the close proximity to the release timing as described in the HGMP and BiOp, no impacts to listed species were anticipated (memo to NOAA, Jeremy Voeltz, 3/8/2022). The disease load and early release of BY20 fish are expected to impact smolt to adult return survival. Age three fish from the BY20 release will return in 2023, providing an early indicator of survival from release.

**Table 1. Annual hatchery program juvenile spring Chinook release dates from Warm Springs NFH into the Warm Springs River. Releases include Warm Springs Stock, Round Butte Stock (A), and Parkdale Stock (B). Round Butte stock fish were identified by a left ventral mark and Parkdale stock fish were identified by a right ventral mark in addition to the AD to distinguish stocks. Data includes marking and tagging information, number of juveniles released, release type forced (F), volitional (V) or accidental (X) and mean juvenile size at release. Data retrieved from CRiS SR80s File: 3/8/2023**

Release Year	Brood Year	Release Dates	Release Temp (°C)	Release Type	Mean Size (Fish/lb)	AD + CWT	AD Only	CWT Retention (%)	Total Released	Annual Total
2013	2011	27-Mar - 10-Apr		V	24	770,451	13,095	99.3	783,546	803,696
	2012	6-May		X	160	19,908	242	98	20,150	
2014	2012	31-Mar - 4-Apr	12.1	F	24	713,563	13,379	98	726,942	726,942
2015	2013	30, 31-Mar	8	F	28	344,834	26,621	93.1	371,455	371,455
2016	2014	30-Mar	10.6	F	22	129,349	3,682	93.5	133,031	133,031
2017	2015	30-Mar	6.7	F	24	396,864	17,451	95.8	414,315	533,714
	2015 (B)	30-Mar	5.6	F	30	112,460	6,939	94.2	119,399**	
2018	2016	29-Mar	7	F	22	533,560	7,881	98.9	541,441	541,441
2019	2017	3, 5-Apr	7.8	F	26	736,730	27,510	96.4	764,240	887,781
	2017 (A)	3-Apr	8.8	F	30	120,045	3,496	97.2	123,541*	
2020	2018	8-Apr	8.3	F	24.8	277,211	3,922	98.6	281,133	647,214
	2018 (A)	8-Apr	8.3	F	24.3	228,470	1,840	99.2	230,310*	
	2018 (B)	8-Apr	8.3	F	23.3	135,798	0	100	135,798**	
2021	2019	8-Apr	6.4	F	21.57	263,732	3,729	98.4	267,461	267,461
2022§	2020	9-Mar§	5.6	F	27.7	242,346	3,862	98.4	246,208	257,953
	2020†	9-Mar§	5.6	F	27.7	11,561	184	98.4	11,745	
Mean by Stock	Warm Springs				24.4	403,646	11,051	97.1	414,697	
	Round Butte (A)				27.1	174,258	2,668	98.2	176,926	
	Parkdale (B)				26.6	124,129	3,470	97.1	127,598	
<b>Total Annual Mean</b>			<b>8.1</b>		<b>24.8</b>	<b>503,688</b>	<b>13,383</b>	<b>97.0</b>		<b>517,072</b>

\*\*Left ventral clip to distinguish as Round Butte Stock

\* Right ventral clip to distinguish as Parkdale Stock

§ Fish released early in 2022 due to BKD.

† Wild x hatchery mix fish

## Off-Station Juvenile Production

### a) PIT Tagging Program

PIT tagging provides real-time tracking data as fish migrate from the Warm Springs River to the Columbia River, over Bonneville Dam (BONN), and to the Pacific Ocean. All data is stored in a regional database managed by Pacific States Marine Fisheries Commission called the Columbia Basin PIT Tag Information System (PTAGIS) and utilized by staff at CRFWCO to estimate juvenile post-release migration and survival, track adult returns, and estimate stray rates. In release years 2013-2022, approximately 15,000 juvenile spring Chinook were tagged annually with PIT tags in late January or early February during their release year from WSNFH (Table 2).

The detection rate of PIT tagged fish at BONN is a function of a) migration survival from release to BONN and b) the detection efficiency of the PIT antenna arrays at the dam. Since 2013, an average 15,321 juveniles have been PIT tagged at and released from WSNFH each year. The mean annual number of detections at BONN is 1,230, a detection rate of 7.9%. Average mean juvenile travel time to BONN after release is approximately 24 days with some juveniles spending up to 121 days between the facility and BONN before migrating downstream. Juveniles travel downstream and pass over BONN as quickly as 12 days or less after release (10<sup>th</sup> percentile). However, the majority of fish (90<sup>th</sup> percentile) pass over BONN within 34 days after release.

At the start of PIT tagging operations in February 2022, the BY20 fish had clinical BKD. The added stress of crowding, handling, and tagging caused increased mortality and PIT tagging operations were cancelled. With less than 3,500 fish tagged and an early release, estimates for juvenile survival in 2022 and future adult returns for BY20 will not be as robust.

**Table 2. The number of juvenile spring Chinook PIT tagged in a given release year and travel times in days (D) to Bonneville Dam (BONN) following release from Warm Springs NFH. Release year is two years after brood year. Migration times to BONN may be underestimated in 2013 due to PIT tagged fish having the option to exit ponds volitionally eight days before being forced into the Warm Springs River. Releases include Warm Springs Stock, Round Butte Stock (A), and Parkdale Stock (B). Data retrieved from PTAGIS:12/2/2022**

Release Year	# PIT Tagged	# Detected at BONN	% Detected	Mean (D)	Median (D)	Range (D)	10th	75th	90th
2013	14,965	1,244	8.3	28	29	(3 - 102)	15	31	35
2014	14,898	1,107	7.4	30	31	(4 - 121)	19	35	41
2015	14,915	1,425	9.6	21	23	(3 - 96)	7	28	33
2016	14,975	1,345	9	19	17	(3 - 118)	8	23	30
2017	9,896	289	2.9	25	27	(3 - 84)	7	34	37
2017 (B)	4,972	95	1.9	34	36	(4 - 56)	15	41	51
2018	14,903	955	6.4	24	26	(4 - 56)	9	33	36
2019	12,887	1,141	8.9	24	23	(3 - 94)	11	30	35
2019 (A)	2,097	160	7.6	30	30	(8 - 48)	22	35	38
2020	7,944	775	9.8	18	17	(4.5 - 121)	10	21	27
2020 (A)	6,566	598	9.1	18	18	(4.5 - 111)	10	21	25
2020 (B)	3,884	353	9.1	18	17	(4.5 - 106)	10	21	27
2021	14,986	1,585	10.6	18	19	(4.5 - 38)	12	21	24
2022	3,424	52	1.5	44	45	(9.5 - 83)	12	57	64
<b>Mean by Stock</b>									
Warm Springs	13,374	1,096	8.1	23	24		11	28	33
Round Butte (A)	4,332	379	8.4	24	24		16	28	32
Parkdale (B)	4,428	224	5.5	26	26		12	31	39
<b>Total Annual Mean*</b>	<b>15,321</b>	<b>1,230</b>	<b>7.9</b>	<b>24</b>	<b>24</b>		<b>12</b>	<b>29</b>	<b>34</b>

\* Mean values do not include 2022 early release year.

## b) Juvenile Survival

PIT tag detection histories are used to estimate the apparent juvenile survival from release at WSNFH downstream to BONN. A PIT tagged downstream migrating juvenile fish can pass BONN using a variety of routes, some of which have PIT tag detection arrays and some of which do not. For example, tagged fish passing through the turbines or through spillways would not be detected, while a fish passing through the juvenile bypass or corner collector could be detected. Since there is not 100% detection capability at BONN, detection probability must be estimated in order to separate out a tagged fish that died before reaching BONN from a tagged fish that was alive but was not detected as it passed BONN. For this analysis, apparent survival from release to BONN was estimated using the live recapture Cormack-Jolly-Seber model in Program MARK.



The model uses encounter histories of tagged fish to estimate the detection probability at BONN and estimate the apparent survival of fish from release to BONN. Survival estimates are reported on a scale from 0.0 to 1.0 (Table 3, Figure 3). To account for minijack returns, all juveniles detected at BONN before June 1 and in the juvenile bypass or corner collector after June 1 of their release year were considered juvenile detections. Minijacks detected in adult ladders after June 30 were considered downstream detections. As a note, the term “apparent survival” is used to indicate that a tagged fish that is alive, but never migrates past BONN, is considered a “mortality” in the model.

Estimated apparent juvenile survival of the Warm Springs NFH spring Chinook for BY11 - BY19 (release years 2013 - 2021) ranged from 0.5 to 0.95 (Table 3; Figure 3). Mean values do not include BY20 due to the early release in 2022.

**Table 3. Juvenile Spring Chinook survival from release at Warm Springs NFH to Bonneville Dam. Release year is two years after brood year. Estimates are median survival and lower and upper credible intervals. The Markov chain Monte Carlo Bayesian parameter estimation method in MARK was used to estimate the variance of the estimated survival. Data retrieved from PTAGIS:12/16/2022**

Release Year	Brood Year	Median Survival	95% Lower	95% Upper
2013	2011	0.71	0.56	0.86
2014	2012	0.64	0.52	0.78
2015	2013	0.5	0.42	0.58
2016	2014	0.6	0.43	0.8
2017	2015	0.66	0.28	1
2018	2016	0.59	0.39	1
2019	2017	0.64	0.44	0.87
2020	2018	0.95	0.71	1
2021	2019	0.57	0.41	0.84
2022	2020	0.12	0.03	0.56
Mean*		0.65	0.46	0.86

\* Mean values do not include 2022 early release year.




Figure 3. Juvenile Spring Chinook survival from release at Warm Springs NFH to Bonneville Dam (Brood Years 2011 - 2020). Estimates are median survival with 95% lower and upper credible intervals.

## Adult Returns: Smolt-to-Adult Survival, Detections, Age Structure, and Harvest Data

### a) Adult Returns

Adult returns to WSNFH are estimated by hatchery personnel and the marking and biosampling crew from CRFWCO. Coded Wire Tag recoveries maintained in the RMIS database are used to estimate the number of harvested adults and spawning ground recoveries (Table 4). Coded Wire Tag codes from fish released in respective brood years are queried in RMIS to determine freshwater and ocean recoveries by fishery. At WSNFH, the number of hatchery returns and harvested adults fluctuates from year to year. Collectively, the facility has produced a mean of 2,280 adults annually since BY06 resulting in a mean smolt-to-adult survival rate (SAR) of 0.42%. This is above the target SAR of 0.39% set from brood years 1978 – 2001 (CTWSRO 2007).

**Table 4. The estimated number of hatchery returns, harvested adults, and fish present on the spawning grounds for spring Chinook released from Warm Springs NFH. Hatchery return estimates include returns to Warm Springs NFH. Strays to non-federal hatcheries are included in the Total # of Adults. Adult returns are based on CWT recovery expansion data from RMIS. Due to delays in reporting to RMIS, CWT recoveries may be adjusted every year for accuracy. Data downloaded from RMIS on 4/18/2023.**

<b>Brood Year</b>	<b>Hatchery Returns</b>	<b>Columbia River Harvest</b>	<b>Ocean Harvest</b>	<b>Spawning Grounds</b>	<b>Total # Adults</b>	<b>Smolt-to-Adult Survival (%)</b>
2006	1,561	387	3	0	1,951	0.56
2007	2,938	508	1	0	3,447	0.59
2008	1,387	415	11	0	1,813	0.26
2009	1,366	73	5	0	1,444	0.27
2010	1,552	787	8	0	2,347	0.49
2011	6,451	1,555	29	0	8,035	1.03
2012	384	1,076	13	0	1,473	0.20
2013	1,350	197	9	0	1,556	0.42
2014	322	83	1	0	406	0.31
2015	313	19	1	0	333	0.06
2016*	903	5	0	0	908	0.17
2017*	1,614	479	1	0	2,094	0.24
<b>Mean</b>	<b>1,762</b>	<b>510</b>	<b>8</b>	<b>0</b>	<b>2,280</b>	<b>0.42</b>

\*Due to delays in reporting to RMIS, returns for brood years 2016 – 2017 are not complete. Mean calculated for Brood Years 2006-2015

An average 722 CWTs have been recovered each year at Warm Springs NFH since 2013 (Table 5). The Warm Springs NFH spring Chinook program accounts for 99 percent of all recoveries; spring Chinook from other programs include Round Butte (0.6%); spring Chinook from other programs account for < 0.5% of all recoveries.

**Table 5. Coded Wire Tag recoveries for all hatchery programs collected at Warm Springs NFH 2013 – 2022. Number of CWT recoveries are not expanded and do not reflect sample or tagging rates. Data retrieved from CRiS files and RMIS: 3/9/2023.**

<b>Return Year</b>	<b>CWT Recoveries</b>	<b>Hatchery Origin</b>	<b>% of Total Annual CWT Return</b>
2013	4	Lookingglass Hatchery	0.5
	1	Npt Hatchery	0.1
	1	Parkdale	0.1
	7	Round Butte Hatchery	0.9
	1	Sawtooth Hatchery	0.1
	798	Warm Springs NFH	98.3
2014	1	Rapid River Hatchery	0.1
	3	Round Butte Hatchery	0.3
	982	Warm Springs NFH	99.6
2015	5	Irrigon Hatchery	0.2
	1	Little White Salmon NFH	0.0
	2	Magic Valley Hatchery	0.1
	3	Round Butte Hatchery	0.1
	2192	Warm Springs NFH	99.5
2016	1	Cle Elum Hatchery	0.3
	1	Cottonwood Cr Pond	0.3
	4	Irrigon Hatchery	1.4
	4	Round Butte Hatchery	1.4
	279	Warm Springs NFH	96.5
2017	1	Irrigon Hatchery	4.3
	22	Warm Springs NFH	95.7
2018	7	Round Butte Hatchery	3.1
	222	Warm Springs NFH	96.9
2019	1	Imnaha Pond	0.3
	1	Klickitat Hatchery (YKFP)	0.3
	3	Round Butte Hatchery	0.9
	443	Warm Springs NFH	98.3
2020	6	Round Butte Hatchery	1.3
	2	Sawtooth Hatchery	0.4
	443	Warm Springs NFH	98.0
2021	807	Warm Springs NFH	99.0
	7	Round Butte Hatchery	1.0
2022	958	Warm Springs NFH	99.7
	3	Round Butte Hatchery	0.3
<b>Mean</b>	<b>722</b>		

## b) Bonneville Dam and Ladder Detections

Spring Chinook adults return and pass Bonneville Dam as early as Mar-20 and as late as Jul-20. The average median Bonneville Dam passage date of PIT tagged Spring Chinook adults (Ages 3, 4, and 5) released from WSNFH is May-09 (Table 6).

**Table 6. Median Bonneville Dam passage date of adult spring Chinook PIT tagged and released from Warm Springs NFH ( $\geq$  Age 3). Confidence limits do not include detections of five fish or fewer per age group to reduce the variability and increase the accuracy of the estimate. Data retrieved from PTAGIS: 3/13/2023**

<b>Return Year</b>	<b>Median Passage Date</b>	<b>First Detection Date</b>	<b>Last Detection Date</b>	<b># of Fish Detected</b>	<b>Bonneville Expansion</b>	<b>95% CI</b>	<b>Hat. Return</b>	<b>Hat. Return/Bonn. Expansion (%)</b>
2013	May-10	Apr-26	Jul-01	58	1,597	(1,087 - 2,304)	1,818	114
2014	May-05	Apr-05	Jun-21	136	5,180	(4,058 - 6,711)	2,478	48
2015	Apr-27	Mar-20	Jul-04	195	10,348	(8,691 - 12,483)	6,635	64
2016	May-07	Apr-06	Jul-01	142	6,724	(5,517 - 8,119)	782	12
2017	May-23	May-04	Jul-20	76	1,620	(1,156 - 2,067)	1,748	108
2018	May-07	Apr-23	Jun-11	46	456	(289 - 520)	260	57
2019	May-09	Apr-30	May-13	9	371	(172 - 772)	391	105
2020	May-07	Apr-22	Jul-13	19	759	(361 - 1,012)	542	71
2021	May-11	Apr-24	Jun-28	56	3,079	(2,047 - 4,897)	1,746	57
2022	May-14	Apr-01	Jul-14	66	2,157	(1,261 - 2,993)	1,224	57
<b>Mean</b>	<b>May-09</b>	<b>Apr-16</b>	<b>Jun-27</b>	<b>80</b>	<b>3,229</b>		<b>1,762</b>	<b>69</b>

Since Return Year 2013, spring Chinook adults ( $\geq$  Age 3) PIT tagged and released from Warm Springs NFH returned to the Warm Springs NFH Ladder as early as Apr-14 and as late as Sep-07 with the average median May-25 (Table 7).

**Table 7. Median passage date at Warm Springs NFH Ladder of adult spring Chinook PIT tagged and released from Warm Springs NFH ( $\geq$  Age 3). Confidence limits do not include detections of five fish or fewer per age group to reduce the variability and increase the accuracy of the estimate. Data retrieved from PTAGIS: 3/13/2023**

<b>Return Year</b>	<b>Median Passage Date</b>	<b>First Detection Date</b>	<b>Last Detection Date</b>	<b># of Fish Detected</b>	<b>Ladder Expansion</b>	<b>95% CI</b>	<b>Hat. Return</b>	<b>Hat. Return /Ladder Expansion (%)</b>
2013	May-29	May-11	Sep-04	47	1,300	(806 - 1,902)	1,818	140
2014	May-22	Apr-24	Jul-27	79	3,015	(2,186 - 4,270)	2,478	82
2015	May-15	Apr-18	Sep-07	138	7,310	(5,933 - 9,123)	6,635	91
2016	May-25	May-03	Sep-06	59	2,788	(2,026 - 3,499)	782	28
2017	Jun-11	May-20	Aug-28	43	926	(571 - 1,274)	1,748	189
2018	May-23	May-12	Aug-26	24	213	(147 - 327)	260	122
2019	May-23	May-19	May-27	4	162	(NA - NA)	391	241
2020	May-26	Apr-14	Aug-16	12	505	(174 - 711)	542	107
2021	May-17	May-01	Jun-09	31	1,791	(1,202 - 2,562)	1,746	97
2022	Jun-01	May-04	Jul-10	34	1,005	(561 - 1,520)	1,224	122
<b>Mean</b>	<b>May-25</b>	<b>May-03</b>	<b>Aug-03</b>	<b>47</b>	<b>1,902</b>		<b>1,762</b>	<b>122</b>



### c) Age Structure

Monitoring adult returns to the hatchery provides information on sex ratios, length information, and age structure (Table 8: brood year; Table 9: return year). Service staff use CWT recoveries and scale sampling to age fish. Since return year 2013, approximately 89% of adults have returned to the facility at Age-4, approximately 10% have returned as jacks at Age-3, and 1% have returned at Age-5. No Age-6 returns have been documented (Table 9). The facility has a mean of 1,782 adult returns each year. There is a goal to have between 2% and 5% of jacks in the broodstock based on the percentage of jacks in the wild population and their estimated contribution during spawning (CTWSRO 2007).

**Table 8. Estimated age structure of hatchery adult spring Chinook returns to WSNFH by brood year for all stocks released at WSNFH. CRiS Age Composition Report run on: 1/31/2023**

<b>Brood Year</b>	<b>Age-3</b>	<b>Age-4</b>	<b>Age-5</b>	<b>Age-6</b>	<b>Total # Adults</b>
2007	553	3,047	59	0	3,659
2008	708	1,249	45	0	2,002
2009	46	1,425	7	0	1,478
2010	348	2,082	39	0	2,469
2011	389	6,207	29	0	6,625
2012	389	748	40	0	1,177
2013	5	1,509	2	0	1,516
2014	199	243	7	0	449
2015	15	355	8	0	378
2016	29	373	3	0	405
2017*	161	1,677	33	NA	NA
2018*	67	1,003	NA	NA	NA
2019*	188	NA	NA	NA	NA
<b>Mean</b>	<b>238</b>	<b>1,660</b>	<b>25</b>	<b>0</b>	<b>2,016</b>

\*Denotes incomplete brood years given that adults have not yet returned to the hatchery.

**Table 9. Total number of hatchery adult spring Chinook returns to WSNFH and estimated age structure by *return year* for all stocks released at WSNFH. CRiS Age Composition Report run on: 1/31/2023**

<b>Return Year</b>	<b>Age-3</b>	<b>Age-4</b>	<b>Age-5</b>	<b>Age-6</b>	<b>Total # Adults</b>
2013	348	1,425	45	0	1,818
2014	389	2,082	7	0	2,478
2015	389	6,207	39	0	6,635
2016	5	748	29	0	782
2017	199	1,509	40	0	1,748
2018	15	243	2	0	260
2019	32	542	8	0	582
2020	161	373	8	0	542
2021	67	1,677	3	0	1,747
2022	188	1,003	33	0	1,224
<b>Mean</b>	<b>179</b>	<b>1,581</b>	<b>21</b>	<b>0</b>	<b>1,782</b>

#### **d) Hatchery Returns by Stock**

Collecting CWTs provides additional information for multiple stocks released on station returning as adults to the hatchery. Each stock has a unique CWT which can be used to calculate the age at return for each fish. The majority of adults from all stocks return to the facility at Age-4 (Table 10, Figure 4). A Chi-square test of independence was performed to examine the relationship between stock and returns to the hatchery. The relationship between these variables was significant, ( $X^2(2, N = 2,535,039) = 182.88, p < 2.2e-16$ ). Returns differ by stock origin, adults from the Warm Springs stock return at a significantly higher percentage in all years (Table 10, Figure 5). For example, for every 100,000 juvenile fish released from each stock, an average of approximately 128 Warm Springs, 63 Round Butte, or 45 Parkdale adults return to the hatchery.

For BY09, the return percentage of Warm Springs stock was 96% greater than the Round Butte stock. For BY15, return percentage of the Warm Springs stock was 111% greater than the Parkdale stock. For BY17, return percentage of the Warm Springs stock was 100% greater than the Round Butte stock. Returns for BY18 are not yet complete and will require an additional year for a complete return analysis. So far, the return percentage of the Warm Springs stock (ages 3 and 4) is 57% greater than the Round Butte stock and 88% greater than the Parkdale stock.

**Table 10: Proportion by age at return and return percentage to WSNFH for Warm Springs, Parkdale, and Round Butte stock releases. Data is based on unexpanded CWT recoveries of fish returning by stock and brood year. Data retrieved from CRFWCO marking files and RMIS: 3/9/2023.**

<b>Stock</b>	<b>Brood Year</b>	<b>Total CWT Released</b>	<b>Total CWT Return</b>	<b>Age-2 % of Return (n)</b>	<b>Age-3 % of Return (n)</b>	<b>Age-4 % of Return (n)</b>	<b>Age-5 % of Return (n)</b>	<b>Return Percentage (Return/Release) (95% CI)</b>
Round Butte	2009	216,162	136	0	2.2% (3)	97.8% (133)	0	0.06% (0.05 - 0.07%)
Warm Springs	2009	311,296	544	0.2% (1)	4.0% (22)	95.0 % (517)	0.7% (4)	0.17% (0.16 - 0.19%)
Parkdale	2015	112,460	20	0	0	90.0% (18)	10.0% (2)	0.02% (0.01 - 0.03%)
Warm Springs	2015	396,864	294	0.3% (1)	4.4% (13)	93.5% (275)	1.7% (5)	0.07% (0.07 - 0.08%)
Round Butte	2017	120,045	38	0	15.8% (6)	78.9% (30)	5.3% (2)	0.03% (0.02 - 0.04%)
Warm Springs	2017	736,730	678	0.4% (3)	17.3% (117)	78.5% (532)	3.8% (26)	0.09% (0.09 - 0.10%)
Parkdale	2018	135,798	93	7.5% (7)	1.1% (1)	91.4% (85)	NA	0.07% (0.06 - 0.08%)
Round Butte	2018	228,471	222	1.8% (4)	2.7% (6)	95.5% (212)	NA	0.10% (0.08 - 0.11%)
Warm Springs	2018	277,213	508	2.8% (14)	4.3% (22)	92.9% (472)	NA	0.18% (0.17 - 0.20%)
Parkdale								0.05%
Round Butte								0.06%
Warm Springs								0.13%




Figure 4. Age-at-return proportion for stocks reared and released from WSNFH, by brood year.




Figure 5. Return rate (percent of release returning to WSNFH) for stocks reared and released from WSNFH, by brood year.

### e) External Mark Accuracy by Stock

In 2018, low Warm Springs stock adult returns led to WSNFH requesting and receiving eggs from both Parkdale Hatchery and Round Butte Hatchery to augment hatchery production. Each stock received a unique CWT code and differential fin marks (ADLV/ADRV/AD) to identify their stock of origin when they returned to the hatchery as adults.

Based on visual identification of adult returns at the hatchery and subsequent reading of coded-wire tags there was an overall 11% error rate in visual identification of stocks by fin marks. Of the total 84 visual call errors, 59 (70%) were likely due to the regeneration of ventral fins (i.e., Round Butte and Parkdale stocks misidentified as Warm Springs stock) (Table 11).

- Error rate was lowest for Warm Springs fish (2%), of the 473 coded-wire tagged Warm Springs fish that returned as adults, 464 (98%) were identified correctly (AD clip); 9 (2%) were misidentified as Round Butte (ADLV clip) and zero were misidentified as Parkdale (ADRV clip).
- Error rate for Round Butte fish was (22%), of the 211 Round Butte coded-wire tagged fish that returned as adults, 165 (78%) were identified correctly (ADLV clip); 8 (4%) were misidentified as Parkdale (ADRV clip) and 38 (18%) were misidentified as Warm Springs (AD clip) fish.
- Error rate was highest for Parkdale fish (34%), Of the 85 Parkdale coded-wire tagged fish that returned as adults, 56 (66%) were identified correctly (ADRV clip); 8 (9%) were misidentified as Round Butte fish (ADLV clip), and 21 (25%) were misidentified as Warm Springs (AD clip) fish.

**Table 11. External mark identification accuracy for BY18 (release year 2020) Warm Springs, Round Butte, and Parkdale stock returning to the WSNFH as adults. Data retrieved from CRiS Biosample File 3/20/2023.**

<b>Stock</b>	<b>Correctly Identified by External Mark</b>	<b>Incorrectly Identified by External Mark</b>	<b>Error Rate (Incorrectly Identified/Total CWT Return)</b>	<b>Incorrectly Identified as Warm Springs Stock</b>
Warm Springs	464	9	2%	NA
Round Butte	165	46	22%	38
Parkdale	56	29	34%	21



## f) Adult Harvest

The WSNFH provides salmon to supplement tribal and sport harvest opportunities in the Deschutes and Columbia Rivers. Estimates of wild and hatchery spring Chinook are based on ODFW and CTWSRO creel surveys (Table 12).

**Table 12. Deschutes harvest estimates of wild and WSNFH spring Chinook salmon 2013 - 2022. Estimates based on ODFW and CTWSRO creel surveys, and estimated proportion of total harvest of Warm Springs NFH and Round Butte Hatchery returns. Dashed line indicates limited Tribal harvest; however, no creel survey took place in 2020 or 2021 due to COVID-19 safety precautions. Data retrieved from ODFW and Round Butte Hatchery hatchery return data 1/4/2023**

Return Year	Wild				WSNFH				Total Harvest
	Adult Sport	Jack Sport	Adult Tribal	Jack Tribal	Adult Sport	Jack Sport	Adult Tribal	Jack Tribal	
2013	0	0	2	0	0	0	40	134	176
2014	0	0	21	0	436	189	124	22	792
2015	0	0	17	0	0	0	365	23	405
2016	0	0	0	0	955	21	643	19	1,638
2017	0	0	0	0	0	0	172	11	183
2018	0	0	0	0	57	3	9	0	69
2019*	0	0	0	0	0	0	22	0	22
2020*	0	0	-	-	0	0	-	-	-
2021*	0	0	-	-	0	0	-	-	-
2022*	0	0	0	0	0	0	28	7	35
<b>Mean</b>	<b>0</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>145</b>	<b>21</b>	<b>175</b>	<b>27</b>	<b>415</b>

\* No Sport Harvest 2019 - 2022; limited Tribal harvest 2020 - 2022.

During the spring Chinook salmon migration period (April 15 - August 30) all fish ascending the fish ladder are diverted into catch and holding ponds. Fish are sorted, and either passed up stream if wild appearing (i.e., no external marks), held for broodstock, or surplused (known hatchery fish) and given to tribal representatives for distribution. Disposition of fish depends on number of returns, their condition, and origin (Tables 13 and 14).

**Table 13. Disposition of hatchery (all stocks) and wild spring Chinook salmon at Warm Springs NFH, 2013 - 2022. Numbers include spring Chinook salmon adults (age 4-5) and jacks (age 3) that were surplus to hatchery broodstock needs before being distributed to the CTWSRO. Number distributed based on hatchery fish-removal-file records of distribution of adult fish that returned to Warm Springs NFH prior to August 1 of each year. Total may include uses not listed. Unresolved data discrepancies in 2019, 2020, and 2021 Fish Removal files. Data retrieved from CRFWCO CRiS Fish Removal file 1/6/2023.**

Return Year	WSNFH Upstream	WSNFH Surplus Donated	WSNFH Surplus Dump	WSNFH Brood	WSNFH Transfer	WSNFH Mortality	WSNFH Total	Wild Upstream	Wild Brood	Mortality	Wild Green	Wild Surplus	Wild Total
2013	63	689	0	482	0	184	1,418	397	0	3	0	0	400
2014	0	996	0	194	35‡	486	1,711	761	0	6	0	0	767
2015	78	4,901	145	615*	0	188	6,635	1,369	0	12	0	0	1,381
2016	0	1,550	3	758†	323§	150	2,784	335	0	13	0	0	348
2017	-	-	-	-	-	80	1,555**	-	-	-	0	0	193
2018	0	5	0	206	0	47	260	247	0	0	0	0	247
2019	0	0	39	259	0	94	394	204	0	0	0	0	204
2020	3	0	70	360	0	79	566	10	35	5	0	0	50
2021	352^	159	73	166	426	570	1,746	67	78	35	3	1	184
2022	0	663	23	191	0	347	1,224	184	29	84	2	0	299
<b>Mean</b>	<b>55</b>	<b>996</b>	<b>39</b>	<b>398</b>	<b>87</b>	<b>222</b>	<b>1,829</b>	<b>392</b>	<b>16</b>	<b>17</b>	<b>0</b>	<b>0</b>	<b>402</b>

\* All fish spawned at Little White Salmon NFH in 2015.

†645 fish spawned at Little White Salmon NFH in 2016.

‡Transferred to ODFW for research.

§Transferred to Round Butte, CTWSRO, PGE.

\*\*No data recorded for other uses of fish in 2017.

^Fish were transferred to CTWSRO (352) or Round Butte Hatchery (12) and outplanted upstream

**Table 14: Adult Spring Chinook passed at WSNFH or trucked upstream each year to spawn naturally. Data from CRiS Fish Removal file 3/14/2023**

Return Year	Origin	Upstream Passage Method	Females	Males	Jacks	Unknown	Total	Annual Total
2012	Wild	Passed	2	5	24	349	380	414
	Hatchery	Passed	0	3	7	24	34	
2013	Wild	Passed	1	1	88	307	397	460
	Hatchery	Passed	0	0	3	60	63	
2014	Wild	Passed	2	6	100	653	761	761
2015	Wild	Passed	0	0	58	1,311	1,369	1,447
	Hatchery	Passed	0	0	28	50	78	
2016	Wild	Passed	0	0	4	331	335	335
2017	Wild	Passed	0	3	14	169	186	186
2018	Wild	Passed	6	15	5	221	247	248
	Hatchery	Passed	0	1	0	0	1	
2019	Wild	Passed	3	5	3	193	204	204
2020	Wild	Passed	7	1	0	2	10	13
	Hatchery	Trucked	0	0	3	0	3	
2021	Wild	Trucked	30	37	0	0	67	419
	Hatchery	Trucked	204	123	1	24	352	
2022	Wild	Trucked	126	45	7	0	178	184
	Wild	Passed	2	1	3	0	6	
	Hatchery	Trucked	0	0	0	0	0	

## 2022 Run Reconstruction

Run reconstruction estimates the number of age 3, age 4, and age 5 fish that returned to the mouth of the Deschutes River for a given brood year. Run reconstruction methods and data can be found in the annual run forecast reports (Lovtang et al. 2011). In 2022, an estimated 299 wild and 3,351 hatchery Warm Springs River spring Chinook are estimated to have returned to the Deschutes River (Tables 15 and 16).

There was limited tribal harvest and no sport harvest in 2020. During the 2020 Deschutes-Sherars Falls Spring Chinook Harvest, it is likely 50 fish may have been harvested. However, there were no spring or fall creel surveys due to COVID-19 safety precautions (Mark Manion, CTWSRO).

**Table 15. Run reconstruction of *wild* spring Chinook salmon from the Warm Springs River, 2022. Run reconstruction performed 1/6/2023, aging for age 3 fish based on hatchery records of jacks (fish < 60cm), ages 4 and 5 were estimated based on historical data (85% of fish > 60cm Age 4)..**

<b>Location</b>	<b>Wild Stock Disposition</b>	<b>Age 3</b>	<b>Age 4</b>	<b>Age 5</b>	<b>Total 4 + 5</b>	<b>Total</b>
To WSNFH	Upstream of WSNFH	10	174	0	174	184
To WSNFH	WSNFH Broodstock	1	28	0	28	29
To WSNFH	DIPS/Jumpouts/killed	4	80	0	80	84
To WSNFH	Surplus/Green	0	2	0	2	2
To WSNFH	<b>Total</b>	<b>15</b>	<b>284</b>	<b>0</b>	<b>284</b>	<b>299</b>
Harvest	Sport	0	0	0	0	0
Harvest	Tribal	0	0	0	0	0
Harvest	<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
Below WSNFH	Spawned	0	0	0	0	0
<b>Total Estimated Return</b>		<b>15</b>	<b>284</b>	<b>0</b>	<b>284</b>	<b>299</b>

Age 3 based on hatchery records of jacks (fish < 60cm), Age 4 and 5 were estimated based on historical data (85% of fish > 60cm Age 4)

**Table 16. Run reconstruction of *hatchery* spring Chinook salmon from the Warm Springs River, 2022. Run reconstruction performed 1/6/2023**

Location	Hatchery Stock Disposition	Age 3	Age 4	Age 5	Total 4 + 5	Total
To WSNFH	Upstream of WSNFH	0	0	0	0	0
To WSNFH	Surplus (Ad Only)	156	169	5	174	330
To WSNFH	Surplus (AdLV or AdRV)	3	350	3	353	356
To WSNFH	Mortality (Ad Only)	30	301	17	318	346
To WSNFH	Mortality (AdLV or AdRV)	0	1	0	1	1
To WSNFH	Spawned/Green (Ad Only)	16	164	9	173	190
To WSNFH	Total to WSNFH (including AdLV and AdRV)	205	985	34	1,019	1,224
To WSNFH	Total to WSNFH (excluding AdLV and AdRV)	202	635	31	666	868
Harvest	Sport	0	0	0	0	0
Harvest	Tribal	7	27	1	28	35
Harvest	Total	7	27	1	28	35
<b>Total Estimated Return</b>		<b>212</b>	<b>1,012</b>	<b>35</b>	<b>1,047</b>	<b>1,259</b>
<b>Total Estimated Return (excluding AdLV and AdRV)</b>		<b>209</b>	<b>662</b>	<b>32</b>	<b>694</b>	<b>903</b>

No sport harvest in 2022, Limited Tribal harvest in 2022.

## 2022 Wild Spring Chinook Management

### Wild Adult Return

The wild Spring Chinook Salmon population in the Warm Springs River has been below the minimum escapement goal of 1,000 adults upstream of Warm Springs NFH in nine out of the last ten years (Table 14). Additionally, pre-spawn mortality of wild fish upstream of the hatchery (defined as the number of fish per redd) has been increasing in the past 10 years (ranging 10 -15 fish per redd). In 2020, the preseason forecast for wild adult (ages 4 and 5) spring Chinook salmon returns to the Deschutes Basin ranged from 50 to 300 total returns. In early 2020, the CTWSRO developed a plan to transport one-third of the wild adults from Warm Springs NFH to the spawning areas in the upper watershed. The hope was that transporting the fish would move them upstream of a possible thermal barrier that was hypothesized to block upstream migrating fish from reaching the spawning grounds. In addition to transporting wild adult fish, the CTWSRO began having discussions with the Service about collecting one-third of the wild fish for on-hatchery spawning and rearing at Warm Springs NFH. The remaining one-third of the fish would be passed upstream of the hatchery as in previous years. However, due to the COVID-19 outbreak in March of 2020 and subsequent working restrictions, only 10 wild fish were passed upstream of the hatchery and the decision was made to collect the remaining 40 wild fish for holding and spawning at the hatchery.

In 2021 and 2022, the wild fish plan was to collect 60 female and 40 male wild adults (as determined by ultrasound) for on-hatchery spawning and rearing; excess wild adults would be trucked upstream of the hatchery to naturally spawn. Broodstock held on station would be injected with antibiotics.

In 2022, a total of 299 wild fish were collected at the hatchery. Of the 115 wild fish held for broodstock, 84 fish died before spawning, two fish were green females, and 29 were spawned (Table 17, Figure 6). Between May 23 and August 11, 178 wild fish were trucked upstream and outplanted by the CTWSRO. An ultrasound was used to distinguish 126 females and 52 males before they were outplanted. An additional six wild fish (2 female, four male) were passed upstream of the hatchery via the outgoing channel.

The wild-origin broodstock were sexed by ultrasound, injected with Draxxin, and held in the broodpond closest to natural sunlight. Spawning took place between August 24<sup>th</sup> and September 13<sup>th</sup> and occurred in six takes. A 2x2 spawning matrix was used to increase the number of family groups and genetic diversity for the supplementation production. With 2x2 spawning, each wild female's eggs were split into 2 buckets, and 2 wild males were spawned with each female (Figure 7). After time to allow for fertilization, the 2 buckets of eggs were recombined and placed into incubation trays.


With the supplementation program bringing wild fish into the hatchery program, it was possible to integrate wild fish into the hatchery population to minimize genetic divergence between Warm Springs River hatchery and wild stocks. Excess milt from each wild male was spawned in a 2x2 spawning matrix with hatchery females (Figure 7). Offspring from these wild x hatchery crosses are tracked and kept separate during incubation and ponding until a decision is made on their final disposition (e.g., leave all unmarked and release into the wild, or use 50% for hatchery production and 50% for unmarked wild release, or use all for hatchery production).

Monitoring wild adult returns at the hatchery provided information on sex ratios, length information, and age structure based on scale samples. Approximately 95% of wild adults collected at the facility were Age-4 (n = 284) and 5% were jacks at Age-3 (n = 15), zero Age-5 fish were documented (Table 18).



**Table 17. Total number of wild adult spring Chinook returns to WSNFH, estimated age structure, and disposition. Data from CRiS Age Composition Report 3/14/2023**

	<b>Total Adult Return</b>	<b>Passed Upstream</b>	<b>Surplused</b>	<b>Green</b>	<b>Mortality</b>	<b>Spawnd</b>
Male	120	56	0	0	45	19
Female	179	128	0	2	39	10
Unknown	0	0	0	0	0	0
<b>Total</b>	<b>299</b>	<b>184</b>	<b>0</b>	<b>2</b>	<b>84</b>	<b>29</b>



*Figure 6. Wild adult spring Chinook returns to WSNFH and their distribution. One hundred eighty four fish were passed upstream, and 29 fish were spawned at the hatchery. An additional 15 hatchery females were spawned with wild males to integrate wild fish genetics into the hatchery population.*

### 2022 Warm Springs NFH Wild Spawning Matrix




Figure 7. A 2x2 spawning matrix was used to increase wild adult spring Chinook family groups and genetic diversity. Each wild female's eggs were spawned with 2 wild males. Excess milt from each wild male was also spawned with two hatchery females in a 2x2 spawning matrix.

**Table 18. Estimated age structure of wild adult spring Chinook returns to Warm Springs NFH based on scale samples of wild returns used for brood stock. Data from CRiS Age Composition Report 3/14/2023.**

Return Year	Age-3	Age-4	Age-5	Total # Adults
2020	5	40	5	50
2021	9	172	3	184
2022	15	284	0	299
<b>Mean</b>	9	151	2	161

### Wild Juvenile Release

Approximately 52,600 BY20 juveniles were released as wild fish (unmarked, untagged offspring of at least one wild origin parent) on March 9, 2022 (Table 19). Fish were released approximately one month early due to an outbreak of BKD among the BY20 population. Plans were made to release the BY20 fish upstream in spring 2022, however, due to concerns with BKD affecting BY21 hatchery fish downstream, BY20 wild juveniles were released at the hatchery. The BY20 wild release included wild x hatchery cross juveniles split 50:50 between hatchery production and wild fish population. The unmarked untagged “wild” portion of the BY20 wild x hatchery fish resulted in approximately 21% of the wild release (11 K of the total 52.6 K).

In May 2022, the CTWS BNR released 85,533 wild BY21 fish (unmarked, untagged offspring of wild origin parents) in locations in the upper Warm Springs River basin instead of transporting to LWSNFH for summer rearing (Table 19). The unmarked wild component of the wild x hatchery cross juveniles (approximately 38 K) were transported to LWSNFH for summer rearing to allow

for potential PIT tagging in spring of 2023, however none of these fish were Pit tagged prior to release in 2023.

**Table 19. Annual wild juvenile spring Chinook release dates. Data includes release location, number released and lifestage. Data retrieved from CRiS SR80s File: 3/8/2023.**

<b>Brood Year</b>	<b>Release Year</b>	<b>Release Date(s)</b>	<b>Release Location</b>	<b>Release Number</b>	<b>Lifestage</b>
2020	2022	3/9	Warm Springs NFH	52,600	Yearling
2021	2022	5/17 – 5/24	Upper Warm Springs River Basin	85,533	Fingerling

## 2023 Run Forecast

### Warm Springs River Wild stock

The mean forecasted adult (Age-4 and Age-5) return ranges from 170 to 410 wild fish to the mouth of the Deschutes River (Table 20). There is approximately a 30%-40% chance that the adult return will be less than 100 fish.

### Warm Springs NFH program

The mean forecasted adult (Age-4 and Age-5) return ranges from 1,400 to 2,200 hatchery fish to the mouth of the Deschutes River (Table 20). There is approximately a 60% to 70% chance that the return will be greater than 1,000 hatchery fish. The forecast is for Warm Springs NFH stock fish only. An additional one to ten Age 5 Round Butte/Parkdale stock fish may be expected to return. Age-3 returns (jacks) will be from the BY20 release, which was released in early March due to BKD outbreak. Low survival of BY20 is expected.

**Table 20. Forecast Model Predictions of Spring Chinook Salmon Returns to the Deschutes River in 2023 based on Hand and Haeseker (2011). Run date as of: 1/5/2023**

<b>Return</b>	<b>Std Reg (All Data)</b>	<b>Std Reg (10 yr Data)</b>	<b>Return Ratio (10 yr Data)</b>	<b>% Age Model (10 yr Data)</b>
Wild Age 4	362	202	300	133
Wild Age 5	48	46	41	37
Wild Total	410	248	341	170
<b>Wild Prob. &gt;100 adults return</b>	<b>60%</b>	<b>57%</b>	<b>64%</b>	<b>52%</b>

<b>Return</b>	<b>Std Reg</b>	<b>LN Reg</b>	<b>Return Ratio (All Data)</b>	<b>% Age Model (All Data)</b>
WSNFH Age 4	1,670	1,580	2,189	1,452
WSNFH Age 5	37	26	35	33
WSNFH Total	1,707	1,606	2,224	1,485
<b>WSNFH Prob. &gt;1,000 adults return</b>	<b>64%</b>	<b>64%</b>	<b>72%</b>	<b>62%</b>

<b>Return</b>	<b>Std Reg</b>	<b>LN Reg</b>	<b>Return Ratio (All Data)</b>	<b>% Age Model (All Data)</b>
RBH Age 4	1,809	1,896	2,378	1,619
RBH Age 5	60	39	90	91
RBH Total	1,869	1,935	2,468	1,710

## Transfers

In recent years, the primary issues related to meeting release goals has been maintaining broodstock health after they have returned to the hatchery, achieving eye-up, and survival until marking. This facility has transferred spring Chinook between other state, tribal and federal hatcheries to make up for loss of eggs, provide relief from high water temperatures, and accommodate power outages due to hatchery construction. WSNFH stock has been transferred to Round Butte State Fish Hatchery and vice versa. Round Butte hatchery stock are the preferred stock for backfilling production shortfalls at WSNFH (Smith et al. 2016; Smith, 2018). Parkdale Hatchery stock have been reared and released from WSNFH; however, the Parkdale program has been trying to develop their own, Hood River, stock and the genetic and phenotypic differences between the Parkdale stock and the WSNFH stock are not fully known. Early results from release of Parkdale stock fish from WSNFH indicated lower adult returns than the WSNFH stock. An evaluation-of the brood year 2018 on-hatchery and off-hatchery performance of the WSNFH, Round Butte, and Parkdale stocks will provide additional information on the similarities and differences. All juvenile or egg transfers from Round Butte Hatchery to Warm Springs NFH are marked with AD+CWT+LV; all juvenile or egg transfers from Parkdale

Hatchery to Warm Springs NFH are marked with AD+CWT+RV. Lastly, both adults and juveniles have been transferred from WSNFH to the Little White Salmon NFH (LWSNFH) for spawning and temporary rearing before being transferred back to WSNFH (Table 21).

Due to elevated water temperatures at WSNFH in 2015 and 2021, the USFWS and CTWSRO-Bureau of Natural Resources (BNR) Hatchery Evaluation Team decided to move fish off station to LWSNFH. Those moves occurred as ‘emergency’ actions under less-than-ideal conditions (air temperatures over 100° F and water temperatures approaching 70° F). In anticipation of the reoccurring increased temperatures in the Warm Springs River, the regular transfer of WSNFH juveniles to LWSNFH began in 2022. The proactive transfer of juvenile fish to another facility between May and September is necessary to avoid chronic and acute thermal stress when temperatures can exceed optimal rearing conditions in the summer months. Rearing fish for the best animal welfare in more favorable conditions is intended maximize survival and reduce the need for multiple antibiotic treatments. During this time, the LWSNFH has available space and conditions for 560,000 fish until adaptive modifications are made to in-basin rearing facilities.

The CTWSRO-BNR decided to release the BY21 wild source subyearling in May of 2022 in locations in the upper Warm Springs River basin instead of transporting the wild source juveniles to LWSNFH for summer rearing. Fish transfer took place over two weeks in mid-May of 2022.

**Table 21. Transfer dates and total number of spring Chinook from three stocks, (WS) Warm Springs, (RB) Round Butte, and (P) Parkdale. Transfer locations were to or from (WS) Warm Springs NFH, (RB) Round Butte State Fish Hatchery, (P) Parkdale Hatchery, and (LW) Little White Salmon NFH.**

Transfer Year	Transfer Dates	Brood Year	Stock	Lifestage	Transfer From	Transfer To	Total # Transferred
2015	Spring		WS	Adults	WS	LW	680
	Spring		WS	Adults	WS	RB	708
	Summer	2014	WS	Juveniles	WS	LW	133,031
	November	2015	WS	Eggs	LW	WS	926,679
	November	2015	WS	Eggs	RB	WS	401,954
2016	Spring	2015	P	Juveniles	P	WS	130,000
	Spring	2015	RB	Juveniles	RB	WS	45,000
	Summer	2015	WS, RB, P	Juveniles	WS	LW	450,000
	July		WS	Adults	WS	LW	NA
	November	2016	WS	Eggs	LW	WS	NA
2017	June		WS	Adults	WS	LW	NA
	Fall	2017	RB	Eggs	RB	WS	NA
2018	September	2018	P	Eggs	P	WS	153,538
	September	2018	RB	Eggs	RB	WS	249,186
2019	April	2018	RB	Juveniles	RB	WS	15,000
2021	May		RB	Adults (AD/LV)	WS	RB	128
	May		WS	Adults (AD)	WS	RB	286
	May -July		WS	Adults (AD)	WS	Upstream Natural Spawning	352
	May- July		WS	Adults (Wild)	WS	Natural Spawning	18
	June	2020	WS	Juveniles	WS	LW	300,000
	Fall	2021	WS	Eggs	WS	WSU	200
	October	2020	WS	Juveniles	LW	WS	300,000
	2022	May	2021	WS	Juveniles	WS	LW
2022	May	2021	WS-Wild	Juveniles	WS	Upstream Release	85,533
	October	2021	WS	Juveniles	LW	WS	200,000
	October	2022	RB	Eggs	RB	WS	60,763
	October	2022	P	Eggs	P	WS	87,830

- **2010** – In early 2007, the water supply to egg trays at WSNFH was inadvertently shut-off and resulted in egg loss. Round Butte stock (BYs 2006 and 2007) were reared and released as juveniles at WSNFH to make up for the loss of eggs. In 2010, these fish returned as age 4 and were not included in Warm Springs broodstock.
- **2011** - Round Butte stock adults returning as ages 4 and 5 were segregated and not included in Warm Springs Broodstock (see 2010 note). WSNFH collected eggs surplus to their production needs. The resulting surplus Warm Springs stock juveniles (approximately 107,000) were marked and released as sub-yearlings into Shitike Creek in spring of 2012. No monitoring was conducted to determine the fate of Shitike releases.
- **2012** - Round Butte stock adults returning as age 5 were segregated and not included in Warm Springs broodstock (see 2010 note).
- **2013** - In 2009, the hatchery spawned Warm Springs NFH stock males (~63) with Round Butte stock females (no data on how many females). Adult returns in 2013 included these Round Butte-Warm Springs NFH stock crosses but they were not used as broodstock.
- **2014** – High broodpond mortality (70%) was due to disease outbreaks.
- **2015** - Hatchery records are incomplete/inconsistent for 2015. Records of adults transferred are different between Warm Springs NFH and Little White NFH. Juvenile records also differ. Warm Springs BY15 release numbers are estimates.
  - April – June approximately 701 Warm Springs stock adults (BY15) were transferred to Round Butte State Fish Hatchery to meet Round Butte program shortfalls.
  - Warm Springs juveniles were moved to LWSNFH due to high temperatures in the Warm Springs River.
  - In July 2015, the remaining Warm Springs stock adults were transferred from WSNFH to Little White Salmon NFH where they were spawned and the eggs moved for incubation to the Spring Creek National Fish Hatchery (SCNFH) until eye-up, and then back to the WSNFH for hatching.
  - At spawning, Little White Salmon NFH transferred 926,679 green eggs to WSNFH. There was an unexplained egg loss with an estimated mortality of 47% (~491,143 eggs survived).
  - In November of 2015, ~450,000 Warm Springs stock fertilized eggs were transferred back to WSNFH from Round Butte State Hatchery. Almost all 92% (>400,000) eggs died approximately 7-10 days after their transfer. USFWS’s Fish Health program did an investigation and write-up ([Thompson and Goodwin 2016](#)); no cause for egg loss was identified (Figure 8).






Figure 8. Transfers of Warm Springs NFH stock, April – November 2015 from Thompson and Goodwin 2016.

- **2016** - Hatchery records are incomplete/inconsistent for BY16 juveniles. Little White Salmon NFH juvenile data indicates fewer fish transferred back to Warm Springs NFH (444,313) than Warm Springs NFH says were released the following year (541,441). Warm Springs NFH release numbers are the same as the tagging numbers, no on-station

mortality data was recorded for Warm Springs NFH. Total release numbers for BY16 are likely over-estimates. Release to adult survival estimates for BY16 are unreliable.

- In the spring of 2016, ~ 130,000 Parkdale stock juveniles and ~45,000 Round Butte stock juveniles were transferred to WSNFH to supplement the 2015 egg loss (all non-WS stock fish given Left Ventral clip). In summer 2016, concerns about electrical power interruption during construction prompted the transfer of 450,000 juveniles to Little White Salmon NFH; ~90,000 juveniles remained at WSNFH. Mortality (marking to release) of fish transferred to Little White Salmon NFH was higher than the fish that remained at WSNFH.
- **2017** - Due to concerns of another year of high egg loss, the hatchery requested surplus BY17 Spring Chinook salmon eggs from Round Butte state hatchery to help them cushion expected high egg mortality. The BY17 Warm Springs NFH stock did not experience high egg mortality, and approximately 764,000 Warm Springs NFH stock juveniles were released in 2019. The additional BY17 Round Butte stock juveniles (~123,000 adipose and left ventral clipped) that were released at Warm springs NFH in 2019 resulted in the exceedance of the Warm Springs HGMP allowance. The one-time release of additional fish was determined not to have a substantial effect on ESA-listed species beyond what was considered in the USFWS BiOp covering the WSNFH program.
- **2018** - Low Warm Springs NFH stock adult returns in 2018 led to WSNFH requesting and receiving eggs from both Parkdale hatchery and Round Butte hatchery to augment hatchery production. These fish were segregated according to their source and received both an adipose clip and left ventral clip (Round Butte) or right ventral clip (Parkdale) to distinguish them from the Warm Springs stock upon their return. At release in 2020, 281,133 WSNFH stock, 135,798 Parkdale stock, and 230,310 Round Butte stock fish were released from BY18 (sr80s.dbf file).
- **2019** - Upon examination of records and better than usual survival, Round Butte State Fish Hatchery had an excess of approximately 15,000 +/- of brood year 2018 Spring Chinook salmon fingerlings. These fish were surplus to the needs and above the carrying capacity of the Round Butte facility and were made available to WSNFH. Warm Springs NFH was under its station release goal of 750,000 for brood year 2018 due to low returns of brood fish and a higher-than-normal post eyed egg loss to a single take of eggs. Warm Springs NFH was already rearing approximately 234,000 Round Butte source fingerlings (from egg transfers) on station from the same brood year. These fish were segregated from the other populations on station and received both an adipose clip and left ventral clip to remain consistent with the program for fish coming from the Round Butte source (Frejie 2019). There are no Warm Springs NFH records indicating the transfer of the ~15,000 Round Butte juveniles, and it is unclear where these fish were placed and reared at Warm Springs NFH.
- **2021**
  - May:

- Warm Springs began transferring all AD/LV Round Butte Stock (reared and released from Warm Springs NFH and returned to Warm Springs NFH) to Round Butte Hatchery to supplement their low returns.
  - Round Butte Hatchery submitted a formal request for 220 Warm Springs Stock adults. 100 (50 pair, 1:1 male:female) for Round Butte Hatchery and 120 (60 pair, 1:1 male:female) for Parkdale Hatchery for a total request of 220 fish. Warm Springs NFH is giving all Warm Springs Fish Ad/LV and AD transferred to Round Butte a dorsal fin punch to help with RMIS reporting. Adults recovered from Round Butte with CWTs must report the correct return location.
  - Hatchery Warm Springs stock fish were trucked upstream by CTWSRO to naturally spawn since all wild fish were being held at the hatchery. All outplanted hatchery adults were PIT tagged and a genetic clip was collected (either a caudal clip or caudal punch). As temperatures warmed in July, the Tribe was willing to take the risk of mortality of the fish because they would add nutrients to the river.
- June:
- The tribe had concerns about holding all wild on station until July 15th, however, USFWS wants to represent entire run in the spawning and must hold some back. Surplus wild fish above what was needed for spawning (60 females and 40 males) were be removed from the holding pond and trucked upstream on 7/14).
  - PGE inquired about passing excess hatchery brood upstream of Pelton Dam for reintroduction program. See Utilization of Excess Broodstock for Reintroduction \_2021.docx
  - One week before the record heat wave on June 26 2021, the Warm Springs HET recommended immediate transfer (within three days of decision) of fish from Warm Springs NFH to Little White Salmon NFH (LWSNFH). Fish held long-term in non-optimal conditions (i.e., chronic and acute stress) are less resistant to disease and treatment regimens are less effective. Morbidity and death will increase with longer exposure to higher water temperatures. Proactive steps considering transporting fish off-station during summer rearing conditions were developed and included in the BY20 AOP (see Appendix for criteria). With very hot temperatures forecasted for the next 10 days, uncertainty as to the July-August temperatures, and all the BY20 wild stock on-station at WSNFH, the Warm Springs HET met to consider transferring juveniles to another location.
    - – Juvenile holding conditions were averaging 65° F for several days with the long-term climate and weather predictions showing a marked increase in air and (assumed) water temperatures as well as (assumed) lower river flows.
    - – Fish were being managed for an acute bacterial infection (*C. shasta*) with topical chemicals and planned medicated feed.

- – There were 15 raceways with approximately 36K fish in each (13 raceways with hatchery or mixed hatchery/wild origin and 2 raceways with wild origin).
  - – Secondary locations were discussed for wild stock (Parkdale Hatchery, an acclimation pond at the Moving Falls Facility on the Hood River, multiple facilities, LWSNFH). After evaluating further logistics (CDL personnel availability, environmental condition, etc.), Hauling priorities were wild fish to truck first and keep separate from the hatchery stock.
- Juvenile fish were trucked to LWSNFH for the summer where temperatures would be lower. The wild fish from 2 raceways were transferred Saturday afternoon. Sunday morning the second load contained three raceways of fish. By Sunday at 10 am, the last three raceways were transported to LWSNFH. On Monday, 6/28, the Warm Springs River temperature was 77.7 F and raceways were 78.9 F. There was some mortality Monday morning, but they are looking better by Wednesday 6/30. Lessons learned are to be proactive, communicate with the Regional Office and the Tribe, and work with other hatcheries for assistance (i.e., vehicles, space, CDL drivers). There is a need to put together a protocol with past knowledge to assist the hatchery manager and the decision-making process.
- CTWSRO Requested an additional 75 (25 females and 50 males) hatchery Warm Springs NFH stock to be transferred to Round Butte to fill Parkdale shortfall. This request was approved because it was from the Tribe and because fish sent to Round Butte Hatchery are more likely to contribute returns to the Deschutes River and ultimately contribute their genetic material to the larger population (compared to fish moved above the Dam).
- Three adult fish requests and an additional request for eggs (WSU research proposal for 100-200 eggs total, green or eyed from a few different females) were approved by FWS.
- July:
  - Eighteen wild fish at Warm Springs NFH were transported upstream and released Wednesday, 7/14. As the hatchery ultrasounded and injected fish that were kept for broodstock, every 3rd female and every 4th or 5th male were not injected with antibiotics and instead moved to the truck to go upstream.
- October:
  - BY20 juveniles were transferred back to Warm Springs.
- **2022**
  - May:
    - 5/9: BY21 juvenile hatchery-origin fish were proactively trucked to LWSNFH for summer rearing where water temperatures would be lower.

- 5/17 – 5/24: The CTWSRO-BNR released the natural-origin BY21 fish in locations in the upper Warm Springs River basin instead of transporting to LWSNFH for summer rearing.
- September:
  - 9/29: BY21 juveniles were transferred back to Warm Springs.
- October:
  - Low Warm Springs NFH stock adult survival in 2022 led to WSNFH requesting and receiving eggs from both Parkdale and Round Butte Hatcheries to augment hatchery production. These fish were segregated according to their source and will receive both an adipose clip and left ventral clip (Round Butte) or right ventral clip (Parkdale) to distinguish them from the Warm Springs NFH stock upon their return and not used as broodstock.

## Other Fish counted and passed above Warm Springs NFH

The number of stray hatchery steelhead counted at the fish ladder at WSNFH increased beginning in 1987 but has decreased since 2003. From 1982 to 1986, stray hatchery steelhead composed a mean of 13.6% (range of 6.6% to 23.0%) of the total number of steelhead counted at the ladder. Between 1987 and 2003, a mean of 50.9% (range of 34.7% to 66.4%) of the steelhead counted were stray hatchery fish (Hand and Olson 2003). In more recent years, stray hatchery fish have decreased to a mean of 11.8% (range of 2% to 24%) of the total number of steelhead counted at the ladder (Table 22).

Except for steelhead, counts of other species of fish are intermittent and may not necessarily reflect total number of fish each year. Numbers of whitefish and suckers passed upstream of the ladder declined between 2018 and 2020 and have increased in recent years. Northern Pikeminnow have always been rare at the ladder. Coho numbers have been above average four of the last five years.

A Lamprey Passage Structure (LPS) was installed at the fish ladder in 2018. Since then, 87 Pacific lamprey have been observed when a total of 7 were counted in the previous seven years combined. The LPS is open and run from March to November each year with the majority of lamprey detected between mid-May and late August. In 2018, adult lamprey were PIT tagged lower in the Warm Springs River and created an opportunity to verify camera detections with known PIT tag detections at the WSNFH fish ladder PIT antenna. In 2019, the LPS was fitted with a camera system triggered by lamprey passing an infrared break beam which is the primary method of lamprey detection. That year, three PIT tagged lamprey were detected at the PIT array, all three detections had corresponding video detections on the camera system. Future PIT tagging of adult lamprey by the CTWSRO will allow for additional truing.

**Table 22. Counts of wild Steelhead, hatchery Steelhead, Fall Chinook (wild and hatchery combined), Coho (wild and hatchery combined), Rainbow Trout, Bull Trout, Whitefish, Northern Pikeminnow, Sucker, and Pacific Lamprey counted at the Warm Springs NFH fish ladder 2013 - 2022. Except for Steelhead, counts are intermittent and may not necessarily reflect total number of fish in a given year. Data retrieved from CRiS Adult Entry File: 4/11/2023**

<b>Year</b>	<b>Wild Steelhead</b>	<b>Hatchery Steelhead</b>	<b>Fall/Summer Chinook</b>	<b>Coho</b>	<b>Rainbow Trout</b>	<b>Bull Trout</b>	<b>Whitefish</b>	<b>Northern Pikeminnow</b>	<b>Sucker</b>	<b>Pacific Lamprey</b>
2013	381	55	4	90	19	9	166	2	697	1
2014	201	40	3	365	18	1	197	0	521	0
2015	373	34	0	3	17	1	783	0	471	0
2016	279	34	0	190	22	1	11	0	239	0
2017	262	8	0	264	7	0	52	0	539	0
2018	60	1	0	2	9	1	13	0	250	13*
2019	24	4	0	443	7	0	1	0	36	25*
2020	52	7	6	747	14	1	1	0	391	4*
2021	85	27	1	556	17	5	108	0	159	20*
2022	6	1	0	605	7	1	52	0	401	25*
<b>Mean</b>	<b>172</b>	<b>21</b>	<b>1</b>	<b>326</b>	<b>14</b>	<b>2</b>	<b>140</b>	<b>0</b>	<b>370</b>	<b>10</b>

\* Pacific Lamprey counted in LPS

## Past M&E Studies

- **2007** – [Comparing Two Methods Used to Mark Juvenile Chinook Salmon: Automated and Manual Marking \(Hand et al. 2010\)](#). A study compared the automated fish-marking trailer to the manual-marking trailer. The automated fish-marking trailer had higher clip quality and tag retention with no increase in rates of injury or marking to release survival.
- **2008** – [Distribution and Survival of Adult Hatchery Spring Chinook Salmon Radio-Tagged and Released Upstream of Warm Springs NFH in 2008 \(Conder et al. 2010\)](#). During the spring and summer of 2008, 35 hatchery-origin spring Chinook salmon were radio-tagged and released upstream of the hatchery. We studied their movement patterns, identified potential holding areas, estimated survival, and approximated their contribution to spawning. Based on tag movements during the spawning period, 60% of the tagged fish survived to spawning and 31% of the radio-tagged hatchery fish contributed to natural spawning.
- **2008** – [Use of Parentage Analysis to Determine Reproductive Success of Hatchery-Origin Spring Chinook Salmon Outplanted into Shitike Creek, Oregon \(Baumsteiger et al. 2008\)](#). In 2002 and 2003, 83 and 265 adult hatchery salmon, respectively, were outplanted into Shitike Creek. The number of (juvenile) offspring attributed to an individual (adult) outplant was variable, ranging from 1 to more than 10. This study shows that under the right conditions, outplanted adult hatchery fish taken from localized hatchery stocks can contribute to the overall juvenile production in a natural stream. Outplanting adult salmon from Warm Springs NFH into Shitike Creek continued through 2005 ([Hand et al. 2005](#)).
- **2010** – [Feasibility of live spawning wild male spring Chinook salmon at Warm Springs NFH, 2010 Report \(Hand et al. 2014b\)](#). We evaluated the feasibility of using live-spawned wild males to provide a genetic contribution to both the hatchery broodstock and natural production by live-spawning five wild males and releasing the fish back into the Warm Springs River. It appeared that live-spawning of wild males may be a feasible method to include wild genetics into the hatchery broodstock while not compromising the overall wild production.
- **2012** – [Effectiveness of an integrated hatchery program: Can genetic-based performance differences between hatchery and wild Chinook salmon be avoided? \(Hayes et al. 2013\)](#). The authors evaluated the performance of fish from hatchery, wild, and crossed populations in hatchery and stream environments. Hatchery fish performed differently than wild fish possibly because they were accustomed to rearing at higher densities in a hatchery setting (domestication) leading to genetic divergence. Future studies are needed to evaluate which hatchery techniques are most useful for reducing performance differences and reducing risk to wild populations.
- **2013** – [An Evaluation of Rearing Densities to Improve Growth and Survival of Hatchery Spring Chinook Salmon \(Olson and Paiya 2013\)](#). For three consecutive brood years (BY2000-02), density treatments consisted of low, medium, and high groups in 57.8-m<sup>3</sup> raceways with approximately 16,000, 24,000, and 32,000 fish/raceway, respectively. Fish

reared at high density exhibited the highest on-hatchery mortality rate during two brood years; however, differences in mortality rate among densities were not significant ( $P = 0.20$ ). In one brood year, adult recovery rates appeared to support the hypothesis that lower initial densities improved post-release survival ( $P < 0.01$ ). All rearing densities utilized in this evaluation were relatively low and may partially explain why more differences were not readily apparent among density groups.

- **2014** – [Pacific lamprey and Bull Trout passage assessment at Warm Springs NFH \(Gallion and Skalicky 2014\)](#). An evaluation at the hatchery indicated significant passage deficiencies for Pacific lamprey which likely delay and limit passage through the fishway. Passage limitations for bull trout through the fishway were not as significant.
- **2014** – [Genetic Composition of the Warm Springs River Chinook Salmon Population Maintained Following Eight Generations of Hatchery Production \(Smith et al. 2014\)](#). The genetic characteristics of the endemic population was examined before (1976–1977) and after (2001–2011) hatchery became operational. Natural-origin Chinook Salmon changed very little over the eight generations. However, differences between hatchery- and natural-origin fish are expected to increase if hatchery operations do not integrate natural-origin fish and incorporate Round Butte Hatchery fish into the broodstock.
- **2014** – [Adult Recovery of Hatchery Spring Chinook Salmon Adipose Fin-Clipped and Coded-Wire-Tagged Using an Automated and Manual Marking Trailer \(Hand et al. 2014a\)](#). At WSNFH, the adult recovery rate for fish marked in the automated trailer was 0.16%, compared with a recovery rate of 0.14% for fish marked in the manual trailer. A fish was 1.17 times more likely to be recovered as an adult at the hatchery if marked in an automated trailer.
- **2015**– [Migratory Behavior of Chinook Salmon Microjacks Reared in Artificial and Natural Environments \(Hayes et al. 2015\)](#). Emigration was evaluated for hatchery age-1 mature males and immature parr. Mature age-1 fish were significantly longer, heavier, and had greater condition factor. These mature age-1 male fish have the potential to contribute to the spawning population but can also represent a loss of productivity.
- **2016** – [Migration Timing and Survival of Warm Springs NFH Juvenile Spring Chinook Salmon in the Deschutes Basin \(Davis et al. 2016\)](#). In 2012, 2013 and 2014, radio-telemetry was used to evaluate where the majority of spring Chinook mortalities occur. Median travel time from WSNFH to Bonneville Dam was 27 days compared to a two-day travel time to the mouth of the Deschutes, suggesting the rate of travel slows from an average 70 rkm/day to 3.5 rkm/day when fish enter the Columbia River.
- **2018** – [Evaluation of adult Pacific Lamprey upstream passage at Warm Springs National Fish Hatchery, 2017 Annual Report \(Barkstedt and Johnsen 2018\)](#). A previous evaluation of both physical structures and adult lamprey passage determined that the barrier dam and fish ladder impeded lamprey upstream migration (Gallion and Skalicky 2014). The Confederated Tribes of Warm Springs Reservation and the USFWS collaborated to design, install, and monitor a LPS. The LPS was installed in 2017, began operation in 2018, and successfully provided passage for 13 adult Pacific Lamprey in its first year



- **2018** – Evaluation of on-hatchery and off-hatchery performance of WSNFH stock, Round Butte stock, and Parkdale stock juveniles reared and released from Warm Springs NFH. A report will be available in fall 2023 after all adults have returned.

## Summary and Future Studies

The WSNFH produces spring Chinook Salmon for tribal harvest in the Deschutes and Columbia Rivers, for on-reservation distribution to tribal members, and for sport fishery. The program is currently 31% below the goal of producing 750,000 juveniles for release. The primary issues related to meeting release goals has been maintaining broodstock health after they have returned to the hatchery, achieving eye-up, and survival until marking. The smolt-to-adult survival rate varies annually and has exceeded its goal five times in the last ten years.

Warm Springs NFH juvenile releases have changed over time and are dependent on environmental and hatchery factors. Since 1991, the spring releases have ranged from March 27 to April 27 (April 3 on average). All the juveniles have been successfully marked with a CWT and adipose fin clip, non-WSNFH stocks have been differentially marked with a ventral clip, and a subsample are PIT Tagged before release.

During the juvenile fish downstream migration season (March to late summer), the Columbia River hydropower system operations are modified to improve in-river conditions for migrating fish. One modification is to spill water and juvenile fish over dam spillways, instead of putting the water through the turbines. Spring spill dates for McNary, John Day, The Dalles, and Bonneville Dams start April 10. Based on PIT tag data since BY05, the fastest hatchery releases reach Bonneville dam in approximately 4 days. These fish likely pass The Dalles Dam 1-2 days prior to reaching Bonneville Dam (see Davis 2016 for data on Deschutes River migration). If the fundamental objective of the hatchery release is to maximize the likelihood of hatchery releases passing through mainstem spillways instead of turbines, the Service recommends hatchery releases should start no more than three days prior to spill (April 7). A less conservative approach, 90% of the fish passing the mainstem dams during spill, would be to start hatchery releases no more than 8 - 9 days prior to spill.

Wild and hatchery fish return to the Warm Springs River from late April through September and are spawned from late August through September dependent on environmental and hatchery factors. Most wild and hatchery fish return to the Warm Springs River by late June. Hatchery spawning has begun as early as August 10 and as late as September 24, the average first date of spawning is August 22. In 2019, the first date of spawning was September 5<sup>th</sup>, four days after the previous latest start of spawning in 2010. Higher intensity daylight LED lights were installed over each pond and simulate the day length needed to cue future spawning. All fish have been spawned between August 31 and September 13 since 2019.

The facility has produced a mean hatchery smolt-to-adult survival rate that can exceed the target SAR of 0.39% at times but it is variable year to year (mean = 0.42 [95% CI 0.22, 0.61]). Due to low wild fish returns (<1,000 fish), wild fish were not regularly incorporated into the hatchery broodstock. The threshold 1,000 returning wild fish was met only once since 2004, so the program has effectively been operated under a segregated paradigm for several generations. Beginning in 2020, the wild fish supplementation program provided an opportunity to

incorporate wild fish into the hatchery broodstock thus improving the hatchery's ability to maintain wild fish genetic characteristics in the hatchery population and minimize genetic divergence between Warm Springs River hatchery and wild fish.

The 2023 forecast for WSNFH stock adult returns (model estimates of 1,485 to 2,224 Age 4 and 5 fish) indicate that it is likely that the hatchery's broodstock needs will be met. Round Butte Hatchery may also meet broodstock needs with model estimates of 1,710 to 2,468 Age 4 and 5 fish. While the wild fish forecast estimate is an improvement over 2020 -2022 (170 - 410 Age 4 and 5 fish), it is however, still concerning and warrants close monitoring and discussions of potential emergency actions. All the forecasts have a high degree of uncertainty, which will necessitate in-season monitoring and readiness to adjust management plans.

To make up for insufficient eggs in 2018, WSNFH received spring Chinook eggs and juveniles from both the Round Butte State Fish Hatchery and CTWSRO's Parkdale Hatchery. These fish were released in 2020 and an evaluation of their on-hatchery and off-hatchery performance will provide additional information on the similarities and differences between these three stocks. Juvenile Parkdale and Round Butte sourced fish released from WSNFH are differentially marked (left or right ventral fin) to distinguish them from Warm Springs broodstock in subsequent years. Ventrally marked fish are excluded from spawning with Warm Springs stock but can still be inadvertently spawned with Warm Springs stock if the ventral fin grows back. For BY18, there was an overall 11% error rate (84 errors of 769 fish) when identifying stocks by fin marks; 70% of the errors (59 of 84) were due to the regeneration of ventral fins (i.e., Round Butte and Parkdale stocks misidentified as Warm Springs stock). Inadvertent inclusion in the hatchery broodstock may increase as the number and frequency of transfers from outside the Warm Springs population increases, potentially posing a genetic risk to the Warm Springs stock (Smith 2018). The majority of adults from all stocks return to the facility at Age-4. However, the percentage of fish that return to the hatchery as adults is significantly greater for the Warm Springs stock. For BY18, the Warm Springs stock returned at a rate 57% greater than the Round Butte stock and 88% greater than the Parkdale stock. Although, no difference in survival of ventrally marked versus unmarked fish at Warm Springs NFH (Olson and Cates, 1999), some research suggests removal of a ventral fin may reduce survival in other salmonids. Fin removal activities increase physiological stress (Sharpe et al. 1998) or may lead to reduced long-term survival (Nicola and Cordone 1973). Future transfers are contingent upon availability and only after consultation and concurrence of CTWSRO and the Service.

Other species of fish collected at the WSNFH fish ladder include wild steelhead, hatchery steelhead, fall Chinook (wild and hatchery), coho (wild and hatchery), rainbow trout, bull trout, whitefish, northern pikeminnow, sucker, and Pacific lamprey. These fish are counted and passed upstream or made available to the CTWSRO. Wild fish counts at WSNFH of summer steelhead have declined and is of concern. The Warm Springs NFH trap and ladder are being monitored to evaluate if it is functioning properly or if there are any infrastructure problems that could cause migrational delay.

## Future M&E Studies

- Annual run reconstruction of wild and hatchery spring Chinook salmon
- Collect data for population monitoring of ESA listed summer steelhead and bull trout
- Monitor other fish passing the hatchery site,

- Rearing and release studies at the hatchery to improve performance,
  - Diet
  - Growth
  - Reduced rearing densities
  - Fish health evaluations
- Explore funding available to continue developing collaborative projects with our partners, especially CTWSRO
- Evaluate performance and ecological interactions of hatchery and wild fish
- Evaluate & implement projects and/or facilities to reduce high water temperature during late spring to early fall juvenile rearing at the hatchery
- Evaluate on-hatchery and off-hatchery performance of BY18 Warm Springs, Round Butte, and Parkdale stocks reared at WSNFH

### Acknowledgements

Data used in this report was downloaded from CRiS maintained at the CRFWCO, RMIS, and PTAGIS. Hatchery personnel at WSNFH collected data on release dates, adult returns, and annual number of juveniles released from the facilities. Marking and biosampling crews from the USFWS documented the number of adult returns as well as adipose fin-clipped, coded-wire tagged, and PIT-tagged juveniles prior to release. Funding for M&E of this hatchery program was provided by the USFWS.

## Appendix A

A summary of on-station hatchery notes for 2022 (i.e., feeding changes, ponding, unique issues, etc.)

- Infrastructure/Tools
  - All adult fish were sexed with the ultrasound unit
  - The SCADA system for the chiller was adjusted to monitor incoming water temperature instead of outgoing water. The set points were changed to maintain cooler water with fewer fluctuations.
  - Security Lighting was possibly left on each night 2020, 2021, and 2022. The switch was turned off in late summer 2022
- Fish Management
  - BY20 Juveniles released early on March 9<sup>th</sup> because due to clinical BKD outbreak. Less than 3,500 fish PIT tagged before release.
  - BY20 Wild fish were released on station instead of upstream to prevent BKD affecting BY21 fish downstream at the hatchery.
  - BY21 Wild fingerlings were released early into upstream tributaries instead of being moved to Little White Salmon NFH for the summer.
  - BY21 fingerlings (hatchery, hatchery wild mix-marked, hatchery wild mix-unmarked) moved to Little White Salmon NFH for the summer to proactively avoid high water temperature.
- Fish Health
  - Formalin barrel was empty and pump malfunctioned, brood ponds were not treated.
  - Draxxin injections were done all at once for the wild fish in the spawning building in July. Fish were held in the cross channel while brood pond was drained and refilled. Cross channel had warmer water and caused additional stress on the fish.
  - There were no issues with ich.
  - Were any batches of BY22 eggs was culled due to BKD?
- Policy
  - The *US v OR* Policy Committee approved the change to the production table to incorporate the wild supplementation program into the existing hatchery program, and to footnote the transfer of ~560,000 juveniles to Little White Salmon during the summer.
- Temperature monitoring
  - Loggers were installed at the three water temperature monitoring sites on 10/31/2022 at the lamprey pump intake in the WSNFH building, hatchery intake in Warm Springs River and hatchery outfall in Warm Springs River. The hatchery outfall loggers were installed in the hatchery outfall structure (44.862323, -121.244032 Google Maps) instead of the Warm Spring River due to access issues. The ambient air temperature logger was installed on the opposite side of the building (44.860996, -121.246136 Google Maps) instead of at the

intake and should provide a better representation of local air temperature. -  
Kristofor Kannarr, Hydrologist, Water Resources Branch

## References

- Archibald, J. 2013. Warm Spring Fish Passage Evaluation Report 2013. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office, Vancouver, WA.
- Barkstedt, J., and A. Johnsen. 2018. Evaluation of adult Pacific Lamprey upstream passage at Warm Springs National Fish Hatchery, 2017 Annual Report. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office, Vancouver, WA.
- Baumsteiger, J., D. M. Hand, D. E. Olson, R. Spateholts, G. FitzGerald, and W. R. Ardren. 2008. [Use of Parentage Analysis to Determine Reproductive Success of Hatchery-Origin Spring Chinook Salmon Outplanted into Shitike Creek, Oregon](#). North American Journal of Fisheries Management.
- Bureau of Indian Affairs (BIA). 1967. Business Lease 3552. Recorded Document Number 145-8165. Warm Springs Indian Reservation of Oregon
- Cates, B. 1992. Warm Springs National Fish Hatchery evaluation and anadromous fish study on the Warm Springs Indian Reservation of Oregon 1975-1989. United States Fish; Wildlife Service, Lower Columbia River Fishery Resource Office, Vancouver, WA.
- Conder, T., D. M. Hand, D. E. Olson, and J. Lovtang. 2010. Distribution and Survival of Adult Hatchery Spring Chinook Salmon Radio-Tagged and Released Upstream of Warm Springs National Fish Hatchery in 2008. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office, Vancouver, WA.
- Confederated Tribes of the Warm Springs Reservation of Oregon, and U.S. Fish and Wildlife Service (USFWS). 2007. Warm Springs National Fish Hatchery: Operational Plan and Implementation Plan 2007-2011.
- Davis, B., J. Poirier, and D. M. Hand. 2016. Migration Timing and Survival of Warm Springs National Fish Hatchery Juvenile Spring Chinook Salmon in the Deschutes Basin, 2012-2014 final Report. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office, Vancouver, WA.
- Frejje, T. 2019. National Fish Hatchery Planned Release, Transfer Schedule or Research Request, 3/20/2019.
- Gallion, D., and J. Skalicky. 2014. Pacific Lamprey and Bull Trout Passage Assessment at Warm Springs National Fish Hatchery. Page 43. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office, Vancouver, WA.
- Hand, D. M., W. R. Brignon, and D. E. Olson. 2014a. [Adult Recovery of Hatchery Spring Chinook Salmon Adipose Fin-Clipped and Coded-Wire-Tagged Using an Automated and Manual Marking Trailer](#). North American Journal of Aquaculture.
- Hand, D. M., W. R. Brignon, D. E. Olson, and J. Rivera. 2010. [Comparing Two Methods Used to Mark Juvenile Chinook Salmon: Automated and Manual Marking](#). North American Journal of Aquaculture.

- Hand, D. M., T. Conder, D. E. Olson, and J. Lovtang. 2014b. Feasibility of live spawning wild male spring Chinook salmon at Warm Springs National Fish Hatchery, 2010 Report. U.S. Fish & Wildlife Service, Columbia River Fisheries Program Office.
- Hand, D. M., and D. E. Olson. 2003. Steelhead Returns to Warm Springs National Fish Hatchery, 1978-2003.
- Hand, D. M., D. E. Olson, R. O. Engle, T. A. Hoffman, R. Spateholts, and G. FitzGerald. 2005. Distribution, Behavior, and Reproductive Success of Outplanted Hatchery Spring Chinook Salmon in Shitike Creek, OR. Progress Report for 2004 and Work Plan for 2005. Confederated Tribes of the Warm Springs Reservation of Oregon; U.S. Fish & Wildlife Service.
- Hand, David M. and Haeseker, Steven L. 2011. Retrospective Analysis of Preseason Run Forecast Models for Warm Springs stock Spring Chinook Salmon in the Deschutes River, Oregon. U.S. Fish & Wildlife Service.  
[https://www.fws.gov/CRFWCO/publications/Analysis\\_of\\_Warm\\_Springs\\_Stock\\_Spring\\_Chinook\\_Salmon\\_Forecasting\\_Methods.pdf](https://www.fws.gov/CRFWCO/publications/Analysis_of_Warm_Springs_Stock_Spring_Chinook_Salmon_Forecasting_Methods.pdf).
- Hayes, M. C., R. R. Reisenbichler, S. P. Rubin, D. C. Drake, K. D. Stenberg, and S. F. Young. 2013. [Effectiveness of an integrated hatchery program: Can genetic-based performance differences between hatchery and wild Chinook salmon be avoided?](#) Canadian Journal of Fisheries and Aquatic Sciences.
- Hayes, M. C., S. P. Rubin, R. R. Reisenbichler, and L. A. Wetzel. 2015. [Migratory behavior of chinook salmon microjacks reared in artificial and natural environments.](#) Journal of Fish and Wildlife Management.
- Integrated Hatchery Operations Team (IHOT). 1995. Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries, Annual Report 1994, Report to Bonneville Power Administration, Contract No. 1992BI60629, Project No. 199204300. Pages 119 electronic pages (BPA Report DOE/BP-60629).
- Lovtang, J., M. W. Gauvin, and D. M. Hand. 2011. Spring Chinook Salmon in the Deschutes River, Oregon, Wild and Hatchery. 1975-2009 returns and 2010 Run Size Forecasts. Confederated Tribes of the Warm Springs Reservation of Oregon, Oregon Department of Fish & Wildlife, U.S. Fish & Wildlife Service, Warm Springs, OR.
- National Marine Fisheries Service (NMFS). 2007. Section 7 Biological Opinion USFWS Artificial Propagation Programs in the Lower Columbia River and Middle Columbia River.
- National Marine Fisheries Service (NMFS). 2018. Section 7 Biological Opinion and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Response.
- Nicola, S. J. and A. J. Cordone, 1973 Effects of fin removal on survival and growth of rainbow trout (*Salmo gairdneri*) in a natural environment, Transactions of the American Fisheries Society, 102, Pages 753– 758.



- Olson D.E., and B.C. Cates. 1999. Differential Performance Of Ventral Fin Clipped And Adipose Fin Clipped/Coded-Wire Tagged Spring Chinook Salmon At Warm Springs National Fish Hatchery, Oregon
- Olson, D. E., and M. Paiya. 2013. [An evaluation of rearing densities to improve growth and survival of hatchery spring Chinook salmon.](#)
- Olson, D. E., and R. Spateholts. 2001. Hatcheries, Harvest and Wild Fish: An Integrated Program at Warm Springs National Fish Hatchery, Oregon. Pages 23–42 52nd annual pacific northwest fish culture conference.
- Olson, D. E., R. Spateholts, M. Paiya, and D. E. Campton. 2004. Salmon hatcheries for the 21st century: A model at Warm Springs National Fish Hatchery. Pages 585–602 in M.J. Nickum, P. M. Mazik, J. G. Nickum, and D. D. MacKinlay, editors. Propagated fish in resource management Symposium. American Fisheries Society, Bethesda, Maryland.
- Reagan, R. E., N. S. Adams, D. W. Rondorf, G. FitzGerald, R. Spateholts, T. A. Hoffman, and D. E. Olson. 2005. Distribution, Migration Behavior, Habitat Use, and Species Interactions of Fall-Released Juvenile Hatchery Spring Chinook Salmon in the Deschutes River, Oregon, 2003 Annual Report for 2003 Columbia River Publications. Columbia River Publications web page.
- Sharpe, C. S., D. A. Thompson, H. L. Blankenship and C. B. Schreck, 1998 Effects of routine handling and tagging procedures on physiological stress responses in juvenile chinook salmon, *Progressive Fish-Culturist*, 60, Pages 81– 87.
- Smith, C. T. 2018. Memo to Mike Clark, Complex Manager, Columbia River Gorge NFH Complex. June 29, 2018.
- Smith, C. T., R. French, J. Lovtang, and D. M. Hand. 2014. [Genetic Composition of the Warm Springs River Chinook Salmon Population Maintained Following Eight Generations of Hatchery Production.](#) Transactions of the American Fisheries Society.
- Smith, C. T., M. Smith, and D. M. Hand. 2016. Population genetic analysis of Parkdale Fish Hatchery Spring Chinook Salmon Final Report. Abernathy Fish Technology Center. Longview, Washington
- Thompson, D., and A. Goodwin. 2016. An Investigation into the Cause of Egg Mortality at Warm Springs National Fish Hatchery during the fall of 2015. U.S. Fish & Wildlife Service.
- U.S. Fish and Wildlife Service (USFWS). 1995. Fish Health Policy and Implementation, 713 FW. U.S. Fish & Wildlife Service Manual (FWM).
- U.S. Fish and Wildlife Service (USFWS). 2004. Aquatic Animal Health Policy. <https://www.fws.gov/policy/713fw1.pdf>.
- U.S. Fish and Wildlife Service (USFWS). 2006. Columbia River Basin, Columbia Plateau Province, Deschutes River Watershed, Warm Springs National Fish Hatchery



Assessments and Recommendations Final Report. U.S. Fish & Wildlife Service, Portland, Oregon. <http://www.fws.gov/pacific/fisheries/Hatcheryreview/team.html>.

U.S. Fish and Wildlife Service (USFWS). 2019. Warm Spring National Fish Hatchery 2019 Draft Annual Operational Plan. U.S. Fish & Wildlife Service, Portland, Oregon.