Bayou Sauvage National Wildlife Refuge

Bird list

This blue goose, designed by J.N. "Ding" Darling, has become a symbol of the National Wildlife Refuge System.

Bayou Sauvage National Wildlife Refuge consists of approximately 25,000 acres and is located entirely within the city limits of New Orleans, making it the nation's largest urban refuge. The refuge contains a variety of wildlife habitat, including bottomland hardwoods, freshwater and brackish marshes, lagoons, canals, borrow pits, and natural bayous, meeting the needs of 340 bird species during various seasons of the year.

For many birds, Bayou Sauvage is the last stop over on their journey along the Mississippi Flyway before continuing on over the Gulf of Mexico. During fall, winter, and summer months, migratory waterfowl populations can exceed 75,000. Hurricane Katrina severely impacted habitat for neotropical migrants, however, with the restoration of the forested ridge, ideally, these birds will once again make Bayou Sauvage part of their annual journey.

Wading birds are present year round and mottled ducks and wood ducks are common summer nesters. The American bald eagle and peregrine falcon can also be found here. The following list contains 255 species, representing observations on and adjacent to Bayou Sauvage National Wildlife Refuge. Sightings of unlisted species should be reported to: Refuge Manager, BS NWR, 61389 Highway 434, Lacombe, Louisiana 70445.

This list conforms to the nomenclature and arrangement established by the American Ornithologists' Union (1998).

Symbols used in this brochure represent the following:

Sp Spring (March-May)

F Fall (September-November)

S Summer (June-August)

W Winter (December-February)

- c Common (certain to be seen in suitable habitat)
- u Uncommon (present but not certain to be seen)
- o Occasional (seen only a few times during a season)
- r Rare (seen at intervals of 2 to 5 years)

Five primary habitats common to the refuge are identified as follows:

F Field O Open Water f Forest M Marsh

A Aerial Flyover

New Species ^ Species of Special Concern *

Bayou Sauvage National Wildlife Refuge 61389 Hwy 434 Lacombe, LA 70445 985/882 2000

U.S. Fish & Wildlife Service 1 800/344 WILD

September 2013







	Sp	Su	F	W	Habitat		Sp	Su	F	W	Habitat
Ducks, Geese, Swans (Anatidae)American Wigeon	U	U	\mathbf{C}		M	New World Vultures (Cathartidae)Turkey Vulture	$_{\mathrm{C}}^{\mathrm{C}}$	$_{\mathrm{C}}^{\mathrm{C}}$	$_{\mathrm{C}}^{\mathrm{C}}$	C C	A
Black-bellied Whistling-Duck ^	\mathbf{C}	\mathbf{C}	0	О	M	Black Vulture	C	C	C	C	A
Blue-winged Teal	\mathbf{C}	R	\mathbf{C}	\mathbf{C}	M	Hawks, Kites, Eagles, and					
Bufflehead	U	D	0	С	M	Allies (Accipitridae) Bald Eagle*	С	C	\mathbf{C}	\mathbf{C}	A
Canada Goose Canvasback	R R	R R	$_{ m C}^{ m R}$	R	М М,О	Broad-winged Hawk	Ü	U	U	C	A
Common Goldeneye	10	10	O	O	M,O	Cooper's Hawk*	Ŭ	U	Ŭ	U	A
Gadwall	U		U	\mathbf{C}	M	Mississippi Kite	C	\mathbf{C}	U		A
Greater Scaup				\mathbf{R}	M,O	Northern Harrier	С	TT	С	С	A
Greater White-fronted Goose	0		0	0	M M	Osprey* Red-shouldered Hawk	$_{ m C}^{ m U}$	U C	$_{ m C}^{ m U}$	U C	A A
Green-winged Teal Hooded Merganser*	0	R	\mathbf{U}	$^{\mathrm{C}}$	M	Red-tailed Hawk	$\overset{\circ}{\mathrm{C}}$	C	$\overset{\circ}{\mathrm{C}}$	C	A
Lesser Scaup	Ŭ		Ŭ	$\tilde{\mathrm{C}}$	M,O	Sharp-shinned Hawk	U		U	Ū	A
Mallard	U		U	\mathbf{C}	M	Swallow-tailed Kite*	U	U			A
Mottled Duck	С	\mathbf{C}	С	C	M	Falcons (Falconidae)					
Northern Pintail Northern Shoveler	U O		$_{\mathrm{O}}^{\mathrm{U}}$	$^{\mathrm{C}}$	M M	American Kestrel*	\mathbf{C}		\mathbf{C}	\mathbf{C}	A
Red-breasted Merganser	Ö		ŏ	$\ddot{\mathrm{C}}$	M,O	Merlin	U		U	U	A
Redhead				\mathbf{C}	M,O	Peregrine Falcon*			R	R	A
Ring-necked Duck	R		R	\mathbf{C}	M,O	Rails, Gallinules, and Coots (Rallidae))				
Ruddy Duck Snow Goose	R U		R U	C O	M,O M	Clapper Rail ^	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	M
Wood Duck	Ö	O	Ö	Ö	M	King Rail	$\stackrel{\text{C}}{\sim}$	$\stackrel{\text{C}}{\sim}$	$\stackrel{\mathbf{C}}{\sim}$	$\stackrel{\text{C}}{\sim}$	M
	Ŭ			Ü	-1-	Virginia Rail Sora	$_{ m C}^{ m C}$	\mathbf{C}	$^{ m C}_{ m C}$	$_{\mathrm{C}}^{\mathrm{C}}$	M M
Loons (Gaviidae)	0		0	C	МО	Black Rail*	C		C	C	M
Common Loon	0		O	\mathbf{C}	M,O	Yellow Rail*				$\check{\mathrm{C}}$	M
Grebes (Podicipedae)						Purple Gallinule ^	R	R	-	-	M
Eared Grebe	R	R	\mathbf{C}	~	M,O	American Coot	$_{\mathrm{C}}^{\mathrm{O}}$	O C	$^{ m C}_{ m C}$	C C	M M
Pied-billed Grebe Horned Grebe	C O	R	$_{ m R}^{ m C}$	C	M,O M,O	Common Moorhen	C	C	C	C	1/1
norned Grebe	U		11	C	W1,O	Plovers (Charadriidae)					
Pelicans (Pelicanidae)	~	~	~		3.5.0	Black-bellied Plover Wilson's Plover ^*	0		O	R	M
American White Pelican Brown Pelican*	C C	$^{ m C}_{ m C}$	$_{\mathrm{C}}^{\mathrm{C}}$	\mathbf{C}	M,O O	Wilson's Plover Semipalmated Plover	R O		0	M R	M
	C	C	C	C	U	Killdeer	$\ddot{\mathrm{C}}$	\mathbf{C}	$\ddot{\mathrm{C}}$	C	F
Cormorants (Phalacrocoracidae)	a	~	~	~	0						
Double-crested Cormorant	С	С	С	С	0	Stilts (Recurvirostridae) Black-necked Stilt	\mathbf{C}	\mathbf{C}	С	U	M
Darters (Anhinga)						Black-necked Still	C	C	C	U	1/1
Anhinga	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	0	Sandpipers, Phalaropes, and					
Frigatebirds (Fregatidae) Magnificent Frigatebird	0	O	0	O	A	Allies (Scolopacidae) American Woodcock*	0		O	O	M,F
wagimicent i rigatebiru	O	O	O	O	11	Dunlin	U		U	U	M
Bitterns, Herons, and Allies (Ardeidae		ъ	ъ		3.5	Greater Yellowlegs	$\ddot{\mathrm{C}}$		Ŭ	Ü	M
American Bittern* Black-crowned Night-Heron	R O	R U	$_{ m R}$	R	M M	Least Sandpiper	U		U	U	M
Cattle Egret	Ü	Ü	U	U	M	Lesser Yellowlegs Long-billed Dowitcher	$_{\mathrm{O}}^{\mathrm{U}}$		\mathbf{U}	U R	M M
Great Blue Heron	$\tilde{\mathrm{C}}$	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{M}	Pectoral Sandpiper	0		Ö	п	M
Great Egret	\mathbf{C}	$\stackrel{ ext{C}}{\sim}$	\mathbf{C}	\mathbf{C}	M	Red Knot ^*	$\ddot{ m R}$		$\ddot{ m R}$		M
Green Heron Least Bittern*	R O	C U	R O	R	M M	Ruddy Turnstone ^	\mathbf{R}		R	\mathbf{R}	M
Little Blue Heron	U	C	U	U	M	Sanderling	R		0	0	M
Reddish Egret*	$\ddot{\mathrm{R}}$	\mathbf{R}	$\ddot{\mathrm{R}}$	\mathbf{R}	\mathbf{M}	Semipalmated Sandpiper* Short-billed Dowitcher	0		0	R	M M
Snowy Egret	C	C	C	\mathbf{C}	M	Solitary Sandpiper*	ŏ		ŏ	10	M
Tricolored Heron	U O	$_{\mathrm{C}}^{\mathrm{C}}$	$_{\mathrm{O}}^{\mathrm{U}}$	U	M M	Spotted Sandpiper	U		U	U	M
Yellow-crowned Night-Heron	U	C	U		1/1	Stilt Sandpiper ^ Western Sandpiper	R U		R U	U	M M
Ibises and Spoonbills						Willet	$\stackrel{ m O}{ m R}$		C	C	M
(Threskiornithidae) White Ibis	С	C	\mathbf{C}	\mathbf{C}	M	Wilson's Snipe ^	Ü		Ŭ	$\ddot{\mathrm{C}}$	M,F
Glossy Ibis*	U	U	U	C	M	Gulle Torne and Chimmara (Laridae)					
White-faced Ibis	Ü	U	U	U	M	Gulls, Terns, and Skimmers (Laridae) Bonaparte's Gull	0		0	О	A
Roseate Spoonbill*		O	O	U	M	Bonaparte 8 Gun Laughing Gull	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	A,M
						Ring-billed Gull	U	\mathbf{R}	U	\mathbf{C}	Á,F
						—Herring Gull Least Tern*	0	R C	\mathbf{U}	U	A A
						ICast ICIII	J	O	J		А

	Sp	Su	F	W	Habitat		Sp	Su	F	W	Habitat
Gull-billed Tern*	\mathbf{R}	\mathbf{R}	R	R	A,M	Jays and Crows (Corvidae)	_				
Caspian Tern*	U	U	U	U	A	Blue Jay American Crow	$_{\mathrm{C}}^{\mathrm{C}}$	$_{ m C}^{ m C}$	C	$_{\mathrm{C}}^{\mathrm{C}}$	$\mathbf{f} \\ \mathbf{f}$
Black Tern Common Tern	O R		$_{ m R}^{ m U}$	$_{ m R}^{ m U}$	A,M A	Fish Crow	C	C	$_{\mathrm{C}}^{\mathrm{C}}$	C	f, M
Forster's Tern	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	M,A						_,
Royal Tern Sandwich Tern ^	U R	$_{ m R}^{ m C}$	$_{ m R}^{ m C}$	$_{ m R}^{ m C}$	A M,A	Larks (Alaudidae) Horned Lark				R	F
Black Skimmer*	R R	R R	к R	к R	M,A M,A					10	1
					,	Swallows (Hirundinidae) Northern Rough-winged					
Pigeons and Doves (Columbidae) Rock Pigeon	O	0	0	0	F	Swallow	U	U	U		\mathbf{f}
Eurasian Collared-Dove	Ŭ	$\overset{\circ}{\mathrm{C}}$	U	U	\mathbf{F}	Purple Martin	$\stackrel{ ext{C}}{\sim}$	\mathbf{C}	U	~	f,A
White-winged Dove ^	C	C	0	O C	F F	Tree Swallow Bank Swallow	$_{ m U}^{ m C}$		C U	\mathbf{C}	A A
Mourning Dove Common Ground-Dove*	C R	$_{ m R}^{ m C}$	$_{ m R}^{ m C}$	R	F	Barn Swallow	Č	\mathbf{C}	$\overset{\circ}{\mathrm{C}}$		A
					-	Cliff Swallow	O		O		A
Cuckoos (Cuculidae) Yellow-billed Cuckoo	U	U	U		\mathbf{f}	Chickadees and Titmice (Paridae)					
Black-billed Cuckoo	$\overset{ m O}{ m R}$	$\overset{\circ}{\mathrm{R}}$	$\overset{ m O}{ m R}$		f	Carolina Chickadee	C	\mathbf{C}	C	C	\mathbf{f}
Typical Owls (Strigidae)						Tufted Titmouse	С	C	С	С	\mathbf{f}
Eastern Screech-Owl	U	U	U	U	\mathbf{f}	Nuthatches (Sittidae)					
Great Horned Owl	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	f	Red-breasted Nuthatch	a	R C	$_{ m C}^{ m R}$	C	$\mathbf{f}\\ \mathbf{f}$
Barred Owl	\mathbf{C}	С	С	С	\mathbf{f}	Brown-headed Nuthatch*	\mathbf{C}	C	C	C	Ι
Barn Owls (Tytonidae)						Wrens (Troglodytidae)				~	
Barn Owl	U	U	U	U	\mathbf{F}	—_House Wren Winter Wren	$_{ m O}^{ m U}$		$_{\mathrm{O}}^{\mathrm{U}}$	$_{ m R}^{ m C}$	$\mathbf{f} \\ \mathbf{f}$
Goatsuckers (Caprimulgidae)						Sedge Wren*	Ŭ		Ŭ	U	M
Common Nighthawk	\mathbf{C}	\mathbf{C}	\mathbf{C}		f,F	Marsh Wren	\mathbf{C}	~	C	C	M
Chuck-will's-widow*	U	U	U		\mathbf{f}	Carolina Wren	\mathbf{C}	C	\mathbf{C}	\mathbf{C}	\mathbf{f}
Swifts (Apodidae)	~	~	~			Kinglets (Regulidae)	С		С	С	${f f}$
Chimney Swift	С	С	С		A	Ruby-crowned Kinglet	C		C	C	1
Hummingbirds (Trochilidae)			~			Gnatchatchers (Sylviidae)	C	C	α	C	c
Ruby-throated Hummingbird Rufous Hummingbird	С	\mathbf{C}	С	R	$\operatorname*{f}$	Blue-gray Gnatcatcher	С	C	С	\mathbf{C}	\mathbf{f}
				11	1	Thrushes (Turdidae)			~		4.5
Kingfishers (Alcedinidae)	C	C	C	C	M	American Robin Eastern Bluebird	$_{ m C}^{ m U}$	$\stackrel{\cup}{\mathrm{C}}$	$^{ m C}_{ m C}$	\mathbf{C}	f,F f,F
Belted Kingfisher	С	С	С	С	IVI	Gray-cheeked Thrush	Ü	O	U	O	\mathbf{f}
Woodpeckers and Allies (Picidae)	TT	т.т	TT	TT	c	Hermit Thrush	U		U	U	$_{c}^{\mathrm{f}}$
Red-Headed Woodpecker Red-Bellied Woodpecker	U U	U U	U U	U U	$\mathbf{f} \\ \mathbf{f}$	Swainson's Thrush Veery	U O		U O		$^{ m f}$
Northern Flicker	U	U	U	U	\mathbf{f}	Wood Thrush*	$\ddot{\mathrm{C}}$	\mathbf{C}	$\overset{\circ}{\mathrm{C}}$		$\mathbf{f}\\ \mathbf{f}$
Yellow-Bellied Sapsucker	U	U	U	U	\mathbf{f}	Mockingbirds and					
Hairy Woodpecker Downy Woodpecker	U U	U U	U U	U U	$_{\rm f}^{\rm f}$	Thrashers (Mimidae)					
Pileated Woodpecker	Ŭ	Ü	Ŭ	Ü	$\dot{\mathbf{f}}$	Brown Thrasher	С	\mathbf{C}	\mathbf{C}	\mathbf{C}	$\mathbf{f} \\ \mathbf{f}$
Tyrant Flycatchers (Tyrannidae)						Gray Catbird Northern Mockingbird	U C	\mathbf{C}	${ m C} \over { m C}$	$_{\mathrm{C}}^{\mathrm{C}}$	$^{ m f}$
Acadian Flycatcher	U	U	U		\mathbf{f}		O	O	O	O	1
Alder Flycatcher	a	C	0		M	Starlings (Sturnidae) European Starling	\mathbf{C}	С	С	\mathbf{C}	F
Eastern Kingbird Eastern Phoebe	$_{\mathrm{C}}^{\mathrm{C}}$	С	C C C	С	$\operatorname*{f}$	•	C	C	C	C	Г
Eastern Wood-Pewee	$_{\mathrm{C}}^{\mathrm{C}}$	\mathbf{C}	$\ddot{\mathrm{C}}$	Ü	\mathbf{f}	Pipits (Motacillidae)	O		O	U	\mathbf{F}
Great Crested Flycatcher	С	\mathbf{C}	U		$_{m{c}}^{\mathrm{f}}$	American Pipit	U		U	U	Г
Least Flycatcher Olive-sided Flycatcher*			O R		$\operatorname*{f}$	Waxwings (Bombycillidae)	a		т.т	a	c
Yellow-bellied Flycatcher			R		\mathbf{f}	Cedar Waxwing	С		U	\mathbf{C}	\mathbf{f}
Shrikes (Laniidae)						Wood-Warblers	**		**		a
Loggerhead Shrike*	O	O	O	O	F	American Redstart* Bay-breasted Warbler	U O	U	U O		$_{f f}$
Vireos (Vireonidae)	C	C	C	C	£	Black-and-white Warbler	Ö		O		$\begin{array}{c} \mathbf{f} \\ \mathbf{f} \\ \mathbf{f} \end{array}$
White-eyed Vireo Yellow-throated Vireo	$_{\mathrm{C}}^{\mathrm{C}}$	$_{\mathrm{C}}^{\mathrm{C}}$	$_{\mathrm{C}}^{\mathrm{C}}$	С	$^{\rm f}_{\rm f}$	Blackburnian Warbler	0		0		f
Blue-headed Vireo	U	-	U	\mathbf{C}	${f f}$	Black-throated Blue Warbler Black-throated	R		R		f
Philadelphia Vireo	R C	\mathbf{C}	$_{ m C}^{ m U}$		$\mathbf{f}\\ \mathbf{f}$	Green Warbler	U		U		\mathbf{f}
Red-eyed Vireo	U	U	U		I						

	Sp	Su	F	W	Habitat		Sp	Su	F	W	Habitat
Blue-winged Warbler*	U		U		\mathbf{f}	Common Grackle	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{f}
Canada Warbler ^*	\mathbf{R}		O		f	Boat-tailed Grackle	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	M
Cerulean Warbler*	R		R		\mathbf{f}	Bronzed Cowbird ^	~	U	~	a	M,f,F
Chestnut-sided Warbler Common Yellowthroat	R C	\mathbf{C}	$_{ m C}^{ m R}$	С	$\mathbf{f}\\\mathbf{M}$	Brown-headed Cowbird Orchard Oriole*	C U	C U	C	С	F,f,M f
Golden-winged Warbler*	$\stackrel{ m C}{ m R}$	C	$^{ m C}_{ m R}$	C	M	Baltimore Oriole	U	Ü	U	U	1 f
Hooded Warbler	Ü	U	Ü		M			O		C	•
Kentucky Warbler*	U	U	U		\mathbf{f}	Frigilline and Cardueline					
Louisiana Waterthrush*	R		R		M	Finches and Allies (Fringillidae)	TT	TT	TT	TT	f IT
Magnolia Warbler	U		U U		f M	—House Finch Pine Siskin	U	U	U	$_{ m R}^{ m U}$	$_{ m f}^{ m f}$
Nashville Warbler Northern Parula	U C	\mathbf{C}	C		f IVI	American Goldfinch	U		U	C	f,F
Northern Waterthrush	ŏ	C	Ü		s	Purple Finch	0		O	O	f
Orange-crowned Warbler	U		U	\mathbf{C}	s						
Ovenbird	U		U	-	S	Old World Sparrows (Passeridae) House Sparrow	U	U	U	U	f,F
Palm Warbler	U	a	\mathbf{C}	$^{ m C}_{ m C}$	F,f	110use Spar10w	U	U	U	U	1,1
Pine Warbler Prairie Warbler*	$_{ m R}^{ m C}$	\mathbf{C}	$_{ m R}^{ m C}$	C	f M	Reporting Your Sightings On eBird					
Prothonotary Warbler*	C	\mathbf{C}	U		M	This online checklist program pro					
Swainson's Warbler*	ŏ	Ŭ	ŏ		M	and Wildlife staff with updated bi					
Tennessee Warbler	U		U		\mathbf{f}	maximizes the utility and accession made each year by both recreation					
Wilson's Warbler	0	ъ	0	\mathbf{R}	M	watchers. The program is one of					
Worm-eating Warbler* Yellow Warbler	O	R O	O	U	f,M M	biodiversity data resources in exi					
Yellow-breasted Chat	U	C	U	U	M	serve as a vital foundation for bet					
Yellow-rumped Warbler	$\ddot{\mathrm{C}}$	O	$\ddot{\mathrm{C}}$	\mathbf{C}	f,M	distribution. The program resour	ces ar	e free	to acc	cess a	and
Yellow-throated Warbler	Ö	O	Ö	$\ddot{\mathrm{R}}$	f,M	are used by a global community of ornithologists, and conservation l			land	mana	agers,
						ormunologists, and conservation	oioiogi	sis.			
Tanagers (Thraupidae) Summer Tanager	U	U	U		${f f}$	Responsible birders should abide b					
Scarlet Tanager	$\overset{ m O}{ m R}$	U	R		\mathbf{f}	Association's Code of Birding Ethic	s. Mor	e infor	matio	n can	be found
					-	at http://www.aba.org1. Promote the welfare of birds	and th		******		
Emberizids (Emberizidae)		т.т			c	1. Fromote the wehare of birds	and u	ieir en	wiron	шеш	J•
Bachman's Sparrow*	U U	U	U U	$_{ m C}^{ m U}$	$^{ m f}$	2. Respect the law, and the right	ts of o	thers.			
Chipping SparrowDark-eyed Junco	U		U	C	F	9 F		1		, .	C 1 1 · 1
Eastern Towhee	$\ddot{\mathrm{C}}$	\mathbf{C}	$\overset{\circ}{\mathrm{C}}$	$\tilde{\mathrm{C}}$	f	3. Ensure that feeders, nest str environments are safe.	ucture	es, and	otnei	r artı	nciai bird
Field Sparrow	O		O	U	\mathbf{F}	environments are sare.					
Fox Sparrow				R	\mathbf{f}	4. Group birding, whether organ	nized o	or imp	romp	tu,	
Grasshopeer Sparrow*	0		0	\mathbf{U}	F F	requires special care.					
Henslow's Sparrow*Lark Sparrow ^*	0		0	U	r F						
Le Conte's Sparrow*	ŏ		ŏ	Ü	F						
Lincoln's Sparrow	U		\mathbf{C}	\mathbf{C}	\mathbf{M}						
Nelson's Sharp-tailed											
Sparrow*	0		0	U	М						
Savannah Sparrow Seaside Sparrow*	C U	U	C U	C U	F, M M						
Song Sparrow	Ü	O	Ü	$\overset{\circ}{\mathrm{C}}$	M						
Swamp Sparrow	$\ddot{\mathrm{C}}$		$\tilde{\mathrm{C}}$	$\ddot{\mathrm{C}}$	M						
Vesper Sparrow	0		0	U	F						
White-crowned Sparrow	0		O U	U	$_{c}^{\mathrm{f}}$						
White-throated Sparrow	U		U	\mathbf{C}	\mathbf{f}						
Cardinals, Saltators, and Allies											
(Cardinalidae)			-	-							
Northern Cardinal	С	\mathbf{C}	С	\mathbf{C}	$\begin{array}{c} \mathbf{f} \\ \mathbf{f} \end{array}$						
Rose-breasted Grosbeak Blue Grosbeak	U U		U U		$^{ m I}$						
Bide Grosbeak Indigo Bunting	Ü	\mathbf{C}	Ü		\mathbf{f}						
Painted Bunting*	O	U	O		\mathbf{f}						
Dickcissel*	\mathbf{R}	\mathbf{R}	\mathbf{R}		\mathbf{F}						
Blackbirds (Icteridae)											
Red-winged Blackbird	\mathbf{C}	\mathbf{C}	\mathbf{C}	\mathbf{C}	M						
Eastern Meadowlark	\mathbf{C}	$\ddot{\mathrm{C}}$	\mathbf{C}	\mathbf{C}	\mathbf{F}						
Rusty Blackbird*	0		O	U	M						
Brewer's Blackbird	O		O	U	F						

