

Recovery Plan

**Tuberculed-blossom
Pearly Mussel
Epioblasma (=Dysnomia)
torulosa torulosa**

**Turgid-blossom Pearly Mussel
Epioblasma (=Dysnomia)
turgidula**

&

**Yellow-blossom Pearly Mussel
Epioblasma (=Dysnomia)
florentina florentina**

RECOVERY PLAN FOR THE

Tuberculed-blossom Pearly Mussel
Epioblasma (=Dysnomia) torulosa torulosa (Rafinesque, 1820)


Turgid-blossom Pearly Mussel
Epioblasma (=Dysnomia) turgidula (Lea, 1858)

Yellow-blossom Pearly Mussel
Epioblasma (=Dysnomia) florentina florentina (Lea, 1857)

Published by

U.S. Fish and Wildlife Service
Atlanta, Georgia

Approved:


Associate Director, U.S. Fish and Wildlife Service

Date:

JAN 25 1985

DISCLAIMER

THIS IS THE COMPLETED RECOVERY PLAN FOR THE TUBERCULED-BLOSSOM, TURGID-BLOSSOM AND YELLOW-BLOSSOM PEARLY MUSSELS. IT HAS BEEN APPROVED BY THE U.S. FISH AND WILDLIFE SERVICE. IT DOES NOT NECESSARILY REPRESENT OFFICIAL POSITIONS OR APPROVALS OF COOPERATING AGENCIES, AND IT DOES NOT NECESSARILY REPRESENT THE VIEWS OF ALL INDIVIDUALS WHO PLAYED A ROLE IN PREPARING THIS PLAN. THIS PLAN IS SUBJECT TO MODIFICATION AS DICTATED BY NEW FINDINGS, CHANGES IN SPECIES STATUS, AND COMPLETION OF TASKS DESCRIBED IN THE PLAN. GOALS AND OBJECTIVES WILL BE ATTAINED AND FUNDS EXPENDED CONTINGENT UPON APPROPRIATIONS, PRIORITIES, AND OTHER CONSTRAINTS.

THE RECOVERY PLANS FOR THE MUSSEL AND FISH SPECIES OF THE TENNESSEE RIVER VALLEY HAVE BEEN DEVELOPED ON A SPECIES-BY-SPECIES BASIS. FOR IMPLEMENTATION PURPOSES, THE PLANS WILL BE CONSOLIDATED ON A WATERSHED BASIS, AND THE NEEDS OF ALL LISTED SPECIES IN THAT SYSTEM WILL BE ADDRESSED.

This plan was prepared under contract by Steven Ahlstedt, Tennessee Valley Authority, Norris, Tennessee.

ADDITIONAL COPIES ARE FOR SALE FROM:

FISH AND WILDLIFE REFERENCE SERVICE
1776 E. JEFFERSON STREET
4TH FLOOR
ROCKVILLE, MARYLAND 20852

301/468-1737, Ext. 326
1-800-582-3421

CONTENTS

	<u>Page</u>
PART I	
INTRODUCTION	1
DISTRIBUTION	3
Historical	3
Present	9
ECOLOGY AND LIFE HISTORY	10
<u>Epioblasma torulosa torulosa</u>	11
<u>Epioblasma turgidula</u>	13
<u>Epioblasma florentina florentina</u>	15
REASONS FOR DECLINE AND/OR EXTINCTION	20
Impoundment	21
Siltation	22
Pollution	23
PART II	
RECOVERY	26
Recovery Objectives	26
Step-Down Outline	27
Narrative Outline	27
BIBLIOGRAPHY	30
PART III	
IMPLEMENTATION SCHEDULE	37
APPENDIX - List of Reviewers	39

TABLE

1. Historical records for <u>Epioblasma (=Dysnomia) torulosa torulosa</u> prior to 1970 with archaeological specimens recorded to 1983	4
--	---

CONTENTS
(continued)

	<u>Page</u>
2. Historical records for <u>Epioblasma (=Dysnomia) turgidula</u> prior to 1970 with archaeological specimens recorded to 1982	7
3. Historical records for <u>Epioblasma (=Dysnomia) florentina</u> <u>florentina</u> prior to 1970 with archaeological specimens recorded to 1982	8

FIGURE

1. Typical Naiad Life Cycle Depicting the Various Stages	19
--	----

PHOTOGRAPHS

1. Photograph of <u>E. t. torulosa</u>	12
2. Photograph of <u>E. turgidula</u>	14
3. Photograph of <u>E. f. florentina</u>	16

PART I
INTRODUCTION

Freshwater mussels (naiades) are known to occur in every temperate and tropical climate. Approximately one-half of the extant species of freshwater mussels occur in North America. These species occur in what has been recognized as six faunal zones or regions which include: (1) North Atlantic Slope, (2) Pacific coastal, (3) Mississippian, (4) Ozarkian, (5) Cumberlandian, and (6) Ohioan Regions (Johnson, 1980). Eastern North America still contains the most diverse freshwater molluscan fauna known in the world. Stansbery (1970) reported that fauna numbers over a thousand species of bivalves and gastropods combined. Unique among all freshwater mussels is a genus of Unionidae Epioblasma (=Dysnomia), known for their high degree of sexual dimorphism. Of this group, eight species reported earlier this century are presumed extinct. The extinction of this group appears to be symptomatic of a general synecological problem that exists between Epioblasma and environmental changes that have occurred in many streams and rivers throughout North America. Stansbery (1971) reports that all members of this group, without exception, are characteristic riffle or shoal species inhabiting parts of those streams which are shallow with sandy-gravel substrate and rapid currents. The elimination of these species has been attributed to impoundments, barge canals, and other flow alteration structures that have eliminated riffle and shoal areas.

Twenty-three freshwater mussels in the United States are listed as endangered by the U.S. Department of Interior. Of these, eight species are members of the genus Epioblasma (=Dysnomia). Three of these

species, (1) tuberculed-blossom pearly mussel Epioblasma torulosa torulosa, (2) turgid-blossom pearly mussel Epioblasma turgidula, and (3) yellow-blossom pearly mussel Epioblasma florentina florentina were all proposed for listing in September, 1975 (Federal Register 40(188):44329-44333) and listed in June, 1976 (Federal Register 41(115):24062-24067). Due to the scarcity of recent information concerning each of these species, recovery plans for all three species will be combined into one document.

Epioblasma t. torulosa was described by Rafinesque (1820) as Amblema torulosa from the Ohio and Kentucky Rivers. For clarification of nomenclatorial changes in this group refer to Bogan and Parmalee (1983). Vanatta (1915) also reported the type specimen as coming from the Kentucky River. All records for this species (including various named forms or subspecies) indicate that it was widespread in the larger rivers of eastern United States and southern Ontario, Canada.

Epioblasma turgidula was described by Lea (1858) as Unio turgidulus from the Cumberland River, Tennessee, and Florence, Alabama (Tennessee River). The female of this species was later described by Reeve (1864) as Unio deviatus (Stansbery, 1976a). Call (1885) further listed the distribution as the Cumberland and Duck Rivers, Tennessee. The Cumberland River, Tennessee, record in the original description is questionable in that the lower Tennessee River was often referred to as the Cumberland River in earlier years (Stansbery, 1976a). Epioblasma turgidula is a medium to large river Cumberlandian type reported from the Ozarks. Ortmann (1925) postulated that this distribution pattern suggests a former connection of the Ozarks with the Cumberland Plateau.

Epioblasma f. florentina was described by Lea (1857) as Unio florentinus from the Tennessee River, Florence and Lauderdale Counties,

Alabama, and the Cumberland River, Tennessee. Epioblasma f. florentina was a large river Cumberlandian type found throughout the Tennessee and Cumberland River drainages. One apparently related headwaters form, Epioblasma walkeri occurs in the headwater tributaries of these drainages while another apparently related form, Epioblasma florentina curtisi, is endemic to the Ozarkian Region of Missouri.

DISTRIBUTION

Historical

Epioblasma torulosa (including forms or subspecies) once was widespread in the Tennessee, Cumberland, Ohio, and St. Lawrence river drainages. The inflated big river form, Epioblasma t. torulosa was found throughout the Tennessee River system including the Elk, Duck, and Nolichucky Rivers. Based on the amount of material observed by the author, Stansbery (1976a), and Bogan and Parmalee (1983) from archaeological Indian shell middens along the banks of the Tennessee River, E. t. torulosa was a relatively common species in the Tennessee River. Epioblasma t. torulosa was also reported from the Cumberland River but must have been very rare based on the lack of records from the Cumberland River for this species. Ortmann (1925) reported hesitantly that E. t. torulosa occurred in the Cumberland River. However, archaeological specimens of E. t. torulosa are reported from the Cumberland by Parmalee et al., (1980) and Bogan and Parmalee (1983). Epioblasma t. torulosa is also reported from the Ohio and Kentucky Rivers, and the Scioto and Kanawha Rivers (Stansbery, 1961, 1972, 1980; Johnson, 1978; Clarke, 1982; Taylor, 1983). Table 1 lists known historical records prior to 1970 with archaeological specimens recorded to 1983.

Table 1. Historical records for Epioblasma (=Dysnomia) torulosa torulosa prior to 1970 with archaeological specimens recorded to 1983.

River	Source
Tennessee River	Rafinesque (1820) Conrad (1834) Lewis (1876) Call (1885) Hinkley (1904, 1906) Ortmann (1918, 1920, 1925) van der Schalie (1939) Stansbery (1964, 1976b) Warren (1975) Parmalee et al. (1980) archaeological specimens Bogan and Parmalee (1983) archaeological specimens
Elk River	Ortmann (1925) van der Schalie (1939) Stansbery (1964) Isom et al. (1973) Johnson (1978) Ahlstedt (1983) relict specimens observed
Duck River	Bogan and Parmalee (1983) specimens collected by Athearn
Nolichucky River	Johnson (1978)
Cumberland River	Ortmann (1926) Parmalee et al. (1980) archaeological specimens Bogan and Parmalee (1983) archaeological specimens
Ohio River	Rafinesque (1820) Call (1900) Ortmann and Walker (1922) Ortmann (1925) Parmalee (1960, 1967) La Roque (1967) Stansbery (1970) Johnson (1978)
Kentucky River	Rafinesque (1820) Vanatta (1915) Danglade (1922) Ortmann and Walker (1922) Johnson (1978, 1980)

Table 1. Continued.

River	Source
Kanawha River, W. Virginia	Stansbery (1972) archaeological specimens Stansbery (1980) Clarke (1982) relict specimens Taylor (1983)
Scioto River, Ohio	Higgins (1858) Stansbery (1961)

Epioblasma turgidula was historically relatively widespread with a disjunct distribution occurring in both the Cumberlandian and Ozarkian Regions (Johnson, 1978). This Cumberlandian form has been reported from the Tennessee River and tributary streams including Shoal and Bear Creeks, Elk, Duck, Holston, Clinch, and Emory Rivers (Ortmann, 1918, 1924, 1925; Stansbery, 1964, 1970, 1971, 1976a; Johnson, 1978). Additional records are also reported from the Cumberland River (Ortmann, 1918; Clench and van der Schalie, 1944; Johnson, 1978). This species is also reported from the Ozark Mountain Region including Spring Creek, Black and White Rivers (Simpson, 1914; Johnson, 1978). Table 2 lists known historical records prior to 1970 with archaeological specimens recorded to 1982.

Epioblasma f. florentina is a Cumberlandian type historically widespread in the Tennessee and Cumberland Rivers and tributaries to the Tennessee River. Epioblasma f. florentina was reported from Hurricane, Limestone, Bear, and Cypress Creeks, all tributary streams to the Tennessee River in northern Alabama (Ortmann, 1925; Bogan and Parmalee, 1983). This species was reported from larger tributary streams of the lower and upper Tennessee River including the Flint, Elk, and Duck Rivers (Marsh, 1885; Isom et al. 1973; Bogan and Parmalee, 1983) and the Holston, Clinch, and Little Tennessee Rivers (Ortmann, 1918; Stansbery, 1970; TVA, 1972a). Specimens were also reported from Citico Creek, a tributary to the Little Tennessee River (Bogan and Parmalee, 1983). Epioblasma f. florentina apparently occurred throughout the Cumberland River (Wilson and Clark, 1914; Ortmann, 1918, 1925; Shoup et al. 1941; Neel and Allen, 1964; Parmalee et al. 1980). Table 3 lists known historical records prior to 1970 with archaeological specimens recorded to 1982.

Table 2. Historical records for *Epioblasma* (=Dysnomia) *turgidula* prior to 1970 with archaeological specimens recorded to 1982.

River	Source
Spring Creek, Arkansas	Simpson (1914)
Black River, Arkansas	Johnson (1978) specimens reported early 1900's
White River, Missouri and Arkansas	Johnson (1978) specimens reported early 1900's
Tennessee River	Lea (1857, 1858) Lewis (1876) Call (1885) Simpson (1900, 1914) Ortmann (1918, 1924) Stansbery (1964) Johnson (1978) Parmalee et al. (1982) archaeological specimens
Shoals Creek, Alabama	Hinkley (1906) Ortmann (1918, 1925) Johnson (1978)
Bear Creek, Alabama	Ortmann (1918, 1924, 1925) Johnson (1978)
Elk River	Stansbery (1970, 1971) Johnson (1978)
Duck River	Marsh (1885) Call (1885) Ortmann (1918, 1924) Stansbery (1976a)
Holston River, Tennessee	Ortmann (1918, 1924) Johnson (1978)
Clinch River, Tennessee	Johnson (1978)
Emory River, Tennessee	Ortmann (1918, 1924)
Cumberland River	Lea (1858) Call (1885) Simpson (1900, 1914) Ortmann (1918) Clench and van der Schalie (1944) Johnson (1978)

Table 3. Historical records for Epioblasma (=Dysnomia) florentina florentina prior to 1970 with archaeological specimens recorded to 1982.

River	Source
Tennessee River	Lea (1857) Hinkley (1906) Lewis (1876) Ortmann (1918, 1925) Parmalee et al. (1982) archaeological specimens Morrison (1942) Simpson (1900, 1914) Stansbery (1964, 1976b)
Flint River, Alabama	Bogan and Parmalee (1983)
Hurricane Creek, Alabama	Bogan and Parmalee (1983)
Limestone Creek, Alabama	Bogan and Parmalee (1983)
Bear Creek, Alabama	Bogan and Parmalee (1983)
Cypress Creek, Alabama	Ortmann (1925)
Elk River	Isom et al. (1973)
Duck River	Marsh (1885)
Holston River	Ortmann (1918)
Little Tennessee River, Tennessee	TVA (1972a) specimens collected by Athearn 1962, 1963, 1965
Citico Creek, Tennessee	Bogan and Parmalee (1983) Athearn collected specimens in 1957
Clinch River	Stansbery (1971)
Cumberland River	Lea (1861) Ortmann (1918, 1925) Wilson and Clark (1914) Shoup et al. (1941) Neel and Allen (1964) Simpson (1900, 1914) Parmalee et al. (1980) archaeological specimens Sickel (1982) relict specimens

Present

Epioblasma t. torulosa was collected fresh-dead (one specimen) in 1969 from the Kanawha River, West Virginia, below Kanawha Falls (Stansbery, 1980). To date, this represents the only record of this species from the Kanawha River. Stansbery reported that no living specimens of E. t. torulosa have ever been collected from the Kanawha River. Recent studies of the Kanawha River by Clarke (1982) and Taylor (1983) found no evidence of E. t. torulosa. Clarke (1982) recommended that a detailed scuba search be conducted below Kanawha Falls to determine this species' status in the Kanawha. Taylor (1983) concluded that this species no longer occurs in the drainage. Stansbery (1976b) and Jenkinson (1981) consider this species to be "probably extinct."

Epioblasma turgidula was last reported in the mid-1960's from the Duck River near Normandy, Tennessee (Stansbery, 1976a). With the completion of Normandy Dam on the Duck River by TVA in 1976, the three locations where specimens of E. turgidula were collected are now either impounded or affected by the operation of Normandy Dam. This species has not been found in recent freshwater mussel surveys conducted on the Duck River (TVA, 1972b, 1979; Ahlstedt, 1980), nor has it been reported from any other stream or river system. Stansbery (1971) and Jenkinson (1981) consider this species to be extinct.

Stansbery (1971, 1976b) reported that E. f. florentina has not been seen or collected in over a half century and thus considers this species extinct. However, Athearn collected specimens of E. f. florentina from the Little Tennessee River in the mid 1960's and from Citico Creek in 1957 (TVA, 1972a; Bogan and Parmalee, 1983).

ECOLOGY AND LIFE HISTORY

Freshwater mussels (naiades), are benthic animals that normally remain buried in the substrate with only the most posterior margin of the shell and siphons exposed to the water column. Freshwater mussels are found in a variety of habitats ranging from mud and sand between bedrock ledges, to rubble and gravel substrates. The majority of freshwater mussel species are usually found in riverine conditions in relatively firm rubble, gravel, and sand substrate swept-free of siltation. Typically, most mussel species are found buried in the substrate in relatively shallow areas called riffles or shoals. Freshwater mussels in the genus Epioblasma are characteristic riffle or shoal species inhabiting those parts of streams which are shallow with sandy gravel substrate and rapid currents (Stansbery, 1971). The eight species presumed extinct were, with few exceptions, recorded from riffles of our largest rivers. This type of habitat has nearly been eliminated by impoundment of the large rivers. Of these eight species, Stansbery lists E. f. florentina and E. turgidula as extinct. Epioblasma f. florentina has not been seen in over half a century and the last known population of E. turgidula in the Duck River were apparently destroyed. Neel and Allen's (1964) survey of the upper Cumberland Basin documented an almost total elimination of the genus Epioblasma (=Dysnomia) presumably due to mine wastes.

Epioblasma torulosa torulosa (photo 1)

Epioblasma t. torulosa is a larger, more inflated form of Epioblasma torulosa gubernaculum (Reeve, 1865) which occurs only in headwater tributaries of the Tennessee River above Knoxville, Tennessee, (as per Ortmann, 1918). Epioblasma t. torulosa exhibited considerable ecophenotypic variation in sculpture and obesity. Ball (1922) demonstrated the relationships between shell inflation and tuberculation with streamsize in this species. Previously, Ortmann (1918) noted that the typical big river form E. t. torulosa from the Tennessee River had a row of prominent knobs across the middle of the shell but, further upstream, the knobs become reduced and finally disappear, thus passing into the form E. t. gubernaculum. Epioblasma t. torulosa also appeared to grade into another form Epioblasma torulosa rangiana (Lea, 1838) of the Ohio River system (Ortmann, 1913; Johnson, 1978; Bogan and Parmalee, 1983). Johnson (1978) provides the following description of the shell:

Shell of medium size, reaching almost 90 mm in length. Outline irregularly ovate, elliptical or obovate. Valves inequilateral, subinflated to inflated, solid. Anterior end regularly rounded, posterior end of male slightly produced; more broadly rounded in females. Ventral margin slightly curved. Dorsal margin curved forming an indistinct angle with the obliquely descending posterior margin. Hinge ligament short. Posterior ridge of the male rather low, narrowly rounded, separated from a similar medial ridge by a broad furrow of varying depth that ends in an emargination between the somewhat produced ridges. Both the ridges and the furrow vary from being smooth to having elevated knobs. The marsupial swelling in the female is sometimes marked by a number of small radial furrows, but the ridges become obscure. The rounded marsupial swelling extended from the middle of the base



PHOTO 1. Epioblasma torulosa torulosa

0 1
cm

to the upper part of the posterior end. Umbos full, somewhat turned over a small lunule, located toward the anterior third of the shell, sculpture feeble and corrugated. Surface of the shell with many distinct growth lines. Periostracum smooth and shiny, tawny, yellowish green, or straw colored, usually with numerous green rays.

Left valve with two triangular pseudocardial teeth; slight interdentum. Two long, almost straight, lateral teeth. Right valve with one large pseudocardial tooth with a smaller tooth before it. One lateral tooth, sometimes with a vestigial tooth below. Umbonal cavity shallow. Anterior and posterior adductor muscle scars and pallial line, well impressed. The shell is thin in the marsupial region of the female especially toward the margin. Nacre white to salmon-red.

Male shells are generally irregularly ovate with a rather wide radial furrow of varying depth ending in a broad sinus, often bluntly pointed behind.

Female shells are generally obovate, larger than the male, having a large, flattened, rounded marsupial swelling extending from the middle of the base to near the upper part of the posterior end, which is thin and often a darker green than the rest of the shell.

Epioblasma turgidula (photo 2)

Epioblasma turgidula is a small-sized Cumberlandian type found both in the Cumberlandian and Ozarkian Plateau regions. As with other members of this genus, this species is sexually dimorphic. Male specimens of E. turgidula were originally described by Lea (1858) while Reeve (1864) later described the female as Unio deviatus. Walker (1910) in his paper on Truncilla first recognized the two sexes as being the same species since he includes Truncilla deviatus in his keys to both male and female specimens. Ortmann (1918) also correctly identified U. deviatus as the female of U. turgidulus. Stansbery (1976a) and Johnson (1978) report that E. turgidula closely resembles E. biemarginata and is superficially similar to all the members of the subgenus Capsaeformis. A detailed description of E. turgidula is given by Johnson (1978) and is presented below:




PHOTO 2. Epioblasma turgidula



Shell rather small, seldom reaching more than 40 mm in length. Outline elliptical, ovate, or obovate. Valves inequilateral, solid, slightly inflated. Anterior and regularly rounded; posterior end of male rather pointed, of female more broadly rounded. Ventral margin curved in males, almost straight in females before the marsupial swelling. Dorsal margin slightly curved forming an indistinct angle with the obliquely descending posterior margin. Hinge ligament short. Posterior ridge of the male, double, somewhat raised, ending posteriorly in a slight biangulation. The ridges of the female fade out on the marsupial swelling. Dorsal slope slightly concave. The male has a rather wide, shallow, radial furrow, which ends in an emargination. The medial ridge is only slightly developed, and both it and the radial furrow are obscured in females by marsupial swelling. Umbos moderately full and elevated, located in the anterior third of the shell; umbonal sculpture not observed. Surface of the shell with irregular growth lines, which are especially strong on the female posteriorly. Periostracum rather shiny, yellowish green, with numerous fine green rays over the entire surface.

Left valve with two small, rough, subcompressed pseudocardinal teeth. No interdentum. Two short, straight, lateral teeth. Right valve with one small, triangular, pseudocardinal tooth, with a second tiny tooth before it that is parallel to the hinge line. One lateral tooth. Umbonal cavities shallow. Anterior adductor muscle scars well impressed; posterior scars faint. Pallial line distinct anteriorly. Nacre bluish white. Shell heavier anteriorly, posterior end of females especially thin and iridescent.

Male shells are elliptical or ovate, with a distinct raised double posterior ridge, which ends in a biangulation near the base. Before the ridge is a wide shallow radial furrow and faint medial ridge.

The female is somewhat obovate, and while the marsupial swelling obliterates the radial furrow and the medial and posterior ridges, there is a tendency for the shell to be somewhat concave in the region of the posterior ridges. The surface of the marsupial swelling is not different from the rest of the shell, the feeble green rays are rather evenly distributed in both sexes.

Epioblasma florentina florentina (photo 3)

Epioblasma f. florentina is a medium-sized Cumberlandian type found both in the Cumberlandian and Ozarkian regions. This species is represented in the Ozarkian region by E. f. curtisi (Utterback, 1915)



♀



♂

0 1
cm

PHOTO 3. Epioblasma florentina florentina

which is being evaluated in a separate recovery plan. E. f. florentina represents the big-river form of this species which may grade into the smaller, headwaters form E. f. walkeri (Wilson and Clark, 1914; Ortmann, 1918 and 1924). As with E. turgidula, E. f. florentina is superficially similar to all the members of the subgenus Capsaeformis. The detailed description of E. f. florentina given by Johnson (1978) is as follows:

Shell of medium size, seldom reaching over 60 mm in length. Outline elliptical or irregularly obovate. Valves somewhat inequilateral, subinflated, subsolid. Anterior end regularly rounded, posterior end of male slightly produced; posterior more broadly rounded in females. Ventral margin slightly but uniformly curved in the male; almost straight in females to the sulcus, behind which the marsupial swelling extends well below the base. Dorsal margin straight, forming an angle with the obliquely descending posterior margin. Hinge ligament short. Posterior ridge of the male, double, but faint, ending in a slight biangulation posteriorly; ridge scarcely visible in females. There is a wide shallow radial depression in front of the full, lower posterior ridge in the male. The considerable marsupial swelling of the female is usually marked by a sulcus before and behind and is sometimes finely toothed on the margin.

Umbos quite full and elevated, located slightly anterior of the middle in the male, and in the anterior third of the shell in the female, umbonal sculpture not observed. Surface of the shell with uneven growth lines. Periostracum subshiny, yellow, honey yellow, brownish yellow, or whitish with numerous green rays more or less uniformly distributed over the entire surface.

Left valve with two chunky triangular pseudocardinal teeth; slight interdentum. Two short slightly curved lateral teeth. Right valve with one large triangular pseudocardinal tooth, usually with a smaller tooth before it that is parallel to the hinge line. One lateral tooth, sometimes with a vestigial tooth below. Umbonal cavities shallow. Anterior adductor muscle scars well impressed, posterior ones shallow. Pallial line distinct anteriorly. Nacre bluish white. Shell heavier anteriorly, posterior end of females especially thin and iridescent.

Male shells irregularly elliptical, with a double posterior ridge that ends in a slight bimargination near the median. There is a wide, shallow, radial depression in front of the full, lower posterior ridge.

Female shells irregularly obovate with a thin, slightly inflated marsupial swelling, which may be considerably produced and extended well below the base, often marked by two distinct sulci, toothed on the margin.

The life histories of E. t. torulosa, E. turgidula, and E. f. florentina are probably similar to those of most naiades and are briefly illustrated in Figure 1. Males produce sperm which are discharged into the surrounding water and dispersed by water currents. Females downstream from the males obtain these sperm during the normal process of siphoning water while feeding and respiration (Stein, 1971). Fertilization of the eggs by sperm occurs within the gills of the female. In the genus Epioblasma fertilized eggs are retained in the posterior section of the outer gills which are modified as brood pouches. Based on the position of the genus Epioblasma (=Dysnomia) in the subfamily Lampsilinae, it is assumed that only the outer two demibranches serve as ovisacs (Heard and Guckert, 1970).

The family Unionidae are separated into two groups based on the length of time glochidia remain in the female (Ortmann, 1919). By Ortmann's definitions, bradytictic bivalves (long-term breeders) breed from midsummer through fall or early winter. Embryos develop in the female over winter and are released the following spring or summer. Tachytictic bivalves (short-term breeders) breed in spring and release glochidia by mid- to late-summer of the same year. Data for other Epioblasma species (Ortmann, 1919; Bogan and Parmalee, 1983) would suggest that Epioblasma t. torulosa, E. turgidula, and E. f. florentina are long-term breeders.




Figure 1. Typical naiad life cycle depicting the various stages. The life cycle for most species of naiades is very similar to that depicted here (Grace and Buchanan 1981).

Glochidial releases by some bradytictic species have also been observed during fall and winter (Zale, 1980). Upon release into the water column, mature glochidia attach to the gills or fins of appropriate host fishes to encyst and metamorphose to the juvenile stage. The glochidia of most Epioblasma species are bean-shaped and hookless. Hookless glochidia typically have a more spoon-shaped delicate shell and are most frequently parasitic on gill filaments of fish (Lefevre and Curtis, 1910; Coker and Surber, 1911). The fish host(s) for E. t. torulosa, E. turgidula, and E. f. florentina are unknown.

REASONS FOR DECLINE AND/OR EXTINCTION

The genus Epioblasma (=Dysnomia) as a whole is becoming rarer because all members of this genus are riverine and typically found in streams which are shallow with sandy-gravel substrate with rapid currents (Stansbery, 1971). Eight Epioblasma species are presumed extinct and were, with few exceptions, recorded from riffles or shoals of our largest rivers. The reasons for the extinction of these species and the decline of the remaining Epioblasma species are not totally understood; however, these species or their fish host(s) may be reacting to levels or types of impacts that are not killing other species in the same habitat. Due to the longevity of most mussel species--up to 50 years--and their sedentary nature, they are especially vulnerable to stream perturbations such as impoundments, siltation, and pollution.

Impoundment

Possibly the single greatest factor contributing to the decline of freshwater mussels, not only in the Tennessee Valley but other regions as well, is the alteration and destruction of stream habitat due to impoundments for flood control, navigation, hydro-electric power production, and recreation. Ortmann's (1918) monograph on the naiads of the upper Tennessee River is the most significant work on that region's freshwater mussel fauna prior to construction of impoundments on many of these streams. At the time of Ortmann's survey, a total of 66 species of mussels occurred in the Tennessee River between Chattanooga and Knoxville, Tennessee. Pardue (1981) reported only 23 species of mussels living in the Tennessee River during a survey conducted in 1978. Publications by Wilson and Clark (1912, 1914) and Neel and Allen (1964) on the mussels of the Cumberland River and its tributaries also offer an excellent historical account of that fauna prior to impoundment and extensive coal mining. A total of 79 species of freshwater mussels was reported by Wilson and Clark (1914) and 59 species were later reported by Neel and Allen (1964).

Since the early 1930's and 1940's, the Tennessee Valley Authority, Aluminum Company of America (Alcoa), and the U.S. Army Corps of Engineers have constructed 51 impoundments throughout the Tennessee and Cumberland River systems alone. Stream impoundments affect species composition by eliminating those species not capable of adapting to reduced flows, altered temperature regimes, and anoxic conditions. Tributary dams typically have hypolimnial discharges that cause the stream below the dam (reservoir tailwater) to differ significantly from both preimpoundment conditions and from upstream river reaches. Hypolimnial discharges

include: altered temperature regimes, extreme water level fluctuations, reduced turbidity, seasonal oxygen deficits, and high concentrations of certain heavy metals (TVA, 1980). Biological responses attributable to these environmental changes typically include reductions in the fish and benthic macroinvertebrate communities (Isom, 1971). Hickman (1937) recorded numerous species of mussels and snails in the vicinity of the Norris Dam construction site prior to the impoundment of that reach of the Clinch River and predicted that the Norris Dam flood control project would have a deteriorating effect on the molluscan fauna. A. R. Cahn (1936) collected 45 mussel species and nine river snail species in the dewatered riverbed following closure of Norris Dam. In a return visit to the area 4 months later, he could not find a single live mussel. Ortman (1925) witnessed the most famous and unique locality for mussels (Muscle Shoals, Alabama), destroyed by the construction of Wilson Dam on the Tennessee River.

Siltation

Siltation is another factor that has severely affected freshwater mussels. In rivers and streams, the greatest diversity and abundance of mussels are usually associated with gravel and/or sand substrates. These substrates are most common in running water (Hynes, 1970). Increased silt transport into our waterways due to strip mining, coal washing, dredging, farming, logging, and road construction are some of the more obvious results of human alteration of the landscape. Hynes (1974) states that there are two major effects of inorganic sediments introduced into aquatic ecosystems. The first is an increase in the turbidity of the water with a consequent reduction in the depth

of light penetration, and the second is a blanketing effect on the substrate. High turbidity levels due to the presence of suspended solids in the water column have a mechanical or abrasive action which can irritate, damage, or clog the gills or feeding structures of mollusks (Loar et al. 1980). Additionally, high levels of suspended solids may reduce or inhibit feeding by filter-feeding organisms, such as mussels, causing nutritional stress and mortality (Loosanoff, 1961). Freshwater mussels are quite long lived and rather sedentary by nature; many species are unable to survive in a layer of silt greater than 0.6 centimeters (Ellis, 1936). Since most freshwater mussels, especially the Cumberlandian types and the genus Epioblasma, are typically riverine species that require clean, flowing water over stable, silt-free rubble, gravel, and sand shoals, the smothering action of silt is often severe. Fuller (1977) reported that siltation associated with poor agricultural practices and deforestation of much of North America was probably the most significant factor impacting mussel communities. Mussel life cycles can be affected indirectly from siltation by impacting host-fish populations by smothering fish eggs or larvae, reducing food availability, or filling of interstitial spaces in gravel and rubble substrate, thus eliminating spawning beds and habitat critical to the survival of young fishes (Loar et al. 1980).

Pollution

A third factor which must be considered is the impact caused by various forms of pollutants. An increasing number of streams throughout the United States receive municipal, agricultural, and industrial waste discharges. The damage suffered varies according to a complex

of interrelated factors, which includes the characteristics of the receiving stream and the nature, magnitude, and frequency of the stresses being applied. The degradation can be so severe and of such duration that the streams are no longer considered valuable in terms of their biological resources (Hill et al. 1974). These areas will not recover if there are residual effects from the pollutants, or if there is an inadequate pool of organisms for recruitment or recolonization (Cairns et al. 1971).

The absence of freshwater mussels can be an indication of environmental disruption only when and where their former presence can be demonstrated (Fuller, 1974). It is very rare that the composition and size of the mussel fauna can be quantitatively and/or qualitatively correlated with a specific disruption, be it chemical or physical (Ingram, 1956). However, some data are available concerning the adverse impacts of some pollutants on freshwater mussels along with other components of the ecosystem. Neel and Allen (1964) reported that coal mine acids in headwater tributaries of the Cumberland River have practically eliminated the most diverse known assemblage of Epioblasma (= Dysnomia) species. A combination of toxic wastes, gravel dredging, and increased fertilizer and pesticide use has reduced the freshwater mussel fauna in the Stones River from 45 to 30 species (Schmidt, 1982). Ortmann (1918), in his studies of the freshwater mussels in the upper Tennessee drainage, reported numerous streams to be already polluted and the mussel fauna gone. These streams included the Powell River, for a certain distance below Big Stone Gap, Virginia (wood extracting plant); the North Fork Holston River for some distance below Saltville, Virginia (salt and plaster of Paris industries); French Broad River at Asheville,

North Carolina; Big Pigeon River from Canton, North Carolina, all the way to its mouth (wood pulp and paper mill); and the Tellico River below Tellico Plains, Tennessee (wood pulp and extracting mill).

PART II
RECOVERY

A. Recovery Objectives

The ultimate goal of this recovery plan is to locate, maintain, and enhance any known populations of E. t. torulosa, E. turgidula, and E. f. florentina. Based on the information provided in this document and the opinions expressed by numerous experts, it appears that all three of these species may already be extinct. Stansbery (1971) listed eight Epioblasma species presumed to be extinct, including E. turgidula and E. f. florentina. In 1976, Stansbery reported that E. t. torulosa may also be extinct. A survey by TVA biologists (Jenkinson, 1981) found no reason to believe that E. f. florentina, E. turgidula, and E. t. torulosa might be found alive. The extinction of these species appears to be symptomatic of a general synecological problem that exists between Epioblasma and chronic environmental changes that have occurred and are continuing to occur in most major river systems. Based on the recent decline for other endangered species of this group, including E. walkeri (Neves, 1984), E. t. gubernaculum (Ahlstedt, 1984), and E. sampsoni (Clarke, 1981), it is highly improbable, if and when live specimens of any one of the three subject species are found that (1) populations can be restored to other rivers within its historic range, or (2) the species can ever recover to the point of delisting. Nevertheless, recovery efforts for each species will be reevaluated when the following criteria are met:

1. A reproducing population of either E. t. torulosa, E. turgidula, or E. f. florentina are found in any stream or river system.
2. Each species and its habitat are protected from present and foreseeable anthropogenic and natural events that may interfere with the survival of the population.

B. Step-Down Outline

Prime Objective: Determine the current status for each of these species, and provide protection for any populations located.

1. Conduct intensive surveys for each species.
 - 1.1 Identify essential habitat and specific areas in need of protection.
 - 1.2 Utilize existing legislation and regulations to protect the species and its habitat.
2. Determine present and foreseeable threats to the species and strive to minimize and/or eliminate them.
3. Assess overall status for species listed in this recovery plan.

C. Narrative Outline

1. Conduct intensive surveys for each species. Intensive dive/float surveys using scuba-equipped divers, snorkelers, and/or commercial mussel fishermen are recommended for attempting to locate living populations of each species:

Epioblasma t. torulosa--Freshwater mussel surveys are recommended for the lower Ohio River below Owensboro, Kentucky, and the Kanawha River, West Virginia, below Kanawha Falls (KRM 95.5-81.0).

Epioblasma turgidula--Freshwater mussel surveys are recommended for Shoal and Bear Creeks, Alabama; Emory River, Tennessee; Spring Creek, Arkansas; Black River, Arkansas; and White River, Missouri and Arkansas.

Epioblasma f. florentina--Freshwater mussel surveys are recommended for Hurricane and Bear Creeks, Alabama; upper Little Tennessee River above Tellico Reservoir; Citico Creek, Tennessee; and the Big South Fork Cumberland River.

- 1.1 Identify essential habitat and specific areas in need of protection. If individuals are found, the extent of the population must be delineated.
- 1.2 Utilize existing legislation and regulations to protect the species and its habitat. Prior to and during implementation of this recovery plan the species will be protected by the full enforcement of existing laws and regulations.
2. Determine present and foreseeable threats to the species and strive to minimize and/or eliminate them. All local, state, and federal development and enforcement agencies and land use groups will be notified of the sensitivity of certain areas with known populations of these species. Since these areas will probably represent the only known location(s) for these species, every effort will be made to prevent any modification or impact which might prove harmful to the species and its habitat. These impacts typically include dredging, strip

mining, industrial development, road and bridge construction, installation of sewage treatment plants and their operation, and the use of herbicides along roads and powerline corridors as well as pesticides and fertilizers for farm crops.

3. Assess overall status of each species. This recovery plan must be evaluated periodically to determine the current status of these 3 species. If freshwater mussel surveys fail to locate live specimens of E. t. torulosa, E. turgidula, and E. f. florentina, these species will be recommended for removal from the federal endangered species list due to extinction.

BIBLIOGRAPHY

- Ahlstedt, S. A. 1980. The molluscan fauna of the Duck River between Normandy and Columbia Dams in central Tennessee. Bull. Amer. Malacol. Union for 1980:60-62.
- _____. 1984. Recovery plan for green-blossom pearly mussel (Epioblasma (=Dysnomia) torulosa gubernaculum, Reeve, 1865). USFWS, Region 4, Atlanta, Georgia. 50 pp.
- _____. 1983. The molluscan fauna of the Elk River in Tennessee and Alabama. Bull. Amer. Malacol. Union for 1983 (1):43-50.
- Anthony, J. G. 1865. Descriptions of new species of North American Unionidae. Am. J. Conch. 1:155-164. pl. 12-16.
- Ball, G. H. 1922. Variation in freshwater mussels. Ecology 3(2):93-121.
- Bates, J. M. 1970. Ohio mussel fisheries study Part 1. Mussel studies. Center for Aquatic Biology, Eastern Michigan Univ. at Ypsilanti. Nat. Tech. Info. Serv. Pub. Com-73-10145. pp. 1-108.
- Bogan, A. and P. Parmalee. 1983. Tennessee's rare mollusks. IN: Tennessee's Rare Wildlife, Final Report: TWRA, Tennessee Department of Conservation and Tennessee Heritage Prog. June 1979. Univ. of TN., Knoxville. 360 pp.
- Cahn, A. R. 1936. The molluscan fauna of the Clinch River below Norris Dam upon completion of that structure. Unpublished TVA report. Norris, TN.
- Cairns, J., Jr., J. S. Crossman, K. L. Dickson, and E. E. Herricks. 1971. The recovery of damaged streams. Assoc. Southeast. Biol. Bull. 18:79-106.
- Call, R. E. 1885. Unionidae of the Mississippi Valley. Bull. Des Moines Acad. Sci. 1(1):5-57.
- _____. 1900. A descriptive illustrated catalogue of the mollusca of Indiana. Indiana Dept. Geol. Nat. Res. Ann. Rept. 24:335-535, 78 pl.
- Clarke, A. H. 1981. Determination of the precise geographical areas occupied by four endangered species of freshwater mollusks. Final report. USFWS, Fed. Bldg. Room 668, Fort Snelling, Twin Cities, MN.

- Clarke, A. H. 1982. Survey of the freshwater mussels of the upper Kanawha River (RM 91-95), Fayette County, West Virginia, with special reference to Epioblasma torulosa (Rafinesque) and Lampsilis abrupta (Say) (= Lampsilis orbiculata (Hildreth), of authors). Final Report. USFWS, One Gateway Center, Room 618, Newton Corner, Mass. 02158. Order No. 50181-0546-2. 45 pp.
- Clench, W. J. and H. van der Schalie. 1944. Notes on naiades from the Green, Salt, and Tradewater Rivers in Kentucky. Michigan Acad. Arts, Sci., Letters. 29:223-228.
- Coker, R. W. and T. Surber. 1911. A note on the metamorphosis of the mussel (Lampsilis laevissimus). Biol. Bull. 20:179-182.
- Conrad, T. A. 1834. New freshwater shells of the United States. Phila., Penn. pp. 1-76.
- Danglade, E. 1922. The Kentucky River and its mussel resources. U.S. Bur. Fish Doc. No. 934. 8 pp.
- Ellis, M. M. 1936. Erosion silt as a factor in aquatic environments. Ecology. 17:29-42.
- Fuller, S. 1974. Clams and mussels (Mollusca: Bivalvia). pp. 215-273. IN: C. W. Hart and S.L.H. Fuller (eds.). Pollution ecology of freshwater invertebrates. Academic Press, New York, New York.
- _____. 1977. Freshwater and terrestrial mollusks. pp. 143-194. IN: J. E. Cooper, S. S. Robinson, and J. B. Funderburg (eds.). Endangered and threatened plants and animals of North Carolina. North Carolina State Museum of Natural History, Raleigh.
- Grace, T. B. and A. C. Buchanan. 1981. Naiades (mussels) of the Lower Osage River, Tavern Creek, and Maries River, Missouri. Missouri Dept. of Conservation. Prepared for U.S. Army Corps of Eng., Kansas City District. 147 pp.
- Heard, W. H. and R. H. Guckert. 1970. A reevaluation of the recent Unionacea (Pelecypoda) of North America. Malacologia. 10(2):333-355.
- Hickman, M. E. 1937. A contribution to mollusca of east Tennessee. Unpublished master's thesis. Dept. of Zool., Univ. of Tennessee, Knoxville. 165 pp. 104 pl.
- Higgins, F. 1858. A catalogue of the shell-bearing species of mollusca of Columbus, Ohio. 1858.
- Hill, D. M., E. A. Taylor, and C. F. Saylor. 1974. Status of faunal recovery in the North Fork Holston River, Tennessee and Virginia. Proc. of the Twenty-eighth Annu. Conf. Southeast Assoc. of Game and Fish Comm. Nov. 17-20, 1974. White Sulphur Springs, WV. pp. 398-414.

- Hinkley, A. A. 1904. List of Alabama shells collected in Oct. and Nov., 1903. *Nautilus*. 18(4):37-45, (5):54-57.
- _____. 1906. Some shells of Mississippi and Alabama. *Nautilus*. 20(3):34-36, 20(4):40-44, 20(5):52-55.
- Hynes, H. B. N. 1970. The ecology of running waters. University of Toronto Press, Toronto, Ontario.
- _____. 1974. The biology of polluted waters. University of Toronto Press, Toronto, Ontario.
- Ingram, W. M. 1956. The use and value of biological indicators of pollution: Fresh water clams and snails. pp. 94-135. IN: C. M. Tarzwell (ed.) Biological problems in water pollution. Taft Sanitary Eng. Center. Cincinnati, OH.
- Isom, B. G. 1971. Effects of storage and mainstream reservoirs on benthic macroinvertebrates in the Tennessee Valley. *Res. Fish, and Limnol. Spec. Publ. No. 8*. Am. Fish. Soc. pp. 179-191.
- _____, P. Yokley, Jr., and C. H. Gooch. 1973. Mussels of the Elk River Basin in Alabama and Tennessee. 1965-67. *Am. Midl. Nat.* 89(2):442-446.
- Jenkinson, J. J. 1981. Endangered or threatened aquatic mollusks of the Tennessee River system. *Bull. Amer. Malacol. Union. for* 1980:43-45.
- Johnson, R. I. 1978. Systematics and zoogeography of Plagiola (=Dysnomia =Epioblasma) an almost extinct genus of freshwater mussels (Bivalvia: Unionidae) from middle North America. *Bull. Mus. Comp. Zool.* 148(6): 239-321.
- _____. 1980. Zoogeography of North American Unionacea (Mollusca: Bivalvia) north of the maximum Pleistocene glaciation. *Bull. Mus. Comp. Zool.* 149(2):189 pp.
- La Rocque, A. 1966-1970. Pleistocene mollusca of Ohio. *Ohio Div. Geol. Surv. Bull. Vol. 62*. 4 pts.
- Lea, I. 1838. Description of new freshwater and land shells. *Trans. Am. Phil. Soc.* 6(1):1-154. pl. 1-24.
- _____. 1857. Descriptions of six new species of Uniones from Alabama. *Proc. Acad. Nat. Sci. Phila.* 9:83.
- _____. 1858. Descriptions of new species of Unio from Tennessee, Alabama, and North Carolina. *Proc. Acad. Nat. Sci. Phila.* 10:40-41.
- _____. 1861. New Unionidae of the United States. *J. Acad. Nat. Sci. Phila.* 5:53-109. pl. 1-18.

- Lefevre, G. and W. C. Curtis. 1910. Experiments in the artificial propagation of freshwater mussels. Bull. U.S. Bur. Fish. 20:615-626.
- Lewis, J. 1876. Fauna of Alabama. Freshwater and land shells. Ala. Geol. Surv. Prog. Rept. 1876:61-100.
- Loar, J. M., L. L. Dye, R. R. Turner, and S. G. Hildebrand. 1980. Analysis of environmental issues related to small-scale hydroelectric development: 1. Dredging. ORNL, Env. Science Div. Publ. No. 1565, Oak Ridge, TN. 134 pp.
- Loosanoff, V. L. 1961. Effects of turbidity on small larval and adult bivalves. Gulf Caribb. Fish. Inst. Univ. Miami Proc. 14:80-95.
- Marsh, P. 1885. List of shells collected in central Tennessee by A. A. Hinkley and P. Marsh with notes on species. Published by author. Alledo, IL. 10 pp.
- Morrison, J. P. E. 1942. Preliminary report on molluscan fauna found in the shell mounds of the Pickwick Landing Basin in the Tennessee River Valley. IN: William S. Webb and David L. DeJarnette, An archaeological survey of Pickwick Basin in the adjacent portions of States of Alabama, Mississippi, and Tennessee. Bur. Am. Ethol. Bull. 129:337-392.
- Neel, J. K. and W. Allen. 1964. The mussel fauna of the upper Cumberland Basin before its impoundment. Malacologia. 1(3):427-459.
- Neves, R. J. 1984. Recovery plan for tan riffle shell mussel (Epioblasma walkeri). USFWS, Region 4, Atlanta, Georgia. 59 pp.
- Ortmann, A. E. 1913. The Alleghenian Divide and its influence upon the freshwater fauna. Proc. Am. Philos. Soc., Phila. 52(210):287-390.
- _____. 1918. The nayades (freshwater mussels) of the upper Tennessee drainage with notes on synonymy and distribution. Proc. Am. Philos. Soc. Phila. 57:521-626.
- _____. 1919. A monograph of the naiades of Pennsylvania. Pt. 3. Systematic account of the genera and species. Mem. Carnegie Mus. 8(1):1-389. pls. 1-21.
- _____. 1920. Correlation of shape and station in freshwater mussels. Proc. Am. Philos. Soc. 19:269-312.
- _____. 1924. The naiad fauna of Duck River in Tennessee. Am. Midl. Nat. (9)1:18-62.
- _____. 1925. The naiad fauna of the Tennessee River system below Walden Gorge. Am. Midl. Nat. 9(7):321-372.
- _____. 1926. The naiades of the Green River drainage in Kentucky. Ann. Carnegie Mus. 17:167-188.

- _____ and B. Walker. 1922. On the nomenclature of certain North American naiades. Occ. Pap. Mus. Zool. Univ. Mich. No. 112. 75 pp.
- Pardue, J. W. 1981. A survey of the mussels (Unionidae) of the upper Tennessee River. 1978. Sterkiana. 71:41-51.
- Parmalee, P. W. 1960. Mussels from the Angel site, Indiana. Nautilus. 74(2):70-75.
- _____. 1967. The freshwater mussels of Illinois. Ill. State Mus. Pop. Series 8:1-108.
- Parmalee, P. W., W. E. Klippel, and A. E. Bogan. 1980. Notes on the pre-historic and present status of the naiad fauna of the middle Cumberland River, Smith County, Tennessee. Nautilus. 94(3):93-105.
- _____. 1982. Aboriginal and modern freshwater mussel assemblages (Pelecypoda: Unionidae) from the Chickamauga Reservoir, Tennessee. Brimleyana. 8:75-90.
- Rafinesque, C. S. 1820. Monographic des coquilles bivalves fluviatiles de la riviere Ohio, contenant douze genres et soixante-huit especies. Ann. Gen. des Sci. Brux. 5:287-322.
- Reeve, L. A. 1843-78. Conchologia Iconica: or illustrations of the shells of molluscous animals. 20 Vols., 2,727 colored plates.
- Schmidt, John. 1982. The freshwater mussels of the Stone's River above J. Persey Priest Reservoir, TN. Master's Thesis, Tenn. Tech. Univ. at Cookeville. 65 pp. Published by U.S. Army Corps of Engineers, Nashville District.
- Shoup, C. S., J. H. Peyton, and G. Gentry. 1941. A limited survey of the Obey River and adjacent streams in Tennessee. Tenn. Dept. Conserv., Div. Game and Fish. Misc. Pub. No. 3. 76 pp.
- Sickel, J. B. 1982. A survey of the freshwater mussels of the lower Cumberland River from Barkley Dam tailwater downstream to the Ohio River. Nashville Dist., U.S. Army Corps of Engineers. 24 pp.
- Simpson, C. T. 1900. Synopsis of the naiades, or pearly freshwater mussels. Proc. U.S. Nat. Mus. 22:501-1044, pl. 18.
- _____. 1914. A descriptive catalogue of the naiades, or pearly freshwater mussels. Bryant Walker. Detroit, MI. 1540 pp.
- Stansbery, D. H. 1961. Species of naiades recorded for central Ohio (Scioto Drainage above Circleville). Ohio State Mus. Zool. at Columbus. 2 pp.
- _____. 1964. The Mussel (Muscle) Shoals of the Tennessee River revisited. Am. Malacol. Union Ann. Rep. 25-28.
- _____. 1970. Eastern freshwater mollusks. (I) The Mississippi and St. Lawrence River systems. IN: A. H. Clarke (ed.). 1970. Rare and endangered mollusks of North America. Malacologia. 13:9-12.

- _____. 1971. Rare and endangered mollusks in eastern United States. IN: S. E. Jorgenson and R. E. Sharp (edg.). Proc. of a symposium on rare and endangered mollusks (naiades) U.S. Bur. Sport Fish. Wildl., U.S. Fish Wildl. Serv., Dept. of Int. Twin Cities, MN. 50 figures. 79 pp.
- _____. 1972. A preliminary list of the naiad shells recovered from the Buffalo site. IN: Bettye J. Broyles, A late 17th century Indian village site (46 Pu 31) in Putnam Co., West Virginia. Rept. of Arch. Inv. No. 5. West Virginia Geol. and Economic Survey, Morganton, WV.
- _____. 1976a. Status of endangered fluviatile mollusks in central North America: *Epioblasma turgidula* (Lea, 1858). Ohio St. Univ. Res. Foundation Rept. to Dept. of Int., Fish and Wildl. Serv., Bur. of Sport Fish. and Wildl. 12 pp.
- _____. 1976b. Naiad mollusks. IN: Endangered and threatened plants and animals of Alabama. Bull. Alabama Mus. Nat. Hist. No. 2:42-52.
- _____. 1980. The naiad mollusks of the Kanawha River below Kanawha Falls with special attention to endangered species (*Bivalvia:Unionida:Unionidae*). Unpublished technical report. Ohio State Univ. Mus. Zool. Columbus, OH. 16 pp.
- Stein, C. B. 1971. Naiad life cycles: their significance in the conservation of the fauna. IN: Rare and endangered mollusks (Naiads) of the U.S. USDOI (Fish and Wildlife Service), Region 3, Federal Building, Fort Snelling. Twin Cities, MN. pp. 19-25.
- Taylor, R. W. 1983. A survey of the freshwater mussels of the Kanawha River. Prepared for Huntington District, Corps of Eng., Huntington, WV. Contract No. DACW69-82-C-0081. 62 pp.
- Tennessee Valley Authority. 1972a. Biological evaluation of the Little Tennessee River Basin Tellico Project. TVA unpublished data. Div. of Env. Res. and Dev. Muscle Shoals, Ala. 15 pp.
- _____. 1972b. Unpublished report, Duck and Buffalo Rivers.
- _____. 1979. An evaluation of mussel populations in the Duck River, Tennessee. May-June 1979. TVA unpublished data. Div. Wat. Res., Fish and Aquatic Ecol. Branch. Norris, TN 37828. 11 pp.
- _____. 1980. The fisheries resource of the Tennessee Valley tailwaters-Tims Ford. TVA unpublished data. Division of Water Resources. Norris, TN. 17 pp.
- Utterback, W. I. 1915. The naiades of Missouri. Am. Midl. Nat. 4(3-8). 200 pp.

- van der Schalie, H. 1939. Additional notes on the naiades (freshwater mussels) of the lower Tennessee River. *Am. Midl. Nat.* 22(2):452-457.
- Vanatta, E. G. 1915. Rafinesque's types of *Unio*. *Proc. Acad. Nat. Sci. Phila.* 67:549-559.
- Walker, B. 1910. Notes on *Truncilla*, with a key to the species. *Nautilus.* 24(7):75-81.
- Warren, R. E. 1975. Prehistoric Unionacean (freshwater mussel) utilization of the Widows Creek site (1Ja305), Northeast Alabama. Unpub. Master's Thesis. Univ. of Neb. Lincoln.
- Wilson, C. B. and H. W. Clark. 1912. The mussel fauna of the Kankakee Basin. Dept. Comm. and Labor. U.S. Bur. Fish Doc. No. 758. 52 pp.
- _____. 1914. The mussels of the Cumberland River and its tributaries. U.S. Bur. of Fish Doc. No. 781. 63 pp.
- Zale, A. V. 1980. The life histories of four freshwater lampsiline mussels (Mollusca: Unionidae) in Big Moccasin Creek, Russel County, Virginia. M.S. Thesis. Virg. Polytech. Inst. State Univ. Blacksburg, VA. 256 pp.

KEY TO IMPLEMENTATION SCHEDULE COLUMNS 1 AND 4

General Category (Column 1):

Information Gathering - I or R (research)

1. Population status
2. Habitat status
3. Habitat requirements
4. Management techniques
5. Taxonomic studies
6. Demographic studies
7. Propagation
8. Migration
9. Predation
10. Competition
11. Disease
12. Environmental contaminant
13. Reintroduction
14. Other information

Management - M

1. Propagation
2. Reintroduction
3. Habitat maintenance and manipulation
4. Predator and competitor control
5. Depradation control
6. Disease control
7. Other management

Acquisition - A

1. Lease
2. Easement
3. Management agreement
4. Exchange
5. Withdrawal
6. Fee title
7. Other

Other - O

1. Information and education
2. Law enforcement
3. Regulations
4. Administration

Priority (Column 4):

- 1 - Those actions absolutely necessary to prevent extinction of the species.
- 2 - Those actions necessary to maintain the species' current population status.
- 3 - All other actions necessary to provide for full recovery of the species.

Tubercled-blossom Pearly Mussel (Epioblasma (=Dysnomia) torulosa torulosa)
 Turgid-blossom Pearly Mussel (Epioblasma (=Dysnomia) turgidula), and Yellow-blossom Pearly Mussel (Epioblasma (=Dysnomia) florentina florentina)
 Part III Implementation Schedule

*1 General Category	Plan Task	Task Number	Priority	Task Duration	Responsible Agency*2		Estimated Fiscal Year Costs *3			Comments/Notes	
					FWS Region	Program	Other	FY 1	FY 2		FY 3
II,2	Conduct intensive surveys for each species.	1	1	3 yr.	3,4,5	SE	THRA, TVA, KDFWR, ADCNR, AGFC, AMHC, MDC, MDR, INPC	25,000	25,000	25,000	*1. See general categories for Implementation Schedules.
I3,M3, 02-4	If populations are found, identify essential habitat and specific areas in need of protection and protect with existing legislation and regulations.	1.1 and 1.2	1 (if populations are found)	Continuous	3,4,5	SE,ES,LE	Same as above	1,000	1,000	1,000	*2. Other agencies' responsibility would be of a cooperative nature or grant program. In some cases contracts could be let to universities or private enterprises.
MI-7, 01	If populations are found, determine present and foreseeable threats and strive to minimize and/or eliminate them.	2	1 (if populations are found)	Unknown	3,4,5	SE,ES	Same as above	Unknown	Unknown	Unknown	*3. Note: Task costs have not been estimated for this plan. This species' present/historic distribution coincides with that of other listed species. Thus, a task aimed at this species will benefit others. Rather than attempting to apportion the costs to each species, recovery tasks will be estimated at a later date when the plans are combined on a watershed basis for implementation.
04	Assess overall status of species.	3	2	Continuous until delisted	3,4,5	SE,ES	Same as above	500	500	500	
Abbreviations:											
SE -	Endangered Species										
ES -	Ecological Services										
LE -	Law Enforcement										
THRA -	Tennessee Wildlife Resources Agency										
THP -	Tennessee Heritage Program										
TVA -	Tennessee Valley Authority										
KDFWR -	Kentucky Department of Fish and Wildlife Resources										
KNPC -	Kentucky Nature Preserves Commission										
WVDNR -	West Virginia Department of Natural Resources										
ADCNR -	Alabama Department of Conservation and Natural Resources										
AGFC -	Arkansas Game and Fish Commission										
AMHC -	Arkansas Natural Heritage Commission										
MDC -	Missouri Department of Conservation										
MDNR -	Missouri Department of Natural Resources										
IDC -	Illinois Department of Conservation										
INPC -	Illinois Nature Preserves Commission										

APPENDIX
List of Reviewers

Mr. Gary Myers
Executive Director
Tennessee Wildlife Resources Agency
Ellington Agricultural Center
Post Office Box 40747
Nashville, Tennessee 37204

Mr. Martin E. Rivers
Environmental Quality Staff
Tennessee Valley Authority
Room 201, Summer Place Building
Knoxville, Tennessee 37902

Mr. Sam Pearsall
Program Coordinator
Tennessee Department of Conservation
Tennessee Heritage Program
701 Broadway
Nashville, Tennessee 37203

Mr. Chuck Cook
The Nature Conservancy
Post Office Box 3017
Nashville, Tennessee 37219

Mr. Ralph Jordan, Jr.
Tennessee Valley Authority
Office of Natural Resources
Forestry Building
Norris, Tennessee 37828

Director
Kentucky Nature Preserves Commission
407 Broadway
Frankfort, Kentucky 40601

Dr. James Henry Wilson
Endangered Species Coordinator
Missouri Department of Conservation
Post Office Box 180
Jefferson City, Missouri 65102

Mr. Harold K. Grimmett
Executive Director
Natural Heritage Commission
Continental Building, Suite 500
Main and Markham
Little Rock, Arkansas 72201

Mr. Keith Guyse
Wildlife Section
Alabama Department of Conservation
and Natural Resources
64 North Union Street
Montgomery, Alabama 36130

Mr. Sam Barkley
Endangered Species Biologist
Arkansas Game and Fish Commission
#2 Natural Resources Drive
Little Rock, Arkansas 72205

Mr. Harold E. Alexander
Endangered Species Coordinator
Arkansas Game and Fish Commission
#2 Natural Resources Drive
Little Rock, Arkansas 72205

Dr. Devere E. Burt, Chairman
Endangered Species and Populations
Committee
Ohio Biological Survey
Director, Cincinnati Museum of
Natural History
1720 Gilbert Avenue
Cincinnati, Ohio 45202

Mr. Peter E. Zurbuch
Assistant Chief
Wildlife Resources
Department of Natural Resources
Post Office Box 67
Elkins, West Virginia 26241

Mr. Michael Sweet
Endangered Species Program
Coordinator
Department of Conservation
Division of Forest Resources and
Natural Heritage
524 South Second
Lincoln Tower Plaza
Springfield, Illinois 62706

Mr. Larry R. Gale, Director
Department of Conservation
P.O. Box 180
Jefferson City, Missouri 65102

Mr. Fred Lafser, Director
 Department of Natural Resources
 P.O. Box 176
 Jefferson City, Missouri 65102

Mr. John McGregor
 Kentucky Department of Fish and
 Wildlife Resources
 #1 Game Farm Road
 Frankfort, Kentucky 40601

Mr. Steven A. Ahlstedt
 Field Operations
 Division of Water Resources
 Forestry Building
 Norris, Tennessee 37828

Mr. Herbert D. Athearn
 Route 5, Box 376
 Cleveland, Tennessee 37311

Mr. John M. Bates
 Ecological Consultants, Inc.
 1900 Dexter Avenue
 Ann Arbor Michigan 48103

Dr. Arthur E. Bogan
 Department of Malacology
 Academy of Natural Sciences
 Nineteenth and the Parkway
 Philadelphia, Pennsylvania 19103

Mr. Alan C. Buchanan
 Missouri Department of Conservation
 Fish and Wildlife Research Center
 110 College Avenue
 Columbia, Missouri 65201

Dr. Arthur H. Clarke
 7 Hawthorne Street
 Mattapoisett, Massachusetts 02739

Mr. George M. Davis
 Academy of Natural Sciences
 19th and the Parkway
 Philadelphia, Pennsylvania 19103

Mr. Steve N. Wilson, Director
 Game and Fish Commission
 #1 Natural Resources Drive
 Little Rock, Arkansas 72205

Ms. Sally D. Dennis
 Center of Environmental Studies
 Virginia Polytechnic Institute
 and State University
 Blacksburg, Virginia 24061

Mr. Samuel L.H. Fuller
 Department of Limnology
 Academy of Natural Sciences
 19th and the Parkway
 Philadelphia, Pennsylvania 19103

Mr. John Jenkinson
 Tennessee Valley Authority
 Evans Building
 Knoxville, Tennessee 37902

Mr. Jack M. Hoffman, Chief
 Fish Division
 Commission of Game
 and Inland Fisheries
 4010 West Broad Street
 Box 11104
 Richmond, Virginia 23230

Mr. Robert V. Davis, Executive Director
 State Water Control Board
 P.O. Box 11143
 Richmond, Virginia 23230

Dr. Richard Neves
 Virginia Cooperative Fishery Unit
 106 Cheatham Hall
 Virginia Polytechnic Institute
 Blacksburg, Virginia 24061

Dr. and Mrs. Wayne C. Starnes
 TVA Forestry Building
 Norris, Tennessee 37828

Mr. Howard Larsen, Regional Director
 U.S. Fish and Wildlife Service
 One Gateway Center
 Newton Corner, Massachusetts 02138

Mr. Anthony J. Campbell
 Executive Director
 Tennessee Conservation League
 1720 West End Avenue
 Suite 600
 Nashville, Tennessee 37203

Virginia Wildlife Federation
Box 1780
Norfolk, Virginia 23501

Dr. R. Don Estes, Leader
Tennessee Cooperative
Fishery Research Unit
Tennessee Technological University
Box 5063
Cookeville, Tennessee 38501

Mr. John Hardcastle
Chapter Chairman
The Nature Conservancy
Capitol Hill Building 114
301 7th Ave., North
Nashville, Tennessee 37219

Mr. Bob Hatcher, Nongame Biologist
Tennessee Wildlife Resources Agency
Ellington Agriculture Center
P.O. Box 40747
Nashville, Tennessee 37204

Mr. D.W. Yambert
Nongame Biologist
Tennessee Wildlife Resources Agency
Route 3, Box 153-A
Talbott, Tennessee 37203

Mr. Dan Eagars, Zoologist
Tennessee Heritage Program
2611 West End Avenue
Nashville, Tennessee 37203

Dr. Tom Ripley, Manager
Office of Natural Resources
Tennessee Valley Authority
Locust Street Building
Knoxville, Tennessee 37902

Mr. Richard Fitz
Tennessee Valley Authority
Locust Street Building
Knoxville, Tennessee 37902

Mr. Charles Gooch
Field Operations
Tennessee Valley Authority
A 251 401 Building
Chattanooga, Tennessee 37401

Dr. William H. Heard
Department of Biology
Florida State University
Tallahassee, Florida 32306

Dr. John C. Hurd
Science Department
La Grange College
La Grange, Georgia 30240

Dr. Marc J. Imlay
Columbia National Fisheries
Research Laboratory
U.S. Fish and Wildlife Service
Route 1,
Columbia, Missouri 65201

Mr. Billy G. Isom
Fisheries and Aquatic Ecology Branch
Division of Water Resources
Tennessee Valley Authority
E & D Building
Muscle Shoals, Alabama 35660

Dr. Eugene P. Keferl
Division of Natural Science
Brunswick Junior College
Brunswick, Georgia 34520

Dr. Paul W. Parmalee
Department of Anthropology
The University of Tennessee
Knoxville, Tennessee 37916

Dr. Hugh J. Porter
Institute of Marine Sciences
University of North Carolina
P.O. Drawer 809
Morehead City, North Carolina 28557

Dr. James B. Sickle
Department of Biology
Murray State University
Murray, Kentucky 40271

Dr. David H. Stansbery
Museum of Zoology
Ohio State University
1813 North High Street
Columbus, Ohio 43210

Dr. Carol B. Stein
Museum of Zoology
Ohio State University
1813 North High Street
Columbus, Ohio 43210

Dr. Edward M. Stern
Department of Biology
University of Wisconsin
at Stevens Point
Stevens Point, Wisconsin 54481

Dr. Fred G. Thompson
Florida State Museum
Museum Road
University of Florida
Gainesville, Florida 32611

Dr. Henry Van der Schalie
15000 Buss Road
Manchester, Michigan 48158

Dr. John D. Williams
Department of Biology
Eastern Kentucky State University
Richmond, Kentucky 23219

Dr. Paul Yokley, Jr.
Department of Biology
University of North Alabama
Florence, Alabama 35630

Mr. David Kenney, Director
Department of Conservation
Lincoln Tower Plaza
524 South Second Street
Springfield, Illinois 62706

Mr. Frank H. Beal, Director
Institute of Natural Resources
Division of Environmental Management
309 West Washington Street
Chicago, Illinois 60606

Ms. Mary Lou Marzuki, Chairman
Nature Preserves Commission
320 South Third Street
Rockford, Illinois 61108

Mr. Paul G. Risser, Chief
State Natural History Survey Division
172 Natural Resources Building
607 East Peabody Drive
Champaign, Illinois 61820

Mr. Carl E. Kays, Commissioner
Department of Fish
and Wildlife Resources
#1 Game Farm Road
Frankfort, Kentucky 40601

Ms. Jackie Swigart, Secretary
Department of Natural Resources
and Environmental Protection
5th Floor, Capital Plaza Tower
Frankfort, Kentucky 40601

Mr. David C. Callaghan, Director
Department of Natural Resources
1800 Washington Street, East
Charleston, West Virginia 25305

Regional Director
U.S. Fish and Wildlife Services
Twin Cities Regional Office
Federal Building, Fort Snelling
Twin Cities, Minnesota 55111

Mr. John Saxon, Chairman
Department of Pollution Control
and Ecology
8001 National Drive
P.O. Box 9583
Little Rock, Arkansas 72219

Ms. Susan Brenholts, Director
Department of Arkansas Natural
and Cultural Heritage
Continental Building, Suite 500
Main and Markham
Little Rock, Arkansas 72201