



**U.S. Fish and Wildlife Service Pacific Region**  
**Columbia River Basin Hatchery Review Team**

**Columbia River Basin, Mountain Snake Province**  
***Clearwater and Salmon River Watersheds***



***Dworshak, Kooskia and Hagerman National Fish Hatcheries***  
***Clearwater, Magic Valley, McCall, and Sawtooth Fish Hatcheries***  
***(Idaho Department of Fish & Game)***

**Assessments and Recommendations**

**Final Report, Appendix A:**  
**All-H Analyzer (AHA) Output for Salmon and Steelhead Stocks**

**March 2011**

*Please cite as:*

U.S. Fish and Wildlife Service (USFWS). 2011. *Dworshak, Kooskia, and Hagerman National Fish Hatcheries and Clearwater, Magic Valley, McCall, and Sawtooth State Fish Hatcheries Assessments and Recommendations - Appendix A: All-H Analyzer (AHA) Output for Salmon and Steelhead Stocks. Final Report, March 2011*. Hatchery Review Team, Pacific Region. U.S. Fish and Wildlife Service, Portland, Oregon. Available at: <http://www.fws.gov/Pacific/fisheries/Hatcheryreview/reports.html>.

## **Appendix A: All-H Analyzer (AHA) Output for Spring/Summer Chinook Salmon and Summer Steelhead Stocks at Dworshak NFH, Kooskia NFH, Hagerman NFH (USFWS), and Clearwater FH, Magic Valley FH, McCall FH, and Sawtooth FH (IDFG), including natural populations in the Clearwater and Salmon rivers, Idaho.**

### *What is AHA?*

*AHA* is an *Excel*-based spreadsheet simulation model that quantifies the mean number of adult salmon or steelhead returning to a watershed after many generations (years) of reproduction and migration based on equilibrium, or near equilibrium, conditions. Recent versions of *AHA* (Versions 6.x and higher) for the Columbia River allocate returning adult fish to one of six locations: (1) a hatchery and other recapture facilities where adult fish are removed; (2) the natural habitat where adult fish spawn; (3) harvest in the marine environment; (4) harvest in the lower Columbia River mainstem downstream from Bonneville Dam; (5) harvest in the upper Columbia River mainstem upstream from Bonneville Dam, and (6) harvest in terminal areas in the general areas where they originated as smolts.

The *AHA* model was originally developed primarily by Lars Mobrand (Mobrand Biometrics, Vashon Island, WA), in collaboration with the Washington Department of Fish and Wildlife (WDFW) and the Northwest Indian Fisheries Commission (NWIFC), as part of the HSRG review<sup>1</sup> of salmon and steelhead hatcheries in Puget Sound and coastal Washington state. Updated versions that summarize results over all populations (Version 12.1 and higher) were developed by Greg Blair (Jones, Stokes, and Mobrand; Vashon Island, WA) in collaboration with Lars Mobrand.

*AHA* is based on the Beverton-Holt spawner-recruit model where habitat *capacity* is the asymptote of the Beverton-Holt curve (i.e., maximum number of adult recruits that the habitat can produce) and *productivity* is the slope of the curve at the origin (i.e., the maximum number of adult recruits per adult spawner [ $R/S$ ] when density-dependent factors or competition can be ignored). The actual model (spreadsheet) consists of several component modules (e.g., natural component module, hatchery component module, genetic fitness module, etc.) where estimated mean values of biological and population dynamic parameters are provided by the user (e.g. mean fecundity of females, estimated egg-to-smolt survival, etc.). The genetic fitness function is based on the model of Ford (2002)<sup>2</sup> and allows the mean fitness of a population (productivity) to change incrementally over time depending on (a) the mean proportion of natural spawners composed of hatchery-origin adults ( $p_{HOS}$ ) relative to the mean proportion of the hatchery broodstock composed of natural-origin adults ( $p_{NOB}$ ), and (b) the number of generations that hatchery-origin fish spawn naturally in the watershed.

A useful index for assessing the genetic impact of a hatchery program on a natural population is the *Proportionate Natural Influence* =  $PNI = p_{NOB}/(p_{NOB} + p_{HOS})$ , where  $PNI$  varies

---

<sup>1</sup> [www.hatcheryreform.org](http://www.hatcheryreform.org)

<sup>2</sup> Ford, M.J. 2002. Selection in captivity during supportive breeding may reduce fitness in the wild. *Conservation Biology* 16: 815-825.

## USFWS Columbia Basin Hatchery Review Team

### *Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

from 0.0 (high hatchery influence, low natural environmental influence) to 1.0 (no hatchery influence). When  $pHOS = pNOB$ , the hatchery and natural environments will have approximately equal influence on the genetic constitutions of both hatchery and wild fish, and  $PNI = 0.5$ . For integrated hatchery populations and natural stocks, the goal is for  $PNI$  to be greater than 0.5 and as close to 1.0 as possible (e.g.,  $> 0.67$ ) as a long-term goal.

The AHA model is currently being used by fishery comanagers in the Pacific Northwest as a “planning tool” to (a) document assumptions and parameters for hatchery and natural populations and (b) compare the outcomes of alternative management strategies and scenarios. Only those scenarios achieving realistic or desired outcomes are considered valid. For example, any scenario that results in extirpation of a stock is considered invalid, where any or all of the four H’s<sup>3</sup> can contribute to stock extirpation.

*For more detail on AHA, see the “Tools” page of the Hatchery Scientific Review Group’s (HSRG’s) website: <http://www.hatcheryreform.us>. An AHA user’s guide, the AHA model, and all AHA datasets for the Columbia River Basin are available for download. See also “HSRG White Paper No. 1” under the publications page for theoretical details regarding  $PNI$  and the relative effects of the hatchery and natural environments on mean population fitness.*

### ***Explanation of Tables A1-A56<sup>4</sup>***

Information in the following AHA output tables (Tables A1-A56) is intended to provide a summary “snapshot” of the potential equilibrium outcomes associated with current hatchery programs and alternatives considered by the Hatchery Review Team. Five sets of simulations or scenarios are presented in each table: (a) current situation and hatchery programs; (b) the potential “no hatchery” condition; (c) the “hatchery solution” proposed by the HSRG; and (d) two alternative scenarios proposed by the HRT. AHA version 13.2.1 was used to generate the outputs in Tables A1-A56.

Output of AHA is displayed in a series of colored bar graphs representing the equilibrium, mean numbers of adult fish (recruits) allocated to each of the six aforementioned locations (hatchery, freshwater habitat, and four harvest locations). In those bar graphs, solid green represents natural-origin fish; solid pink represents hatchery-origin fish, pink diagonal hash bars represent hatchery-origin fish in excess of comanager goals, and pink vertical bars represent hatchery-origin fish that have been selected at least one generation in the natural environment (e.g. as occurs in a genetically-integrated hatchery program).

The graph in the lower-left portion of each table shows the realized mean values of  $pNOB$  and  $pHOS$  for each of the five scenarios (Current, No Hatchery, etc.). The diagonal lines represent combination values of  $pNOB$  and  $pHOS$  that yield the same value of  $PNI$ . Symbols on the  $PNI$  graph correspond with each of the scenarios shown as bar graphs (see also the *Components of*

---

<sup>3</sup> *Habitat, Harvest, Hydropower, and Hatcheries are commonly referred to as the “four H’s”.*

<sup>4</sup> *Parameter estimates used to generate the following tables have not all been verified and should be considered preliminary. However, their values are based on the best information available, and the general results presented in the following tables are not expected to change significantly as the parameter estimates are verified and updated.*

## USFWS Columbia Basin Hatchery Review Team

### *Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011*

*this Report* section for definitions of biological significance, population viability, habitat, and harvest ratings).

The outputs presented in Tables A1-A56 are intended to be viewed electronically on a desktop computer where portions of the tables can be “zoomed in” for clarity. The Review Team acknowledges that those tables are difficult to read as printed paper copies.

*AHA* data files in V.13.2.1 are stored as “roll-up” files for all populations of each salmonid species in the Columbia River Basin. The specific parameterized versions of the “roll-up” files and *AHA* (V.13.2.1) used to generate the outputs in Tables A1-A56 are available upon request from the Chair of the Hatchery Review Team. Some familiarity with *AHA* is required, and expected, before implementing the software. The software and baseline data files for Tables A1-A56 can be downloaded from the HSRG website under the “Tools” tab at [www.hatcheryreform.us](http://www.hatcheryreform.us).

# USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

## **Tables A1-A9: AHA outputs for the Clearwater River Major Population Group (MPG) of spring-summer Chinook salmon**

# USFWS Columbia Basin Hatchery Review Team

Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011

## Parameters and assumptions for spring/summer Chinook in the Clearwater River Major Population Group (MPG)

Productivity and survival parameters for hatchery origin spring Chinook within the Clearwater River basin (Table 1A) were obtained from the HSRG, as provided by Idaho Department of Fish and Game (IDFG).

**Table 1A. Mean values of reproductive and survival parameters assumed in the AHA analyses for hatchery-origin spring/summer Chinook released within the Clearwater River MPG.**

Hatchery/ Release location <i>Population/Stock</i>	Female Fecundity (No. of eggs)	Within- hatchery survival (egg-to- smolt)	Post-release survival (%) (smolt-to-adult)	Recruit/spawner (R/S; adults)
Dworshak NFH <i>Dworshak NFH Spring Chinook</i>	4,500	0.79	0.58%	7.0
Kooskia NFH <i>Kooskia NFH Spring Chinook</i>	4,500	0.79	0.58%	7.0
Clearwater FH/ Powell Satellite, Lochsa River <i>Lochsa River Spring Chinook</i>	4,200	0.90	0.44%	6.0
Clearwater FH, NPTH/ Lower Selway River, <i>Lochsa River Spring Chinook</i>	4,500	Smolts = 0.79 Parr = 0.85	Smolts = 0.26% Smolt+parr=0.14%	Smolts = 3.2 Parr = 0.7
Clearwater Hatchery/ Upper Selway River, <i>Lochsa River Spring Chinook</i>	4,500	Parr = 0.85	Parr = 0.07%	Parr = 0.9
Clearwater Hatchery/ Crooked and Red River Satellites <i>S.F. Clearwater R. Spring Chinook</i>	4,200	0.70	0.37%	4.0
Nez Perce Tribal Hatchery <i>NPTH Spring Chinook</i>	4,500	0.79	0.35%	4.2
Nez Perce Tribal Hatchery/ Newsome Creek <i>Newsome Creek Spring Chinook</i>	4,200	Smolts = 0.70 Pre-smolts = 0.79	Smolts = 0.37% Pre-smolts= 0.09%	Smolts = 4.0 Pre-smolts = 1.0
Nez Perce Tribal Hatchery/ Lolo Creek <i>Lolo Creek Spring Chinook</i>	4,500	Smolts = 0.70 Pre-smolts = 0.79	Smolts = 0.37% Pre-smolts= 0.08%	Smolts = 4.0 Pre-smolts = 1.0

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

### *Current programs, HSRG solutions, and HRT alternatives*

The HRT modeled *Alternative 1* (current hatchery programs with implementation of all HRT recommendations) and the Team's preferred alternatives for each of the spring Chinook hatchery programs conducted within the Clearwater River basin. Each population release site consists of five scenarios: *Current* program, *No Hatchery* scenario, *HSRG Solution*, *HRT Alternative 1*, and the *HRT Preferred Alternative* (Table 2A). The *HRT Preferred Alternative* assumes that reestablishment of naturally spawning populations is the principal short-term goal of spring Chinook hatchery programs in the Lochsa, Selway, and upper South Fork Clearwater rivers.



# USFWS Columbia Basin Hatchery Review Team

## Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011

**Table 2A. Summary of the number of hatchery-origin (HOR) spring Chinook smolts, pre-smolts, and parr released in the Clearwater River under four strategies modeled in AHA Tables A1-A9: (1) *Current* programs are based on the LSRCP Cooperators 2008 Annual Operating Plan for the Clearwater River, (2) the *HSRG Solution*, (3) *HRT Alternative 1* (current programs with implementation of HRT recommendations), and (4) the *HRT Preferred Alternative* which emphasizes reestablishment of naturally spawning populations in the Lochsa, Selway, and upper S.F. Clearwater rivers and emphasizes harvest in the mainstem Clearwater River. The names of hatchery stocks/populations are in italics (e.g., “*Dworshak*”, “*Kooskia*”, etc.) and are identified by the facility/location where adults are trapped for broodstock. The Hatchery Review Team did not specifically review programs associated with the Nez Perce Tribal Hatchery (e.g., Lolo Creek, Newsome Creek), but the AHA outputs for those programs are presented here as part of the data set packages available from the HSRG ([www.hatcheryreform.us](http://www.hatcheryreform.us)).**

Release location	Current	HSRG Solution	HRT Alternative 1	HRT Preferred Alternative
Dworshak NFH, N.F. Clearwater River (AHA Table A1)	• 1.05 M <i>Dworshak</i> smolts	No recommendation (same as Current)	Same as current	1.45M <i>Dworshak</i> smolts
Kooskia NFH, Clear Creek (AHA Table A2)	• 600K <i>Kooskia</i> smolts	No recommendation (same as Current)	• 500K <i>Kooskia</i> smolts	Terminate on-station rearing; • 600K <i>Kooskia</i> smolts reared at Clearwater Hatchery.
S.F. Clearwater River, Red and Crooked rivers (AHA Table A3)	• 1.1M <i>SF Clearwater</i> smolts	No recommendation (same as Current)	Same as Current	• 300K <i>Red/Crooked R.</i> smolts ( <i>pNOB</i> = 0.67)
Newsome Creek (AHA Table A4)	• 30K <i>Newsome Cr.</i> pre-smolts • 45K <i>NPTH</i> pre-smolts	• 75K <i>Newsome Cr.</i> smolts ( <i>pNOB</i> = 1.0)	No recommendation (Same as HSRG)	No recommendation (Same as HSRG)
Lolo Creek (AHA Table A5)	• 85K <i>Lolo Cr.</i> pre-smolts • 65K <i>NPTH</i> pre-smolts	• 100K <i>Lolo Cr.</i> smolts ( <i>pNOB</i> = 1.0)	No recommendation (Same as HSRG)	No recommendation (Same as HSRG)
Nez Perce Tribal Hatchery ( <i>NPTH</i> ) (AHA Table A6)	• 125K <i>NPTH</i> smolts	No recommendation (same as Current)	No recommendation (same as Current)	No recommendation (same as Current)
Lochsa River at Powell Pond (AHA Table A7)	• 400K <i>Lochsa</i> smolts	• 700K <i>Lochsa</i> smolts	• 400K <i>Lochsa</i> smolts	• 300K <i>Lochsa</i> smolts ( <i>pNOB</i> = 0.75)
Lower Selway River, including Meadow Cr. (AHA Table A8)	• 300K <i>Lochsa</i> smolts • 400K <i>Dworshak</i> parr (released into Meadow Creek)	No recommendation (same as Current)	• 300K <i>Lochsa</i> smolts	• 200K <i>Lochsa</i> smolts
Upper Selway River (AHA Table A9)	• 300K <i>Lochsa</i> parr	• 100K <i>Selway</i> smolts ( <i>pNOB</i> = 1.0)	• 300K <i>Lochsa</i> parr	• 200K <i>Lochsa</i> parr



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A2. AHA output for Kooskia NFH spring Chinook, Clear Creek, and Middle Fork Clearwater River

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Clearwater

Spring Chinook

6A-Clearwater\_Middle Fork Clearwater

Spring Chinook (Kooskia-Hatchery)

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Reintroduction: Harvest

Segregated hatchery

Not listed; excluded from ESU

Reintroduction: Conservation

Natural Reproduction Only

Not listed; excluded from ESU

Reintroduction: Harvest

Segregated hatchery

Not listed; excluded from ESU

Reintroduction: Harvest

Segregated hatchery

Not listed; excluded from ESU

Reintroduction: Harvest

Segregated hatchery

Not listed; excluded from ESU

Habitat

Productivity (Adult)

Min NOR Escape

Smolt Productivity

Ad. Capacity

% Kelt

Sm. Capacity

0.01

0

1

1

0.01

0

1

1

No Hatchery

0.01

0

1

1

HSRG Solution

0.01

0

1

1

HRT Alternative #1

0.01

0

1

1

HRT Preferred Alternative

0.01

0

1

1

Hydro

Ocean Surv

Juv Passage Surv.

Adjusted Productivity

Baseline SAR

Vary? (Y/N)

Adult Passage

Adj. Capacity

0.092

0.008

1.00

0

0.029

0.008

0.34

0

0.029

0.008

0.34

0

0.029

0.008

0.34

0

0.029

0.008

0.34

0

0.029

0.008

0.34

0

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

HORs

NORs

HORs

NORs

HORs

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.300

0.083

0.413

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

400

600,652

90%

10%

7.0

y

400

600,652

90%

10%

7.0

y

400

600,652

90%

10%

7.0

y

400

600,652

90%

10%

7.0

y

400

600,652

90%

10%

7.0

y

400

600,652

90%

10%

7.0

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Habitat

Hydro

Harvest


Hatchery


Current Habitat


Current Hydro


Current Harvest


Realized Spawning Composition











Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

80

80%

0.81

0.50

0.50

12

0.5

0.5%

0.8%

80

80%

0.81

1.00

0.99

30

1.00

0.8%

80

80%

0.81

0.50

0.50

12

1.00

0.5%

0.8%

80

80%

0.81

0.50

0.50

12

1.00

0.5%

0.8%

80

80%

0.81

0.50

0.50

12

1.00

0.5%

0.8%

NOR Escapement

HoS Total Escapement (includes strays)

Max

Min

Ave

0

0

0

559

97

183

0

0

0

559

97

183

0

0

0

467

81

153

559

97

183

USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A3. AHA output for South Fork Clearwater River Spring Chinook, including Red and Crooked rivers (Clearwater Anadromous Fish Hatchery)

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Clearwater

Spring Chinook

4-Clearwater\_South Fork Clearwater Spring Chinook

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Reintroduction: Harvest and Conservation

Integrated Hatchery

Not listed; excluded from ESU

Reintroduction: Harvest and Conservation

Natural Reproduction only

Not listed; excluded from ESU

Reintroduction: Harvest and Conservation

Segregated Hatchery

Not listed; excluded from ESU

Reintroduction: Harvest and Conservation

Segregated Hatchery

Not listed; excluded from ESU

Reintroduction: Conservation

Integrated Hatchery

Not listed; excluded from ESU

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative #1

HRT Preferred Alternative

Productivity (Adult)

Min NOR Escape

Smolt Productivity

Ad. Capacity

% Kelt

Sm. Capacity

1.30

1

161

1,875

231,905

1.30

1

161

1,875

231,905

1.30

1

161

1,875

231,905

1.30

1

161

1,875

231,905

1.30

1

161

1,875

231,905

1.30

1

161

1,875

231,905

Hydro

Ocean Surv

Juv Passage Surv.

Adjusted Productivity

Baseline SAR

Vary? (Y/N)

Adult Passage

Adj. Capacity

0.092

1.00

1.30

0.008

1.00

1,875

y

0.82

1,875

0.029

0.34

1.30

0.008

0.82

1,875

y

0.82

1,875

0.029

0.34

1.30

0.008

0.82

1,875

y

0.82

1,875

0.029

0.34

1.30

0.008

0.82

1,875

y

0.82

1,875

0.029

0.34

1.30

0.008

0.82

1,875

y

0.82

1,875

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

HORs

NORs

HORs

NORs

HORs

0.010

0.020

0.050

0.100

0.170

0.010

0.080

0.080

0.350

0.455

0.010

0.020

0.050

0.100

0.170

0.010

0.080

0.080

0.350

0.455

0.010

0.020

0.050

0.100

0.170

0.010

0.080

0.080

0.350

0.455

0.010

0.020

0.050

0.100

0.170

0.010

0.080

0.080

0.350

0.455

0.010

0.020

0.050

0.100

0.170

0.010

0.080

0.080

0.350

0.455

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

pHOS-Goal

Cons/Harv/Both

Int/Seg/Step/None

Local

Imported

Export Goal/Realized

% to Nat. Spawn

Recruits/Spawner

Fitness? [Y / N]

Both

Int

Cons

None

Both

Seg

Both

Seg

Both

Seg

Cons

Int

832

13

90%

4.0

1,100,736

10%

y

1,100,736

22

10%

y

1,100,736

16

10%

y

300,861

27

90%

4.0

300,861

27

90%

4.0

300,861

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro


Current Harvest


Current Habitat

Current Hydro

HRT -Pref.

Realized Spawning Composition





Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

80

80%

0.81

0.50

0.99

16

0.5

0.3%

0.8%

Max

Min

Ave

301

13

93

661

59

186

685

119

224

666

116

218

1,571

-2

311

80

80%

0.81

1.00

0.50

30

<







# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A4. AHA output for Newsome Creek Spring Chinook, S.F. Clearwater River (Nez Perce Tribal Hatchery)

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Reintroduction: Conservation		Reintroduction: Conservation		Reintroduction: Conservation		Reintroduction: Conservation		Reintroduction: Conservation												
Clearwater		Spring Chinook		Harv&Hatchery Strategy:		Integrated + supplementation		Natural Reproduction only		Integrated + supplementation		Integrated + supplementation		Integrated + supplementation												
4-Clearwater_South Fork				Population Recovery Designation:		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU												
Clearwater_Newsome Creek Spring																										
Chinook				Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative												
Habitat	Productivity (Adult)	Ad. Capacity	1.30	625	1.30	625	1.30	625	1.30	625	1.30	625	1.30	625												
	Min NOR Escape	% Kelt	1		1		1		1		1		1													
	Smolt Productivity	Sm. Capacity	161	77,302	161	77,302	161	77,302	161	77,302	161	77,302	161	77,302												
Hydro	Ocean Surv	Baseline SAR	0.092	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008												
	Juv Passage Surv.	Vary? (Y/N)	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82												
	Adjusted Productivity	Adj. Capacity	1.30	625	1.30	625	1.30	625	1.30	625	1.30	625	1.30	625												
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010												
	Harv - L. Mainstem	NORs		HORs	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020												
	Harv - U. Mainstem	NORs		HORs	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050												
	Harv - Terminal	NORs		HORs	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005												
	Total Exploitation Rate	NORs		HORs	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083												
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	10%		pNOB	pHOS	100%		pNOB	pHOS	100%													
	Purpose		PNI	pHOS-Realized	0.08	43%			0.64	57%			0.64	57%												
	Type		Cons/Harv/Both	Int/Seg/Step/None	Cons	Int	Cons	None	Cons	Int	Cons	Int	Cons	Int												
	Broodstock by Source		Local	Imported	22	35	75,411		57	75,279	57	75,279	57	75,279												
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release			11			20		14		29												
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	50%	50%		100%		100%		100%		100%												
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	1.0	y		y	4.0	y	4.0	y	4.0	y												
	Update																									
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat												
		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro												
		Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		HRT-Pref.												
		Hatchery																								
Realized Spawning Composition																										
     																										
Relative Hatchery Optimum ->					80		80		80		80		80													
Weir Factor ->																										
Relative Reproductive Success (HOS) ->					80%		80%		80%		80%		80%													
Initial Fitness Factor (A) ->					0.81		0.81		0.81		0.81		0.81													
Fitness Factor after 100 generations (B) ->					0.50		1.00		0.77		0.77		0.77													
Average Fitness Factor (100 Generations) ->					0.50		0.99		0.78		0.78		0.78													
Generations until average fitness is reached ->					20		30		10		10		10													
Minimum Hatchery Program (% of BS Goal) ->																										
"Fitness Floor" ->					0.5																					
Calculated Hatchery SAR ->					0.1%				0.3%		0.3%		0.3%													
Calculated Natural SAR ->					0.8%		0.8%		0.8%		0.8%		0.8%													
					Max	Min	Ave	Max	Min	Ave	Max	Min	Ave													
NOR Escapement					146	10	41	239	19	89	724	17	154	714												
HoS Total Escapement (includes strays)					122	21	39				772	134	252	755												
											755	131	247	801												
														139												
														262												

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A5. AHA output for Lolo Creek Spring Chinook, Clearwater River (Nez Perce Tribal Hatchery)

All H Analyzer (AHA)												
Apply & Recalculate												
Population (Ctrl + Page Up/Down to Scroll)												
Subregion/Subbasin			Species/Race			Population Management Intent:			Reintroduction: Conservation			
Clearwater			Spring Chinook			Harv&Hatchery Strategy:			Integrated + Supplementation			
5-Clearwater_Lolo Creek Spring Chinook			Population Recovery Designation:			Not listed; excluded from ESU			Not listed; excluded from ESU			
			Historic			Current			No Hatchery			
			HSRG Solution			HRT Alternative #1			HRT Preferred Alternative			
Habitat	Productivity (Adult)		Ad. Capacity		1.30		1,500		1.30		1,500	
	Min NOR Escape		% Kelt		1		1		1		1	
	Smolt Productivity		Sm. Capacity		161		185,524		161		185,524	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)		0.092		0.008		y		0.029	
	Juv Passage Surv.		Adult Passage		1.00		1.00		0.34		0.82	
	Adjusted Productivity		Adj. Capacity		1.30		1,500		1.30		1,500	
Harvest	Harv - Marine		NORs		HORs		0.010		0.010		0.010	
	Harv - L. Mainstem		NORs		HORs		0.020		0.020		0.020	
	Harv - U. Mainstem		NORs		HORs		0.050		0.050		0.050	
	Harv -Terminal		NORs		HORs		0.005		0.005		0.005	
	Total Exploitation Rate		NORs		HORs		0.083		0.083		0.083	
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB	
	Purpose		PNI		pHOS-Realized		Cons		Int		Cons	
	Type		Cons/Harv/Both		Int/Seg/Step/None		64		48		100,990	
	Broodstock by Source		Local		Imported		76		100,990		76	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Smolt Release		22		16		30	
	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		1.0		y		4.0	
Patterns	Update		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat	
			Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro	
			Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest	
			Hatchery									

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A6. AHA output for Nez Perce Tribal Hatchery Spring Chinook, Clearwater River

All H Analyzer (AHA)														
Apply & Recalculate														
Population (Ctrl + Page Up/Down to Scroll)														
Subregion/Subbasin		Species/Race		Population Management Intent:		Reintroduction: Harvest and Conservation		Reintroduction: Harvest and Conservation		Reintroduction: Harvest and Conservation		Reintroduction: Harvest and Conservation		
Clearwater		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		
6C-Clearwater_Lower Mainstem_Spring Chinook (NPTH-Hatchery)		Population Recovery Designation:		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative		
Habitat	Productivity (Adult)	Ad. Capacity	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0		
	Min NOR Escape	% Kelt	1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	0	1	0	1	0	1	0	1	0	1		
Hydro	Ocean Surv	Baseline SAR	0.092	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008		
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82		
	Adjusted Productivity	Adj. Capacity	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0		
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010		
	Harv - L. Mainstem	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010		
	Harv - U. Mainstem	NORs		HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140		
	Harv -Terminal	NORs		HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030		
	Total Exploitation Rate	NORs		HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182		
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI		pHOS-Realized		Harv		100%		Harv		100%	
	Type		Cons/Harv/Both		Int/Seg/Step/None		Cons		None		Cons		None	
	Broodstock by Source		Local		Imported		83		120		83		120	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays		120		10%		120		10%	
Patterns	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		90%		10%		90%		10%	
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		4.2		y		4.2		y	
	Update		Habitat		Current Habitat									
			Hydro		Current Hydro									
			Harvest		Current Harvest									
		Hatchery												

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011

Table A7. AHA output for Lochsa River Spring Chinook (Powell Satellite Facility and Clearwater Fish Hatchery)

All H Analyzer (AHA)										Apply & Recalculate																																																																																																															
Population (Ctrl + Page Up/Down to Scroll)																																																																																																																									
Subregion/Subbasin		Species/Race		Population Management Intent:		Reintroduction: Harvest and Conservation		Reintroduction: Conservation		Reintroduction: Harvest and Conservation		Reintroduction: Harvest and Conservation		Reintroduction: Conservation																																																																																																											
Clearwater		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery + Imports		Natural Reproduction Only		Segregated Hatchery		Segregated Hatchery		Integrated Hatchery																																																																																																											
1-Clearwater_Lochsa Spring Chinook				Population Recovery Designation:		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU		Not listed; excluded from ESU																																																																																																											
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative																																																																																																													
Habitat	Productivity (Adult)	Ad. Capacity	1.30	940	1.30	940	1.30	940	1.30	940	1.30	940	1.30	940																																																																																																											
	Min NOR Escape	% Kelt	1		1		1		1		1		1																																																																																																												
	Smolt Productivity	Sm. Capacity	161	116,265	161	116,262	161	116,262	161	116,262	161	116,262	161	116,262																																																																																																											
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.008	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y																																																																																																										
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82																																																																																																										
	Adjusted Productivity	Adj. Capacity		1.30	940	1.30	940	1.30	940	1.30	940	1.30	940	1.30	940																																																																																																										
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010																																																																																																										
	Harv - L. Mainstem		NORs	HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020																																																																																																										
	Harv - U. Mainstem		NORs	HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050																																																																																																										
	Harv - Terminal		NORs	HORs	0.050	0.200	0.050	0.200	0.050	0.200	0.050	0.200	0.050	0.200	0.050																																																																																																										
	Total Exploitation Rate		NORs	HORs	0.124	0.330	0.124	0.330	0.124	0.330	0.124	0.330	0.124	0.330	0.124																																																																																																										
Hatchery	Broodstock Composition		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS		pNOB																																																																																																										
	Purpose		PNI		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB																																																																																																										
	Type		Cons/Harv/Both		pHOS-Realized		Cons		None		Cons		None		Cons																																																																																																										
	Broodstock by Source		Local		Imported		Both		Seg		Both		Seg		Both																																																																																																										
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Smolt Release		236		401,436		412		700,812		236																																																																																																										
Patterns	Destination for HOR Returns		90%		10%		80%		20%		80%		20%		30%																																																																																																										
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		6.0		y		6.0		y		6.0																																																																																																										
	Update		Habitat		Current Habitat		Habitat		Current Habitat		Habitat		Current Habitat		Habitat																																																																																																										
			Hydro		Current Hydro		Hydro		Current Hydro		Hydro		Current Hydro		Hydro																																																																																																										
			Harvest		Current Harvest		Harvest		Current Harvest		Harvest		Current Harvest		Harvest																																																																																																										
Realized Spawning Composition																																																																																																																									
<table><tr><td>Relative Hatchery Optimum -&gt;</td><td>80</td><td>80</td><td>80</td><td>80</td><td>80</td></tr><tr><td>Weir Factor -&gt;</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Relative Reproductive Success (HOS) -&gt;</td><td>80%</td><td>80%</td><td>80%</td><td>80%</td><td>80%</td></tr><tr><td>Initial Fitness Factor (A) -&gt;</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td></tr><tr><td>Fitness Factor after 100 generations (B) -&gt;</td><td>0.50</td><td>1.00</td><td>0.50</td><td>0.50</td><td>0.63</td></tr><tr><td>Average Fitness Factor (100 Generations) -&gt;</td><td>0.50</td><td>0.99</td><td>0.50</td><td>0.50</td><td>0.66</td></tr><tr><td>Generations until average fitness is reached -&gt;</td><td>17</td><td>30</td><td>15</td><td>16</td><td>38</td></tr><tr><td>Minimum Hatchery Program (% of BS Goal) -&gt;</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>"Fitness Floor" -&gt;</td><td>0.5</td><td></td><td></td><td></td><td></td></tr><tr><td>Effective pHOS -&gt;</td><td>0.47</td><td></td><td>0.59</td><td>0.53</td><td>0.74</td></tr><tr><td>pNOB -&gt;</td><td></td><td></td><td></td><td></td><td>0.37</td></tr><tr><td>Calculated Hatchery SAR -&gt;</td><td>0.4%</td><td></td><td>0.4%</td><td>0.4%</td><td>0.4%</td></tr><tr><td>Calculated Natural SAR -&gt;</td><td>0.8%</td><td></td><td>0.8%</td><td>0.8%</td><td>0.8%</td></tr><tr><td colspan="6">NOR Escapement</td></tr><tr><td>Max</td><td>371</td><td>33</td><td>105</td><td>262</td><td>17</td></tr><tr><td>Min</td><td>358</td><td>62</td><td>117</td><td>1,172</td><td>203</td></tr><tr><td>Ave</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>HoS Total Escapement (includes strays)</td><td></td><td></td><td></td><td></td><td></td></tr></table>														Relative Hatchery Optimum ->	80	80	80	80	80	Weir Factor ->						Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%	Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81	Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.63	Average Fitness Factor (100 Generations) ->	0.50	0.99	0.50	0.50	0.66	Generations until average fitness is reached ->	17	30	15	16	38	Minimum Hatchery Program (% of BS Goal) ->						"Fitness Floor" ->	0.5					Effective pHOS ->	0.47		0.59	0.53	0.74	pNOB ->					0.37	Calculated Hatchery SAR ->	0.4%		0.4%	0.4%	0.4%	Calculated Natural SAR ->	0.8%		0.8%	0.8%	0.8%	NOR Escapement						Max	371	33	105	262	17	Min	358	62	117	1,172	203	Ave						HoS Total Escapement (includes strays)					
Relative Hatchery Optimum ->	80	80	80	80	80																																																																																																																				
Weir Factor ->																																																																																																																									
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%																																																																																																																				
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81																																																																																																																				
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.63																																																																																																																				
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.50	0.50	0.66																																																																																																																				
Generations until average fitness is reached ->	17	30	15	16	38																																																																																																																				
Minimum Hatchery Program (% of BS Goal) ->																																																																																																																									
"Fitness Floor" ->	0.5																																																																																																																								
Effective pHOS ->	0.47		0.59	0.53	0.74																																																																																																																				
pNOB ->					0.37																																																																																																																				
Calculated Hatchery SAR ->	0.4%		0.4%	0.4%	0.4%																																																																																																																				
Calculated Natural SAR ->	0.8%		0.8%	0.8%	0.8%																																																																																																																				
NOR Escapement																																																																																																																									
Max	371	33	105	262	17																																																																																																																				
Min	358	62	117	1,172	203																																																																																																																				
Ave																																																																																																																									
HoS Total Escapement (includes strays)																																																																																																																									





# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports – March 2011


Table A9. AHA output for Upper Selway River Spring Chinook, Lochsa River Stock (Clearwater Fish Hatchery)


All H Analyzer (AHA)												
Apply & Recalculate												
Population (Ctrl + Page Up/Down to Scroll)												
Subregion/Subbasin		Species/Race		Population Management Intent:		Reintroduction: Conservation		Reintroduction: Conservation		Reintroduction: Harvest and Conservation		
Clearwater		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery + Supplementation		Natural Reproduction only		Integrated Hatchery + Supplementation		
3-Clearwater_Upper Selway Spring Chinook				Population Recovery Designation:		Not listed: excluded from ESU		Not listed: excluded from ESU		Not listed: excluded from ESU		
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		
Habitat	Productivity (Adult)	Ad. Capacity	1.30	600	1.30	600	1.30	600	1.30	600	1.30	600
	Min NOR Escape	% Kelt	1		1		1		1		1	
	Smolt Productivity	Sm. Capacity	161	74,442	161	74,210	161	74,210	161	74,210	161	74,210
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.008	y	0.029	0.008	y	0.029	0.008	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34
	Adjusted Productivity	Adj. Capacity		1.30	600	1.30	600	1.30	600	1.30	600	600
Harvest	Harv - Marine			NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem			NORs	HORs	0.020	0.020	0.020	0.020	0.020	0.020	0.020
	Harv - U. Mainstem			NORs	HORs	0.050	0.050	0.050	0.050	0.050	0.050	0.050
	Harv -Terminal			NORs	HORs	0.005	0.005	0.005	0.005	0.005	0.005	0.005
	Total Exploitation Rate			NORs	HORs	0.083	0.083	0.083	0.083	0.083	0.083	0.083
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB	
	Purpose		PNI		pHOS-Realized		Both		Cons		None	
	Type		Cons/Harv/Both		Int/Seg/Step/None		Seg		None		Seg	
	Broodstock by Source		Local		Imported		186		300,516		68	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Smolt Release		10		100%		16	
	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		100%		100%		100%	
Patterns	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		0.9		y		y	
	Hatchery		Hatchery		Hatchery		Hatchery		Hatchery		Hatchery	
	Hatchery		Hatchery		Hatchery		Hatchery		Hatchery		Hatchery	
Update	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Hatchery		Hatchery		Hatchery		Hatchery		Hatchery	


700


5


Realized Spawning Composition

















Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.74	0.50	0.50
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.75	0.50	0.50
Generations until average fitness is reached ->	16	30	22	16	16
Minimum Hatchery Program (% of BS Goal) ->					
"Fitness Floor" ->	0.5				
Calculated Hatchery SAR ->	0.1%				
Calculated Natural SAR ->	0.8%				
NOR Escapement	459	46	125	394	108
HoS Total Escapement (includes strays)	552	95	180	430	141

**Tables A10-A17: *AHA* outputs for Clearwater River Major Population Group (MPG) of summer steelhead**

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

### *Parameters and assumptions for summer steelhead in the Clearwater River Major Population Group (MPG)*

Productivity and survival parameters for hatchery-origin steelhead released within the Clearwater River basin (Table 3A) were obtained from the HSRG, as provided by Idaho Department of Fish and Game (IDFG).

**Table 3A. Mean values of reproductive and survival parameters assumed in the AHA analyses for hatchery-origin steelhead released within the Clearwater River MPG.**

<b>Hatchery/ Release location Population/Stock</b>	<b>Female Fecundity (No. of eggs)</b>	<b>Within- hatchery survival (egg-to-smolt)</b>	<b>Post-release survival (%) (smolt-to- adult)</b>	<b>Recruit/spawner (R/S; adults)</b>
Dworshak NFH <i>Dworshak NFH Steelhead</i>	6,848	0.74	0.78%	35.0
Dworshak NFH/ Clear Creek, Kooskia NFH <i>Dworshak NFH Steelhead</i>	6,848	0.74	0.78%	35.0
Dworshak NFH, Clearwater FH/ S.F. Clearwater River, Red House <i>Dworshak NFH Steelhead</i>	6,848	0.74	0.78%	35.0
Dworshak NFH, Clearwater FH/ Upper S.F. Clearwater River <i>Dworshak NFH Steelhead</i>	6,848	0.74	0.78%	35.0
Clearwater FH/ Lolo Creek <i>Dworshak NFH Steelhead</i>	6,848	0.74	0.78%	35.0

### *Current programs, HSRG solutions, and HRT alternatives*

The HRT modeled *Alternative 1* and its preferred alternatives for each of the steelhead hatchery programs conducted within the Clearwater River basin. Each population release site consists of five scenarios: *Current program*, *No Hatchery* scenario, *HSRG Solution*, *HRT Alternative 1*, and the *HRT Preferred Alternative* (Table 4A). The *HRT Preferred Alternative* assumes that reestablishment of naturally spawning populations is the principal goal of hatchery programs in the upper South Fork Clearwater River, including natural spawning locations in Crooked River, Red River, American River, Newsome Creek, Meadow Creek, and Mill Creek. The *HRT Preferred Alternative* emphasizes harvest in the lower S.F. Clearwater River (Red House release site) and mainstem Clearwater River downstream from Clear Creek and Kooskia NFH.

# USFWS Columbia Basin Hatchery Review Team

## Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011

**Table 4A. Summary of the number of hatchery-origin (HOR) steelhead smolts released in the Clearwater River under four strategies modeled in AHA Tables A10-A17: (1) *Current* programs based on the LSRCP Cooperators 2008 Annual Operating Plan for the Clearwater River, (2) the *HSRG Solution*, (3) *HRT Alternative 1* (current programs with implementation of HRT recommendations), and (4) *HRT Preferred Alternative*. All hatchery stocks are considered “*B-run*”, and are primarily of Dworshak NFH origin. Release sites in the upper S.F. Clearwater River were pooled in the analyses (Red River, Crooked River, Newsome Creek, American River, Meadow Creek, and Mill Creek). The names of hatchery stocks/populations are in italics (e.g., “*Dworshak*”) and are identified by the facility/location where adults are trapped for broodstock.**

Release location	Current	HSRG Solution	HRT Alternative 1	HRT Preferred Alternative
Dworshak NFH, N.F. Clearwater River (AHA Table A10)	• 1.20M <i>Dworshak</i> smolts	No recommendation (same as Current)	Same as Current	Same as Current
Kooskia NFH, Clear Creek (AHA Table A11)	• 300K <i>Dworshak</i> smolts	• 300K <i>Kooskia</i> smolts	Same as HSRG	Same as HSRG
Lower S.F. Clearwater River at Red House (AHA Table A12)	• 665K <i>Dworshak</i> smolts	• 1.16M <i>Dworshak</i> smolts	• 1.4M <i>Dworshak</i> smolts	• 1.085M <i>Dworshak</i> smolts (release 315K <i>Dworshak</i> smolts in the Little Salmon River)
Upper S.F. Clearwater River, (AHA Table A13)	• 735K <i>Dworshak</i> smolts	• 124K <i>Upper S.F.</i> smolts ( <i>pNOB</i> = 1.0)	Same as HSRG	Same as HSRG
Lolo Creek (AHA Table A14)	• 50K <i>Dworshak</i> smolts	• 50K <i>Lolo Cr.</i> smolts ( <i>pNOB</i> = 1.0)	No recommendation (Same as HSRG)	No recommendation (Same as HSRG)

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011


Table A10. AHA output for Dworshak NFH B-run Summer Steelhead, North Fork Clearwater River


All H Analyzer (AHA)															
Apply & Recalculate															
Population (Ctrl + Page Up/Down to Scroll)															
Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Harvest and Conservation		Harvest and Conservation		Harvest and Conservation		Harvest and Conservation	
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery + outplanting		Natural Reproduction only		Segregated Hatchery + outplanting		Segregated Hatchery + outplanting		Segregated Hatchery + outplanting	
5-Clearwater_NF Clearwater Summer Steelhead (B-Run-Hatchery)				Population Recovery Designation:		N/A		N/A		N/A		N/A		N/A	
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative	
Habitat	Productivity (Adult)		Ad. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0
	Min NOR Escape		% Kelt	1		1		1		1		1		1	
	Smolt Productivity		Sm. Capacity	1	1	1	1	1	1	1	1	1	1	1	1
Hydro	Ocean Surv		Baseline SAR	0.098	0.018	y		0.048	0.018	y		0.048	0.018	y	
	Juv Passage Surv.		Adult Passage	1.00	1.00			0.49	0.77			0.49	0.77		
	Adjusted Productivity		Adj. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0
Harvest	Harv - Marine		NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem		NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - U. Mainstem		NORs		HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140
	Harv -Terminal		NORs		HORs	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600
	Total Exploitation Rate		NORs		HORs	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal										
	Purpose		PNI		pHOS-Realized		100%				100%				100%
	Cons/Harv/Both		Int/Seg/Step/None			Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg
	Broodstock by Source		Local	484	Imported	1,500	1,500			484	1,500	1,500	1,500	484	1,500
	Broodstock Exported (from HOR Surplus)		Export Goal/Realized	90%	Strays <td>90%</td> <td>10%</td> <td></td> <td>100%</td> <td>90%</td> <td>10%</td> <td>90%</td> <td>10%</td> <td>90%</td> <td>10%</td>	90%	10%		100%	90%	10%	90%	10%	90%	10%
Patterns	Destination for HOR Returns		% to Hatchery	35.0	% to Nat. Spawn.		y		y		y		y		y
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]										
	Update														
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.															
Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat	
Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro	
Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest	
Hatchery															

480

7

Realized Spawning Composition





Relative Hatchery Optimum ->		80	80	80	80	80							
Weir Factor ->													
Relative Reproductive Success (HOS) ->		25%	25%	25%	25%	25%							
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->		0.50	1.00	0.50	0.50	0.50							
Average Fitness Factor (100 Generations) ->		0.50	0.99	0.50	0.50	0.50							
Generations until average fitness is reached ->		12	30	12	12	12							
Minimum Hatchery Program (% of BS Goal) ->		0.5											
Effective pHOS ->		1.00	1.00	1.00	1.00	1.00							
"Fitness Floor" ->													
Calculated Hatchery SAR ->		1.4%											
Calculated Natural SAR ->		1.8%											
NOR Escapement		Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)		1,946	338	637				1,946	338	637	1,946	338	637

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A11. AHA output for Clear Creek B-run Summer Steelhead, Middle Fork Clearwater River (Dworshak NFH)


All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest mitigation		Harvest mitigation		Harvest mitigation		Harvest mitigation		Harvest mitigation	
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery	
6A-Clearwater_Lower Clearwater Summer Steelhead (B-Run-Hatchery)		Population Recovery Designation:		N/A		N/A		N/A		N/A		N/A		N/A	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative			
Habitat	Productivity (Adult)	Ad. Capacity	2.00	100	2.00	100	2.00	100	2.00	100	2.00	100	2.00	100	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	50	2,500	110	5,522	110	5,522	110	5,522	110	5,522	110	5,522	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.098	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity	0.91	45	2.00	100	2.00	100	2.00	100	2.00	100	2.00	100	
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem		NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	
	Harv -Terminal		NORs	HORs	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	
	Total Exploitation Rate		NORs	HORs	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized		74%				81%		56%		56%	
	Cons/Harv/Both		Int/Seg/Step/None		Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg	
	Broodstock by Source		Local	Imported	122	302,936			122	302,936	122	302,936	122	302,936	
	Broodstock by Source		Export Goal/Realized	Smolt Release											
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	45%	55%		100%	45%	55%	90%	10%	90%	10%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	20.0	y		y	35.0	y	35.0	y	35.0	y	
	Update		Habitat	Hydro	Harvest	Hatchery									

Realized Spawning Composition



Relative Hatchery Optimum ->		80	80	80	80	80							
Weir Factor ->													
Relative Reproductive Success (HOS) ->		25%	25%	25%	25%	25%							
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->		0.50	1.00	0.50	0.50	0.50							
Average Fitness Factor (100 Generations) ->		0.50	0.99	0.50	0.50	0.50							
Generations until average fitness is reached ->		13	30	13	15	15							
Minimum Hatchery Program (% of BS Goal) ->		0.5	Effective pHOS ->	0.74	Effective pHOS ->	0.81							
"Fitness Floor" ->		0.5	pNOB ->		pNOB ->								
Calculated Hatchery SAR ->		0.8%			1.4%								
Calculated Natural SAR ->		1.8%			1.8%								
NOR Escapement		Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)		153	22	45	109	12	34	167	25	50	114	13	32
		1,541	267	504				2,697	468	883	490	85	160

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A12. AHA output for Lower South Fork Clearwater River *B-run* Summer Steelhead (Dworshak NFH and Clearwater FH)


All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest	
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery	
3-Clearwater_SF Clearwater Summer Steelhead (B-Run)				Population Recovery Designation:		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing	
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative	
Habitat	Productivity (Adult)	Ad. Capacity	1.50	350	1.50	350	1.50	350	1.50	350	1.50	350	1.50	350	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	75	17,500	83	19,326	83	19,326	83	19,326	83	19,326	83	19,326	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.018		0.048	0.018	y	0.048	0.018	y	0.048	0.018	y	
	Juv Passage Surv.	Adult Passage				0.49	0.77		0.49	0.77		0.49	0.77		
	Adjusted Productivity	Adj. Capacity	1.36	317	1.50	350	1.50	350	1.50	350	1.50	350	1.50	350	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	
	Harv -Terminal	NORs	HORs	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.030	
	Total Exploitation Rate	NORs	HORs	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.182	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
			PNI	pHOS-Realized		85%				90%		91%		89%	
	Purpose	Type	Cons/Harv/Both	Int/Seg/Step/None	Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg	
	Broodstock by Source		Local	Imported	268	665,467			467	1,159,600	564	1,400,458	438	1,087,592	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		268				240		168		168	
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
Patterns	Update		Habitat	Hydro	Harvest	Hatchery									

Realized Spawning Composition



Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	25%	80%	25%	25%	25%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.50										
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.50	0.50	0.50										
Generations until average fitness is reached ->	13	30	13	13	13										
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.85	Effective pHOS ->	0.90										
"Fitness Floor" ->		pNOB ->		pNOB ->											
Calculated Hatchery SAR ->	1.4%			1.4%	1.4%										
Calculated Natural SAR ->	1.8%			1.8%	1.8%										
NOR Escapement	Max 573	Min 86	Ave 172	Max 136	Min 12	Ave 52	Max 613	Min 97	Ave 189	Max 623	Min 101	Ave 193	Max 608	Min 96	Ave 186
HoS Total Escapement (includes strays)	11,594	2,013	3,796				19,509	3,388	6,389	23,189	4,027	7,594	18,122	3,147	5,935



USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A13. AHA output for Upper South Fork Clearwater River B-run Summer Steelhead (Dworshak NFH and Clearwater FH)







All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation and Harvest		Conservation		Conservation and Harvest		Conservation		Conservation				
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Integrated Hatchery		Integrated Hatchery		Integrated Hatchery				
3-Clearwater_SF Clearwater_Crooked River Summer Steelhead (B-Run)		Population Recovery Designation:		Contributing		Contributing		Contributing		Contributing		Contributing		Contributing				
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative						
Habitat	Productivity (Adult)	Ad. Capacity	1.50	400	1.50	400	1.50	400	1.50	400	1.50	400	1.50	400	400			
	Min NOR Escape	% Kelt	1		1		1		1		1		1					
	Smolt Productivity	Sm. Capacity	75	20,000	83	22,087	83	22,087	83	22,087	83	22,087	83	22,087	22,087			
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.018		0.048	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y	
	Juv Passage Surv.	Adult Passage			0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.77	
	Adjusted Productivity	Adj. Capacity		1.36	362	1.50	400	1.50	400	1.50	400	1.50	400	1.50	400	1.50	400	
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem		NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	
	Harv -Terminal		NORs	HORs	0.030	0.200	0.030	0.200	0.030	0.200	0.030	0.200	0.030	0.200	0.030	0.200	0.200	
	Total Exploitation Rate		NORs	HORs	0.182	0.326	0.182	0.326	0.182	0.326	0.182	0.326	0.182	0.326	0.182	0.326	0.326	
Hatchery	Broodstock Composition		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Type		Cons/Harv/Both		pHOS-Realized		Cons		None		Cons		Int		Cons		Int	
	Broodstock by Source		Local		Smolt Release		50		123,906		50		123,906		50		123,906	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays		300		61		20%		80%		20%		80%	
Patterns Update	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		35.0		100%		35.0		80%		35.0		80%	
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		35.0		y		35.0		y		35.0		y	
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		
		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		
		Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		
		Hatchery																

Realized Spawning Composition



Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	25%	80%	80%	80%	80%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.70	0.70	0.70										
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.71	0.71	0.71										
Generations until average fitness is reached ->	13	30	35	35	35										
Minimum Hatchery Program (% of BS Goal) ->	0.5														
Effective pHOS ->	0.90	0.90	0.83	0.83	0.83										
"Fitness Floor" ->	0.5														
Calculated Hatchery SAR ->	1.4%	1.4%	1.4%	1.4%	1.4%										
Calculated Natural SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%										
NOR Escapement	704	113	217	155	14	59	723	63	181	724	63	182	723	63	181
Hos Total Escapement (includes strays)	23,910	4,153	7,830				3,398	590	1,112	3,436	596	1,125	3,386	588	1,108

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A14. AHA output for Lolo Creek A+B-run Summer Steelhead (Clearwater Fish Hatchery)

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Harvest and Conservation		Harvest and Conservation		Harvest and Conservation	
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery + supplementation		Natural Reproduction only		Integrated Hatchery + supplementation		Integrated Hatchery + supplementation		Integrated Hatchery + supplementation	
4-Clearwater_Lolo Summer Steelhead (A+B-Run)		Population Recovery Designation:		Contributing		Contributing		Contributing		Contributing		Contributing		Contributing	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative			
Habitat	Productivity (Adult)	Ad. Capacity	2.00	200	2.00	200	2.00	200	2.00	200	2.00	200	2.00	200	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	100	10,000	110	11,043	110	11,043	110	11,043	110	11,043	110	11,043	
Hydro	Ocean Surv	Baseline SAR	0.098	0.018	0.048	0.018	0.048	0.018	0.048	0.018	0.048	0.018	0.048	0.018	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity	1.81	181	2.00	200	2.00	200	2.00	200	2.00	200	2.00	200	
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs		HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	
	Harv -Terminal	NORs		HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
	Total Exploitation Rate	NORs		HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized											
	Cons/Harv/Both		Int/Seg/Step/None	Smolt Release											
	Broodstock by Source		Local	Imported	20	49,662			20	49,413	20	49,413	20	49,413	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		139				107		102		85	
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	10%	90%		100%		100%		100%	50%	50%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	35.0	y	y	y	35.0	y	35.0	y	35.0	y	
	Update		Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Hydro		Harvest		Hatchery				
		Current Habitat		Current Hydro		Current Harvest									



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A15. AHA output for Lower Clearwater River A-run Summer Steelhead

All H Analyzer (AHA)



Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation																																																																																																								
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only																																																																																																								
6-Clearwater_Lower Clearwater Summer Steelhead (A-Run)		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary																																																																																																								
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative																																																																																																										
Habitat	Productivity (Adult)	Ad. Capacity	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430																																																																																																								
	Min NOR Escape	% Kelt	1		1		1		1		1		1																																																																																																									
	Smolt Productivity	Sm. Capacity	261	71,500	261	71,511	261	71,511	261	71,511	261	71,511	261	71,511																																																																																																								
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.020	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020																																																																																																							
	Juv Passage Surv.	Adult Passage		0.49	0.49	0.77		0.49	0.49	0.77		0.49	0.49	0.77																																																																																																								
	Adjusted Productivity	Adj. Capacity	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430	5.21	1,430																																																																																																								
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010																																																																																																								
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010																																																																																																								
	Harv - U. Mainstem		NORs	HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060																																																																																																								
	Harv -Terminal		NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030																																																																																																								
	Total Exploitation Rate		NORs	HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106																																																																																																								
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS																																																																																																								
	Purpose		PNI	pHOS-Realized	None	1%	None	1%	None	1%	None	1%	None	1%																																																																																																								
	Type		Cons/Harv/Both	Int/Seg/Step/None	None	None	None	None	None	None	None	None	None	None																																																																																																								
	Broodstock by Source		Local	Imported																																																																																																																		
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		58		76		40		40		40																																																																																																								
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%																																																																																																								
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	35.0	y	35.0	y	35.0	y	35.0	y	35.0	y																																																																																																								
	Update		Habitat	Hydro	Harvest																																																																																																																	
Realized Spawning Composition																																																																																																																						
																																																																																																																						
																																																																																																																						
<table><tr><td>Relative Hatchery Optimum -&gt;</td><td>80</td><td>80</td><td>80</td><td>80</td><td>80</td></tr><tr><td>Weir Factor -&gt;</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Relative Reproductive Success (HOS) -&gt;</td><td>25%</td><td>25%</td><td>25%</td><td>25%</td><td>25%</td></tr><tr><td>Initial Fitness Factor (A) -&gt;</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td></tr><tr><td>Fitness Factor after 100 generations (B) -&gt;</td><td>0.91</td><td>1.00</td><td>0.86</td><td>0.95</td><td>0.95</td></tr><tr><td>Average Fitness Factor (100 Generations) -&gt;</td><td>0.92</td><td>0.99</td><td>0.89</td><td>0.95</td><td>0.95</td></tr><tr><td>Generations until average fitness is reached -&gt;</td><td>18</td><td>30</td><td>11</td><td>25</td><td>25</td></tr><tr><td>Minimum Hatchery Program (% of BS Goal) -&gt;</td><td>0.5</td><td></td><td></td><td></td><td></td></tr><tr><td>Effective pHOS -&gt;</td><td>0.01</td><td>0.01</td><td>0.02</td><td>0.01</td><td>0.01</td></tr><tr><td>"Fitness Floor" -&gt;</td><td>0.5</td><td></td><td></td><td></td><td></td></tr><tr><td>Calculated Hatchery SAR -&gt;</td><td>3.5%</td><td></td><td></td><td>3.5%</td><td>3.5%</td></tr><tr><td>Calculated Natural SAR -&gt;</td><td>2.0%</td><td>2.0%</td><td>2.0%</td><td>2.0%</td><td>2.0%</td></tr><tr><td>NOR Escapement</td><td>Max 3,501</td><td>Min 528</td><td>Ave 1,056</td><td>Max 3,695</td><td>Min 567</td><td>Ave 1,115</td><td>Max 3,393</td><td>Min 504</td><td>Ave 1,025</td><td>Max 3,585</td><td>Min 546</td><td>Ave 1,081</td><td>Max 3,585</td><td>Min 546</td><td>Ave 1,081</td></tr><tr><td>HOS Total Escapement (includes strays)</td><td>176</td><td>31</td><td>58</td><td>234</td><td>41</td><td>76</td><td>123</td><td>21</td><td>40</td><td>123</td><td>21</td><td>40</td><td>123</td><td>21</td><td>40</td></tr></table>															Relative Hatchery Optimum ->	80	80	80	80	80	Weir Factor ->						Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%	Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81	Fitness Factor after 100 generations (B) ->	0.91	1.00	0.86	0.95	0.95	Average Fitness Factor (100 Generations) ->	0.92	0.99	0.89	0.95	0.95	Generations until average fitness is reached ->	18	30	11	25	25	Minimum Hatchery Program (% of BS Goal) ->	0.5					Effective pHOS ->	0.01	0.01	0.02	0.01	0.01	"Fitness Floor" ->	0.5					Calculated Hatchery SAR ->	3.5%			3.5%	3.5%	Calculated Natural SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%	NOR Escapement	Max 3,501	Min 528	Ave 1,056	Max 3,695	Min 567	Ave 1,115	Max 3,393	Min 504	Ave 1,025	Max 3,585	Min 546	Ave 1,081	Max 3,585	Min 546	Ave 1,081	HOS Total Escapement (includes strays)	176	31	58	234	41	76	123	21	40	123	21	40	123	21	40
Relative Hatchery Optimum ->	80	80	80	80	80																																																																																																																	
Weir Factor ->																																																																																																																						
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%																																																																																																																	
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81																																																																																																																	
Fitness Factor after 100 generations (B) ->	0.91	1.00	0.86	0.95	0.95																																																																																																																	
Average Fitness Factor (100 Generations) ->	0.92	0.99	0.89	0.95	0.95																																																																																																																	
Generations until average fitness is reached ->	18	30	11	25	25																																																																																																																	
Minimum Hatchery Program (% of BS Goal) ->	0.5																																																																																																																					
Effective pHOS ->	0.01	0.01	0.02	0.01	0.01																																																																																																																	
"Fitness Floor" ->	0.5																																																																																																																					
Calculated Hatchery SAR ->	3.5%			3.5%	3.5%																																																																																																																	
Calculated Natural SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%																																																																																																																	
NOR Escapement	Max 3,501	Min 528	Ave 1,056	Max 3,695	Min 567	Ave 1,115	Max 3,393	Min 504	Ave 1,025	Max 3,585	Min 546	Ave 1,081	Max 3,585	Min 546	Ave 1,081																																																																																																							
HOS Total Escapement (includes strays)	176	31	58	234	41	76	123	21	40	123	21	40	123	21	40																																																																																																							

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A16. AHA output for Lochsa River B-run Summer Steelhead

All H Analyzer (AHA)															
Apply & Recalculate															
Population (Ctrl + Page Up/Down to Scroll)															
Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only	
1-Clearwater_Lochsa Summer Steelhead (B-Run)		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative			
Habitat	Productivity (Adult)	Ad. Capacity	2.50	2,000	2.50	2,000	2.50	2,000	2.50	2,000	2.50	2,000	2.50	2,000	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	125	100,000	138	110,434	138	110,434	138	110,434	138	110,434	138	110,434	
Hydro	Ocean Surv	Baseline SAR	0.098	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity	2.26	1,811	2.50	2,000	2.50	2,000	2.50	2,000	2.50	2,000	2.50	2,000	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	
	Harv -Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
	Total Exploitation Rate	NORs	HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized	None	5%	None	5%	None	5%	None	5%	None	5%	
	Cons/Harv/Both		Int/Seg/Step/None	None	None	None	None	None	None	None	None	None	None	None	
	Broodstock by Source		Local	Imported											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		20		32		10		10		10	
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	35.0	y	35.0	y	35.0	y	35.0	y	35.0	y	
	Update		Habitat	Hydro	Harvest										
			Hatchery												
Realized Spawning Composition															
															
															
Relative Hatchery Optimum -> 80															
Weir Factor ->															
Relative Reproductive Success (HOS) -> 25%															
Initial Fitness Factor (A) -> 0.81															
Fitness Factor after 100 generations (B) -> 0.98															
Average Fitness Factor (100 Generations) -> 0.97															
Generations until average fitness is reached -> 32															
Minimum Hatchery Program (% of BS Goal) -> 0.5															
Effective pHOS -> 0.01															
"Fitness Floor" ->															
Calculated Hatchery SAR -> 1.2%															
Calculated Natural SAR -> 1.8%															
NOR Escapement															
HoS Total Escapement (includes strays)															

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A17. AHA output for B-run Selway River Summer Steelhead

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation																																																																																																								
Clearwater		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only																																																																																																								
2-Clearwater_Selway Summer Steelhead (B-Run)		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary																																																																																																								
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative #1		HRT Preferred Alternative																																																																																																										
Habitat	Productivity (Adult)	Ad. Capacity	2.50	2,500	2.50	2,500	2.50	2,500	2.50	2,500	2.50	2,500	2.50	2,500																																																																																																								
	Min NOR Escape	% Kelt	1		1		1		1		1		1																																																																																																									
	Smolt Productivity	Sm. Capacity	125	125,000	138	138,046	138	138,046	138	138,046	138	138,046	138	138,046																																																																																																								
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.098	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y																																																																																																							
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77																																																																																																							
	Adjusted Productivity	Adj. Capacity		2.26	2,264	2.50	2,500	2.50	2,500	2.50	2,500	2.50	2,500	2.50	2,500																																																																																																							
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010																																																																																																							
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010																																																																																																							
	Harv - U. Mainstem		NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140																																																																																																							
	Harv -Terminal		NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030																																																																																																							
	Total Exploitation Rate		NORs	HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182																																																																																																							
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS																																																																																																						
	Purpose		PNI	pHOS-Realized	5%	5%	5%	5%	5%	5%	5%	5%	5%	5%	5%	5%																																																																																																						
	Type		Cons/Harv/Both	Int/Seg/Step/None	None	None	None	None	None	None	None	None	None	None	None	None																																																																																																						
	Broodstock by Source		Local	Imported																																																																																																																		
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		19		30		8		8		8		8																																																																																																						
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%																																																																																																						
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	35.0	y	35.0	y	35.0	y	35.0	y	35.0	y	35.0	y																																																																																																						
	Update		Habitat	Hydro	Harvest																																																																																																																	
Realized Spawning Composition																																																																																																																						
<table><tr><td>Relative Hatchery Optimum -&gt;</td><td>80</td><td>80</td><td>80</td><td>80</td><td>80</td></tr><tr><td>Weir Factor -&gt;</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Relative Reproductive Success (HOS) -&gt;</td><td>25%</td><td>25%</td><td>25%</td><td>25%</td><td>25%</td></tr><tr><td>Initial Fitness Factor (A) -&gt;</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td><td>0.81</td></tr><tr><td>Fitness Factor after 100 generations (B) -&gt;</td><td>0.99</td><td>1.00</td><td>0.97</td><td>1.00</td><td>1.00</td></tr><tr><td>Average Fitness Factor (100 Generations) -&gt;</td><td>0.98</td><td>0.99</td><td>0.96</td><td>0.99</td><td>0.99</td></tr><tr><td>Generations until average fitness is reached -&gt;</td><td>33</td><td>30</td><td>30</td><td>32</td><td>32</td></tr><tr><td>Minimum Hatchery Program (% of BS Goal) -&gt;</td><td>0.5</td><td></td><td></td><td></td><td></td></tr><tr><td>Effective pHOS -&gt;</td><td>0.00</td><td>0.00</td><td>0.01</td><td>0.00</td><td>0.00</td></tr><tr><td>"Fitness Floor" -&gt;</td><td>0.5</td><td></td><td></td><td></td><td></td></tr><tr><td>Calculated Hatchery SAR -&gt;</td><td>1.2%</td><td>1.2%</td><td>1.2%</td><td>1.2%</td><td>1.2%</td></tr><tr><td>Calculated Natural SAR -&gt;</td><td>1.8%</td><td>1.8%</td><td>1.8%</td><td>1.8%</td><td>1.8%</td></tr><tr><td>NOR Escapement</td><td>Max 3,842</td><td>Min 479</td><td>Ave 1,116</td><td>Max 3,911</td><td>Min 489</td><td>Ave 1,136</td><td>Max 3,754</td><td>Min 472</td><td>Ave 1,094</td><td>Max 3,893</td><td>Min 485</td><td>Ave 1,130</td><td>Max 3,893</td><td>Min 485</td><td>Ave 1,130</td></tr><tr><td>HOS Total Escapement (includes strays)</td><td>57</td><td>10</td><td>19</td><td>91</td><td>16</td><td>30</td><td>25</td><td>4</td><td>8</td><td>25</td><td>4</td><td>8</td><td>25</td><td>4</td><td>8</td></tr></table>															Relative Hatchery Optimum ->	80	80	80	80	80	Weir Factor ->						Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%	Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81	Fitness Factor after 100 generations (B) ->	0.99	1.00	0.97	1.00	1.00	Average Fitness Factor (100 Generations) ->	0.98	0.99	0.96	0.99	0.99	Generations until average fitness is reached ->	33	30	30	32	32	Minimum Hatchery Program (% of BS Goal) ->	0.5					Effective pHOS ->	0.00	0.00	0.01	0.00	0.00	"Fitness Floor" ->	0.5					Calculated Hatchery SAR ->	1.2%	1.2%	1.2%	1.2%	1.2%	Calculated Natural SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%	NOR Escapement	Max 3,842	Min 479	Ave 1,116	Max 3,911	Min 489	Ave 1,136	Max 3,754	Min 472	Ave 1,094	Max 3,893	Min 485	Ave 1,130	Max 3,893	Min 485	Ave 1,130	HOS Total Escapement (includes strays)	57	10	19	91	16	30	25	4	8	25	4	8	25	4	8
Relative Hatchery Optimum ->	80	80	80	80	80																																																																																																																	
Weir Factor ->																																																																																																																						
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%																																																																																																																	
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81																																																																																																																	
Fitness Factor after 100 generations (B) ->	0.99	1.00	0.97	1.00	1.00																																																																																																																	
Average Fitness Factor (100 Generations) ->	0.98	0.99	0.96	0.99	0.99																																																																																																																	
Generations until average fitness is reached ->	33	30	30	32	32																																																																																																																	
Minimum Hatchery Program (% of BS Goal) ->	0.5																																																																																																																					
Effective pHOS ->	0.00	0.00	0.01	0.00	0.00																																																																																																																	
"Fitness Floor" ->	0.5																																																																																																																					
Calculated Hatchery SAR ->	1.2%	1.2%	1.2%	1.2%	1.2%																																																																																																																	
Calculated Natural SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%																																																																																																																	
NOR Escapement	Max 3,842	Min 479	Ave 1,116	Max 3,911	Min 489	Ave 1,136	Max 3,754	Min 472	Ave 1,094	Max 3,893	Min 485	Ave 1,130	Max 3,893	Min 485	Ave 1,130																																																																																																							
HOS Total Escapement (includes strays)	57	10	19	91	16	30	25	4	8	25	4	8	25	4	8																																																																																																							

**Tables A18-A39: *AHA* outputs for Salmon River Summer Steelhead**

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011*

### *Parameters and assumptions for Salmon River steelhead*

Steelhead smolts from several hatchery populations are currently released into the Salmon River drainage. These hatchery populations include Sawtooth Hatchery steelhead (*A-run*), Pahsimeroi Hatchery steelhead (*A-run*), Oxbow Hatchery steelhead (*A-run*), Dworshak NFH steelhead (*B-run*), Squaw Creek *B-run* hatchery steelhead representing the offspring/descendants of Dworshak NFH steelhead returning to Squaw Creek, and East Fork “Naturals” steelhead (*A/B run*) which are the offspring/descendants of natural-origin steelhead trapped in the East Fork Salmon River. Each of these steelhead stocks has characteristic mean female fecundities, mean egg-to-smolt survivals depending on hatchery, mean smolt-to-adult survival rates, and mean numbers adult recruits per adult spawner (*R/S*). Values of these various parameters for generating *AHA* Tables A18-A39 are presented in Table 5A.

**Table 5A. Mean values of reproductive and survival parameters assumed in the *AHA* analyses for hatchery-origin steelhead released within the Salmon River drainage (data obtained from the HSRG). Some of the values below represent updated estimates obtained by the HRT.**

<b>Hatchery/ Release location <i>Population/Stock</i></b>	<b>Fecundity (No. of eggs)</b>	<b>Within- hatchery egg-to-smolt survival</b>	<b>Salmon River smolt-to- adult survival (%)</b>	<b>Recruit/spawner (<i>R/S</i>; adults)</b>
Hagerman NFH, Magic Valley FH, Niagara Springs FH/ Salmon River basin <i>Oxford, Pahsimeroi, and Sawtooth Hatchery A-run Steelhead</i>	4,912	0.75	0.70%	12.9
Hagerman NFH, Magic Valley FH/ Salmon River basin <i>Dworshak NFH B-run Steelhead</i>	6,848	0.56	0.14%	2.5
Clearwater FH/ Little Salmon River <i>Dworshak NFH B-run Steelhead</i>	6,848	0.75	0.14%	3.4
Magic Valley FH, Niagara Springs FH/ Salmon River basin <i>Localized B-run Steelhead</i>	6,000	0.67	0.40%	7.8
Magic Valley FH/ E.F. Salmon River, Current <i>East Fork “Naturals” steelhead</i>	6,000	0.70	0.19%	4.0
Magic Valley FH/ Salmon River basin, under HSRG and HRT recommendations <i>East Fork “Naturals” steelhead</i>	6,000	0.70	0.34%	7.1

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

### *Current programs, HSRG solutions, and HRT alternatives*

Tables A18-A39 simulate published *HSRG solutions* and two sets of HRT alternatives referred to as *HRT Alternatives A* and *HRT Alternatives B*. *HRT Alternatives A* and *B* are identical except *HRT Alternative A* models a localized hatchery population of *B-run* steelhead on the Pahsimeroi River, whereas *HRT Alternative B* terminates all *B-run* hatchery programs and replaces those released numbers of fish with *A-run* steelhead (Table 6A).

The HSRG solutions and the two HRT sets of alternatives modeled here terminate the annual transfer and outplanting of *Dworshak NFH B-run steelhead* smolts in the East Fork Salmon River and at Squaw Creek. Instead, the *HSRG Solution* and *HRT Alternative “A”* scenarios are based on the development of localized hatchery populations of *B-run* steelhead for the upper Salmon River. The *HSRG Solution* models this “localized” hatchery population for the East Fork Salmon River, assuming construction of a new weir in the East Fork immediately upstream from its confluence with the mainstem Salmon River. *HRT Alternative A* models a localized hatchery broodstock on the Pahsimeroi River because trapping facilities are already in place there. In this context, the HSRG solution can be viewed as a potential “long-term recommendation”, while *HRT Alternative A* can be viewed as a “short-term recommendation”.

The *HSRG Solution* and HRT alternatives for the Little Salmon River hatchery programs maintain the current release numbers of both *A-run* and *Dworshak NFH B-run steelhead*. However, the HRT is recommending that Clearwater FH, rather than Hagerman NFH or Magic Valley FH, rear *Dworshak B-run steelhead* released in the Little Salmon River because of the greater in-hatchery survivals and recruit-per-spawners associated with rearing at Clearwater FH.

The HRT further recommends that all *Sawtooth A-run steelhead* be reared at Hagerman NFH and that all *Pahsimeroi A-run steelhead* be reared at Magic Valley and Niagara Springs FH. Currently, approximately 1.45M and 1.59M steelhead smolts are reared annually at Hagerman NFH and Magic Valley FH, respectively. The “HSRG Solutions” in Tables A19-A39 assume that those numbers would remain unchanged. The two sets of HRT alternatives would reduce those numbers slightly to 1.40 and 1.55M smolts, respectively. Further reductions may be necessary in response to declining water supplies at both facilities.



**Table 6A. Summary of the number of hatchery-origin (HOR) steelhead smolts released in the Salmon River under four strategies modeled in AHA Tables A18-A39: (1) *Current* programs based on the LSRCP Cooperators 2008 Annual Operating Plan for the Salmon River, (2) the *HSRG Solution*, (3) *HRT Alternative A* (development of localized hatchery populations of *B-run* steelhead) and *HRT Alternative B* (A-run steelhead programs only). The names of hatchery stocks/populations are in italics (e.g., “*Oxbow*”, “*Pahsimeroi*”) and are identified by the facility/location where adults are trapped for broodstock. Hatcheries where juvenile fish are reared to the smolt stage prior to release are shown in parentheses. The *HSRG solution* assumes construction of new weirs in the lower Lemhi River and the lower East Fork Salmon River.**

Release location: Program	Current	HSRG Solution	HRT Alternative A	HRT Alternative B
Little Salmon River: <i>A-run</i> (AHA Tables A18, A19)	<ul style="list-style-type: none"> <li>• 275K <i>Oxbow</i> smolts (Niagara Springs FH)</li> <li>• 170K <i>Pahsimeroi</i> smolts (Niagara Springs FH)</li> <li>• 200K <i>Pahsimeroi</i> smolts (Hagerman NFH)</li> </ul>	No recommendation (same as Current)	<ul style="list-style-type: none"> <li>• 275K <i>Oxbow</i> smolts (Niagara Springs FH)</li> <li>• 370K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	Same as HRT Alt. A
Little Salmon River: <i>B-run</i> (AHA Table A20)	<ul style="list-style-type: none"> <li>• 215K <i>Dworshak</i> smolts (Magic Valley FH)</li> <li>• 100K <i>Dworshak</i> smolts (Hagerman NFH)</li> </ul>	No recommendation (same as Current)	<ul style="list-style-type: none"> <li>• 315 K <i>Dworshak</i> smolts (Clearwater FH)</li> </ul>	Same as HRT Alt. A
Salmon River at Red Rock: <i>A-run</i> (AHA Tables A28, A29)	<ul style="list-style-type: none"> <li>• 120K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	New conservation weir in lower Lemhi River	Terminate smolt releases at Red Rock site	Same as HRT Alt. A
Salmon River at Shoup Bridge: <i>A-run</i> (AHA Tables A30, A31)	<ul style="list-style-type: none"> <li>• 80K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	No recommendation (same as Current)	<ul style="list-style-type: none"> <li>• 120K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	Same as HRT Alt. A
Salmon River at Colston Corner: <i>A-run</i> (AHA Table A30, A31)	<ul style="list-style-type: none"> <li>• 140K <i>Pahsimeroi</i> or <i>Sawtooth</i> smolts (Magic Valley FH)</li> </ul>	No recommendation (same as Current)	<ul style="list-style-type: none"> <li>• 180K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	Same as HRT Alt. A
Pahsimeroi River at hatchery weir: <i>A-run</i> (AHA Table A30, A31)	<ul style="list-style-type: none"> <li>• 830K <i>Pahsimeroi</i> smolts (Niagara Springs FH)</li> <li>• 30K <i>Pahsimeroi</i> smolts (Magic Valley FH)</li> </ul>	No recommendation (same as Current)	<ul style="list-style-type: none"> <li>• 900K <i>Pahsimeroi</i> smolts (Niagara Springs FH)</li> </ul>	<ul style="list-style-type: none"> <li>• 1.15M <i>Pahsimeroi</i> smolts (Niagara Springs FH)</li> </ul>
Pahsimeroi River at hatchery weir: <i>B-run</i> (AHA Table A32)	No program	Same as Current	<ul style="list-style-type: none"> <li>• 250K <i>localized Dworshak</i> smolts, <i>Pahsimeroi</i> broodstock. (Magic Valley FH)</li> </ul>	Same as Current
Salmon River at	<ul style="list-style-type: none"> <li>• 120K <i>Pahsimeroi</i> or <i>Sawtooth</i></li> </ul>			

**USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011**

McNabb Point (AHA Table A33)	A-run smolts (Magic Valley FH)	• 120K <i>East Fork “F2” HOR Naturals</i> (Magic Valley FH)	Same as HSRG Solution	Same as HSRG Solution
Salmon River at Tunnel Rock (AHA Table A33)	• 60K <i>Pahsimeroi</i> or <i>Sawtooth</i> A-run smolts (Magic Valley FH)	• 60K <i>East Fork “F2” HOR Naturals</i> (Magic Valley FH)	Same as HSRG Solution	Same as HSRG Solution
E.F. Salmon River at existing weir (RM 18): <i>E.F. “Naturals”</i> (AHA Table A34)	• 60K <i>E.F. Naturals</i> (pNOB = 0.5) (Magic Valley FH)	Terminate releases at existing weir	• 150K <i>E.F. Naturals</i> (pNOB = 0.75), released upstream of weir (Magic Valley FH)	Same as HSRG solution
E.F. Salmon River at mouth (RM 1.0): <i>E.F. “Naturals”</i> (AHA Tables A33, A34)	No program	• Construct new weir at mouth • 100K <i>E.F. Naturals</i> (pNOB = 0.75) (Magic Valley FH)	• 325K <i>East Fork “F2” HOR Naturals</i> (Magic Valley FH)	• Construct new weir at mouth • 150K <i>E.F. Naturals</i> (pNOB = 0.75) (Magic Valley FH) • 325K <i>East Fork “F2” HOR Naturals</i> (Magic Valley FH)
E.F. Salmon River at mouth (RM 1): <i>B-run</i> (AHA Table A35)	• 225K <i>Dworshak</i> smolts (Magic Valley FH) • 100K <i>Dworshak</i> smolts (Hagerman NFH)	• Construct new weir at mouth • 325K <i>localized Dworshak</i> smolts, E.F. broodstock (Magic Valley FH)	No program	No program
Squaw Creek and Pond: <i>B-run</i> (AHA Tables A36, A37)	• 60K <i>localized Dworshak</i> smolts, Squaw Cr. broodstock (Magic Valley FH) • 250K <i>Dworshak</i> smolts (Magic Valley FH)	• 270K <i>E.F. Salmon River localized Dworshak</i> smolts (Magic Valley FH)	• 60K <i>localized Dworshak</i> smolts, Squaw Cr. broodstock (Magic Valley FH)	No program
Slate Creek: <i>A-run</i> (AHA Tables A38, A39)	• 90K <i>Pahsimeroi</i> smolts (Magic Valley FH)	No recommendation (same as Current)	• 90K <i>Sawtooth</i> smolts (Hagerman NFH)	Same as HRT Alt. A
Yankee Fork, Salmon River: <i>A-run</i> (AHA Tables A38, A39)	• 240K <i>Sawtooth</i> smolts (Hagerman NFH) • 90K <i>Sawtooth</i> smolts (Magic Valley FH)	No recommendation (same as Current)	• 330K <i>Sawtooth</i> smolts (Hagerman NFH)	Same as HRT Alt. A
Valley Creek: <i>A-run</i> (AHA Tables A38, A39)	• 50K <i>Sawtooth</i> smolts (Magic Valley FH)	No recommendation (same as Current)	• 50K <i>Sawtooth</i> smolts (Hagerman NFH)	Same as HRT Alt. A
Salmon River at Sawtooth Hatchery weir: <i>A-run</i> (AHA Tables A38, A39)	• 810K <i>Sawtooth</i> smolts (Hagerman NFH)	No recommendation (same as Current)	• 930K <i>Sawtooth</i> smolts (Hagerman NFH)	Same as HRT Alt. A

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A18. AHA output for Little Salmon River summer steelhead (A-run ). Hatchery influence on this population is from outplanted Dworshak NFH steelhead (B-run), and outplanted Oxbow and Pahsimeroi Hatchery A-run steelhead (see Tables A19 and A20).**

All H Analyzer (AHA)																
Apply & Recalculate																
Population (Ctrl + Page Up/Down to Scroll)																
Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest		
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery		
12-Salmon_Little Salmon Summer Steelhead (A-Run)		Population Recovery Designation:		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing		
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B				
Habitat	Productivity (Adult)	Ad. Capacity	3.60	474	3.60	474	3.60	474	3.60	474	3.60	474	3.60	474		
	Min NOR Escape	% Kelt	1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	90	11,850	180	23,704	180	23,704	180	23,704	180	23,704	180	23,704		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.098	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity		1.80	237	3.60	474	3.60	474	3.60	474	3.60	474	3.60	474	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs	HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	
	Harv -Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
	Total Exploitation Rate	NORs	HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI	pHOS-Realized	Cons		None		Harv		Seg		Harv		Seg	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Harv		Seg		Harv		Seg		Harv		Seg	
	Broodstock by Source		Local	Imported	Harv		Seg		Harv		Seg		Harv		Seg	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays	Harv		Seg		Harv		Seg		Harv		Seg	
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	Harv		Seg		Harv		Seg		Harv		Seg	
Patterns	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	Harv		Seg		Harv		Seg		Harv		Seg	
	Update		Habitat	Hydro	Harvest	Hatchery		Hatchery		Hatchery		Hatchery		Hatchery		Hatchery
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.																
549																
1																
Realized Spawning Composition																
Relative Hatchery Optimum -> 80																
Weir Factor ->																
Relative Reproductive Success (HOS) -> 80%																
Initial Fitness Factor (A) -> 0.81																
Fitness Factor after 100 generations (B) -> 0.50																
Average Fitness Factor (100 Generations) -> 0.50																
Generations until average fitness is reached -> 14																
Minimum Hatchery Program (% of BS Goal) -> 0.5																
Effective pHOS -> 0.62																
Effective pHOS -> 0.62																
Effective pHOS -> 0.62																
Effective pHOS -> 0.62																
Effective pHOS -> 0.62																
Calculated Hatchery SAR -> 2.0%																
Calculated Natural SAR -> 2.0%																
NOR Escapement																
HoS Total Escapement (includes strays)																
Max Min Ave																
857 129 258																
5,154 895 1,688																
1,103 158 323																
857 129 258																
5,154 895 1,688																
5,162 897 1,691																
857 129 258																
5,162 897 1,691																

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A19. AHA output for *Oxbow* and *Pahsimeroi* Hatchery A-run steelhead released into the Little Salmon River.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Population Management Intent:

Harvest

Conservation

Harvest

Harvest

Harvest

Salmon

Summer Steelhead

Harv&Hatchery Strategy:

Segregated Hatchery - Outplanting

Natural Reproduction only

Segregated Hatchery - Outplanting

Segregated Hatchery - Outplanting

Segregated Hatchery - Outplanting

TZA-Salmon-Little Salmon Summer Steelhead (A-Run-Pahsimeroi-Oxbow-Hatchery)

Population Recovery Designation:

Not listed - excluded from DPS

N/A

Not listed - excluded from DPS

Not listed - excluded from DPS

Not listed - excluded from DPS

Historic

Current

No Hatchery

HSRG Solution

HRT Alternatives A

HRT Alternatives B

Habitat

Productivity (Adult)

Ad. Capacity

0.00

0

0.00

0

0.00

0

0.00

0

0.00

0

Min NOR Escape

% Kelt

1

1

1

1

1

Smolt Productivity

Sm. Capacity

0

0

0

1

0

1

0

1

0

1

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

0.092

0.020

y

0.053

0.020

y

0.053

0.020

y

0.053

0.020

y

0.053

0.020

y

Juv Passage Surv.

Adult Passage

1.00

1.00

0.49

0.77

0.49

0.77

0.49

0.77

0.49

0.77

0.49

0.77

Adjusted Productivity

Adj. Capacity

0.00

0

0.00

0

0.00

0

0.00

0

0.00

0

0.00

0

Harvest

Harv - Marine

NORs

HORs

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

Harv - L. Mainstem

NORs

HORs

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

0.010

Harv - U. Mainstem

NORs

HORs

0.060

0.060

0.060

0.060

0.060

0.060

0.060

0.060

0.060

0.060

0.060

0.060

Harv - Terminal

NORs

HORs

0.030

0.600

0.030

0.600

0.030

0.600

0.030

0.600

0.030

0.600

0.030

0.600

Total Exploitation Rate

NORs

HORs

0.106

0.631

0.106

0.631

0.106

0.631

0.106

0.631

0.106

0.631

0.106

0.631

Hatchery

Broodstock Composition

pNOB-Goal

pHOS-Goal

pNOB

pHOS

pNOB

pHOS

pNOB

pHOS

pNOB

pHOS

pNOB

pHOS

Purpose

Type

Cons/Harv/Both

Int/Seg/Step/None

Harv

100%

Cons

None

Harv

100%

Harv

100%

Harv

100%

Harv

100%

Broodstock by Source

Local

Imported

350

644,700

350

644,700

350

644,700

350

644,700

Brood Exported (from HOR Surplus)

Export Goal/Realized

Smolt Release

100%

100%

100%

100%

100%

Destination for HOR Returns

% to Hatchery

% to Nat. Spawn.

Productivity of Hatchery Fish

Recruits/Spawner

Fitness? [Y / N]

12.9

y

y

12.9

y

12.9

y

12.9

y

12.9

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro


Current Harvest


Current Habitat


Current Hydro


Current Harvest


Realized Spawning Composition











Relative Hatchery Optimum ->	80	80	80	80	80							
Weir Factor ->												
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%							
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->	0.19	1.00	0.19	0.19	0.19							
Average Fitness Factor (100 Generations) ->	0.24	0.99	0.24	0.24	0.24							
Generations until average fitness is reached ->	43	30	43	43	43							
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS ->	1.00	Effective pHOS ->	1.00							
"Fitness Floor" ->		pNOB ->		pNOB ->								
Calculated Hatchery SAR ->	0.7%			0.7%								
Calculated Natural SAR ->	2.0%		2.0%	2.0%								
NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)	5,669	984	1,856	5,669	984	1,856	5,669	984	1,856	5,669	984	1,856

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A20. AHA output for Dworshak NFH steelhead (B-run ) released into the Little Salmon River.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery - Outplanting		Natural Reproduction only		Segregated Hatchery - Outplanting		Segregated Hatchery - Outplanting		Segregated Hatchery - Outplanting	
12B-Salmon_Little Salmon Summer				Population Recovery Designation:		Not listed - excluded from DPS		N/A		Not listed - excluded from DPS		Not listed - excluded from DPS		Not listed - excluded from DPS	
Steelhead (B-Run-Dworshak-Hatchery)				Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B	
Habitat	Productivity (Adult)	Ad. Capacity	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	0	0	0	1	0	1	0	1	0	1	0	1	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y
	Juv Passage Surv.	Adult Passage		1.00	1.00		0.49	0.77		0.49	0.77		0.49	0.77	
	Adjusted Productivity	Adj. Capacity		0.00	0		0.00	0		0.00	0		0.00	0	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010	
	Harv - U. Mainstem	NORs	HORs	0.140	0.140		0.140	0.140		0.140	0.140		0.140	0.140	
	Harv -Terminal	NORs	HORs	0.030	0.600		0.030	0.600		0.030	0.600		0.030	0.600	
	Total Exploitation Rate	NORs	HORs	0.182	0.663		0.182	0.663		0.182	0.663		0.182	0.663	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose	Type	PNI	pHOS-Realized	Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg	
	Broodstock by Source		Local	Imported	164	314,460			164	314,460	123	315,864	123	315,864	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release											
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	2.5	y		y	2.5	y	3.4	y	3.4	y	
Patterns Update	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat	Hydro	Harvest	Hatchery	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest

7/91

3

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.19	1.00	0.19	0.19	0.19										
Average Fitness Factor (100 Generations) ->	0.24	0.99	0.24	0.24	0.24										
Generations until average fitness is reached ->	43	30	43	43	43										
Minimum Hatchery Program (% of BS Goal) ->															
"Fitness Floor" ->															
Effective pHOS ->	1.00	1.00	1.00	1.00	1.00										
Effective pNOB ->															
Calculated Hatchery SAR ->	0.1%		0.1%	0.1%	0.1%										
Calculated Natural SAR ->	1.8%		1.8%	1.8%	1.8%										
NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)	471	81	154	471	81	154	480	83	157	480	83	157	480	83	157

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A21. AHA output for South Fork Salmon River steelhead (B-run ).

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation				
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only				
13-Salmon_South Fork Summer Steelhead (B-Run)				Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary				
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B				
Habitat	Productivity (Adult)	Ad. Capacity	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115		
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	75	27,875	166	61,568	166	61,568	166	61,568	166	61,568	166	61,568	166	61,568		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.77
	Adjusted Productivity	Adj. Capacity		1.36	505	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	3.00	1,115	1,115
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - U. Mainstem	NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140
	Harv -Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030
	Total Exploitation Rate	NORs	HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS
	Purpose		PNI	pHOS-Realized	Cons	1%	Cons	1%	Cons	1%	Cons	1%	Cons	1%	Cons	1%	Cons	1%
	Type		Cons/Harv/Both	Int/Seg/Step/None	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None
	Broodstock by Source		Local	Imported	Smolt Release													
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		14												
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%		100%		100%
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	y	y	y	y	y	y	y	y	y	y	y	y	y	y
	Update		Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat	Current Habitat
		Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro	Current Hydro
		Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest
		Hatchery																

548

4

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.98	1.00	0.98	0.98	0.98										
Average Fitness Factor (100 Generations) ->	0.97	0.99	0.97	0.97	0.97										
Generations until average fitness is reached ->	31	30	31	31	31										
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.01	Effective pHOS ->	0.01										
"Fitness Floor" ->		pNOB ->		pNOB ->											
Calculated Hatchery SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%										
Calculated Natural SAR ->															
NOR Escapement	Max 2,006	Min 267	Ave 579	Max 2,058	Min 275	Ave 594	Max 2,006	Min 267	Ave 579	Max 2,006	Min 267	Ave 579	Max 2,006	Min 267	Ave 579
HoS Total Escapement (includes strays)	43	7	14	43	7	14	43	7	14	43	7	14	43	7	14



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A23. AHA output for Chamberlain Creek steelhead (A-run ).

All H Analyzer (AHA)

Apply & Recalculate


Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation				
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only				
15-Salmon Chamberlain Summer Steelhead (A-Run)		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary				
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B						
Habitat	Productivity (Adult)	Ad. Capacity	3.00	399	3.00	399	3.00	399	3.00	399	3.00	399	3.00	399	3.00	399		
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	75	9,975	150	19,954	150	19,954	150	19,954	150	19,954	150	19,954	150	19,954		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y
	Juv Passage Surv.	Adult Passage		1.00		1.00	0.49		0.77	0.49		0.77	0.49		0.77	0.49		0.77
	Adjusted Productivity	Adj. Capacity		1.50		199	3.00		399	3.00		399	3.00		399	3.00		399
Harvest	Harv - Marine	NORs	HORs	0.010		0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010		0.010
	Harv - L. Mainstem	NORs	HORs	0.010		0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010		0.010
	Harv - U. Mainstem	NORs	HORs	0.060		0.060	0.060		0.060	0.060		0.060	0.060		0.060	0.060		0.060
	Harv -Terminal	NORs	HORs	0.030		0.030	0.030		0.030	0.030		0.030	0.030		0.030	0.030		0.030
	Total Exploitation Rate	NORs	HORs	0.106		0.106	0.106		0.106	0.106		0.106	0.106		0.106	0.106		0.106
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS		
	Purpose		PNI	pHOS-Realized	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None		
	Broodstock by Source		Local	Imported	Smolt Release													
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		2		2		2		2		2		2		
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%		100%		
Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y		y		y		y			
Patterns	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat	Hydro	Harvest	Hatchery												
	Update																	

546

6

Realized Spawning Composition



Relative Hatchery Optimum ->	80	80	80	80	80							
Weir Factor ->												
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%							
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->	1.00	1.00	1.00	1.00	1.00							
Average Fitness Factor (100 Generations) ->	0.99	0.99	0.99	0.99	0.99							
Generations until average fitness is reached ->	32	30	32	32	32							
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.00	Effective pHOS ->	0.00							
"Fitness Floor" ->		pNOB ->		pNOB ->								
Calculated Hatchery SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%							
Calculated Natural SAR ->												
NOR Escapement	Max 848	Min 116	Ave 245	Max 848	Min 116	Ave 245	Max 848	Min 116	Ave 245	Max 848	Min 116	Ave 245
HoS Total Escapement (includes strays)	6	1	2	6	1	2	6	1	2	6	1	2



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A24. AHA output for Lower Middle Fork Salmon River steelhead (B-run ).

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only	
16-Salmon_Lower Middle Fork Salmon				Population Recovery Designation:		Primary		Primary		Primary		Primary	
Summer Steelhead (B-Run)				Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A	
Habitat		Productivity (Adult)		Ad. Capacity		2.50		1,587		2.50		1,587	
		Min NOR Escape		% Kelt		1				1			
		Smolt Productivity		Sm. Capacity		63		39,675		138		87,629	
Hydro		Ocean Surv		Baseline SAR		Vary? (Y/N)		0.048		0.018		Y	
		Juv Passage Surv.		Adult Passage				0.49		0.77			
		Adjusted Productivity		Adj. Capacity		1.13		620		2.50		1,587	
Harvest		Harv - Marine		NORs		HORs		0.010		0.010			
		Harv - L. Mainstem		NORs		HORs		0.010		0.010			
		Harv - U. Mainstem		NORs		HORs		0.140		0.140			
		Harv -Terminal		NORs		HORs		0.030		0.030			
		Total Exploitation Rate		NORs		HORs		0.182		0.182			
Hatchery		Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB	
		Purpose		PNI		pHOS-Realized		Cons		None		Cons	
		Type		Cons/Harv/Both		Int/Seg/Step/None		Cons		None		Cons	
		Broodstock by Source		Local		Imported							
		Brood Exported (from HOR Surplus)		Export Goal/Realized		Smolt Release		10		10		10	
		Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		100%		100%		100%	
		Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		y		y		y	
Patterns		Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Hydro		Current Habitat		Current Hydro		Current Habitat	
Update				Harvest		Hatchery		Current Harvest		Current Harvest		Current Harvest	

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80							
Weir Factor ->												
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%							
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->	0.99	1.00	0.99	0.99	0.99							
Average Fitness Factor (100 Generations) ->	0.98	0.99	0.98	0.98	0.98							
Generations until average fitness is reached ->	33	30	33	33	33							
Minimum Hatchery Program (% of BS Goal) ->	0.5											
"Fitness Floor" ->												
Calculated Hatchery SAR ->												
Calculated Natural SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%							
NOR Escapement	Max 2,450	Min 305	Ave 711	Max 2,483	Min 311	Ave 721	Max 2,450	Min 305	Ave 711	Max 2,449	Min 305	Ave 711
HoS Total Escapement (includes strays)	31	5	10				31	5	10	31	5	10

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A25. AHA output for Upper Middle Fork Salmon River steelhead (B-run).

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation		
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		
17-Salmon_Upper Middle Fork Salmon				Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		
Summer Steelhead (B-Run)						Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		
Habitat	Productivity (Adult)	Ad. Capacity	2.50	1,667	2.50	1,667	2.50	1,667	2.50	1,667	2.50	1,667	2.50	1,667		
	Min NOR Escape	% Kelt	1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	63	41,675	138	92,047	138	92,047	138	92,047	138	92,047	138	92,047		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018		
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77			
	Adjusted Productivity	Adj. Capacity		1.13	755	2.50	1,667	2.50	1,667	2.50	1,667	2.50	1,667			
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010			
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010				
	Harv - U. Mainstem	NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140					
	Harv -Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030						
	Total Exploitation Rate	NORs	HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182						
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI	pHOS-Realized	Cons		None		Cons		None		Cons		None	
	Broodstock by Source		Local	Imported	Smolt Release											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays												
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.			100%				100%					
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]			y				y					
Patterns	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat			
			Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro			
			Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest			
	Update		Hatchery													

Realized Spawning Composition

Relative Hatchery Optimum ->		80	80	80	80	80
Weir Factor ->						
Relative Reproductive Success (HOS) ->		80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->		1.00	1.00	1.00	1.00	1.00
Average Fitness Factor (100 Generations) ->		0.99	0.99	0.99	0.99	0.99
Generations until average fitness is reached ->		30	30	30	30	30
Minimum Hatchery Program (% of BS Goal) ->			Effective pHOS ->	Effective pHOS ->	Effective pHOS ->	Effective pHOS ->
"Fitness Floor" ->		0.5	pNOB ->	pNOB ->	pNOB ->	pNOB ->
Calculated Hatchery SAR ->			1.8%	1.8%	1.8%	1.8%
Calculated Natural SAR ->						
NOR Escapement		Max	Min	Ave	Max	Min
HoS Total Escapement (includes strays)		2,608	326	758	2,608	326

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A26. AHA output for Panther Creek steelhead (A-run ).

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only	
18-Salmon_Panther Creek Summer Steelhead (A-Run)		Population Recovery Designation:		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B			
Habitat	Productivity (Adult)	Ad. Capacity	2.00	428	2.00	428	2.00	428	2.00	428	2.00	428	2.00	428	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	50	10,700	100	21,403	100	21,403	100	21,403	100	21,403	100	21,403	
Hydro	Ocean Surv	Baseline SAR	0.092	0.020	0.053	0.020	0.053	0.020	0.053	0.020	0.053	0.020	0.053	0.020	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity	1.00	214	2.00	428	2.00	428	2.00	428	2.00	428	2.00	428	
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs		HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	
	Harv -Terminal	NORs		HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
	Total Exploitation Rate	NORs		HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	
	Broodstock by Source		Local	Imported	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release	2		2		2		2		2		
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	100%		100%		100%		100%		100%		
Patterns Update	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	y		y		y		y		y		
	Habitat		Current Habitat												
	Hydro		Current Hydro												
Harvest		Current Harvest													
Hatchery															

543

9

Realized Spawning Composition

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.99	1.00	0.99	0.99	0.99										
Average Fitness Factor (100 Generations) ->	0.99	0.99	0.99	0.99	0.99										
Generations until average fitness is reached ->	33	30	33	33	33										
Minimum Hatchery Program (% of BS Goal) ->	0.5														
"Fitness Floor" ->	0.5														
Calculated Hatchery SAR ->															
Calculated Natural SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%										
NOR Escapement	Max 599	Min 70	Ave 179	Max 606	Min 71	Ave 181	Max 599	Min 70	Ave 179	Max 599	Min 70	Ave 179	Max 599	Min 70	Ave 179
HoS Total Escapement (includes strays)	6	1	2	6	1	2	6	1	2	6	1	2	6	1	2

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A27. AHA output for North Fork Salmon River steelhead (A-run ).


All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction Only		Natural Reproduction only		Natural Reproduction only	
19-Salmon_North Fork Salmon Summer Steelhead (A-Run)				Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary	
						Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A	
Habitat	Productivity (Adult)	Ad. Capacity	3.00	226	3.00	226	3.00	226	3.00	226	3.00	226	3.00	226	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	75	5,650	150	11,302	150	11,302	150	11,302	150	11,302	150	11,302	
Hydro	Ocean Surv.	Baseline SAR	0.092	0.020	0.053	0.020	0.053	0.020	0.053	0.020	0.053	0.020	0.053	0.020	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity	1.50	113	3.00	226	3.00	226	3.00	226	3.00	226	3.00	226	
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - U. Mainstem	NORs		HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	
	Harv -Terminal	NORs		HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
	Total Exploitation Rate	NORs		HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized	Cons	None	Cons	None	Cons	None	Cons	None	Cons	None	
	Broodstock by Source		Local	Imported	Smolt Release										
	Brood Exported (from HOR Surplus)		Export Goal/Realized	% to Hatchery		10		100		10		11		12	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y		y		y	
Patterns Update	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat	Current Habitat			Current Habitat			Current Habitat			Current Habitat		
			Hydro	Current Hydro			Current Hydro			Current Hydro			Current Hydro		
			Harvest	Current Harvest			Current Harvest			Current Harvest			Current Harvest		

Realized Spawning Composition



Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.84	1.00	0.84	0.82	0.76
Average Fitness Factor (100 Generations) ->	0.87	0.99	0.87	0.86	0.82
Generations until average fitness is reached ->	8	30	8	6	3
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.02	Effective pHOS ->	0.02
"Fitness Floor" ->		pNOB ->		pNOB ->	0.03
Calculated Hatchery SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%
Calculated Natural SAR ->					
NOR Escapement	Max 416	Min 55	Ave 122	Max 482	Min 66
HoS Total Escapement (includes strays)	30	5	10	30	5

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A28. AHA output for Lemhi River steelhead (A-run ). This population is influenced by hatchery fish from the release of *Pahsimeroi* Hatchery A-run steelhead smolts in the mainstem Salmon River, primarily at the Red Rock site (currently 120,000 smolts; see Table A29).**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest + Conservation		Conservation		Harvest + Conservation		Conservation		Conservation				
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		HOR outplants at Red Rock, MS Salmon		Natural Reproduction only		HORs + weir on Lemhi River		Natural Reproduction only		Natural Reproduction only				
20-Salmon_Lemhi Summer Steelhead (A-Run)		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary				
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B						
Habitat	Productivity (Adult)	Ad. Capacity	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139		
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	45	28,475	90	56,959	90	56,959	90	56,959	90	56,959	90	56,959	90	56,959		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	
	Adjusted Productivity	Adj. Capacity		0.90	569	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	1.80	1,139	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010		
	Harv - U. Mainstem	NORs	HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060		
	Harv - Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030		
	Total Exploitation Rate	NORs	HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106		
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS		
	Purpose		PNI	pHOS-Realized	5%	34%	5%	6%	5%	11%	5%	13%	5%	13%	5%	13%		
	Cons/Harv/Both		Int/Seg/Step/None	Harv	Seg	Cons	None	Harv	Seg	Cons	None	Cons	None	Cons	None	Cons		
	Broodstock by Source		Local Imported	Smolt Release														
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		390				39		49				56		
Patterns	Destination for HOR Returns		% to Nat. Spawn.		100%		100%		100%		100%		100%		100%			
	Productivity of Hatchery Fish		Recruits/Spawner		y		y		y		y		y		y			
	Update		Fitness? [Y / N]															
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat				
		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro				
		Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest				
		Hatchery		Current Hatchery		Current Hatchery		Current Hatchery		Current Hatchery		Current Hatchery		Current Hatchery				

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->			90%		
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.50
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.64	0.57	0.55
Generations until average fitness is reached ->	19	30	59	42	
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.34	Effective pHOS ->	0.06
"Fitness Floor" ->		pNOB ->		pNOB ->	
Calculated Hatchery SAR ->					
Calculated Natural SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%
NOR Escapement	646	63	187	1,303	147
Hos Total Escapement (includes strays)	1,191	207	390		
				644	15
				118	20
				39	
				149	26
				49	
				170	29
				56	

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A29. AHA output for *Pahsimeroi Hatchery A-run steelhead* released/outplanted in the mainstem Salmon River at Red Rock site, downstream from the mouth of the Lemhi River.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Salmon

Summer Steelhead

20A-Salmon\_Lemhi Summer Steelhead (A-Run-Pahsimeroi Hatchery)

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Harvest

Conservation

Harvest

Conservation

Conservation

HOR outplants at Red Rock

Natural Reproduction only

HORs + weir on Lemhi River

Natural Reproduction only

Natural Reproduction only

Not listed; excluded from DPS

N/A

Not listed; excluded from DPS

N/A

N/A

Historic

Current

No Hatchery

HSRG Solution

HRT Alternatives A

HRT Alternatives B

Habitat

Productivity (Adult)

Min NOR Escape

Smolt Productivity

Ad. Capacity

% Kelt

Sm. Capacity

0.00

1

0

0

0.00

1

0

0.00

1

0

0.00

1

0

Hydro

Ocean Surv

Baseline SAR

Juv Passage Surv.

Adjusted Productivity

Vary? (Y/N)

Adult Passage

Adj. Capacity

0.092

0.020

1.00

0.00

0.053

0.020

0.49

0.00

0.053

0.020

0.49

0.00

0.053

0.020

0.49

0.00

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

HORs

0.010

0.010

0.060

0.030

0.106

0.010

0.010

0.060

0.030

0.106

0.010

0.010

0.060

0.030

0.106

0.010

0.010

0.060

0.030

0.106

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

Local

Imported

Export Goal/Realized

% to Hatchery

Recruits/Spawner

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

Smolt Release

Strays

% to Nat. Spawn.

Fitness? [Y / N]

pNOB

pHOS

100%

Harv

Seg

Cons

None

65

119,730

100%

12.9

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest


Hatchery


Current Habitat


Current Hydro


Current Harvest

Realized Spawning Composition









Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

80

25%

0.81

0.50

0.50

12

0.5

0.7%

2.0%

0

1,052

80

25%

0.81

1.00

0.99

30

0.7%

2.0%

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A30. AHA output for Pahsimeroi River steelhead (A-run).** This population is influenced primarily by the release of *Pahsimeroi Hatchery A-run steelhead* smolts at the hatchery weir on the Pahsimeroi River.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation and Harvest		Conservation		Conservation and Harvest		Conservation and Harvest		Conservation and Harvest	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery	
21-Salmon_Pahsimeroi Summer Steelhead (A-Run)				Population Recovery Designation:		Contributing		Contributing		Contributing		Contributing		Contributing	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B			
Habitat	Productivity (Adult)	Ad. Capacity	1.65	1,029	1.65	1,029	1.65	1,029	1.65	1,029	1.65	1,029	1.65	1,029	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	41	25,725	83	51,459	83	51,459	83	51,459	83	51,459	83	51,459	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.098	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77
	Adjusted Productivity	Adj. Capacity		0.82	514	1.65	1,029	1.65	1,029	1.65	1,029	1.65	1,029	1.65	1,029
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - U. Mainstem		NORs	HORs	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060
	Harv - Terminal		NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030
	Total Exploitation Rate		NORs	HORs	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106	0.106
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized		10%		10%		10%		10%		10%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Both	None	Cons	None	Both	None	Both	None	Both	None	
	Broodstock by Source		Local	Imported											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		86		82		93		109		100	
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y		y		y	
	Update		Habitat	Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	
		Harvest	Hatchery	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest		

464

13

Realized Spawning Composition

Relative Hatchery Optimum ->		80	80	80	80	80							
Weir Factor ->													
Relative Reproductive Success (HOS) ->		80%	80%	80%	80%	80%							
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->		0.50	1.00	0.50	0.50	0.50							
Average Fitness Factor (100 Generations) ->		0.51	0.99	0.51	0.51	0.50							
Generations until average fitness is reached ->		27	30	27	26	22							
Minimum Hatchery Program (% of BS Goal) ->		0.5	Effective pHOS ->	0.25	Effective pHOS ->	0.25							
"Fitness Floor" ->			pNOB ->		pNOB ->								
Calculated Hatchery SAR ->													
Calculated Natural SAR ->		2.0%	2.0%	2.0%	2.0%	2.0%							
		Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
NOR Escapement		204	17	63	929	101	309	208	17	61	203	18	66
HoS Total Escapement (includes strays)		262	45	86				250	44	82	285	49	93

262	45	86				250	44	82	285	49	93	331	58	109
-----	----	----	--	--	--	-----	----	----	-----	----	----	-----	----	-----

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A31. AHA output for *Pahsimeroi Hatchery A-run steelhead* released into the *Pahsimeroi River* at the hatchery weir (currently 860,000 smolts), and into the mainstem Salmon River at Shoup Bridge (80,000 smolts) and at Colston Corner (140,000 smolts).**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		0		Harvest		Harvest		Harvest		
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery: A-run only		
21A-Salmon_Pahsimeroi Summer Steelhead (A-Run-Pahsimeroi-Hatchery)		Population Recovery Designation:		Not listed; excluded from DPS		N/A		Not listed; excluded from DPS		Not listed; excluded from DPS		Not listed; excluded from DPS		Not listed; excluded from DPS		
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B				
Habitat	Productivity (Adult)		Ad. Capacity		0.00		0		0.00		0		0.00		0	
	Min NOR Escape		% Kelt		1		1		1		1		1		1	
	Smolt Productivity		Sm. Capacity		0		1		0		1		0		1	
Hydro	Ocean Surv. Baseline SAR		Vary? (Y/N)		0.092		0.020		y		0.053		0.020		y	
	Juv Passage Surv.		Adult Passage		1.00		1.00		0.49		0.77		0.49		0.77	
	Adjusted Productivity		Adj. Capacity		0.00		0		0.00		0		0.00		0	
Harvest	Harv - Marine		NORs		HORS		0.010		0.010		0.010		0.010		0.010	
	Harv - L. Mainstem		NORs		HORS		0.010		0.010		0.010		0.010		0.010	
	Harv - U. Mainstem		NORs		HORS		0.060		0.060		0.060		0.060		0.060	
	Harv -Terminal		NORs		HORS		0.030		0.600		0.030		0.600		0.030	
	Total Exploitation Rate		NORs		HORS		0.106		0.631		0.106		0.631		0.106	
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB	
	Purpose		Type		Cons/Harv/Both		Harv		Seg		Cons		None		Harv	
	Broodstock by Source		Local		Imported		Smolt Release		586		1,079,411		586		1,079,411	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays		75%		25%		75%		25%		75%	
	Destination for HOR Returns		Recruits/Spawner		Fitness? [Y / N]		12.9		y		12.9		y		12.9	
Patterns	Update		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat	
			Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro	
			Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest	



Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80														
Weir Factor ->																			
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%														
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81														
Fitness Factor after 100 generations (B) ->	0.19	1.00	0.19	0.19	0.19														
Average Fitness Factor (100 Generations) ->	0.24	0.99	0.24	0.24	0.24														
Generations until average fitness is reached ->	43	30	43	43	43														
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS ->	1.00	Effective pHOS ->	1.00														
"Fitness Floor" ->		pNOB ->		pNOB ->															
Calculated Hatchery SAR ->	0.7%			0.7%															
Calculated Natural SAR ->	2.0%			2.0%															
NOR Escapement	0	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave						
HoS Total Escapement (includes strays)	2,372	412	777	0	0	0	0	0	0	0	2,372	412	777	2,640	458	864	3,186	553	1,043



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A32. AHA output for localized B-run summer steelhead smolts (derived from Dworshak NFH stock) released into the Pahsimeroi River. The modeled program represents an annual release of 250,000 smolts derived from adult fish returning to the weir on the Pahsimeroi River.**

All H Analyzer (AHA)															
Apply & Recalculate															
Population (Ctrl + Page Up/Down to Scroll)															
Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		B-run outplants elsewhere in Salmon Riv.		Natural Reproduction only		B-run program in E.F.		Segregated Hatchery		No B-run program in upper Salmon Riv.	
21B-Salmon_Pahsimeroi Summer Steelhead (B-Run-Pahsimeroi-Hatchery)				Population Recovery Designation:		N/A		N/A		N/A		Not listed; excluded from DPS		N/A	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B			
Habitat	Productivity (Adult)	Ad. Capacity	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1
	Smolt Productivity	Sm. Capacity	0	0	0	1	0	1	0	1	0	1	0	1	0
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.020	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y
	Juv Passage Surv.	Adult Passage		1.00	1.00		0.49	0.77		0.49	0.77		0.49	0.77	
	Adjusted Productivity	Adj. Capacity		0.00	0		0.00	0		0.00	0		0.00	0	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010	
	Harv - U. Mainstem	NORs	HORs	0.140	0.140		0.140	0.140		0.140	0.140		0.140	0.140	
	Harv - Terminal	NORs	HORs	0.030	0.600		0.030	0.600		0.030	0.600		0.030	0.600	
	Total Exploitation Rate	NORs	HORs	0.182	0.663		0.182	0.663		0.182	0.663		0.182	0.663	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized											
	Type		Cons/Harv/Both	Int/Seg/Step/None	Harv	None	Cons	None	Harv	None	Harv	None	Harv	None	
	Broodstock by Source		Local	Imported											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays											
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y		y		y	
	Update		Habitat	Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	Current Habitat	Current Hydro	
		Harvest		Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest	Current Harvest		
		Hatchery													
950															
15															
<div><div>Realized Spawning Composition</div><div></div></div>															
Relative Hatchery Optimum ->				80		80		80		80		80		80	
Weir Factor ->															
Relative Reproductive Success (HOS) ->				25%		25%		25%		25%		25%		25%	
Initial Fitness Factor (A) ->				0.81		0.81		0.81		0.81		0.81		0.81	
Fitness Factor after 100 generations (B) ->				1.00		1.00		1.00		0.19		1.00		1.00	
Average Fitness Factor (100 Generations) ->				0.99		0.99		0.99		0.24		0.99		0.99	
Generations until average fitness is reached ->				30		30		30		43		30		30	
Minimum Hatchery Program (% of BS Goal) ->						Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->	
"Fitness Floor" ->						pNOB ->		pNOB ->		pNOB ->		pNOB ->		pNOB ->	
Calculated Hatchery SAR ->										0.4%					
Calculated Natural SAR ->				1.8%		1.8%		1.8%		1.8%		1.8%		1.8%	
				Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
NOR Escapement					0	0	0	0	0	0	0	0	0	0	0
HoS Total Escapement (includes strays)										100	17	32			



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011


**Table A34. AHA output for East Fork Salmon River steelhead (A/B-run). The HSRG solution and HRT Alternative B assume construction of a new weir at RM 1.0 in the East Fork Salmon River. HRT Alternative A assumes releases are moved upstream of the current weir at RM 18.**


All H Analyzer (AHA)										Apply & Recalculate													
Population (Ctrl + Page Up/Down to Scroll)																							
Subregion/Subbasin			Species/Race			Population Management Intent:			Conservation			Conservation			Conservation + Harvest			Conservation + Harvest			Conservation + Harvest		
Salmon			Summer Steelhead			Harv&Hatchery Strategy:			Integrated Hatchery			Natural reproduction only			Stepping Stone Hatchery Program			Stepping Stone Hatchery Program			Stepping Stone Hatchery Program		
22-Salmon_East Fork Salmon Summer Steelhead						Population Recovery Designation:			Primary			Primary			Primary			Primary			Primary		
						Historic			Current			No Hatchery			HSRG Solution			HRT Alternatives A			HRT Alternatives B		
Habitat	Productivity (Adult)	Ad. Capacity	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048			
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1		1		1				
	Smolt Productivity	Sm. Capacity	38	26,200	75	52,408	75	52,408	75	52,408	75	52,408	75	52,408	75	52,408	75	52,408	75	52,408			
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y		
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77			
	Adjusted Productivity	Adj. Capacity		0.75	524	1.50	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048	1.50	1,048			
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010			
	Harv - L. Mainstem	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010				
	Harv - U. Mainstem	NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140				
	Harv -Terminal	NORs	HORs	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030				
	Total Exploitation Rate	NORs	HORs	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182	0.182				
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	50%	30%	50%	30%	50%	30%	50%	30%	50%	30%	50%	30%	50%	30%	50%	30%			
	Purpose		PNi	pHOS-Realized	0.53	44%	0.53	44%	0.66	39%	0.59	53%	0.60	51%	0.60	51%	0.60	51%	0.60	51%			
			Cons/Harv/Both	Int/Seg/Step/None	Cons	Int	Cons	Int	Cons	Int	Cons	Int	Cons	Int	Cons	Int	Cons	Int	Cons	Int			
	Broodstock by Source		Local	Imported	29	60,900	29	60,900	48	100,800	72	151,200	72	151,200	72	151,200	72	151,200	72	151,200			
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays	20%	80%	20%	80%	86	269	160	466	160	295	160	295	160	295	160	295			
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%	75%	25%	50%	50%	50%	50%	50%	50%	50%	50%	50%	50%			
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	4.0	y	4.0	y	7.1	y	7.1	y	7.1	y	7.1	y	7.1	y	7.1	y			
	Update																						
Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.			Habitat			Current Habitat			Current Habitat			Current Habitat			Current Habitat			Current Habitat			Current Habitat		
			Hydro			Current Hydro			Current Hydro			Current Hydro			Current Hydro			Current Hydro			Current Hydro		
			Harvest			Current Harvest			Current Harvest			Current Harvest			Current Harvest			Current Harvest			Current Harvest		
			Hatchery																				

467

16

Realized Spawning Composition





Relative Hatchery Optimum ->	80	80	80	80	80									
Weir Factor ->														
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%									
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81									
Fitness Factor after 100 generations (B) ->	0.67	1.00	0.79	0.73	0.73									
Average Fitness Factor (100 Generations) ->	0.70	0.99	0.80	0.74	0.75									
Generations until average fitness is reached ->	38	30	9	29	28									
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS -> 0.44	Effective pHOS -> 0.39	Effective pHOS -> 0.53	Effective pHOS -> 0.51									
"Fitness Floor" -> 0.5		pNOB -> 0.25	pNOB -> 0.38	pNOB -> 0.38	pNOB -> 0.38									
Calculated Hatchery SAR ->	0.2%													
Calculated Natural SAR ->	2.0%	2.0%	0.3%	0.3%	0.3%									
			Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
NOR Escapement			969	88	262	407	35	155	789	42	202	1,138	69	272
HoS Total Escapement (includes strays)			1,965	340	643				1,058	184	346	2,135	370	699



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A36. AHA output for localized B-run hatchery steelhead released into Squaw Creek. The HSRG Solution imports localized B-run steelhead annually from a new B-run hatchery population propagated from adults that return to the East Fork Salmon River.**


All H Analyzer (AHA)


Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest	
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Local Segregated Broodstock		Natural Reproduction only		Local broodstock at E.F. Salmon R.		Broodstock at Pahsimeroi FH		No B-run program in upper Salmon River	
23C-Salmon_Upper Salmon Summer Steelhead (Upper Salmon B-Run Program)				Population Recovery Designation:		Not listed; excluded from DPS		N/A		Not listed; excluded from DPS		Not listed; excluded from DPS		N/A	
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B	
Habitat	Productivity (Adult)		Ad. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0
	Min NOR Escape		% Kelt	1		1		1		1		1		1	
	Smolt Productivity		Sm. Capacity	0	0	1	1	1	1	1	1	1	1	1	1
Hydro	Ocean Surv		Baseline SAR	0.092	0.018	y	0.048	0.018	y	0.048	0.018	y	0.048	0.018	y
	Juv Passage Surv.		Adult Passage	1.00	1.00	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77	0.49	0.77
	Adjusted Productivity		Adj. Capacity	0.00	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - U. Mainstem		NORs	HORs	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140	0.140
	Harv -Terminal		NORs	HORs	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.030	0.600	0.600
	Total Exploitation Rate		NORs	HORs	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.182	0.663	0.663
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB
	Purpose		PNI	pHOS-Realized		100%				100%				100%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg	Harv
	Broodstock by Source		Local	Imported	30	59,153			134	269,340	30	59,153			
	Broodstock Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release	25%	75%		100%		100%	25%	75%		75%	75%
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	7.1	y		y	7.1	y	7.1	y	7.1	y	y
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]											
	Update		Habitat	Hydro	Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat
				Hatchery											

Realized Spawning Composition





Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	25%	25%	25%	25%	25%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.19	1.00	0.19	0.19	1.00
Average Fitness Factor (100 Generations) ->	0.24	0.99	0.24	0.24	0.99
Generations until average fitness is reached ->	43	30	43	43	30
Minimum Hatchery Program (% of BS Goal) ->					
"Fitness Floor" ->					
Effective pHOS ->	1.00	1.00	1.00	1.00	1.00
Effective pNOB ->					
Calculated Hatchery SAR ->	0.4%		0.4%	0.4%	0.4%
Calculated Natural SAR ->	1.8%	1.8%	1.8%	1.8%	1.8%
NOR Escapement	Max	Min	Ave	Max	Min
HOS Total Escapement (includes strays)	183	29	59	1,092	189

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A37. AHA output for Dworshak NFH B-run steelhead outplanted into Squaw Creek. The HSRG Solution and HRT Alternative A would terminate these outplants (as would HRT Alternative B) and develop a localized hatchery population at the upper Salmon River basin.**


All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:			Harvest			Conservation			Harvest			Harvest																		
Salmon		Summer Steelhead		Harv&Hatchery Strategy:			Outplanting from Dworshak NFH			Natural Reproduction only			Local broodstock at E.F. Salmon R			Local Broodstock at Pahsimeroi P																		
23B-Salmon_Upper Salmon Summer Steelhead (B-Run Dworshak-Hatchery)					Population Recovery Designation:			N/A			N/A			N/A			B-run steelhead program in upper Salmon R																	
					Historic			Current			No Hatchery			HSRG Solution			HRT Alternatives A			HRT Alternatives B														
Habitat	Productivity (Adult)		Ad. Capacity		0.01			0			0.01			0			0.01			0			0.01			0								
	Min NOR Escape		% Kelt		1						1						1						1											
	Smolt Productivity		Sm. Capacity		0			0			1			1			1			1			1			1								
Hydro	Ocean Surv		Baseline SAR		0.092			0.018			y			0.048			0.018			y			0.048			0.018			y					
	Juv Passage Surv.		Adult Passage		1.00			1.00			0.49			0.77			0.49			0.77			0.49			0.77								
	Adjusted Productivity		Adj. Capacity		0.00			0			0.01			0			0.01			0			0.01			0								
Harvest	Harv - Marine		NORs		HORs			0.010			0.010			0.010			0.010			0.010			0.010			0.010								
	Harv - L. Mainstem		NORs		HORs			0.010			0.010			0.010			0.010			0.010			0.010			0.010								
	Harv - U. Mainstem		NORs		HORs			0.140			0.140			0.140			0.140			0.140			0.140			0.140								
	Harv -Terminal		NORs		HORs			0.030			0.600			0.030			0.600			0.030			0.600			0.030			0.600					
	Total Exploitation Rate		NORs		HORs			0.182			0.663			0.182			0.663			0.182			0.663			0.182			0.663					
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB			pHOS			pNOB			pHOS			pNOB			pHOS			pNOB			pHOS						
	Purpose		Type		PNI		pHOS-Realized		100%																									
	Cons/Harv/Both		Int/Seg/Step/None		Harv			Seg			Cons			None			Harv			None			Harv			None								
	Broodstock by Source		Local		Imported		Smolt Release		130			249,267																						
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays																													
	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		25%			75%			100%			100%			100%			100%			100%									
Patterns	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		2.5			y			y			2.5			y			2.5			y			2.5			y			
	Update		Habitat		Hydro		Harvest		Current Habitat			Current Hydro			Current Harvest			Current Habitat			Current Hydro			Current Harvest			Current Habitat			Current Hydro			Current Harvest	

Realized Spawning Composition



Relative Hatchery Optimum ->		80		80		80		80		80			
Weir Factor ->													
Relative Reproductive Success (HOS) ->		25%		25%		25%		25%		25%			
Initial Fitness Factor (A) ->		0.81		0.81		0.81		0.81		0.81			
Fitness Factor after 100 generations (B) ->		0.19		1.00		1.00		1.00		1.00			
Average Fitness Factor (100 Generations) ->		0.24		0.99		0.99		0.99		0.99			
Generations until average fitness is reached ->		43		30		30		30		30			
Minimum Hatchery Program (% of BS Goal) ->				Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->			
"Fitness Floor" ->				pNOB ->		pNOB ->		pNOB ->		pNOB ->			
Calculated Hatchery SAR ->		0.1%				0.1%				0.1%			
Calculated Natural SAR ->		1.8%		1.8%		1.8%		1.8%		1.8%			
		Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
NOR Escapement		0	0	0	0	0	0	0	0	0	0	0	0
HoS Total Escapement (includes strays)		280	48	91									

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A38. AHA output for upper Salmon River steelhead (A-run ) including Yankee Fork, Valley Creek, Slate Creek, and Squaw Creek.


All H Analyzer (AHA)


Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Harvest and Conservation		Harvest and Conservation		Harvest and Conservation		
Salmon		Summer Steelhead		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery		
23-Salmon_Upper Salmon Summer Steelhead (A-Run)				Population Recovery Designation:		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing		
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternatives A		HRT Alternatives B		
Habitat	Productivity (Adult)	Ad. Capacity	1.50	1,283	1.50	1,283	1.50	1,283	1.50	1,283	1.50	1,283	1.50	1,283		
	Min NOR Escape	% Kelt	1		1		1		1		1		1			
	Smolt Productivity	Sm. Capacity	38	32,075	75	64,168	75	64,168	75	64,168	75	64,168	75	64,168		
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.020	y	0.053	0.020	y	0.053	0.020	y	0.053	0.020	y	
	Juv Passage Surv.	Adult Passage		1.00	1.00		0.49	0.77		0.49	0.77		0.49	0.77		
	Adjusted Productivity	Adj. Capacity		0.75	641		1.50	1,283		1.50	1,283		1.50	1,283		
Harvest	Harv - Marine	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010		
	Harv - L. Mainstem	NORs	HORs	0.010	0.010		0.010	0.010		0.010	0.010		0.010	0.010		
	Harv - U. Mainstem	NORs	HORs	0.060	0.060		0.060	0.060		0.060	0.060		0.060	0.060		
	Harv -Terminal	NORs	HORs	0.030	0.030		0.030	0.030		0.030	0.030		0.030	0.030		
	Total Exploitation Rate	NORs	HORs	0.106	0.106		0.106	0.106		0.106	0.106		0.106	0.106		
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI	pHOS-Realized	Both		45%		Both		44%		Both		45%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Both		Seg		Cons		None		Both		Seg	
	Broodstock by Source		Local	Imported	727		100%		648		100%		699		100%	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release	y		y		y		y		y		y	
Destination for HOR Returns		% to Nat. Spawn.		100%		100%		100%		100%		100%		100%		
Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	y		y		y		y		y		y		
Patterns Update	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat	Hydro	Harvest	Hatchery	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat	Current Hydro	Current Harvest	Current Habitat

Realized Spawning Composition





Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->		25%	25%		
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.50
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.50	0.50	0.50
Generations until average fitness is reached ->	17	30	17	17	18
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.45	Effective pHOS ->	0.44
"Fitness Floor" ->	0.5	pNOB ->		pNOB ->	
Calculated Hatchery SAR ->					
Calculated Natural SAR ->	2.0%	2.0%	2.0%	2.0%	2.0%
NOR Escapement	Max 802	Min 77	Ave 223	Max 821	Min 84
HoS Total Escapement (includes strays)	2,221	386	727	1,980	344





**Table A40-A45: AHA outputs for the South Fork Salmon River Major  
Population Group (MPG) of spring/summer Chinook salmon**

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

### *Parameters and assumptions for spring/summer Chinook in the South Fork Salmon River*

The South Fork Salmon River MPG for spring/summer Chinook includes the Little Salmon River and Rapid River watersheds, the South Fork Salmon River and tributaries, and the intervening section of the mainstem Salmon River. Hatchery programs currently operate on Rapid River (*Rapid River Hatchery Spring Chinook*), the South Fork Salmon River, and Johnson Creek, a tributary to the East Fork of the South Fork Salmon River.

**Table 7A. Mean values of reproductive and survival parameters assumed in the AHA analyses for hatchery-origin spring/summer Chinook released within the South Fork Salmon River MPG. Parameter values were obtained from the HSRG as provided by IDFG.**

<b>Hatchery/ Release location Population/Stock</b>	<b>Fecundity (No. of eggs)</b>	<b>Within- hatchery egg-to-smolt survival</b>	<b>Salmon River smolt-to- adult survival (%)</b>	<b>Recruit/spawner (R/S; adults)</b>
Rapid River FH <i>Rapid River Hatchery Spring Chinook</i>	4,100	0.85	0.47%	6.0
McCall Hatchery/ South Fork Salmon River <i>South Fork Salmon River Summer Chinook</i>	4,300	0.80	0.73%	6.0
McCall Hatchery/ Johnson Creek <i>Johnson Creek Summer Chinook</i>	4,300	0.85	0.45%	6.0

### *Current programs, HSRG solutions, and HRT alternatives*

Of the three principal hatchery programs conducted within the South Fork Salmon River MPG, the HRT reviewed only the South Fork Salmon River summer Chinook program that operates under the LSRCP. AHA analyses for *Rapid River Hatchery Spring Chinook* and *Johnson Creek Summer Chinook* represent those reported by the HSRG. The HRT specifically modeled *HRT Alternatives 2, 4, and 5* for the South Fork Salmon River Summer Chinook program.

*Alternative 2* represents an integrated hatchery program where the current program would be transitioned from the current segregated program (1.0M smolts,  $PNI = 0$ ) to an integrated program that would include natural-origin fish in the broodstock ( $pNOB = 50\%$ ) to achieve a value of  $PNI \geq 0.5$ . The HRT assumed that the current location of the weir is capable of trapping only 50% of returning hatchery-origin adults relative to the total available spawning habitat available in the South Fork Salmon River (i.e., Poverty Flats downstream of the weir and Stolle Meadows upstream of the weir). Based on the productivity and capacity parameters of the Beverton-Holt curve ( $r = R/S_{max} = 3.0$  and  $C = \text{max. number of adult recruits} = 2,140 \text{ adults}$ ), the HRT's analysis indicates that the South Fork Salmon River could support an integrated hatchery program with a maximum release of 490,000

## USFWS Columbia Basin Hatchery Review Team

### *Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011*

hatchery-origin smolts for the entire S.F. population to maintain  $PNI > 0.5$ , a reduction of approximately 50% from the current program. The program would include (a) the trapping and spawning of approximately 500 adults for producing 490,000 smolts and (b) the spawning of approximately 300 additional adults ( $n=294$  “exported” adults assuming a 2:1 male:female sex ratio) to yield 300,000 eyed eggs for the Dollar Creek egg-box program of the Shoshone-Bannock Tribes.

*Alternative 4* represents a “stepping-stone” hatchery program composed of two broodstocks each year. The first broodstock would be integrated with the natural population that spawns in Stolle Meadows upstream of the weir. Smolt offspring of this first broodstock would be tagged but not marked (no adipose fin clip) at the time of release. The second broodstock would use hatchery-origin adults from the first broodstock to produce  $F2$  hatchery-origin fish for harvest. The AHA analyses suggest that the first integrated broodstock could yield 175,000 smolts with  $pNOB = 0.5$ . The second broodstock would yield 825,000 smolts that would be marked with an adipose fin clip. Adult returns from this second broodstock would be sufficient to provide 300,000 eyed eggs (approximately 200 adults, assuming a 1:1 sex ratio) for the Dollar Creek egg box program. The HRT assumed that the current weir is capable of trapping 90% of all hatchery-origin adults intercepted at the weir relative to the naturally spawning population upstream of the weir. *Alternative 4* is the HRT’s “preferred alternative”.

*Alternative 5* is similar to *Alternative 4* except the first broodstock of the stepping stone program would be integrated relative to the entire naturally spawning population in the South Fork (i.e., Stolle Meadows and Poverty Flat). *HRT Alternative 5* and the *HSRG Solution* are similar except (a) the HSRG assumed a female fecundity of 4,700 eggs compared to the assumption of 4,300 eggs by the HRT, (b) the HSRG assumed that the current weir could trap 95% of returning hatchery-origin adults, whereas the HRT assumed only 50% of returning adults could be trapped relative to the entire S.F. Salmon River (i.e., 50% of the adults spawn downstream of the current weir), and (c) the HSRG was able to achieve desired values of  $PNI$  with  $pNOB = 30\%$ , whereas the HRT analysis indicates a required value of  $pNOB = 67\%$  because of the large numbers of hatchery-origin fish that would be expected to spawn downstream of the weir, thus inflating the expected value of  $pHOS$ . These latter analyses yield a release of 250,000 unmarked (but tagged) smolts from the first “integrated” broodstock and 750,000 marked smolts from the second “production” broodstock.

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

**Table 8A. Summary of the number of hatchery-origin (HOR) spring/summer Chinook smolts released into the geographic area of the South Fork Salmon River Major Population Group (MPG) under four strategies modeled in AHA Tables A40-A45: (1) *Current* programs based on the LSRCP Cooperators 2008 Annual Operating Plan for the Salmon River, (2) *HRT Alternative 5* which is similar to the *HSRG Solution*, (3) *HRT Alternative 1* (current programs with implementation of HRT recommendations), and (4) *HRT Alternative 4*, which is the HRT’s “preferred alternative”. The names of hatchery stocks/populations are in italics (e.g., “*Rapid River*”) and are identified by the facility/location where adults are trapped for broodstock.**

Release location (Hatchery)	Current	S.F. Salmon River HRT Alternative 5 (HSRG Solution)	S.F. Salmon River, HRT Alternative 2	S.F. Salmon River, HRT Alternative 4
Rapid River (Rapid River FH) (AHA Tables A40, A41)	<ul style="list-style-type: none"> <li>• 2.7M <i>Rapid River Hatchery</i> smolts</li> </ul>	No recommendation (same as Current)	No recommendation (same as Current)	No recommendation (same as Current)
S.F. Salmon River (McCall FH) (AHA Tables A42, A43; see also AHA Table A44)	<ul style="list-style-type: none"> <li>• 1.0M <i>SF Salmon River</i> smolts (<i>pNOB</i> = 0)</li> <li>• 300K <i>SF Salmon River</i> eyed eggs to Shoshone-Bannock Tribes for egg box program</li> </ul>	<ul style="list-style-type: none"> <li>• 250K <i>SF Salmon River</i> smolts (<i>pNOB</i> = 0.67)</li> <li>• 750K <i>SF Salmon River F2</i> smolts (<i>pNOB</i> = 0)</li> <li>• 300K <i>SF Salmon River F2</i> eyed eggs to Shoshone-Bannock Tribes for egg box program</li> </ul>	<ul style="list-style-type: none"> <li>• 490K <i>S.F. Salmon River</i> smolts (<i>pNOB</i> = 0.5)</li> <li>• 300K <i>SF Salmon River</i> eyed eggs to Shoshone-Bannock Tribes for egg box program</li> </ul>	<ul style="list-style-type: none"> <li>• 175K <i>SF Salmon River</i> smolts (<i>pNOB</i> = 0.5)</li> <li>• 825K <i>SF Salmon River F2</i> smolts (<i>pNOB</i> = 0)</li> <li>• 300K <i>SF Salmon River F2</i> eyed eggs to Shoshone-Bannock Tribes for egg box program</li> </ul>
Johnson Creek (McCall FH) (AHA Table A45)	<ul style="list-style-type: none"> <li>• 100K <i>Johnson Cr.</i> smolts</li> </ul>	No recommendation (same as Current)	No recommendation (same as Current)	No recommendation (same as Current)

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A40: AHA output for spring/summer Chinook in the Little Salmon River. *Rapid River Hatchery Spring Chinook* are modeled in Table A41. HRT Alternatives refer to the South Fork Salmon River summer Chinook program.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest				
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery				
10-Salmon_Little Salmon Spring-Summer Chinook		Adult Passage		Population Recovery Designation:		Stabilizing		Stabilizing		Stabilizing		Stabilizing		Stabilizing				
Chinook		Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2		HRT Alternative 4						
Habitat	Productivity (Adult)		Ad. Capacity	1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250			
	Min NOR Escape		% Kelt	1		1		1		1		1		1				
	Smolt Productivity		Sm. Capacity	33	31,250	161	154,603	161	154,603	161	154,603	161	154,603	161	154,603			
Hydro	Ocean Surv		Baseline SAR	Vary? (Y/N)	0.098	0.040	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y		
	Juv Passage Surv.		Adult Passage		1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82		
	Adjusted Productivity		Adj. Capacity		1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250	1.30	1,250		
Harvest	Harv - Marine		NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010			
	Harv - L. Mainstem		NORs		HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080			
	Harv - U. Mainstem		NORs		HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080			
	Harv -Terminal		NORs		HORs	0.080	0.500	0.080	0.500	0.080	0.500	0.080	0.500	0.080	0.500			
	Total Exploitation Rate		NORs		HORs	0.152	0.581	0.152	0.581	0.152	0.581	0.152	0.581	0.152	0.581			
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS	
	Purpose		PNI		pHOS-Realized		59%		59%		59%		59%		59%		59%	
	Type		Cons/Harv/Both		Int/Seg/Step/None		Harv		None		Harv		None		Harv		None	
	Broodstock by Source		Local		Imported													
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays		484		477		474		466		466		466	
Patterns	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		100%		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		y		y		y		y		y		y	
	Update		Hatchery															

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.50	0.50
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.50	0.50	0.50
Generations until average fitness is reached ->	15	30	15	15	15
Minimum Hatchery Program (% of BS Goal) ->					
"Fitness Floor" ->	0.5				
Calculated Hatchery SAR ->					
Calculated Natural SAR ->	0.8%	0.8%	0.8%	0.8%	0.8%
NOR Escapement	994	100	268	261	14
HoS Total Escapement (includes strays)	1,479	257	484	1,457	253

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A41. AHA output for Rapid River Hatchery Spring Chinook released at the hatchery into the Rapid River (see also Table A40). HRT Alternatives refer to the South Fork Salmon River summer Chinook program.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest		Conservation		Harvest		Harvest		Harvest	
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Segregated Hatchery		Segregated Hatchery	
10A-Salmon_Little Salmon Spring Chinook		Population Recovery Designation:		N/A		N/A		N/A		N/A		N/A		N/A	
(Rapid River-Hatchery)		Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2		HRT Alternative 4			
Habitat	Productivity (Adult)	Ad. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1
	Smolt Productivity	Sm. Capacity	0	0	1	1	1	1	1	1	1	1	1	1	1
Hydro	Ocean Surv	Baseline SAR	0.098	0.040	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34
	Adjusted Productivity	Adj. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem	NORs		HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020
	Harv - U. Mainstem	NORs		HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050
	Harv -Terminal	NORs		HORs	0.080	0.500	0.080	0.500	0.080	0.500	0.080	0.500	0.080	0.500	0.080
	Total Exploitation Rate	NORs		HORs	0.152	0.581	0.152	0.581	0.152	0.581	0.152	0.581	0.152	0.581	0.152
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB
	Purpose		PNI	pHOS-Realized		100%		100%		100%		100%		100%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Harv	Seg	Cons	None	Harv	Seg	Harv	Seg	Harv	Seg	Harv
	Broodstock by Source		Local	Imported	1,745	2,736,596			1,745	2,736,596	1,745	2,736,596	1,745	2,736,596	1,745
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release	220	220			220	220	220	220	220	220	220
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.	90%	10%		100%	90%	10%	90%	10%	90%	10%	90%
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	6.0	y	6.0	y	6.0	y	6.0	y	6.0	y	6.0
	Update	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat	Current Habitat			Current Habitat		Current Habitat		Current Habitat		HRT Alt-B	
				Hydro	Current Hydro			Current Hydro		Current Hydro		Current Hydro		Current Hydro	
				Harvest	Current Harvest			Current Harvest		HSRG Harvest		Current Harvest		HSRG Harvest	
		Hatchery													

485

2

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80							
Weir Factor ->												
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%							
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->	0.19	1.00	0.19	0.19	0.19							
Average Fitness Factor (100 Generations) ->	0.24	0.99	0.24	0.24	0.24							
Generations until average fitness is reached ->	43	30	43	43	43							
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS ->	1.00	Effective pHOS ->	1.00							
"Fitness Floor" ->		pNOB ->		pNOB ->								
Calculated Hatchery SAR ->	0.4%	0.4%	0.4%	0.4%	0.4%							
Calculated Natural SAR ->	0.8%	0.8%	0.8%	0.8%	0.8%							
NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)	1,494	259	489	1,494	259	489	1,494	259	489	1,494	259	489

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A42. AHA output for South Fork Salmon River summer Chinook. The HSRG solution and HRT Alternative 5 are essentially identical. HRT Alternatives 4 and 5 represent “stepping-stone” programs. Outputs for the “production” broodstocks are presented in Table A43.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Conservation		Conservation and Harvest		Conservation	
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Stepping Stone Hatchery		Integrated Hatchery		Stepping Stone Hatchery	
11-Salmon_SF Salmon Summer Chinook				Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary	
						Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2	
Habitat	Productivity (Adult)		Ad. Capacity	3.00	2,150	3.00	2,150	3.00	2,150	3.00	2,150	3.00	2,150	3.00	1,075
	Min NOR Escape		% Kelt	1		1		1		1		1		1	
	Smolt Productivity		Sm. Capacity	300	215,000	371	265,918	371	265,918	371	265,918	371	265,918	371	132,959
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.098	0.010	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y
	Juv Passage Surv.		Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82
	Adjusted Productivity		Adj. Capacity	3.00	2,150	3.00	2,150	3.00	2,150	3.00	2,150	3.00	2,150	3.00	1,075
Harvest	Harv - Marine		NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem		NORs		HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.020
	Harv - U. Mainstem		NORs		HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.050
	Harv -Terminal		NORs		HORs	0.080	0.300	0.080	0.300	0.080	0.080	0.080	0.300	0.080	0.080
	Total Exploitation Rate		NORs		HORs	0.152	0.413	0.152	0.413	0.152	0.152	0.152	0.413	0.152	0.152
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB	pHOS		pNOB	pHOS		pNOB	pHOS	
	Purpose		PNI		pHOS-Realized			60%							
	Type		Cons/Harv/Both		Int/Seg/Step/None		Harv	Seg		Cons	None		Both	Int	
	Broodstock by Source		Local	979	Smolt Release			1,000,224							
	Brood Exported (from HOR Surplus)		Imported	294	Export Goal/Realized		294	3		245	250,235		480	490,406	
Patterns	Destination for HOR Returns		% to Hatchery	50%	% to Nat. Spawn.			100%		485	909		294	3	
	Productivity of Hatchery Fish		Recruits/Spawner	6.0	Fitness? [Y / N]			y		50%	50%		50%	50%	
Update		Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		HRT Alt-B	
		Hydro		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro	
		Harvest		Harvest		Current Harvest		Current Harvest		HSRG Harvest		Current Harvest		HSRG Harvest	
		Hatchery													

459

3

Realized Spawning Composition

Relative Hatchery Optimum ->		80	80	80	80	80										
Weir Factor ->																
Relative Reproductive Success (HOS) ->		80%	80%	80%	80%	80%										
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->		0.50	1.00	0.70	0.63	0.73										
Average Fitness Factor (100 Generations) ->		0.50	0.99	0.71	0.66	0.74										
Generations until average fitness is reached ->		15	30	31	38	31										
Minimum Hatchery Program (% of BS Goal) ->		0.5	Effective pHOS -> 0.60	Effective pHOS -> 0.55	Effective pHOS -> 0.50	Effective pHOS -> 0.36										
"Fitness Floor" ->			pNOB ->	pNOB ->	pNOB ->	pNOB ->										
Calculated Hatchery SAR ->		0.6%														
Calculated Natural SAR ->		0.8%														
NOR Escapement		Max 3,432	Min 487	Ave 1,008	Max 4,218	Min 570	Ave 1,218	Max 4,001	Min 423	Ave 1,066	Max 3,261	Min 195	Ave 759	Max 1,714	Min 138	Ave 426
HoS Total Escapement (includes strays)		5,879	1,021	1,925				4,898	850	1,604	2,887	501	945	915	158	299

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A43. AHA output for the F2 “production” broodstocks associated with *HRT Alternatives 4* and 5 in the South Fork Salmon River.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Harvest		Conservation and Harvest		Conservation and Harvest		
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery only		Natural Reproduction only		Segregated Stepping Stone Broodstock		Integrated Hatchery only		Segregated Stepping Stone Broodstock		
11A-Salmon_SF Salmon Summer Chinook (McCall-Hatchery)		Population Recovery Designation:		N/A		N/A		N/A		N/A		N/A		N/A		
		Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2		HRT Alternative 4				
Habitat	Productivity (Adult)	Ad. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1	
	Smolt Productivity	Sm. Capacity	0	0	1	1	1	1	1	1	1	1	1	1	1	
Hydro	Ocean Surv	Baseline SAR	0.096	0.040	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	
	Adjusted Productivity	Adj. Capacity	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0.01	0	0	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	
	Harv - U. Mainstem	NORs	HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	
	Harv -Terminal	NORs	HORs	0.080	0.300	0.080	0.300	0.080	0.300	0.080	0.300	0.080	0.300	0.080	0.300	
	Total Exploitation Rate	NORs	HORs	0.152	0.413	0.152	0.413	0.152	0.413	0.152	0.413	0.152	0.413	0.152	0.413	
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB	
	Purpose		PNI		pHOS-Realized		None		None		None		None		None	
	Type		Cons/Harv/Both		Int/Seg/Step/None		None		None		Harv		Seg		Harv	
	Broodstock by Source		Local		Imported		194		485		194		750,780		533	
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Strays		50%		50%		50%		100%		90%	
Patterns	Destination for HOR Returns		% to Nat. Spawn.		100%		100%		100%		100%		100%		10%	
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		y		y		y		y		y	
	Update		Habitat		Hydro		Current Habitat		Current Hydro		Current Habitat		Current Hydro		HRT Alt-B	
		Harvest		Hatchery		Current Harvest		Current Harvest		HSRG Harvest		Current Harvest		HSRG Harvest		

Realized Spawning Composition

Realized Spawning Composition charts showing PNI, pHOS, and pNOB for various hatcheries and harvest scenarios. The charts include a 3D surface plot and several bar charts comparing different scenarios.

Relative Hatchery Optimum ->	80	80	80	80	80							
Weir Factor ->												
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%							
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81							
Fitness Factor after 100 generations (B) ->	1.00	1.00	0.19	1.00	0.19							
Average Fitness Factor (100 Generations) ->	0.99	0.99	0.24	0.99	0.24							
Generations until average fitness is reached ->	30	30	43	30	43							
Minimum Hatchery Program (% of BS Goal) ->												
"Fitness Floor" ->												
Calculated Hatchery SAR ->	0.8%	0.8%	0.4%	0.8%	0.4%							
Calculated Natural SAR ->												
NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
HoS Total Escapement (includes strays)	0	0	0	0	0	0	2,907	505	952	639	111	209



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A44. AHA output for Secesh River spring Chinook. HRT Alternatives refer to South Fork Salmon River summer Chinook program.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only	
12-Salmon_Secesh Spring Chinook		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary	
		Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2		HRT Alternative 4			
Habitat	Productivity (Adult)	Ad. Capacity	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	41	33,750	200	166,972	200	166,972	200	166,972	200	166,972	200	166,972	
Hydro	Ocean Surv	Baseline SAR	0.092	0.040	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	
	Adjusted Productivity	Adj. Capacity	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	1.62	1,350	
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs	HORs	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	
	Harv - U. Mainstem	NORs	HORs	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	
	Harv -Terminal	NORs	HORs	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	
	Total Exploitation Rate	NORs	HORs	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized	None	20%	None	None	None	14%	None	10%	None	1%	
	Broodstock by Source		Local	Imported											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		23		16		13		5			
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%			
Patterns	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	y	y	y	y	y	y	y	y	y	y	
	Update		Habitat	Hydro	Harvest	Hatchery	Current Habitat	Current Hydro	Current Harvest	HRT Alt-B	Current Habitat	Current Hydro	Current Harvest	Current Habitat	

Realized Spawning Composition

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A45. AHA output for Johnson Creek summer Chinook. HRT Alternatives refer to South Fork Salmon River summer Chinook program.

All H Analyzer (AHA)

Apply & Recalculate


Population (Ctrl + Page Up/Down to Scroll)


Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
S.F. Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Integrated Hatchery		Natural Reproduction only		Integrated Hatchery		Integrated Hatchery		Integrated Hatchery	
13-Salmon_EF-SF Johnson Creek Summer Chinook				Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary	
				Historic		Current		No Hatchery		HRT Alt. 5; HSRG Sol.		HRT Alternative 2		HRT Alternative 4	
Habitat	Productivity (Adult)	Ad. Capacity	1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	36	42,500	179	210,261	179	210,261	179	210,261	179	210,261	179	210,261	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.040	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82
	Adjusted Productivity	Adj. Capacity		1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700	1.45	1,700
Harvest	Harv - Marine	NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem	NORs	HORs	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020
	Harv - U. Mainstem	NORs	HORs	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050
	Harv -Terminal	NORs	HORs	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005	0.005
	Total Exploitation Rate	NORs	HORs	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083	0.083
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	100%	36%	100%	36%	100%	36%	100%	36%	100%	35%	
	Purpose		PNI	pHOS-Realized	0.74	36%	0.74	36%	0.74	36%	0.74	36%	0.74	35%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	Both	Int	None	None	Both	Int	Both	Int	Both	Int	
	Broodstock by Source		Local	Imported	62	101,810			62	101,810	62	101,810	62	101,810	
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		22				15		12		4	
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%		100%	
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]	6.0	Y	6.0	Y	6.0	Y	6.0	Y	6.0	Y	
	Update		Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		HRT Alt-B
		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro	
		Harvest		Current Harvest		Current Harvest		HSRG Harvest		Current Harvest		Current Harvest		HSRG Harvest	
		Hatchery													


456


6


Realized Spawning Composition














Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.87	1.00	0.87	0.87	0.88
Average Fitness Factor (100 Generations) ->	0.86	0.99	0.86	0.86	0.87
Generations until average fitness is reached ->	35	30	35	35	35
Minimum Hatchery Program (% of BS Goal) ->	0.5	Effective pHOS ->	0.36	Effective pHOS ->	0.36
"Fitness Floor" ->		pNOB ->	0.50	pNOB ->	0.50
Calculated Hatchery SAR ->	0.4%	0.4%	0.4%	0.4%	0.4%
Calculated Natural SAR ->	0.8%	0.8%	0.8%	0.8%	0.8%
NOR Escapement	Max 2,224	Min 178	Ave 571	Max 1,078	Min 109
HoS Total Escapement (includes strays)	2,224	178	571	1,078	109

**Table A46-A56: AHA outputs for the Upper Salmon River Major  
Population Group (MPG) of spring/summer Chinook salmon**

## USFWS Columbia Basin Hatchery Review Team

*Lower Snake Federal Hatcheries Assessments and Recommendations Reports—March 2011*

### *Parameters and assumptions for spring/summer Chinook in the Upper Salmon River Major Population Group (MPG)*

The Upper Salmon River MPG for spring/summer Chinook includes the mainstem Salmon River and all tributaries upstream of, and including, the North Fork Salmon River. The Interior Columbia Technical Recovery Team (ICTRT) identified nine demographically independent populations within this MPG (AHA Tables A46-A56). Two hatchery populations are also present within this MPG: *Pahsimeroi Hatchery Summer Chinook* and *Sawtooth Hatchery Spring Chinook*.

**Table 9A. Mean values of reproductive and survival parameters assumed in the AHA analyses for hatchery-origin spring/summer Chinook released within the Upper Salmon River MPG. These parameter values were obtained from the HSRG, as provided by IDFG.**

<b>Hatchery/ Release location Population/stock</b>	<b>Fecundity (No. of eggs)</b>	<b>Within- hatchery egg-to-smolt survival</b>	<b>Salmon River smolt-to- adult survival (%)</b>	<b>Recruit/spawner (R/S; adults)</b>
Pahsimeroi FH <i>Pahsimeroi Hatchery Summer Chinook</i>	5,000	0.80	0.39%	6.0
Sawtooth FH <i>Sawtooth Hatchery Spring Chinook - Current</i>	4,900	0.80	0.13%	2.0
Sawtooth FH <i>Sawtooth Hatchery Spring Chinook: HSRG Solution and HRT Alternatives A and B</i>	4,900	0.80	0.18%	2.5
Sawtooth FH/ <i>E.F. Salmon River Spring/Summer Chinook, Yankee Fork Spring Chinook Valley Creek Spring Chinook</i>	4,900	0.80	0.18%	2.5

### *Current programs, HSRG solutions, and HRT alternatives*

The HRT reviewed one hatchery program in the upper Salmon River basin: the spring Chinook program at Sawtooth FH (Table 10A). The intent of the current program at the Sawtooth FH is to release 1.0M *Sawtooth Hatchery Spring Chinook* smolts annually into the mainstem Salmon River at the hatchery weir. Currently, the *Sawtooth Hatchery Spring Chinook* stock is propagated as a segregated population ( $pNOB = 0$ ).

The *HSRG solution* involved the establishment of a second, conservation hatchery program derived from natural-origin fish trapped at the Sawtooth Hatchery weir with a mean equilibrium value of  $pNOB = 67\%$  and an annual release of 200,000 smolts. The existing segregated hatchery population would be maintained with a maximum release of 1.2M smolts. The *HSRG Solution* requires the

## USFWS Columbia Basin Hatchery Review Team

### *Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011*

maintenance of two independent hatchery populations. The HRT adopted this *HSRG Solution* as *HRT Alternative 4*.

*HRT Alternative 2* would convert the existing segregated hatchery program to an integrated hatchery program. *AHA* modelling suggests that current habitat conditions would restrict an integrated hatchery program to a maximum release of 500,000 smolts, with  $pNOB = 0.5$ .

*HRT Alternative 3* would convert the existing segregated hatchery program to a stepping stone hatchery program composed of two broodstocks where returning adults from the first “integrated” broodstock would be used as parents for the second “production” broodstock. *AHA* modeling suggests that current habitat conditions would restrict each of the two broodstocks in a stepping-stone program to a maximum release of 375,000 smolts each, with a combined maximum release of 750,000 smolts, with  $pNOB = 0.5$  for the first broodstock, and  $pNOB = 0$  for the second broodstock. These latter results are consistent with *HSRG* conclusions that a stepping stone program would not be able to meet the total desired release of 1.0M smolts under current habitat conditions. The HRT favored *Alternative 3* over *Alternative 4* (i.e., the *HSRG* solution) because of the large numbers of hatchery-origin spring Chinook that currently spawn in the mainstem Salmon River downstream from the hatchery weir and the presumed inability to preclude large numbers of hatchery-origin fish from a “segregated” hatchery program from spawning naturally in the mainstem Salmon River. The HRT believed this latter risk could be reduced via a “stepping stone” program, albeit with a 25% reduction in the maximum number of smolts available for release.

The HRT also examined alternatives for initiating hatchery programs on the East Fork Salmon River, Yankee Fork, and Valley Creek. Two scenarios were modeled for each population. A small integrated conservation program of 52,000 smolts ( $pNOB = 1.0$ ) and a large segregated harvest program of 1.0M smolts (*HRT Alternatives A* and *B*, respectively) were each modeled for the E.F. Salmon River and Valley Creek. The former integrated programs of 52,000 smolts were the sizes that maximized the predicted number of natural-origin recruits returning to each of the two streams. This latter number of smolts also maximized the number of natural-origin recruits in Yankee Fork. In all three streams, productivity and capacity constraints inhibited the long-term conservation benefit of hatchery-origin fish spawning naturally. The *AHA* analyses also suggest that large segregated hatchery programs in Valley Creek and the East Fork Salmon River would not be able to meet *LSRCP* mitigation goals without large numbers of hatchery-origin fish escaping fisheries and potentially spawning naturally, thereby increasing biological risks to *ESA* listed natural populations under current harvest regimes. Based on those analyses, the HRT modeled two integrated programs for Yankee Fork, each with an annual release of 52,000 smolts, but with  $pNOB = 1.0$  and  $0.5$ , respectively.

# USFWS Columbia Basin Hatchery Review Team

## Lower Snake Federal Hatcheries Assessments and Recommendations Reports–March 2011

**Table 10A. Summary of the number of hatchery-origin (HOR) spring/summer Chinook smolts released into the geographic area of the Upper Salmon River MPG (upstream of the confluence of the Middle Fork Salmon River) under four strategies modeled in AHA Tables A46-A56: (1) Current programs based on the LSRCP Cooperators 2008 Annual Operating Plan for the Salmon River, (2) the HSRG Solution, and (3) two HRT Alternatives.** [Note: The analyses for the E.F. Salmon River, Yankee Fork, and Valley Creek make no explicit assumptions regarding the location of weirs in each of those three streams. The results of the AHA analyses are based on the Beverton-Holt habitat parameters of productivity and capacity, as reported by the HSRG. Those parameters refer to the entire stream and, hence, implicitly assume that weirs would be located downstream from all natural spawning areas. With respect to the E.F. Salmon River, the analysis implicitly assumes that all spawning areas for spring Chinook are upstream of the current weir, or the weir would be moved downstream from its present location to a location on the East Fork that is downstream from all natural spawning areas.]

Release location (Hatchery)	Current	HSRG Solution	HRT Alternative A	HRT Alternative B
Pahsimeroi River (Pahsimeroi FH) (AHA Tables A48, A49)	• 1.0M <i>Pahsimeroi River</i> smolts ( <i>pNOB</i> = 0)	• 285K <i>Pahsimeroi River</i> smolts ( <i>pNOB</i> = 0.75) • 1.0M <i>Pahsimeroi River F2</i> smolts ( <i>pNOB</i> = 0)	No recommendation (same as HSRG Solution)	No recommendation (same as HSRG Solution)
E.F. Salmon River (AHA Table A51))	No program	No recommendation (Same as Current)	• 52K <i>E.F. Salmon River</i> smolts ( <i>pNOB</i> = 1.0)	• 1.0M <i>E.F. Salmon River</i> smolts ( <i>pNOB</i> = 0)
Yankee Fork (AHA Table A52)	No program	No recommendation (same as Current)	<b><u>HRT Alternative 1a</u></b> • 52K <i>Yankee Fork</i> smolts ( <i>pNOB</i> = 1.0)	<b><u>HRT Alternative 1b</u></b> • 52K <i>Yankee Fork</i> smolts ( <i>pNOB</i> = 0.5)
Valley Creek (AHA Table A53)	No program	No recommendation (same as Current)	• 52K <i>Valley Creek</i> smolts ( <i>pNOB</i> = 1.0)	• 1.0 M <i>Valley Creek</i> smolts ( <i>pNOB</i> = 0)
Upper Salmon River (Sawtooth FH) (AHA Tables A54, A55)	• 1.0M <i>Sawtooth</i> smolts ( <i>pNOB</i> = 0)	<b><u>HRT Alternative 4</u></b> • 200K <i>Upper Salmon River</i> smolts ( <i>pNOB</i> = 0.67) • 1.2M <i>Sawtooth</i> smolts ( <i>pNOB</i> = 0)	<b><u>HRT Alternative 2</u></b> • 500K <i>Upper Salmon River</i> smolts ( <i>pNOB</i> = 0.5)	<b><u>HRT Alternative 3</u></b> • 375K <i>Upper Salmon River</i> smolts ( <i>pNOB</i> = 0.5) • 375K <i>Upper Salmon River F2</i> smolts ( <i>pNOB</i> = 0)

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A46. AHA output for North Fork Salmon River spring Chinook.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

24-Salmon\_NF Salmon River Spring

Chinook

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Habitat

Productivity (Adult)

Ad. Capacity

Min NOR Escape

% Kelt

Smolt Productivity

Sm. Capacity

1.60

550

1

13,750

1.60

550

1

68,026

1.60

550

1

68,026

1.60

550

1

68,026

1.60

550

1

68,026

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

Adj. Capacity

0.092

0.040

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

Harvest

Harv - Marine

NORs

HORs

0.010

0.010

Harv - L. Mainstem

NORs

HORs

0.020

0.080

Harv - U. Mainstem

NORs

HORs

0.050

0.080

Harv -Terminal

NORs

HORs

0.005

0.100

Total Exploitation Rate

NORs

HORs

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

Local

Imported

Export Goal/Realized

Smolt Release

% to Hatchery

% to Nat. Spawn.

Recruits/Spawner

Fitness? [Y / N]

pNOB

pHOS

30%

None

None

None

None

28%

None

None

None

None

28%

None

None

None

None

22%

None

None

24

100%

y

100%

y

18

100%

y

19

100%

y

10

100%

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

HSRG

Current Habitat

Current Hydro

Current Harvest

HRT-A

Current Habitat

Current Hydro


Current Harvest


HRT-B

534

1

Realized Spawning Composition





Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

80

80%

0.81

0.50

0.50

21

0.5

0.8%

0.8%

146

14

46

502

55

168

122

11

39

123

11

39

57

10

19

31

5

10

80

80%

0.81

1.00

0.99

30

0.30

0.8%

0.8%

56

10

18

57

10

19

31

5

10

80

80%

0.81

0.50

0.50

22

0.28

0.8%

0.8%

56

10

18

57

10

19

31

5

10

80

80%

0.81

0.50

0.50

22

0.28

0.8%

0.8%

56

10

18

57

10

19

31

5

10

80

80%

0.81

0.50

0.52

0.22

0.8%

0.8%

56

10

18

57

10

19

31

5

10

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A47. AHA output for Lemhi River spring Chinook.

All H Analyzer (AHA)

Apply & Recalculate







Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Conservation	
Upper Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only	
25-Salmon_Lemhi River Spring Chinook		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary	
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative A		HRT Alternative B			
Habitat	Productivity (Adult)	Ad. Capacity	1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	33	97,500	162	482,363	162	482,363	162	482,363	162	482,363	162	482,363	
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.040	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82
	Adjusted Productivity	Adj. Capacity		1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900	1.31	3,900
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	Harv - L. Mainstem		NORs	HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020
	Harv - U. Mainstem		NORs	HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050
	Harv -Terminal		NORs	HORs	0.005	0.100	0.005	0.100	0.005	0.100	0.005	0.100	0.005	0.100	0.005
	Total Exploitation Rate		NORs	HORs	0.083	0.246	0.083	0.246	0.083	0.246	0.083	0.246	0.083	0.246	0.083
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	
	Purpose		PNI	pHOS-Realized		30%				27%		27%		15%	
	Type		Cons/Harv/Both	Int/Seg/Step/None	None	None	None	None	None	None	None	None	None	None	
	Broodstock by Source		Local	Imported											
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Smolt Release		24		18		19		10			
	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%			
Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y		y		y		
Patterns	Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.		Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		Current Habitat		
	Update		Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		Current Hydro		
		Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest		Current Harvest			
		Hatchery						HSRG		HRT-A		HRT-B			

453

2

Realized Spawning Composition



Relative Hatchery Optimum ->		80	80	80	80	80
Weir Factor ->						
Relative Reproductive Success (HOS) ->		80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->		0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->		0.50	1.00	0.50	0.50	0.50
Average Fitness Factor (100 Generations) ->		0.51	0.99	0.52	0.52	0.57
Generations until average fitness is reached ->			30			39
Minimum Hatchery Program (% of BS Goal) ->			Effective pHOS ->	Effective pHOS ->	Effective pHOS ->	Effective pHOS ->
"Fitness Floor" ->		0.5	pNOB ->	pNOB ->	pNOB ->	pNOB ->
Calculated Hatchery SAR ->						
Calculated Natural SAR ->		0.8%	0.8%	0.8%	0.8%	0.8%
		Max	Min	Ave	Max	Min
NOR Escapement		214	10	46	1,551	116
HoS Total Escapement (includes strays)		74	13	24		



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A48. AHA output for Pahsimeroi River summer Chinook.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Upper Salmon R

Species/Race

Spring Chinook

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Harvest and Conservation

Segregated Hatchery

Primary

Conservation

Natural Reproduction only

Primary

Harvest and Conservation

Stepping Stone Integrated Hatchery

Primary

Harvest and Conservation

Stepping Stone Integrated Hatchery

Primary

Harvest and Conservation

Stepping Stone Integrated Hatchery

Primary

26-Salmon\_Pahsimeroi Summer Chinook

Habitat

Productivity (Adult)

Ad. Capacity

Min NOR Escape

Smolt Productivity

1.70

3,200

1

80,000

43

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

Adj. Capacity

0.092

0.040

y

1.00

1.00

1.70

3,200

0.029

0.008

y

0.34

0.82

1.70

3,200

0.029

0.008

y

0.34

0.82

1.70

3,200

0.029

0.008

y

0.34

0.82

1.70

3,200

0.029

0.008

y

0.34

0.82

1.70

3,200

0.029

0.008

y

0.34

0.82

1.70

3,200

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

NORs

NORs

NORs

NORs

NORs

HORs

HORs

HORs

HORs

HORs

HORs

0.010

0.020

0.050

0.080

0.152

0.010

0.080

0.080

0.100

0.246

0.010

0.020

0.050

0.080

0.152

0.010

0.020

0.050

0.080

0.152

0.010

0.020

0.050

0.080

0.152

0.010

0.020

0.050

0.080

0.152

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

Local

Imported

Smolt Release

Export Goal/Realized

% to Hatchery

Recruits/Spawner

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

Strays

% to Nat. Spawn.

Fitness? [Y / N]

pNOB

37%

Harv

Seg

526

999,400

23

90%

10%

6.0

y

pNOB

75%

Cons

Step

150

285,000

478

90%

10%

6.0

y

pNOB

75%

Cons

Step

150

285,000

478

90%

10%

6.0

y

pNOB

75%

Cons

Step

150

285,000

478

90%

10%

6.0

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

HSRG

Current Habitat

Current Hydro


HRT-A

Current Habitat

Current Hydro

HRT-B

Realized Spawning Composition



Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

80

80%

0.81

0.50

0.50

18

0.5

0.3%

0.8%

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

0.37

0.33

0.38

0.33

0.38

0.33

0.38

0.33

0.38

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

NOR Escapement

HoS Total Escapement (includes strays)

1,362

128

398

2,550

273

863

3,561

269

920

3,561

269

921

3,540

267

916

881

152

288

1,719

298

563

1,720

299

563

1,694

294

554

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A49. AHA output for Pahsimeroi Hatchery summer Chinook resulting from the second segregated broodstock of a two-stage “stepping stone” hatchery program, as proposed by the HSRG.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

26A-Salmon\_Pahsimeroi Summer Chinook (Pahsimeroi Hatchery)

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Harvest

Segregated Hatchery

N/A

Conservation

Natural Reproduction only

N/A

Harvest

Stepping Stone Segregated Hatchery

N/A

Harvest

Stepping Stone Segregated Hatchery

N/A

Harvest

Stepping Stone Segregated Hatchery

N/A

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Habitat

Productivity (Adult)

Min NOR Escape

Smolt Productivity

Ad. Capacity

% Kelt

Sm. Capacity

0.00

0

0.00

0

0.00

0

1

1

1

1

1

1

0

0

0

1

0

1

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

Adj. Capacity

0.092

0.040

y

1.00

1.00

0.00

0

0.029

0.008

y

0.34

0.82

0.00

0

0.029

0.008

y

0.34

0.82

0.00

0

0.029

0.008

y

0.34

0.82

0.00

0

0.029

0.008

y

0.34

0.82

0.00

0

0.029

0.008

y

0.34

0.82

0.00

0

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

HORs

NORs

HORs

NORs

HORs

0.010

0.080

0.080

0.080

0.100

0.246

0.010

0.080

0.080

0.080

0.100

0.246

0.010

0.080

0.080

0.080

0.100

0.246

0.010

0.080

0.080

0.080

0.100

0.246

0.010

0.080

0.080

0.080

0.100

0.246

0.010

0.080

0.080

0.080

0.100

0.246

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

pHOS-Goal

Cons/Harv/Both

Int/Seg/Step/None

Local

Imported

Smolt Release

Strays

% to Hatchery

% to Nat. Spawn.

Recruits/Spawner

Fitness? [Y / N]

Harv

Seg

None

None

Harv

Step

Harv

Step

Harv

Step

Harv

Step

Harv

Step

526

999,400

526

999,400

526

999,400

526

999,400

526

999,400

526

999,400

50%

50%

50%

50%

50%

50%

50%

50%

50%

50%

50%

50%

6.0

y

6.0

y

6.0

y

6.0

y

6.0

y

6.0

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Realized Spawning Composition

1.00

0.80

0.60

0.40

0.20

0

0.1

0.2

0.3

0.4

0.5

0.6

0.7

0.8

0.9

1.00

pNOB

pHOS

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

2,000

1,800

1,600

1,400

1,200

1,000

800

600

400

200

0

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

80

80%

0.81

1.00

0.99

30

0.5

80

80%

0.81

1.00

0.99

30

Effective pHOS ->

pNOB ->

0.8%

0

0

0

80

80%

0.81

0.50

0.50

12

Effective pHOS ->

pNOB ->

0.3%

0.8%

0

0

0

80

80%

0.81

0.50

0.50

12

Effective pHOS ->

pNOB ->

0.3%

0.8%

0

0

0

80

80%

0.81

0.50

0.50

12

Effective pHOS ->

pNOB ->

0.3%

0.8%

0

0

0

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

4,054

704

1,327

4,054

704

1,327

4,054

704

1,327

4,054

704

1,327

4,054

704

1,327

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A50. AHA output for spring Chinook spawning in lower section of upper Salmon River, from mouth of Lemhi River to Redfish Lake Creek.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

27-Salmon\_Lower Salmon Mainstem Spring Chinook

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Habitat

Productivity (Adult)

Ad. Capacity

Min NOR Escape

% Kelt

Smolt Productivity

Sm. Capacity

1.50

2,000

1

50,000

186

247,366

186

247,366

186

247,366

186

247,366

186

247,366

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

Adj. Capacity

0.092

0.040

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

Harvest

Harv - Marine

NORs

HORs

Harv - L. Mainstem

NORs

HORs

Harv - U. Mainstem

NORs

HORs

Harv -Terminal

NORs

HORs

Total Exploitation Rate

NORs

HORs

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

Local

Imported

Smolt Release

Export Goal/Realized

Strays

% to Hatchery

% to Nat. Spawn.

Recruits/Spawner

Fitness? [Y / N]

pNOB

pHOS

22%

None

None

24

100%

y

pNOB

pHOS

30%

None

None

61

100%

y

pNOB

pHOS

20%

None

None

21

100%

y

pNOB

pHOS

23%

None

None

25

100%

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

HSRG

Current Habitat

Current Hydro

HRT-A

Current Habitat


Current Hydro


HRT-B


636


5


Realized Spawning Composition














Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

Max

Min

Ave

356

14

68

74

13

24

Max

Min

Ave

1,457

152

514

Max

Min

Ave

367

32

112

186

32

61

Max

Min

Ave

388

13

67

64

11

21

Max

Min

Ave

350

15

68

76

13

25

80

80

80%

0.81

1.00

0.99

30

0.22

0.30

0.20

0.23

0.8%







0.8%

0.8%

0.8%

0.8%

**Realized Spawning Composition**



Relative Hatchery Optimum ->				80			80			80			80			80								
Weir Factor ->																								
Relative Reproductive Success (HOS) ->				80%			80%			80%			80%			80%								
Initial Fitness Factor (A) ->				0.81			0.81			0.81			0.81			0.81								
Fitness Factor after 100 generations (B) ->				0.50			1.00			0.50			0.50			0.50								
Average Fitness Factor (100 Generations) ->				0.52			0.99			0.50			0.53			0.52								
Generations until average fitness is reached ->							30			22						30								
Minimum Hatchery Program (% of BS Goal) ->							Effective pHOS -> 0.22			Effective pHOS -> 0.30			Effective pHOS -> 0.20			Effective pHOS -> 0.23								
"Fitness Floor" ->				0.5			pNOB ->			pNOB ->			pNOB ->			pNOB ->								
Calculated Hatchery SAR ->				0.8%						0.8%						0.8%								
Calculated Natural SAR ->										0.8%						0.8%								
				Max			Min			Ave			Max			Min			Ave					
NOR Escapement				356			14			68			1,457			152			514					
HoS Total Escapement (includes strays)				74			13			24						186			32			61		
																64			11			21		
																76			13			25		

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A51. AHA output for East Fork Salmon River spring/summer Chinook. HRT Alternative A represents a small, integrated conservation program, and HRT Alternative B represents a large segregated program in partial support for achieving LSRCP mitigation goals.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Conservation		Conservation		Conservation		Conservation		Harvest	
Upper Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Natural Reproduction only		Natural Reproduction only		Natural Reproduction only		Integrated Hatchery + Supplementation		Segregated Hatchery	
28-Salmon_East Fork Salmon River Spring-Summer Chinook		Population Recovery Designation:		Contributing		Contributing		Contributing		Contributing		Contributing		Contributing	
				Historic		Current		No Hatchery		HSRG Solution		HRT Alternative A		HRT Alternative B	
Habitat	Productivity (Adult)	Ad. Capacity	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	
	Min NOR Escape	% Kelt	1		1		1		1		1		1		
	Smolt Productivity	Sm. Capacity	37	37,500	181	185,524	181	185,524	181	185,524	181	185,524	181	185,524	
Hydro	Ocean Surv	Baseline SAR	0.092	0.040	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	0.029	0.008	
	Juv Passage Surv.	Adult Passage	1.00	1.00	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	
	Adjusted Productivity	Adj. Capacity	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	1.46	1,500	
Harvest	Harv - Marine	NORs		HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
	Harv - L. Mainstem	NORs		HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	
	Harv - U. Mainstem	NORs		HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	
	Harv - Terminal	NORs		HORs	0.005	0.100	0.005	0.100	0.005	0.100	0.005	0.100	0.005	0.100	
	Total Exploitation Rate	NORs		HORs	0.083	0.246	0.083	0.246	0.083	0.246	0.083	0.246	0.083	0.246	

Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS	pNOB	pHOS
	Purpose		PNI	pHOS-Realized	None	28%	None	None	None	29%	0.85	18%	Harv	39%
	Type		Cons/Harv/Both	Int/Seg/Step/None	None	None	None	None	None	None	Cons	Int	Harv	Seg
	Broodstock by Source		Local	Imported							30	52,744	567	1,000,189
	Brood Exported (from HOR Surplus)		Export Goal/Realized	Strays		24				27		28		19
Patterns	Destination for HOR Returns		% to Hatchery	% to Nat. Spawn.		100%		100%		100%		100%	90%	10%
	Productivity of Hatchery Fish		Recruits/Spawner	Fitness? [Y / N]		y		y		y	2.5	y	2.5	y
	Update		Habitat	Hydro	Harvest									

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rollout application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Hatchery	Habitat	Marine	L. Mainstem	U. Mainstem	Terminal
Current Habitat					
Current Hydro					
Current Harvest					

Hatchery	Habitat	Marine	L. Mainstem	U. Mainstem	Terminal
Current Habitat					
Current Hydro					
Current Harvest					

Hatchery	Habitat	Marine	L. Mainstem	U. Mainstem	Terminal
Current Habitat					
Current Hydro					
Current Harvest	HSRG				

Hatchery	Habitat	Marine	L. Mainstem	U. Mainstem	Terminal
Current Habitat					
Current Hydro					
Current Harvest	HRT-A				

Hatchery	Habitat	Marine	L. Mainstem	U. Mainstem	Terminal
Current Habitat					
Current Hydro					
Current Harvest	HRT-B				

454

6

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80
Weir Factor ->					
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.50	0.95	0.50
Average Fitness Factor (100 Generations) ->	0.51	0.99	0.51	0.93	0.50
Generations until average fitness is reached ->	27	30	26	44	18
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS ->	Effective pHOS ->	Effective pHOS ->	Effective pHOS ->
"Fitness Floor" ->	0.5	pNOB ->	pNOB ->	pNOB ->	pNOB ->
Calculated Hatchery SAR ->		0.8%	0.8%	0.8%	0.1%
Calculated Natural SAR ->					0.8%
NOR Escapement	Max	Min	Ave	Max	Min
HoS Total Escapement (includes strays)	187	13	51	980	100
	74	13	24		

Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave
178	14	55	178	14	55	1,318	95	388	595	55	174
83	14	27	83	14	27	317	55	104	422	73	138

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A52. AHA output for Yankee Fork spring Chinook. HRT Alternatives A and B represent small integrated conservation programs to increase the overall recruit per spawner ratio for the population as a whole. Two different broodstock strategies are modeled.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

Population Management Intent:

Conservation

Natural Reproduction only - Captive Rearing

Harv&Hatchery Strategy:

Stabilizing

Population Recovery Designation:

Stabilizing

29-Salmon\_Yankee Fork Spring Chinook

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Habitat

Productivity (Adult)

Min NOR Escape

Smolt Productivity

Ad. Capacity

% Kelt

Sm. Capacity

1.45

1

36

600

15,000

1.45

1

179

600

74,210

1.45

1

179

600

74,210

1.45

1

179

600

74,210

1.45

1

179

600

74,210

Hydro

Ocean Surv

Juv Passage Surv.

Adjusted Productivity

Baseline SAR

Adult Passage

Adj. Capacity

0.092

1.00

1.45

0.040

1.00

600

y

0.029

0.34

1.45

0.008

0.82

600

y

0.029

0.34

1.45

0.008

0.82

600

y

0.029

0.34

1.45

0.008

0.82

600

y

0.029

0.34

1.45

0.008

0.82

600

y

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv -Terminal

Total Exploitation Rate

NORs

NORs

NORs

NORs

NORs

HORs

HORs

HORs

HORs

HORs

0.010

0.020

0.050

0.005

0.083

0.010

0.080

0.080

0.100

0.246

0.010

0.020

0.050

0.005

0.083

0.010

0.080

0.080

0.100

0.246

0.010

0.020

0.050

0.005

0.083

0.010

0.080

0.080

0.100

0.246

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

PNI

Cons/Harv/Both

Local

Export Goal/Realized

% to Hatchery

Recruits/Spawner

pHOS-Goal

pHOS-Realized

Int/Seg/Step/None

Imported

Smolt Release

Strays

% to Nat. Spawn.

Fitness? [Y / N]

None

34%

None

24

100%

y

None

None

None

100%

y

None

46%

None

114

100%

y

100%

0.72

Cons

30

2.5

40%

Int

52,744

81

100%

y

50%

0.55

Cons

30

2.5

41%

Int

52,821

83

75%

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

HSRG

Current Habitat

Current Hydro

HRT-A

Current Habitat


Current Hydro


HRT-B

45/7

7

Realized Spawning Composition





Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

80

80%

0.81

0.50

0.50

20

0.5

0.8%

80

80%

0.81

1.00

0.99

30

Effective pHOS ->

pNOB ->

0.8%

80

80%

0.81

0.50

0.50

17

Effective pHOS ->

pNOB ->

0.8%

80

80%

0.81

0.85

0.85

34

Effective pHOS ->

pNOB ->

0.1%

80

80%

0.81

0.69

0.72

32

Effective pHOS ->

pNOB ->

0.1%

NOR Escapement

HoS Total Escapement (includes strays)

Max

Min

Ave

127

11

38

74

13

24

381

39

139

389

38

108

348

60

114

766

51

190

479

83

157

616

49

164

427

74

140

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A53. AHA output for Valley Creek spring Chinook. HRT Alternative A represents a small, integrated conservation program, and HRT Alternative B represents a large segregated program in partial support for achieving LSRCP mitigation goals.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

30-Salmon\_Valley Spring Chinook

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Natural Reproduction only

Contributing

Conservation

Integrated Hatchery

Contributing

Harvest

Segregated Hatchery

Contributing

Habitat

Productivity (Adult)

Ad. Capacity

Min NOR Escape

Smolt Productivity

1.55

800

1

20,000

1.55

800

1

98,946

1.55

800

1

98,946

1.55

800

1

98,946

1.55

800

1

98,946

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

0.092

0.040

y

1.00

1.00

1.55

800

0.029

0.008

y

0.34

0.82

1.55

800

0.029

0.008

y

0.34

0.82

1.55

800

0.029

0.008

y

0.34

0.82

1.55

800

0.029

0.008

y

0.34

0.82

1.55

800

Harvest

Harv - Marine

Harv - L. Mainstem

Harv - U. Mainstem

Harv - Terminal

Total Exploitation Rate

NORs

HORs

NORs

HORs

NORs

HORs

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

0.010

0.010

0.020

0.080

0.050

0.080

0.005

0.100

0.083

0.246

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

pHOS-Goal

PNI

pHOS-Realized

Cons/Harv/Both

Int/Seg/Step/None

Local

Imported

Export Goal/Realized

Strays

% to Hatchery

% to Nat. Spawn.

Recruits/Spawner

Fitness? [Y / N]

pNOB

pHOS

None

28%

None

None

None

None

24

100%

y

pNOB

pHOS

None

28%

None

None

22

100%

y

pNOB

pHOS

100%

0.80

25%

Cons

Int

30

52,744

21

100%

2.5

y

pNOB

pHOS

Harv

Seg

577

1,001,113

13

50%

50%

2.5

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

HSRG

Current Habitat

Current Hydro


HRT-A

Current Habitat

Current Hydro

HRT-B

Realized Spawning Composition



Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

1000

900

800

700

600

500

400

300

200

100

0

0

100

200

300

400

500

600

700

800

900

1000

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

1000

900

800

700

600

500

400

300

200

100

0

0

100

200

300

400

500

600

700

800

900

1000

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

1000

900

800

700

600

500

400

300

200

100

0

0

100

200

300

400

500

600

700

800

900

1000

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

1000

900

800

700

600

500

400

300

200

100

0

0

100

200

300

400

500

600

700

800

900

1000

Hatchery

Habitat

Marine

L. Mainstem

U. Mainstem

Terminal

1000

900

800

700

600

500

400

300

200

100

0

0

100

200

300

400

500

600

700

800

900

1000

Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

80

80%

0.81

0.50

0.50

22

0.5

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

0.28

0.28

0.28

0.28

0.63

0.8%

0.8%

0.8%

0.1%

0.1%

NOR Escapement

HOS Total Escapement (includes strays)

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

155

14

49

657

70

225

146

13

46

851

55

228

1,033

121

285

74

13

24

68

12

22

297

51

97

1,891

313

611

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A54. AHA output for upper Salmon River spring Chinook, including Sawtooth Hatchery, upstream from Redfish Lake Creek. HRT Alternative A represents a large integrated program, and HRT Alternative B represents a two-stage stepping stone program.**

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Conservation		Harvest and Conservation		Harvest and Conservation										
Upper Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Integrated Hatchery		Integrated Hatchery		Stepping Stone - Integrated Hatchery										
31-Salmon_Upper Salmon Mainstem Spring Chinook		Population Recovery Designation:		Primary		Primary		Primary		Primary		Primary		Primary										
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative A		HRT Alternative B												
Habitat	Productivity (Adult)	Ad. Capacity	1.80	2,000	1.80	2,000	1.80	2,000	1.80	2,000	1.80	2,000	1.80	2,000	1.80	2,000								
	Min NOR Escape	% Kelt	1		1		1		1		1		1		1									
	Smolt Productivity	Sm. Capacity	45	50,000	223	247,366	223	247,366	223	247,366	223	247,366	223	247,366	223	247,366								
Hydro	Ocean Surv	Baseline SAR	Vary? (Y/N)	0.092	0.040	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y	0.029	0.008	y						
	Juv Passage Surv.	Adult Passage		1.00	1.00	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82	0.34	0.82						
	Adjusted Productivity	Adj. Capacity		1.80	2,000		1.80	2,000		1.80	2,000		1.80	2,000		1.80	2,000							
Harvest	Harv - Marine		NORs	HORs	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010						
	Harv - L. Mainstem		NORs	HORs	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080	0.020	0.080						
	Harv - U. Mainstem		NORs	HORs	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080	0.050	0.080						
	Harv - Terminal		NORs	HORs	0.080	0.100	0.080	0.100	0.080	0.100	0.080	0.100	0.080	0.100	0.080	0.100	0.080	0.100						
	Total Exploitation Rate		NORs	HORs	0.152	0.246	0.152	0.246	0.152	0.246	0.152	0.246	0.152	0.246	0.152	0.246	0.152	0.246						
Hatchery	Broodstock Composition		pNOB-Goal	pHOS-Goal	pNOB		pHOS		pNOB		pHOS		pNOB		pHOS		pNOB		pHOS					
	Purpose	Type	PNI	pHOS-Realized	Harv		Seg		None		None		67%		27%		50%		40%					
	Broodstock by Source		Cons/Harv/Both	Int/Seg/Step/None	Cons		Int		Cons		Int		Both		Int		Cons		Step					
	Brood Exported (from HOR Surplus)		Local	Imported	532		1,001,011		106		199,393		267		499,207		200		376,320					
	Destination for HOR Returns		Export Goal/Realized	Strays	75%		25%		50%		117		25%		75%		200		164		47			
Patterns	Productivity of Hatchery Fish		% to Hatchery	% to Nat. Spawn.	Recruits/Spawner		Fitness? [Y / N]		75%		25%		50%		75%		50%		50%					
	Update		Habitat	Hydro	Harvest	Hatchery	Current Habitat		Current Hydro		Current Harvest		Current Habitat		Current Hydro		Current Harvest		Current Habitat		Current Hydro		Current Harvest	

Realized Spawning Composition

Relative Hatchery Optimum ->	80	80	80	80	80										
Weir Factor ->															
Relative Reproductive Success (HOS) ->	80%	80%	80%	80%	80%										
Initial Fitness Factor (A) ->	0.81	0.81	0.81	0.81	0.81										
Fitness Factor after 100 generations (B) ->	0.50	1.00	0.85	0.61	0.66										
Average Fitness Factor (100 Generations) ->	0.50	0.99	0.85	0.66	0.70										
Generations until average fitness is reached ->	18	30	34	40	39										
Minimum Hatchery Program (% of BS Goal) ->		Effective pHOS -> 0.38	Effective pHOS -> 0.27	Effective pHOS -> 0.50	Effective pHOS -> 0.40										
"Fitness Floor" -> 0.5		pNOB ->	pNOB ->	pNOB -> 0.33	pNOB -> 0.26										
Calculated Hatchery SAR ->	0.1%			0.1%	0.1%										
Calculated Natural SAR ->	0.8%	0.8%		0.8%	0.8%										
NOR Escapement	Max 1,134	Min 110	Ave 322	Max 1,897	Min 209	Ave 615	Max 2,003	Min 147	Ave 533	Max 1,555	Min 12	Ave 352	Max 1,316	Min 35	Ave 334
HoS Total Escapement (includes strays)															

# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

**Table A55. AHA output for segregated broodstock portions of Sawtooth Hatchery spring Chinook. The HSRG proposed separate integrated and segregated programs. The HRT proposed coupling these programs as a two-stage stepping stone program (HRT Alternative B).**

All H Analyzer (AHA)																																																																																																																																																																																																																			
Apply & Recalculate																																																																																																																																																																																																																			
Population (Ctrl + Page Up/Down to Scroll)																																																																																																																																																																																																																			
Subregion/Subbasin		Species/Race		Population Management Intent:		Harvest and Conservation		Conservation		Harvest		Harvest and Conservation		Harvest																																																																																																																																																																																																					
Upper Salmon R		Spring Chinook		Harv&Hatchery Strategy:		Segregated Hatchery		Natural Reproduction only		Segregated Hatchery		Integrated Hatchery		Stepping Stone Segregated Hatchery																																																																																																																																																																																																					
31A-Salmon_Upper Salmon Mainstem				Population Recovery Designation:		N/A		N/A		N/A		N/A		N/A																																																																																																																																																																																																					
Spring Chinook (Sawtooth Hatchery)																																																																																																																																																																																																																			
		Historic		Current		No Hatchery		HSRG Solution		HRT Alternative A		HRT Alternative B																																																																																																																																																																																																							
Habitat	Productivity (Adult)		Ad. Capacity		0.00		0		0.00		0		0.00		0																																																																																																																																																																																																				
	Min NOR Escape		% Kelt		1				1				1																																																																																																																																																																																																						
	Smolt Productivity		Sm. Capacity		0		0		1		0		1		0																																																																																																																																																																																																				
Hydro	Ocean Surv		Baseline SAR		0.092		0.040		y		0.029		0.008		y																																																																																																																																																																																																				
	Juv Passage Surv.		Adult Passage		1.00		1.00		0.34		0.82		0.34		0.82																																																																																																																																																																																																				
	Adjusted Productivity		Adj. Capacity		0.00		0		0.00		0		0.00		0																																																																																																																																																																																																				
Harvest	Harv - Marine		NORs		HORs		0.010		0.010		0.010		0.010		0.010																																																																																																																																																																																																				
	Harv - L. Mainstem		NORs		HORs		0.020		0.080		0.020		0.080		0.020																																																																																																																																																																																																				
	Harv - U. Mainstem		NORs		HORs		0.050		0.080		0.050		0.080		0.050																																																																																																																																																																																																				
	Harv -Terminal		NORs		HORs		0.080		0.100		0.080		0.100		0.080																																																																																																																																																																																																				
	Total Exploitation Rate		NORs		HORs		0.152		0.246		0.152		0.246		0.152																																																																																																																																																																																																				
Hatchery	Broodstock Composition		pNOB-Goal		pHOS-Goal		pNOB		pHOS		pNOB		pHOS		pNOB																																																																																																																																																																																																				
	Purpose		PNI		pHOS-Realized		None		None		100%		1.00		100%																																																																																																																																																																																																				
	Type		Cons/Harv/Both		Int/Seg/Step/None		Both		Seg		Harv		Seg		Both																																																																																																																																																																																																				
	Broodstock by Source		Local		Imported						638		1,200,461		200																																																																																																																																																																																																				
	Brood Exported (from HOR Surplus)		Export Goal/Realized		Smolt Release						75%		25%		75%																																																																																																																																																																																																				
Patterns	Destination for HOR Returns		% to Hatchery		% to Nat. Spawn.		100%		100%		25%		100%		25%																																																																																																																																																																																																				
	Productivity of Hatchery Fish		Recruits/Spawner		Fitness? [Y / N]		y		y		y		y		y																																																																																																																																																																																																				
	Update		Habitat		Current Habitat						Current Habitat																																																																																																																																																																																																								
			Hydro		Current Hydro						Current Hydro																																																																																																																																																																																																								
			Harvest		Current Harvest						HSRG																																																																																																																																																																																																								
		Hatchery																																																																																																																																																																																																																	
Realized Spawning Composition																																																																																																																																																																																																																			
<table><tr><td>Relative Hatchery Optimum -&gt;</td><td>80</td><td></td><td></td><td>80</td><td></td><td></td><td>80</td><td></td><td></td><td>80</td><td></td><td></td><td>80</td><td></td></tr><tr><td>Weir Factor -&gt;</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Relative Reproductive Success (HOS) -&gt;</td><td>80%</td><td></td><td></td><td>80%</td><td></td><td></td><td>80%</td><td></td><td></td><td>80%</td><td></td><td></td><td>80%</td><td></td></tr><tr><td>Initial Fitness Factor (A) -&gt;</td><td>0.81</td><td></td><td></td><td>0.81</td><td></td><td></td><td>0.81</td><td></td><td></td><td>0.81</td><td></td><td></td><td>0.81</td><td></td></tr><tr><td>Fitness Factor after 100 generations (B) -&gt;</td><td>1.00</td><td></td><td></td><td>1.00</td><td></td><td></td><td>0.19</td><td></td><td></td><td>1.00</td><td></td><td></td><td>0.19</td><td></td></tr><tr><td>Average Fitness Factor (100 Generations) -&gt;</td><td>0.99</td><td></td><td></td><td>0.99</td><td></td><td></td><td>0.24</td><td></td><td></td><td>0.99</td><td></td><td></td><td>0.24</td><td></td></tr><tr><td>Generations until average fitness is reached -&gt;</td><td>30</td><td></td><td></td><td>30</td><td></td><td></td><td>43</td><td></td><td></td><td>30</td><td></td><td></td><td>43</td><td></td></tr><tr><td>Minimum Hatchery Program (% of BS Goal) -&gt;</td><td></td><td></td><td></td><td></td><td></td><td>Effective pHOS -&gt;</td><td></td><td>Effective pHOS -&gt;</td><td></td><td>Effective pHOS -&gt;</td><td></td><td>Effective pHOS -&gt;</td><td></td><td>Effective pHOS -&gt;</td></tr><tr><td>"Fitness Floor" -&gt;</td><td></td><td></td><td></td><td></td><td></td><td>pNOB -&gt;</td><td></td><td>pNOB -&gt;</td><td></td><td>pNOB -&gt;</td><td></td><td>pNOB -&gt;</td><td></td><td>pNOB -&gt;</td></tr><tr><td>Calculated Hatchery SAR -&gt;</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Calculated Natural SAR -&gt;</td><td>0.8%</td><td></td><td></td><td>0.8%</td><td></td><td></td><td>0.1%</td><td></td><td></td><td>0.8%</td><td></td><td></td><td>0.1%</td><td></td></tr><tr><td>NOR Escapement</td><td>Max</td><td>Min</td><td>Ave</td><td>Max</td><td>Min</td><td>Ave</td><td>Max</td><td>Min</td><td>Ave</td><td>Max</td><td>Min</td><td>Ave</td><td>Max</td><td>Min</td><td>Ave</td></tr><tr><td>HoS Total Escapement (includes strays)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>1,024</td><td>177</td><td>335</td><td>0</td><td>0</td><td>0</td><td>321</td><td>55</td><td>105</td></tr></table>															Relative Hatchery Optimum ->	80			80			80			80			80		Weir Factor ->															Relative Reproductive Success (HOS) ->	80%			80%			80%			80%			80%		Initial Fitness Factor (A) ->	0.81			0.81			0.81			0.81			0.81		Fitness Factor after 100 generations (B) ->	1.00			1.00			0.19			1.00			0.19		Average Fitness Factor (100 Generations) ->	0.99			0.99			0.24			0.99			0.24		Generations until average fitness is reached ->	30			30			43			30			43		Minimum Hatchery Program (% of BS Goal) ->						Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->	"Fitness Floor" ->						pNOB ->		pNOB ->		pNOB ->		pNOB ->		pNOB ->	Calculated Hatchery SAR ->															Calculated Natural SAR ->	0.8%			0.8%			0.1%			0.8%			0.1%		NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	HoS Total Escapement (includes strays)	0	0	0	0	0	0	1,024	177	335	0	0	0	321	55	105
Relative Hatchery Optimum ->	80			80			80			80			80																																																																																																																																																																																																						
Weir Factor ->																																																																																																																																																																																																																			
Relative Reproductive Success (HOS) ->	80%			80%			80%			80%			80%																																																																																																																																																																																																						
Initial Fitness Factor (A) ->	0.81			0.81			0.81			0.81			0.81																																																																																																																																																																																																						
Fitness Factor after 100 generations (B) ->	1.00			1.00			0.19			1.00			0.19																																																																																																																																																																																																						
Average Fitness Factor (100 Generations) ->	0.99			0.99			0.24			0.99			0.24																																																																																																																																																																																																						
Generations until average fitness is reached ->	30			30			43			30			43																																																																																																																																																																																																						
Minimum Hatchery Program (% of BS Goal) ->						Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->		Effective pHOS ->																																																																																																																																																																																																					
"Fitness Floor" ->						pNOB ->		pNOB ->		pNOB ->		pNOB ->		pNOB ->																																																																																																																																																																																																					
Calculated Hatchery SAR ->																																																																																																																																																																																																																			
Calculated Natural SAR ->	0.8%			0.8%			0.1%			0.8%			0.1%																																																																																																																																																																																																						
NOR Escapement	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave	Max	Min	Ave																																																																																																																																																																																																				
HoS Total Escapement (includes strays)	0	0	0	0	0	0	1,024	177	335	0	0	0	321	55	105																																																																																																																																																																																																				



# USFWS Columbia Basin Hatchery Review Team: Lower Snake Assessments and Recommendations Reports–March 2011

Table A56. AHA output for Panther Creek spring Chinook.

All H Analyzer (AHA)

Apply & Recalculate

Population (Ctrl + Page Up/Down to Scroll)

Subregion/Subbasin

Species/Race

Upper Salmon R

Spring Chinook

32-Salmon\_Panther Creek Spring Chinook (Extirpated)

Population Management Intent:

Harv&Hatchery Strategy:

Population Recovery Designation:

Conservation-Reintroduction

Habitat Restoration

Stabilizing

Conservation-Reintroduction

Habitat Restoration

Stabilizing

Conservation-Reintroduction

Habitat Restoration

Stabilizing

Conservation-Reintroduction

Habitat Restoration

Stabilizing

Conservation-Reintroduction

Habitat Restoration

Stabilizing

Historic

Current

No Hatchery

HSRG Solution

HRT Alternative A

HRT Alternative B

Habitat

Productivity (Adult)

Ad. Capacity

Min NOR Escape

% Kelt

Smolt Productivity

Sm. Capacity

0.10

1,200

1

30,000

0.10

1,200

1

148,419

0.10

1,200

1

148,419

0.10

1,200

1

148,419

0.10

1,200

1

148,419

Hydro

Ocean Surv

Baseline SAR

Vary? (Y/N)

Juv Passage Surv.

Adult Passage

Adjusted Productivity

Adj. Capacity

0.092

0.040

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

0.029

0.008

y

Harvest

Harv - Marine

NORs

HORs

Harv - L. Mainstem

NORs

HORs

Harv - U. Mainstem

NORs

HORs

Harv -Terminal

NORs

HORs

Total Exploitation Rate

NORs

HORs

0.010

0.010

0.020

0.020

0.050

0.050

0.005

0.005

0.083

0.083

0.010

0.010

0.020

0.020

0.050

0.050

0.005

0.005

0.083

0.083

0.010

0.010

0.020

0.020

0.050

0.050

0.005

0.005

0.083

0.083

Hatchery

Broodstock Composition

Purpose

Type

Broodstock by Source

Brood Exported (from HOR Surplus)

Destination for HOR Returns

Productivity of Hatchery Fish

pNOB-Goal

pHOS-Goal

PNI

pHOS-Realized

Cons/Harv/Both

Int/Seg/Step/None

Local

Imported

Smolt Release

Export Goal/Realized

Strays

% to Hatchery

% to Nat. Spawn.

Recruits/Spawner

Fitness? [Y / N]

pNOB

pHOS

94%

None

None

24

100%

y

pNOB

pHOS

94%

None

None

18

100%

y

pNOB

pHOS

94%

None

None

19

100%

y

pNOB

pHOS

94%

None

None

10

100%

y

Patterns

Update

Patterns shown here are assigned to the population from the AHA or ROL file. Red cells indicate patterns are not assigned to this population/scenario in the rolup application. The shown pattern needs to be added to the application, or a different pattern should be assigned to this population in the ROL/AHA file.

Habitat

Hydro

Harvest

Hatchery

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

Current Harvest

Current Habitat

Current Hydro

HSRG

Current Habitat

Current Hydro


HRT-A


Current Habitat

Current Hydro

HRT-B

Realized Spawning Composition





Relative Hatchery Optimum ->

Weir Factor ->

Relative Reproductive Success (HOS) ->

Initial Fitness Factor (A) ->

Fitness Factor after 100 generations (B) ->

Average Fitness Factor (100 Generations) ->

Generations until average fitness is reached ->

Minimum Hatchery Program (% of BS Goal) ->

"Fitness Floor" ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Effective pHOS ->

pNOB ->

Calculated Hatchery SAR ->

Calculated Natural SAR ->

NOR Escapement

HoS Total Escapement (includes strays)

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

Max

Min

Ave

**Pacific Region Fishery Resources**  
**911 NE 11<sup>th</sup> Avenue**  
**Portland, OR 97232**  
**503/872.2763**  
**E-Mail: [Don\\_Campton@fws.gov](mailto:Don_Campton@fws.gov)**

**U.S. Fish and Wildlife Service**  
**[www.fws.gov](http://www.fws.gov)**

**For Columbia River Basin Hatchery Review Information**  
**[www.fws.gov/pacific/Fisheries/Hatcheryreview/](http://www.fws.gov/pacific/Fisheries/Hatcheryreview/)**

**The mission of the U.S. Fish and Wildlife Service is working with others to conserve, protect and enhance fish, wildlife, plants and their habitats for the continuing benefit of the American people.**

**March 2011**

