U.S. Fish & Wildlife Service

Grass Kelp (*Ulva flexuosa*)

Ecological Risk Screening Summary

U.S. Fish & Wildlife Service, February 2014
Revised, March 2016, November 2017
Web Version, 12/12/2019



Photo: Sarka Martinez. Licensed under Creative Commons BY-NC 4.0.

1 Native Range and Status in the United States

Native Range

From Sturtevant et al. (2012):

"Unknown. This species is widespread. It occurs around the world in inland and/or coastal waters of Latin America and the Caribbean, Asia, Europe, Australia, the U.S.A., Mexico, and various islands in the Pacific Ocean (Kowalski 1975, Trono 1975, Webber 1975, Riouall 1976, Sivalingam 1977, Morton 1978, Vaughan 1978, Bird and McIntosh 1979, Lobel and Ogden 1981, Kies and Dworsky 1982, Rodriguez de Rios and Lobo 1984, Grant and Prasad 1985, Guner et al. 1985, Ho 1987, Hadi et al. 1989, Khotimchenko 1993, Beach et al. 1995, Martinez-Murillo and Aladro-Lubel 1996, Magnusson 1997, Schories et al. 1997, Fernandez et al. 1998,

Woolcott and King 1999, Hodgson and McDermid 2000, Tabudravu et al. 2002, Matik-Skoko et al. 2004, Lourenco et al. 2006, Sahoo et al. 2006).

Ulva flexuosa is considered a cosmopolitan species that has a worldwide distribution (Lougheed and Stevenson 2004)."

From Mareš et al. (2011):

"Most importantly, we were able to identify all studied populations from European inland waters as *U. flexuosa*. This result was strongly supported by both morphological and molecular approaches. We consequently could confirm previous observations exclusively based on morphology by Wærn (1952), Bliding (1963), Marvan et al. (1997), Lederer et al. (1998), Sitkowska (1999), Skácelová (2004), Messyasz and Rybak (2009), and Kaštovský et al. (2010), who reported *U. flexuosa* from the Czech Republic, Poland, and Sweden. For comparison, we sampled some of the important localities noted by these authors, for example, the river Saxån in Sweden, where Bliding collected original material for his monograph, or the classical Czech site in the Soos salt marsh, from which *Ulva* has been reported since the 19th century (Hansgirg 1892, Brabez 1941). Further evidence was gained by the examination of herbarium specimens. Historical collections from the Czech Republic, Hungary, and Romania starting in the 1930s (originally classified as *U. intestinalis*) demonstrably belonged to the same taxon as our recent samples, which was proved by analysis of both morphological and molecular data. Using *rbc*L sequencing, […]"

"Based on our results, *U. flexuosa* should not be considered as alien or invasive in inland European waters. There is now convincing evidence that the species has occurred in this area at least since collection of the type specimen of *U. flexuosa* subsp. *pilifera* (Kützing 1856). Our observations of eight historic specimens (two from Hungary, one from Romania, and five from the Czech Republic) indicate that many previous reports of freshwater *U. intestinalis* should be regarded as *U. flexuosa* in a modern systematic view, as was already proposed by Wærn (1952), Bliding (1963), and Kaštovský et al. (2010). As our data set is limited, we cannot exclude the possible occurrence of other taxa, but we believe that *U. flexuosa* is most likely the prevailing species of Ulva in European freshwaters."

Status in the United States

From Sturtevant et al. (2012):

"*Ulva flexuosa* was recorded for the first time in 2003 in Muskegon Lake, Michigan, where the Muskegon River empties into Lake Michigan. Within the same drainage, it has also been recorded from White Lake and Mona Lake (Lougheed 2005, Lougheed and Stevenson 2004)."

"The Great Lakes range of *U. flexuosa* is currently limited to Muskegon, Mona and White Lake. However, as it is able to form blooms in Muskegon Lake—which has a lower conductivity than Lake Erie—there is potential for it to thrive in other locations in the Great Lakes (Lougheed and Stevenson 2004)."

"Established where recorded."

Means of Introductions in the United States

From Sturtevant et al. (2012):

"*Ulva flexuosa* was very likely transported to the Great Lakes basin in ballast water or on ship hulls (Lougheed and Stevenson 2004)."

Remarks

Various sources list *Ulva flexuosa* (e.g., Guiry in World Register of Marine Species) or *Enteromorpha flexuosa* (e.g., ITIS) as the senior synonym. *Ulva and Enteromorpha* have been treated as separate genera in the past, however, genetic analysis suggests that they are congeneric and as *Ulva* was the first name used in the literature, *Enteromorpha* should be reduced to a synonym status (Shimada et al. 2003). Taxonomy at the Family level and above is the same for both genera. This ERSS follows current scientific understanding that *Ulva flexuosa* is the valid name for this species and *Enteromorpha flexuosa* is a synonym. Information searches were conducted using both names. There are multiple subspecies known, the specific number of which depends on the source being referenced (e.g., 5 subspecies given by Guiry (2015), 2 subspecies given in ITIS, 4 subspecies and 4 varieties in GBIF Secretariat (2019)).

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From Guiry (2015):

"*Ulva flexuosa* Wulfen, 1803"

"Classification: Biota > Plantae (Kingdom) > Viridaeplantae (Subkingdom) > Chlorophyta (Phylum) > Ulvophyceae (Class) > Ulvales (Order) > Ulvaceae (Family) > *Ulva* (Genus) > *Ulva flexuosa* (Species)"

"Status accepted"

"Synonymised names […] *Enteromorpha flexuos*a (Wulfen) J.Agardh, 1883"

Size, Weight, and Age Range

From Sturtevant et al. (2012):

"Size: < 30 cm long and 0.5-1.0 cm in diameter (Sahoo et al. 2003)"

From Hill (2001):

"*E. flexuosa* grows to 20 cm but generally tends to be smaller."

Environment

From Sturtevant et al. (2012):

"*Ulva flexuosa* is primarily a marine species but can tolerate freshwater conditions and is considered euryhaline. It is fast growing and prefers littoral zones with some wave action. In the Lake Michigan watershed, it occurs in mesotrophic to eutrophic aquatic habitats. Its establishment could have been spurred by increased salt concentrations from industrial sources and de-icing salts since the 1850s, as well as a lack of native grazing pressure."

"*Ulva flexuosa* is known from diverse habitats around the world and has been recorded in salinities ranging from 0.5–34‰."

From Hill (2001):

"Optimum reproduction temperature is under 30° C (Mairh, Pandey and Tewari 1986) in waters with a pH of approximately 8.2."

From Mareš et al. (2011):

"Our chemical analyses from several sampling localities indicate that *U. flexuosa* prefers environments with a higher content of dissolved ions (especially chloride and sulfate) than normally present in average freshwater (Wetzel 2001)."

Climate/Range

From Hill (2001):

"Growth of this species in outdoor ponds in India (Mairh, Pandey and Tewari 1986) showed that *E. flexuosa* was able to sustain growth in water temperatures as high as 30° C. Favorable growth was maintained at temperatures ranging from 15.5 - 30° C.

At 33° C, bleaching of a few branches begins to occur, and above this temperature, adverse effects on growth increase as more and more branches become bleached."

Distribution Outside the United States

Native

From Sturtevant et al. (2012):

"Unknown. This species is widespread. It occurs around the world in inland and/or coastal waters of Latin America and the Caribbean, Asia, Europe, Australia, the U.S.A., Mexico, and various islands in the Pacific Ocean (Kowalski 1975, Trono 1975, Webber 1975, Riouall 1976, Sivalingam 1977, Morton 1978, Vaughan 1978, Bird and McIntosh 1979, Lobel and Ogden 1981, Kies and Dworsky 1982, Rodriguez de Rios and Lobo 1984, Grant and Prasad 1985, Guner et al. 1985, Ho 1987, Hadi et al. 1989, Khotimchenko 1993, Beach et al. 1995, Martinez-Murillo and Aladro-Lubel 1996, Magnusson 1997, Schories et al. 1997, Fernandez et al. 1998,

Woolcott and King 1999, Hodgson and McDermid 2000, Tabudravu et al. 2002, Matik-Skoko et al. 2004, Lourenco et al. 2006, Sahoo et al. 2006).

Ulva flexuosa is considered a cosmopolitan species that has a worldwide distribution (Lougheed and Stevenson 2004)."

From Mareš et al. (2011):

"Most importantly, we were able to identify all studied populations from European inland waters as *U. flexuosa*. This result was strongly supported by both morphological and molecular approaches. We consequently could confirm previous observations exclusively based on morphology by Wærn (1952), Bliding (1963), Marvan et al. (1997), Lederer et al. (1998), Sitkowska (1999), Skácelová (2004), Messyasz and Rybak (2009), and Kaštovský et al. (2010), who reported *U. flexuosa* from the Czech Republic, Poland, and Sweden. For comparison, we sampled some of the important localities noted by these authors, for example, the river Saxån in Sweden, where Bliding collected original material for his monograph, or the classical Czech site in the Soos salt marsh, from which *Ulva* has been reported since the 19th century (Hansgirg 1892, Brabez 1941). Further evidence was gained by the examination of herbarium specimens. Historical collections from the Czech Republic, Hungary, and Romania starting in the 1930s (originally classified as *U. intestinalis*) demonstrably belonged to the same taxon as our recent samples, which was proved by analysis of both morphological and molecular data. Using *rbc*L sequencing, […]"

"Based on our results, *U. flexuosa* should not be considered as alien or invasive in inland European waters. There is now convincing evidence that the species has occurred in this area at least since collection of the type specimen of *U. flexuosa* subsp. *pilifera* (Kützing 1856). Our observations of eight historic specimens (two from Hungary, one from Romania, and five from the Czech Republic) indicate that many previous reports of freshwater *U. intestinalis* should be regarded as *U. flexuosa* in a modern systematic view, as was already proposed by Wærn (1952), Bliding (1963), and Kaštovský et al. (2010). As our data set is limited, we cannot exclude the possible occurrence of other taxa, but we believe that *U. flexuosa* is most likely the prevailing species of Ulva in European freshwaters."

Introduced

The native, and therefore also the invasive, range of *Ulva flexuosa* is uncertain (Sturtevant et al. 2012).

From Heesch et al. (2009):

"Based on the criteria used in this study, we have concluded that of the 19 species of *Ulva*, at least four are probable introductions [to New Zealand] (*U. armoricana*, *U. californica*, *U. flexuosa* and *U. lactuca*)."

Kirkendale et al. (2013) lists *Ulva flexuosa* as possibly introduced to Australia based on a lack of genetic diversity in the samples collected from Australia.

Means of Introduction Outside the United States

Information on the specific means of introduction for *Ulva flexuosa* outside the United States could not be found.

From Hill (2001):

"The ecological success of *E. flexuosa* and other pioneering species is in part attributed to the readily available pool of motile unicells that are able to rapidly colonize new areas. The chance for successful settlement of these cells is greatly enhanced because gametes and zoospores of this species remain viable for 10 or more days due to their ability to photosynthesize, often achieving photosynthetic rates approaching those of the adult thalli (Beach et al. 1989)."

Short Description

From Sturtevant et al. (2012):

"*Ulva flexuosa* has a smooth, tubular thallus that is linear and uniform in width, with numerous branches in the lower third. It can have one to several fronds. *Ulva flexuosa* ranges from translucent dark to light green, with chloroplasts taking up part to most of the outer region of its cells. It has a very small rhizoidal region (Woolcott and King 1999). In general, its subspecies are differentiated by their degree of branching: *U. flexuosa* subsp. *flexuosa* tends to be unbranched, while *Ulva flexuosa* subsp. *paradoxa* has branched filaments and is typically more slender (Sahoo et al. 2003). Additional characteristics are summarized below.

Ulva flexuosa subsp. *flexuosa* is yellow-green with tubular fronds, cylindrical stalks, and a blade expanding above the stalk which ends in a rounded tip. It is attached to the substrate by a small, round basal disc (Sahoo et al. 2003). *Ulva flexuosa* subsp. *flexuosa* has sparse branching, one to two pyrenoids (for carbon dioxide fixation), and quadrangular or rectangular cells arranged in approximately regular rows (Mareš et al. 2011).

Ulva flexuosa subsp. *paradoxa* has an intensely branched stalk with rectangular cells arranged in long rows. These rows are somewhat unordered in the central parts of broader thalli, thinning to a single row of cells at the ends of branchlets (Mareš et al. 2011)."

Biology

From Sturtevant et al. (2012):

"To reproduce, dioecious gametophytes produce male and female gametes that have two flagellae, while sporophytes produce zoospores with four flagellae (Koeman and van den Hoek 1984). Zoospores can be stored in the dark for over 50 days and still germinate with a 35% survival rate, suggesting that transport in ballast water is possible (Beach et al. 1995, Kolwalkar et al. 2007). Swarmers may be released with increasing exposure at night and greater desiccation, whereas vegetative growth may typically occur with shorter exposure times (Mairh et al. 1985)."

"In Muskegon Lake [Michigan], the two subspecies co-occur in littoral zone macrophyte beds, although *U. flexuosa* subsp. *paradoxa* has been recorded to comprise 61% of the macroalgae

present by biovolume and *U. flexuosa* subsp. *flexuosa* only 31%. In White Lake [Michigan], *Ulva* sp. has been known to comprise 5% of the macroalgae present, and in Mona Lake [Michigan], 20% (Lougheed and Stevenson 2004)."

"It has been recorded in tidal pools, on mudflats, in sea dikes, in polyhaline man-made waters, in oligohaline ponds, in drainage ditches, on rocks, and in warm calm waters below low tide (Kapraun 1970, Koeman and van den Hoek 1984, Hadi et al. 1989, Sahoo et al. 2003)."

From Hill (2001):

"*E. flexuosa* is highly fecund (Beach et al. 1989), with propagule release via both mitotic spores and meiotic gametes occurring on a daily basis in the lower latitudes. Release of reproductive cells (spores and gametes) into the intertidal zone is driven by tidal and lunar rhythms (Smith 1947, Christie and Evans 1962, Beach et al. 1995). Spores and gametes of this species are photosynthetically competent upon release into the water column, with unicells remaining motile for up to 11 days. Both gametes and zoospores have higher cellular respiration rates when compared to parental tissues, presumably due to their motility. Additionally, the photosynthetic rate for gametes and zoospores of *E. flexuosa* and several other pioneering species is substantially higher than photosynthetic rates observed in the reproductive cells of later successional genera such as the kelps (Phaeophyta). Thus, motility coupled with photosynthetic ability increase the ecological success in settlement and recruitment of this species."

Human Uses

From Hill (2001):

"*E. flexuosa* has some commercial importance due to its antibacterial activity against *Mycobacterium tuberculosis* (Mairh, Pandey and Tewari 1986). This algal species has also been investigated for its potential value as a bioindicator of trace metal pollution in coastal waters (Sivalingam 1978)."

From Sturtevant et al. (2012):

"*Ulva flexuosa* is used as a bioindicator for metal contamination, including lead and iron (Ho 1987, Tabudravu et al. 2002). *Ulva flexuosa* was also found to be an economically efficient species for bioabsorption in industrial settings. It could be used as an eco-friendly alternative for wastewater treatment in dye manufacturing, tannery, textile, and cosmetic industries (Silvasamy et al. 2012)."

Diseases

Information on potential diseases of *Ulva flexuosa* was not found.

Threat to Humans

From Sturtevant et al. (2012):

"*Ulva flexuosa* is known to foul a variety of permanent man-made structures and boats. This can lead to expensive repairs of fouled boat motors. There are even reports of it growing on boats treated with antifouling paint (Kolwalker et al. 2007, Lougheed and Stevenson 2004)."

3 Impacts of Introductions

Ulva flexuosa was unknown within the Great Lakes basin before it was discovered in 2003 (Sturtevant et al. 2012). It is therefore considered to be introduced to the basin.

From Lougheed and Stevenson (2004):

"In addition, *Enteromorpha* [*Ulva flexuosa*] appears to be a poor host for epiphytic algal growth […]. Epiphytic diatom densities were significantly lower […] on both subspecies of *Enteromorpha* [*U. flexuosa*] (3.5 ± 1.5 and 1.9 ± 1.0 cells mm^{-2}) than on *Cladophora* (1,286 ± 319 cells mm^{-2}) and *Oedogonium* (677 ± 280 cells mm^{-2}). Epiphytic diatom growth on *Zygnema* and *Spirogyra* was close to zero, and was not significantly difference from that observed on *Enteromorpha* [*U. flexuosa*]. These results concur with those of Lowe *et al.* (1982) who found that epiphyte density on *Cladophora* (1,320 cell mm^{-2}) was 1,000-fold higher than on the mucilaginous-coated cell walls of the exotic alga *Bangia atropurpurea*. This is a concern in systems where *Enteromorpha* [*U. flexuosa*] is dominant and epiphytic diatoms may be important components of the food web."

Other reports of impacts from blooms of *Ulva flexuosa* were from areas that may comprise the native range of the species and so were not included here.

4 Global Distribution



Figure 1. Known global distribution of *Ulva flexuosa*. Map from GBIF Secretariat (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.

Additional observations from Czech Republic, Finland, Germany, Hungary, Italy, Norway, Poland, Romania, Scotland, Slovakia, and Sweden are given in Mareš et al. (2011).

5 Distribution Within the United States



Figure 2. Known distribution of *Ulva flexuosa* in the United States. Map from BISON (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.



Figure 3. Known distribution of *Ulva flexuosa* in Hawaii. Map from BISON (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.



Figure 4. Additional known distribution of *Ulva flexuosa* (listed under the synonym *Enteromorpha flexuosa*) in the United States. Map from BISON (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.



Figure 5. Additional known distribution of *Ulva flexuosa* (listed under the synonym *Enteromorpha flexuosa*) in Hawaii. Map from BISON (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.



Figure 6. Known distribution of *Ulva flexuosa* (listed under the synonym *Enteromorpha flexuosa*) in the Northern Mariana Islands. Map from BISON (2019). Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis.



Figure 7. Known distribution of *Ulva flexuosa* in the Great Lakes Basin. Map from Sturtevant et al. (2012).

6 Climate Matching

Summary of Climate Matching Analysis

The climate match for *Ulva flexuosa* was high along the Atlantic, Pacific, and Gulf coasts, where the species thought to be native, and in the Great Lakes region, where *U. flexuosa* is introduced. There were patches of high match inland in the west, where the species is not known to be present. The climate match was medium for the rest of the contiguous United States, except in scattered locations throughout the west and in a portion of the Plains States bordering Canada, where the match was low. The Climate 6 score (Sanders et al. 2018; 16 climate variables; Euclidean distance) for the contiguous United States was 0.756, high. (Scores of 0.103 or greater are classified as high.) All States had high individual Climate 6 scores except North Dakota and South Dakota, which had medium scores.



Figure 8. RAMP (Sanders et al. 2018) source map showing weather stations selected as source locations (red) and non-source locations (gray) for *Ulva flexuosa* climate matching. Source locations from Mareš et al. (2011), Sturtevant et al. (2012), BISON (2019) and GBIF Secretariat (2019). Selected source locations are within 100 km of one or more species occurrences, and do not necessarily represent the locations of occurrences themselves. Because the climate matching analysis (section 6) is not valid for marine waters, no marine occurrences were used in the climate matching analysis. Points that look to be in marine locations were verified to be located on islands.



Figure 9. Map of RAMP (Sanders et al. 2018) climate matches for *Ulva flexuosa* in the contiguous United States based on source locations reported by Mareš et al. (2011), Sturtevant et al. (2012), BISON (2019), and GBIF Secretariat (2019). 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
0.000≤X≤0.005	Low
0.005<X<0.103	Medium
≥0.103	High

7 Certainty of Assessment

The certainty of assessment is high. *Ulva flexuosa* is globally distributed. A potentially long history of invasiveness is the reason for the cryptogenic classification of this organism. A record of known introduction and establishment was found with documentation of a negative impact from peer-review literature. There is sufficient information on *U. flexuosa* to satisfy the elements of this assessment.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Ulva flexuosa is a cryptogenic aquatic macroalgae that is globally distributed. *U. flexuosa* primarily a marine and coastal species but it is tolerant to freshwater conditions and a variety of climates. It has uses as a bioindicator of contamination, antibacterial materials, and bioremediation. The history of invasiveness is high. It has been reported in Lake Michigan starting in 2003, with no previous records in the Great Lakes basin, with negative impacts to abundance of epiphytic diatoms. The climate match with the contiguous United States is high. The climate match was high in coastal areas where this species is possibly native and in the Great Lakes basin where it has been introduced. It was also high in parts of the west that do not have any reports of this species' presence. The certainty of assessment is high. The overall risk assessment category is high.

Assessment Elements

- **History of Invasiveness (Sec. 3): High**
- **Climate Match (Sec. 6): High**
- **Certainty of Assessment (Sec. 7): High**
- **Remarks/Important additional information** Referred to in the literature frequently as the synonym *Enteromorpha flexuosa*. There are many subspecies and varieties.
- **Overall Risk Assessment Category: High**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

BISON. 2019. Biodiversity Information Serving Our Nation (BISON). United States Geological Survey. Available: https://bison.usgs.gov. (December 2019).

GBIF Secretariat. 2019. GBIF backbone taxonomy: *Ulva flexuosa* Wulfen. Global Biodiversity Information Facility, Copenhagen. Available: https://www.gbif.org/species/5273507. (December 2019).

Guiry, M. D. 2015. *Ulva flexuosa*. *In* World Register of Marine Species. Available: http://www.marinespecies.org/aphia.php?p=taxdetails&id=234468. (March 2017).

Heesch, S., J. E. S. Broom, K. F. Neill, T. J. Farr, J. L. Dalen, and W. A. Nelson. 2009. *Ulva*, *Umbraulva* and *Gemina*: genetic survey of New Zealand taxa reveals diversity and introduced species. European Journal of Phycology 44(2):143–154.

Hill, K. 2001. *Enteromorpha flexuosa*. Smithsonian Marine Station at Fort Pierce. Available: http://www.sms.si.edu/irlspec/Entero_flexuo.htm. (March 2016).

ITIS (Integrated Taxonomic Information System). 2014. *Enteromorpha flexuosa*. Integrated Taxonomic Information System, Reston, Virginia. Available: https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=6544#null. (February 2014).

Kirkendale, L., G. W. Saunders, and P. Winberg. 2013. A molecular survey of *Ulva* (Chlorophyta) in temperate Australia reveals enhanced levels of cosmopolitanism. Journal of Phycology 49:69–81.

Lougheed, V. L., and R. J. Stevenson. 2004. Exotic marine macroalgae (*Enteromorpha flexuosa*) reaches bloom proportions in a coastal lake of Lake Michigan. Journal of Great Lakes Research 30(4):538–544.

Mareš, J., E. Leskinen, M. Sitkowska, O. Skácelová, and J. Blomster. 2011. True identity of the European freshwater *Ulva* (Chlorophyta, Ulvophyceae) revealed by a combined molecular and morphological approach. Journal of Phycology 47:1177–1192.

Sanders, S., C. Castiglione, and M. Hoff. 2018. Risk assessment mapping program: RAMP, version 3.1. U.S. Fish and Wildlife Service.

Shimada, S., M. Hiraoka, S. Nabata, M. Iima, and M. Masuda. 2003. Molecular phylogenetic analyses of the Japanese *Ulva* and *Enteromorpha* (Ulvales, Ulvophyceae), with special reference to the free-floating *Ulva*. Phycological Research 51:99–108.

Sturtevant, R. A., M. M. McCarthy, and A. Fusaro. 2012. *Ulva* (*Enteromorpha*) *flexuosa* subsp. *flexuosa* and *flexuosa* subsp. *paradoxa*. U.S. Geological Survey, Nonindigenous Aquatic Species Database, Gainesville, Florida, and NOAA Great Lakes Aquatic Nonindigenous Species Information System, Ann Arbor, Michigan. Available: http://nas.er.usgs.gov/queries/GreatLakes/FactSheet.aspx?SpeciesID=2726. (February 2014).

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Beach, et al. 1989. [Source material did not give full citation for this reference.]

Beach, K. S., C. M. Smith, T. Michael, and H. W. Shin. 1995. Photosynthesis in reproductive unicells of *Ulva fasciata* and *Enteromorpha flexuosa*: implications for ecological success. Marine Ecology Progress Series 125(1–3):229–237.

Bliding, C. 1963. A critical survey of European taxa in Ulvales. Part I: Capsosiphon, Percursaria, Blidingia, Enteromorpha. Opera Botanica 8:1–160.

Bird, K. T., and R. P. McIntosh. 1979. Notes on the marine algae of Guatemala. Revista de Biologia Tropical 27(2):163–170.

Brabez, R. 1941. Zur Kenntnis der Algenflora des Franzensbader und Sooser Thermenbereiches. Beihefte zum Botanischen Centralblatt A61:137–236.

Christie, A. O., and L. V. Evans. 1962. Periodicity in the liberation of gametes and zoospores of *Enteromorpha intestinalis* Link. Nature 193:193–194.

Fernandez, O. A., K. J. Murphy, A. L. Cazorla, M. R. Sabbatini, M. Lazzari, J. C. J. Domaniewski, and J. H. Irigoyen. 1998. Interrelationships of fish and channel environmental conditions with aquatic macrophytes in an Argentine irrigation system. Hydrobiologia 380:15–25.

Grant, J. M., and D. Prasad. 1985. Chemical composition of some useful seaweeds of Jamaica, West Indies. Journal of Phycology 21(supplement):15.

Guner, H., V. Aysel, A. Sukatar, and M. Ozturk. 1985. The flora of the Aegean Sea, Turkey. I. Cyanophyta, Chlorophyta, Phaeophyta, and Angiospermae. DOGA Bilim Dergisi Seri A Temel Bilimler 9(2):272–282.

Hadi, R., A. M. Hadi, K. M. Bahram, and A. A. S. Hassan. 1989. The new recorded species of *Enteromorpha* in Baghdad area, Iraq. Bulletin of the Iraq Natural History Museum 8(2):163–172.

Hansgirg, A. 1892. Prodromus der Algenflora von Bo¨hmen. 2. Arch. Naturwiss. Landesdurchforsch. Böhmen 8:1–268.

Ho, Y. B. 1987. Metals in 19 intertidal macroalgae in Hong Kong waters. Marine Pollution Bulletin 18(10):564–566.

Hodgson, L. M., and J. K. McDermid. 2000. Marine plants of Pohnpei and Ant Atoll: Chlorophyta, Phaeophyta, and Magnoliophyta. Micronesica 32(2):289–307.

Kapraun, D. F. 1970. Field and cultural studies of *Ulva* and *Enteromorpha* in the vicinity of Port Aransas, Texas. Contributions in Marine Science 15:205–285.

Kaštovský, J., T. Hauer, J. Mareš, M. Krautová, T. Bešta, J. Komárek, B. Desortová, et al. [Source material did not give full list of authors.] 2010. A review of the alien and expansive species of freshwater cyanobacteria and algae in the Czech Republic. Biological Invasions 12:3599–625.

Khotimchenko, S. V. 1993. Fatty acids of green macrophytic algae from the Sea of Japan. Phytochemistry (Oxford) 32(5):1203–1207.

Kies, L., and N. Dworsky. 1982. Species composition and biomass of floating and dried algal mats in shallow ponds from the vicinity of Pevestorf District of Luechow, Dannenberg, Lower Saxony, West Germany. Mitteilungen aus dem Institut für Allgemeine Botanik, Hamburg 18:71–84.

Koeman, R. P. T., and C. van den Hoek. 1984. The taxonomy of *Enteromorpha* Chlorophyceae in the Netherlands. 3. The sections Flexuosae and Clathratae and an addition to the section Proliferae. Cryptogamie Algologie 5(1):21–61.

Kolwalkar, J. P., S. S. Sawant, and V. K. Dhargalkar. 2007. Fate of *Enteromorpha flexuosa* (Wolfen) J. Agardh and its spores in darkness: implications for ballast water management. Aquatic Botany 86(1):86–88.

Kowalski, W. 1975. Occurrence of the species of a marine green alga *Enteromorpha* in the Szczecin, Pomerania inland waters. Fragmenta Floristica et Geobotanica 21(4):527–536.

Kützing, F. T. 1856. Tabulae phycologicae oder Abbildungen der Tange. Bd. 6. W. Köhne, Nordhausen.

Lederer, F., A. Gardavský, A. Lukešová, K. Kubečková, R. Čápová, E. Lodrová, and K. Trojánková. 1998. Biodiverzita a ekologie sinic a řas minerálních pramenů a rašelinit na ú zemí NPR Soos a v okolí Františkových Lázní a Mariánských Lázní. Pages 68–107 *in* F. Lederer and Z. Chocholoušková, editors. Flóra a vegetace minerálních pramenů a rašeliništNPR Soos. Sborník katedry biologie PeF ZČU. PeF ZČU, Plzeň.

Lobel, P. S., and J. C. Ogden. 1981. Foraging by the herbivorous parrotfish *Sparisoma radians*. Marine Biology (Berlin) 64(2):173–184.

Lougheed, V. L. 2005. An invasive macroalga (*Enteromorpha flexuosa*) in Muskegon Lake. International Association for Great Lakes Research, Conference and Program Abstracts 48:115.

Lourenco, S. O., E. Barbarino, A. Nascimento, J. N. P. Freitas, and G. S. Diniz. 2006. Tissue nitrogen and phosphorus in seaweeds in a tropical eutrophic environment: what a long term study tells us. Journal of Applied Phycology 18(3–5):389–398.

Magnusson, G. 1997. Diurnal measurements of Fv/Fm used to improve productivity estimates in macroalgae. Marine Biology (Berlin) 130(2):203–208.

Mairh, Pandey, and Tewari. 1986. [Source did not give full citation for this reference.]

Mairh, O. P., R. S. Pandey, A. Tewari, R. M. Oza, and H. V. Joshi. 1985. Seasonal phenomena, growth, and swarmer liberation rhythm in *Enteromorpha flexuosa* (Chlorophyceae occurring on the coast of Okha, India). Proceedings of the Indian National Science Academy Part B Biological Sciences 51(3):381–385.

Martinez-Murillo, M. E., and M. A. Aladro-Lubel. 1996. Sessile ciliates epibiontic of algae in the Tamiahua Lagoon, Veracruz, Mexico. Tropical Ecology 37(2):265–272.

Marvan, P., V. Keršner, and J. Komárek. 1997. Invazní sinice a řasy. Pages 13–19 *in* P. Pyšek and K. Prach, editors. Invazní rostliny v české flóře. Zprávy Čes. Bot. Spol. 14.Česká Botanická Společnost, Praha.

Matik-Skoko, S., B. Antolic, and M. Kraljevic. 2004. Ontogenetic and seasonal feeding habits of the annular seabream (*Diplodus annularis* L.) in *Zostera* sp. beds, eastern Adriatic Sea. Journal of Applied Ichthyology 20(5):376–381.

Messyasz, B., and A. Rybak. 2009. The distribution of green algae species from the *Ulva* genera (syn. *Enteromorpha*; Chlorophyta) in Polish inland waters. Oceanological and Hydrobiological Studies 38:121–38.

Morton, O. 1978. Some interesting records of algae from Ireland. Irish Naturalists' Journal 19(7):240–242.

Riouall, R. 1976. Quantitative study of macrophytic algae of loose substratum from Prevost Pond, Herault, France. Naturalia Monspeliensia 26:73–94.

Rodriguez De Rios, N., and M. Lobo. 1984. 2 additions to the marine alga flora of Venezuela. Ernstia 25:1–7.

Sahoo, D., P. Baweja, and N. Kushwah. 2006. Developmental studies in *Porphyra vietnamensis*: a high-temperature resistant species from the Indian coast. Journal of Applied Phycology 18(3–5):279–286.

Sahoo, D., N. Sahu, and D. Sahoo. 2003. A critical survey of seaweed diversity of Chilika Lake, India. Algae - Inchon 18(1):1–12.

Schories, D., A. Albrech, and H. Lotze. 1997. Historical changes and inventory of macroalgae from Koenigshafen Bay in the northern Wadden Sea. Helgolaender Meeresuntersuchungen 51(3):321–341.

Silvasamy, A., S. Nethaji, and L. L. J. Lalli-Nisha. 2012. Equilibrium, kinetic and thermodynamic studies on the biosorption of reactive acid dye on *Enteromorpha flexuosa* and *Gracilariacorticata*. Environment Science Pollution Research 19:1687–1695.

Sitkowska, M. 1999. Dwa nowe stanowiska *Enteromorpha flexuosa* subsp. *pilifera* (Chlorophyta) w Polsce. Fragm. Florist. Geobot. Ser. Pol. 6:301–4.

Sivalingam, P. M. 1977. Marine algal distribution in Penang Island, Malaysia. Bulletin of the Japanese Society of Phycology 25(4):202–209.

Sivalingam, P. M. 1978. Biodeposited trace metals and mineral content studies of some tropical marine algae. Botanica Marina 21:327–330.

Skácelová, O. 2004. Flóra sinic a řas tůní v inundačních pásmech řek. Doctoral dissertation. University of South Bohemia, Českě Budějovice, Czech Republic.

Smith, G. M. 1947. On reproduction of some Pacific coast species of *Ulva*. American Journal of Botany 34:80–87.

Tabudravu, J. N., P. Gangaiya, S. Sotheeswaran, and G. R. South. 2002. *Enteromorpha flexuosa* (Wulfen) J. Agardh (Chlorophyta: Ulvales): evaluation as an indicator of heavy metal contamination in a tropical estuary. Environmental Monitoring and Assessment 75(2):201–213.

Trono, G. C., Jr. 1975. The marine benthic algae of Bulusan and vicinity province of Sorsogon, Philippines. Part 1. Introduction and Chlorophyta. Kalikasan 4(1):23–41.

Vaughan, F. A. 1978. Food habits of the sea bream *Archosargus rhomboidalis* and comparative growth on plant and animal food. Bulletin of Marine Science 28(3):527–536.

Wærn, M. 1952. Rocky-shore algae in the Öregund Archipelago. Acta Phytogeogr. Suecica 30:1–298.

Webber, E. E. 1975. Recent additions to the marine algal flora of Nahant, Massachusetts, USA. Rhodora 77(811):430–433.

Wetzel, R. G. 2001. Limnology. Lake and River Ecosystems. Academic Press, San Diego, California.

Woolcott, G. W., and R. J. King. 1999. *Ulva* and *Enteromorpha* (Ulvales, Ulvophyceae, Chlorophyta) in eastern Australia: comparison of morphological features and analyses of nuclear rDNA sequence data. Australian Systematic Botany 12(5):709–725.