

***Acestrorhynchus pantaneiro* (a fish, no common name)**

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, March 2014

Revised, January 2018

Web Version, 6/12/2018



Photo: © Ictiologia UNEMAT-Cáceres. Licensed under Creative Commons (CC BY-NC).
Available: http://eol.org/data_objects/33467871 (January 2018).

1 Native Range, and Status in the United States

Native Range

From Froese and Pauly (2018):

“South America: Paraguay, Paraná, Uruguay, La Plata, [Brazil, Paraguay, Uruguay, Argentina] and Mamoré River basins [Bolivia, Brazil].”

From Eschmeyer et al. (2018):

“Distribution: Argentina, Bolivia, Brazil, Paraguay and Uruguay.”

Status in the United States

This species has not been reported as introduced or established in the United States.

From Seriously Fish (2018):

“This species is a member of the putative ‘*A. lacustris* group’ of closely-related species within the genus alongside *A. abbreviatus*, *A. lacustris* and *A. pantaneiro* [*sic*]. [...] Several members of the group are occasionally available in the trade but unless collection data is known are virtually impossible to tell apart.”

Means of Introductions in the United States

This species has not been reported as introduced or established in the United States.

Remarks

From Artioli et al. (2013):

“This species differs from the other of the group [*A. lacustris*] in the presence of 31 to 35 longitudinal series of scales around the caudal peduncle, the tip of the pectoral fin not reaching the pelvic fin origin, 25 to 30 longitudinal series of scales between the lateral line and dorsal fin origin, and 15 to 17 series of scales between the lateral line and the anal fin origin (Menezes 1992).”

According to Meurer and Zaniboni-Filho (2012), saicanga and peixe-cachorro are common names of *Acestrorhynchus pantaneiro*.

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2018):

“Kingdom Animalia
Subkingdom Bilateria
Infrakingdom Deuterostomia
Phylum Chordata
Subphylum Vertebrata
Infraphylum Gnathostomata
Superclass Osteichthyes
Class Actinopterygii
Subclass Neopterygii
Infraclass Teleostei
Superorder Ostariophysii

Order Characiformes
Family Acestrorhynchidae
Genus Acestrorhynchus
Species *Acestrorhynchus pantaneiro*

“Taxonomic Status: valid”

Size, Weight, and Age Range

From Froese and Pauly (2018):

“Max length : 35.2 cm TL male/unsexed; [Zaniboni Filho et al. 2004]; max. published weight: 396.00 g [Zaniboni Filho et al. 2004]”

Environment

From Froese and Pauly (2018):

“Freshwater; benthopelagic.”

Climate/Range

From Froese and Pauly (2018):

“Tropical”

Distribution Outside the United States

Native

From Froese and Pauly (2018):

“South America: Paraguay, Paraná, Uruguay, La Plata, and Mamoré River basins.”

From Eschmeyer et al. (2018):

“Distribution: Argentina, Bolivia, Brazil, Paraguay and Uruguay.”

Introduced

From Neuhaus et al. (2016):

“*Acestrorhynchus pantaneiro* is an invasive species in the Patos Lagoon system in South Brazil.”

Means of Introduction Outside the United States

From Artiolo et al. (2013):

“Regarding the introduction of *A. pantaneiro* into the Tramandaí River system [Brazil], we discarded the hypothesis of escapes through drainage from culture tanks, as suggested by Saccol-Pereira et al. (2006) for the Patos system.”

“The first record of the *A. pantaneiro* in the Tramandaí River system at Fortaleza lagoon suggests that invasion started by the south, perhaps facilitated by the short distance between the northeast portion at Patos Lagoon basin and Tramandaí system [...].”

Short Description

From Artiolo et al. (2013):

“The specimens have a round black humeral spot and a distinct spot at the base of the caudal fin, the general pattern in the *A. lacustris* species group, which includes *A. pantaneiro*.”

Biology

From Froese and Pauly (2018):

“Feeds on fish [Zaniboni Filho et al. 2004].”

From Neuhaus et al. (2016):

“This species apparently develops better in lentic or semi-lentic conditions. The success in lentic environments may be associated with the fact that the species is an opportunistic piscivore that tends to feed on the most abundant species and reproduces by multiple spawning without parental care (Meurer and Zaniboni-Filho 2012).

From Meurer and Zaniboni-Filho (2012):

“*Acestrorhynchus pantaneiro* in the present study demonstrated opportunistic reproductive and feeding behavior (multiple spawning, ability to prey different fish species, and tolerate distinct hydrological conditions). Additionally, the species presents a long reproductive period and consumed the most abundant fish species.”

Human Uses

From Froese and Pauly (2018):

“Fisheries: commercial”

From Seriously Fish (2018):

“This species is a member of the putative ‘*A. lacustris* group’ of closely-related species within the genus alongside *A. abbreviatus*, *A. lacustris* and *A. pantaneiro* [*sic*]. [...] Several members of the group are occasionally available in the trade but unless collection data is known are virtually impossible to tell apart.”

Diseases

From Chemes and Takemoto (2011):

“Family Bucephalidae (Poche, 1907)

Rhipidocotyle santanaensis (Lunaschi, 2004)

Host: *Acestrorhynchus pantaneiro* (Menezes, 1992)

Site of infection: caeca

Location: Paraná River (Santa Ana City, Corrientes Province)

Refs.: Lunaschi (2004)”

No OIE reportable diseases have been documented for this species.

Threat to Humans

From Froese and Pauly (2017):

“Harmless”

3 Impacts of Introductions

From Neuhaus et al. (2016):

“Introductions and invasions of piscivorous predator species can impact not only the species richness of native fish community but may also cause changes over all trophic levels of the ecosystem by eliminating fish with particular diet traits as, for example, zooplanktivorous species (Pinto-Coelho et al., 2008). The results of the present study show that *A. pantaneiro* competes principally with *O. robustus* for food resources. The congeners *O. robustus* and *O. jenynsii*, which developed simpatrically [*sic*], partitioned their food niches during the process of co-evolution, to decrease the overlap and, consequently, competition for the same food resources. Niche theory suggests that the high food niche overlap of both species may cause future niche partitioning of *A. pantaneiro* and *O. robustus* or, as consequence of competition, a decrease in abundance of one of them. Due to the fact that *A. pantaneiro* is dispersing in all directions along the tributaries of the Patos Lagoon and the Tramandaí River basin, further studies are recommended to understand the interaction of this species with other native species of these river systems.”

From Meurer and Zaniboni-Filho (2012):

“[...] *A. pantaneiro* is well adapted to colonize impounded areas. [...] *A. pantaneiro* changed from 0.06% of the total fish biomass per unit of effort before the construction of Itá Dam to 7.80% after the formation of the reservoir (Zaniboni-Filho et al., 2008). The increase in 130 times in biomass shows its success in the colonization of this reservoir, which can be attributed to the reproductive and food strategies of the species. The great food availability created by the proliferation of small-sized species, a common characteristic of newly formed reservoirs, probably propitiated an abundance of food for *A. pantaneiro*.”

4 Global Distribution



Figure 1. Map of known global distribution of *Acestrorhynchus pantaneiro*, reported from South America. Map from GBIF Secretariat (2017).

5 Distribution Within the United States

This species has not been reported as introduced or established in the United States.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match (Sanders et al. 2014; 16 climate variables; Euclidean Distance) was highest in Florida, high in the coastal Southeast and along the coasts of the Gulf Coast states. Medium match occurred in the remainder of the Southeast. Low matches occurred in the northern Great Lakes states, New England, and the northern and western United States. Climate 6 match indicated that the contiguous U.S. has a medium climate match. The range for a medium climate match is from 0.005 to 0.103; Climate 6 match of *Acestrorhynchus pantaneiro* is 0.077.




Figure 2. RAMP (Sanders et al. 2014) source map showing weather stations in South America selected as source locations (red; Brazil, Bolivia, Paraguay, Uruguay, Argentina) and non-source locations (gray) for *Acestrorhynchus pantaneiro* climate matching. Source locations from GBIF Secretariat (2017).




Figure 3. Map of RAMP (Sanders et al. 2014) climate matches for *Acestorhynchus pantaneiro* in the contiguous United States based on source locations reported by GBIF Secretariat (2017). 0= Lowest match, 10=Highest match.

The “High”, “Medium”, and “Low” climate match categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 < X < 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

Information on the biology and distribution of *A. pantaneiro* is available. Where introduced, *A. pantaneiro* may compete with native piscivorous fishes for food as shown through one study of dietary overlap, but it is unclear if that is enough to cause harm to the native species at a population level. Certainty of this assessment is low.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Acestrorhynchus pantaneiro is a freshwater fish species native to South America. This species is fished commercially and may appear in the aquarium trade. *A. pantaneiro* feeds on other fish and reproduces by multiple spawning. In the Patos Lagoon in Brazil, introduced *A. pantaneiro* may compete with native piscivorous fishes for food, as shown in one dietary study; however, there is no evidence that *A. pantaneiro* has negatively impacted populations of native piscivores in that system. More research is needed to fully understand the impacts from introductions of this species; absence of this research makes the certainty of this assessment low. Climate match with the contiguous United States is medium. Overall risk posed by this species is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): None Documented**
- **Climate Match (Sec.6): Medium**
- **Certainty of Assessment (Sec. 7): Low**
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

Artiolo, L. G. S, P. C. Neto, M. Renata, and C. B. Fialho. 2013. First record of the non-native species *Acestrorhynchus pantaneiro* Menezes, 1992 (Characiformes, Acestrorhynchidae) in the Tramandaí River System, Rio Grande do Sul, Brazil. *Pan-American Journal of Aquatic Sciences* 8(1):51-54.

Chemes, S. B., and R. M. Takemoto. 2011. Diversity of parasites from Middle Paraná system freshwater fishes, Argentina. *International Journal of Biodiversity and Conservation* 3(7):249-266.

Encyclopedia of Life (EOL). 2017. *Acestrorhynchus pantaneiro*. Available: http://eol.org/data_objects/30901643. Photo license available: <http://creativecommons.org/licenses/by-nc/3.0/legalcode>. (January 2018).

Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2018. Catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (January 2018).

Froese, R. and D. Pauly, editors. 2018. *Acestrorhynchus pantaneiro* Menezes, 1992. FishBase. Available: <https://www.fishbase.de/summary/Acestrorhynchus-pantaneiro.html>. (June 2018).

- GBIF Secretariat. 2017. GBIF backbone taxonomy: *Acestrorhynchus pantaneiro* (Menezes 1992). Global Biodiversity Information Facility, Copenhagen. Available: <http://www.gbif.org/species/2355576> (January 2018).
- ITIS (Integrated Taxonomic Information System). 2018. *Acestrorhynchus pantaneiro* (Menezes 1992). Available: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=640370 (January 2018).
- Meurer, S. and E. Zaniboni-Filho. 2012. Reproductive and feeding biology of *Acestrorhynchus pantaneiro* Menezes, 1992 (Osteichthyes: Acestrorhynchidae) in areas under the influence of dams in the upper Uruguay River, Brazil. *Neotropical Ichthyology* 10(1):159-166.
- Neuhaus, E. B., D. A. Antonetti, and U. H. Schulz. 2016. The invasive fish *Acestrorhynchus pantaneiro* Menezes, 1992 in Southern Brazil: Occurrence and food niche overlap with two native species. *Journal of Applied Ichthyology* 32:1107-1112
- Sanders, S., C. Castiglione, and M. H. Hoff. 2014. Risk Assessment Mapping Program: RAMP. U.S. Fish and Wildlife Service.
- Seriously Fish. 2018. *Acestrorhynchus pantaneiro* Menezes, 1992. Available: <https://www.seriouslyfish.com/species/acestrorhynchus-pantaneiro/>. (June 2018).

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

- Lunaschi, L. I. 2004. Two news species of bucephalids (Digenea: Bucephalidae) parasitic in freshwater fishes of Argentina. *Parasitology International* 53:229-234.
- Pinto-Coelho, R. M., J. F. Bezerra-Neto, F. Miranda, T. G. Mota, R. Resck, A. M. Santos, and F. A. R. Barbosa. 2008. The inverted trophic cascade in tropical plankton communities: Impacts of exotic fish in the Middle Rio Doce lake district, Minas Gerais, Brazil. *Brazilian Journal of Biology* 68:1025-1037.
- Saccol-Pereira, A., P. C. C. Milani, and C. B. Fialho. 2006. Primeiro registro de *Acestrorhynchus pantaneiro* Menezes, 1992 (Characiformes, Acestrorhynchidae) no sistema da laguna dos Patos, Rio Grande do Sul, Brasil. *Biota Neotropica* 6(3).
- Zaniboni Filho, E., S. Meurer, O. A. Shibatta, and A. P. de Oliverira Nuñer. 2004. Catálogo ilustrado de peixes do alto Rio Uruguai. Editora da UFSC: Tractebel Energia, Florianopolis, Brazil.

Zaniboni-Filho, E., A. P. O. Nuñez, D. A. Reynalte-Tataje, S. Hermes-Silva, and S. Meurer. 2008. Alterações espaciais e temporais da estrutura da comunidade de peixes em decorrência da implantação do reservatório de Itá (alto rio Uruguai). Pages 21-48 *in* E. Zaniboni-Filho, and A. P. O. Nuñez, editors. Reservatório de Itá. Estudos ambientais, desenvolvimento de tecnologias de cultivo e conservação da ictiofauna. Editora da UFSC, Florianópolis, Brazil.