

Hydrilla (*Hydrilla verticillata*) – monoecious biotype

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, Web Version – 9/13/2017



Photo: USGS. Licensed under Public Domain.

1 Native Range and Status in the United States

Native Range

From Jacono et al. (2011):

“Korea appears the likely origin for the monoecious type (Madeira et al. 1997).”

From CABI (2015):

“It is thought to be native but is relatively rare in Europe (Preston and Croft, 1997), sufficiently so that it is protected in Lithuania (Balevicius, 1998). It occurs in certain areas in Poland and Belarus, and has been found in solitary lakes in Ireland (Preston and Croft, 1997).”

Status in the United States

From Jacono et al. (2011):

“Populations north of South Carolina are essentially monoecious (having both male and female flowers on the same plant) (Mадiera et al. 2000). They set some fertile seed, and depend on tubers for overwintering.”

“The most abundant aquatic plant in Florida public waters (Schardt 1994), seventy percent of Florida's freshwater drainage basins contain waterbodies infested with hydrilla (FLDEP 1988-94 and other data). Less common throughout Alabama, although widespread in the Mobile Delta and northern portion of the Mobile Bay (Zolcynski 1997); common at Coffeerville, Aliceville and Oak Mountain reservoirs of central Alabama (D. Powell, Alabama Power Company, pers. comm. 1996), and well established at Guntersville, and other northern impoundments on the Tennessee River (Bates and Smith 1994). Extending along the Tombigbee River from Aliceville Reservoir, AL into eastern Mississippi (E. Dibble, Mississippi State University [sic], pers. com. 1998). Also, reestablished [sic] in the Tennessee-Tombigbee Waterway, at Beckner boat ramp and in the old bendway, (D. Franks, Mississippi Dept. Wildlife, Fisheries and Parks, pers. comm. 2000). Reported in southeastern Tennessee from the middle Tennessee River drainage, where herbicide and natural decline in Lake Chickamauga have precluded recent collection; yet, still found downstream, as scattered stands, in Nickajack Reservoir (Tennessee Valley Authority 1990, D. Webb, Tennessee Valley Authority, pers. com. 1997). Know primarily in the southwestern drainages of Georgia; problematic at Lake Seminole for over a decade (Eubanks 1996). Occurring, otherwise, at a few smaller waterbodies [sic] in Georgia's upper Ocmulgee drainage and at the Strom Thurmond Reservoir on the upper Savannah River (L. Ager, Georgia Department of Natural Resources, pers. com. 1998). Expanding over 32,000 acres in Lakes Marion and Moultrie, South Carolina following the 1989 hurricane, Hugo (Roach et al 1993). Problematic at six additional reservoirs in South Carolina (S. de Kozlowski, South Carolina Department of Natural Resources, pers. comm. 1997), within the Seneca, Saluda, Wateree, Four Hole Swamp, and Cooper drainages. Covering approximately 5,800 acres in North Carolina, primarily at eastern sites, including reservoirs in the upper Neuse River drainage, yet reported from as far west as Buncombe county, in the western mountains (NCDWR 1996). Established in the Potomac, Rappahannock, Anna, Chickahominy, and Appomattox Rivers of Virginia's coastal plain; extending into piedmont Virginia, at several reservoirs (E. Steinkoenig and J. Kauffman, Virginia Game and Inland Fisheries; J. Tate, Virginia Department of Agriculture and Consumer Services; pers. comm. 1998). Although reduced in abundance, continuing to dominate beds of submerged vegetation in the tidal freshwater reaches of the Potomac River on the Virginia/Maryland border (Orth et al. 1996). Established in Maryland at marsh creeks and rivers on the western and northeastern shores of the Chesapeake Bay, including the Pautuxent River, where it has become the most abundant plant species (M. Naylor, Maryland Dept. of Natural Resources, pers. comm. 2000; Orth et al. 1996; Posey et al. 1993). Also common at reservoirs in mid-Maryland, especially those draining the Patuxent River, and in the Ohio drainage of far western Maryland, at Deep Creek Lake (M. Naylor, Maryland Department of Natural Resources, pers. comm. 2000). Discovered recently at three sites in Pennsylvania in scattered stands in the Schuylkill River, downtown Philadelphia (P. Madeira, USDA/ARS, Aquatic Weed Control Research, pers. comm. 1996); at Highland Lake, a 28 acre impoundment on Southwick Creek (Colangelo 1998); and at Lake Nockamixon, where plants are abundant in a drowned portion of

Haycock Creek [E. Zacharias and A. Schuyler 8345 (PH), 1998]. Common in Delaware ponds (AREC 1995), especially in southern Sussex County where it has spread to nearly 1200 acres in 14 ponds and portions of the Nanticoke River (C. Martin, Delaware Dept. of Natural Resources, 1999). Present since the late 1980s at two ponds in southeastern Connecticut (Balcom 1997), where plants were originally misidentified due to the absence of midrib teeth (Les 1996). In Massachusetts, found in 2001 in a Cape Cod pond (B. Hellquist pers. comm. 2001) and new to a Pembroke-area pond as of 2008. Found in Maine in 2002 in the Saco drainage at Pickerel Pond, York County, where plants are established and abundant throughout the pond (R. Bouchard, ME DEP, pers. comm. 2003). First documented in 2003 for the state of New Jersey in the Lower Delaware drainage at Lake Mallard in the Pinelands National Reserve (Sullivan s.n. DOV, FLAS). This small lake is one of several connected ponds. As of April 2003, hydrilla had not been detected in any of the other linked ponds (G. Sullivan, Allied Biological, Inc., pers. comm. 2003). Found in Indiana's Lake Manitou in 2006, prompting closure to public boating access and chemical treatment (D. Keller, Indiana DNR, pers. comm. 2006). As of June of 2008, the treatments have significantly reduce hydrilla growth and tuber production, prompting a limited re-opening of one boat ramp. Found in a privately-owned artificial pond in Marinette County, Wisconsin in 2007. State and county agencies are working to identify the possible introduction pathway and eradication plan.

Most recently identified (2008) in three lakes in Suffolk and Orange Counties, New York where authorities plan treatments including herbicides and grass carp (S. Kishbaugh, NYSDEC Division of Water, pers. comm. 2008) and in three counties in eastern Kentucky.

West of Mississippi River

Appearing more frequently at lakes and reservoirs in Louisiana, especially along Highway 1 as it stretches diagonally across the state, in canals and bayous of the Atchafalaya Basin, and through the coastal marsh region south of Interstate 10 (C. Biggar, Louisiana Department of Wildlife and Fisheries, pers. comm. 1998). Also present at Lake Bruin (indicated by the dot in the eastern Tensas drainage) which is disjunct from other infested regions in Louisiana. Known from over 80 Texas reservoirs, residing in drainages that extend from north-central to eastern Texas, and south to the Rio Grande, at the Mexico border (Helton and Hartman 1997; E. Reyes, U.S. Fish and Wildlife Service, pers. comm. 1998). Eradicated in the mid 1980s from two ponds in Phoenix, Arizona, where no new infestations have been found (E. Hall, Arizona Dept. of Agriculture, pers. comm. 1996). Eradicated from private ponds and several reservoirs in nine California counties; presently occurring at less than 50 sites in Imperial, Tulare, Madera, Mariposa, Calaveras, Yuba, Lake and Shasta Counties (PPDC 1997). Recent infestations in Clear Lake threaten the highly productive Sacramento/San Joaquin River Delta area (Anderson 1996). In 1995, it was discovered in two interconnected Washington lakes, east of Puget Sound, where early biomass measurements reflect high growth potential in the northwestern climate (K. Hamel, Washington State University, pers. comm. 1995). Following successive years of herbicide treatment, only two isolated plants were found in one of these lakes in 2006 and none in 2007. New to the state of Arkansas, established in Lake Ouachita and DeGray Lake, in the Ouachita Headwaters and the Upper Ouachita drainages, respectively. First identified in Lake Ouachita in 1999 during creel surveys. Lake Ouachita is a 42,000 acre lake, with an approximate hydrilla infestation size of 4,000 acres. DeGray Lake is nearly half the size of Lake Ouachita and has 3-4 confirmed acres

of hydrilla infested waters. Hydrilla has been verified growing in waters of 7.3 m in depth with expected growth to be in the 9.1 - 10.7 m level in the near future due to light penetration [sic] to 12.2 m in depth and to the sedimentary composition of the lake. Lake Ouachita personnel are working on a project to initiate biological control efforts in the lake (R. Stokes, USACE, Manager, Lake Ouachita, pers. comm. 2003). Hydrilla was found in late 2007, growing with *Pistia stratiotes* in a geothermal area near Bruneau, Idaho, upstream of the CJ Strike Reservoir and the Snake River. Surveys have found it seven miles downstream from the hot spring area. It was also found in June 2008 in a suburban ditch in West Boise; this ditch remains warm through winter thanks to flow from geothermal wells and drains to the Boise River. Both populations have been determined to be dioecious, hopefully suggesting infestations are limited to geothermally influenced warm-water areas (T. Woolf, Idaho Dept of Agriculture, pers. comm. 2008). Most recently found in a park pond in Kansas, where eradication efforts will begin in 2009 (J. Goeckler, Kansas Department of Wildlife & Parks, pers. comm. 2008).”

From USDA (2017):

Country/State	Common Name	Classification
United States	hydrilla	Noxious weed
Alabama	hydrilla	Class A noxious weed
Arizona	hydrilla, Florida-elodea	Prohibited noxious weed
California	hydrilla	A list (noxious weeds)
	hydrilla	Noxious aquatic weed
	hydrilla	Quarantine
Colorado	hydrilla	A list (noxious weeds)
Connecticut	hydrilla	Invasive, banned
Florida	hydrilla, Florida-elodea	Prohibited aquatic plant, Class 1
Maine	hydrilla	Invasive aquatic plant
Massachusetts	hydrilla	Prohibited
Nevada	hydrilla	Noxious weed
New Mexico	hydrilla	Class A noxious weed
North Carolina	waterthyme	Class A noxious weed
Oregon	hydrilla	“A” designated weed
	hydrilla	Quarantine
South Carolina	hydrilla	Invasive aquatic plant
	hydrilla	Plant pest
Texas	hydrilla	Noxious plant
Vermont	hydrilla	Class A noxious weed
Washington	hydrilla	Class A noxious weed
	hydrilla	Wetland and aquatic weed quarantine

Means of Introductions in the United States

From Jacono et al. (2011):

“The monoecious strain was a separate introduction, first found decades later in the Potomac Basin (Environmental Laboratory 1985).”

“Hydrilla is mainly introduced to new waters as castaway fragments on recreational boats, their motors and trailers and in live wells. Stem pieces root in the substrate and develop into new colonies, commonly beginning near boat ramps. Once established, boat traffic continues to shatter and spread hydrilla throughout the waterbody. Both types propagate primarily by stem fragmentation, although axillary buds (turions) and subterranean tubers are also important. Tubers are resistant to most control techniques (Schardt 1994) and may be viable as a source of reinfestation [sic] for years (Van and Steward 1990).

Hydrilla may be unknowingly transplanted into private ponds as a contaminant on watergarden plants. It is often found spreading after extensive 2,4-D use in public waters once heavily populated with Eurasian water-milfoil (*Myriophyllum spicatum*) (Bates and Smith 1994).”

From NANSF (2013):

“Monoecious hydrilla may have been first documented within the US in Delaware, and then in Washington, DC and the Potomac River in 1980 (Haller 1982, Steward et al. 1984). Steward et al. (1984) suggested that it was possibly introduced during caging and transplanting experiments where hydrilla, mistakenly identified as *Elodea canadensis*, was transplanted from Kenilworth Gardens in Washington, DC. Lilypons Water Gardens, located in Adamstown, Maryland, may have been the source of the Kenilworth Gardens infestation, as they were a popular commercial supplier of aquatic ornamental plants in the area, and Haller (1982) reported seeing a hydrilla-like plant with tubers there during a 1980 visit. Monoecious hydrilla was first documented in North Carolina in 1980 after reports of dense, weedy vegetation in Umstead Lake in the mid to late 1970’s (Figure 2 [in source material]). The Delaware and North Carolina invasions seem to be relatively synchronous and it is possible that early monoecious hydrilla spread in the US was facilitated by contaminated aquatic plant (esp. water lily) shipments from Lilypons (Madeira et al. 2000) or other sources.”

“The monoecious biotype spread and became the dominant hydrilla biotype found in the Mid-Atlantic States (Kay 1992; Langeland 1996). In 2000, Madeira et al. showed that monoecious hydrilla was found in drainages of the Atlantic Basin from central Georgia up to Pennsylvania and Connecticut. It was not found in Gulf Basin drainages, but was sporadically located in the Pacific basin, in California and Washington (Madeira et al. 2000). Monoecious hydrilla is also found in the Interior Basin in Maryland and North Carolina.”

Remarks

From CABI (2015):

“*H. verticillata* is a submerged plant that has rapid growth and a highly effective survival strategy that makes it one of the most troublesome aquatic weeds of water bodies in the world. It forms dense masses, outcompeting native plants and interfering with many uses of waterways. It can be spread by water flow, waterfowl and recreational activities and is sold as an aquarium plant. In the USA it has been listed as Federal Noxious Weed since 1976; its import is prohibited in Western Australia and Tasmania, and it is on the EPPO alert list.”

From NANSP (2013):

“Langeland (1996) suggested that monoecious hydrilla could spread as far north as southern Canada, based on its range in Europe. Les et al. (1997) compiled the northernmost hydrilla distributions worldwide and Balciunas and Chen (1993) provided a comparison of January temperatures in North America to those in Asia where hydrilla has been documented. Based on reported worldwide hydrilla distribution and climate patterns, there are vast areas in North America at risk of invasion by hydrilla (Figure 3 [in source material]).”

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2015):

“Kingdom Plantae
Subkingdom Viridiplantae
Infrakingdom Streptophyta
Superdivision Embryophyta
Division Tracheophyta
Subdivision Spermatophytina
Class Magnoliopsida
Superorder Lilianae
Order Alismatales
Family Hydrocharitaceae
Genus *Hydrilla*
Species *Hydrilla verticillata* (L. f.)Royle”

“Taxonomic Status: Current Standing: accepted”

Size, Weight, and Age Range

From Jacono et al. (2011):

“Stems grow up to 9 m in length; leaves are 6-20 mm long and 2-4 mm wide.”

Environment

From GISD (2011):

“*H. verticillata* is found in freshwater but can tolerate salinities of up to 7% salinity of seawater. It has been found in springs, lakes, marshes, ditches, rivers, and tidal zones. It can grow in relatively low light and CO₂ conditions.”

From CABI (2015):

“In the tropics, *H. verticillata* is described as tolerant of a wide variety of water conditions, from acidic and oligotrophic to eutrophic or brackish; it thrives on many kinds of pollution and tolerates a great deal of disturbance (Cook and Lüönd, 1982), although increasing salinity appears to limit its dispersal (Rout et al., 1998; Mataraza et al., 1999; Rout and Shaw, 2001). Due to its tolerance of low light conditions (White et al., 1996), it is capable of growing in water up to 7 m deep (Yeo et al., 1984). [...] In temperate regions, it grows in alkaline, moderately calcareous, mesotrophic or slightly eutrophic waters (Preston and Croft, 1997), richer in SO₄, but generally poorer in Na, K and Cl than those of *Elodea canadensis* (Klosowski and Tomaszewicz, 1997). It also appears to occur more often as scattered stands within more diverse aquatic plant communities (Klosowski and Tomaszewicz, 1997; Balevicius, 1998).”

Climate/Range

From GISD (2011):

“*H. verticillata* prefers temperatures between 20 and 27 degrees Celsius.”

From NANSF (2013):

“Optimal growth and survival for the dioecious type is found in warmer climates, while the monoecious form is better suited for more temperate climates with lower temperatures and shorter growing seasons (Ames et al. 1986; Van 1989; Madeira et al. 2000; Netherland 1997; Steward et al. 1987). Dioecious hydrilla typically thrives all year in the warm waters of the southern US, while monoecious hydrilla dies back completely in the winter and acts as a herbaceous perennial (Harlan et al. 1985).”

Distribution Outside the United States

Native

From Jacono et al. (2011):

“Korea appears the likely origin for the monoecious type (Madeira et al. 1997).”

From CABI (2015):

“It is thought to be native but is relatively rare in Europe (Preston and Croft, 1997), sufficiently so that it is protected in Lithuania (Balevicius, 1998). It occurs in certain areas in Poland and Belarus, and has been found in solitary lakes in Ireland (Preston and Croft, 1997).”

Introduced

From GISD (2011):

“The dioecious and the monoecious plant are now found on every continent except Antarctica.”

From Zhuang (2013):

“Introduced:

Austria; Germany; Hungary; Italy; Spain (Canary Is., Spain (mainland)); United States

Present - origin uncertain:

Latvia; Poland”

From CABI (2015):

“On the African continent it occurs around Lake Victoria and Lake Tanganyika in the Rift Valley of East Africa, while it has also been reported from Mozambique and a few isolated places in West Africa and, in 2006, from South Africa.”

Means of Introduction Outside the United States

From GISD (2011):

“Floating vegetation/debris: Plant fragments dispersed by river flow.

Ignorant possession: Shipments of water lilies have been found contaminated with Hydrilla.

Pet/aquarium trade: Sold as an aquarium plant.”

Short Description

From Jacono et al. (2011):

“Submersed perennial herb. Rooted, with long stems that branch at the surface where growth becomes horizontal and dense mats form. Small, pointed leaves are arranged in whorls of 4 to 8. Leaves have serrated margins and one or more sharp teeth under the midrib (see Godfrey and Wooten 1979). Development of these features may vary with location, age, and water quality (Kay 1992).”

From CABI (2015):

“*H. verticillata* is a submerged, monoecious or dioecious perennial. Its stems are branched, about 1 mm thick and up to 3 m long; the internodes are 3 to 50 mm long. The sessile leaves are formed in whorls at the nodes; there are 3-8, sometimes up to 12 leaves in a whorl. The leaves are 7-40 mm long, linear to lanceolate, with a conspicuous midrib. They have sharply toothed margins and spines on the vein on the lower side of the leaves; a few teeth may also be formed on this vein. These leaf characteristics are commonly used to distinguish *H. verticillata* from similar submerged plants in the Hydrocharitaceae, like *Egeria* and *Elodea* spp.

The inflorescences are unisexual, arising from spathes situated in the leaf axils, each flower has three sepals and three petals. All six perianth parts are clear or translucent green (the sepals usually slightly reddish). The male spathe is about 1.5 mm long, solitary in the leaf axils, somewhat spiny. The female spathe is about 5 mm long, solitary in the leaf axils. There are three petals, three stamens and three styles. The ovary is cylindrical to narrowly conical and is enclosed in the base of a hypanthium; the style is as long as the hypanthium and there are three stigmas. For further information, see Cook et al. (1974) and Aston (1977).

The fruit is cylindrical, about 7 mm long and 1.5 mm wide. It contains 2-7 oblong-elliptic seeds. For further information, see Cook and Lüönd (1982); Swarbrick et al. (1981); and Yeo et al. (1984).”

Biology

From GISD (2011):

“*H. verticillata* reproduces mostly by asexual vegetative fragmentation (from stem fragments), but it also grows new plants from tubers and underground tubers and reproduces sexually with flowers. One *H. verticillata* tuber can lead to the production of 5,000 new tubers per square m. It spreads faster in flowing water habitats because the fragments are more efficiently dispersed.

Tubers and turions can survive ice cover, drying, ingestion, and regurgitation by waterfowl. Tubers may remain viable in the sediment for several years.”

From CABI (2015):

“*H. verticillata* is a submerged plant which is rooted by means of filiform, adventitious roots. The stems, which consist of distinct nodes and internodes, are branched and approach or touch the surface of the water. The internodes tend to elongate in flowing water. The flowers are unisexual, arising from spathes situated in the leaf axils, each flower has three sepals and three petals. All six perianth parts are clear or translucent green (the sepals usually slightly reddish). The ovary is enclosed in the base of a hypanthium, the style is as long as the hypanthium and there are three stigmas. Due to an elongation of the hypanthium, the female flower ascends to the surface of the water. The perianth segments remain closed over the stigmas during this movement and retain a bubble of air above them. The perianth segments open to form a wide funnel which floats with its rim just at the water surface, its walls holding back the water and preventing wetting of the stigmas. The male flower becomes detached from the plant and subsequently rises to the surface of the water where the perianth segments uncurl. The anthers dehisce explosively and spread pollen for some 20 cm around the open flower. Pollination occurs via the air.

H. verticillata spreads horizontally by means of branches which grow over the bottom of a waterbody. Vertical branches and roots are produced at nodes on these runners. Vegetative multiplication is also possible by means of fragmentation, i.e. pieces of branches which have become detached are able to form new, rooted plants, if they come into contact with a favourable substratum. In the USA, hydrilla grows optimally at 20-27°C.

It is capable of surviving conditions unfavourable for growth, by producing two types organ capable of remaining dormant for extended periods. These structures are respectively formed in the axil of a leaf (generally described as axillary turions, turions or green turions) and at the tip of branches which grow into the hydrosol (generally described as subterranean turions, brown turions or tubers). (Turions can be defined as short, specialized shoots of aquatic plants in which food material is stored and which eventually become detached from the parent plant). The axillary turions are stalked and cylindrical or slightly conical in shape. The subterranean turions

are boat-shaped and covered by whorls of tough and fleshy scale leaves. For further information on these turions, see the Description section. As many as 1000 (Pieterse, 1981) to 6000 (USDA, 2011) subterranean turions may be produced per square metre in one growing season and remain viable for over 4 years (USDA, 2011). In Florida, USA, the average number of subterranean turions varies from 36 to 207 per m² and the average number of axillary turions from 5 to 90 per m² (Sutton and Portier, 1985). In areas where *H. verticillata* dies during the winter, the formation of turions occurs mainly in the autumn. Axillary turions are frequently formed on free-floating fragments. The formation of subterranean turions is stimulated by short days (Steward and Van, 1987).

There have been numerous studies into the biology of turion production; the most useful of these is a comprehensive review (Netherland, 1997). Additional studies have dealt with the effects of photoperiod on turion development (Steward, 1997; Steward, 2000); factors affecting turion formation (Langeland et al., 1996); the size of turions (Spencer and Ksander, 1995); and the timing of plant development from turions (Spencer and Ksander, 1995; Spencer and Ksander, 1997).

H. verticillata may be either monoecious or dioecious. Its rapid vegetative growth and, as a consequence, the formation of large clones, questions whether strains which produce only male or female flowers are able to reproduce effectively by sexual means. In California and the Gulf States of the USA, and in Europe, there is no seed formation because only female flowers are produced.”

From NANSP (2013):

“Pesacreta (1990) examined carbohydrate allocation in monoecious hydrilla and found that the majority of starch accumulation occurred in plant shoots when exposed to short photoperiods. Starch levels in tubers were found to decrease mostly in the first two weeks after sprouting (Pesacreta 1990). Pesacreta (1990) also found that monoecious hydrilla displayed enhanced fragmentation after 8 weeks of high temperatures (32° C).”

“Monoecious hydrilla tubers have a very high germination rate in laboratory trials, often greater than 90% (Harlan et al. 1985, Van and Steward 1990, personal experience). However, while monoecious hydrilla tubers readily germinate when removed from sediment, when left undisturbed in situ the germination rate is much lower (Van and Steward 1990). Carter et al. (1987) found that monoecious hydrilla tubers require a chilling period prior to sprouting which may prevent sprouting the same year of formation. Monoecious hydrilla tubers have been shown to remain in undisturbed soil for more than 4 years after production (Van and Steward 1990), and six year tubers have still been viable in North Carolina (unpublished data). There appears to be an environmental factor imposed dormancy which prevents depletion of tuber populations. Axillary turions will germinate within one year or not at all (Van and Steward 1990). Nawrocki et al. (2011) also found that monoecious hydrilla tubers have multiple axillary buds preformed within dormant tubers that can produce secondary shoots, even after terminal shoot removal.”

From Jacono et al. (2011):

“Freshwater lakes, ponds, rivers, impoundments and canals.”

Human Uses

From GISD (2011):

“Pet/aquarium trade: Sold as an aquarium plant.”

From Zhuang (2013):

“A dried powder from the plant has be [sic] used as detergent in the treatment of abscesses, burns and wounds. It has been used as an ornamental plant.”

Diseases

No OIE reportable diseases were reported.

From CABI (2015):

“Epiphytic cyanobacteria found on hydrilla are thought to be the agents producing a toxin that causes avian vacuolar myelinopathy (AVM) a disease that has killed at least 100 bald eagles (*Haliaeetus leucocephalus*) and thousands of American coots (*Fulica americana*) since 1994 in locations from Texas to North Carolina, USA (Wilde et al., 2005). The incidence of AVM is likely to increase as *H. verticillata* spreads.”

Threat to Humans

From CABI (2015):

“It can also result in reduced water flow and stagnant pools which become habitats for mosquito larvae. A case study on the social impact of invasion of a lake in Guatemala by hydrilla has been produced by Binemelis et al. (2007).”

From GISD (2011):

“Apart from interfering with fishing, boat motors can become tangled with them and swimming areas choked. *H. verticillata* often slows or clogs rivers, irrigation ditches, and flood control canals, creating stagnant water that is prime mosquito breeding habitat. Dense stands can even cause flooding, alter water quality by decreasing oxygen levels and increasing pH and water temperature.”

3 Impacts of Introductions

From Jacono et al. (2011):

“Once established, hydrilla results in an array of ecosystem disruptions. Changes often begin with its invasion of deep, dark waters where most plants can not grow. Hydrilla grows aggressively and competitively, spreading through shallower areas and forming thick mats in surface waters that block sunlight penetration to native plants below (van Dijk 1985). In the southeast, hydrilla effectively displaces beneficial native vegetation (Bates and Smith 1994) such as wild-celery (*Vallisneria americana*) and coontail (*Ceratophyllum demersum*) (van Dijk 1985; Rizzo et al. 1996).

It has been shown to alter the physical and chemical characteristics of lakes. Colle and Shireman (1980) found sportfish reduced in weight and size when hydrilla occupied the majority of the water column, suggesting that foraging efficiency was reduced as open water space and natural vegetation gradients were lost. Stratification of the water column (Schmitz et al. 1993; Rizzo et al. 1996), decreased oxygen levels (Pesacreta 1988), and fish kills (Rizzo et al. 1996) have been documented. Changes in water chemistry may also be implicated in zooplankton and phytoplankton declines (Schmitz and Osborne 1984; Schmitz et al. 1993).

Hydrilla seriously affects water flow and water use. Infestations in the Mobile Delta are reducing flow in small tidal streams and creating a backwater habitat (J. Zolcynski pers. comm. 1998). Its heavy growth commonly obstructs boating, swimming and fishing in lakes and rivers and blocks the withdrawl [sic] of water used for power generation and agricultural irrigation.”

From GISD (2011):

“*H. verticillata* competes with native plants by growing to the water surface and forming dense mats that totally exclude sunlight from other plants, which in turn can significantly reduce aquatic plant and animal biodiversity. Large populations of *H. verticillata* may affect fish size and population levels where predatory fish cannot hunt effectively within the thick mats. The dense mats also affect recreational activities. Apart from interfering with fishing, boat motors can become tangled with them and swimming areas choked. *H. verticillata* often slows or clogs rivers, irrigation ditches, and flood control canals, creating stagnant water that is prime mosquito breeding habitat. Dense stands can even cause flooding, alter water quality by decreasing oxygen levels and increasing pH and water temperature.”

From CABI (2015):

“*H. verticillata* poses a potential threat to areas outside its native habitats; this has been demonstrated in the USA and the Panama Canal area. As *H. verticillata* is introduced to the New World as an aquarium plant, legislative measures should be taken worldwide to restrain this trade.”

“Due to its rapid growth and a highly effective survival strategy, *H. verticillata* is one of the most troublesome aquatic weeds in the world. It rapidly outcompetes other plant species and forms dense masses, which may completely fill the volume of waterbodies. Consequently, the

often multifunctional use of canals, rivers and lakes becomes seriously hampered by infestations of the weed.

Harmful effects of *H. verticillata* include: impeding the movement of irrigation and drainage water; hindering navigation and recreational use of the water; physical interference with hydro-electric schemes and fisheries; competition with native plants; impacts on native fauna; reductions in size and weight of sport fish (Colle and Shireman, 1980 in Jacono et al., 2011); and the creation of favourable habitats for organisms which cause or transmit disease.

Although it is increasingly troublesome in its original habitat in South-East Asia and Australia, particularly in man-made lakes and irrigation canals, its impact is most significant where it is introduced. This applies, in particular, to the USA, where it was introduced in Florida in the early 1950s (Schardt, 1995). The costs of controlling *H. verticillata* in Florida were reported to be \$200 per ha per year (Haller, 1995) when an area of more than 12,000 ha were heavily infested in the state. Useful summaries of economic and ecological costs due to *H. verticillata* are provided by the Northeast Aquatic Nuisance Species Panel (for the USA) and by Hofstra and Champion (2006; for New Zealand)."

"Epiphytic cyanobacteria found on hydrilla are thought to be the agents producing a toxin that causes avian vacuolar myelinopathy (AVM) a disease that has killed at least 100 bald eagles (*Haliaeetus leucocephalus*) and thousands of American coots (*Fulica americana*) since 1994 in locations from Texas to North Carolina, USA (Wilde et al., 2005). The incidence of AVM is likely to increase as *H. verticillata* spreads."

From NANSP (2013):

"Monoecious hydrilla is an extremely tolerant and competitive plant. It can establish and then displace native plants. Monoecious hydrilla can persist alone and competitively with *Elodea canadensis* (Michx.) in flowing systems like streams and waterways in New Zealand (Hofstra et al. 2010). Spencer and Ksander (2000) showed the strong competitive ability of monoecious hydrilla mixed with American pondweed, and Meadows and Richardson (2012) found that monoecious hydrilla out-competed four other submersed aquatic plants (Eurasian watermilfoil (*Myriophyllum spicatum* L.; invasive), curly leaf pondweed [*Potamogeton crispus* L.; (invasive), *Elodea canadensis* Michx. (native), and *Vallisneria americana* Michx. (native)] in a mesocosm trial."

4 Global Distribution



Figure 1. Known global distribution of both biotypes of *Hydrilla verticillata*. Map from GBIF (2013).

Global biotype distribution data was not available. Jacono et al. (2011) and CABI (2015) state that the populations on the Korean peninsula and some in Europe are of the monoecious biotype, but more detailed information was not available.

5 Distribution Within the United States




Figure 2. Distribution of monoecious *Hydrilla verticillata* by HUC 8 in the United States. Map developed by U.S. Fish and Wildlife Service with data from U.S. Geological Survey.

6 Climate Matching

Summary of Climate Matching Analysis 1: Both Biotypes

The climate match for *Hydrilla verticillata*, both biotypes, was high across the country. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the Continental U.S. was 0.841, high, and all states had an individually high climate match.




Figure 3. RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red) and non-source locations (grey) for *Hydrilla verticillata* climate matching. Source locations are for both biotypes of *Hydrilla verticillata* GBIF (2013).




Figure 4. Map from RAMP (Sanders et al. 2014) of a current climate match for *Hydrilla verticillata* in the continental United States based on source locations reported by GBIF (2013). 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 < X < 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

Future Climate Matches for Both Biotypes

Climate matches for *Hydrilla verticillata* using Representative Concentration Pathways (RCPs) for predicting future climates based on human produced greenhouse gases (IPCC 2014). RCP2.6 predicts a scenario where emission of greenhouse gases is reduced, RCP4.5 predicts continued levels of emission from present, and RCP8.5 predicts a future climate based on rising levels of emissions. Climate matches for each RCP were modeled at two generational time steps, 2050 and 2070. As the climate scenarios predict warmer temperatures, the climate match for the monoecious biotype of *Hydrilla verticillata* lowers (Figure 9).




Figure 5. RAMP (Sanders et al. 2014) climate match using RCP2.6 at the 2050 and 2070 time steps. Climate 6 scores remained high.




Figure 6. RAMP (Sanders et al. 2014) climate match using RCP4.5 at the 2050 and 2070 time steps. Climate 6 scores remained high, with localized areas of medium matches in western states.




Figure 7. RAMP (Sanders et al. 2014) climate match using RCP8.5 at the 2050 and 2070 time steps. Climate 6 scores remained high, with local areas of medium matches in western states.

Summary of Climate Matching Analysis 2: Monoecious Distribution in the United States

The climate match for monoecious *Hydrilla verticillata* was high mainly in the eastern United States and select portions of the West Coast. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the Continental U.S. was 0.525, high, and was high in Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, Washington D.C., Wisconsin, and Wyoming.




Figure 8. RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red) and non-source locations (grey) for monoecious *Hydrilla verticillata* climate matching. Source locations from U.S. Geological Survey.




Figure 9. Map from RAMP (Sanders et al. 2014) of a current climate match for monoecious *Hydrilla verticillata* in the continental United States based on source locations from U.S. Geological Survey. 0 = Lowest match, 10 = Highest match.

Future Climate Matches for the monoecious biotype

Climate matches for monoecious *Hydrilla verticillata* using Representative Concentration Pathways (RCPs) for predicting future climates based on human produced greenhouse gases (IPCC 2014). RCP2.6 predicts a scenario where emission of greenhouse gasses is reduced, RCP4.5 predicts continued levels of emission from present, and RCP8.5 predicts a future climate based on rising levels of emissions. Climate matches for each RCP were modeled at two generational time steps, 2050 and 2070.




Figure 10. RAMP (Sanders et al. 2014) climate match using RCP2.6 at the 2050 and 2070 time steps. Climate 6 scores remained high.




Figure 11. RAMP (Sanders et al. 2014) climate match using RCP4.5 at the 2050 and 2070 time steps. Climate 6 scores remained high.




Figure 12. RAMP (Sanders et al. 2014) climate match using RCP8.5 at the 2050 and 2070 time steps. Climate 6 scores remained high, with local areas of medium matches in western states in the 2050 time step and the western and southern states in the 2070 time step.

7 Certainty of Assessment

The certainty of this assessment is high. There is a wealth of information available about *Hydrilla verticillata* and its impacts. There is some information about the distribution of the two different biotypes in the continental U.S., however distributions of the different biotypes around the world were not available at the time of this assessment. There is a possibility that the climate match would change if a more specific global distribution of the biotypes was known. The information provided above states that the two biotypes have different optimal temperatures. It is the author's opinion that any change in source locations for the climate match would still result in a high climate match for monoecious *Hydrilla verticillata* in the continental United States.

8 Risk Assessment

Summary of Risk to the Contiguous United States

The history of invasiveness is high. *Hydrilla verticillata* is found on virtually every continent with negative impacts. The climate match is high when considering both biotypes and just the monoecious biotype. The climate match for the current climate conditions and for all future scenarios resulted in a high match for both comparisons. The certainty of assessment is high. The overall risk assessment is high.

Assessment Elements

- **History of Invasiveness (Sec. 3): High**
- **Climate Match (Sec. 6): High**
- **Certainty of Assessment (Sec. 7): High**
- **Remarks/Important additional information No additional remarks.**
- **Overall Risk Assessment Category: High**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

CABI. 2015. *Hydrilla verticillata*. In Invasive Species Compendium. CAB International, Wallingford, UK. Available: <http://www.cabi.org/isc/datasheet/28170>. (February 2015).

GBIF (The Global Biodiversity Information Facility). 2013. GBIF backbone taxonomy. Available: <http://www.gbif.org/species/5329570>. (March 2015).

GISD (Global Invasive Species Database), Invasive Species Specialist Group. 2011. Available: <http://issg.org/database/species/ecology.asp?si=272&fr=1&sts=sss&lang=EN>. (February 2015).

IPCC (Intergovernmental Panel on Climate Change). 2014. Scenario Process for AR5. Available: http://sedac.ipcc-data.org/ddc/ar5_scenario_process/RCPs.html. (January 2015).

- ITIS (Integrated Taxonomic Information System). 2015. Available:
http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=38974. (February 2015).
- Jacono, C. C., M. M. Richardson, and V. Howard Morgan. 2011. *Hydrilla verticillata*. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available:
<http://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=6>. (February 2015).
- Madeira, P. T., C. C. Jacono, and T. K. Van. 2000. Monitoring hydrilla using two RAPD procedures and the Nonindigenous Aquatic Species database. *Journal of Aquatic Plant Management* 38:33-40.
- Northeast Aquatic Nuisance Species Panel (NANSP). 2013. Monoecious hydrilla – a review of the literature.
- Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk assessment mapping program: RAMP. U. S. Fish and Wildlife Service.
- Zhuang, X. 2013. *Hydrilla verticillata*. The IUCN Red List of Threatened Species. Version 2014.3. Available: <http://www.iucnredlist.org/details/full/167871/0>. (February 2015).
- USDA, NRCS. 2017. *Hydrilla verticillata* (L. f.) Royle waterhyme. The PLANTS database. National Plant Data Team, Greensboro, North Carolina. Available:
<http://plants.usda.gov/core/profile?symbol=AZMI>. (March 2017).

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

- Ames, et al. 1986. [Source did not give full citation for this reference.]
- Anderson, L. 1996. Eradicating California's hydrilla. *Aquatic Nuisance Species Digest* 1(3):25-33.
- AREC (Aquatic Resources Education Center). 1995. List of aquatic plants found in Delaware ponds 1973-1995. Aquatic Resources Education Center, Division of Fish and Wildlife, Delaware Department of Natural Resources and Environmental Control, Smyrna, Delaware.
- Aston, H. I. 1977. *Aquatic plants of Australia*. Melbourne University Press, Melbourne, Australia.
- Balciunas and Chen. 1993. [Source did not give full citation for this reference.]

- Balcom, N., editor. 1997. Hydrilla in CT update. *Aquatic Exotic News* 4(1):4. Northeast Sea Grant Network, Connecticut Sea Grant, Groton.
- Balevicius, A. 1998. The vegetation of lakes in Veisiejai Regional Park. *Botanica Lithuanica* 4(3):267-284.
- Bates, A. L., and C. S. Smith. 1994. Submersed plant invasions and declines in the southeastern United States. *Lake and Reservoir Management* 10(1):53-55.
- Binemelis, et al. 2007. [Source did not give full citation for this reference.]
- Carter, V., N. B. Rybicki, and C. L. Schulman. 1987. Effect of salinity and temperature on germination of monoecious hydrilla propagules. *Journal of Aquatic Plant Management* 25: 54-57.
- Colangelo, P. A. 1998. Fish and Boat Commission: triploid grass carp permit applications. *Pennsylvania Bulletin* 28(40):4992.
- Colle, D. E., and J. V. Shireman. 1980. Coefficients of condition for largemouth bass, bluegill, and redear sunfish in hydrilla-infested lakes. *Transactions of the American Fisheries Society* 109:521-531.
- Cook, C. D. K., B. J. Gut, E. M. Rix, J. Schneller, and M. Seitz. 1974. *Water plants of the world: a manual for the identification of the genera of freshwater macrophytes*. The Hague, The Netherlands.
- Cook, C. D. K., and R. Lüönd. 1982. A revision of the genus *Hydrilla* (Hydrocharitaceae). *Aquatic Botany* 13(4):485-504
- Environmental Laboratory. 1985. Monoecious Hydrilla in the Potomac River. Miscellaneous Paper MP A-85-5. US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
- Eubanks, M. J. 1996. Lake Seminole hydrilla action plan: an integrated approach. Abstracts 36 Annual Meeting, The Aquatic Plant Management Society, July 14-17, Burlington, Vermont.
- FLDEP (Florida Department of Environmental Protection), Bureau of Aquatic Plant Management. 1988-1994. Florida aquatic plant surveys, electronic data. Bureau of Aquatic Plant Management, Florida Department of Environmental Protection, Tallahassee.
- Godfrey, R. K., and J. W. Wooten. 1979. *Aquatic and wetland plants of southeastern United States, Monocotyledons*. University of Georgia, Athens.
- Haller, W. T. 1982. Hydrilla goes to Washington. *Aquatics* 4(4):6-7.

- Haller, W. T. 1995. Hydrilla control - past, present and future. *Aquatics* 17:6-8.
- Harlan, S. M., G. J. Davis, and G. J. Pesacreta. 1985. Hydrilla in three North Carolina lakes. *Journal of Aquatic Plant Management* 23:68-71. 14
- Helton, R., and L. Hartmann. 1997. Statewide aquatic vegetation survey summary, 1996 report. Statewide Freshwater Fisheries Monitoring and Management Program, Federal Aid in Sport Fish Restoration Act, Project F-30-R-21. Inland Fisheries Division, Jasper, Texas.
- Hofstra, D. E., and P. D. Champion. 2006. Organism consequence assessment: *Hydrilla verticillata*. Hamilton, New Zealand: National Institute for Water and Atmospheric Research.
- Hofstra, D. E., P. D. Champion, and J. S. Clayton. 2010. Predicting invasive success of *Hydrilla verticillata* (L.f) Royle in flowing water. *Hydrobiologia* 656:213-219.
- Kay, S. H. 1992. Hydrilla: a rapidly spreading aquatic weed in North Carolina. Publication AG-449, North Carolina Cooperative Extension Service, North Carolina State University.
- Klosowski, S., and H. Tomaszewicz. 1997. Sociology and ecology of *Hydrillettum verticillatae* and *Elodeetum canadensis* (Pign. 1953) Pass. 1964 in north-eastern Poland. *Tuexenia* 17:125-136.
- Langeland, K. A. 1996. Hydrilla tuber formation in response to single and sequential bensulfuron methyl exposures at different times. *Hydrobiologia* 340(1/3):247-251.
- Les, D. H. 1996. *Hydrilla verticillata* threatens New England. *Aquatic Exotic News* 3(1):1-2.
- Les, D. H., L. J. Mehrhoff, M. A. Cleland, and J. D. Gable. 1997. *Hydrilla verticillata* (Hydrocharitaceae) in Connecticut. *Journal of Aquatic Plant Management* 35:10-14.
- Madeira, P. T., Van, D. Steward, and R. Schnell. 1997. Random amplified polymorphic DNA analysis of the phenetic relationships among world-wide accessions of *Hydrilla verticillata*. *Aquatic Botany* 59:217-236.
- Madeira, P. T., C. C. Jacono, and T. K. Van. 2000. Monitoring hydrilla using two RAPD procedures and the Nonindigenous Aquatic Species Database. *Journal of Aquatic Plant Management* 38:33-40.
- Mataraza, L. K., J. B. Terrell, A. B. Munson, and D. E. Canfield Jr. 1999. Changes in submersed macrophytes in relation to tidal storm surges. *Journal of Aquatic Plant Management* 37:3-12.
- Meadows, S. T., and R. J. Richardson. 2012. Competition of monoecious hydrilla with other submersed macrophytes. Proceedings of the 52nd Annual Meeting Aquatic Plant Management Society (APMS), July 22-25, Salt Lake City, Utah.

- Nawrocki, J. J., S. T. Hoyle, J. Scott, and R. J. Richardson. 2011. Proceedings of the 51st Annual Meeting Aquatic Plant Management Society (APMS), 24-27 July, Baltimore, Maryland. Abstract 37-38.
- NCDWR (North Carolina Division of Water Resources). 1996. Economic and environmental aspects of N.C. aquatic weed infestations. North Carolina Department of Environment, Health, and Natural Resources.
- Netherland, M. D. 1997. Turion ecology of hydrilla. *Journal of Aquatic Plant Management* 35:1-10.
- Orth, R. J., J. F. Nowak, G. F. Anderson, and J. R. Whiting. 1996. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1995. U.S. Environmental Protection Agency, Annapolis, Maryland.
- Pesacreta, G. 1988. Water chemistry from North Carolina piedmont impoundments with Hydrilla (*Hydrilla verticillata* (L.f.)Royle). Doctoral dissertation. North Carolina State University, Raleigh.
- Pesacreta, G. J. 1990. Pilot study: carbohydrate allocation in hydrilla biotypes. Miscellaneous Paper A-90-2. US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
- Pieterse, A. H. 1981. *Hydrilla verticillata* - a review. *Abstracts on Tropical Agriculture* 7(6):9-34.
- Posey, M. H., C. Wigand, and J. C. Stevenson. 1993. Effects of an introduced aquatic plant, *Hydrilla verticillata*, on benthic communities in the Upper Chesapeake Bay. *Estuarine, Coastal and Shelf Science* 37:539-555.
- PPDC (Plant Pest Diagnostic Center). 1997. Weed eradication database. Integrated Pest Control Branch, Division of Plant Industry, California Department of Food and Agriculture, Sacramento.
- Preston, C. D., and J. M. Croft. 1997. Aquatic plants in Britain and Ireland. Harley, Colchester, UK.
- Rizzo, W. M., R. G. Boustany, and D. R. Meaux. 1996. Ecosystem changes in a subtropical Louisiana lake due to invasion by Hydrilla. Abstract in: From Small Streams to Big Rivers, Society of Wetland Scientists 17th Annual Meeting, June 9-14, 1996, Kansas City, Missouri.
- Roach, H., J. Inabinet, and J. Tuten. 1993. Long term plan for aquatic plant management in Lake Marion and Lake Moultrie South Carolina. Santee Cooper Environmental Resources Division, Moncks Corner, South Carolina.

- Rout, N. P., and B. P. Shaw. 2001. Salt tolerance in aquatic macrophytes: possible involvement of the antioxidative enzymes. *Plant Science* 160(3):415-423.
- Rout, N. P., S. B. Tripathi, and B. P. Shaw. 1998. Effect of salinity on chlorophyll and proline contents in three aquatic macrophytes. *Biologia Plantarum* 40(3):453-458.
- Schardt, J. 1994. 1994 Florida aquatic plant survey report. Florida Department of Environmental Protection, Bureau of Aquatic Plant Management, Tallahassee.
- Schardt, J. 1995. Hydrilla reaches crisis levels in Florida waters. *Aquatics* 17:10-12.
- Schmitz, D. C., and J. A. Osborne. 1984. Zooplankton densities in a Hydrilla infested lake. *Hydrobiologia* 111:127-132.
- Schmitz, D. J., J. D. Schardt, A. J. Leslie, F. A. Dray, Jr., J. A. Osborne, and B. V. Nelson. 1993. The ecological impact and management history of three invasive alien aquatic plant species in Florida. Pages 173-194 in B. N. McKnight, editor. *Biological pollution: the control and impact of invasive exotic species*. Indian Academy of Science, Indianapolis, Indiana.
- Spencer, D. F., and G. G. Ksander. 1995. Differential effects of the microbial metabolite, acetic acid, on sprouting of aquatic plant propagules. *Aquatic Botany* 52(1-2):107-119.
- Spencer, D. F., and G. G. Ksander. 1997. Dilute acetic acid exposure enhances electrolyte leakage by *Hydrilla verticillata* and *Potamogeton pectinatus* tubers. *Journal of Aquatic Plant Management* 35:25-30.
- Spencer, D. F., and G. G. Ksander. 2000. Interactions between American pondweed and monoecious hydrilla grown in mixtures. *Journal of Aquatic Plant Management* 38:5-13.
- Steward, K. K. 1997. Influence of photoperiod on tuber production in various races of *Hydrilla (Hydrilla verticillata)*. *Hydrobiologia*, 354:57-62.
- Steward, K. K. 2000. Influence of photoperiod on vegetative propagule production in three turion-producing races of *Hydrilla verticillata* (L.f.) Royle. *Hydrobiologia* 432(1/3):1-8.
- Steward, K. K., and T. K. Van. 1987. Comparative studies of monoecious and dioecious hydrilla (*Hydrilla verticillata*) biotypes. *Weed Science* 35(2):204-210.
- Steward, K. K., T. K. Van, V. Carter, and A. H. Pieterse. 1984. Hydrilla invades Washington, DC and the Potomac. *American Journal of Botany* 71(1):162-163
- Sutton, D. L., and K. M. Portier. 1985. Density of tubers and turions of hydrilla in south Florida. *Journal of Aquatic Plant Management* 23:64-67

- Swarbrick, J. T., C. M. Finlayson, and A. J. Cauldwell. 1981. The biology of Australian weeds. 7. *Hydrilla verticillata* (L.f.) Royle. *Journal of the Australian Institute of Agricultural Science* 47(4):183-190.
- Tennessee Valley Authority. 1990. Final environmental assessment: demonstration of use of grass carp in management of aquatic plants in Guntersville Reservoir. TVA/RDG/EQS-90/3, Muscle Shoals, Alabama.
- USDA. 2011. *Hydrilla verticillata* Royle. Draft Federal Noxious Weed List Fact Sheet. Unpublished, available from USDA.
- Van, T. K. 1989. Differential responses to photoperiods in monoecious and dioecious *Hydrilla verticillata*. *Weed Science* 37:552-556.
- Van, T., and K. Steward. 1990. Longevity of monoecious hydrilla propagules. *Journal of Aquatic Plant Management* 28:74-76.
- Van Dijk, G. 1985. *Vallisneria* and its interactions with other species. *Aquatics* 7(3):6-10.
- White, A., J. B. Reiskind, and G. Bowes. 1996. Dissolved inorganic carbon influences the photosynthetic responses of *Hydrilla* to photoinhibitory conditions. *Aquatic Botany* 53(1/2):3-13.
- Wilde, S. B., T. M. Murphy, C. P. Hope, S. K. Habrun, J. Kempton, A. Birrenkott, F. Wiley, W. W. Bowerman, and A. J. Lewitus. 2005. Avian vacuolar myelinopathy linked to exotic aquatic plants and a novel cyanobacterial species. *Environmental Toxicology* 20(3):348-353.
- Yeo, R. R., R. H. Falk, and J. R. Thurston. 1984. The morphology of hydrilla (*Hydrilla verticillata*) (L.f.) Royle. *Journal of Aquatic Plant Management* 22:1-17.
- Zolczynski, J. 1997. Mobile Delta submersed aquatic vegetation survey, 1994. Fisheries Section, Game and Fish Division, Alabama Department of Conservation and Natural Resources, Spanish Fort.