

Midas Cichlid (*Amphilophus citrinellus*)

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, web version – 3/30/2018



Photo: Tanya Dewey. Licensed under Creative Commons BY-NC-SA 3.0 Unported. Available: http://animaldiversity.org/collections/contributors/tanya_dewey/reddevilcichlid2/. (May 2017).

1 Native Range and Status in the United States

Native Range

From Froese and Pauly (2014):

“Central America: Atlantic slope of Nicaragua and Costa Rica (San Juan River drainage, including Lakes Nicaragua, Managua, Masaya and Apoyo).”

Status in the United States

From Nico and Neilson (2011):

“First collected in July 1980, the species is now established in the Black Creek Canal system, Dade County, Florida (Anderson et al. 1984; Loftus and Kushlan 1987). It was taken from the Mowry (C-103) Canal, Miami-Dade County, in July 1992 (W. Smith-Vaniz, pers. comm.). It is relatively common there as well as in other connected canals such as C-103N and C-102 (W.F. Loftus, pers. comm.). There is an unconfirmed report from Bivens Arm Lake in Gainesville, Alachua County, March 1985 (J. Miller, pers. comm.); this record is not supported by a voucher specimen and the report is considered somewhat suspect. A voucher specimen exists [sic] for Hillsborough County from 1976 in Six-mile Creek. Collections of specimens from other south Florida canals. The species has been collected from the University of Florida campus (Hill and Cichra 2005). The first report from Hawaii was of six specimens found dead, possibly from fungal infections, in Wahiawa Reservoir on Oahu, in February 1991 (Devick 1991); the species is now documented as being firmly established in Wahiawa Reservoir, and it also may be in other localities on Oahu (M. Yamamoto, pers. comm.). One fish was taken from Dodgeville Pond in Attleboro, Bristol County, Massachusetts, in July 1990 (Hartel 1992; Cardoza et al. 1993). One fish was found dead in Silver Lake in North St. Paul, Ramsey County, Minnesota (K. Schmidt, pers. comm.). In Puerto Rico, the fish has become established in the Canaboncito River, Canabon River, and La Plata River, and the Guajataca, Loiza, Dos Bocas, and La Plata Reservoirs (F. Grana, pers. comm.), but no voucher specimens exist for these locations.”

“Established in Florida, Hawaii, and Puerto Rico. Failed in Massachusetts and Minnesota.”

Means of Introductions in the United States

From Nico and Neilson (2011):

“Introductions into Florida were via aquarium or fish farm releases. In other states, introductions most likely represent aquarium releases.”

Remarks

No additional remarks.

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2014):

“Synonym(s): *Cichlasoma citrinellum* (Günther, 1864)

Kingdom Animalia

Subkingdom Bilateria

Infrakingdom Deuterostomia

Phylum Chordata

Subphylum Vertebrata
Infraphylum Gnathostomata
Superclass Osteichthyes
Class Actinopterygii
Subclass Neopterygii
Infraclass Teleostei
Superorder Acanthopterygii
Order Perciformes
Suborder Labroidei
Family Cichlidae
Genus *Amphilophus* Agassiz, 1859
Species *Amphilophus citrinellus* (Günther, 1864)”

From Eschmeyer et al. (2017):

“*citrinellus*, *Heros* Günther [A.] 1864:153 [Proceedings of the Zoological Society of London 1864 (pt 1) (art. 3) (for 22 Mar. 1864) [...]] Lake Nicaragua, Nicaragua. Syntypes: BMNH 1864.1.26.201-203 (3). •Valid as *Cichlosoma citrinellum* (Günther 1864) -- (Bussing 1987:195 [...]). •Valid as *Cichlasoma citrinellum* (Günther 1864) -- (Page & Burr 1991:332 [...], Conkel 1993:83 [...], Fuller et al. 1999:420 [...], Nelson et al. 2004:151 [...]). •Valid as *Amphilophus citrinellus* (Günther 1864) -- (Kullander & Hartel 1997:195 [...], Bussing 1998:312 [...], Burgess 2000:54 [...], Kullander in Reis et al. 2003:610 [...], Stauffer & McKaye 2002:8 [...], Mundy 2005:418 [...], Stauffer et al. 2008:117 [...], Scharpf 2009:3 [...] as *citrinellum*, Oldfield 2009:1 [...], Page & Burr 2011:610 [...], Page et al. 2013:156 [...], Kottelat 2013:373 [...], Angulo et al. 2013:1004 [...], Řičan et al. 2016:37 [...]). **Current status:** Valid as *Amphilophus citrinellus* (Günther 1864). Cichlidae: Cichlinae.”

Size, Weight, and Age Range

From Froese and Pauly (2014):

“Max length: 24.4 cm SL male/unsexed; [Kullander 2003]”

Environment

From Froese and Pauly (2014):

“Freshwater; benthopelagic. [...]; 23°C - 33°C [assumed to be recommended aquarium temperature range] [Conkel 1993]; [...]”

Climate/Range

From Froese and Pauly (2014):

“Tropical; [...]; 15°N - 8°N”

Distribution Outside the United States

Native

From Froese and Pauly (2014):

“Central America: Atlantic slope of Nicaragua and Costa Rica (San Juan River drainage, including Lakes Nicaragua, Managua, Masaya and Apoyo).”

Introduced

From Froese and Pauly (2014):

“Introduced to Taiwan, Puerto Rico, Singapore, Philippines”

From Nico and Neilson (2011):

“Midas cichlid, along with red devil cichlid (*A. labiatus*) were introduced to, and are established in, Queensland, Australia in 1992 (Lintermans 2004).”

From FAO (2014):

“Introduced to Philippines from unknown.
Introduced to Singapore from unknown.”

From Kwik et al. (2013):

“*A. citrinellus*, the dominant species in the surveyed urban ponds, has also established itself in Thailand and Australia (Wilson 2005; Nico et al. 2007) [...]”

Means of Introduction Outside the United States

From FAO (2014):

“Reasons of Introduction: ornamental”

From Corfield et al. (2008:82):

“No explicit risk beyond concerns about releases by aquarists and escapes from garden ponds during flood events. The number of fish sold in Australia is relatively small and this species is of medium importance to the industry (Table 7.1 [in original source]). The risk of spread by humans is low.”

Short Description

From Froese and Pauly (2014):

“This species coloration is mostly bright orange to orange-red in adults; mature males are of larger size, longer fins and with a distinct hump on their heads; aquarists usually refer to the lighter-colored, thin-lipped form as this species.”

Biology

From Froese and Pauly (2014):

“Lives in box-cut canals with rocky vertical sides, crevices used for spawning and protection of the young [Page and Burr 1991]. Found in lakes; uncommon in the rivers but will penetrate the lower river valleys where the water is slow flowing or tranquil [Conkel 1993]. Omnivorous, eating mostly aufwuchs, snails and small fishes [Conkel 1993]; also feeds on insect larvae, worms and other bottom-dwelling organisms [Yamamoto and Tagawa 2000]. Majority of this fish has normal cryptic coloration (black, gray or brown), matching the substrate for camouflage and survival purposes. About 10 % of this species is xanthomorphic, undergoing a color metamorphosis at varying stages of growth [Conkel 1993].”

“Spawn preferentially on the ceiling of natural caves [Lavery 1991]. Deposit eggs on hard substrates, such as rocks or logs; both parents guarding the eggs and the fry for several weeks [Yamamoto and Tagawa 2000]. 300-1000 eggs [Baensch and Riehl 1985].”

From Nico and Neilson (2011):

“Primarily omnivorous, opportunistically consuming algae, insect larvae, benthic invertebrates, and fishes (Barlow 1976). Midas cichlids are substrate spawners, with reproduction generally occurring during the rainy season with breeding pairs defending a small territory around some form of cover (Noakes and Barlow 1973; Barlow 1976). Parental care is provided through protection of eggs and schooling fry from predators and through 'contacting' behavior, where fry will consume dermal mucus of the parents as part of their diet (Noakes and Barlow 1973; Barlow 1976).”

“Generally found in lakes and estuaries over a variety of substrate types; uncommon to rare in rivers and streams. Usually associated with some sort of structure or cover (e.g., rocky outcrops, logs).”

Human Uses

From Froese and Pauly (2014):

“An experimental fish being used for behavioral studies [Robins et al. 1991].”

“Fisheries: of no interest; aquarium: commercial”

Diseases

No records of OIE reportable diseases found.

From Froese and Pauly (2014):

“*Procamallanus* Infection 13, Parasitic infestations (protozoa, worms, etc.)
Sciadicleithrum Infection, Parasitic infestations (protozoa, worms, etc.)”

Threat to Humans

From Froese and Pauly (2014):

“Harmless”

3 Impacts of Introductions

No records of documented impacts from introductions were found. The following outlines potential impacts.

From Nico and Neilson (2011):

“Impact of Introduction: Unknown.”

From Kwik et al. (2013):

“The results also appear to show that the fish community consisted of introduced species with the dominant species including two feral species (*Amphilophus citrinellus* and *Oreochromis mossambicus*) with established breeding populations (i.e. presence of breeding pits and size classes ranging from juveniles to egg brooding adults).”

“*A. citrinellus*, the dominant species in the surveyed urban ponds, has also established itself in Thailand and Australia (Wilson 2005; Nico et al. 2007), but it is uncertain what impacts this species might have on the native fishes of Singapore if it spreads into reservoirs or natural stream or swamp systems.”

From Corfield et al. (2008:82):

“The red devil [*Amphilophus citrinellus*] is included in the proposed grey list for ornamental fish indicating that it is of some concern but that more information is required (NRMMC 2006). Currently known to be present in only one location (Ross River, northern Queensland).”

4 Global Distribution




Figure 1. Known global distribution of *Amphilophus citrinellus*. Map from Froese and Pauly (2014).



Figure 2. Known global distribution of *Amphilophus citrinellus*. Map from GBIF Secretariat (2014).

Locations in Massachusetts and Minnesota are from failed introductions (Nico and Neilson 2011). The location in India could not be verified as from an established population. None of these locations were used as source points in the climate match.




Figure 3. *Amphilophus citrinellus* population in Australia (Corfield et al. 2008).

5 Distribution Within the United States



Figure 4. Known distribution of *Amphilophus citrinellus* in the United States. Map from USGS NAS Database (Nico and Neilson 2011).

Locations in Massachusetts and Minnesota are from failed introductions and were not used as source points in the climate match.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match for *Amphilophus citrinellus* was very high for Florida and a portion of Texas' Gulf Coast. There was medium match to areas along the southern Atlantic Coast, Gulf Coast, and parts of southern Arizona and California. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the contiguous U.S. was 0.044, medium, and Florida, Georgia, North Carolina, and South Carolina had individually high climate matches.




Figure 5. RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red) and non-source locations (grey) for *Amphilophus citrinellus* climate matching. Source locations from Nico and Neilson (2011), Froese and Pauly (2014), and GBIF Secretariat (2014).




Figure 6. Map of RAMP (Sanders et al. 2014) climate matches for *Amphilophus citrinellus* in the contiguous United States based on source locations reported by Nico and Neilson (2011), Froese and Pauly (2014), and GBIF Secretariat (2014). 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X < 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

The certainty of this assessment is medium. There was adequate information available for *Amphilophus citrinellus*. Records of introductions were found. The information found for impacts of introductions did not document actual impacts.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Amphilophus citrinellus is a freshwater fish native to Central America. The history of invasiveness is not documented. There are records of introductions, with some establishing populations, going as far back as the 1970s. No information on documented impacts was found, just potential impacts. *A. citrinellus* is present in the pet trade but at a low volume. There is already an established population in Florida and the climate match indicates that there is suitable climate to support the spread of this species. The climate match is medium. The certainty of assessment is medium. The overall risk assessment category is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): None Documented**
- **Climate Match (Sec. 6): Medium**
- **Certainty of Assessment (Sec. 7): Medium**
- **Remarks/Important additional information** No additional remarks.
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

Corfield, J., B. Diggles, C. Jubb, R. M. McDowall, A. Moore, A. Richards, and D. K. Rowe. 2008. Review of the impacts of introduced ornamental fish species that have established wild populations in Australia. Report. Department of the Environment, Water, Heritage, and the Arts, Australia.

Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2017. Catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (May 2017).

FAO (Fisheries and Agriculture Organization of the United Nations). 2014. Database on introductions of aquatic species. FAO, Rome. Available: <http://www.fao.org/fishery/introsp/search/en>. (April 2014).

Froese, R., and D. Pauly, editors. 2014. *Amphilophus citrinellus* (Günther, 1864). FishBase. Available: <http://www.fishbase.org/summary/Amphilophus-citrinellus.html>. (April 2014).

GBIF Secretariat. 2014. GBIF backbone taxonomy: *Amphilophus citrinellus* (Günther, 1864). Global Biodiversity Information Facility, Copenhagen. Available: <http://www.gbif.org/species/2370102>. (April 2014).

Google Inc. 2011. Google Earth (Version 7.0.3.8542) [Software]. Available: <http://www.google.com/intl/en/earth/index.html>. (April 2014).

ITIS (Integrated Taxonomic Information System). 2014. *Amphilophus citrinellus* (Günther, 1864). Integrated Taxonomic Information System, Reston, Virginia. Available: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=648192. (April 2014).

Kwik, J. T. B., Z. Y. Kho, B. S. Quek, H. H. Tan, and D. C. J. Yeo. 2013. Urban stormwater ponds in Singapore: potential pathways for spread of alien freshwater fishes. *BioInvasions Records* 2(3):239–245.

Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk assessment mapping program: RAMP. U.S. Fish and Wildlife Service.

Nico, L., and M. Neilson. 2011. *Amphilophus citrinellus*. U.S. Geological Survey, Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <http://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=442>. (April 2014).

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Anderson, R. S., R. J. Metzger, and P. L. Shafland. 1984. Establishment of the Midas cichlid, *Cichlasoma citrinellum*, in Florida. *Florida Scientist* 47:263–267.

Angulo, A., C. A. Garita-Alvarado, W. A. Bussing, and M. I. López. 2013. Annotated checklist of the freshwater fishes of continental and insular Costa Rica: additions and nomenclatural revisions. *Check List* 9(5):987–1019.

Baensch, H. A., and R. Riehl. 1985. *Aquarien atlas. Band 2*. Mergus, Verlag für Natur-und Heimtierkunde GmbH, Melle, Germany.

Barlow, G. W. 1976. The Midas cichlid in Nicaragua. Pages 333–358 in T. B. Thorson, editor. *Investigations of the ichthyofauna of Nicaraguan lakes*. University of Nebraska, Lincoln.

Burgess, W. E. 2000. The *Cichlasoma* story. *Herichthys*, the break-up. *Tropical Fish Hobbyist* 48(11):44–54.

Bussing, W. A. 1998. Peces de las aguas continentales de Costa Rica. *Revista de Biología Tropical* 46(Supplement 2):1–468.

- Cardoza, J. E., G. S. Jones, T. W. French, and D. B. Halliwell. 1993. Exotic and translocated vertebrates of Massachusetts, 2nd edition. Fauna of Massachusetts Series 6. Massachusetts Division of Fisheries and Wildlife, Publication 17223-110-200-11/93-C.R, Westborough.
- Conkel, D. 1993. Cichlids of North and Central America. T.F.H. Publications.
- Devick, W. S. 1991. Disturbances and fluctuations in the Wahiawa Reservoir ecosystem. Project F-14-R-15, Job 4, Study I. Division of Aquatic Resources, Hawaii Department of Land and Natural Resources.
- Fuller, P. L., L. G. Nico, and J. D. Williams. 1999. Nonindigenous fishes introduced into inland waters of the United States. American Fisheries Society, Special Publication 27, Bethesda, Maryland.
- Günther, A. 1864. Report of a collection of fishes made by Messrs. Dow, Godman, and Salvin in Guatemala. Part first. Proceedings of the Zoological Society of London 1864(1)(3):144–154.
- Hartel, K. E. 1992. Non-native fishes known from Massachusetts freshwaters. Harvard University, Museum of Comparative Zoology, Fish Department, Occasional Report, Cambridge, Massachusetts.
- Hill, J. E., and C. E. Cichra. 2005. Eradication of a reproducing population of convict cichlids, *Cichlasoma nigrofasciatum* (Cichlidae), in north-central Florida. Florida Scientist 68:65–74.
- Kottelat, M. 2013. The fishes of the inland waters of Southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. Raffles Bulletin of Zoology Supplement 27:1–663.
- Kullander, S. O. 2003. Cichlidae (Cichlids). Pages 605–654 in R. E. Reis, S. O. Kullander, and C. J. Ferraris, Jr., editors. Checklist of the freshwater fishes of South and Central America. EDIPUCRS, Porto Alegre, Brazil.
- Kullander, S. O., and K. E. Hartel. 1997. The systematic status of cichlid genera described by Louis Agassiz in 1859: *Amphilophus*, *Baiodon*, *Hypsophrys* and *Parachromis* (Teleostei: Cichlidae). Ichthyological Exploration of Freshwaters 7(3):193–202.
- Lavery, R. J. 1991. Physical factors determining spawning site selection in a Central American hole nester, *Cichlasoma nigrofasciatum*. Environmental Biology of Fishes 31(2):203–206.
- Lintermans, M. 2004. Human-assisted dispersal of alien freshwater fish in Australia. New Zealand Journal of Marine and Freshwater Research 38:481–501.

- Loftus, W. F., and J. A. Kushlan. 1987. Freshwater fishes of southern Florida. *Bulletin of the Florida State Museum of Biological Science* 31(4):255.
- Mundy, B. C. 2005. Checklist of the fishes of the Hawaiian Archipelago. *Bishop Museum Bulletins in Zoology* 6:1–703.
- Nelson, J. S., E. J. Crossman, H. Espinosa Pérez, L. T. Findley, C. R. Gilbert, R. N. Lea, and J. D. Williams. 2004. Common and scientific names of fishes from the United States, Canada, and Mexico, 6th edition. American Fisheries Society, Special Publication 29, Bethesda, Maryland.
- Nico, L. G., W. H. Beamish, and P. Musikasinthorn. 2007. Discovery of the invasive Mayan cichlid fish “*Cichlasoma*” *urophthalmus* (Günther, 1862) in Thailand, with comments on other introductions and potential impacts. *Aquatic Invasions* 2:197–214.
- Noakes, D. L. G., and G. W. Barlow. 1973. Ontogeny of parent-contacting in young *Cichlasoma citrinellum* (Pisces, Cichlidae). *Behaviour* 46(3/4):221–255.
- NRMMC (Natural Resource Management Ministerial Council). 2006. A strategic approach to the management of ornamental fish in Australia. Department of Agriculture, Fisheries and Forestry, Australia.
- Oldfield, R. G. 2009. Captive breeding observations support the validity of a recently described cichlid species in Lake Apoyo, Nicaragua. *Occasional Papers of the Museum of Zoology University of Michigan* 741:1–16.
- Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes of North America north of Mexico. Houghton Mifflin, Boston.
- Page, L. M., and B. M. Burr. 2011. Peterson field guide to freshwater fishes of North America north of Mexico, 2nd edition). *Freshwater fishes of North America*.
- Page, L. M., H. Espinosa-Pérez, L. D. Findley, C. R. Gilbert, R. N. Lea, N. E. Mandrak, R. L. Mayden, and J. S. Nelson. 2013. Common and scientific names of fishes from the United States, Canada, and Mexico, 7th edition. American Fisheries Society, Special Publication 34, Bethesda, Maryland.
- Reis, R. E., S. O. Kullander, and C. J. Ferraris, Jr., editors. 2003. Check list of the freshwater fishes of South and Central America. CLOFFSCA. EDIPUCRS, Porto Alegre, Barzil.
- Říčan, O., L. Piálek, K. Dragová, and J. Novák. 2016. Diversity and evolution of the Middle American cichlid fishes (Teleostei: Cichlidae) with revised classification. *Vertebrate Zoology* 66(1):1–102.

- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott. 1991. Common and scientific names of fishes from the United States and Canada. American Fisheries Society, Special Publication 20, Bethesda, Maryland.
- Scharpf, C. 2009. Annotated checklist of North American freshwater fishes, including subspecies and undescribed forms. Part V: Sciaenidae ... [through] Achiridae (plus supplemental material). *American Currents* 35(1):1–32.
- Stauffer, J. R., Jr., J. K. McCrary, and K. E. Black. 2008. Three new species of cichlid fishes (Teleostei: Cichlidae) from Lake Apoyo, Nicaragua. *Proceedings of the Biological Society of Washington* 121(1):117–129.
- Stauffer, J. R., Jr., and K. R. McKaye. 2002. Descriptions of three new species of cichlid fishes (Teleostei: Cichlidae) from Lake Xiloá, Nicaragua. *Cuadernos de investigación de la UCA, Universidad Centroamericana* 12:1–18.
- Wilson, G. G. 2005. Impact of invasive exotic fishes on wetland ecosystems in the Murray-Darling Basin. Pages 45–60 *in* B. Phillip, editor. *Native fish and wetlands in the Murray-Darling Basin workshop*. Murray-Darling Basin Commission, Canberra, Australia.
- Yamamoto, M. N., and A. W. Tagawa. 2000. *Hawai'i's native and exotic freshwater animals*. Mutual Publishing, Honolulu, Hawaii.