

Tilapia walteri

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, June 2015

Photo not available.

1 Native Range, and Status in the United States

Native Range

From Froese and Pauly (2015):

“Africa: Cavally and Nipoué (= Cess) Rivers in Côte d'Ivoire and Liberia; also reported from the upper St. John River in Liberia [Teugels and Thys van den Audenaerde 1992, 2003].”

Status in the United States

This species has not been reported in the U.S.

Means of Introductions in the United States

This species has not been reported in the U.S.

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2015):

“Kingdom Animalia
Subkingdom Bilateria
Infrakingdom Deuterostomia
Phylum Chordata
Subphylum Vertebrata
Infraphylum Gnathostomata
Superclass Osteichthyes
Class Actinopterygii
Subclass Neopterygii
Infraclass Teleostei
Superorder Acanthopterygii
Order Perciformes
Suborder Labroidei
Family Cichlidae
Genus *Tilapia*
Species *Tilapia walteri* Thys van den Audenaerde, 1968”

“Taxonomic Status: valid”

Size, Weight, and Age Range

From Froese and Pauly (2015):

“Max length : 27.0 cm TL male/unsexed; [Teugels and Thys van den Audenaerde 1991]”

Environment

From Froese and Pauly (2015):

“Freshwater; benthopelagic.”

Climate/Range

From Froese and Pauly (2015):

“Tropical; 24°C - 26°C [Baensch and Riehl 1995]; 8°N - 5°N”

Distribution Outside the United States

Native

From Froese and Pauly (2015):

“Africa: Cavally and Nipoué (= Cess) Rivers in Côte d'Ivoire and Liberia; also reported from the upper St. John River in Liberia [Teugels and Thys van den Audenaerde 1992, 2003].”

Introduced

No introductions of this species have been reported.

Means of Introduction Outside the United States

No introductions of this species have been reported.

Short description

From Froese and Pauly (2015):

“Dorsal spines (total): 15 - 16; Dorsal soft rays (total): 12-13; Anal spines: 3; Anal soft rays: 9 - 10. Diagnosis: lower pharyngeal bone about as long as broad, with anterior lamella shorter than toothed area; median pharyngeal teeth not broadened; upper part of caudal fin yellow-orange, lower part greyish; dorsal fin with 15-16 spines and 12-14 soft rays; 9-13 lower gill-rakers [Teugels and Thys van den Audenaerde 2003].”

Biology

No information available.

Human uses

From Awaïss and Lalèyè (2010):

“This species is harvested for human consumption.”

Diseases

From le Roux and Avenant-Oldewage (2010):

“*C[ichlidogyrus] aegypticus* Ergens 1981 ... Other host/s: *Tilapia walteri*
C. arthracanthus Paperna 1960 ... Other host/s: *Tilapia walteri*
C. cubitus Dossou 1982 ... Other host/s: *Tilapia walteri*
C. digitatus Dossou 1982 ... Other host/s: *Tilapia walteri*
C. ergensi Dossou 1982 ... Other host/s: *Tilapia walteri*
C. gallus Pariselle and Euzet 1995, syn. of *C. aegypticus* Ergens 1981 sensu Dossou (1982) ...
Other host/s: *Tilapia walteri*
C. tiberianus Paperna 1960 ... Other host/s: *Tilapia walteri*
C. yanni Pariselle and Euzet 1996 ... Other host/s: *Tilapia walteri*”

No OIE-notifiable diseases have been reported for this species.

Threat to humans

From Froese and Pauly (2015):

“Harmless”

3 Impacts of Introductions

No introductions of this species have been reported.

4 Global Distribution



Figure 1. Global distribution of *T. walteri*. Map from GBIF (2015).

5 Distribution within the United States

This species has not been reported in the U.S.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match (Sanders et al. 2014; 16 climate variables; Euclidean Distance) was low throughout the contiguous U.S., reflected in a Climate 6 proportion of 0.0. The range for a low climate match is 0.000 to 0.005.




Figure 2. RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red) and non-source locations (gray) for *T. walteri* climate matching. Source locations from GBIF (2015).




Figure 3. Map of RAMP (Sanders et al. 2014) climate matches for *T. walteri* in the continental United States based on source locations reported by GBIF (2015). 0= Lowest match, 10=Highest match. Counts of climate match scores are tabulated on the left.

7 Certainty of Assessment

Little information is available on the biology of *T. walteri* and it has not become established outside its native range. The certainty of this assessment is high because the lack of information about this species precludes any assessment other than “uncertain” risk.

8 Risk Assessment

Summary of Risk to the Continental United States

Tilapia walteri is a benthopelagic cichlid native to river systems in Côte d’Ivoire and Liberia. It has not been reported as introduced outside of this location. Because *T. walteri* has no history of invasiveness, it is currently impossible to know what impacts *T. walteri* might have if introduced to the U.S. Climate match to the contiguous U.S. is low. Overall risk posed by this species is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3):** Uncertain
- **Climate Match (Sec.6):** Low
- **Certainty of Assessment (Sec. 7):** High
- **Overall Risk Assessment Category: Uncertain**

DRAFT

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

Awaïss, A., and P. Lalèyè. 2010. *Tilapia walteri*. The IUCN Red List of Threatened Species, version 2015.2. Available: <http://www.iucnredlist.org/details/181693/0>. (June 2015).

Froese, R., and D. Pauly, editors. 2015. *Tilapia walteri* Thys van den Audenaerde, 1968. FishBase. Available: <http://www.fishbase.org/summary/2490>. (June 2015).

Global Biodiversity Information Facility (GBIF). 2015. GBIF backbone taxonomy: *Tilapia walteri* Thys van den Audenaerde, 1968. Global Biodiversity Information Facility, Copenhagen. Available: <http://www.gbif.org/species/2370717>. (June 2015).

Integrated Taxonomic Information System (ITIS). 2015. *Tilapia walteri* Thys van den Audenaerde, 1968. Integrated Taxonomic Information System, Reston, Virginia. Available: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=648986. (June 2015).

Le Roux, L. E., and A. Avenant-Oldewage. 2010. Checklist of the fish parasitic genus *Cichlidogyrus* (Monogenea), including its cosmopolitan distribution and host species. *African Journal of Aquatic Science* 35(1):21-36.

Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk Assessment Mapping Program: RAMP. US Fish and Wildlife Service.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Baensch, H. A., and R. Riehl. 1995. *Aquarien Atlas*, volume 4. Mergus Verlag GmbH, Verlag für Natur-und Heimtierkunde, Melle, Germany.

Teugels, G. G., and D. F. E. Thys van den Audenaerde. 1991. *Tilapia*. Pages 482-508 in J. Daget, J.-P. Gosse, G. G. Teugels, and D. F. E. Thys van den Audenaerde, editors. *Check-list of the freshwater fishes of Africa (CLOFFA)*, volume 4. ISNB, Brussels; MRAC, Tervuren, Belgium; and ORSTOM, Paris.

Teugels, G. G., and D. F. E. Thys van den Audenaerde. 1992. *Cichlidae*. Pages 714-779 in C. Levêque, D. Paugy, and G. G. Teugels, editors. *Faune des poissons d'eaux douces et saumâtres d'Afrique de l'Ouest*, volume 2. Coll. Faune Tropicale n° 28. Musée Royal de l'Afrique Centrale, Tervuren, Belgium, and O.R.S.T.O.M., Paris.

Teugels, G. G., and D. F. E. Thys van den Audenaerde. 2003. Cichlidae. Pages 521-600 in D. Paugy, C. Lévêque and G. G. Teugels, editors. The fresh and brackish water fishes of West Africa, volume 2. Coll. faune et flore tropicales 40. Institut de recherche de développement, Paris, France, Muséum national d'histoire naturelle, Paris, France and Musée royal de l'Afrique Central, Tervuren, Belgium.

DRAFT