Stage I Assessment Report, Volume 1 — Injury Assessment: Kalamazoo River Environment

Final Report

Michigan Department of Environmental Quality,
Michigan Attorney General,
U.S. Fish and Wildlife Service,
and
National Oceanic and Atmospheric Administration

Prepared by:

Stratus Consulting Inc.
PO Box 4059
Boulder, CO 80306-4059
(303) 381-8000

March 15, 2005
SC10087

Contents

List of Figures vii
List of Tables xi

Chapter 1 Introduction 1-1

1.1 Natural Resource Damage Assessment Process 1-1
1.2 Public Review and Comment 1-2
1.3 Background Information on the Assessment Area 1-3
1.4 Potentially Responsible Parties 1-6
1.5 Spatial and Temporal Extent of the Stage I Assessment 1-7
1.6 Organization of the Stage I Injury Assessment Report 1-7

Chapter 2 PCB Releases and Pathways 2-1

2.1 PCBs 2-1
2.2 PCB Releases from PRP Facilities 2-2
2.2.1 History of PCB use at PRP facilities 2-3
2.2.2 Evidence of PCB releases from Allied facilities 2-6
2.2.3 Evidence of PCB releases from Georgia-Pacific facilities 2-11
2.2.4 Evidence of PCB releases from Simpson Plainwell facilities 2-17
2.2.5 Evidence of PCB releases from Fort James facilities 2-21
2.3 PCB Transport Pathways 2-23
2.4 PCB Contamination and Persistence in the KRE 2-27
2.4.1 PCB contamination in the KRE 2-27
2.4.2 Persistence of PCBs in the KRE 2-33
2.5 Conclusions: PCB Releases, Pathways, and Contamination 2-36

Chapter 3 Injuries to Surface Water 3-1

3.1 Injury Definitions 3-1
3.2 Stage I Injury Assessment Approach 3-2
3.3 Drinking Water Standard Exceedences 3-2
3.3.1 Data sources 3-2
3.3.2 Regulatory criteria and standards 3-3
3.3.3 Results 3-5

3.4 Water Quality Criteria Exceedences 3-8
3.4.1 Data sources 3-8
3.4.2 Regulatory criteria and standards 3-8
3.4.3 Results 3-9
3.5 Conclusions 3-12

Chapter 4 Injuries to Sediment 4-1

4.1 Injury Definitions 4-1
4.2 Stage I Injury Assessment Approach 4-1
4.3 Comparison to Concentrations Affecting Benthic Invertebrates 4-1
4.3.1 Data sources 4-2
4.3.2 Toxicity reference value derivation 4-3
4.3.3 Results 4-5
4.4 Dietary Exposure to Mink 4-7
4.4.1 Data sources 4-8
4.4.2 Toxicity reference value derivation 4-8
4.4.3 Results 4-9
4.5 Temporal Extent 4-11
4.6 Conclusions 4-12

Chapter 5 Injuries to Wildlife Services: Fish Consumption Advisories 5-1

5.1 Injury Definitions 5-1
5.2 Stage I Injury Assessment Approach 5-1
5.3 History of KRE Fish Consumption Advisories 5-1
5.3.1 Summary of fish consumption advisories 5-1
5.3.2 Specific descriptions of the fish consumption advisories 5-4
5.3.3 Regulatory criteria and standards 5-8
5.3.4 EPA supplementary fish consumption advisories 5-9
5.4 Exceedences of the Michigan FCA Trigger Levels 5-11
5.4.1 Data sources 5-11
5.4.2 Results 5-14
5.5 Exceedences of the FDA Tolerance Level for PCBs 5-33
5.5.1 Data sources 5-33
5.5.2 Regulatory criteria and standards 5-33
5.5.3 Results 5-34
5.6 Conclusions 5-41

Chapter 6 Injuries to Fish and Aquatic Invertebrates 6-1

6.1 Injury Definitions 6-2
6.2 Stage I Injury Assessment Approach 6-3
6.3 TCDD-Equivalents in Fish Eggs 6-6
6.3.1 Data sources 6-6
6.3.2 Concentrations in eggs causing toxicity 6-7
6.3.3 Results 6-9
6.4 Total PCBs in Smallmouth Bass Livers 6-11
6.4.1 Data sources 6-12
6.4.2 Toxicological benchmarks 6-13
6.4.3 Results 6-14
6.5 Smallmouth Bass Condition, Biochemical, and Histopathological Status 6-17
6.5.1 Data sources 6-17
6.5.2 Summary of results 6-18
6.6 Effects on Benthic Invertebrates 6-22
6.7 Conclusions 6-23

Chapter 7 Injuries to Wildlife 7-1

7.1 Injury Definitions 7-9
7.2 Stage I Injury Assessment Approach 7-10
7.3 Mallard Duck Breast Tissue Concentrations 7-15
7.3.1 Data sources 7-15
7.3.2 Regulatory criteria and standards 7-15
7.3.3 Results 7-16
7.4 PCBs in Bird Diet 7-17
7.4.1 Data sources 7-17
7.4.2 Toxicity reference value derivation 7-18
7.4.3 Results 7-26
7.5 Bald Eagle Reproductive Success 7-29
7.5.1 Data sources 7-29
7.5.2 Benchmarks and injury thresholds 7-30
7.5.3 Results 7-34
7.6 PCB Concentrations in Eggs of Other Birds 7-39
7.6.1 Data sources 7-39
7.6.2 Toxicological benchmarks 7-40
7.6.3 Results 7-43
7.7 Total PCBs in Mammalian Diets 7-49
7.7.1 Data sources 7-49
7.7.2 Toxicological benchmarks 7-51
7.7.3 Results 7-51

7.8 Mink Trapping Success and Total PCBs in Mink Tissue 7-54
7.8.1 Data sources 7-54
7.8.2 Toxicity reference value derivation 7-54
7.8.3 Results 7-55
7.9 Total PCBs in Small Mammal, Shrew, and Muskrat Tissue 7-59
7.9.1 Data sources 7-59
7.9.2 Toxicity reference value derivation 7-60
7.9.3 Results 7-60
7.10 Bioaccumulation of PCBs from Floodplain Soils 7-62
7.10.1 Data sources 7-63
7.10.2 Toxicity reference value derivation 7-64
7.10.3 Results 7-65
7.10.4 Conclusions of the floodplain soil evaluation 7-80
7.11 Conclusions 7-81

Chapter 8 Indirect Injuries 8-1

8.1 Introduction 8-1
8.2 Stage I Assessment Approach 8-2
8.3 No Further Action 8-2
8.4 Access/Deed Restrictions 8-3
8.5 Bank Stabilization 8-3
8.6 Maintaining Dam Structures 8-6
8.7 Sediment Capping 8-7
8.8 Sediment Removal and Disposal 8-9
8.9 Conclusions 8-10

Chapter 9 Conclusions 9-1

References R-1

Figures

1.1 Key features of the Allied Paper, Inc./Portage Creek/Kalamazoo River Superfund Site and surrounding environment 1-4
1.2 Location of current and former PRP paper mill facilities 1-5

2.1 Biphenyl molecular structure 2-1
2.2 Typical fate of paper mill waste during the period when carbonless copy paper contained PCBs 2-5
2.3 The Allied paper mills and surrounding facilities 2-7
2.4 The Georgia-Pacific paper mill and surrounding facilities 2-12
2.5 Simplified flow chart of the fate of Georgia-Pacific paper waste, 1954 to 2000 2-13
2.6 The Simpson Plainwell paper mill and surrounding facilities 2-18
2.7 The Fort James mills and surrounding facilities 2-22
2.8 1994-1995 tributary loadings of PCBs to Lake Michigan 2-24
2.9 Spatial distribution of 1994-1995 tributary loadings of PCBs to Lake Michigan 2-25
2.10 PCB concentrations in surface sediments and floodplain soils from Morrow Lake to the Otsego City Dam 2-28
2.11 PCB concentrations in surface sediments and floodplain soils from the Otsego City Dam to the Lake Allegan Dam 2-29
2.12 PCB concentrations in surface sediments and floodplain soils from the Lake Allegan Dam to the mouth of the Kalamazoo River at Lake Michigan 2-30
2.13 KRE natural resources in which PCB contamination has been documented 2-31
2.14 PCB exposure pathways 2-34

3.1 Locations of reach designations for surface water samples 3-4
3.2 Surface water total PCB concentrations in Portage Creek and the Kalamazoo River from 1985 to 1987 compared to drinking water standards 3-6
3.3 Surface water total PCB concentrations in Portage Creek and the Kalamazoo River from 2000 and 2001 compared to drinking water standards 3-7
3.4 Surface water total PCB concentrations in Portage Creek and the Kalamazoo River from 1985 to 1987 compared to water quality criteria 3-10
3.5 Surface water total PCB concentrations in Portage Creek and the Kalamazoo River in 2000 and 2001 compared to water quality criteria 3-11

4.1 Locations of reach designations for sediment samples 4-4
4.2 Depth-weighted mean PCB concentrations in surficial sediment from cores collected in Portage Creek and the Kalamazoo River compared to consensus-based sediment effect concentrations 4-6
4.3 Depth-weighted mean PCB concentrations in surficial sediment from cores collected in Portage Creek and the Kalamazoo River compared to site-specific TRV for mink 4-10

5.1 History of fish consumption advisories in the Kalamazoo River and Lake Michigan south of Frankfort 5-3
5.2 Reaches of the Kalamazoo River as designated by MDCH for fish consumption advisories, and distribution of RI/FS ABSAs within these reaches 5-13
5.3 Total PCBs in fillets of carp collected upstream of Morrow Dam, 1971-2000 5-15
5.4 Total PCBs in fillets of species not currently under an advisory collected upstream of Morrow Dam, 1971-2000 5-16
5.5 Total PCBs in fillets of carp collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-17
5.6 Total PCBs in fillets of catfish collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-18
5.7 Total PCBs in fillets of largemouth bass collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-19
5.8 Total PCBs in fillets of smallmouth bass collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-20
5.9 Total PCBs in fillets of white sucker collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-21
5.10 Total PCBs in fillets of other species under a general “all other species” advisory collected between Morrow Dam and Lake Allegan Dam, 1971-2000 5-22
5.11 Total PCBs in fillets of carp collected downstream of Lake Allegan Dam, 1971-2000 5-23
5.12 Total PCBs in fillets of catfish collected downstream of Lake Allegan Dam, 1971-2000 5-24
5.13 Total PCBs in fillets of largemouth bass collected downstream of Lake Allegan Dam, 1971-2000 5-25
5.14 Total PCBs in fillets of northern pike collected downstream of Lake Allegan Dam, 1971-2000 5-26
5.15 Total PCBs in fillets of smallmouth bass collected downstream of Lake Allegan Dam, 1971-2000 5-27
5.16 Total PCBs in fillets of trout collected downstream of Lake Allegan Dam, 1971-2000 5-28

5.17 Total PCBs in fillets of walleye collected downstream of Lake Allegan Dam, 1971-2000 5-29
5.18 Total PCBs in fillets of white sucker collected downstream of Lake Allegan Dam, 1971-2000 5-30
5.19 Total PCBs in fillets of yellow perch collected downstream of Lake Allegan Dam, 1971-2000 5-31
5.20 Total PCBs in fillets of other species under a general "all other species" advisory collected downstream of Lake Allegan Dam, 1971-2000 5-32

6.1 Mean and SD of concentrations of estimated total PCBs ww in smallmouth bass livers collected in the Kalamazoo River in 1995 and total PCBs in walleye livers collected in the Lower Fox River and Green Bay, Wisconsin, in 1996 and 1997 6-16

7.1 Wetland types within the former Trowbridge impoundment 7-2
7.2 Total PCBs in whole body fish samples collected in the Kalamazoo River downstream of PRP facilities compared to TRVs for embryomortality effects in birds 7-27
7.3 Total PCBs in whole body fish samples collected in the Kalamazoo River downstream of PRP facilities compared to TRVs for sublethal effects in birds 7-28
7.4 Probability of bald eagles in inland Michigan and Wisconsin and Green Bay producing no young or one or more young in relation to egg PCB concentrations, 1986-1997 7-32
7.5 Total PCB concentrations in bird eggs collected in Allegan State Game Area and other locations along the Kalamazoo River 7-48
7.6 TCDD-eq concentrations based on PCBs and dioxins/furans measured in individual bird eggs in the KRE compared to TRVs for egg embryomortality 7-50
7.7 Total PCBs in whole body fish samples collected in the Kalamazoo River downstream of PRPs compared to TRVs for reproductive effects in mink 7-53
7.8 Total PCB concentrations in mink carcasses collected in reference and assessment locations 7-57
7.9 PCB concentrations in mink livers from reference and assessment locations compared to injury threshold ranges 7-58
7.10 Surface soil PCB concentrations in the former Plainwell impoundment compared to the soil threshold for injury to robins 7-66
7.11 Surface soil PCB concentrations in the former Plainwell impoundment compared to the soil threshold for injury to owls 7-67
7.12 Surface soil PCB concentrations in the former Plainwell impoundment compared to the soil threshold for injury to foxes 7-68
7.13 Surface soil PCB concentrations in the former Plainwell impoundment compared to the soil threshold for injury to mice 7-69

7.14 Surface soil PCB concentrations in the Otsego City impoundment compared to the soil threshold for injury to robins 7-71
7.15 Surface soil PCB concentrations in the Otsego City impoundment compared to the soil threshold for injury to owls 7-72
7.16 Surface soil PCB concentrations in the former Otsego impoundment compared to the soil threshold for injury to robins 7-73
7.17 Surface soil PCB concentrations in the former Otsego impoundment compared to the soil threshold for injury to owls 7-74
7.18 Surface soil PCB concentrations in the former Otsego impoundment compared to the soil threshold for injury to foxes 7-75
7.19 Surface soil PCB concentrations in the former Trowbridge impoundment compared to the soil threshold for injury to robins 7-76
7.20 Surface soil PCB concentrations in the former Trowbridge impoundment compared to the soil threshold for injury to owls 7-77
7.21 Surface soil PCB concentrations in the former Trowbridge impoundment compared to the soil threshold for injury to foxes 7-78
7.22 Surface soil PCB concentrations in the former Trowbridge impoundment compared to the soil threshold for injury to mice 7-79

8.1 Shoreline stabilized with riprap 8-4
8.2 Sheetpile along the Kalamazoo River 8-4
8.3 Cross section of sediment cap over contaminated sediment 8-8
8.4 Mechanical dredge removing sediment 8-9
8.5 Suction-style hydraulic dredge 8-10

Tables

2.1 Demonstrated PCB releases from Allied Facilities 2-8
2.2 PCB concentrations in residuals, sediment, and soil at the Allied Operating Unit 2-10
2.3 PCB concentrations in groundwater beneath Georgia-Pacific disposal areas, 1988-1990 2-15
2.4 PCB concentrations in residuals/soils of Georgia-Pacific disposal areas..... 2-16
2.5 PCB concentrations in Simpson Plainwell Mill effluent 2-19
2.6 PCB concentrations in residuals and soil samples collected at the Simpson Plainwell Mill and the 12th Street Disposal Area, 1987-1994 2-20
2.7 PCBs detected in discharge from Fort James mills 2-22
2.8 Estimated annual PCB load in Portage Creek and locations along the Kalamazoo River, 1994 2-23
2.9 Sediment area, volume, and mass estimates in reaches of the Kalamazoo River and Portage Creek..... 2-26
2.10 Former impoundment size and PCB mass estimates 2-27
2.11 Maximum total PCB concentrations measured in biota in the KRE 2-32

3.1 Approaches to evaluate injury to surface water..... 3-2
3.2 Reach designations for surface water samples 3-3
3.3 Drinking water standards for determining injury to surface water..... 3-5
3.4 Water quality criteria for determining injury to surface water 3-8

4.1 Approaches to evaluate injury to sediment..... 4-2
4.2 Reach designations for sediment samples 4-3
4.3 SECs for invertebrate exposure to PCBs in freshwater sediment..... 4-5
4.4 Exceedences of consensus-based MEC and EEC by depth-weighted mean PCB concentrations in surficial sediment of Portage Creek and the Kalamazoo River..... 4-7
4.5 Site-specific sediment thresholds for the protection of mink 4-9
4.6 Exceedences of site-specific TRV for mink by depth-weighted mean PCB concentrations in surficial sediment of Portage Creek and the Kalamazoo River..... 4-11

5.1 Approaches to evaluate injury to human use and consumption of fish 5-2
5.2 MDCH fish consumption advisories for the Kalamazoo River, 1979-2004..... 5-4
5.3 MDCH fish consumption advisories for Lake Michigan, south of Frankfort, 1977-2004 5-6
5.4 1993 advisory levels recommended by the Great Lakes Fish Consumption Advisory Task Force..... 5-8
5.5 FCA trigger levels used by the State of Michigan since 1977..... 5-9

5.6 Comparison of MDCH and EPA fish consumption advisories for Lake Michigan south of Frankfort, 1997 5-10
5.7 Summary of PCB tissue concentrations and frequency of exceedences of FDA tolerance levels for fish in the Kalamazoo River between Battle Creek and Morrow Dam 5-35
5.8 Summary of PCB tissue concentrations and frequency of exceedences of FDA tolerance levels for fish in the Kalamazoo River between Morrow Dam and Lake Allegan Dam, and in Portage Creek 5-36
5.9 Summary of PCB tissue concentrations and frequency of exceedences of FDA tolerance levels for fish in the Kalamazoo River downstream of Lake Allegan Dam 5-38

6.1 Overview of adverse effects in fish caused by exposure to PCBs 6-3
6.2 Approaches to evaluate injury to aquatic biota 6-5
6.3 Toxic equivalency factors of PCB congeners, PCDDs, and PCDFs relative to TCDD in fish eggs 6-8
6.4 Effect endpoints based on TCDD concentrations in fish eggs 6-9
6.5 TCDD-equivalents of PCBs and PCDDs/PCDFs in fish eggs 6-10
6.6 PCB concentrations in livers of adult fish associated with adverse effects in laboratory studies 6-13
6.7 Mean and SD of concentrations of total PCBs in walleye livers collected in the Lower Fox River and Green Bay, Wisconsin, in 1996 and 1997, and associated prevalence of foci of cellular alteration and hepatic tumors 6-15
6.8 Total PCB concentrations in smallmouth bass livers from D Avenue, Kalamazoo 6-15
6.9 PCB concentrations in liver and fillet of smallmouth bass from the Kalamazoo River, 1995 6-18
6.10 Summary of body condition and biochemical results for KRE smallmouth bass and correlations with tissue PCB concentrations 6-19
6.11 Summary of selected histopathology results of KRE smallmouth bass study, 1995 6-21

7.1 Wildlife species observed in the Kalamazoo River Basin that utilize wetland habitat, and their protection status 7-3
7.2 Overview of adverse effects in birds caused by exposure to PCBs 7-11
7.3 Overview of adverse effects in mammals caused by exposure to PCBs 7-12
7.4 Approaches to evaluate injury to wildlife 7-14
7.5 Total PCB concentrations in breast tissue of mallard ducks, 1988 7-16
7.6 Sublethal effects of PCB exposure to birds 7-19
7.7 Effects of PCB exposure on egg or sperm production 7-21
7.8 Effects of PCB exposure on avian offspring sublethal endpoints 7-22
7.9 Effects of adult PCB exposure on egg hatchability, young fledged, or egg fertility 7-23
7.10 Summary of MDEQ ERA dietary-based HQ results for birds 7-29

7.11 Bald eagle egg toxicity levels identified from comparisons of regional productivities and contaminant concentrations..... 7-31
7.12 Bald eagle productivity in the Kalamazoo River area, 1990-2003..... 7-34
7.13 Comparison of bald eagle nesting success in the Kalamazoo River to other locations..... 7-35
7.14 Concentrations of total PCB and DDE in bald eagle eggs 7-36
7.15 PCB and DDE concentrations in plasma of nestling bald eagles from the Kalamazoo River and other locations in Michigan and Wisconsin..... 7-37
7.16 Egg total PCB concentrations causing adverse effects..... 7-40
7.17 Avian toxic equivalency factors of PCB, dioxin, and furan congeners relative to TCDD..... 7-42
7.18 Bird egg TCDD-eq concentrations causing adverse effects 7-44
7.19 Summary of risks to terrestrial wildlife from the ERA..... 7-52
7.20 Mink trapping success, 1993-1994 7-56
7.21 Total PCB concentrations in small mammals and shrews..... 7-61
7.22 Total PCB concentrations in muskrat livers 7-62
7.23 Total PCB concentrations in whole body muskrat samples collected between 2000 and 2002..... 7-62
7.24 Site-specific soil PCB threshold concentrations for evaluating injury to selected wildlife species..... 7-65
7.25 Summary of Stage I Assessment conclusions regarding toxicological injuries to wildlife..... 7-82

8.1 Remedial options for the Kalamazoo River..... 8-2

9.1 Summary of Stage I Injury Assessment conclusions..... 9-2