



## Status of Lake Sturgeon in the Great Lakes

Indicator #125

### Overall Assessment

Status: **Mixed**  
 Trend: **Improving**  
 Rationale: **There are remnant populations in each basin of the Great Lakes, but few of these populations are large. Much progress has been made in recent years learning about population status in many tributaries. Confirmed observations and captures of lake sturgeon are increasing in all lakes. Stocking is contributing to increased abundance in some areas. There remains a need for information on some remnant spawning populations. Little is known about the juvenile life stage. In many areas habitat restoration is needed because spawning and rearing habitat has been destroyed or altered or access to it has been blocked.**

### Lake-by-Lake Assessment

#### Lake Superior

Status: Mixed  
 Trend: Improving or Undetermined  
 Rationale: Lake sturgeon abundance shows an increasing trend in a few remnant populations and where stocked in the Ontonagon and St. Louis rivers. Lake sturgeon currently reproduce in at least 10 of 22 known historic spawning tributaries.

#### Lake Michigan

Status: Mixed  
 Trend: Improving and Undetermined  
 Rationale: Remnant populations persist in at least nine tributaries having unimpeded connections to Lake Michigan. Successful reproduction has been documented in seven rivers, and abundance has increased in a few rivers in recent years. Active rehabilitation has been initiated through rearing assistance in one remnant population, and reintroductions have been initiated in three rivers.

#### Lake Huron

Status: Mixed  
 Trend: Improving and Undetermined  
 Rationale: Current lake sturgeon spawning activity is limited to five tributaries, four in Georgian Bay and the North Channel and one in Saginaw Bay. Abundant stocks of mixed sizes are consistently captured in the North Channel, Georgian Bay, southern Lake Huron and Saginaw Bay.

#### Lake Erie

Status: Mixed  
 Trend: Improving and Undetermined  
 Rationale: Lakewide incidental catches since 1992 indicate a possible improvement in their status in Lake Erie. Spawning occurs in three known locations in the basin, all located in the connecting waters between Lakes Huron and Erie. The Huron-Erie Corridor supports a robust population of all age classes. The Western basin of Lake Erie, the Detroit River east of Fighting Island, the North Channel of the St. Clair River and Anchor Bay in Lake St. Clair appear to be nursery areas for juveniles and foraging areas for adults.

## Lake Ontario

Status: Mixed

Trend: Improving

Rationale: Lakewide incidental catches since 1995 indicate a possible improvement in their status. Spawning occurs in the Niagara River, Trent River, and possibly the Black River. There are sizeable populations within the St. Lawrence River system. Stocking for restoration began in 1995 in New York.

## Purpose

- To assess the presence and abundance of lake sturgeon in the Great Lakes and their connecting waterways and tributaries
- To infer the health and status of the nearshore benthivore fish community that does, could or should include lake sturgeon

## Ecosystem Objective

Conserve, enhance or rehabilitate self-sustaining populations of lake sturgeon where the species historically occurred and at a level that will permit all state, provincial and federal delistings of classifications that derive from degraded or impaired populations, e.g., threatened, endangered or at risk species. Lake sturgeon is identified as an important species in the Fish Community Goals and Objectives for each of the Great Lakes. Lake Superior has a lake sturgeon rehabilitation plan, and many of the Great Lakes states have lake sturgeon recovery or rehabilitation plans which call for increasing numbers of lake sturgeon beyond current levels.

## State of the Ecosystem


### Background

Lake sturgeon (*Acipenser fulvescens*) were historically abundant in the Great Lakes with spawning populations using many of the major tributaries, connecting waters, and shoal areas across the basin. Prior to European settlement of the region, they were a dominant component of the nearshore benthivore fish community, with populations estimated in the millions in each of the Great Lakes (Baldwin *et al.* 1979). In the mid- to late 1800s, they contributed significantly as a commercial species ranking among the five most abundant species in the commercial catch (Baldwin *et al.* 1979, Fig. 1).

The decline of lake sturgeon populations in the Great Lakes was rapid and commensurate with habitat destruction, degraded water quality, and intensive fishing associated with settlement and development of the region. Sturgeon were initially considered a nuisance species of little value by European settlers, but by the mid-1800s, their value as a commercial species began to be recognized and a lucrative fishery developed. In less than 50 years, their abundance had declined sharply, and since 1900, they have remained a highly depleted species of little consequence to the commercial fishery. Sturgeon is now extirpated from many tributaries and waters where they once spawned and flourished (Figs. 2 and 3). They are considered rare, endangered, threatened, or of watch or special concern status by the various Great Lakes fisheries management agencies. Their harvest is currently prohibited or highly regulated in most waters of the Great Lakes.

## Status of Lake Sturgeon

Efforts continue by many agencies and organizations to gather information on remnant spawning populations in the Great Lakes. Most sturgeon populations continue to sustain themselves at a small fraction of their historical abundance. In many systems, access to spawning habitat has been blocked, and other habitats have been altered. However, there are remnant populations in each basin of the Great Lakes, and some of these populations are large in number (tens of thousands of fish, Fig. 3). Genetic analysis has shown that Great Lakes populations are regionally structured and show significant diversity within and among lakes (DeHaan *et al.* 2006, Welsh *et al.* 2008).



### Lake Superior

The fish community of Lake Superior remains relatively intact in comparison to the other Great Lakes (Bronte *et al.* 2003). Historic and current information indicates that at least 22 Lake Superior tributaries supported spawning lake sturgeon populations (Harkness and Dymond 1961, Auer 2003, Quinlan 2007). Lake sturgeon currently reproduce in at least 10 tributaries. Populations in the Sturgeon River, Michigan, and Bad River, Wisconsin, meet rehabilitation plan criteria for self-sustaining populations (Auer 2003, Auer and Baker 2007, Great Lakes Indian Fish and Wildlife Commission (GLIFWC) and USFWS unpublished data, Quinlan 2007). Improvements in assessment techniques and stocking have resulted in increases in estimated lakewide abundance (Auer and Baker 2007, Schram 2007, and GLIFWC unpublished data). Using hydroacoustic technology, the estimated number of lake sturgeon in the annual spawning run at Sturgeon River, Michigan increased by nearly 100 individuals to range from 350 to 400 adults (Auer and Baker 2007). Genetic analysis has shown that lake sturgeon populations in Lake Superior are significantly different from those in the other Great Lakes (Welsh *et al.* 2008).

Annual assessments were established in key embayments and nearshore waters including Pigeon Bay, Minnesota/Ontario, and in Keweenaw Bay and near the Ontonagon River, Michigan. Habitat (substrate type and water depth) for adult and juvenile fish was geo-referenced and quantified using hydroacoustics in the Kaministiquia River, Ontario (Biberhofer and Prokopec 2005) and Bad River (Cholwek *et al.* 2005). Habitat preference of stocked sturgeon is being studied in the Ontonagon and St. Louis Rivers using radio telemetry (Fillmore 2003, 1854 Authority unpublished data). Due to potential for overexploitation, sport fishing regulations in Ontario waters have been changed to eliminate harvest. There remains a prohibition of commercial harvest of lake sturgeon in Lake Superior. Regulation of recreational and subsistence/home use harvest in Lake Superior varies by agency.

Despite limited progress, challenges remain. Spawning runs are absent in 12 of 22 historic spawning tributaries, and only two populations meet targets identified in the 2003 Rehabilitation Plan. Overall, lake sturgeon abundance remains a small fraction of historical abundance, estimated at 870,000 (Hay-Chmielewski and Whelan 1997) and basic abundance and biological data are unavailable for many stocks.

### Lake Michigan

Sturgeon populations in Lake Michigan continue to sustain themselves at a small fraction of their historical abundance. An optimistic estimate of the lakewide adult abundance is less than 10,000 fish, well below 1% of the most conservative estimates of historic abundance (Hay-Chmielewski and Whelan 1997). Remnant populations currently are known to spawn in waters of at least nine tributaries having unimpeded connections to Lake Michigan (Schneeberger *et al.* 2005, Elliott 2008). Two rivers, the Menominee and Peshtigo, appear to support annual spawning runs of 200 or more adults, and five rivers, the Manistee, Muskegon, Grand, Fox and Oconto, appear to support annual spawning runs of between 25 and 75 adults. Successful reproduction has been documented in all seven of these rivers, and age 0 juveniles can be captured regularly in several of these rivers. Recent recruitment estimates have been made from research efforts in the Peshtigo River indicating that in some years, several hundred fall recruits are produced from that system (Caroffino *et al.* 2007), and research and assessment efforts in the Manistee and Muskegon rivers indicate significant recruitment from those systems as well (Smith, MDNR, personal communication). In addition, abundance of spawners in some rivers appears to have increased in the last decade, indicating that increased recruitment may have been occurring for several years in some rivers. Two other rivers, the Manistique and Kalamazoo, appear to have annual spawning runs of less than 20 fish. Some lake sturgeon have been observed during spawning times in a few other Lake Michigan tributaries such as the St. Joseph and Millecoquins, and near some shoal areas where sturgeon are thought to have spawned historically. It is not known if or how regularly spawning occurs in these systems, and their reproductive status is uncertain. A large self-sustaining population exists in the Lake Winnebago system upstream of the lower Fox River.

Active management in the form of reintroduction stocking and rearing assistance has been implemented in seven Lake Michigan basin tributaries. Commencing in 2005, lake sturgeon have been reared from eggs using streamside rearing facilities and stocked as fingerlings into the Milwaukee, Manitowoc, Cedar, and Whitefish Rivers where sturgeon have been considered extirpated for some time. Over the next 25 years, these reintroductions are intended to rebuild self-sustaining populations that use these rivers to spawn. A streamside rearing facility also is being used to increase the survival of naturally produced larvae in the Manistee River since 2003 (Holtgren *et al.* 2007). Stocking also has been conducted in the upper Menominee River and in the Winnebago system for several years. Though limited recreational harvest is allowed in both the upper Menominee River and the Winnebago system, no harvest is allowed from other Lake Michigan tributaries or from Lake Michigan. Habitat evaluations have been conducted in many sturgeon tributaries within the Lake Michigan basin (Daugherty *et al.* 2009), and improvements in flow conditions and increased planning for fish passage at barriers have the potential to continue to improve habitat conditions in several tributaries.


## Lake Huron

Lake sturgeon populations continue to be well below historical levels. Spawning has been identified in the Garden, Mississauga and Spanish Rivers in the North Channel, in the Nottawasaga River in Georgian Bay and in the Rifle River in Saginaw Bay. Adult spawning populations for each of these river systems are estimated to be in the tens of individuals and are well below rehabilitation targets (Hay-Chmielewski and Whelan 1997, Holey *et al.* 2000). Research in the Saginaw River Watershed in 2005–2007 indicated that lake sturgeon are no longer spawning in that watershed, although sufficient spawning habitat does exist below the Dow Dam on the Tittabawsee River and below the Hamilton Dam on the Flit River. Research is ongoing on the St. Mary's River system and it is unclear if lake sturgeon are using the system for spawning or if the fish are staging there prior to traveling up adjacent tributaries to spawn. The project is ongoing and will continue through 2009. Similar research is being planned for the Rifle River, one of Michigan's last unimpeded rivers. Barriers on Michigan's remaining tributaries to Lake Huron continue to be a major impediment to successful rehabilitation.

Stocks of lake sturgeon in Lake Huron are monitored primarily through the volunteer efforts of commercial fishers cooperating with the various resource management agencies. To date the combined efforts of researchers in U.S. and Canadian waters have resulted in over 6,600 sturgeon tagged in Saginaw Bay, southern Lake Huron, Georgian Bay and the North Channel, with relatively large stocks of mixed sizes being captured at each of these general locations. Tag recoveries and telemetry studies indicate that lake sturgeon are moving within and between jurisdictional boundaries and between lake basins, supporting the need for more cooperative management between the states and between the United States and Canada. As of June 2008, recreational harvest of lake sturgeon in Ontario waters of Lake Huron have been reduced to a zero catch and possession limit and the Ontario Ministry of Natural Resources is moving to a zero harvest limit with the commercial fisheries. Recreational and commercial harvest remains closed in U.S. waters. Traditional use of lake sturgeon by Aboriginal groups in Ontario varies by location and the Ontario Ministry of Natural Resources and Department of Fisheries and Oceans are working with them in an effort to manage those stocks in a sustainable manner.

## Lake Erie

Lake sturgeon populations continue to be well below historical levels with the exception of the stocks located in the connecting waters between Lakes Huron and Erie. Spawning has been identified at two locations in the St. Clair River and at one location in the Detroit River (Manny and Kennedy 2002). Tag recovery data and telemetry research indicate that a robust lake sturgeon stock of approximately 15,000 to 20,000 fish reside in the North Channel of the St. Clair River and Lake St. Clair (Mike Thomas, MDNR, personal communication). The North Channel of the St. Clair River, Anchor Bay in Lake St. Clair, the Detroit River (east of Fighting Island), and the Western basin of Lake Erie have been identified as nursery areas as indicated by consistent catches in commercial and survey fishing gears. In the Central and Eastern basins of Lake Erie, lake sturgeon are scarcer with only occasional catches of sub-adult or adult lake sturgeon in commercial fishing nets and none in research nets. A botulism-related die off in 2001 and 2002 near Buffalo indicate a possible decline in population




abundance of lake sturgeon in the Central and Eastern basin of Lake Erie. However, anglers and divers continue to report sturgeon sightings in the upper Niagara River and into the lake. Survey work conducted in 2005 and 2006 indicated that no lake sturgeon spawning is taking place in the Maumee River (OH) although spawning and nursery habitat would support a reintroduced population (Boase, unpublished data). Research efforts will continue to focus on identifying new spawning locations, genetic difference between stocks, habitat requirements, and migration patterns. In U.S. waters of the St. Clair River and Lake St. Clair, recreational harvest of lake sturgeon is allowed with the following restrictions: only fish between 105 and 125 cm (42-50 in) may be harvested, one fish per season, with an open season from July 16–September 30 and a catch and release season from October 1–November 30. No recreational harvest is allowed in any of the other U.S. or Canadian jurisdictional waters of Lake Erie or the connecting waters (St. Clair and Detroit rivers and Lake St. Clair). The Ontario Ministry of Natural Resources is working to close the one remaining lake sturgeon commercial fishery located in Lake St. Clair which should be implemented by 2009. Traditional use of lake sturgeon by Aboriginal groups in Ontario varies by location and the Ontario Ministry of Natural Resources and Department of Fisheries and Oceans are working with them in an effort to manage those stocks in a sustainable manner.

## Lake Ontario

Lake Ontario has lake sturgeon spawning activity documented in two major tributaries (Niagara River and Trent River) and suspected in at least one more (Black River) on an infrequent basis. There is no targeted assessment of lake sturgeon in Lake Ontario, but incidental catches in research nets have occurred since 1997 (Ontario Ministry of Natural Resources 2004) and 1995 (Eckert 2004), indicating a possible improvement in population status. Age analysis of lake sturgeon captured in the lower Niagara River indicates successful reproduction in the mid-1990s. The New York State Department of Environmental Conservation initiated a stocking program in 1995 to recover lake sturgeon populations. Lake sturgeon has been stocked in the St. Lawrence River and some of its tributaries, inland lakes in New York, and the Genesee River. There are sizeable populations within the St. Lawrence River system, most notably Lac St. Pierre and the Des Prairies and St. Maurice Rivers. However, access is inhibited for many of the historical spawning grounds in tributaries by small dams and within the St. Lawrence River by the Moses-Saunders Dam.

## Pressures

Low numbers or lack of fish (where extirpated) is itself a significant impediment to recovery in many spawning areas. Barriers that prevent lake sturgeon from moving into tributaries to spawn are a major problem. Predation on of eggs and newly hatched lake sturgeon by non-native predators may also be a problem. The genetic structure of remaining populations has been studied by university researchers and fishery managers, and this information will be used to guide future management decisions. With



the collapse of the Caspian Sea sturgeon populations, black market demand for sturgeon caviar could put tremendous pressure on Great Lakes lake sturgeon populations. An additional concern for lake sturgeon in many of the Great Lakes is the ecosystem changes that are resulting from high densities of invasive species such as dreissenid mussels and round gobies, and the presumed related spread of botulism Type E which has produced die-offs of lake sturgeon in most years since 2001.

## Management Implications

Lake sturgeon is an important native species that is listed in the Fish Community Goals and Objectives for all of the Great Lakes. Many of the Great Lakes states and provinces either have or are developing lake sturgeon management plans promoting the need to inventory, protect and restore the species to greater levels of abundance.

While overexploitation removed millions of adult fish, habitat degradation and alteration eliminated traditional spawning grounds. Current work is underway by state, federal, tribal, provincial and private groups to document active spawning sites, assess habitat condition and availability of good habitat, and determine the genetics of remnant Great Lakes lake sturgeon populations.

Several meetings and workshops have been held focusing on identifying the research and assessment needs to further rehabilitation of lake sturgeon in the Great Lakes (Holey *et al.* 2000, Zollweg *et al.* 2003, Quinlan *et al.* 2005, Boase *et al.* 2008) and a significant amount of research and assessment directed towards these needs has occurred in the last 10 years. Among these is the research to better define the genetic structuring of Great Lakes lake sturgeon populations, and genetics-based rehabilitation plans are being developed to help guide reintroduction and rehabilitation efforts being implemented across the Great Lakes. Research into new fish passage technologies that will allow safe upstream and downstream passage around barriers to migration also have been underway for several years. Many groups are continuing to work to identify current lake sturgeon spawning locations in the Great Lakes, and studies are being initiated to identify habitat preferences and recruitment levels for juvenile lake sturgeon (ages 0 to 2). Several agencies are also working in concert on reintroduction and rearing assistance programs to strengthen and reintroduce lake sturgeon into various waters where populations are lacking or at risk from further declines.

## Comments from the author(s)

Research and development are needed to determine ways for lake sturgeon to pass man-made barriers on rivers. In addition, there are significant, legal, logistical, and financial hurdles to overcome in order to restore degraded spawning habitats in connecting waterways and tributaries to the Great Lakes. More monitoring is needed to determine the current status of Great Lakes lake sturgeon populations, particularly the juvenile life stage. Cooperative efforts between law enforcement and fishery managers are required as world pressure on sturgeon stocks will result in the need to protect large adult lake sturgeon in the Great Lakes.

## Assessing Data Quality

Data Characteristics	Strongly Agree	Agree	Neutral or Unknown	Disagree	Strongly Disagree	Not Applicable
1. Data are documented, validated, or quality-assured by a recognized agency or organization		X				
2. Data are traceable to original sources	X					
3. The source of the data is a known, reliable and respected generator of data	X					
4. Geographic coverage and scale of data are appropriate to the Great Lakes basin	X					
5. Data obtained from sources within the U.S. are comparable to those from Canada		X				
6. Uncertainty and variability in the data are documented and within acceptable limits for this indicator report		X				
Clarifying Notes:						

## Acknowledgments

Authors: (2008)

Robert Elliott, U.S. Fish and Wildlife Service (USFWS), Green Bay National Fish and Wildlife Conservation Office, New Franken, WI 54229

Henry Quinlan, U.S. Fish and Wildlife Service (USFWS), Ashland National Fish and Wildlife Conservation Office, Ashland, WI 54806

James Boase, U.S. Fish and Wildlife Service (USFWS), Alpena National Fish and Wildlife Conservation Office, Alpena, MI, 49707

Betsy Trometer, U.S. Fish and Wildlife Service (USFWS), Lower Great Lakes National Fish and Wildlife Conservation Office, Amherst, NY 14228

## Sources

- Auer, N.A. ed., 2003. *A lake sturgeon rehabilitation plan for Lake Superior*. Great Lakes Fishery Commission Misc. Publ. 2003-02.
- Auer, N.A., and Baker, E.A. 2007. Assessment of lake sturgeon spawning stocks using fixed-location, split-beam sonar technology. *J. Applied Ichth.* Vol. 23:113-121.
- Baldwin, N.S., Saalfeld, R.W., Ross, M.A., and Buettner, H.J. 1979. *Commercial fish production in the Great Lakes 1867-1977*. Great Lakes Fishery Commission Technical Report 3.
- Biberhofer, J., and Prokopec, C.M. 2005. *Description and quantification of the submerged aquatic substrates in the lower Kaministiquia River*. Draft Report. National Water Resource Institute, Environment Canada.
- Boase, J., Elliott, R., Quinlan, H., and Trometer, B. 2005. Proceedings of the third Great Lakes lake sturgeon coordination meeting, November 29-30, 2006. Sault Ste. Marie, MI.
- Bronte, C.R., Ebener, M.P., Schreiner, D.R., DeVault, D.S., Petzold, M.M., Jensen, D.A., Richards, C., and Lozano, S.J. 2003. Fish community changes in Lake Superior, 1970-2000. *Can. J. Fish. Aquat. Sci.* 60:1552-1574.
- Caroffino, D.C., Sutton, T.M., Elliott, R.F., and Donofrio, M.C. 2007. *Abundance and mortality of early life stages of lake sturgeon in the Peshtigo River, Wisconsin*. 68<sup>th</sup> Midwest Fish and Wildlife Conference, Madison, WI. December 9-12, 2007.
- Cholwek, G., Yule, D., Eitrem, M., Quinlan, H., and Doolittle, T. 2005. *Mapping potential lake sturgeon habitat in the lower Bad River complex*. U.S.G.S. Lake Superior Biol. Station report. 21 pp.
- DeHaan, P., Libants, S., Elliott, R.F., and Scribner, K.T. 2006. Genetic population structure of remnant lake sturgeon populations in the upper Great Lakes basin. *Trans. Amer. Fish. Soc.* 135:1478-1492.
- Daugherty, D. J., Sutton, T.M. and Elliott, R.F. 2009. Suitability modeling of lake sturgeon habitat in five northern Lake Michigan tributaries: implications for population rehabilitation. *Rest. Ecol.* 17(2):245-257.
- Eckert, T.H. 2004. Summary of 1976-2003 Warm Water Assessment. In *New York State Department of Environmental Conservation. Lake Ontario Annual Report 2003*. Bureau of Fisheries, Lake Ontario Unit and St. Lawrence River Unit. Cape Vincent and Watertown, NY.
- Elliott, R.F. Status and trends in lake sturgeon. In D.F. Clapp, and W. Horns eds., 2008. *The state of Lake Michigan in 2005*. Great Lakes Fish. Comm. Spec. Pub.
- Harkness, W.J., and Dymond, J.R. 1961. *The lake sturgeon: The history of its fishery and problems of conservation*. Ontario Dept. of Lands and Forests, Fish and Wildl. Branch. 120 pp.
- Hay-Chmielewski, E.M., and Whelan, G.E. 1997. *Lake sturgeon rehabilitation strategy*. Michigan Department of Natural Resources Fisheries Division: Special Report Number 18, Ann Arbor, MI.
- Holey, M.E., Baker, E.A., Thuemler, T.F., and Elliott, R.F. 2000. *Research and assessment needs to restore Lake Sturgeon in the Great Lakes: results of a workshop sponsored by the Great Lakes Fishery Trust*. Lansing, MI.

Holtgren, J.M., Paquet, A.J., and Fajfer, S. 2007. Design of a portable streamside rearing facility for lake sturgeon. *N. Amer. J. Aquacult.* 69:317–323.

Manny, B.A., and Kennedy, G.W. 2002. Known lake sturgeon (*Acipenser fulvescens*) spawning habitat in the channel between Lakes Huron and Erie in the Laurentian Great Lakes. *J. Appl. Ichth.* 18:486-490.

Ontario Ministry of Natural Resources. 2004. *Lake Ontario Fish Communities and Fisheries: 2003 Annual Report of the Lake Ontario Management Unit*. Ontario Ministry of Natural Resources, Picton, ON.

Quinlan, H. 2007. Lake sturgeon. In *The state of Lake Superior in 2000*. ed Mark P. Ebener. Great Lakes Fish. Comm. Spec. Pub. 07-02, pp. 29-32.

Quinlan, H., Elliott, R., Zollweg, E., Bryson, D., Boase, J., and Weisser, J. 2005. Proceedings of the second Great Lakes lake sturgeon coordination meeting, November 9-10, 2004. Sault Ste. Marie, MI.

Schram, S.T. 2007. *Dispersal of stocked lake sturgeon in Wisconsin waters of Lake Superior*. Wis. Dept. of Nat. Res. Fish Manage. Report No. 152. October 2007.

Schneeberger, P.J., Elliott, R.F., Jonas, J.L., and Hart, S. 2005. Benthivores. In *The state of Lake Michigan in 2000*. M.E. Holey, and T.N. Trudeau, eds., Great Lakes Fish. Comm. Spec. Pub. 05-01, pp. 25-32.

Thomas, M.V., and Haas, R.C. 2002. Abundance, age structure, and spatial distribution of lake sturgeon, *Acipenser fulvescens*, in the St. Clair system. *J. Appl. Ichth.* 18:495-501.

U.S. Fish and Wildlife Service. Ashland Fishery Resource Office, USFWS, 2800 Lake Shore Drive, Ashland, Wisconsin, 54806, unpublished data.

Welsh, A., Hill, T., Quinlan, H., Robinson, C., May, B. 2008. Genetic assessment of lake sturgeon population structure in the Laurentian Great Lakes. *N. Amer. J. of Fish. Manage.* Vol. 28: 572-591.

Zollweg, E.C., Elliott, R.F., Hill, T.D., Quinlan, H.R., Trometer, E., and Weisser, J.W. eds., 2003. Great Lakes Lake Sturgeon Coordination Meeting. In Proceedings of the December 11-12, 2002 Workshop, Sault Ste. Marie, MI.

## **Last Updated**

*State of the Great Lakes 2009*