

**Table 4: Northern Rocky Mountain States minimum fall wolf population and breeding pairs,\* 1979-2001**

Minimum fall wolf population:

Year:	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01
<u>Recovery Area:</u>																							
NW MONTANA	2	1	2	8	6	6	13	15	10	14	12	33	29	41	55	48	66	70	56	49	63	64	84
YELLOWSTONE																	21	40	86	112	118	177	218
CENTRAL IDAHO																	14	42	71	114	141	192	261
TOTAL	2	1	2	8	6	6	13	15	10	14	12	33	29	41	55	48	101	152	213	275	322	433	563

Breeding pairs:

Year:	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01
<u>Recovery Area:</u>																							
NW MONTANA								1	2	1	1	3	2	4	4	5	6	7	5	5	6	6	7
YELLOWSTONE																	2	4	9	6	8	14	13
CENTRAL IDAHO																		3	6	10	10	10	14
TOTAL								1	2	1	1	3	2	4	4	5	8	14	20	21	24	30	34

\* By the standards of the Rocky Mountain gray wolf recovery plan and wolf reintroduction environmental impact statement, a breeding pair is defined as an adult male and an adult female wolf, accompanied by 2 pups that survived at least until Dec 31. Recovery goals call for 10 breeding pairs per area, or a total of 30 breeding pair distributed through the 3 areas, for 3 years.

Breeding pair count was increased from 23 to 24 for the year 1999, after it was discovered, in 2000, that the Boulder Pack (NWMT) had bred and raised pups the previous year.

Breeding pair count was increased from 28 to 30 for the year 2000, after it was discovered, in 2001, that the Gravelly Pack (GYE) and the Gold Fork Pack (CID) had bred and raised pups the previous year.