

STATUS OF THE DESERT TORTOISE AND ITS CRITICAL HABITAT

Status of the Desert Tortoise

The Service listed the desert tortoise as threatened in 1990 [55 Federal Register (FR) 12178]. The threats described in the listing rule and both recovery plans (Service 1994, 2011) continue to affect the species. The most apparent threats to the desert tortoise are those that result in mortality and permanent habitat loss across large areas, such as urbanization and large-scale renewable energy projects, and those that fragment and degrade habitats, such as proliferation of roads and highways, off-highway vehicle activity, and habitat invasion by non-native invasive plant species.

We remain unable to quantify how threats affect desert tortoise populations. The assessment of the original recovery plan emphasized the need for a better understanding of the implications of multiple, simultaneous threats facing desert tortoise populations and of the relative contribution of multiple threats on demographic factors (i.e., birth rate, survivorship, fecundity, and death rate; Tracy *et al.* 2004).


In recognition of the absence of specific and recent information on the location of habitable areas of the Mojave Desert, especially at the outer edges of this area, Nussear *et al.* (2009) developed a quantitative, spatial habitat model for the desert tortoise north and west of the Colorado River. The model incorporates environmental variables such as precipitation, geology, vegetation, and slope and is based on occurrence data of desert tortoises from sources spanning more than 80 years, including data from the 2001 to 2008 range-wide monitoring surveys. The model predicts the relative potential for desert tortoises to be present in any given location, given the combination of habitat variables at that location in relation to areas of known occupancy throughout the range; calculations of the amount of desert tortoise habitat in the 5-year review (Service 2010) and in this biological opinion use a threshold of 0.5 or greater predicted value for potential desert tortoise habitat. The model does not account for anthropogenic effects to habitat and represents the potential for occupancy by desert tortoises absent these effects.

To understand better the relationship of threats to populations of desert tortoises and the most effective manner to implement recovery actions, the Desert Tortoise Recovery Office developed a spatial decision support system that models the interrelationships of threats to desert tortoises and how those threats affect population change. The spatial decision support system describes the numerous threats that desert tortoises face, explains how these threats interact to affect individual animals and habitat, and how these effects in turn bring about changes in populations. For example, we have long known that the construction of a transmission line can result in the death of desert tortoises and loss of habitat. We have also known that common ravens, known predators of desert tortoises, use transmission line pylons for nesting, roosting, and perching and that the access routes associated with transmission lines provide a vector for the introduction and spread of invasive weeds and facilitate increased human access into an area. Increased human access can accelerate illegal collection and release of desert tortoises and their deliberate maiming and killing, as well as facilitate the spread of other threats associated with human presence, such as vehicle use, garbage and dumping, and invasive plants (Service 2011). Changes in the abundance of native plants because of invasive weeds can compromise the physiological health of desert tortoises, making them more vulnerable to drought, disease, and predation. The spatial decision support system allows us to map threats across the range of the desert tortoise and model the intensity of stresses that these multiple and combined threats place on desert tortoise populations.

The following map depicts the 12 critical habitat units of the desert tortoise, linkages between conservation areas for the desert tortoise and the aggregate stress that multiple, synergistic threats place on desert tortoise populations, as modeled by the spatial decision support system. Conservation areas include designated critical habitat and other lands managed for the long-term conservation of the desert tortoise (e.g., the Desert Tortoise Natural Area, Joshua Tree National Park, and the Desert National Wildlife Refuge).

Recovery Plan

The Service (1994, 2011) has issued an initial recovery plan and a revised recovery plan for the desert tortoise. The 1994 recovery plan recommended that a scientifically credible monitoring plan be developed to determine that the population exhibit a statistically significant upward trend or remain stationary for at least 25 years and that enough habitat would be protected within a recovery unit or the habitat and populations be managed intensively enough to ensure long-term viability. Because both minimum population densities and minimum population numbers need to be considered to ensure recovery, the Service further recommended that reserves be at least 1,000 square miles. Smaller reserves that provide high-quality, secure habitat for 10,000 to 20,000 adult desert tortoises should provide comfortable persistence probabilities for the species well into the future when populations are well above minimum viable density (e.g., 30 or more adults per square mile) and lambdas can be maintained (see page C54 of Service 1994). Conversely, populations with densities below approximately 10 adults per square mile (3.9 per square kilometer) are in danger of extinction (see page 32 of Service 1994).



“Adult” desert tortoise connotes reproductive maturity. Desert tortoises may become reproductive

at various sizes. The Service based its 2010 survey protocol on the methodology used in range-wide sampling but erred in citing 160 millimeters as the size below which surveyors' ability to detect desert tortoises decreases. In range-wide sampling, the Service uses 180 millimeters as its cut-off length for counting desert tortoises, at least in part because the Styrofoam models used for training are 180 millimeters in length. The Service intends to revise the survey protocol and will use 180 millimeters in the revised version. For the purposes of this biological opinion, we have used the term "adult" to indicate reproductive status and "large" to indicate animals larger than 180 millimeters to conform to the Service's protocols for range-wide sampling and pre-project surveys.

The revised recovery plan for the desert tortoise (Service 2011) lists three objectives and associated criteria to achieve delisting. The first objective is to maintain self-sustaining populations of desert tortoises within each recovery unit into the future; the criterion is that the rates of population change (λ) for desert tortoises are increasing (i.e., $\lambda > 1$) over at least 25 years (i.e., a single generation), as measured by extensive, range-wide monitoring across conservation areas within each recovery unit, and by direct monitoring and estimation of vital rates (recruitment, survival) from demographic study areas within each recovery unit.

The second objective addresses the distribution of desert tortoises. The goal is to maintain well-distributed populations of desert tortoises throughout each recovery unit; the criterion is that the distribution of desert tortoises throughout each conservation area increase over at least 25 years.

The final objective is to ensure that habitat within each recovery unit is protected and managed to support long-term viability of desert tortoise populations. The criterion is that the quantity of desert tortoise habitat within each conservation area be maintained with no net loss until population viability is ensured.

The revised recovery plan (Service 2011) also recommends connecting blocks of desert tortoise habitat, such as critical habitat units and other important areas to maintain gene flow between populations. Linkages defined using least-cost path analysis (Averill-Murray *et al.* 2013) illustrate a minimum connection of habitat for desert tortoises between blocks of habitat and represent priority areas for conservation of population connectivity. The previous map in this biological opinion illustrates that, across the range, desert tortoises in areas under the highest level of conservation and management remain subject to numerous threats, stresses, and mortality sources.

Five-Year Review

Section 4(c)(2) of the Endangered Species Act requires the Service to conduct a status review of each listed species at least once every 5 years. The purpose of a 5-year review is to evaluate whether the species' status has changed since it was listed (or since the most recent 5-year review); these reviews, at the time of their completion, provide the most up-to-date information on the range-wide status of the species. For this reason, we are appending the 5-year review of the status of the desert tortoise (Appendix C; Service 2010) to this biological opinion and are incorporating it by reference to provide most of the information needed for this section of the biological opinion. The following paragraphs provide a summary of the relevant information in the 5-year review.

In the 5-year review, the Service discusses the status of the desert tortoise as a single distinct population segment and provides information on the Federal Register notices that resulted in its listing and the designation of critical habitat. The Service also describes the desert tortoise's

ecology, life history, spatial distribution, abundance, habitats, and the threats that led to its listing (i.e., the five-factor analysis required by section 4(a)(1) of the Endangered Species Act). In the 5-year review, the Service concluded by recommending that the status of the desert tortoise as a threatened species be maintained.

With regard to the status of the desert tortoise as a distinct population segment, the Service concluded in the 5-year review that the recovery units recognized in the original and revised recovery plans

(Service 1994 and 2011, respectively) do not qualify as distinct population segments under the Service's distinct population segment policy (61 FR 4722; February 7, 1996). We reached this conclusion because individuals of the listed taxon occupy habitat that is relatively continuously distributed, exhibit genetic differentiation that is consistent with isolation-by-distance in a continuous-distribution model of gene flow, and likely vary in behavioral and physiological characteristics across the area they occupy as a result of the transitional nature of, or environmental gradations between, the described subdivisions of the Mojave and Colorado deserts.

In the 5-year review, the Service summarizes information with regard to the desert tortoise's ecology and life history. Of key importance to assessing threats to the species and to developing and implementing a strategy for recovery is that desert tortoises are long lived, require up to 20 years to reach sexual maturity, and have low reproductive rates during a long period of reproductive potential. The number of eggs that a female desert tortoise can produce in a season is dependent on a variety of factors including environment, habitat, availability of forage and drinking water, and physiological condition. Predation seems to play an important role in clutch failure. Predation and environmental factors also affect the survival of hatchlings. The Service notes in the 5-year review that the combination of the desert tortoise's late breeding age and a low reproductive rate challenges our ability to recover the species.

Since the completion of the 5-year review, the Service has issued several biological opinions that affect large areas of desert tortoise habitat because of numerous proposals to develop renewable energy within its range. These biological opinions concluded that proposed solar plants were not likely to jeopardize the continued existence of the desert tortoise primarily because they were located outside of critical habitat and desert wildlife management areas that contain most of the land base required for the recovery of the species. The proposed actions also included numerous measures intended to protect desert tortoise during the construction of the projects, such as translocation of affected individuals. In aggregate, these projects would result in an overall loss of approximately 48,041 acres of habitat of the desert tortoise. We also predicted that the project areas supported up to 4,363 desert tortoises; we concluded that most of these individuals were small desert tortoises, that most large individuals would likely be translocated from project sites, and that most mortalities would be small desert tortoises that were not detected during clearance surveys. To date, 637 desert tortoises have been observed during construction of projects; most of these individuals were translocated from work areas, although some desert tortoises have been killed (see Appendix D). The mitigation required by the Bureau and California Energy Commission, the agencies permitting these facilities, resulted in the acquisition of private land and funding for the implementation of various actions that are intended to promote the recovery of the desert tortoise. These mitigation measures are consistent with recommendations in the recovery plans for the desert tortoise; many of the measures have been derived directly from the recovery plans and the Service supports their implementation. We expect that, based on the best available scientific information, they will result in conservation benefits to the desert tortoise; however, it is

difficult to assess how desert tortoise populations will respond because of the long generation time of the species.

In August 2016, the Service (2016) issued a biological opinion to the Bureau for the land use plan amendment under the Desert Renewable Energy Conservation Plan. The land use plan amendment addressed all aspects of the Bureau's management of the California Desert Conservation Area; however, the Service and Bureau agreed that only those aspects related to the construction, operation, maintenance, and decommissioning of renewable energy facilities were likely to adversely affect the desert tortoise. The land use plan amendment resulted in the designation of approximately 388,000 acres of development focus areas where the Bureau would apply a streamlined review process to applications for projects that generate renewable energy; the Bureau estimated that approximately 11,290 acres of modeled desert tortoise habitat within the development focus areas would eventually be developed for renewable energy. The Bureau also adopted numerous conservation and management actions as part of the land use plan amendment to further reduce the adverse effects of renewable energy development on the desert tortoise.

The land use plan amendment also increased the amount of land that the Bureau manages for conservation (e.g., areas of critical environmental concern, National Conservation Lands, etc.) from 6,118,135 to 8,689,669 acres (Bureau 2015a); not all of the areas subject to increased protection are within desert tortoise habitat. The Bureau will also manage lands outside of development focus areas according to numerous conservation and management actions; these conservation and management actions are more protective of desert tortoises than direction contained in the previous land use plan. The Service (2016) concluded that the land use plan amendment was not likely to jeopardize the continued existence of the desert tortoise and would benefit its recovery.

In addition to the biological opinions issued for solar development within the range of the desert tortoise, the Service (2012) also issued a biological opinion to the Department of the Army (Army) for the use of additional training lands at Fort Irwin. As part of this proposed action, the Army translocated approximately 650 adult desert tortoises from 18,197 acres of the southern area of Fort Irwin, which had been off-limits to training, to lands south of the base that are managed by the Bureau and the Army. The Army would also use an additional 48,629 acres that lie east of the former boundaries of Fort Irwin; much of this parcel is either too mountainous or too rocky and low in elevation to support numerous desert tortoises.

The Service also issued a biological opinion to the Department of the Navy (Navy) that considered the effects of the expansion of the Marine Corps Air Ground Combat Center at Twentynine Palms (Service 2017). We concluded that the Navy's proposed action, the use of approximately 167,982 acres of public and private land for training, was not likely to jeopardize the continued existence of the desert tortoise. Most of the expansion area lies within the Johnson Valley Off-highway Vehicle Management Area. As part of this proposed action, the Navy translocated 978 adult desert tortoises from the expansion area to four recipient sites to the north and east of the expansion area (Henen 2018). The Lucerne-Ord and Siberia sites are entirely within Bureau-managed lands, and the Rodman-Sunshine Peak North and Cleghorn sites overlap Bureau-managed lands and lands managed by the Navy. The Lucerne-Ord site lies within the Ord-Rodman desert tortoise critical habitat unit.

The incremental effect of the larger actions (i.e., solar development, the expansions of Fort Irwin and the Marine Corps Air Ground Combat Center) on the desert tortoise is unlikely to be positive,

despite the numerous conservation measures that have been (or will be) implemented as part of the actions. The tortoises that were translocated by the Navy from the Johnson Valley Off-highway Vehicle Management Area were moved into populations that were below the Service's established minimum viable density, to attempt to augment these populations and make them more viable in the long-term. The acquisition of private lands as mitigation for most of these actions increases the level of protection afforded these lands; however, these acquisitions do not create new habitat and Federal, State, and privately managed lands remain subject to most of the threats and stresses we discussed previously in this section. Although land managers have been implementing measures to manage these threats and we expect, based on the best available scientific information, that such measures provide conservation benefits to the desert tortoise, we have been unable, to date, to determine whether the expected benefits of the measures have yet been realized, at least in part because of the low reproductive capacity of the desert tortoise. Therefore, the conversion of habitat into areas that are unsuitable for this species continues the trend of constricting the desert tortoise into a smaller portion of its range.

The Service also issued a biological opinion to the Marine Corps that considered the effects of the expansion of the Marine Corps Air Ground Combat Center at Twentynine Palms (Service 2017). We concluded that the Marine Corps' proposed action, the use of approximately 167,982 acres of public and private land for training, was not likely to jeopardize the continued existence of the desert tortoise. Most of the expansion area lies within the Johnson Valley Off-highway Vehicle Management Area. As part of this proposed action, the Marine Corps removed 929 desert tortoises from the expansion area (Hoffmann 2017).

The incremental effect of the larger actions (i.e., solar development, the expansions of Fort Irwin and the Marine Corps Air Ground Combat Center) on the desert tortoise is unlikely to be positive, despite the numerous conservation measures that have been (or will be) implemented as part of the actions. The acquisition of private lands as mitigation for most of these actions increases the level of protection afforded these lands; however, these acquisitions do not create new habitat and Federal, State, and privately managed lands remain subject to most of the threats and stresses we discussed previously in this section. Although land managers have been implementing measures to manage these threats and we expect, based on the best available scientific information, that such measures provide conservation benefits to the desert tortoise, we have been unable, to date, to determine whether the expected benefits of the measures have yet been realized, at least in part because of the low reproductive capacity of the desert tortoise. Therefore, the conversion of habitat into areas that are unsuitable for this species continues the trend of constricting the desert tortoise into a smaller portion of its range.

As the Service notes in the 5-year review (Service 2010), "(t)he threats identified in the original listing rule continue to affect the (desert tortoise) today, with invasive species, wildfire, and renewable energy development coming to the forefront as important factors in habitat loss and conversion. The vast majority of threats to the desert tortoise or its habitat are associated with human land uses." Oftedal's work (2002 in Service 2010) suggests that invasive weeds may adversely affect the physiological health of desert tortoises. Using captive neonate and yearling desert tortoises, Drake *et al.* (2016) found that individuals "eating native forbs had better body condition and immune functions, grew more, and had higher survival rates (>95%) than (desert) tortoises consuming any other diet"; health and body condition declined in individuals fed only grasses (native or non-native). Current information indicates that invasive species likely affect a large portion of the desert tortoise's range. Furthermore, high densities of weedy species increase the likelihood of wildfires; wildfires, in turn, destroy native species and further the spread of

invasive weeds.

Drake *et al.* (2015) “compared movement patterns, home-range size, behavior, microhabitat use, reproduction, and survival for adult desert tortoises located in, and adjacent to, burned habitat” in Nevada. They noted that the fires killed many desert tortoises but found that, in the first 5 years post- fire, individuals moved deeper into burned habitat on a seasonal basis and foraged more frequently in burned areas (corresponding with greater production of annual plants and herbaceous perennials in these areas). Production of annual plants upon which desert tortoises feed was 10 times greater in burned versus unburned areas but was dominated by non-native species [e.g., red brome (*Bromus rubens*)] that frequently have lower digestibility than native vegetation. During years six and seven, the movements of desert tortoises into burned areas contracted with a decline in the live cover of a perennial forage plant that rapidly colonizes burned areas. Drake *et al.* (2015) did not find any differences in health or survivorship for desert tortoises occupying either habitat (burned or unburned) during this study or in reproduction during the seventh year after the fire.

Climate change is likely to affect the prospects for the long-term conservation of the desert tortoise. For example, predictions for climate change within the range of the desert tortoise suggest more frequent and/or prolonged droughts with an increase of the annual mean temperature by 3.5 to 4.0 degrees Celsius. The greatest increases will likely occur in summer [June-July-August mean increase of as much as 5 degrees Celsius (Christensen *et al.* 2007 in Service 2010)]. Precipitation will likely decrease by 5 to 15 percent annually in the region; with winter precipitation decreasing by up to 20 percent and summer precipitation increasing by up to 5 percent. Because germination of the desert tortoise’s food plants is highly dependent on cool- season rains, the forage base could be reduced due to increasing temperatures and decreasing precipitation in winter. Although drought occurs routinely in the Mojave Desert, extended periods of drought have the potential to affect desert tortoises and their habitats through physiological effects to individuals (i.e., stress) and limited forage availability. To place the consequences of long-term drought in perspective, Longshore *et al.* (2003) demonstrated that even short-term drought could result in elevated levels of mortality of desert tortoises. Therefore, long-term drought is likely to have even greater effects, particularly given that the current fragmented nature of desert tortoise habitat (e.g., urban and agricultural development, highways, freeways, military training areas, etc.) will make recolonization of extirpated areas difficult, if not impossible.

Core Criteria for the Jeopardy Determination

When determining whether a proposed action is likely to jeopardize the continued existence of a species, we are required to consider whether the action would “reasonably be expected, directly or indirectly, to reduce appreciably the likelihood of both the survival and recovery of a listed species in the wild by reducing the reproduction, numbers, or distribution of that species” (50 CFR 402.02). Although the Service does not explicitly address these metrics in the 5-year review, we have used the information in that document and more recent information to summarize the status of the desert tortoise with respect to its reproduction, numbers, and distribution.

Reproduction

In the 5-year review, the Service notes that desert tortoises increase their reproduction in high rainfall years; more rain provides desert tortoises with more high quality food (i.e., plants that are higher in water and protein), which, in turn, allows them to lay more eggs. Conversely, the physiological stress associated with foraging on food plants with insufficient water and nitrogen

may leave desert tortoises vulnerable to disease (Ofstedal 2002 in Service 2010), and the reproductive rate of diseased desert tortoises is likely lower than that of healthy animals. Young desert tortoises also rely upon high- quality, low-fiber plants (e.g., native annual plants) with nutrient levels not found in the invasive weeds that have increased in abundance across its range (Ofstedal *et al.* 2002; Tracy *et al.* 2004). Compromised nutrition of young desert tortoises likely represents an effective reduction in reproduction by reducing the number of animals that reaches adulthood; see previous information from Drake *et al.* (2016). Consequently, although we do not have quantitative data that show a direct relationship, the abundance of weedy species within the range of the desert tortoise has the potential to affect the reproduction of desert tortoises and recruitment into the adult population in a negative manner.

Various human activities have introduced numerous species of non-native invasive plants into the California desert. Routes that humans use to travel through the desert (paved and unpaved roads, railroads, motorcycle trails, etc.) serve as pathways for new species to enter habitat of the desert tortoise and for species that currently occur there to spread. Other disturbances of the desert substrate also provide invasive species with entry points into the desert. The following map depicts the potential for these species to invade habitat of the desert tortoise. The reproductive capacity of the desert tortoise may be compromised to some degree by the abundance and distribution of invasive weeds across its range; the continued increase in human access across the desert likely continues to facilitate the spread of weeds and further affect the reproductive capacity of the species.

Numbers

In the 5-year review, the Service discusses various means by which researchers have attempted to determine the abundance of desert tortoises and the strengths and weaknesses of those methods. Due to differences in area covered and especially to the non-representative nature of earlier sample sites, data gathered by the Service's current range-wide monitoring program cannot be reliably compared to information gathered through other means at this time.



Data from small-scale study plots (e.g., 1 square mile) established as early as 1976 and surveyed primarily through the mid-1990s indicate that localized population declines occurred at many sites across the desert tortoise’s range, especially in the western Mojave Desert; spatial analyses of more widespread surveys also found evidence of relatively high mortality in some parts of the range (Tracy *et al.* 2004). Although population densities from the local study plots cannot be extrapolated to provide an estimate of the number of desert tortoises on a range-wide basis, historical densities in some parts of the desert exceeded 100 adults in a square mile (Tracy *et al.* 2004). The Service (2010) concluded that “appreciable declines at the local level in many areas, which coupled with other survey results, suggest that declines may have occurred more broadly.”

The range-wide monitoring that the Service initiated in 2001 is the first comprehensive attempt to determine the densities of desert tortoises in conservation areas across their range. The Desert Tortoise Recovery Office (Service 2015a) used annual density estimates obtained from this sampling effort to evaluate range-wide trends in the density of desert tortoises over time. (All references to the density of desert tortoises are averages. Some areas support higher densities and some lower; desert tortoises are not distributed in uniform densities across large areas.) This analysis indicates that densities in the Northeastern Mojave Recovery Unit have increased since 2004, with the increase apparently resulting from increased survival of adults and sub-adults moving into the adult size class. The analysis also indicates that the populations in the other four

recovery units are declining; the following table depicts the estimated numbers of desert tortoises within conservation areas in each recovery unit and the rates of population change. Surveys did not include the steepest slopes in these desert tortoise conservation areas; however, the model developed by Nussear *et al.* (2009) generally rates steep slopes as less likely to support desert tortoises. Densities in the Joshua Tree and Piute Valley conservation areas within the Colorado Desert Recovery Unit seem to be increasing, although densities in the recovery unit as a whole continue to decline.

Recovery Units	2004	2014	Change	Percentage of Change
Western Mojave	35,777	17,644	-18,133	-51
Colorado Desert	67,087	42,770	-24,317	-36
Northeastern Mojave	4,920	18,220	+13,300	+270
Eastern Mojave	16,165	5,292	-10,873	-67
Upper Virgin River	2,397	1,760	-637	-27
Total	126,346	85,686	-40,660	-32

In the previous summary of the results of range-wide sampling (Service 2014), we extrapolated the densities obtained within conservation areas (e.g., desert wildlife management area, Desert Tortoise Research Natural Area, Joshua Tree National Park) to all modeled habitat of the desert tortoise. This extrapolation may have exaggerated the number of desert tortoises because we applied the values for areas where densities are generally highest (i.e., the conservation areas) to areas where desert tortoises exist in very low densities (e.g., the Antelope Valley). We are also aware of a few areas where the density of desert tortoises outside of conservation areas is higher than inside.

To further examine the status of desert tortoise populations over time, we compared the densities of desert tortoises in the Western Mojave Recovery Unit between 2004 and 2014 (see Service 2015a). In 2004, desert tortoise conservation areas surveyed in the Western Mojave Recovery Unit supported an average density of approximately 5.7 adults per square kilometer (14.8 per square mile). In contrast, surveys in the same areas in 2014 indicated that densities had decreased to 2.8 adults per square kilometer (7.3 per square mile). This decline in densities is consistent with decreases in density of populations in all recovery units over the same time period, with the exception of the Northeastern Mojave Recovery Unit. In fact, historical survey data from numerous plots in the Western Mojave Recovery Unit during the late 1970s and early 1980s suggest that adult desert tortoise densities ranged from 50 to 150 per square mile (Tracy *et al.* 2004).

To further assess the status of the desert tortoise, the Desert Tortoise Recovery Office (Service 2015) used multi-year trends from the best-fitting model describing loge-transformed density of adult animals per square kilometer. In 2014, 3 of the 5 recovery units supported densities below 3.9 adult animals per square kilometer [Western Mojave (2.8), Eastern Mojave (1.5), and Colorado Desert (3.7); see table 10 in Service 2015b], which is the minimum density recommended to avoid extinction in the 1994 recovery plan. The Northeastern Mojave Recovery Unit supported 4.4 adult desert tortoises per square kilometer and the Upper Virgin River Recovery Unit, which is by far the smallest recovery unit, supported 15.3 adults per square kilometer.

Allison (2014) evaluated changes in size distribution of desert tortoises since 2001. In the Western Mojave and Colorado Desert recovery units, the relative number of juveniles to adults indicates that juvenile numbers are declining faster than adults. In the Eastern Mojave, the number of

juvenile desert tortoises is also declining, but not as rapidly as the number of adults. In the Upper Virgin River Recovery Unit, trends in juvenile numbers are similar to those of adults; in the Northeastern Mojave Recovery Unit, the number of juveniles is increasing, but not as rapidly as are adult numbers in that recovery unit. Juvenile numbers, like adult densities, are responding in a directional way, with increasing, stable, or decreasing trends, depending on the recovery unit where they are found.

In this context, we consider “juvenile” desert tortoises to be animals smaller than 180 millimeters in length. The Service does not include juveniles detected during range-wide sampling in density estimations because they are more difficult to detect and surveyors frequently do not observe them during sampling. However, this systematic range-wide sampling provides us with an opportunity to compare the proportion of juveniles to adults observed between years.

Distribution

Prior to 1994, desert tortoises were extirpated from large areas within their distributional limits by urban and agricultural development (e.g., the cities of Barstow and Lancaster, California; Las Vegas, Nevada; and St. George, Utah; etc.; agricultural areas south of Edwards Air Force Base and east of Barstow), military training (e.g., Fort Irwin, Leach Lake Gunnery Range), and off-road vehicle use (e.g., portions of off-road management areas managed by the Bureau and unauthorized use in areas such as east of California City, California).

Since 1994, urban development around Las Vegas has likely been the largest contributor to habitat loss throughout the range. Desert tortoises have been essentially removed from the 18,197-acre southern expansion area at Fort Irwin (Service 2012). The development of large solar facilities has also reduced the amount of habitat available to desert tortoises. No solar facilities have been developed within desert tortoise conservation areas, such as desert wildlife management areas, although such projects have occurred in areas that the Service considers important linkages between conservation areas (e.g., Silver State South Project in Nevada).

The following table depicts acreages of habitat (as modeled by Nussear *et al.* 2009, using only areas with a probability of occupancy by desert tortoises greater than 0.5 as potential habitat) within the recovery units of the desert tortoise and of impervious surfaces as of 2006 (Fry *et al.* 2011); calculations are by Darst (2014). Impervious surfaces include paved and developed areas and other disturbed areas that have zero probability of supporting desert tortoises. All units are in acres.

Recovery Units	Modeled Habitat	Impervious Surfaces (percentage)	Remaining Modeled Habitat
Western Mojave	7,585,312	1,989,843 (26)	5,595,469
Colorado Desert	4,950,225	510,862 (10)	4,439,363
Northeastern Mojave	3,012,293	386,182 (13)	2,626,111
Eastern Mojave	4,763,123	825,274 (17)	3,937,849
Upper Virgin River	231,460	84,404 (36)	147,056
Total	20,542,413	3,796,565 (18)	16,745,848

The Service (2010) concluded, in its 5-year review, that the distribution of the desert tortoise has not changed substantially since the publication of the original recovery plan in 1994 in terms of the

overall extent of its range. Since 2010, we again conclude that the species' distribution has not changed substantially in terms of the overall extent of its range, although desert tortoises have been removed from several thousand acres because of solar development and military activities.

Status of Critical Habitat of the Desert Tortoise

The Service designated critical habitat for the desert tortoise in portions of California, Nevada, Arizona, and Utah in a final rule published February 8, 1994 (59 FR 5820). The Service designates critical habitat to identify the key biological and physical needs of the species and key areas for recovery and to focus conservation actions on those areas. Critical habitat is composed of specific geographic areas that contain the biological and physical features essential to the species' conservation and that may require special management considerations or protection. These features, which include space, food, water, nutrition, cover, shelter, reproductive sites, and special habitats, are called the physical and biological features of critical habitat. The specific physical and biological features of desert tortoise critical habitat are: sufficient space to support viable populations within each of the six recovery units and to provide for movement, dispersal, and gene flow; sufficient quality and quantity of forage species and the proper soil conditions to provide for the growth of these species; suitable substrates for burrowing, nesting, and overwintering; burrows, caliche caves, and other shelter sites; sufficient vegetation for shelter from temperature extremes and predators; and habitat protected from disturbance and human-caused mortality.

Critical habitat of the desert tortoise would not be able to fulfill its conservation role without each of the physical and biological features being functional. As examples, critical habitat would not function properly if a sufficient amount of forage species were present but human-caused mortality was excessive; an area with sufficient space to support viable populations within each of the six recovery units and to provide for movement, dispersal, and gene flow would not function properly without adequate forage species.

The final rule for designation of critical habitat did not explicitly ascribe specific conservation roles or functions to the various critical habitat units. Rather, it refers to the strategy of establishing recovery units and desert wildlife management areas recommended by the recovery plan for the desert tortoise, which had been published as a draft at the time of the designation of critical habitat, to capture the "biotic and abiotic variability found in desert tortoise habitat" (59 FR 5820, see page 5823). Specifically, we designated the critical habitat units to follow the direction provided by the draft recovery plan (Service 1993) for the establishment of desert wildlife management areas. The critical habitat units in aggregate are intended to protect the variability that occurs across the large range of the desert tortoise; the loss of any specific unit would compromise the ability of critical habitat as a whole to serve its intended function and conservation role.

Despite the fact that desert tortoises do not necessarily need to move between critical habitat units to complete their life histories, both the original and revised recovery plans discuss the importance of these critical habitat units and connectivity between them for the recovery of the species. Although it determined that linkages between critical habitat units did not meet the definition of critical habitat, the Service (1994) recommended the identification of buffer zones and linkages for smaller desert tortoise conservation areas to aid in overall recovery efforts; however, land management agencies have generally not established such areas.

We did not designate the Desert Tortoise Natural Area and Joshua Tree National Park in California and the Desert National Wildlife Refuge in Nevada as critical habitat because they are

“primarily managed as natural ecosystems” (59 FR 5820, see page 5825) and provide adequate protection to desert tortoises. Since the designation of critical habitat, Congress increased the size of Joshua Tree National Park and created the Mojave National Preserve. A portion of the expanded boundary of Joshua Tree National Park lies within critical habitat of the desert tortoise; portions of other critical habitat units lie within the boundaries of the Mojave National Preserve.

Within each critical habitat unit, both natural and anthropogenic factors affect the function of the physical and biological features of critical habitat. As an example of a natural factor, in some specific areas within the boundaries of critical habitat, such as within and adjacent to dry lakes, some of the physical and biological features are naturally absent because the substrate is extremely salty; desert tortoises do not normally reside in such areas. Comparing the acreage of desert tortoise habitat as depicted by Nussear *et al.*'s (2009) model to the gross acreage of the critical habitat units demonstrates quantitatively that the entire area within the boundaries of critical habitat likely does not support the physical and biological features. In the following table, the acreage for modeled habitat is for the area in which the probability that desert tortoises are present is greater than 0.5. The acreages of modeled habitat do not include loss of habitat due to human-caused impacts. The difference between gross acreage and modeled habitat is 653,214 acres; that is, approximately 10 percent of the gross acreage of the designated critical habitat is not considered modeled habitat. All units are acres.

Critical Habitat Unit	Gross Acreage	Modeled Habitat
Superior-Cronese	766,900	724,967
Fremont-Kramer	518,000	501,095
Ord-Rodman	253,200	184,155
Pinto Mountain	171,700	144,056
Piute-Eldorado	970,600	930,008
Ivanpah Valley	632,400	510,711
Chuckwalla	1,020,600	809,319
Chemehuevi	937,400	914,505
Gold Butte-Pakoon	488,300	418,189
Mormon Mesa	427,900	407,041
Beaver Dam Slope	204,600	202,499
Upper Virgin River	54,600	46,441
Totals	6,446,200	5,792,986

Human activities can have obvious or more subtle effects on the physical and biological features of critical habitat. The grading of an area and subsequent construction of a building removes physical and biological features; this action has an obvious effect on critical habitat. The revised recovery plan identifies human activities such as urbanization and the proliferation of roads and highways as threats to the desert tortoise and its habitat; these threats are examples of activities that have a clear effect on the physical and biological features of critical habitat.

Condition of the Physical and Biological Features of Critical Habitat

We have included the following paragraphs from the revised recovery plan for the desert tortoise (Service 2011) to demonstrate that other anthropogenic factors affect the physical and biological features of critical habitat in more subtle ways. All references are in the revised recovery plan (i.e., in Service 2011); we have omitted some information from the revised recovery plan where the

level of detail was unnecessary for the current discussion.

Surface disturbance from [off-highway vehicle] activity can cause erosion and large amounts of dust to be discharged into the air. Recent studies on surface dust impacts on gas exchanges in Mojave Desert shrubs showed that plants encrusted by dust have reduced photosynthesis and decreased water-use efficiency, which may decrease primary production during seasons when photosynthesis occurs (Sharifi *et al.* 1997). Sharifi *et al.* (1997) also showed reduction in maximum leaf conductance, transpiration, and water-use efficiency due to dust. Leaf and stem temperatures were also shown to be higher in plants with leaf-surface dust. These effects may also impact desert annuals, an important food source for [desert] tortoises.

[Off-highway vehicle] activity can also disturb fragile cyanobacterial-lichen soil crusts, a dominant source of nitrogen in desert ecosystems (Belnap 1996). Belnap (1996) showed that anthropogenic surface disturbances may have serious implications for nitrogen budgets in cold desert ecosystems, and this may also hold true for the hot deserts that [desert] tortoises occupy. Soil crusts also appear to be an important source of water for plants, as crusts were shown to have 53 percent greater volumetric water content than bare soils during the late fall when winter annuals are becoming established (DeFalco *et al.* 2001). DeFalco *et al.* (2001) found that non-native plant species comprised greater shoot biomass on crusted soils than native species, which demonstrates their ability to exploit available nutrient and water resources. Once the soil crusts are disturbed, non-native plants may colonize, become established, and out-compete native perennial and annual plant species (D'Antonio and Vitousek 1992; DeFalco *et al.* 2001). Invasion of non-native plants can affect the quality and quantity of plant foods available to desert tortoises. Increased presence of invasive plants can also contribute to increased fire frequency.

Proliferation of invasive plants is increasing in the Mojave and Sonoran deserts and is recognized as a substantial threat to desert tortoise habitat. Many species of non-native plants from Europe and Asia have become common to abundant in some areas, particularly where disturbance has occurred and is ongoing. As non-native plant species become established, native perennial and annual plant species may decrease, diminish, or die out (D'Antonio and Vitousek 1992). Land managers and field scientists identified 116 species of non-native plants in the Mojave and Colorado deserts (Brooks and Esque 2002).

Increased levels of atmospheric pollution and nitrogen deposition related to increased human presence and combustion of fossil fuels can cause increased levels of soil nitrogen, which in turn may result in significant changes in plant communities (Aber *et al.* 1989). Many of the non-native annual plant taxa in the Mojave region evolved in more fertile Mediterranean regions and benefit from increased levels of soil nitrogen, which gives them a competitive edge over native annuals. Studies at three sites within the central, southern, and western Mojave Desert indicated that increased levels of soil nitrogen can increase the dominance of non-native annual plants and promote the invasion of new species in desert regions. Furthermore, increased dominance by non-native annuals may decrease the diversity of native annual plants, and increased biomass of non-native annual grasses may increase fire frequency (Brooks 2003).

This summary from the revised recovery plan (Service 2011) demonstrates how the effects of human activities on habitat of the desert tortoise are interconnected. In general, surface disturbance causes increased rates of erosion and generation of dust. Increased erosion alters additional habitat outside of the area directly affected by altering the nature of the substrate, removing shrubs, and possibly destroying burrows and other shelter sites. Increased dust affects photosynthesis in the

plants that provide cover and forage to desert tortoises. Disturbed substrates and increased atmospheric nitrogen enhance the likelihood that invasive species will become established and out-compete native species; the proliferation of weedy species increases the risk of large-scale fires, which further move habitat conditions away from those that are favorable to desert tortoises.

The following paragraphs generally describe how the threats described in the revised recovery plan affect the physical and biological features of critical habitat of the desert tortoise.

Sufficient space to support viable populations within each of the six recovery units and to provide for movement, dispersal, and gene flow.

Urban and agricultural development, concentrated use by off-road vehicles, and other activities such as development of transmission lines and pipelines completely remove habitat. Although we are aware of local areas within the boundaries of critical habitat that have been heavily disturbed, we do not know of any areas that have been disturbed to the intensity and extent that the function of this physical and biological feature has been compromised. To date, the largest single loss of critical habitat is the use of 18,197 acres of additional training land in the southern portion of Fort Irwin.

The widening of existing freeways likely caused the second largest loss of critical habitat. Despite these losses of critical habitat, which occur in a linear manner, the critical habitat units continue to support sufficient space to support viable populations within each of the six recovery units.

In some cases, major roads likely disrupt the movement, dispersal, and gene flow of desert tortoises. State Route 58 and Highway 395 in the Fremont-Kramer Critical Habitat Unit, Fort Irwin Road in the Superior-Cronese Critical Habitat Unit, and Interstate 10 in the Chuckwalla Critical Habitat Unit are examples of large and heavily travelled roads that likely disrupt movement, dispersal, and gene flow. Roads that have been fenced and provided with underpasses may alleviate this fragmentation to some degree; however, such facilities have not been in place for sufficient time to determine whether they will eliminate fragmentation.

The threats of invasive plant species described in the revised recovery plan generally do not result in the removal of this physical and biological feature because they do not convert habitat into impervious surfaces, as would urban development.

Sufficient quality and quantity of forage species and the proper soil conditions to provide for the growth of these species.

This physical and biological feature addresses the ability of critical habitat to provide adequate nutrition to desert tortoises. As described in the revised recovery plan and 5-year review, grazing, historical fire, invasive plants, altered hydrology, drought, wildfire potential, fugitive dust, and climate change/temperature extremes contribute to the stress of “nutritional compromise.” Paved and unpaved roads through critical habitat of the desert tortoise provide avenues by which invasive native species disperse; these legal routes also provide the means by which unauthorized use occurs over large areas of critical habitat. Nitrogen deposition from atmospheric pollution likely occurs throughout all the critical habitat units and exacerbates the effects of the disturbance of substrates. Because paved and unpaved roads are so widespread through critical habitat, this threat has diminished the value of critical habitat for conservation of the desert tortoise throughout its range, to some degree. See the Status of the Desert Tortoise section of this biological opinion for a

map that depicts the routes by which invasive weeds have access to critical habitat; the routes shown on the map are a subset of the actual number of routes that cross critical habitat of the desert tortoise.

Suitable substrates for burrowing, nesting, and overwintering.

Surface disturbance, motor vehicles traveling off route, use of off-highway vehicles management areas, off-highway vehicles events, unpaved roads, grazing, historical fire, wildfire potential, altered hydrology, and climate change leading to shifts in habitat composition and location, storms, and flooding can alter substrates to the extent that they are no longer suitable for burrowing, nesting, and overwintering. Erosion caused by these activities can alter washes to the extent that desert tortoise burrows placed along the edge of a wash, which is a preferred location for burrows, could be destroyed. We expect that the area within critical habitat that is affected by off-road vehicle use to the extent that substrates are no longer suitable is relatively small in relation to the area that desert tortoises have available for burrowing, nesting, and overwintering; consequently, off-road vehicle use has not had a substantial effect on this physical and biological feature.

Most livestock allotments have been eliminated from within the boundaries of critical habitat. Of those that remain, livestock would compact substrates to the extent that they would become unsuitable for burrowing, nesting, and overwintering only in areas of concentrated use, such as around watering areas and corrals. Because livestock grazing occurs over a relatively small portion of critical habitat and the substrates in most areas within livestock allotments would not be substantially affected, suitable substrates for burrowing, nesting, and overwintering remain throughout most of the critical habitat units.

Burrows, caliche caves, and other shelter sites.

Human-caused effects to burrows, caliche caves, and other shelter sites likely occur at a similar rate as effects to substrates for burrowing, nesting, and overwintering for the same general reasons. Consequently, sufficient burrows, caliche caves, and other shelter sites remain in the critical habitat units.

Sufficient vegetation for shelter from temperature extremes and predators.

In general, sufficient vegetation for shelter from temperature extremes and predators remains throughout critical habitat. In areas where large fires have occurred in critical habitat, many of the shrubs that provide shelter from temperature extremes and predators have been destroyed; in such areas, cover sites may be a limiting factor. The proliferation of invasive plants poses a threat to shrub cover throughout critical habitat as the potential for larger and more frequent wildfires increases.

In 2005, wildfires in Nevada, Utah, and Arizona burned extensive areas of critical habitat (Service 2010). Although different agencies report slightly different acreages, the following table provides an indication of the scale of the fires.

Critical Habitat Unit	Total Area Burned (acres)	Percent of the Critical Habitat Unit Burned
Beaver Dam Slope	53,528	26

Gold-Butte Pakoon	65,339	13
Mormon Mesa	12,952	3
Upper Virgin River	10,557	19

The revised recovery plan notes that the fires caused statistically significant losses of perennial plant cover, although patches of unburned shrubs remained. Given the patchiness with which the physical and biological features of critical habitat are distributed across the critical habitat units and the varying intensity of the wildfires, we cannot quantify precisely the extent to which these fires disrupted the function and value of the critical habitat.

Habitat protected from disturbance and human-caused mortality.

In general, the Federal agencies that manage lands within the boundaries of critical habitat have adopted land management plans that include implementation of some or all of the recommendations contained in the original recovery plan for the desert tortoise (see pages 70 to 72 of Service 2010). To at least some degree, the adoption of these plans has resulted in the implementation of management actions that are likely to reduce the disturbance and human-caused mortality of desert tortoises. For example, these plans resulted in the designation of open routes of travel and the closure (and, in some cases, physical closure) of unauthorized routes. Numerous livestock allotments have been relinquished by the permittees and cattle no longer graze these allotments. Because of these planning efforts, the Bureau has proposed the withdrawal of some areas of critical habitat from mineral entry (79 FR 51190; the withdrawal of 10,094.03 acres of public lands within the Superior-Cronese Critical Habitat Unit). Because of actions on the part of various agencies, many miles of highways and other paved roads have been fenced to prevent desert tortoises from wandering into traffic and being killed. The Service and other agencies of the Desert Managers Group in California are implementing a plan to remove common ravens that prey on desert tortoises and to undertake other actions that would reduce subsidies (i.e., food, water, sites for nesting, roosting, and perching, etc.) that facilitate their abundance in the California Desert (Service 2008). The Bureau’s (2016) land use plan amendment for the Desert Renewable Energy Conservation Plan increased the amount of land under protective status and adopted conservation and management actions that furthered the Bureau’s goals for these areas.

Despite the implementation of these actions, disturbance and human-caused mortality continue to occur in many areas of critical habitat (which overlap the Bureau’s areas of critical environmental concern for the most part and are the management units for which most data are collected) to the extent that the value of critical habitat for the conservation of the desert tortoise is, to some degree, diminished. For example, many highways and other paved roads in California remain unfenced. Hughson and Darby (2011) noted that as many as 10 desert tortoises are reported killed annually on paved roads within Mojave National Preserve. Because carcasses on roads are quickly removed by scavengers or destroyed by other vehicles, we expect that far more desert tortoises are killed on roads than are reported.

Unauthorized off-road vehicle use continues to disturb habitat and result in loss of vegetation within the boundaries of critical habitat; although we have not documented the death of desert tortoises as a direct result of this activity, it likely occurs. Additionally, the habitat disturbance caused by this unauthorized activity exacerbates the spread of invasive plants, which displace native plants that are important forage for the desert tortoise, thereby increasing the physiological stress faced by desert tortoises.

Finally, the Bureau will not allow the development of renewable energy facilities on public lands within the boundaries of areas of critical environmental concern and National Conservation Lands (which largely correspond to the boundaries of critical habitat). Counties have not specifically restricted the development of renewable energy facilities on private lands within the boundaries of areas of critical environmental concern and National Conservation Lands. However, the checkerboard pattern of land ownership would likely necessitate that the Bureau consider issuance of a right-of-way for such a facility, which likely decreases the potential for such proposals in the future.

Summary of the Status of Critical Habitat of the Desert Tortoise

As noted in the 5-year review and revised recovery plan for the desert tortoise (Service 2010, 2011), critical habitat of the desert tortoise is subject to landscape-level impacts in addition to the site-specific effects of individual human activities. On the landscape level, atmospheric pollution is increasing the level of nitrogen in desert substrates; the increased nitrogen exacerbates the spread of invasive plants, which outcompete the native plants necessary for desert tortoises to survive. As invasive plants increase in abundance, the threat of large wildfires increases; wildfires have the potential to convert the shrubland-native annual plant communities upon which desert tortoises depend to a community with fewer shrubs and more invasive plants. In such a community, shelter and forage would be more difficult for desert tortoises to find.

Invasive plants have already compromised the value of critical habitat for the conservation of the desert tortoise to some degree with regard to the second physical and biological feature (i.e., sufficient quality and quantity of forage species and the proper soil conditions to provide for the growth of these species). These effects likely extend to the entirety of critical habitat; given the numerous routes by which invasive plants can access critical habitat and the large spatial extent that is subject to nitrogen from atmospheric pollution.

The value of critical habitat has been diminished to some degree with regard to the last physical and biological feature (i.e., habitat protected from disturbance and human-caused mortality) as a result of the wide variety of human activities that continues to occur within its boundaries. These effects result from the implementation of discrete human activities and are thus more site-specific in nature.

Although the remaining physical and biological features have been affected to some degree by human activities, these impacts have not, to date, appreciably diminished the value of the critical habitat units for the conservation of the desert tortoise. We have reached this conclusion primarily because the effects are localized and thus do not affect the value of large areas of critical habitat for the conservation of the desert tortoise.

Land managers have undertaken actions to improve the status of critical habitat. For example, as part of its efforts to offset the effects of the use of additional training maneuver lands at Fort Irwin (Service 2004), the Department of the Army acquired the private interests in the Harper Lake and Cronese Lakes allotments, which are located within critical habitat in the Western Mojave Recovery Unit; as a result, cattle have been removed from these allotments. Livestock have been removed from numerous other allotments through various means throughout the range of the desert tortoise. The retirement of allotments assists in the recovery of the species by eliminating disturbance to the physical and biological features of critical habitat by cattle and range improvements.

APPENDIX D

Solar projects for which the U.S. Fish and Wildlife Service has issued biological opinions or incidental take permits.

The following table summarizes information regarding the solar projects that have undergone formal consultation with regard to the desert tortoise. In the Citations column, a single reference indicates that the acres of desert tortoise habitat and number of desert tortoises are estimates from the biological opinion; when the column includes two citations, the first is for the acreage of habitat and the estimated number of desert tortoises from the biological opinion and the second is for number of desert tortoises that were found onsite prior to or during construction.

Project and Recovery Unit	Acres of Desert Tortoise Habitat	Desert Tortoises Estimated₁	Desert Tortoises Observed₂	Citations³
Eastern Mojave				
Ivanpah Solar Electric Generating System	3,582	1,136	175 ⁷	Service 2011a, Davis 2014
Stateline	1,685	947	55	Service 2013a, Ironwood 2014
Silver State North – NV	685	14 ⁶	4	Service 2010a, Cota 2013
Silver State South – NV	2,427 ⁴	1,020 ⁴	152	Service 2013a, Cota 2014
Amargosa Farm Road – NV	4,350	4 ⁶	-	Service 2010e
Nevada Solar One - NV	400	5 ⁵	5 ⁵	Burroughs 2012, 2014
Copper Mountain North - NV	1,504	10 ⁵	3 ⁵	Service 2011b, 2013b; NewFields Environmental and Engineering, LLC 2014
Copper Mountain - NV	380	5 ⁵	5 ⁵	Burroughs 2012, 2014
Townsite - NV	905	4 ⁸	- ⁵	Service 2014a
Techren Boulder City - NV	2,291	15 ⁹	- ⁵	Service 2012a
Valley Electric Association - NV	80	4	4 ¹⁰	Service 2015a
Western Mojave				
Abengoa Harper Lake	Primarily in abandoned agricultural fields	4 ⁶	-	Service 2011c
Chevron Lucerne Valley	516	10	-	Service 2010b
Cinco	500	53	2	Service 2015b, Daitch 2015
Soda Mountain	1,726	78	-	Service 2015c
Northeastern Mojave				
Res Americas Moapa Solar Energy Center - NV	954	95	-	Service 2014b
Moapa K Road Solar	2,141	186	157	Service 2012b, Burroughs 2013
Playa Solar	1,760	258	77	Service 2015d, Ironwood Consulting 2016

Invenergy Harry Allen Solar	594	242	-	Service 2015d
NV Energy Dry Lake Solar Energy Center	751	45	-	Service 2015d
NV Energy Dry Lake Solar Energy Center at Harry Allen	55	15	-	Service 2015d
Aiya Solar	672	91	-	Service 2015e
Mountainview	146	5	5	Wise 2018
Colorado				
Genesis	1,774	8	0	Service 2010c, Fraser 2014a
Blythe	6,958	30	0	Service 2010d, Fraser 2014b
Desert Sunlight	4,004	56	7	Service 2011d, Fraser 2104a
McCoy	4,533	15	0	Service 2013c, Fraser 2014b
Desert Harvest	1,300	5	-	Service 2013d
Rice	1,368	18	1	Service 2011e, Fraser 2014a
Total	48,041	4,363	637	

¹The numbers in this column are not necessarily comparable because the methodologies for estimating the numbers of desert tortoises occasionally vary between projects. When available, we included an estimate of the numbers of small desert tortoises.

²This column reflects the numbers of desert tortoises observed within project areas. It includes translocated animals and those that were killed by project activities. Project activities may result in the deaths of more desert tortoises than are found. Dashes represent projects for which we have no information at this point; some projects had not broken ground at the time of this biological opinion.

³The first citation in this column is for both the acreage and the estimate of the number of desert tortoises. The second is for the number of desert tortoises observed during construction of the project; where only one citation is present, construction has not begun or data are unavailable at this time.

⁴These numbers include Southern California Edison's Primm Substation and its ancillary facilities.

⁵These projects occurred under the Clark County Multi-species Habitat Conservation Plan; the provisions of the habitat conservation plan do not require the removal of desert tortoises. We estimate that all six projects combined will affect fewer than 50 desert tortoises.

⁶These estimates do not include smaller desert tortoises.

⁷In the table attached to the electronic mail, the number of desert tortoises translocated from the project site is represented by the total number of translocated animals minus the number of animals born in the holding pens.

⁸The estimate of the number of desert tortoises is from the portion of the project on WAPA land (20.39 acres). The remaining lands are covered by the Clark County Multi-species Habitat Conservation Plan; see footnote 5.

⁹The estimate of the number of desert tortoises is from both Bureau (104 acres) and private (2,200 acres) land. The remaining lands are covered by the Clark County Multi-species Habitat Conservation Plan; see footnote 5.

¹⁰Of the 80-acre project site, 76.4 acres were left intact (there was crushing and mowing of vegetation but no blading) with openings along the bottom of the fence. After project completion, four tortoises were released back into the solar facility on September 25, 2017. One adult has left and re-entered the facility twice and the one juvenile has remained within the facility.

LITERATURE CITED

- Burroughs, M. 2012. Electronic mail. Information on solar projects in desert tortoise habitat in Nevada for which the Service has issued biological opinions. Dated April 26. Fish and wildlife biologist, Southern Nevada Fish and Wildlife Office, U.S. Fish and Wildlife Service. Las Vegas, Nevada.
- Burroughs, M. 2013. Electronic mail. Comments on the draft biological opinion for the Stateline and Silver State Solar South projects, San Bernardino County, California, and Clark County, Nevada (Stateline: 2800(P), CACA-048669, CAD090.01; Silver State South: 6840 (NV-052)) (Stateline: 8-8-13-F-43; Silver State South: 84320-2010-F-0208- R003). Dated September 23. Fish and wildlife biologist, Southern Nevada Fish and Wildlife Office, U.S. Fish and Wildlife Service. Las Vegas, Nevada.
- Burroughs, M. 2014. Electronic mails. Status of solar projects in Nevada. Dated January 27. Fish and wildlife biologist, Southern Nevada Fish and Wildlife Office, U.S. Fish and Wildlife Service. Las Vegas, Nevada.
- Cota, M. 2013. Electronic mail. Comments on the draft biological opinion for the Stateline and Silver State Solar South projects, San Bernardino County, California, and Clark County, Nevada (Stateline: 2800(P), CACA-048669, CAD090.01; Silver State South: 6840 (NV-052)) (Stateline: 8-8-13-F-43; Silver State South: 84320-2010-F-0208-R003). Dated September 18. Wildlife biologist, Pahrump Field Office, Bureau of Land Management. Las Vegas, Nevada.
- Cota, M. 2014. Electronic mail. Number of desert tortoises found on the Silver State South Project site. Dated November 25. Wildlife biologist, Pahrump Field Office, Bureau of Land Management. Las Vegas, Nevada.
- Daitch, D. 2015. Electronic mail. Notifications of desert tortoises found on the Cinco Solar project. Dated November 6. Rincon Consultants. Monterey, California.
- Davis, D. 2014. Electronic mail. ISEGS master tortoise list, October 2014. Dated November 3. Environmental specialist III, Ivanpah Solar Thermal, Nipton, California.
- Fraser, J. 2014a. Electronic mail. Number of desert tortoises found on the Genesis and Desert Sunlight solar sites. Dated January 28. Fish and wildlife biologist, Palm Springs Fish and Wildlife Office, U.S. Fish and Wildlife Service. Palm Springs, California.
- Fraser, J. 2014b. Electronic mail. Number of desert tortoises found on the Blythe and McCoy solar sites. Dated November 5. Fish and wildlife biologist, Palm Springs Fish and Wildlife Office, U.S. Fish and Wildlife Service. Palm Springs, California.
- Ironwood Consulting, Inc. 2014. Annual compliance report: April - December 2014. Biological resources. Stateline Solar Farm. Bureau of Land Management case file number: CACA-48669. San Bernardino County, California.
- Ironwood Consulting, Inc. 2016. Desert tortoise 2016 third quarter and final report. Playa Solar Project, Clark County, Nevada. File No. 84320-2015-F-0139. Las Vegas, Nevada.

- NewFields Environmental and Engineering, LLC. 2014. Field Contact Representative and Biological Monitoring Post Construction Report for the Copper Mountain Solar 3 Gen-Tie Line and Access Road. Dated July 2014. Prepared for Bureau of Land Management and U.S. Fish and Wildlife Service. Prepared by NewFields, Las Vegas, Nevada.
- U.S. Fish and Wildlife Service. 2010a. Formal consultation for the Silver State Solar Project (NextLight Renewable Power, LLC), Clark County, Nevada. File No. 84320-2010-F- 0208. Dated September 16. Memorandum to Field Manager, Pahrump Field Office, Bureau of Land Management, Las Vegas, Nevada. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.
- U.S. Fish and Wildlife Service. 2010b. Revised biological opinion for the Lucerne Valley Chevron Solar Project, San Bernardino County, California (3031 (P) CA-680.33) (8-8-10-F- 61R). Memorandum to Field Manager, Barstow Field Office, Bureau of Land Management, Barstow, California. Dated September 29. From Field Supervisor, Ventura Fish and Wildlife Office. Ventura, California.
- U.S. Fish and Wildlife Service. 2010c. Biological opinion on the Genesis Solar Energy Project, Riverside County, California. Memorandum to Field Manager, Palm Springs South Coast Field Office, Bureau of Land Management, Palm Springs, California. Dated November 2. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.
- U.S. Fish and Wildlife Service. 2010d. Biological opinion on the Blythe Solar Power Plant, Riverside County, California. Memorandum to Field Manager, Palm Springs South Coast Field Office, Bureau of Land Management, Palm Springs, California. Dated October 8. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.
- U.S. Fish and Wildlife Service. 2010e. Formal and informal consultation under section 7 of the Endangered Species Act for the Amargosa Farm Road Solar Energy Project, Nye County, Nevada. File nos. 84320-2010-F-0315 and 84320-2010-1-0316. Memorandum to Field Manager, Pahrump Field Office, Bureau of Land Management, Las Vegas, Nevada. Dated November 1. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.
- U.S. Fish and Wildlife Service. 2011a. Biological opinion on BrightSource Energy's Ivanpah Solar Electric Generating System Project, San Bernardino County, California [CACA- 48668, 49502, 49503, 49504] (8-8-10-F-24R). Dated June 10. Memorandum to District Manager, California Desert District, Bureau of Land Management, Moreno Valley, California. From Field Supervisor, Ventura Fish and Wildlife Office. Ventura, California.
- U.S. Fish and Wildlife Service. 2011b. Biological opinion on Copper Mountain North Solar Project, Boulder City, Clark County, Nevada. Memorandum to Field Manager, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada; Director, Public Works, City of Boulder City, Boulder City, Nevada; Adaptive Management Coordinator, Desert Conservation Program, Las Vegas, Nevada; Supervisory Biologist – Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. Dated December 29. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.

- U.S. Fish and Wildlife Service. 2011c. Biological opinion on the Mojave Solar, LLC's Mojave Solar Project, San Bernardino County, California (8-8-11-F-3). Letter sent to Director of Environmental Compliance, Loan Guarantee Program, Department of Energy, Washington, D.C. and Field Manager, Barstow Field Office, Bureau of Land Management, Barstow, California. Dated March 17. From Field Supervisor, Ventura Fish and Wildlife Office. Ventura, California.
- U.S. Fish and Wildlife Service. 2011d. Biological opinion on the Desert Sunlight Solar Farm Project, Riverside County, California. Memorandum to Field Manager, Palm Springs South Coast Field Office, Bureau of Land Management, Palm Springs, California. Dated July 6. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.
- U.S. Fish and Wildlife Service. 2011e. Biological opinion on the Rice Solar Energy Project, Riverside County, California. Dated July 27. Letter to John, Holt, Environmental Manager, Desert Southwest Customer Service Region, Western Area Power Administration, Phoenix, Arizona. From Jim A. Bartel, Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.
- U.S. Fish and Wildlife Service. 2012a. Biological opinion for Techren Boulder City Solar Project, Boulder City, Clark County, Nevada. Dated December 28, 2012. Memorandum to Field Manager, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada; Director, Public Works, City of Boulder City, Boulder City, Nevada; Adaptive Management Coordinator, Desert Conservation Program, Las Vegas, Nevada; Supervisory Biologist – Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.
- U.S. Fish and Wildlife Service. 2012b. Biological opinion for the K Road Moapa Solar Project, Moapa River Indian Reservation, Clark County, Nevada. Memorandum to Superintendent, Southern Paiute Agency, Bureau of Indian Affairs. St. George, Utah. Dated March 7. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.
- U.S. Fish and Wildlife Service. 2013a. Biological opinion for the Stateline Solar and Silver State Solar South Projects, San Bernardino County, California, and Clark County, Nevada. Dated September 30. Memorandum to Field Manager, Needles Field Office, Bureau of Land Management, Needles California, and Assistant Field Manager, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada. From Acting Field Supervisor, Ventura Fish and Wildlife Office. Ventura, California.
- U.S. Fish and Wildlife Service. 2013b. Biological opinion reinitiation of Copper Mountain North Solar Project, Boulder City, Clark County, Nevada. Memorandum to Field Manager, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada; Director, Public Works, City of Boulder City, Boulder City, Nevada; Adaptive Management Coordinator, Desert Conservation Program, Las Vegas, Nevada; Supervisory Biologist – Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. Dated August 8. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.
- U.S. Fish and Wildlife Service. 2013c. Biological opinion on the McCoy Solar Power Project, Riverside County, California. Dated March 6. Memorandum to Field Manager, California Desert District Office, Bureau of Land Management, Moreno Valley, California. From Field

Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.

U.S. Fish and Wildlife Service. 2013d. Biological opinion on the Desert Harvest Solar Project, Riverside County, California [CACA 044919]. Dated January 15. Memorandum to Field Manager, Palm Springs-South Coast Field Office, Bureau of Land Management, Moreno Valley, California. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.

U.S. Fish and Wildlife Service. 2014a. Biological opinion for the Townsite Solar Transmission Project. Dated July 24. Memorandum to Environmental Manager, Western Area Power Administration, U.S. Department of Energy, Phoenix, Arizona; Supervisory Biologist - Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.

U.S. Fish and Wildlife Service. 2014b. Biological opinion for Res Americas Moapa Solar Energy Center, Moapa River Indian Reservation, Clark County, Nevada. Dated January 21. Memorandum to Superintendent, Southern Paiute Agency, Bureau of Indian Affairs, St. George, Utah; Chairman, Moapa Band of Paiutes, Moapa, Nevada; Assistant Field Manager of Renewable Resources, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada; Supervisory Biologist – Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. From State Supervisor, Nevada Fish and Wildlife Office. Reno, Nevada.

U.S. Fish and Wildlife Service. 2015a. Biological opinion for the Valley Electric Association's Community Solar Project Low-Effect Habitat Conservation Plan. Dated October 8. Memorandum to Assistant Regional Director, Ecological Services, Sacramento, California; Supervisory Biologist – Habitat, Nevada Department of Wildlife, Las Vegas, Nevada. From Field Supervisor, Southern Nevada Fish and Wildlife Office. Las Vegas, Nevada.

U.S. Fish and Wildlife Service. 2015b. Biological opinion for the RE Barren Ridge 1 LLC's RE Cinco Generation Intertie Line and RE Cinco Solar Project, Kern County, California (2831-03 (P) CACA-53735 CAD000.06). Dated February 11. Memorandum to Field Manager, Ridgecrest Field Office, Bureau of Land Management, Ridgecrest, California, and Deputy Regional Director, Region 8, U.S. Fish and Wildlife Service, Sacramento, California. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.

U.S. Fish and Wildlife Service. 2015c. Biological opinion for the Soda Mountain Solar Project, San Bernardino County, California [2831-03(CP), CACA-49584, CAD000.06/CAD080]. Dated January 13. Memorandum to District Manager, California Desert District, Bureau of Land Management, Moreno Valley, California. From Field Supervisor, Carlsbad Fish and Wildlife Office. Carlsbad, California.

U.S. Fish and Wildlife Service. 2015d. Biological opinion for four solar energy projects in the Dry Lake Solar Energy Zone. Dated May 1. Memorandum to Assistant Field Manager of Natural Resources, Las Vegas Field Office, Bureau of Land Management, Las Vegas, Nevada. From Field Supervisor, Southern Nevada Fish and Wildlife Office. Las Vegas, Nevada.

U.S. Fish and Wildlife Service. 2015e. Biological opinion for the Aiya Solar Energy Project. Dated December 18. Memorandum to Deputy Regional Director – Trust Services, Western Regional Office, Bureau of Indian Affairs, Phoenix, Arizona. From Field Supervisor,

Southern Nevada Fish and Wildlife Office. Las Vegas, Nevada.

Wise, C. 2018. Electronic mail. Status of solar projects in Nevada. Dated June 28. Fish and wildlife biologist, Southern Nevada Field Office, U.S. Fish and Wildlife Service. Las Vegas, Nevada.