



fish protein to the adequate and properly balanced diet necessary for human welfare, regardless of area or financial status.

During Monday's session was the admonition, especially from the Australians, that the world still had a lot of pasture upon which to provide animal protein. This was supplemented by those who pointed out the ability of cattle to utilize feed which non-cud chewing animals would not touch and that the value of this food would be greatly increased by proper additions of fish protein.

Details of experiments with poultry feeding, corn-soybean diets with fish supplements and the use of fish meal and fish solubles in animal feeding were presented.

Those on the program of the use of fishery products in animal feeding included: A. M. Leroy, Institut National Agronomique, Paris, France; Knut Breirem, Asmund Ekern, Thor Homb, Harald Hvidsten and Ola Ulvesli, all of the Institute of Animal Nutrition, Vollebekk, Norway; R. Braude, National Institute for Research in Dairying, University of Reading, Shinfield, England; J. Baelum, National Research Institute for Animal Husbandry, Copenhagen, Denmark; Philip J. Schaible, Fur Animal Project, Department of Poultry Science, Michigan State University, East Lansing, Michigan, U.S.A.; T. D. Rummels, University of Delaware, Newark, Delaware, U.S.A.; D. G. Snyder, Bureau of Commercial Fisheries, Fishery Technology Laboratory, College Park, Maryland, U.S.A.; L. M. Potter, Department of Poultry Science, Virginia Polytechnic Institute, Blacksburg, Virginia, U.S.A.; W. J. Pudelkiewicz, Lorna Webster, L. D. Matterson, all of the University of Connecticut, Storrs, Connecticut, U.S.A.; J. R. Couch, Department of Poultry Science, Texas Agricultural and Mechanical College System, College Station, Texas, U.S.A.; A. A. Camp, Texas Agricultural Experiment Station, Gonzales, Texas, U.S.A.; C. R. Creger, Department of Biochemistry and Nutrition, Texas Agricultural and Mechanical College System, College Station, Texas, U.S.A.; H. M. Scott, W. F. Dean, A. Aguilera, R. E. Smith, all of the University of Illinois, Urbana, Illinois, U.S.A.; Shigeo Murayama, Masaaki Yanase, both of Tokai Regional Fisheries Research Laboratory, Fisheries Agency, Ministry of Agriculture and Forestry, Tokyo, Japan; K. J. Carpenter, School of Agriculture, University of Cambridge, Cambridge, England; C. H. Lea, Low Temperature Research Station, Cambridge, England; R. G. Warner, C. F. Bassett, R. P. Abernathy, all of the Department of Animal Husbandry, Cornell University and U.S. Fur Animal Experiment Station, Ithaca, New York, U.S.A.

X X X