

NWIPlus

Geospatial Database for Watershed-Level Functional Assessment

The U.S. Fish and Wildlife Service has added descriptors to the National Wetlands Inventory to strengthen the database's applications. The new tool, NWIPlus, can help wetland professionals formulate conservation strategies and assess the impact of wetland losses and gains on watershed functions.

BY RALPH TINER

Much government attention has focused on creating methods for site-specific analysis of wetland functions to evaluate the impacts of proposed development and predict the condition of wetlands through probabilistic sampling. The U.S. Fish and Wildlife Service (FWS) has been developing techniques to use its National Wetlands Inventory (NWI) data to predict wetland functions at the landscape-level, especially for watersheds. The expansion and availability of geospatial data in digital form and advances in geographic information system (GIS) technology over the last 20 years have made it possible to integrate various datasets to: (1) improve NWI mapping; (2) expand wetland classification beyond the standard NWI classification (Cowardin et al. 1979); and (3) use the expanded database to predict wetland functions. The purpose of this article is to introduce additional descriptors for wetland mapping and an inventory-based method for landscape-level functional assessment and also highlight current applications of this method by the NWI and others.

Geospatial Technology Improves the NWI

When the FWS initiated the NWI in the mid-1970s, GIS technology was in its infancy. The focus of the NWI was on producing hard-copy wetland maps. In preparing these maps, the NWI utilized 1:24,000 U.S. Geological Survey topographic maps as the base to display wetlands and deepwater habitats identified through conventional photo-interpretation techniques. When viewing the maps, people could see the size, shape, and type of wetlands situated in a particular location on the landscape and their connection to other wetlands and waters.

Ralph Tiner is the Regional Wetland Coordinator for the U.S. Fish and Wildlife Service's Northeast Region in Hadley, Massachusetts.

For example, a person could see if the wetland was contiguous with a lake, river, stream, or pond, or whether it was "geographically isolated" (i.e., completely surrounded by upland) and could use this information to gain insight into likely functions. While this could easily be done for a particular wetland or small group of wetlands, it was virtually impossible to visualize the landscape context of all wetlands within a watershed or larger geographic area, without pasting together a large number of maps in some way.

With the growth of GIS technology, the NWI began to digitize its data, creating a geospatial database containing information on the size, shape, classification, and location of mapped wetlands and deepwater habitats. This database made it possible to generate statistics on the extent of wetlands and deepwater habitats for geographic areas of varying dimensions, including entire states. The first state summary of NWI data was produced in 1985 for the state of New Jersey (Tiner 1985). Today, the NWI digital data are available for approximately 70 percent of the conterminous United States and 30 percent of Alaska, and users can produce online custom maps for areas of interest (Tiner 2009).

The capabilities of geospatial technology also made it possible to incorporate data created by other agencies and organizations into the wetland interpretation and mapping process. Now, the results of soil surveys, state wetland inventories, aquatic bed surveys, stream mapping, and even on-the-ground delineations in digital geospatial formats can be readily accessed and added to the NWI database. As digital geospatial data became more available and GIS technology advanced to become a desktop tool for scientists and planners, it became possible to use NWI data for watershed and regional analyses. It soon became evident that additional information needed to be added to the classification to predict a wide range of wetland functions for such areas.




Figure 2: Portion of wetland function map for a watershed in Saratoga County, New York.

increase the functionality of wetland inventory databases (FGDC Wetlands Subcommittee 2009).

NWIPlus for Functional Assessment

The impetus for the LLWW descriptors was to provide a convenient and consistent means of using NWI data to predict wetland functions for watersheds. Correlations between the parameters in the NWIPlus database and a number of wetland functions have been developed in working with scientists from various agencies in the Northeast (Tiner 2003b). To date, 11 functions can be predicted by this expert-designed assessment method: (1) surface water detention; (2) streamflow maintenance; (3) nutrient transformation; (4) sediment and particulate retention; (5) carbon sequestration; (6) shoreline stabilization; (7) coastal storm surge detention; (8) provision of fish and shellfish habitat; (9) provision of waterfowl and waterbird habitat; (10) provision of habi-

tat for other wildlife; and (11) conservation of biodiversity. The latter function emphasizes regionally significant wetland types as well as locally uncommon types based on NWI results. At this time, it does not incorporate data from other sources such as state natural heritage programs that could be added by others. The emphasis is on using NWIPlus to generate a preliminary watershed-based assessment of wetland functions. It is a starting point for wetland evaluation, not the end point. The correlations will be updated as needed and have, in one known case, been modified by a state (i.e., Montana) to incorporate local knowledge of wetlands into the functional relationships and to emphasize functions important to the state.

The watershed assessment approach using NWIPlus has been called Watershed-based Preliminary Assessment of Wetland Functions (W-PAWF), since it produces a first-cut evaluation based on map information. It is an inventory-based assessment method that evaluates every mapped wetland based on properties contained in the NWIPlus database. It applies general knowledge about wetlands and their functions to produce a watershed overview highlighting wetlands predicted to perform certain functions at high or moderate levels. It does not account for the opportunity that a wetland has to provide a higher level of function resulting from a certain land-use practice upstream or the presence of certain structures or land uses downstream. For example, two wetlands of equal size and like vegetation may be in the right landscape position to retain sediments. One may be downstream of a landclearing operation that has generated considerable suspended sediments in the water column, while the other is downstream from an undisturbed forest. The first wetland is trapping more sediment due to increased suspended sediment, while the second wetland is not doing so at the same rate. W-PAWF is designed to reflect the potential of a wetland to provide a function. It also does not consider the condition of the adjacent upland (e.g., level of outside disturbance) or the actual water quality of the associated water body, which may be regarded as important metrics for assessing the “health” or condition of individual wetlands.

The final product of this inventory-based assessment is a report containing a narrative description of the study area, methods, wetland types and functions plus a series of thematic maps showing wetlands by type (NWI, landscape position, landform, and water flow path), and wetlands of significance for each of 11 functions (Figure 2). Accompanying digital geospatial data are also available for other analyses. For examples of reports, go to the FWS’ Conservation library, at <http://library.fws.gov/WetlandPublications.html> or to the NWI website (<http://www.fws.gov/wetlands/>; click on the “documents search engine” icon and then type in “wetland characterization” or “functional assessment”). Documents containing the maps are quite large and may take a few moments to download.

Applications of NWIPlus Data

The first applications of NWIPlus data were done in Massachusetts to produce watershed analyses of wetland functions with special attention placed on predicting likely functions of potential wetland restoration sites (Tiner 1997). These data were the foundation for the state’s watershed-based restoration planning to help target restoration in areas providing significant benefits for flood protection, water quality improvement, or habitat.

To date, NWIPlus data have been used to: (1) better characterize wetlands and relate wetlands to water bodies and various functions (e.g., Tiner et al. 2007, Newlon and Burns 2009); (2) increase public awareness of the functions of wetlands and understanding that all wetlands are not alike in either form or function (e.g., New York City Department of Environmental Protection 2009, Homsey 2009); (3) assist agencies in developing wetland conservation strategies (e.g., Berner 2009, Martin 2008, Maryland Department of Environment 2006); (4) help agencies evaluate the effect of recent court decisions on wetlands (e.g., isolated wetlands, Vance 2009); (5) stratify wetlands for research purposes (e.g., Jacobs et al. 2009); (6) estimate the effect of wetland trends on the capacity of watersheds to provide wetland services (e.g., Kudray and Schemm 2008); and (7) assess the cumulative effect of historic wetland loss on functions (e.g., Tiner 2005, Fizzell 2007). Referring to NWIPlus maps of Maine's Casco Bay watershed, which show wetlands of significance by function, William Honachefsky, in his 1999 book, *Ecologically Based Municipal Land Use Planning*, stated that "these newest maps allow decisionmakers and land planners to better assess the true worth of individual wetlands" and that such information should be incorporated in local municipal master plans. The Center for Watershed Protection's report, "Using Local Watershed Plans to Protect Wetlands," mentioned the value of LLW/W descriptors in estimating wetland functions for managing wetlands at the watershed level, and suggested that these types of assessments should be part of any wetland inventory (Capiella et al. 2006).

NWIPlus data have been generated for numerous areas in the country. It has become a standard practice for updating NWI data in the Northeast and will be considered for updates elsewhere depending on regional priorities and budgets. Using these data, W-PAWF has been applied or will be applied to numerous areas in the country by the FWS. Various states have begun using these LLW/W descriptors in their inventories to create an NWIPlus database. For example, the state of Montana is applying these attributes to aid in assessing wetland functions for its watersheds (e.g., Kudray and Schemm 2008, Newlon and Burns 2009). The states of Michigan and Delaware have applied the descriptors to wetland inventory projects. Minnesota has included these descriptors in their requirements for updating NWI data (Minnesota Department of Natural Resources 2009). Other states where wetland inventories are underway and that are considering the application of LLW/W descriptors include Georgia, Kansas, New Mexico, and Oklahoma.

Conclusion

Adding LLW/W descriptors to the NWI database creates a more functional and powerful database—NWIPlus. It can be used in formulating wetland conservation strategies

to help prioritize wetlands for acquisition, restoration, or strengthened protection, as an educational tool to improve the public's understanding of wetland functions, and as a cumulative impact assessment tool to estimate the impact of wetland losses and gains on watershed functions. By creating NWIPlus data, the results of wetland inventories can better describe the variability between and among wetlands and include watershed-based or landscape-level wetland functional assessments. For more than three decades, NWI maps have been used by various levels of government in compiling natural resource inventories, watershed planning, and improving wetland protection. Now, by enhancing NWI data and using it for wetland functional assessment, they have a more valuable tool at their disposal for resource conservation and management. ■

REFERENCES

- Berner, Jason T., *Alternative Futures for Headwater Stream and Wetland Landscapes in the Upper Delaware Basin, New York, USA* (2009) (unpublished masters thesis, University of Illinois at Urbana-Champaign), available at http://www.catskillstreams.org/pdfs/del_wetland_thesis.pdf.
- BRINSON, MARK M., U.S. ARMY ENGINEER WATERWAYS EXPERIMENT STATION, *WETLAND RESEARCH PROGRAM TECHNICAL REPORT WRP-DE-4, A HYDROGEOMORPHIC CLASSIFICATION FOR WETLANDS* (1993), available at <http://el.erdc.usace.army.mil/wetlands/pdfs/wrpde4.pdf>.
- CAPIELLA, KAREN ET AL., CENTER FOR WATERSHED PROTECTION, *ARTICLE TWO OF THE WETLANDS AND WATERSHEDS ARTICLE SERIES, USING LOCAL WATERSHED PLANS TO PROTECT WETLANDS* (2006), available at http://www.cwp.org/Resource_Library/Special_Resource_Management/wetlands.htm.
- COWARDIN, LEWIS M. ET AL., U.S. FISH AND WILDLIFE SERVICE, *CLASSIFICATION OF WETLANDS AND DEEPWATER HABITATS OF THE UNITED STATES* (1979), available at http://library.fws.gov/FWS-OBS/79_31.pdf.
- DIVISION OF ECOLOGICAL RESOURCES, MINNESOTA DEPARTMENT OF NATURAL RESOURCES, *REQUIREMENTS FOR THE NATIONAL WETLANDS INVENTORY UPDATE OF MINNESOTA* (2009), available at http://files.dnr.state.mn.us/eco/wetlands/nwi_requirements_v5.pdf.
- DIVISION OF HABITAT AND RESOURCE CONSERVATION, U.S. FISH AND WILDLIFE SERVICE, *STATUS REPORT ON THE NATIONAL WETLANDS INVENTORY: 2009* (Ralph W. Tiner ed.) (2009), available at http://www.fws.gov/wetlands/_documents/gOther/StatusReportNWIProgram2009.pdf.
- FEDERAL GEOGRAPHIC DATA COMMITTEE WETLANDS SUBCOMMITTEE, *WETLANDS MAPPING STANDARD, FGDC DOCUMENT NO. FGDC-STD-015-2009* (2009), available at http://www.fgdc.gov/standards/projects/FGDC-standards-projects/wetlands-mapping/2009-08%20FGDC%20Wetlands%20Mapping%20Standard_final.pdf.
- FIZZELL, CHAD A., SOUTHWEST MICHIGAN PLANNING COMMISSION, *ASSESSING CUMULATIVE LOSS OF WETLAND FUNCTIONS IN THE PAW PAW WATERSHED USING ENHANCED NATIONAL WETLANDS INVENTORY DATA* (2007), available at http://www.swmpc.org/downloads/pprw_WetlandFunctionAssmnt.pdf.
- Homsey, Andrew, *Development of Hydro-Geomorphologic Modifiers for Delaware's Wetlands in the Delaware Estuary* (2009) (poster presentation at 2009 Delaware Estuary Science and Environmental Summit, Partnership for the Delaware Estuary Inc.), available at http://www.delawareestuary.org/scienceandresearch/science_conf/SC09/2009%20PDF%20Presentations/Session%206/ARHomsey_Poster.pdf.

Resources

The NWI can be viewed online at <http://www.fws.gov/wetlands/>.

Continued on page 23

This May, in the spirit of American Wetlands Month, let us celebrate our coastal wetlands . . . connecting us all with their beauty, their ability to protect our communities from some of nature's most destructive forces, and their vital support of water quality, fish, and wildlife that enhance our lives, no matter how far inland we live or travel. We have a shared responsibility to conserve coastal wetlands so that future generations can also enjoy all the benefits these vibrant and important places have to offer. To learn more about American Wetlands Month and activities in your area, or for additional resources to explore, protect, and restore wetlands, visit www.epa.gov/owow/wetlands/awm. ■

REFERENCES

- Nathan L. Bindoff et al., 2007: *Observations: Oceanic Climate Change and Sea Level*, in *CLIMATE CHANGE 2007: THE PHYSICAL SCIENCE BASIS* (S. Solomon et al., eds., Cambridge Univ. Press 2008).
- KRISTEN M. CROSSETT ET AL., NATIONAL OCEAN SERVICE, NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, COASTAL TRENDS REPORT SERIES, POPULATIONS TRENDS ALONG THE COASTAL UNITED STATES: 1980-2008 (2004), available at http://oceanservice.noaa.gov/programs/mb/pdfs/coastal_pop_trends_complete.pdf.
- DONALD W. FIELD ET AL., NATIONAL OCEAN SERVICE, NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, COASTAL WETLANDS OF THE UNITED STATES: AN ACCOUNTING OF A VALUABLE RESOURCE (1991), available at <http://dnrweb.dnr.state.md.us:8080/FullDisp?itemid=00012267>.
- Factoids About Louisiana's Coastal Wetlands*, LaCoast website, at <http://www.lacoast.gov/education/factoid%20old.htm> (last visited Mar. 22, 2010).
- K.A. LELLIS-DIBBLE ET AL., NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, NOAA TECHNICAL MEMORANDUM NMFS-F/SOP-90, ESTUARINE FISH AND SHELLFISH SPECIES IN U.S. COMMERCIAL AND RECREATIONAL FISHERIES: ECONOMIC VALUE AS AN INCENTIVE TO PROTECT AND RESTORE ESTUARINE HABITAT (2008), available at <http://spo.nwr.noaa.gov/tm/TM90.pdf>.
- LINWOOD PENDLETON, RESTORE AMERICA'S ESTUARIES, THE ECONOMIC AND MARKET VALUE OF COASTS AND ESTUARIES: WHAT'S AT STAKE? (2008), available at <http://www.nmfs.noaa.gov/habitat/habitatconservation/publications/Economic%20and%20Market%20Value%20of%20Coasts%20and%20Estuaries.pdf>.
- SUSAN-MARIE STEDMAN & TOM E. DAHL, NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, NATIONAL MARINE FISHERIES SERVICE AND DEPARTMENT OF THE INTERIOR, FISH AND WILDLIFE SERVICE, COASTAL WATERSHEDS OF THE EASTERN UNITED STATES: STATUS AND TRENDS OF WETLANDS 1998 TO 2004 (2008), available at http://www.fws.gov/Wetlands/_documents/gSandT/NationalReports/StatusTrendsWetlandsCoastalWatershedsEasternUS1998to2004.pdf.
- JAMES TITUS, CLIMATE CHANGE SCIENCE PROGRAM SYNTHESIS AND ASSESSMENT PRODUCT 4.1, U.S. ENVIRONMENTAL PROTECTION AGENCY, COASTAL SENSITIVITY TO SEA-LEVEL RISE: A FOCUS ON THE MID-ATLANTIC REGION (2009), available at <http://www.epa.gov/climatechange/effects/coastal/sap4-1.html>.
- U.S. FISH & WILDLIFE SERVICE, COASTAL ECOSYSTEMS PROGRAM, BRANCH OF COASTAL AND WETLAND RESOURCES, DIVISION OF HABITAT CONSERVATION (1995), at 48.
- STATUS REPORT ON THE NATIONAL WETLANDS INVENTORY PROGRAM: 2009, U.S. FISH AND WILDLIFE SERVICE, DIVISION OF HABITAT AND RESOURCE CONSERVATION (Ralph Tiner, ed., 2009), available at http://library.fws.gov/wetlands/nwi_status_2009.pdf.
- TINER, RALPH W., *Assessing Cumulative Loss of Wetland Functions in the Nanticoke River Watershed Using Enhanced National Wetlands Inventory Data*, 25 WETLANDS (2) 405-19 (2005), available at http://library.fws.gov/Wetlands/TINER_WETLANDS25.pdf.
- TINER, RALPH W., NATIONAL WETLANDS INVENTORY PROGRAM, U.S. FISH AND WILDLIFE SERVICE, CORRELATING ENHANCED NATIONAL WETLANDS INVENTORY DATA WITH WETLAND FUNCTIONS FOR WATERSHED ASSESSMENTS: A RATIONALE FOR NORTHEASTERN U.S. WETLANDS (2003b), available at http://library.fws.gov/Wetlands/corelate_wetlandsNE.pdf.
- TINER, RALPH W., NATIONAL WETLANDS INVENTORY PROGRAM, U.S. FISH AND WILDLIFE SERVICE, DICHOTOMOUS KEYS AND MAPPING CODES FOR WETLAND LANDSCAPE POSITION, LANDFORM, WATER FLOW PATH, AND WATERBODY TYPE DESCRIPTORS (2003a), available at <http://library.fws.gov/Wetlands/dichotomouskeys0903.pdf>.
- TINER, RALPH W., NATIONAL WETLANDS INVENTORY PROGRAM, U.S. FISH AND WILDLIFE SERVICE, WETLANDS OF NEW JERSEY (1985), available at http://library.fws.gov/Wetlands/NJ_wetlands85.pdf.
- TINER, RALPH W., *Piloting a More Descriptive NWI*, 19 NAT'L WETLANDS NEWSL. (Sept.-Oct.) 14-16 (1997).
- TINER, RALPH W. & HERBERT C. BERGQUIST, NATIONAL WETLANDS INVENTORY PROGRAM, U.S. FISH AND WILDLIFE SERVICE, THE HACKENSACK RIVER WATERSHED, NEW JERSEY/NEW YORK: WETLAND CHARACTERIZATION, PRELIMINARY ASSESSMENT OF WETLAND FUNCTIONS, AND REMOTELY-SENSED ASSESSMENT OF NATURAL HABITAT INTEGRITY (2007), available at <http://library.fws.gov/Wetlands/HackensackRiverWatershed07.pdf>.
- VANCE, LINDA K., MONTANA NATURAL HERITAGE PROGRAM, GEOGRAPHICALLY ISOLATED WETLANDS AND INTERMITTENT/EPHEMERAL STREAMS IN MONTANA: EXTENT, DISTRIBUTION, AND FUNCTIONS (2009), available at http://mtnhp.org/Reports/Isolated_Wetlands.pdf.

Resources

Learn more about how coastal communities can protect their environmental resources, at www.coastalsmartgrowth.noaa.gov and www.epa.gov/smart-growth/sg-coastal.html.

Information about the Workgroup's regional reviews can be read at www.epa.gov/owow/wetlands and www.nmfs.noaa.gov/habitat/habitatprotection/wetlands/index.htm.

For information about American Wetlands Month and events in your area, visit www.epa.gov/owow/wetlands/awm.

NWIPlus, from page 7

- HONACHEFSKY, W.B., ECOLOGICALLY BASED MUNICIPAL LAND USE PLANNING (Lewis Publishers 1999), at 256.
- JACOBS, AMY ET AL., WATERSHED ASSESSMENT SECTION, DELAWARE DEPARTMENT OF NATURAL RESOURCES AND ENVIRONMENTAL CONTROL, VOLUME 1: NONTIDAL WETLANDS. WETLAND CONDITION OF THE INLAND BAYS WATERSHED (2009), available at <http://www.dnrec.delaware.gov/Admin/DelawareWetlands/Documents/IB%20Wetland%20Report%20Vol%201%20.pdf>.
- KUDRAY, GREGORY M. & TOM SCHEMM, MONTANA NATURAL HERITAGE PROGRAM, WETLANDS OF THE BITTERROOT VALLEY: CHANGE AND ECOLOGICAL FUNCTIONS (2008), available at http://mtnhp.org/reports/Bitterroot_Change.pdf.
- Martin, Jill W. A Community's Watershed Moment, Escanaba, Michigan (presentation at Association of State Wetlands Managers' Wetlands and Global Climate Change Conference) (Sept. 16-18, 2008), available at http://www.aswm.org/calendar/wetlands2008/pp/wiese_martin.pdf.
- MARYLAND DEPARTMENT OF THE ENVIRONMENT, PRIORITIZING SITES FOR WETLAND RESTORATION, MITIGATION, AND PRESERVATION IN MARYLAND: DORCHESTER COUNTY (2006), available at <http://www.mde.state.md.us/assets/document/wetlandswaterways/DO.pdf>.
- MARYLAND DEPARTMENT OF THE ENVIRONMENT, PRIORITY AREAS FOR WETLAND RESTORATION, PRESERVATION, AND MITIGATION IN MARYLAND'S COASTAL BAYS (2004), available at http://www.mde.state.md.us/assets/document/wetlandswaterways/CB_all.pdf.
- NATURAL RESOURCES MANAGEMENT SECTION, NEW YORK CITY DEPARTMENT OF ENVIRONMENTAL PROTECTION, WETLANDS IN THE WATERSHEDS OF THE NEW YORK CITY WATER SUPPLY SYSTEM (2009), at 26.
- NEWLON, KAREN R. & MEGHAN D. BURNS, MONTANA NATURAL HERITAGE PROGRAM, WETLANDS OF THE FLATHEAD VALLEY: CHANGE AND ECOLOGICAL FUNCTIONS (2009), available at http://mtnhp.org/Reports/Flathead_Change.pdf.