

# **2010 Bull Trout Redd Monitoring in the Wallowa Mountains**



**Prepared by: Gretchen Sausen  
U.S. Fish and Wildlife Service  
La Grande Field Office  
March 2011**

# **2010 Bull Trout Redd Monitoring in the Wallowa Mountains**

**Prepared by: Gretchen Sausen  
U.S. Fish and Wildlife Service  
La Grande Field Office  
March 2011**

## **ABSTRACT**

Bull trout were listed as threatened under the Endangered Species Act in 1998 due to declining populations. The U. S. Fish and Wildlife Service (Service) recommends monitoring populations in subbasins where little is known, including the Grande Ronde and Imnaha subbasins. Spawning survey data is important for determining relative abundance and distribution trends in bull trout populations. This report summarizes the 2010 bull trout spawning data collected in the Wallowa Mountains of NE Oregon and compares this with past years' data. Bull trout spawning surveys have been conducted on similar index areas for selected Grande Ronde and Imnaha River streams from 1999 to 2010. Surveys were conducted by the Oregon Department of Fish and Wildlife (ODFW), U.S. Forest Service (USFS), Service, Nez Perce Tribe (NPT), Idaho Power, fisheries consultants, and volunteers. Objectives of the survey included; locate bull trout spawning areas, determine redd characteristics, determine bull trout timing of spawning, collect spawning density data, determine and compare the spatial distribution of redds along the Lostine River in 2005 through 2010, and over time use all of the data to assess local bull trout population trends and the long-term recovery of bull trout. Timing of spawning, total redds, redd sizes, and redd locations are documented in the report. The local bull trout populations were relatively stable for the survey period (1999-2010) but there was a downward trend in redd counts in 2009 and 2010. The Imnaha population is one of the strongholds within the Imnaha Subbasin. The Lostine River and Bear Creek contain brook trout and hybridization is likely occurring (the Lostine has had documented bull trout and brook trout pairing up for spawning).

## **ACKNOWLEDGMENTS**

The Service has, for the past seven years, provided staff time necessary for the coordination, implementation, and analysis and report summarization of this project. Prior to Service support of this project, the Wallowa-Whitman National Forest, Wallowa Mountains Office (WMO) had secured previous years' funding and support, and was responsible for the coordination, implementation, and analysis and report summarization.

This project would not have been possible without the dedication, hard work, funding, and assistance provided by all the partners. Due to Oregon Watershed Enhancement Board (OWEB) funding in 2007-2009 and recently in 2010-2011; Del Sol Wilderness Adventures provided an invaluable service in 2008 through 2010 by providing a horse/mule packer to pack our gear in and out of the Upper Imnaha to conduct our annual spawning survey in that drainage. The WMO in 2005 through 2007 and other

commercial packers in past years provided this service (2007 funding was provided by OWEB). I would like to thank the partners in 2007 - 2010 which included; the WMO, ODFW, OWEB, Grande Ronde Model Watershed, NPT, the Service, NMFS, Anderson Perry and Associates (Anderson Perry), Idaho Power, consultants, and volunteers. In 2010, Sherry Ann Larson volunteered for two days (1 day) on Bear Creek and one day on Deer Creek. Special thanks to the people who walked the streams (including surveys conducted in 2007-2010), helped with scheduling surveys and surveyors, provided access to private property, packed us into remote areas to survey, assisted with the OWEB grants, assisted in mapping, or summarized the data. These included: Gary Miller (Service), Brad Smith (formerly ODFW), Alan Miller (WMO), John Hollenbeak (WMO – horse/mule packer), Barry Cox, Shay Mann, Mary Hawkins, Paul Arentsen, Morgan Jenkins (Del Sol Wilderness Adventures Horse/Mule Packers and Winding Waters River Expeditions), Jeff Oveson (Grande Ronde Model Watershed), Coby Menton (Grande Ronde Model Watershed), Mary Estes (Grande Ronde Model Watershed), Ken Bronec (consultant), Peter Cleary (NPT), Pat Leach (private access to Lostine River Ranch), Nancy Clarke (private access to Lostine River upstream of Six Mile Bridge), Stewart and Susan Coleman (private access to Lostine river near acclimation site), (Jeff Nehls (NPT), Marika Dobos (ODFW), Eric Shoudel (ODFW), Sherry Ann Larson (volunteer), Ashley Davidson (ODFW), Jeff Yanke (ODFW), Ian Wilson (NPT), Nick Albrecht (ODFW), Kyle Bratcher (ODFW), Jim Trainer (Idaho Power), Rick Wilkison (Idaho Power), Steve Brink (Idaho Power), Dave Bright (NPT), Suzanne Anderson (Service), Lynne Price (consultant), Mac Huff (consultant), and Jim Harbeck (NPT).

## INTRODUCTION

Bull trout were listed as threatened under the Endangered Species Act in 1998 due to declining populations. The Service recommends monitoring populations in subbasins where little is known including the Grande Ronde and Imnaha subbasins (USFWS 2002). Spawning survey data is important for determining relative abundance and distribution trends in bull trout populations. A minimum of 15 years is needed for determining bull trout population trends (Maxwell 1999). Without adequate funding, it had been difficult to find sufficient numbers of experienced bull trout surveyors and packers for surveys in the back-country, and to obtain adequate supplies to get the work accomplished. OWEB funding for the project supported the continued survey of bull trout spawning areas in years 2007 through 2010 in the Wallowa Mountains of northeast Oregon. Bull Trout redd counts (spawning surveys) have been conducted annually on the Wallowa Valley, Hells Canyon National Recreation Areas (HCNRA), and Eagle Cap districts of the USFS and along some sections of private property of the Lostine River by the Service, ODFW, USFS, NPT, contractors, and volunteers for the past nine to ten years.

Objectives of the bull trout spawning surveys included:

- Locate bull trout spawning areas.
- Determine redd characteristics.
- Determine bull trout timing of spawning.
- Collect spawning density data.
- Map the location of the bull trout spawning reaches.
- Determine and compare the spatial distribution of redds along the Lostine River in 2005 through 2010. Collect UTM spatial redd data on Big Sheep, Lick Creek, and Middle Imnaha to compare after a few years of data collection.

- Assess population trends for local bull trout populations.
- Use this information for helping assess the long-term recovery of bull trout.

## LOCATION

The Service and several partners conducted bull trout spawning surveys in 2010 on selected streams in the Grande Ronde and Imnaha Sub-Basins. Stream systems surveyed in 2010 for bull trout redds included; the Lostine River, Bear and Goat Creeks, the Imnaha River, Big Sheep Creek and Lick Creek (Figure 1). An exploratory survey was conducted on Deer Creek in 2010 (commencing at the newly installed culvert at Rd.8270 to 0.8 miles upstream).



Fluvial Bull Trout, Photo by Joel Satore, National Geographic Stock Photos with Wade Fredenberg




Figure 1. Wallowa Mountain Bull Trout Redd Survey Areas.

## METHODS

Spawning surveys for bull trout require as many as ten to twelve people in one day (for complete surveys on large rivers) to complete the surveys during the spawning time. Surveyors walk the rivers through the selected “index areas” to locate the bull trout redds.

This project is part of a larger effort in NE Oregon and SE Washington that occurs during spawning (September – October). Due to the lack of available experienced surveyors to conduct these surveys, we have had to increase our survey days on the accessible sections of the Lostine and Imnaha Rivers to two days (conducting half of the survey length one day and the other half the following day). Surveys were conducted twice (mid and late bull trout spawning season) on the Lostine River, Big Sheep Creek, Lick Creek, Middle Imnaha (Blue Hole to Indian Crossing) and Bear and Goat Creeks. One-time surveys were conducted late in the spawning season in 2010, on the Upper Imnaha River and tributaries, due to access and funding limitations. An exploratory survey was conducted on Deer Creek in 2010 for 0.8 miles of survey. Appendix B, Table 1 compares survey data and survey frequency for 1999-2010 bull trout spawning surveys on selected Grande Ronde and Imnaha River streams. Stream miles surveyed (not including repeat surveys) for the above streams totaled 41.3 in 2007, 46.3 in 2008, 41.8 in 2009, and 41.1 in 2010.

Total redd numbers are all redds documented, and not necessarily comparable river miles (refer to Appendix B, Table 2a-2d for comparable reaches and redd counts for those sections).

The survey protocol (in addition to repeat surveys, or one-time late surveys where feasible) included; 1) visits to known bull trout redds and review of survey form prior to redd count survey, 2) experienced bull trout redd count surveyor(s) paired with inexperienced surveyor (on the job training), 3) bull trout redds measured, data recorded, and redds flagged during survey, and 4) all stream flagging removed post surveys.



**Bull Trout Spawning Trio on Redd, Lostine River (Photo courtesy of USFS).**

Data recorded during the bull trout spawning surveys included; 1) date of survey, 2) stream location, 3) size of redds, 4) visibility of redds, 5) number of redds, and 6) approximate number and sizes of bull trout observed during surveys. In past years, reach locations (upstream and downstream boundary UTM coordinates) were documented. In 2009 and 2010, in addition to the above, bull trout redd UTM locations on the Lostine River, Big Sheep Creek, Lick Creek, and Middle Imnaha within the “index areas” also was documented. Information collected during the bull trout spawning surveys is compiled by the Service and made available to other agencies (i.e., this report).



**Shelly Schmidt and Sue Brady of Anderson Perry, Inc surveying a bull trout redd on the Lostine River, 2008.**

## **RESULTS**

### **Location of Bull Trout Spawning Habitat Areas Surveyed**

Bull trout spawning surveys have been conducted on similar index areas for selected Grande Ronde and Imnaha River streams from 1999 to 2010. During these years, bull trout spawning areas have been established (in particular, the Lostine and Imnaha Rivers) for these streams. Redd characteristics also have been measured on these streams. The Middle Imnaha (Imnaha River from the fish weir below Gumboot confluence to Indian Crossing) was not surveyed in 2005 through 2010, (this area was surveyed in 1999 to 2004 and is considered bull trout spawning habitat). This portion of known bull trout spawning habitat on the Imnaha was not surveyed for the past six years because of limited funding, a lack of experienced surveyors, and a minimal number of redds documented in this area in past years. Bear Creek and Goat Creek were surveyed as in past years but the survey area of Bear Creek and Goat Creek increased (1.9-3.8 total miles between 1999-2006 versus 7.2 in 2007) and the survey frequency increased (generally once from 1999-2006 and twice in 2007). Although the survey area increased in 2007, the redd numbers did not increase substantially, especially in the lowermost survey reaches. In 2008, 2009, and 2010 the surveys were conducted twice in the spawning season and an additional 1.4 miles of Bear Creek was surveyed upstream of the comparable reach for a total of 3.2 miles. In 2010, an exploratory survey for 0.8 miles of stream was conducted on Deer Creek.



**Jim Trainer (Idaho Power) surveying from log on the Blue Hole to Indian Crossing, Imnaha River Survey, 2010**

### **Timing of Bull Trout Spawning**


Timing of bull trout spawning for our surveyed streams, in general, is approximately September 1 through October 15, and as early as August 15 in the Imnaha River system. The Lostine River has been very consistent or predictable (with commencement of spawning documented in 2006 as early as the first week in September), with the Imnaha (being a much larger system) being less predictable. Mary Edwards, fisheries biologist with the Nez Perce Tribe, observed bull trout and brook trout spawning near the acclimation facility as early as late July 2010. Mary was underwater snorkeling at the time of her observation. This early bull spawning has not been observed in this area prior to this observation. Bull trout surveys conducted in September/October 2010 did not confirm any bull trout redds in this location; but that could have been due to the high density of Chinook spawning that occurred in the area. The above information is based on documentation during bull trout spawning surveys and chinook surveys where bull trout were spawning.

We are not exactly certain of when spawning commences and ends within the Upper Imnaha (within the mainstem, North Fork, South Fork, and Cliff Creek [a resident tributary]). There are questions as to what time of the year, dependent on annual flows, bull trout pass over the falls. Some years we have seen fluvial size bull trout spawning in the South Fork Imnaha in mid-late September/early October and in recent years we have not. ODFW observed large fluvial bull trout spawning in South Fork Imnaha in mid-August 2005 (B. Knox, ODFW, pers. comm., 2005). ODFW has observed fluvial bull trout spawning as early as mid-August (during chinook surveys) below the Imnaha falls and as late as early October (during our bull trout surveys) in several years (B. Smith, ODFW, pers. comm., 2005). More years of observation and survey data are needed to understand bull trout spawning and adult movement in the Imnaha Sub-Basin, and Big Sheep and Bear Creek Watersheds.

## Total Number of Bull Trout Redds

### *Lostine River*

Refer to Appendix B, Table 3a and 3b for bull trout redd count summary data for 2010. Thirty six total bull trout redds for 10.1 miles of survey (including Pole Bridge to Six Mile Bridge) were documented in 2010 on the Lostine River. The Pole Bridge to Six Mile Bridge section has not been surveyed every year. The following data for the Lostine River compares consistently surveyed index areas on the Lostine River (8.5 miles) from 1999 to 2010 excluding the Pole Bridge to Six Mile section (Figure 2). The Lostine River had a low of 22 redds in 2002 to a high of 70 redds in 2003. Redd numbers decreased again in 2004 but had been increasing through 2008. Redd numbers decreased in 2010. The twelve-year average from 1999 to 2010 for the Lostine River is 38 redds, approximately 54 percent of the 2003 total. The highest bull trout redd numbers (“the bread and butter”) within the Lostine River has consistently been observed and recorded in the headwaters, from Shady Campground to Bowman [approximately River Mile (RM) 24.5 to RM 22]. In 2010, as in most years, the densities were greatest in the uppermost reach (Shady to French Camp) which is upstream of the chinook spawning index areas.



**Figure 2. Comparison of bull trout surveyed redds and index redds (comparable miles) documented from 1999 to 2010 on the Lostine River.**

### *Bear Creek*

Seven total bull trout redds for 3.2 miles of survey were documented in 2010 on Bear Creek (including Goat Creek). The following data for Bear Creek compares consistently surveyed index areas on Bear Creek and Goat Creek (1.9 miles) from 1999 to 2010 (Figure 5). Redd counts on Bear Creek and Goat Creek had a low of 5 redds in 2000 to a high of 12 total redds in 2001, and decreasing to 7 redds in 2010. The twelve-year average from 1999 to 2010 is 9 redds for Bear and Goat Creeks. Bear Creek/Goat Creek spawning data collected from 1999 to 2006 is restricted in scope due to access and funding limitations, and 2007 was expanded to help identify total spawning area for bull trout in Bear and Goat Creeks. Although the survey area increased in 2007, the redd

numbers did not increase substantially, especially in the lowermost survey reaches. In 2008 and 2010, the surveys were conducted twice in the spawning season and an additional 1.4 miles of Bear Creek was surveyed upstream of the comparable reach for a total of 3.2 miles. The highest bull trout redd counts for the survey sections on Bear/Goat Creeks have been recorded in Goat Creek (mouth to waterfall, RM 0 to RM 0.9), except in 2008 where more redds were documented in Bear Creek than in Goat Creek.




Figure 5. Comparison of bull trout surveyed redds and index redds (comparable miles) documented from 1999 to 2010 on Bear Creek (including Goat Creek).

The local bull trout population in the Lostine and Bear Creek surveys appears to be relatively stable for the survey period (2000-2010), with a downward trend in the Lostine River portion of the population in 2010.

### Deer Creek

In 2010, an exploratory survey was conducted on Deer Creek for approximately 0.8 miles of stream. The survey was located at the Deer Creek culvert at Rd.8270 and upstream approximately 0.8 miles. The survey was conducted twice in 2010 and a total of 12 resident bull trout redds were documented.

### Imnaha River

One hundred and thirty three total bull trout redds for 19.4 miles of survey were documented in 2010 on the Imnaha River (from Indian Crossing to Blue Hole and upstream). Indian Crossing to Blue Hole was surveyed twice (mid to late spawning season) and upstream areas were surveyed once (mid spawning season) in 2010. The following data for the Imnaha River compares consistently surveyed index areas on the Imnaha River (17.5 miles) from 2001 to 2010 (Figure 3). The ten-year average from 2001 to 2010 was 193 redds for the Imnaha River system. Total redd numbers on the Imnaha ranged from 101-262 within that period. The highest bull trout redd counts for the Imnaha River from 2001 to 2010 was recorded in the Upper Imnaha from Blue Hole to Cliff Creek, including Upper Imnaha tributaries. In 2006 and 2007, there was a

significant shift in documented spawning distribution from past years. In 2006 through 2008, the majority of the spawning bull trout were located from the Imnaha falls to Indian Crossing, whereas, in past years the distribution had higher numbers above the Blue Hole (two miles upstream of Indian Crossing), as well as in the upper tributary streams (S.F. and N.F. Imnaha).

In 2010, the Middle Fork Imnaha River, above the falls to forks, for approximately 0.6 miles had an exploratory survey conducted to determine bull trout and redds presence or absence. The habitat was excellent but no bull trout or redds were documented.

In 2009, Cliff Creek a resident bull trout tributary to S.F. Imnaha had the greatest density of redds at 164 redds but in 2010 the redd count in this stream decreased to 45. This lower count in the resident Cliff created a large decrease in the total count for the Imnaha count. Eighty percent of the total redds counted on the Imnaha in 2009 were from Cliff and in 2010 this percentage decreased to 41 percent. In 2010, 65 redds (59%) of the redds documented on the Imnaha were fluvial/resident redds as compared to 42 redds (20%) in 2009.




Figure 3. Comparison of bull trout surveyed redds and index redds (comparable miles) documented from 2001 to 2010 on the Imnaha River.

### *Big Sheep Creek*

Twenty total bull trout redds for 9.6 miles of survey were documented in 2010 on Big Sheep Creek and Lick Creek. These areas were surveyed twice in 2010, mid to late spawning season. The following data for Big Sheep Creek compares consistently surveyed index areas on Big Sheep and Lick Creek (7.6 miles) from 2000 to 2010 (Figure 4). The eleven-year average from 2000 to 2010 was 18 redds for the Big Sheep system. Total redd numbers within the Big Sheep system ranged from 8-34 within that period. Redd surveys for bull trout in the Big Sheep system have been limited in miles of survey (8.4 to 14.1 miles from 2000 to 2010) and in frequency, (2000-2001 surveys were conducted once late season, and in 2002, 2003, and 2005-2010, surveys were conducted

twice, mid and late season, except for lower Lick survey that was monitored only once in 2010 due to lack of surveyors). In 2004, the survey was conducted once late season for Big Sheep and twice, mid and late season for Lick Creek.




Figure 4. Comparison of bull trout surveyed redds and index redds (comparable miles) documented from 2000 to 2010 on Big Sheep and Lick Creeks.

The local bull trout populations in Big Sheep and Lick Creek appear to be relatively stable for the survey period (2000-2010), with a downward trend in the Imnaha River population in 2010. If you subtract the resident cliff creek population from the Upper Imnaha redd counts, there is a significant downward trend in both 2009 and 2010. A minimum of 15 years is needed to determine population trends.

The Imnaha population is one of the strongholds within the Imnaha Subbasin as it has multiple age classes, contains fluvial fish, has an anadromous prey base, has connectivity with the Snake River, and bull trout are distributed throughout the habitat. Primary spawning activity on the Imnaha River has been documented to occur in the headwaters which lie within wilderness. Both fluvial and resident life history forms are present. The Imnaha River is rated at low risk of extinction, and Big Sheep is rated “of special concern” (Buchanan et al. 1997).


## Sizes of Bull Trout Redds

Bull trout redds were measured using the same methodology in 2004 through 2010 and comparison of bull trout redd sizes (mean redd area ( $m^2$ )) for these years is illustrated below. There is a relationship between size of female salmonid and size of redd; large fish make large redds (Bjornn and Reiser 1991; P. Sankovich, Service, pers. comm., 2006). In addition, length/frequency distributions of mature resident bull trout and mature fluvial bull trout do not overlap, therefore, there is little overlap in size of redds (P. Sankovich, pers. comm., 2006).

### *Lostine/Bear/Goat*

Figure 6 compares bull trout redd sizes for the Lostine River, Bear Creek, and Goat Creek in 2004-2010. Mean redd area ( $m^2$ ) ranged from 0.9-1.4 for the Lostine, 0.3-0.9 for Goat Creek, and 0.2-1.0 for Bear Creek. The Bear Creek sample area was expanded in miles surveyed in 2007 (in past years the redd area was smaller and more typical of resident redds, in 2007 the redd area was larger and more typical of fluvial size bull trout redds. Brook trout are thought to be abundant in Bear Creek (due to historical stocking in the headwater lakes). Bull trout redds were not observed or documented in Bear Creek within the index area in 2006. In 2010, one bull trout redd was documented in Bear Creek. This solo redd was much larger than typical, 4.8 mean redd area, and was found among Chinook redds. The survey crew stated that this redd or two redds were superimposed among Chinook redds. It was reported as one fluvial bull trout redd. The Lostine River contains brook trout, but for most survey years, we had not observed brook trout spawning with bull trout. This observation changed in 2008 where they appeared to be spawning together and hybrid fish were observed. Mary Edwards, fisheries biologist with the Nez Perce Tribe, observed bull trout and brook trout spawning near the acclimation facility as early as late July 2010 while underwater snorkeling. Bull trout surveys conducted in September/October 2010 did not confirm any bull trout redds in this location.

To date, genetic sampling results of bull trout/brook trout in the Lostine and in Bear Creek has not occurred. Goat Creek is limited in available spawning habitat, but it appears to be the best available spawning habitat for fluvial fish in the Bear/Goat Creek system during drought years. (Several miles of upper Bear Creek were dry due to low snowpack and summer drought conditions). It appears from the data in 2004-2010 that redds in Goat Creek were a combination of resident and fluvial fish, as the redd sizes were midway between the Bear Creek (resident size) and the Lostine (fluvial size) (except for the fluvial-size redds in Bear Creek in 2007 - 2010. Two bull trout that were observed occupying a redd on Goat Creek in 2004 were <12 inches (<300 ml) and were on a redd that had an area of  $0.3 m^2$ ; therefore, this size of redd is resident (related to the fish size). More years of data collection on these streams should help us better understand the resident and fluvial life histories of bull trout in this area, relative to fish and redd sizes.



**Figure 6. Comparison of bull trout redd sizes [mean redd area (m<sup>2</sup>)] for Lostine River, Bear, and Goat Creeks sampled during bull trout spawning surveys, 2004-2010.**

Footnote: Bull trout redds were not observed in the index area of Bear Creek in 2006 and one extremely large fluvial redd observed in 2010 (not included on graph); (more of an outlier).

### *Imnaha and Big Sheep*

Figure 7 compares bull trout redd sizes for the sampled streams in the Imnaha system in years 2004-2010. Mean redd area (m<sup>2</sup>) ranged from 0.3-1.0 for Lick Creek, 0.1-0.8 for Big Sheep, 0.4-2.6 for Middle Imnaha, 1.2-1.8 for Upper Imnaha, 0.3-1.2 for N.F. Imnaha, 0.7-1.3 for S.F. Imnaha, and 0.1-0.4 for Cliff Creek. Middle Imnaha had only two redds observed in 2005 and three redds in 2004. In 2006-2008, Middle Imnaha had a large distribution of redds in this survey area (sample size = 18 for 2006 and 2007 and 28 for 2008) compared to past years. Middle Imnaha in 2004 and 2006 had primarily fluvial size redds with resident redds documented in 2005 and a mix of fluvial and resident redds in 2007 and 2008. As shown in Figure 7 below, very large redds (produced by large fluvial bull trout) were documented on the Middle Imnaha in 2008. Redd numbers decreased in the Middle Imnaha in 2009 and 2010, to 2 redds for 2009 and 4 for 2010; with a primarily fluvial component. Cliff Creek is a known resident system with a waterfall near the mouth. The survey on Cliff Creek in 2004 included a large fluvial size redd near the confluence with the S.F. Imnaha and therefore the mean redd size was higher than in 2005 and 2006 when no fluvial redds were observed in Cliff Creek below the waterfall. In 2010 two resident size redds were recorded on Cliff Creek below the falls. Lick Creek and Imnaha had an overlap of bull trout and chinook redds, which may make differentiation between the two sometimes difficult. Lick Creek data in 2004-2008 appeared to be a combination of resident and fluvial redds (with more resident redds documented in 2004, 2007-2010 and fluvial redds documented in 2005 and 2006). Big Sheep contained resident redds all years, and Upper Imnaha and S.F. Imnaha contained a majority of fluvial redds. North Fork Imnaha appears to have had more resident redds in 2004, 2007, and 2008, more fluvial size redds in 2005 and 2010, and a mix of fluvial and resident in 2006 and 2009. Cliff Creek is a known resident system with a mean redd size of 0.2 m<sup>2</sup> in 2010. Mean redd size was greater, a size of

0.3 m<sup>2</sup> in 2004 as a result of a fluvial redd near the mouth and potential superimposition of redds above the barrier. 2007 mean redd size included a few larger redds below the falls (near the mouth) which are presumed to be fluvial redds. 2008 mean redd size was greatest above the falls, likely superimpositions of redds above the barrier. In 2008 not all the redds were measured above the falls (31 of 52 total redds, 59.6%) and all redds were measured below the falls (1 of 1, 100%). The sample size in 2008 was large enough to get a good estimate of sizes of redds above and below the falls and at the same time completing the survey in a reasonable amount of time (while redds were still visible during daylight). Refer to Tables 4a and 4b in Appendix B for additional information on 2010 bull trout redd characteristics.





Figure 7. Comparison of bull trout redd sizes [mean redd area (m<sup>2</sup>)] for sampled streams in the Imnaha Subbasin, 2004-2010.

### Bull Trout Redd Distribution on the Lostine River

The bull trout spawning surveys on the Lostine River in years 2005 through 2010 included collection of UTM coordinate data on the spatial distribution of the bull trout redds observed along the Lostine River. Bull trout redds on the Lostine River (as well as in other surveyed streams) were often arranged in complexes (several redds in close proximity to each other). Redds were primarily located in the French Camp to Shady Falls and Bowman to French Camp reaches along the Lostine River in both 2005 and 2006; and mainly in the French Camp to Shady Falls reach in 2007 and 2010 and several miles (approximately 10 miles) downstream of these reaches at the Six Mile Bridge to Pole Bridge reach. High site fidelity is documented by the overlapping of bull trout redds from 2005-2010, especially in the upper reaches of the Lostine River. The distribution of bull trout redds in 2009 was documented to be further upstream than in past years. In 2010, redd distribution was spread over the majority of the survey area (except the lower Acclimation to Lostine River Ranch reach) but had the highest numbers in the upper two reaches (Shady to French and French to Bowman) and the Pole Bridge reach. In 2010, few redds were documented in the Williamson to Walla Walla Reach (similar to past

years except in 2008 and 2009, where several redds were documented in this location). In 2010, one redd was observed in the Lundquist to Acclimation reach.

Sections of the Lostine River not surveyed due to insufficient spawning gravels (boulder and cobble dominated substrate) and difficult access include; Bowman to Walla Walla (approximately 2.2 miles) and Williamson to Pole Bridge (approximately 3.5 miles). Downstream of Westside Ditch on the Lostine River (approximately 9 miles) is private property that is not surveyed due to lack of bull trout spawning gravels, higher stream temperatures, and low flows associated with irrigation withdrawal.



**Figure 8. Map of the Lostine River showing bull trout redd survey reaches and bull trout redd locations in years 2006 through 2010.**



## DISCUSSION

A combination of low flows (due to drought conditions) and severe streambed scour activity in the spring seasons of 2006-2010 created limited available habitat for spawning bull trout in the fall seasons of 2006-2010. This is noted by the author as most evident in the Lostine/Bear and Imnaha systems. During drought conditions, as in 2006-2010, Bear Creek had limited spawning habitat for fluvial bull trout due to the subbing out of flows. It is assumed that there is habitat upstream of the index area above subbing flows for resident bull trout. In 2009 and 2010, the author observed Bear Creek (from Standley trail crossing to Goat Creek confluence) having limited spawning habitat and more rearing habitat due to low flows (concentrated on one or both sides of a large floodplain and large woody debris creating pool habitat in these areas). In 2007, surveyors noted a chinook redd (obvious because of the huge size and large substrate) in lower Goat Creek which was probably a result of extremely low flows and warm temperatures in Bear Creek in 2007. Several side channels on the Imnaha were no longer flowing in 2006-2009, and this was evident as well during the chinook surveys on these streams. This likely had an effect on densities and distribution of redds for both bull trout and chinook species in these systems. In 2010, the Upper Imnaha had more flow than in 2009, and algae on the rocks (potentially didymo or rock snot) that was observed in 2009, disappeared in 2010. There appeared to be some severe scour flows that likely scoured away the algae. Stream walking on the S.F. Imnaha was much easier in 2010 as a result.

The interactions between fluvial bull trout and chinook salmon during spawning time (in particular, where spawning locations and timing may overlap) are not well understood. The author estimates that these salmonid species (in our local streams, including the Lostine, Imnaha, Lick Creek, and Bear Creek) compete for available spawning gravels. Fluvial bull trout migrate upstream to cooler waters and where there is less competition with larger chinook salmon for holding and spawning areas. In 2010, the spring/summer chinook counts in the Grande Ronde and Imnaha systems were very high. In the Lostine, the unexpanded chinook redd counts had a high of 696 total redds compared to 258 in 2009. This is the highest count from eleven years of data (2000-2010) as reported in the 2010 chinook spawning ground summary results (Feldhaus 2010). Ninety of these redds (13 percent) were documented in the French Camp to Bowman Trailhead Reach. One hundred and thirty one (19 percent) were reported in the Walla Walla to Williamson Reach. Eleven chinook redds (2 percent) were documented in the Pole Bridge to Six Mile Bridge Reach. In 2010, the highest numbers of chinook redds were recorded in the lower bull trout reaches; Six Mile Bridge to Acclimation had one hundred and sixty eight redds (24 percent); and Acclimation to Lostine River Ranch had two hundred and twenty two (32 percent). The Shady Falls to French Camp is not a reach that has been annually surveyed for chinook. In 2010, a few chinook redds were noted in this reach during bull trout spawning surveys; most likely due to the large run of fish, they spawned further upstream. The high chinook run in the Lostine likely affected numbers and distribution of bull trout spawning in the Lostine and potentially elsewhere such as Blue Hole to Indian Crossing Reach of the Imnaha.

The Imnaha, in 2006-2008, had a large distribution of bull trout redds that extended further downstream than in past years (Blue Hole to Indian Crossing, a two mile section).

This may be due to limited passage of fluvial bull trout at the falls and potential bull trout/chinook interactions that are not well understood. In 2008, the redd size was larger; therefore, larger fluvial fish were building redds in this area. As mentioned in the previous reports, additional research on timing in the Imnaha and especially passage above and below the falls would be beneficial in our understanding of bull trout in this system. In 2009 and 2010, the Imnaha redd distribution extended further upstream (similar to pre-2006) and total redd counts (excluding resident Cliff Creek) were down in 2009 and 2010, compared to previous years (with 2010 having a slightly higher count than 2009, if you subtract Cliff Creek resident population data). In 2009, Cliff Creek bull trout surveyors documented that a recent avalanche deposited several pieces of wood into the channel. High flows in 2009 channelized the creek bed and created large jams. In 2010, surveyors on Cliff Creek documented bull trout that were still creating redds, so the count is likely lower than the final count. This is one of the disadvantages to a one time survey, but due to access, cost, and lack of experienced surveyors late in the season, the Upper Imnaha survey is likely to remain a one-time survey.

In 2010, the Middle Fork Imnaha River, above the falls to forks, for approximately 0.6 miles, had an exploratory survey conducted to determine bull trout and redds presence or absence. The habitat was excellent but no bull trout or redds were reported.



**Middle Fork Imnaha habitat above the falls**



**Middle Fork Imnaha habitat above the falls, gravel substrate but no fish**

The Middle Fork of Big Sheep (a one mile section of stream in the headwaters) was surveyed in 2008 per recommendation by Service researchers. They have documented several bull trout in that area post their survey of the area utilizing electrofishing techniques. It appears to be excellent rearing habitat, but in 2008 there were very limited spawning gravels and/or use of existing gravels by spawning resident bull trout. This survey was not repeated in 2009 or 2010.

Quartz Creek, a tributary to Lick Creek also had an experimental survey completed in 2008 per Service researcher recommendations. There appears to be bull trout spawning and rearing habitat in this stream, as six redds were reported in a one mile survey reach of stream. The stream is very healthy, i.e. excellent canopy cover/streamside vegetation; therefore, the stream is not recommended to survey on an annual basis due to difficult access and a problem in detecting redds due to the amount of cover. This survey was not conducted in 2009 or 2010.

In late July 2010, after a heavy rainstorm (on July 27<sup>th</sup>), Doc Creek, a tributary to Bear Creek contributed large quantities of mud to downstream Bear Creek. This reach of Bear Creek is downstream of bull trout index spawning areas in Bear and tributary Goat Creek. Bill Knox (ODFW) sent an email on July 30, 2010 concerning the mudflow in Bear Creek and documented large quantities of mud and a few dead fish at the site. No bull trout were confirmed dead from this mud event, but dead fish could have been buried by the mud. US Forest Service personnel flew over upper Doc Creek in early August 2010 and reported that the mud did not appear to originate from one location (debris flow scenario) as one would expect from such large quantities of mud, but instead from several small rills across the previous year's Big Sheep Ridge wildfire area. (Rill erosion is caused by a concentration of flowing water). This area had some areas of bare soil and extensive riling was evident in these areas. It is highly likely that bull trout migrating

upstream to spawning areas from this reach may have been affected by this mudflow event.



**Bear Creek upstream of trail Bridge near boundary camp-ground with muddy stream bottom from August 2010 debris flow in Doc Creek (an upstream trib.)**



**Close-up of mud filled substrate at Bear Creek below trail bridge, 2010**

Deer Creek had an experimental survey conducted in 2009. No redds or bull trout were located. The surveyors recommended continuing this survey in 2010 upstream of the Rd 8920 culvert for approximately one mile. A recent brushed wildfire fire line near the creek made walking the creek much easier than through the dense jungle-like vegetation downstream. The recommendation was to survey this stream perhaps not every year, but often enough to get some trend information. In 2010, an experimental survey was conducted twice from the Rd. 8920 culvert to 0.8 miles upstream and proved to be worth the walk. Twelve resident size redds and thirty nine bull trout were reported in this Deer Creek survey reach. The plan is to continue to survey this reach, dependent on available surveyors.

Overall, future needs for this project include continued funding and support from all involved parties for conducting and reporting bull trout redd counts in the Wallowa Mountains. A phase II of this project, OWEB Proposal: Phase II Wallowa Mountains Bull Trout Redd Monitoring (2010-2011) was submitted for funding in October 2009 and received funding for 2010 and 2011.

At least 15 years (consecutive years) of bull trout redd data are needed for trend data (Maxwell 1999) and for bull trout recovery data needs. Caution must be exercised in using the above bull trout spawning data for adult population trends until at least 15 years of data have been collected. However, we are getting much closer to this goal of a minimum of 15 years!

## CONCLUSION

The local bull trout populations surveyed appear to be relatively stable for the survey period (1999-2008) with some recent downward trend in 2009 and 2010.

The Imnaha population is one of the strongholds within the Imnaha Subbasin as it has multiple age classes, contains fluvial fish, has an anadromous prey base and connectivity with the Snake River, and bull trout are distributed throughout the habitat. Primary spawning activity on the Imnaha River has been documented to occur in the headwaters which lie within wilderness. Both fluvial and resident life history forms are present. The Imnaha River is rated at low risk of extinction, and Big Sheep is rated “of special concern” (Buchanan et al. 1997).

The Lostine River is considered a moderately-strong population within the Grande Ronde Subbasin. Our results are consistent with Buchanan et al. (1997). Lostine River and Bear Creek contain brook trout and the degree of hybridization is unknown; although recent photos and documentation by Mary Edwards (NPT) suggests bull trout likely paring with brook trout and hybridization more likely than past information has shown. Limited redd count data is available on Bear Creek and this portion of the Lostine River/Bear Creek local population has been listed as a special concern by Ratliff and Howell (1992). Future genetic analysis of bull trout and brook trout is recommended to help determine the significance of this threat.



Male adult fluvial bull trout near redd on the Lostine River near Turkey Flat, 2009




Bull trout caught by angling, Kyle Bratcher ODFW, 2010

## REFERENCES

- Al-Chokhachy, R., P. Budy, and H. Schaller. 2005. Understanding the significance of redd Counts: a comparison between two methods for estimating the abundance of and monitoring bull trout populations. *North American Journal of Fisheries Management* 25:1505-1512.
- Bellerud, B.L., S. Gunkel, A.R. Hemmingsen, D.V. Buchanan, and P.J. Howell. 1997. Bull Trout Life History, Genetics, Habitat Needs, and Limiting Factors in Central and Northeast Oregon. 1996 Annual Report. Project Number 95-54. Bonneville Power Administration, Portland, OR.
- Bjornn, T.C., and D.W. Reiser. 1991. Habitat requirements of salmonids in streams. *American Fisheries Society Special Publications* 19:139-179.
- Bonar, S.A., M. Divens, and B. Bolding. 1997. Methods for sampling the distribution and abundance of bull trout/Dolly Varden. Washington Department of Fish and Wildlife Research Report No. RAD97-05. Olympia, WA. 48 p.
- Buchanan, D.V., M.L. Hanson, and R.M. Hooton. 1997. Status of Oregon's Bull Trout. Oregon Department of Fish and Wildlife, Portland, Oregon.
- Dunham, J., B. Rieman, and K. Davis. 2001. Sources and Magnitude of Sampling Error in Redd Counts for Bull Trout. *North American Journal of Fisheries Management* 21:343-352.
- Ecovista. 2004. Salmon Subbasin Management Plan. For the Nez Perce Tribe Watershed Division and Shoshone Bannock Tribe as part of Northwest Power and Conservation Council's Fish and Wildlife Program.
- Ecovista. 2004a. Imnaha Subbasin Plan. Plan includes Assessment, Inventory, and Management Plan. For Nez Perce Tribe as part of Northwest Power and Conservation Council's Fish and Wildlife Program.
- Feldhaus, Joseph. 2010. 2010 Grande Ronde and Imnaha River Basin chinook salmon spawning ground survey results. ODFW Research Office, La Grande, OR 97850. December 30, 2010.
- Knox, B. 2010. Email concerning mud flow in Bear Creek from July 27, 2010 rainstorm. ODFW, Enterprise, Oregon. July 30, 2010.
- Krueger, C. C., and D. J. Decker. 1993. The process of fisheries management. Pages 33-54 in C. C. Kohler and W. A. Hubert, editors. *Inland fisheries management in North America*. American Fisheries Society, Bethesda, Maryland.
- Maxwell, B.A. 1999. A power analysis on the monitoring of bull trout stocks using redd counts. *North American Journal of Fisheries Management*. 19: 860-866.

- Mendel, G., M. Gembala, J. Trump, and C. Fulton. 2006. Baseline Assessment of Salmonids in Tributaries of the Snake and Grande Ronde Rivers in Southeast Washington. 2005 Annual Report. Washington Department of Fish and Wildlife, Dayton, WA.
- Nowak, C. M. and 25 co-authoring agencies. 2004. Grande Ronde Subbasin Plan. Prepared for the Northwest Power and Conservation Council.
- Ratliff, D.E. and P.J. Howell. 1992. The status of bull trout populations in Oregon. Pages 10 to 17 In: P.J. Howell and D.V. Buchanan, eds. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon.
- Rieman, B.E. and D.L. Myers 1997. Use of redd counts to detect trends in bull trout (*Salvelinus confluentus*) populations. *Conservation Biology* 11:1015-1018.
- Sankovich, P. m., S.L. Gunkel, A.R. Hemmingsen, I.A. Tattam, and P.J. Howell. 2003. Migratory patterns, structure, abundance, and status of bull trout populations from subbasins in the Columbia Plateau. 2002 Annual Report. Project 199405400. Bonneville Power Administration, Portland, OR.
- Starcevich, S.J., S. Jacobs, and P.J. Howell. 2005. Migratory Patterns, Structure, Abundance, and Status of Bull Trout Populations from Subbasins in the Columbia Plateau and Blue Mountain Provinces. 2004 Annual Report. Project 199405400. Bonneville Power Administration, Portland, OR.
- Taper, M.L., D.F. Staples, and B.B. Shepard. 2005. Observer Error Structure in Bull Trout Redd Counts in Montana Stream: Implications for Inference on True Redd Numbers. *North American Journal of Fisheries Management*, in-review. Presented as a Power Point Presentation at ScCS meeting in Glacier Park Montana, September 7-9, 2005.
- U.S. Department of the Interior, Fish and Wildlife Service. 2002. Chapter 11, Grande Ronde River Recovery Unit, Oregon and Washington and Chapter 12, Imnaha-Snake Rivers Recovery Unit. In: U.S. Fish and Wildlife Service. Bull Trout (*Salvelinus confluentus*) Draft Recovery Plan. Portland, Oregon.
- U.S. Fish and Wildlife Service. 1999 – 2010. Unpublished bull trout spawning survey data on file at La Grande Field Office, U.S. Fish and Wildlife Service, La Grande, Oregon.

**APPENDIX A – PHOTOS TAKEN AT SURVEY LOCATIONS**



**Lostine River at a bend in the river and view up the Wallowa Mountains**



**Bear Creek**



**Western Toad near Goat Creek, 2010**



**From left to right, Ashley Davidson (ODFW), Lynne Price (consultant), and Eric Shoudel (ODFW) post Bear Creek survey, 2010**



**2010 Upper Imnaha survey crew, author on left, and to her right; Ian Wilson (NPT), Ken Bronec (consultant), Jeff Yanke (ODFW), Lynne Price (consultant), and Brad Smith (consultant). Alan Miller not in photo.**



**Upper Imnaha Survey, Imnaha River, 2010**



**Bull trout fluvial redds on stream margin in the Upper Imnaha Falls to Lower Imnaha Falls Reach, Imnaha River , 2010**



**Ken Bronce and Lynne Price (consultants) surveying the Lostine River and enjoying their day!**



**Kyle Bratcher (ODFW) last survey day on Lostine River, retrieving several Chinook survey flags while surveying for bull trout redds, 2010**



**Ian Wilson (NPT) surveying S.F. Imnaha, past scour flows evident by eroding stream banks, 2010**



**Dead brook trout observed on Lostine River, 2010**



**A Little Brown Bat observed by surveyors flying (during sunny daylight hours) and drowning post dipping into the SF Innaha at Blue Creek trail crossing**

**APPENDIX B – TABLES**

**Table 1 – Bull Trout Spawning Surveys and Survey Frequencies for selected Grande Ronde River and Imnaha River Streams, 1999-2010**

<b>Stream</b>	<b>Year</b>	<b>Dates</b>	<b>Survey Frequency</b>	<b>Total Redds</b>	<b>Total Miles</b>	<b>Total Redds/Mile</b>
Lostine River	1999	9/16,9/23,10/12	3 Times	39	9.75	4.0
	2000	9/21,9/28,10/12	3 Times	38	13.74	2.8
	2001	9/17-18,10/11-12	Twice	43	14.4	3.0
	2002	9/23-24,10/7-8	Twice	22	10.7	2.1
	2003	9/23-24,10/6-7	Twice	71	10.5	6.8
	2004	9/14-15,10/5-6	Twice	26	8.5	3.1
	2005	9/15, 9/21-22, 10/3-10/4	Twice, and 3 Times in Turkey Flat and Shady Campground areas	32	10.5	3.0
	2006	9/14, 9/20-21, 10/2-10/4	Twice, and 3 Times in Turkey Flat and Shady Campground areas	45	10.5	4.3
	2007	9/19-9/20, 10/3-10/4	Twice	47	10.1	4.7
	2008	9/17, 10/1-10/2, 10/9-10/10	Twice, and 3 Times in Turkey Flat and Shady Campground	53	10.1	5.3
	2009	9/11, 9/23-24, 10/7-10/8	Twice, and 3 Times in French Camp to Bowman	41	10.1	5.2
	2010	9/22, 10/6, 10/7	Twice	36	10.1	3.6
Bear Creek (including Goat Cr)	1999	9/7,9/22	Once Bear, Twice Goat	6	1.8	3.3
	2000	10/18	Once	5	1.8	2.8
	2001	10/16	Once	12	2.3	5.2
	2002	10/15	Once	7	2.3	3.0
	2003	10/16	Once	17	3.8	4.5
	2004	10/1	Once	11	2.3	4.8
	2005	10/11	Once	16	2.8	5.7
	2006	10/10	Once	9	1.9	4.7
	2007	9/17, 10/9	Twice, Exploratory (more reaches than past years)	11.0	7.2	1.5
	2008	9/30, 10/7	Twice	20	3.2	6.3

	2009	9/22, 10/13	Twice	8	3.2	2.5
	2010	9/16, 10/13	Twice	7	3.2	2.2
Deer Creek	2009	10/14	Once	0	1.5	0
	2010	9/17, 10/4	Twice	12	0.8	15

**Table 1 (Continued)– Bull Trout Spawning Surveys and Survey Frequencies for Selected Grande Ronde River and Imnaha River Streams, 1999-2010**

<b>Stream</b>	<b>Year</b>	<b>Dates</b>	<b>Survey Frequency</b>	<b>Total Redds</b>	<b>Total Miles</b>	<b>Total Redds/Mile</b>
Imnaha River	1999	9/20,28,10/11	Middle = Thrice	14	15.2	0.9
(excluding Big Sheep)	2000	9/20,22,25,26,27,10/11	Upper = Once, Middle = Twice	92	29.1	3.2
	2001	9/20,21,10/1,2,3,9,10	Upper = Once, Middle = Twice	291	31.3	9.3
	2002	9/25,26,9/30,10/1-2,10/10-11	Upper = Once, Middle = Twice	113	30.5	3.7
	2003	9/25-26,9/29-30,10/1,10/8-9	Upper = Once, Middle = Twice	266	31.6	8.4
	2004	9/15-9/16,9-27,28,29,10/7-8	Upper = Once, Middle = Twice	293	31.5	9.3
Middle=Blue Hole to Indian	2005	9/26-28, 10/7	Once Upper and Middle	276	19.4	14.2
2005-2010	2006	9/25-28,10/5	Upper = Once, Middle = Twice	186	19.4	9.6
	2007	9/24-27, 10/5	Upper = Once, Middle = Twice	284	19.4	14.6
	2008	9/22-25, 10/6	Upper = Once, Middle = Twice	190	19.4	9.8
	2009	9/28-30, 10/1, 10/9	Upper = Once, Middle = Twice	235	19.4	12.1

**Table 1 (Continued)– Bull Trout Spawning Surveys and Survey Frequencies for Selected Grande Ronde River and Imnaha River Streams, 1999-2010**

<b>Stream</b>	<b>Year</b>	<b>Dates</b>	<b>Survey Frequency</b>	<b>Total Redds</b>	<b>Total Miles</b>	<b>Total Redds/Mile</b>
	2010	9/27-29, 9/30, 10/8	Upper = Once, Middle = Twice	133	19.4	6.9
Big Sheep	1999	9/21,29,30,10/18,19	Once	20	14.2	1.4
[(including Lick,	2000	10/13-10/16	Once	12	8.4	1.4
and Salt).	2001	10/14,10/17	Once	24	8.4	2.9
Salt Cr. not	2002	9/30,10/1,10/14-15	Twice	41	9.3	4.4
surveyed	2003	9/22,10/14	Twice	18	9.3	1.9
post 2003,						
and Upper	2004	9/20,9/28-9/29,9/30,10/4,10/19	Once Big Sheep, Twice Lick	43	14.1	3.0
Big						
Sheep	2005	9/19-20, 10/6	Twice	16	8.6	1.9
exploratory						
in 2004].						
	2006	9/19, 10/4	Twice	18	7.6	2.4
	2007	9/21, 10/2	Twice	27	8.6	3.1
	2008	9/29, 10/1, 10/10	Twice except once exploratory Quartz creek tributary to Lick and once Lick Creek 39 Rd to meadow	30	9.1	3.3
	2009	9/16, 10/6	Twice	20	7.6	2.6
	2010	9/15, 10/5	Twice except once Lick Creek 39 Rd. to Meadow	20	7.6	2.6

**Table 2a–Bull Trout Spawning Surveys for the Lostine River Comparing 1999 to 2010 Surveys**

Stream	Survey Years											
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>Lostine</b>												
	<b>Redds Surveyed</b>											
<b>Reaches (miles surveyed)</b>												
Lundquist Bridge to OC Ranch (2.8)	1	0	2	3	3	5	0	5	4	5	0	1
Williamson to Walla Walla (2.2 miles)	0	2	1	0	6	1	3	0	2	13	8	2
Bowman to French Camp (1.6 miles)	18	19	16	11	18	3	9	9	5	12	7	6
French Camp to Shady Falls (1.5)	20	12.0	23	8	43	17	12	22	31	20	23	21
<b>Lostine Total Redds (Comparable Reaches)</b>	<b>39</b>	<b>33</b>	<b>42</b>	<b>22</b>	<b>70</b>	<b>26</b>	<b>24</b>	<b>36</b>	<b>42</b>	<b>50</b>	<b>38</b>	<b>30</b>
<b>Lostine Total Miles of Comparable Stream</b>	8.1	8.1	8.1	8.1	8.1	8.1	8.1	8.1	8.1	8.1	8.1	8.1
<b>Lostine Redds/Mile Comparable Stream</b>	<b>4.6</b>	<b>3.9</b>	<b>4.9</b>	<b>2.6</b>	<b>8.2</b>	<b>3.1</b>	<b>2.8</b>	<b>4.2</b>	<b>5.2</b>	<b>6.2</b>	<b>4.7</b>	<b>3.7</b>
<b>Total Redds For Year</b>	39	38.0	43.0	22.0	71.0	26.0	32.0	45	47	53	41	36
<b>Total Miles Surveyed For Year</b>	9.8	13.7	14.4	10.7	10.5	8.5	10.5	10.5	10.1	10.1	10.1	10.1
<b>Total Redds/Mile For Year</b>	4.0	2.8	3.0	2.1	6.8	3.1	3.0	4.3	4.6	5.2	4.1	3.6

Notes: The Lostine was surveyed three times in 1999 and 2000. During survey years 2001-2010, the Lostine was surveyed twice (except Shady Campground and Turkey Flat areas were surveyed three times in 2005, 2006, and 2008 and French Camp to Bowman was surveyed three times in 2009). Dates of Lostine bull trout spawning surveys generally commenced as early as the second or third week in September and the last survey was conducted in the first or second week in October.

**Table 2b – Bull Trout Spawning Surveys for Bear and Goat Creeks Comparing 1999 – 2010 Surveys**

Stream	Survey Years											
	99	00	01	02	03	04	05	06	07	08	09	10
<b>Bear Creek</b>			<b>Redds</b>	<b>Surveyed</b>								
<b>Reaches (miles surveyed)</b>												
Bear: Goat Confluence to Wilderness Boundary (1mile)	0	2	3	1	2	3	5	0	1	6	1	1
Goat Creek: Mouth to Falls (0.9)	8	3	9	6	9	8	6	9	7	4	5	6
<b>Bear (and Goat) Total Redds (Comparable Reaches)</b>	<b>8</b>	<b>5</b>	<b>12</b>	<b>7</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>6</b>	<b>7</b>
<b>Bear Creek Total Miles of Comparable Stream</b>	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9
<b>Bear Creek Redds/Mile Comparable Stream</b>	<b>4.2</b>	<b>2.6</b>	<b>6.3</b>	<b>3.7</b>	<b>5.8</b>	<b>5.8</b>	<b>5.8</b>	<b>4.7</b>	<b>4.2</b>	<b>5.3</b>	<b>3.2</b>	<b>3.2</b>
<b>Total Redds For Year</b>	8	5	12	7	17	12	16	9	11	20	8	7
<b>Total Miles Surveyed For Year</b>	1.9	1.9	2.3	2.3	3.8	2.3	2.8	1.9	7.2	3.2	3.2	3.2
<b>Total Redds/Mile For Year</b>	4.2	2.6	5.2	3.0	4.5	5.2	5.7	4.7	1.5	6.3	2.5	2.2

Notes: These surveys were conducted once from 1999-2006, usually late in the spawning season, the first or second week in October [except in 1999, surveys were conducted in September (on 9/7 and 9/22)]. In 2007, the surveys included several additional “experimental” miles and were conducted twice in the spawning season, once in mid-September and once in early October. In 2008 - 2010, the surveys were conducted twice in the spawning season and an additional 1.4 miles of Bear Creek was surveyed upstream of the comparable reach.

**Table 2c – Bull Trout Spawning Surveys for the Innaha River, Comparing 2001 – 2010 Surveys**

Stream	Survey Years									
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>Innaha River</b>	<b>Redds Surveyed</b>									
<b>Reaches (miles surveyed)</b>										
<b>South Fork Innaha and tributaries:</b>										
Cliff Creek, mouth to 2.5 miles (2.5 miles)	96	22	57	65	61	17	93	52	164	45
South Fork Innaha, NF to Soldier (1.5 miles)	6	7	14	12	44	9	30	6	7	2
South Fork Innaha, Soldier to Cliff (3.1 miles)	33	18	37	29	55	26	37	15	8	27
<b>North Fork Innaha:</b>										
North Fork, above Middle Fork (4.1 miles)	49	18	40	68	39	18	30	17	2	4
North Fork, below Middle Fork to mouth (2.1 miles)	2	8	15	9	21	6	7	5	3	12
Middle Fork, mouth to falls (0.8 miles)	12	0	12	6	24	7	17	8	7	5
<b>Innaha River:</b>										
Innaha River, NF to Falls (0.6 miles)	0	3	5	1	2	3	2	1	0	0
Innaha River, Falls to lower falls (0.8 miles)	41	18	35	40	13	37	28	12	13	11
Innaha River, Blue Hole to Indian Crossing (2.0 miles)	8	7	9	3	2	18	18	28	2	4
<b>Innaha Total Redds (Comparable Reaches)</b>	<b>247</b>	<b>101</b>	<b>224</b>	<b>233</b>	<b>261</b>	<b>141</b>	<b>262</b>	<b>144</b>	<b>206</b>	<b>110</b>
<b>Innaha Total Miles of Comparable Stream</b>	17.5	17.5	17.5	17.5	17.5	17.5	17.5	17.5	17.5	17.5
<b>Innaha Redds/Mile Comparable Stream</b>	<b>14.1</b>	<b>5.8</b>	<b>12.8</b>	<b>13.3</b>	<b>14.9</b>	<b>8.1</b>	<b>15.0</b>	<b>8.23</b>	<b>11.8</b>	<b>6.3</b>
<b>Total Redds For Year</b>	269	103	293	286	276	186	284	184	235	133
<b>Total Miles Surveyed For Year</b>	19.4	18.3	42.8	41.2	19.4	19.4	19.4	19.4	19.4	19.4
<b>Total Redds/Mile For Year</b>	13.9	5.6	6.8	6.9	14.2	9.6	14.6	9.5	12.1	6.9

Notes:

All reaches except Blue Hole to Indian Crossing were surveyed once in 2001 to 2010.

The Blue Hole to Indian Crossing reach was surveyed twice, in mid September and October from 2001-2010, except in 2005 it was surveyed once.

**Table 2d – Bull Trout Spawning Surveys for Big Sheep Creek and Lick Creek, Comparing 2001 – 2010 Surveys**

Stream	Survey Years										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>Big Sheep Creek (including Lick Creek)</b>		<b>Redds</b>	<b>Surveyed</b>								
<b>Reaches (miles surveyed)</b>											
Big Sheep, canal to 39 rd. (1.9 miles)	2	6	17	2	3	5	6	12	3	2	8
Lick Creek, Meadow to 39 rd. (1.5 miles)	0	6	3	0	1	3	5	3	4	5	7
Lick Creek, 39 rd. to Quartz Creek (4.2 miles)	8	5	14	8	4	1	7	12	15	13	5
<b>Big Sheep Total Redds (Comparable Reaches)</b>	<b>10</b>	<b>17</b>	<b>34</b>	<b>10</b>	<b>8</b>	<b>9</b>	<b>18</b>	<b>27</b>	<b>22</b>	<b>20</b>	<b>20</b>
<b>Big Sheep Creek Total Miles of Comparable Stream</b>	7.6	7.6	7.6	7.6	7.6	7.6	7.6	7.6	7.6	7.6	7.6
<b>Big Sheep Creek Redds/Mile Comparable Stream</b>	<b>1.3</b>	<b>2.2</b>	<b>4.5</b>	<b>1.3</b>	<b>1.1</b>	<b>1.2</b>	<b>2.4</b>	<b>4.6</b>	<b>2.9</b>	<b>2.6</b>	<b>2.6</b>
<b>Total Redds For Year</b>	12	24	41	18	43	16	18	27	30	20	20
<b>Total Miles Surveyed For Year</b>	8.4	8.4	9.3	9.3	14.1	8.6	7.6	8.6	9.6	7.6	7.6
<b>Total Redds/Mile For Year</b>	1.4	2.9	4.4	1.9	3.0	1.9	2.4	3.1	3.1	2.6	2.6

Notes: Survey frequency varied by year, surveys were conducted once in mid to late October in years 2000 and 2001 for both Big Sheep and Lick Creek, and surveys were conducted twice, once in September and once in October in years 2002-2010, except for Big Sheep which was surveyed once in 2004; and Lick Creek (Meadow to 39 Rd.) was surveyed once in 2010.

Bull Trout Spawning Surveys  
For the Imnaha River, 2010  
USFWS, La Grande Field Office

Page 1 of 2

Imnaha Basin Stream Reach, Section	Date(s)	Kilometers Surveyed	Miles Surveyed	Occ	Redds			Per km	Per Mile	Total But obs	Bull Trout Observed (mm)			
					Unocc	Total					<6"(150mm)	<12" (~300mm)	<18"(450mm)	>18"(450mm)
<b>Upper Imnaha System</b>														
<b>South Fork Tributaries</b>														
Cliff Cr., mouth to 3.6 km	28-Sep	4.0	2.5	20.0	25.0	45.0	11.3	18.0	75.0	15.0	60.0	0.0	0.0	
<b>South Fork Tributaries Total</b>		<b>4.0</b>	<b>2.5</b>	<b>20.0</b>	<b>25.0</b>	<b>45.0</b>	<b>11.3</b>	<b>18.0</b>	<b>75.0</b>	<b>15.0</b>	<b>60.0</b>	<b>0.0</b>	<b>0.0</b>	
<b>North Fork</b>														
M.F., upstm of falls to forks (exploratory survey)	28-Sep	1.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Middle Fork., mouth to falls	28-Sep	1.3	0.8	0.0	5.0	5.0	3.8	6.2	2.0	1.0	1.0	0.0	0.0	
N. Fk., above M. Fk. (reach 3-7) 8.0	27-Sep	6.6	4.1	0.0	4.0	4.0	0.6	1.0	1.0	1.0	0.0	0.0	0.0	
N. Fk., below M. Fk. (reach 1-2)	27-Sep	3.4	2.1	2.0	10.0	12.0	3.5	5.7	4.0	0.0	0.0	4.0	0.0	
<b>North Fork Total</b>		<b>11.3</b>	<b>7.0</b>	<b>2.0</b>	<b>19.0</b>	<b>21.0</b>	<b>1.9</b>	<b>3.0</b>	<b>7.0</b>	<b>2.0</b>	<b>1.0</b>	<b>4.0</b>	<b>0.0</b>	
<b>South Fork</b>														
S. Fk., North Fork to Soldier Cr.	27-Sep	2.4	1.5	0.0	2.0	2.0	0.8	1.3	0.0	0.0	0.0	0.0	0.0	
S. Fk., Soldier to Cliff Cr.	28-Sep	5.0	3.1	0.0	27.0	27.0	5.4	8.7	1.0	1.0	0.0	0.0	0.0	
<b>South Fork Total</b>		<b>7.4</b>	<b>4.6</b>	<b>0.0</b>	<b>29.0</b>	<b>29.0</b>	<b>3.9</b>	<b>6.3</b>	<b>1.0</b>	<b>1.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	
<b>Upper Imnaha</b>														
Upper Imnaha (Falls to North Fork)	29-Sep	1.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Upper Imnaha Falls to lower falls	29-Sep	1.3	0.8	3.0	8.0	11.0	8.5	13.6	14.0	0.0	0.0	10.0	4.0	
Falls downstream .67 mi. to beg. of gorge	29-Sep	1.1	0.7	3.0	13.0	16.0	14.5	23.4	9.0	0.0	0.0	1.0	8.0	
Lower end of gorge to next gorge (.25 mi)	29-Sep	0.4	0.2	0.0	3.0	3.0	7.5	12.1	0.0	0.0	0.0	0.0	0.0	
Canyon above slide to canyon just above slide	29-Sep	1.5	0.9	0.0	4.0	4.0	2.7	4.3	0.0	0.0	0.0	0.0	0.0	
<b>Upper Imnaha Total</b>		<b>5.3</b>	<b>3.3</b>	<b>6.0</b>	<b>28.0</b>	<b>34.0</b>	<b>6.4</b>	<b>10.3</b>	<b>23.0</b>	<b>0.0</b>	<b>0.0</b>	<b>11.0</b>	<b>12.0</b>	
<b>Upper Imnaha System Total</b>														
		<b>28.0</b>	<b>17.4</b>	<b>28.0</b>	<b>101.0</b>	<b>129.0</b>	<b>4.6</b>	<b>7.4</b>	<b>106.0</b>	<b>18.0</b>	<b>61.0</b>	<b>15.0</b>	<b>12.0</b>	
<b>Imnaha Basin Total (Page 2)</b>		<b>15.5</b>	<b>9.6</b>	<b>2.0</b>	<b>22.0</b>	<b>24.0</b>	<b>1.5</b>	<b>2.5</b>	<b>44.0</b>	<b>19.0</b>	<b>12.0</b>	<b>2.0</b>	<b>11.0</b>	
<b>Imnaha Basin Total Pages 1 &amp; 2)</b>		<b>43.5</b>	<b>27.0</b>	<b>30.0</b>	<b>123.0</b>	<b>153.0</b>	<b>0.5</b>	<b>0.8</b>	<b>150.0</b>	<b>37.0</b>	<b>73.0</b>	<b>17.0</b>	<b>23.0</b>	

Bull Trout Spawning Surveys  
 For the Imnaha River, 2010  
 USFWS, La Grande Field Office

Imnaha Basin Stream Reach, Section	Date(s)	Kilometers Surveyed	Miles Surveyed	Occ	Redds		Per km	Per Mile	Total But obs	Bull Trout Observed (mm)			
				Unocc	Total				<6"(150mm)	<12" (~300mm)	<18"(450mm)	>18"(450mm)	
<b>Middle Imnaha</b>													
Blue Hole to Indian Crossing	30-Sep	3.2	2.0	0.0	1.0	1.0	0.3	0.5	12.0	0.0	0.0	1.0	11.0
Blue Hole to Indian Crossing	8-Oct			0.0	3.0	3.0	0.9	1.5	1.0	0.0	0.0	1.0	0.0
<b>Middle Imnaha Total</b>		<b>3.2</b>	<b>2.0</b>	<b>0.0</b>	<b>4.0</b>	<b>4.0</b>	<b>1.3</b>	<b>2.0</b>	<b>13.0</b>	<b>0.0</b>	<b>0.0</b>	<b>2.0</b>	<b>11.0</b>
<b>Big Sheep System</b>													
Big Sheep, Canal to Rd. 39	15-Sep	3.1	1.9	1.0	4.0	5.0	1.6	2.6	8.0	1.0	7.0	0.0	0.0
Big Sheep, Canal to Rd. 39	5-Oct			1.0	2.0	3.0	1.0	1.6	7.0	5.0	2.0	0.0	0.0
Lick Cr. Meadow to 39 rd.	5-Oct	2.4	1.5	0.0	7.0	7.0	2.9	4.7	1.0	1.0	0.0	0.0	0.0
Lick Cr. 39 Rd. to Quartz Cr.	15-Sep	6.8	4.2	0.0	4.0	4.0	0.6	1.0	5.0	2.0	3.0	0.0	0.0
Lick Cr. 39 Rd. to Quartz Cr.	5-Oct			0.0	1.0	1.0			10.0	10.0	0.0	0.0	0.0
<b>Big Sheep System Total</b>		<b>12.3</b>	<b>7.6</b>	<b>2.0</b>	<b>18.0</b>	<b>20.0</b>	<b>1.6</b>	<b>2.6</b>	<b>31.0</b>	<b>19.0</b>	<b>12.0</b>	<b>0.0</b>	<b>0.0</b>
<b>Imnaha Basin Total (Page 1)</b>													
		<b>28.0</b>	<b>17.4</b>	<b>28.0</b>	<b>101.0</b>	<b>129.0</b>	<b>4.6</b>	<b>7.4</b>	<b>106.0</b>	<b>18.0</b>	<b>61.0</b>	<b>15.0</b>	<b>12.0</b>
<b>Imnaha Basin Total (Page 2)</b>													
		<b>15.5</b>	<b>9.6</b>	<b>2.0</b>	<b>22.0</b>	<b>24.0</b>	<b>1.5</b>	<b>2.5</b>	<b>44.0</b>	<b>19.0</b>	<b>12.0</b>	<b>2.0</b>	<b>11.0</b>
<b>Imnaha Basin Total Pages 1 &amp; 2)</b>													
		<b>43.5</b>	<b>27.0</b>	<b>30.0</b>	<b>123.0</b>	<b>153.0</b>	<b>0.5</b>	<b>0.8</b>	<b>150.0</b>	<b>37.0</b>	<b>73.0</b>	<b>17.0</b>	<b>23.0</b>

Bull Trout Spawning Surveys  
For Some Grande Ronde Tributaries, 2010  
USFWS, La Grande Field Office

Page 1 of 1

Grande Ronde Basin Stream Reach, Section	Date(s)	Kilometers Surveyed	Miles Surveyed	Occ	Redds			Per km	Per Mile	Total But obs	Bull Trout Observed (mm)			
					Unocc	Total					<6"(150mm)	<12" (~300mm)	<18"(450mm)	>18"(450mm)
<b>Bear Creek</b>														
Goat Cr (Mouth to Falls)	16-Sep	1.4	0.9	4.0	2.0	6.0	4.1	6.7	20.0	1.0	9.0	5.0	5.0	
Goat Cr (Mouth to Falls)	13-Oct			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bear Creek (Standley Trail to USFS Cabin)	16-Sep	1.4	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bear Creek (Standley Trail to USFS Cabin)	13-Oct			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bear Creek (USFS Cabin to Goat Creek)	16-Sep	0.8	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bear Creek (USFS Cabin to Goat Creek)	13-Oct			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bear Creek (Goat Creek to Wilderness Boundary)	16-Sep	1.4	0.9	0.0	1.0	1.0	0.7	1.1	3.0	2.0	1.0	0.0	0.0	
Bear Creek (Goat Creek to Wilderness Boundary)	13-Oct			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Bear Creek Total</b>		<b>5.1</b>	<b>3.2</b>	<b>4.0</b>	<b>3.0</b>	<b>7.0</b>	<b>1.4</b>	<b>2.2</b>	<b>23.0</b>	<b>3.0</b>	<b>10.0</b>	<b>5.0</b>	<b>5.0</b>	
<b>Lostine River</b>														
Lundquist Bridge to OC Ranch	22-Sep	4.4	2.8	0.0	0.0	0.0	0.0	0.0	2.0	0.0	1.0	0.0	1.0	
Lundquist Bridge to OC Ranch	7-Oct			0.0	1.0	1.0	0.2	0.4	0.0	0.0	0.0	0.0	0.0	
Pole Bridge to 6 Mile Bridge	22-Sep	3.2	2.0	0.0	5.0	5.0	1.6	2.5	2.0	0.0	1.0	1.0	0.0	
Pole Bridge to 6 Mile Bridge	6-Oct			0.0	1.0	1.0	0.3	0.5	1.0	0.0	1.0	0.0	0.0	
Williamson to Walla Walla	22-Sep	3.5	2.2	0.0	2.0	2.0	0.6	0.9	0.0	0.0	0.0	0.0	0.0	
Williamson to Walla Walla	7-Oct			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bowman to French Camp	22-Sep	2.6	1.6	0.0	2.0	2.0	0.8	1.3	0.0	0.0	0.0	0.0	0.0	
Bowman to French Camp	6-Oct			1.0	3.0	4.0	1.5	2.5	10.0	8.0	2.0	0.0	0.0	
French Camp to Shady Falls	22-Sep	2.4	1.5	1.0	8.0	9.0	3.8	6.0	5.0	1.0	1.0	0.0	3.0	
French Camp to Shady Falls	6-Oct			0.0	12.0	12.0	5.0	8.0	1.0	0.0	1.0	0.0	0.0	
<b>Lostine River Total</b>		<b>16.2</b>	<b>10.1</b>	<b>2.0</b>	<b>34.0</b>	<b>36.0</b>	<b>2.2</b>	<b>3.6</b>	<b>21.0</b>	<b>9.0</b>	<b>7.0</b>	<b>1.0</b>	<b>4.0</b>	
<b>Deer Creek</b>														
Rd 8270 Road Crossing upstream 0.8 miles	17-Sep	1.3	0.8	1.0	10.0	11.0	8.5	13.8	38.0	28.0	10.0	0.0	0.0	
Rd 8270 Road Crossing upstream 0.8 miles	4-Oct			0.0	1.0	1.0	0.8	1.0	1.0	1.0	0.0	0.0	0.0	
<b>Deer Creek Total</b>		<b>1.3</b>	<b>0.8</b>	<b>1.0</b>	<b>11.0</b>	<b>12.0</b>	<b>9.3</b>	<b>15.0</b>	<b>39.0</b>	<b>29.0</b>	<b>10.0</b>	<b>0.0</b>	<b>0.0</b>	
<b>Grande Ronde Basin Total</b>														



**Table 4a –  
Summary of Measured Bull Trout Redds, Grande Ronde River Basin  
Lostine River and Goat Creek 2010**

Stream	n*1		Length (M)	Width (M)	Area (m2)	Length/Width ratio
Lostine	36	mean	1.1	0.7	0.9	1.8
		sd	0.5	0.3	0.7	0.7
		max	2.5	0.7	3	3.6
		min	0.3	0.2	0.1	0.6
Goat	6	mean	1.0	0.4	0.4	2.4
		sd	0.2	0.2	0.3	0.5
		max	1.1	1.0	1.1	2.4
		min	0.7	0.4	0.4	2.4

\*n = number of redds observed and measured (sample size).

Bear Creek had one redd documented in 2010; therefore is not included in table 4a.

**Table 4b – Summary of Measured Bull Trout Redds, Imnaha River Basin 2010**

Stream	n*1		Length (M)	Width (M)	Area (m2)	Length/Width ratio
Lick Creek	12	mean	.7	.5	.4	1.4
		sd	0.4	0.2	0.4	0.4
		max	1.3	0.9	1.1	2.1
		min	0.2	0.2	0.1	0.8
Big Sheep Creek	8	mean	0.7	0.3	0.2	1.9
		sd	0.2	0.1	0.1	0.3
		max	0.8	0.4	0.3	2.4
		min	0.4	0.2	0.1	1.3
Middle Imnaha	4	mean	1.5	0.7	1.1	2.1
		sd	0.7	0.2	0.8	0.7
		max	2.5	0.9	2.4	2.7
		min	1.1	0.4	0.4	1.5
Upper Imnaha	34	mean	1.7	0.8	1.4	2.2
		sd	0.6	0.2	0.8	0.8
		max	3.0	1.3	3.4	4.3
		min	0.7	0.3	0.3	1.1
N.F. Imnaha	20	mean	1.1	0.6	0.8	1.9
		sd	0.5	0.2	0.5	0.7
		max	2.4	1.2	1.8	4.8
		min	0.4	0.2	0.1	1.2
S.F. Imnaha	29	mean	1.1	0.5	0.7	2.3
		sd	0.6	0.2	0.6	1.0
		max	2.3	1.3	3.0	5.5
		min	0.3	0.2	0.1	0.8
Cliff Creek	41	mean	0.6	0.4	0.2	1.8
		sd	0.2	0.1	0.1	0.4
		max	1.0	0.7	0.6	3.0
		min	0.3	0.1	0.1	1.0

Footnote 1: n = number of redds observed and measured (sample size).