

Draft

Upper Snake Recovery Unit

Implementation Plan

for

Bull Trout Recovery Plan

June 2015

Prepared by

U.S. Fish and Wildlife Service

Idaho Fish and Wildlife Office

Boise, Idaho

Table of Contents

Introduction	E-1
Current Status of Bull Trout in the Upper Snake Recovery Unit.....	E-8
Factors Affecting Bull Trout in the Upper Snake Recovery Unit	E-10
Ongoing Upper Snake Recovery Unit Conservation Measures (Summary)	E-17
Research, Monitoring, and Evaluation	E-18
Recovery Measures Narrative	E-19
Implementation Schedule for the Upper Snake Recovery Unit.	E-56
References	E-79
Appendix I. Core Area Technical Meeting Summaries	E-81
Appendix II. Core Area Summaries	E-88

List of Figures

Figure E-1. Map of the Upper Snake Recovery Unit for bull trout.	E-3
---	-----

List of Tables

Table E-1. Bull trout Population Summary by Major Geographic Regions within Upper Snake Recovery Unit.....	E-4
Table E-2. Summary of Bull Trout Status by Core Area within Upper Snake Recovery Unit..	E-5
Table E-3. Primary Threats for the Upper Snake Recovery Unit (by Core Area)	E-13
Table E-4. Upper Snake Recovery Unit Implementation Schedule	E-56

Upper Snake Recovery Unit

Draft Recovery Unit Implementation Plan

Introduction

This draft recovery unit implementation plan (RUIP) describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the Upper Snake Recovery Unit, including estimates of time required and cost. This document supports and complements the Draft Revised Recovery Plan for the Coterminous U.S. Population of Bull Trout (USFWS 2014), which described recovery criteria and a general range-wide recovery strategy for the species, but deferred detailed discussion of species status and recovery actions within each of the six recovery units to RUIPs developed in coordination with State, Federal, Tribal, and other conservation partners. After we have received public comment on the draft RUIPs, we will incorporate changes as appropriate and release a final version in conjunction with the final recovery plan.

The Upper Snake Recovery Unit includes portions of central Idaho, northern Nevada, and eastern Oregon. Major drainages include the Salmon River, Malheur River, Jarbidge River, Little Lost River, Boise River, Payette River, and the Weiser River. The Upper Snake Recovery Unit contains 22 bull trout core areas (Figure E-1) within 7 geographic regions or major watersheds: Salmon River (10 core areas, 123 local populations), Boise River (2 core areas, 29 local populations), Payette River (5 core areas, 25 local populations), Little Lost River (10 local populations), Malheur River (2 core areas, 8 local populations), Jarbidge River (6 local populations), and Weiser River (5 local populations) (Table E-1). The Upper Snake Recovery Unit includes a total of 206 local populations, with almost 60 percent being present in the Salmon River watershed.

Three major bull trout life history expressions are present in the Upper Snake Recovery Unit. The Upper Salmon River, Deadwood River, Anderson Ranch, Arrowrock, Opal Lake, and Lake Creek core areas currently support adfluvial¹ populations of bull trout. All remaining core areas contain resident² populations and most have fluvial³ populations. Large areas of intact habitat exist primarily in the Salmon drainage, as this is the only drainage in the Upper Snake Recovery Unit that still flows directly into the Snake River; most other drainages no longer have direct connectivity due to irrigation uses or instream barriers. Bull trout in the Salmon basin

¹ Adfluvial: Life history pattern of spawning and rearing in tributary streams and migrating to lakes or reservoirs to mature.

² Resident: Life history pattern of residing in tributary streams for the fish's entire life without migrating.

³ Fluvial: Life history pattern of spawning and rearing in tributary streams and migrating to larger rivers to mature.


share a genetic past with bull trout elsewhere in the Upper Snake Recovery Unit. Historically, the Upper Snake Recovery Unit is believed to have largely supported the fluvial life history form; however, many core areas are now isolated or have become fragmented watersheds, resulting in replacement of the fluvial life history with resident or adfluvial forms. The Weiser River, Squaw Creek, and North Fork Payette River core areas contain only resident populations of bull trout.

Current Status of Bull Trout in the Upper Snake Recovery Unit

The breakdown of core areas by geographic region and the overall status of bull trout populations within the Upper Snake Recovery Unit are summarized in Tables E-1 and E-2. A description of bull trout status within each geographic region follows below; descriptions of current bull trout status and distribution for each individual core area are given in Appendix II below.

Salmon River

The Salmon River basin represents one of the few basins that are still free-flowing down to the Snake River. The core areas in the Salmon River basin do not have any major dams and a large extent (approximately 89 percent) is federally managed, with large portions of the Middle Fork Salmon River and Middle Fork Salmon River - Chamberlain core areas occurring within the Frank Church River of No Return Wilderness. Most core areas in the Salmon River basin contain large populations with many occupied stream segments. The Salmon River basin contains 10 of the 22 core areas in the Upper Snake Recovery Unit and contains the majority of the occupied habitat. Over 70 percent of occupied habitat in the Upper Snake Recovery Unit occurs in the Salmon River basin as well as 123 of the 206 local populations. Connectivity between core areas in the Salmon River basin is intact; therefore it is possible for fish in the mainstem Salmon to migrate to almost any Salmon River core area or even the Snake River. Connectivity within Salmon River basin core areas is mostly intact except for the Pahsimeroi River and portions of the Lemhi River. The Upper Salmon River, Lake Creek, Opal Lake core areas contain adfluvial populations of bull trout while the remaining core areas contain fluvial populations while only the Lemhi contains strictly resident populations. Most core areas appear to have increasing or stable trends but trends are not known in the Pahsimeroi, Lake Creek, or Opal Lake core areas. The Idaho Department of Fish and Game reported trend data from seven of the 10 core areas. This trend data indicated that populations were stable or increasing in the Upper Salmon River, Lemhi River, Middle Salmon River-Chamberlain, and the South Fork Salmon River (IDFG 2005, 2008, 2014). Trends were stable or decreasing in the Little-Lower Salmon River, Middle Fork Salmon River, and the Middle Salmon River-Panther (IDFG 2005, 2008, 2014).



E-3

Figure E-1. Map of the Upper Snake Recovery Unit for bull trout.

Table E-1. Bull trout population summary by major geographic regions within the Upper Snake Recovery Unit

Geographic Region	# Core Areas	# Local Populations	% Local Pops In Recovery Unit	Occupied Habitat in Recovery Unit	% Federally Managed (USFWS 2002a, 2004a)
Salmon River	10	123	59.7%	70%	89%
Boise River	2	29	14.1%	12%	62%
Payette River	5	25	12.1%	<9%	60%
Little Lost River	1	10	4.9%	<3%	89%
Malheur River	2	8	3.9%	<3%	60%
Jarbidge River	1	6	2.9%	3%	89%
Weiser River	1	5	2.4%	<2%	44%

Table E-2. Summary of bull trout status by core area within the Upper Snake Recovery Unit.

Core Area	Population Status (IDFG 2005)	Population Status (IDFG 2008)	Population Status (IDFG 2014)	Trends	# Local Pops	% Local Pops	Primary Threats Identified
Little-Lower Salmon River	Increasing	Increasing	Stable	Stable – Increasing/Decreasing	6	2.9%	No
South Fork Salmon River	Increasing	Increasing	Increasing	Increasing	27	13.1%	No
Middle Salmon River-Chamberlain	Decreasing	Stable	Increasing	Stable	9	4.4%	No
Middle Fork Salmon River	Increasing	Decreasing	Decreasing	Stable – Technical Partners	28	13.6%	No
Middle Salmon River-Panther	Decreasing	Decreasing		Stable – Technical Partners	18	8.7%	No
Lemhi River	Increasing	Increasing	Stable-Increasing	Stable- Increasing	6	2.9%	No
Pahsimeroi River				<i>Unknown</i>	9	4.4%	Yes

Core Area	Population Status (IDFG 2005)	Population Status (IDFG 2008)	Population Status (IDFG 2014)	Trends	# Local Pops	% Local Pops	Primary Threats Identified
Upper Salmon River	Increasing	Increasing	Increasing	Increasing	18	8.7%	No
Opal Lake				Stable – Technical Partners – Unknown	1	0.5%	No
Lake Creek				Stable – Technical Partners – Unknown	1	0.5%	No
Anderson Ranch			Increasing	Increasing	11	5.3%	No
Arrowrock				<i>Unknown</i>	18	8.7%	Yes
Squaw Creek				<i>Unknown</i>	4	1.9%	Yes
North Fork Payette River				<i>Unknown</i>	1	0.5%	Yes
Middle Fork Payette River				<i>Unknown</i>	3	1.5%	Yes
Deadwood River				<i>Unknown</i>	6	2.9%	Yes

Core Area	Population Status (IDFG 2005)	Population Status (IDFG 2008)	Population Status (IDFG 2014)	Trends	# Local Pops	% Local Pops	Primary Threats Identified
Upper South Fork Payette River				<i>Unknown</i>	11	5.3%	Yes
Little Lost River	Decreasing	Increasing	Stable- Decreasing	Stable – Technical Partners	10	4.9%	No
North Fork Malheur River				Decreasing– Technical Partners	5	2.4%	Yes
Malheur River (Upper)				Decreasing– Technical Partners	3	1.5%	Yes
Jarbidge River				Stable – Technical Partners - Unknown	6	2.9%	No
Weiser River			Increasing	Increasing	5	2.4%	No

Idaho Department of Fish and Game (IDFG) 2005 – Bull Trout Status Review and Assessment in the State of Idaho

IDFG 2008 – Distribution, Abundance, and Population Trend of Bull Trout in Idaho

IDFG 2014 – Bull Trout Trends in Abundance and Probabilities of Persistence in Idaho

Trends – based on IDFG reports and Technical Partner Input (Appendix I)

Boise River

In the Boise River basin, two large dams are impassable barriers to upstream fish movement: Anderson Ranch Dam on the South Fork Boise River, and Arrowrock Dam on the mainstem Boise River. Fish in Anderson Ranch Reservoir have access to the South Fork Boise River upstream of the dam. Fish in Arrowrock Reservoir have access to the North Fork Boise River, Middle Fork Boise River, and lower South Fork Boise River. The Boise River basin contains 2 of the 22 core areas in the Upper Snake Recovery Unit. The core areas in the Boise River basin account for roughly 12 percent of occupied habitat in the Upper Snake Recovery Unit and contain 29 of the 206 local populations. Approximately 90 percent of both Arrowrock and Anderson Ranch core areas are federally owned; most lands are managed by the Forest Service, with some portions occurring in designated wilderness areas. Both the Arrowrock core area and the Anderson Ranch core area are isolated from other core areas. Both core areas contain fluvial bull trout that exhibit adfluvial characteristics and numerous resident populations. The Idaho Department of Fish and Game in 2014 determined that the Anderson Ranch core area had an increasing trend while trends in the Arrowrock core area is unknown (IDFG 2014).

Payette River

The Payette River basin contains three major dams that are impassable barriers to fish: Deadwood Dam on the Deadwood River, Cascade Dam on the North Fork Payette River, and Black Canyon Reservoir on the Payette River. Only the Upper South Fork Payette River and the Middle Fork Payette River still have connectivity, the remaining core areas are isolated from each other due to dams. Both fluvial and adfluvial life history expression are still present in the Payette River basin but only resident populations are present in the Squaw Creek and North Fork Payette River core areas. The Payette River basin contains 5 of the 22 core areas and 25 of the 206 local populations in the recovery unit. Less than 9 percent of occupied habitat in the recovery unit is in this basin. Approximately 60 percent of the lands in the core areas are federally owned and the majority are managed by the Forest Service. Trend data are lacking and the current condition of the various core areas is unknown, but there is concern due to the current isolation of three (North Fork Payette River, Squaw Creek, Deadwood River) of the five core areas; the presence of only resident local populations in two (North Fork Payette River, Squaw Creek) of the five core areas; and the relatively low numbers present in the North Fork core area.

Jarbidge River

The Jarbidge River core area contains two major fish barriers along the Bruneau River: the Buckaroo diversion and C. J. Strike Reservoir. Bull trout are not known to migrate down to the Snake River. There is one core area in the basin, with populations in the Jarbidge River; this

watershed does not contain any barriers. Approximately 89 percent of the Jarbidge core area is federally owned. Most lands are managed by either the Forest Service or Bureau of Land Management. A large portion of the core area is within the Bruneau-Jarbidge Wilderness area. A tracking study has documented bull trout population connectivity among many of the local populations, in particular between West Fork Jarbidge River and Pine Creek. Movement between the East and West Fork Jarbidge River has also been documented; therefore both resident and fluvial populations are present. The core area contains six local populations and 3 percent of the occupied habitat in the recovery unit. Trend data are lacking within this core area.

Little Lost River

The Little Lost River basin is unique in that the watershed is within a naturally occurring hydrologic sink and has no connectivity with other drainages. A small fluvial population of bull trout may still exist, but it appears that most populations are predominantly resident populations. There is one core area in the Little Lost basin, and approximately 89 percent of it is federally owned by either the Forest Service or Bureau of Land Management. The core area contains 10 local populations and less than 3 percent of the occupied habitat in the recovery unit. The current trend condition of this core area is likely stable, with most bull trout residing in Upper Sawmill Canyon (IDFG 2014).

Malheur River

The Malheur River basin contains major dams that are impassable to fish. The largest are Warm Springs Dam, impounding Warm Springs Reservoir on the mainstem Malheur River, and Agency Valley Dam, impounding Beulah Reservoir on the North Fork Malheur. The dams result in two core areas that are isolated from each other and from other core areas. Local populations in the two core areas are limited to habitat in the upper watersheds. The Malheur River basin contains 2 of the 22 core areas and 8 of the 206 local populations in the recovery unit. Fluvial and resident populations are present in both core areas. This basin contains less than 3 percent of the occupied habitat in the recovery unit, and approximately 60 percent of lands in the two core areas are federally owned. Trend data indicates that populations are declining in both core areas.

Weiser River

The Weiser River basin contains local populations that are limited to habitat in the upper watersheds. The Weiser River basin contains only a single core area that consists of 5 of the 206 local populations in the recovery unit. Local populations occur in only three stream complexes in the upper watershed: 1) Upper Hornet Creek, 2) East Fork Weiser River, and 3) Upper Little Weiser River. These local populations include only resident life histories. This basin contains

less than 2 percent of the occupied habitat in the recovery unit, and approximately 44 percent of lands are federally owned. Trend data from the Idaho Department of Fish and Game indicate that the populations in the Weiser core area are increasing (IDFG 2014) but it is considered vulnerable because local populations are isolated and likely do not express migratory life histories.

Changes to Recovery Unit

Two notable changes to core area descriptions and boundaries within the Upper Snake Recovery Unit have occurred since the 2002 Draft Recovery Plan for Bull Trout. These changes include the removal of the Lucky Peak core area and the splitting of the Malheur core area. The Lucky Peak core area that was identified in 2002 has since been determined to be a population sink with limited reproduction. Genetic testing has determined that the individuals in Lucky Peak are identical to individuals from Arrowrock, and review of information indicates that the current population in the Lucky Peak core area is sustained artificially through entrainment (USFWS 2008). Thus, we have concluded it should no longer be identified as a core area. In addition, based on updated genetic information (DeHaan *et al.* 2007), two genetically distinct groups of bull trout exist within the Malheur River basin. Therefore, the former Malheur core area in Oregon was divided into two separate core areas, the Upper Malheur core area and the North Fork Malheur core area.

Factors Affecting Bull Trout in the Upper Snake Recovery Unit

Most threats to bull trout, as described in various documents including State plans (*e.g.*, Montana Bull Trout Restoration Team 2000; Batt 1996), the draft recovery plans (USFWS 2002a, 2002b, 2004b, 2004c), the critical habitat rules (USFWS 2002a, 2004a, 2010), the updated Bull Trout Core Area Templates (USFWS 2005b, 2008), the Bull Trout Core Area Conservation Status Assessment (USFWS 2005c), and the 2014/2015 Technical Partner Meetings (Appendix I), fall into the category of destruction, modification, or curtailment of habitat. Most of these impacts (*e.g.*, dewatering, sedimentation, thermal modification, and water quality degradation) are human-caused and are a consequence of specific land and water management activities.

For the purpose of this draft RUIP, we are identifying three broad threat categories: 1) Habitat Threats, 2) Demographic Threats, and 3) Nonnative Fish Threats. Habitat Threats are those that impact bull trout habitat (habitat fragmentation and degradation resulting from upland/riparian land management and instream impacts), Demographic Threats are those that impact individuals or populations (connectivity impairment and small population size), while

Nonnative Fish Threats result from effects of introduced fish species or their management that impact individuals or populations (competition, predation, and hybridization).

Habitat Threats and Demographic Threats are likely the major limiting factors for bull trout in the Upper Snake Recovery Unit. These factors affect individuals and local populations as well as habitat for the species. Although in some basins reservoirs formed by dams have allowed bull trout to express adfluvial life histories, dams, irrigation diversions, and road crossings have also formed impassable barriers to fish movement within the basins, further fragmenting habitats and isolating bull trout. Land management activities that degrade aquatic and riparian habitats by altering stream flows and riparian vegetation, such as water diversions, past and current mining operations, timber harvest and road construction, and improper grazing practices, have negatively affected bull trout in several areas of the recovery unit.

Bull trout are also subject to negative interactions with nonnative brook trout in some streams. Brook trout populations are prevalent throughout the Upper Snake Recovery Unit; this species has been identified as a significant threat to bull trout in some core areas. In some local populations and core areas, bull trout abundance appears to be related to brook trout competition and hybridization. Low abundance of bull trout appears to be related to high road density, sedimentation, passage barriers, and brook trout.

Primary Threats

In the revised draft recovery plan (USFWS 2014) we updated the known threats identified in the previous draft recovery plans (USFWS 2002a, 2002b, 2004b, 2004c), with specific focus on threats at the individual core area level, where threats operatively impact bull trout local populations and limit their recovery potential. A threat was considered a primary threat if the threat affected the persistence of a local population and eventually the persistence of a core area. In December 2014, the Service also invited technical partners to several meetings to discuss the threats to bull trout in each of the 22 core areas in the Upper Snake Recovery Unit, with the intent of identifying primary threats (Appendix I).

We evaluated whether a threat should be considered a primary threat by considering information from technical partners, the current status and distribution of populations, known trend information, and existing conservation measures. We weighted information from technical partners heavily in our evaluations. Core areas that all partners determined had no primary threats were identified as having no primary threats. In certain core areas technical partners determined that there were no primary threats based on best professional assessment of core area condition and environment (*e.g.*, remote wilderness regions with few known environmental impacts) although trend data may have been negative (see Table E-2, Middle Fork Salmon River

and Middle Salmon River - Panther) or not available (see Table E-2, Lake Creek, Opal Lake, and Jarbidge). For all other core areas the Service has identified potential primary threats that are present as identified by technical partners. A draft list of potential primary threats is provided in Table E-3 below. For this draft Recovery Unit Implementation Plan we have attempted to be inclusive, and we invite the public and partners to comment on this list of primary threats. A summary of meeting discussions is provided in Appendix I and Appendix II.

Climate Change

Climate change may exacerbate already identified threats to bull trout habitat such as warming water temperatures, but we are unaware of unique or different threats posed in the near-term. Our strategy for addressing climate change is to reduce or remove these already identified threats and to collaborate with partners to develop a range-wide climate vulnerability assessment to ensure we manage climate change impacts to bull trout with the greatest certainty. The identification of core areas and watersheds that are most likely to maintain habitats suitable for bull trout over the foreseeable future under probable climate change scenarios will also help guide the allocation of bull trout conservation resources to improve the likelihood of success.

Table E-3. Primary Threats for the Upper Snake Recovery Unit (by Core Area)

<u>Geographic Region</u> Core Area – Complex <i>Core Area - Simple</i>	Number of Local Populations	PRIMARY THREATS ¹		
		<i>Habitat</i>	<i>Demographic</i>	<i>Nonnatives</i>
<u>Salmon River Geographic Region</u>				
Little-Lower Salmon River	6	None	None	None
South Fork Salmon River	27	None	None	None
Middle Salmon River-Chamberlain	9	None	None	None
Middle Fork Salmon River	28	None	None	None
Middle Salmon River-Panther	19	None	None	None
Lemhi River	6	None	None	None
Pahsimeroi River	9	Instream Impacts (1.2) Dewatering, Altered Flow	Connectivity Impairment (2.1) Fish Passage Issues	
Upper Salmon River	18	None	None	None
<i>Opal Lake</i>	1	None	None	None
<i>Lake Creek</i>	1	None	None	None

<u>Geographic Region</u> Core Area – Complex Core Area - Simple	Number of Local Populations	PRIMARY THREATS¹		
		<i>Habitat</i>	<i>Demographic</i>	<i>Nonnatives</i>
<u>Boise River Geographic Region</u>				
Anderson Ranch	11	None	None	None
Arrowrock	18	Instream Impacts (1.2) Altered flows (water management)	Connectivity Impairment (2.1) Fish Passage Issues Forage Fish Availability (2.4) Water Management	Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)
<u>Payette River Geographic Region</u>				
Squaw Creek	4	Upland/Riparian Land Management (1.1) Livestock Grazing	Connectivity Impairment (2.1) Fish Passage Issues	Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)
North Fork Payette River	1		Connectivity Impairment (2.1) Fish Passage Issues Small Population Size (2.3) Genetic, Demographic Stochasticity	Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)
Middle Fork Payette River	3			Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)
Deadwood River	6		Connectivity Impairment (2.1) Fish Passage Issues	Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)

<u>Geographic Region</u> Core Area – Complex Core Area - Simple	Number of Local Populations	PRIMARY THREATS ¹		
		<i>Habitat</i>	<i>Demographic</i>	<i>Nonnatives</i>
			Water Management	
Upper South Fork Payette River	11		Connectivity Impairment (2.1) Fish Passage Issues	Nonnative fishes (3.1) Predation/Species Competition, Hybridization (brook trout)
<u>Little Lost River Geographic Region</u>				
Little Lost River	10	None	None	None
<u>Malheur River Geographic Region</u>				
North Fork Malheur River	5	Upland/Riparian Land Management (1.1) Forest Management Practices, Livestock Grazing Instream Impacts (1.2) Water Management <u>Water Quality:</u> Forest Management Practices, Livestock Grazing	Connectivity Impairment (2.1) Entrainment, Dewatering, Temperature Barriers	Nonnative fishes (3.1) Potential for Invasion

<u>Geographic Region</u> Core Area – Complex Core Area - Simple	Number of Local Populations	PRIMARY THREATS¹		
		<i>Habitat</i>	<i>Demographic</i>	<i>Nonnatives</i>
Upper Malheur River	3	Upland/Riparian Land Management (1.1) Forest Management Practices (legacy and current), Livestock Grazing <u>Water Quality:</u> Forest Management Practices (legacy and current), Livestock Grazing	Connectivity Impairment (2.1) Entrainment, Fish Passage Issues, Dewatering, Temperature Barriers Small Population Size (2.3) Genetic, Demographic Stochasticity	Nonnative fishes (3.1) Competition, Hybridization
<u>Jarbidge River Geographic Region</u>				
Jarbidge River	6	None	None	None
<u>Weiser River Geographic Region</u>				
Weiser River	5	None	None	None

¹ **Primary Threat:** Factors known or likely (i.e., non-speculative) to negatively impact bull trout populations at the core area level, and accordingly require management actions to assure bull trout persistence to a degree necessary that bull trout will not be at risk of extirpation within that core area in the foreseeable future (50 years).

Ongoing Upper Snake Recovery Unit Conservation Measures (Summary)

Since the listing of bull trout, numerous conservation measures have been and continue to be implemented within the Upper Snake Recovery Unit. These measures are being undertaken by a wide variety of local and regional partnerships, including State fish and game agencies, State and Federal land management and water resource agencies, Tribal governments, power companies, watershed working groups, water users, ranchers, and landowners. In many cases, these bull trout conservation measures incorporate or are closely interrelated with work being done for recovery of salmon and steelhead, which are limited by many of the same threats.

Many restoration projects have been implemented from local funds as well as Bonneville Power Administration funds in predominantly anadromous drainages. Bonneville Power Administration has also funded projects in the Malheur River to support Tribal efforts in recovering bull trout. The Bureau of Reclamation has been implementing various projects within the Malheur and Boise watersheds to better understand the impacts of their operations on bull trout populations.

The numerous localized fish habitat restoration projects in the Salmon River basin that are implemented by Federal, State, and private partners (U.S. Forest Service, Bureau of Land Management, Upper Salmon Basin Watershed Program) should continue and be expanded if possible, to protect and maintain the currently robust population. The Upper Salmon Basin Watershed Program has implemented over 500 projects since 1993 to increase instream flow and improve fish habitat across the Salmon River headwaters, Lemhi River, and Pahsimeroi River watersheds. The program, coordinated by the Idaho Governor's Office of Species Conservation, works with cooperating private landowners to develop restoration projects and obtain funding from Bonneville Power Administration and other agencies. Projects have included removal of migration barriers to provide fish access to 75 miles (121 km) of stream, screening of 249 irrigation diversions, instream habitat improvement in 494 miles (795 km) of stream, and riparian habitat restoration over 352 miles (566 km) of stream (158 miles [254 km] fenced). Projects have benefited bull trout, salmon, and other salmonid species.

The Forest Service and Bureau of Land Management have updated their Land and Resource Management Plans and Resource Management Plans to incorporate conservation measures that protect both local populations and habitat used by bull trout. Numerous passage projects have also increased the amount of habitat as well as improved connectivity throughout the recovery unit. Both these Federal agencies have areas within the Upper Snake with special designations such as Wild and Scenic River (Jarbridge River) or Wilderness Designation (Frank

Church Wilderness of No Return in the Salmon River and the Jarbidge Wilderness in the Jarbidge River). Both of these designations afford protection for bull trout and its habitat.

In southwestern Idaho, the National Forests, Idaho Department of Fish and Game, and cooperating private landowners should continue to implement upland and stream habitat restoration actions. Fish passage barriers within the following core areas (*e.g.*, Arrowrock, Squaw Creek, North Fork Payette, and Deadwood core areas) should be evaluated and addressed to improve bull trout population connectivity. The Idaho Department of Lands also implement conservation measures to ensure that fish passage is provided on vegetative management projects that warrant it.

Research, Monitoring, and Evaluation

The Upper Snake Recovery Unit currently lacks trend data in most core areas and there is a need to collect more information to determine whether populations are stable or increasing. The Idaho Department of Fish and Game and Oregon Department of Fish and Wildlife have collected trend data in only 12 of the 22 core areas in the recovery unit. Only 8 of the 12 indicated a positive trend (Table E-2) and decreasing trends were observed in some core areas that we have otherwise identified as likely to be stable without primary threats (Middle Fork Salmon and Middle Salmon-Panther). While many parts of the range are stable or increasing, other areas do not have any information regarding trends. Bull trout trends are unknown within the entire Payette River geographic region, while the Salmon River geographic region has a robust amount of information.

Recovery Measures Narrative

Below is a list of actions that are needed to ensure that bull trout are recovered within the Upper Snake Recovery Unit.

Salmon River Geographic Region

Little-Lower Salmon River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery. Monitor John Day Creek and Slate Creek.
- Manage for demographic stochasticity by ensuring local populations contain more than 50 to 100 reproductive individuals and manage for environmental stochasticity with populations containing 1000 to 10,000 individuals, when practicable. Focus on additional survey efforts in smaller watersheds such as John Day Creek, Slate Creek, Lake-Lower Salmon, and Partridge Creek.
- Implement brook trout removal efforts wherever feasible and biologically supported. Consider monitoring populations in John Day Creek and Slate Creek.

- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

South Fork Salmon River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnatives

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery (South Fork Salmon River, Upper East Fork South Fork Salmon River, Lake Creek to Loon Lake, Sugar, Krassel-Indian, Curtis, Johnson [headwaters to mouth], and Cow-Oompaul creeks).
- Clean up mine waste at active, inactive, and orphan sites (Cinnibar and Stibnite Mine) (Meadow Creek and Blowout Creek).
- Implement brook trout removal efforts wherever feasible and biologically supported.
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Middle Salmon River - Chamberlain Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnatives

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery (Warren, Upper Hourse, Wind, Big Mallard, Witsher, Upper Meadow, and Upper Crooked creeks).
- Clean up mine waste at active, inactive, and orphan sites (Warren, Falls, Lake, and Upper Crooked creeks).
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Middle Fork Salmon River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery (Headwaters to Salmon River, Headwaters to Fall Creek, Bear Valley, Elk Creek, and Lower Camas Creek).
- Implement brook trout removal efforts wherever feasible and biologically supported.
- Identify barriers for bull trout and implement tasks to provide passage.
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Middle Salmon River - Panther Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery (Big Deer, Hughes, McKim, Musgrove, Moose, Hull, Hughes, Lick, Upper Horse, Squaw, Pine, Opal, Porphyry, Dahlenega Creeks).
- Clean up mine waste at active, inactive, and orphan sites (Blackbird Mine and Bear Track Mine).
- Identify barriers for bull trout and implement tasks to provide passage.
- Implement brook trout removal efforts wherever feasible and biologically supported.
- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas (North Fork River, Red, Twelve/Lake watershed, Hat Creek, Deep-Moyer watershed, Napias watershed).
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Lemhi River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Identify barriers for bull trout and implement tasks to provide passage.
- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas (Hayden watershed, Little Eightmile, Canyon, Reservoir, Upper Texas, and Little Timber creeks).
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Pahsimeroi River Core Area

1. Actions to Address Habitat Threats

1.1 Upland/Riparian Land Management

1.2 Instream Impacts

1.2.1 Restore streams that are partially or completely dewatered. Improve instream flows.

1.3 Water Quality

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.2 Fisheries Management

2.3 Small Population Size

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.

Upper Salmon River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Implement brook trout removal efforts wherever feasible and biologically supported.
- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas (East Fork Salmon, Morgan, Squaw, Challis, Grandiw, Slate, Big Lake, Boulder, and Valley Creeks).
- Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River Geographic Region.

Opal Lake Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Protect, restore, and maintain suitable habitat conditions for bull trout. Maintain or improve water quality in bull trout core areas.

Lake Creek Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Protect, restore, and maintain suitable habitat conditions for bull trout. Maintain or improve water quality in bull trout core areas.

Boise River Geographic Region

Anderson Ranch Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery.
- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.
- Identify barriers for bull trout and implement tasks to provide passage.
- Implement brook trout removal efforts wherever feasible and biologically supported.

- Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.

Arrowrock Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.1.2 Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.

2.2 Fisheries Management

2.3 Small Population Size

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

Payette River Geographic Region

Squaw Creek Core Area

1. Actions to Address Habitat Threats

1.1 Upland/Riparian Land Management

- 1.1.1 Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.

1.2 Instream Impacts

1.3 Water Quality

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

- 2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.2 Fisheries Management

2.3 Small Population Size

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

- 3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

North Fork Payette Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.2 Fisheries Management

2.3 Small Population Size

2.3.1 Manage for demographic stochasticity by ensuring local populations contain more than 50 to 100 reproductive individuals and manage for environmental stochasticity with populations containing 1000 to 10,000 individuals, when practicable.

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

Middle Fork Payette Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None.

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

- 3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

- 4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

Deadwood River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.1.2 Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.

2.2 Fisheries Management

2.3 Small Population Size

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

Upper South Fork Payette River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

2.1 Connectivity Impairment

2.1.1 Identify barriers for bull trout and implement tasks to provide passage.

2.2 Fisheries Management

2.3 Small Population Size

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

3.1.1 Implement brook trout removal efforts wherever feasible and biologically supported.

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.
Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

None

Little Lost River Geographic Region

Little Lost River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Identify barriers for bull trout and implement tasks to provide passage.
- Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.

Malheur River Geographic Region

North Fork Malheur Core Area

1. Actions to Address Habitat Threats

1.1. Upland/Riparian Land Management

Forest Management Practices

- #### 1.1.1 Restore canopy and riparian cover and native vegetation in all bull trout spawning, rearing, and migration areas. Horse Creek, Swamp Creek, Sheep Creek, Flat Creek, Elk Creek, Little Crane Creek, Crane Creek, and the North Fork Malheur River have suppressed woody vegetation and loss of effective shade. Emphasis should also be put on the Little Malheur River as bull trout have been recently captured there and it has the potential to provide spawning and rearing habitat. This component is vital to restoring not only shade but also natural instream processes, hydrologic function, and thermal regimes.

- 1.1.2 Implement stream restoration projects in degraded stream reaches. Review habitat information to identify and prioritize opportunities for stream restoration in Horse Creek, Swamp Creek, Sheep Creek, Flat Creek, Elk Creek, Little Crane Creek, Crane Creek, the North Fork Malheur River, and Little Malheur River. Design and implement projects based on findings.
- 1.1.3 Provide a reliable source of large hardwood beaver forage. Beaver have disappeared from much of their historical range. Beavers initiate and maintain critical watershed processes important to water retention, sediment sequestration, cold water storage, and flood plain connectivity. The re-establishment of these processes in the riverscape is critical to the recovery of bull trout and their habitat. The current lack of hardwoods in riparian habitats and the necessary structure to support beaver dam construction are one of the factors limiting the recolonization of the Upper Malheur River by beaver. Grazing pressure on riparian communities is detrimental to re-establishing these critical riparian hardwoods. Implement activities to encourage riparian shrub and hardwood communities to re-establish in an effort to encourage beaver to naturally recolonize and restore the riverscape. Consider providing large support material to jump start beaver dam construction.
- 1.1.4 Evaluate and implement actions to encourage beaver recolonization. To assist in re-establishing functional riparian communities, Federal, Tribal and State resource managers should identify and implement measures to increase beaver abundance where feasible and biologically supportable. Reduction in beaver trapping pressures, increases in active releases, and utilization of beaver control structures should be considered where effective and appropriate.

Livestock Grazing

- 1.1.5 Reduce grazing impacts. While recognizing that no livestock grazing would likely achieve recovery of habitat and populations more rapidly, the following measures would allow for livestock grazing occurring while habitat and populations recover at less than a near-natural rate of recovery. Livestock grazing within riparian areas proximate to bull trout critical habitat should be limited to light utilization and minimal bank disturbance. Based on current and best available science, threshold indicators should be monitored utilizing the Multiple Indicator Monitoring Method (Burton *et al.* 2011). Threshold indicators when measured for early to mid-season should not exceed:

Indicator	Foraging / Migration / Overwintering	Spawning / Rearing	Comments
Bank Alteration	Less than 20%	Less than 15%	<ul style="list-style-type: none"> • Monitor within a week of the cows coming off the pasture. • Burton <i>et al.</i> 2011 • Bengeyfield 2006
Stubble Height	6" (Early season) 8" (Late season)	8" (Early season) 10" (Late season)	<ul style="list-style-type: none"> • Goss 2013 (MS Thesis) • Clary and Webster (1989)
Browse	Light (21 to 40%)	Slight (0 to 20%)	<ul style="list-style-type: none"> • Burton <i>et al.</i> 2011

To further aid in the recovery of bull trout and minimize the potential for redd trampling, no livestock grazing should occur within sections of streams that are designated as Spawning/Rearing (USFWS 2010) after August 15 to the following spring. Removing livestock use after August 15 should also aid in the recovery of woody shrubs which provide shade and stability to stream channels. These streams include: Horse Creek, Swamp Creek, Sheep Creek, Flat Creek, Elk Creek, Little Crane Creek, Crane Creek, and the North Fork Malheur River. Little Malheur River is severely degraded and currently considered unoccupied during the peak of the summer due to stream temperatures. Special emphasis should be placed on restoring this tributary to support bull trout.

In conjunction with the above; other measures can be used to minimize grazing impacts which include fencing, changes in timing, rest, rest rotation, off-site watering and salting. Federal land management agencies should implement PACFISH/INFISH standards and guidelines for livestock grazing, as appropriate. Priority sites within the North Fork Malheur River include the following Federal allotments: Spring Creek allotment, North Fork allotment, Flag Prairie allotment, and Ott allotment, all of which have some stream temperature, riparian habitat, and channel complexity problems.

- 1.1.6 Curtail unauthorized livestock use on U.S. Forest Service property. Implement regulations designed to reduce and eliminate violations of grazing permits and unauthorized grazing. Any cattle, sheep, goat, hog, or equine not considered wild and free roaming that is grazing without a permit is considered unauthorized by 36 CFR 222.20(b)(13).

1.2. Instream Impacts

Water Management

- 1.2.1 Maintain a conservation pool in Beulah Reservoir to provide adequate overwinter rearing habitat for adult and sub-adult migratory bull trout. Reduction in reservoir levels, sometimes to run-of-the-river, can negatively affect temperature and prey fish availability, greatly influencing growth and survival of adfluvial bull trout in the reservoir (Petersen and Kofoot 2002). Implement Terms and Conditions from the 2005 Service Biological Opinion to maintain a conservation pool in Beulah Reservoir minimizing the frequency and extent of a draw down during years in which snow pack and/or stream flow is insufficient to exceed irrigation needs.

1.3. Water Quality

Forest Management Practices and Livestock Grazing

- 1.3.1 Maintain or improve effective shade to achieve water quality objectives as outlined in the TMDL (Total Maximum Daily Load) recommending no reduction of effective shade. Management activities should allow for recovery of effective shade based on site potential vegetation. The Oregon Department of Environmental Quality completed the Malheur Basin TMDL and Water Quality Management Plan in September 2010. The Malheur National Forest is slated to complete a water quality recovery plan in 2015. Follow recommendations and measures presented in these plans. Implement action 1.1.1 to help improve effective shade.

Agriculture Practices

- 1.3.2 Cool irrigation returns and run-off. Diversions or runoff warmer than the receiving water should be cooled when possible before allowing it to enter the receiving system (e.g., subterranean pipes).

2. Actions to Address Demographic Threats

2.1. Connectivity Impairment

Entrainment & Fish Passage Issues

- 2.1.1 Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers. Complete an inventory of unscreened irrigation diversions within the basin. Known high priorities for screening include diversions on the North Fork Malheur River.

Diversions on Forest Service property have either been screened or closed; if re-opened install appropriate screens to prevent entrainment of bull trout.

- 2.1.2 Reduce occurrence of spill at Agency Valley Dam to minimize entrainment of bull trout below the dam and provide upstream passage when entrainment occurs. Bull trout are entrained over the spillway of Agency Valley Dam when spill occurs. Once entrained, there is no existing facility for fish to return upstream to the reservoir. Due to elevated summer stream temperatures and low flows caused by irrigation withdrawal habitat conditions downstream of the dam are not suitable for bull trout survival. Implement the Terms and Conditions of the 2005 Service Biological Opinion to continue all existing efforts to limit the use of the spillway, minimize duration and quantity of spill, and trap and return bull trout that are entrained when the spillway is used back to Beulah Reservoir or the North Fork Malheur River above the dam.
- 2.1.3 Identify and remove barriers to juvenile and adult passage. Log weirs, culverts, legacy structures and other aquatic organism passage barriers impede juvenile and adult passage and prevent movement between spawning, rearing and overwinter habitats. Culverts for the National Forest road NF-13 where it crosses Swamp and Sheep creeks are a priority.

Dewatering

- 2.1.4 Improve and secure instream flows. Restore connectivity and opportunities for migration by securing instream flows and/or water rights. Improve irrigation efficiency. In addition, implement stream restoration actions identified under Recovery Action 1. Benefits of stream restoration will include raising the water table and restoring natural instream flow, providing more water during summer and late fall.

Temperature Barriers

- 2.1.5 Eliminate thermal barriers by maintaining or improving riparian vegetation communities, providing shade to streams, and increasing instream flow. Current bull trout distribution and movement is impeded by thermal barriers between spawning and rearing habitats. During the summer months, water temperature in the North Fork Malheur River between Beulah Reservoir and Crane Creek is considered a barrier to movement, and additional thermal barriers upstream of Crane Creek may occur in some years. Implement actions designed to cool warm water

temperatures, increase flows, and improve hydrologic function as detailed under Recovery Action 1.

2.2. Fisheries Management

2.3. Small Population Size

2.4. Forage Fish Availability

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fishes

Potential for Invasion

- 3.1.1 Survey and monitor the North Fork watershed for the presence of brook trout. Currently, brook trout are absent in the North Fork Malheur River. The illegal introduction and subsequent invasion of brook trout into the North Fork Malheur River would negatively impact bull trout populations through competition and hybridization and could quickly become one of the most significant threats to bull trout in the basin. Periodically and regularly survey the North Fork Malheur River watershed, within and outside the distribution of bull trout, for the presence of brook trout. Consider using e-DNA methodology as a low cost means to monitor the basin for brook trout occupancy.
- 3.1.2 Prioritize the removal of brook trout in adjacent basins. The presence of brook trout in high densities in the Upper Malheur River basin poses a direct threat to the bull trout in the North Fork Malheur core area. The risk of illegal inter-basin transfer is potentially very high. The removal or control of brook trout in the Upper Malheur River basin will decrease this risk considerably.
- 3.1.3 Implement actions that will ensure the expression of a migratory life history. Impacts of brook trout to bull trout populations appear to be most significant for populations of primarily resident fish. Bull trout populations containing large migratory individuals manage to maintain despite the presence of brook trout. In addition, large fish are more fecund, have great productivity, and can out-compete smaller brook trout for food and space resources. Actions specific to fostering a migratory life history include 1.2.1 and 2.1.2 to ensure suitable overwintering habitat in Beulah Reservoir, and those under 2.1 to maintain passable migratory corridors.

- 3.1.4 Implement brook trout removal efforts when brook trout are detected in the North Fork Malheur core area. Immediately conduct brook trout eradication efforts, when and if detected, in the North Fork Malheur Basin to prevent dispersal and colonization of the species.
- 3.1.5 Develop and implement an educational effort to address problems and consequences of unauthorized fish introductions. Target areas where inter-basin transfer of brook trout from adjacent systems would be most likely.

4. Monitoring and Evaluation

4.1 Habitat

Monitoring

- 4.1.1 Continue to monitor temperature, water quality, water quantity, and riparian condition to evaluate the effectiveness of habitat restoration actions and to provide the ability to detect trends in these metrics.

4.2 Demographic

Research

- 4.2.1 Further define bull trout distribution and habitat use in the core area. Research is needed to determine the extent to which bull trout express a fluvial, and potentially adfluvial, life history in the North Fork Malheur River.

Monitoring

- 4.2.2 Continue maintenance and operation of fish screens on all diversions. Constant monitoring and maintenance is necessary to keep fish screens operating properly.
- 4.2.3 Continue ongoing population monitoring efforts within the basin. Maintain current long term redd count datasets assessing abundance and distribution of spawning migratory bull trout. If necessary, bolster the monitoring program with new protocols and methodologies consistent with other programs statewide. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

4.3.1 Implement bass and crappie monitoring efforts in Beulah Reservoir.

Current data demonstrates multiple year classes of both nonnative species. Implementation of a conservation pool will likely allow both species to expand. Predation by bass could occur on bull trout and competition for prey would occur by both species.

Conservation Recommendations

- Promote interagency collaboration and coordination on bull trout recovery actions by supporting existing bull trout working groups or the formation of new bull trout working groups where they do not exist. While working groups may be facilitated by any interested stakeholder, most often they are organized and facilitated by the Service, a State agency, U.S. Forest Service, or a Tribal entity. Although the Service has no guidelines for format or process, existing working groups are largely informal, are organized at various scales (*e.g.*, core area, river basin, geographic region, or recovery unit) and generally meet at least annually.
- Identify and address sediment sources in North Fork Malheur Basin affecting bull trout. Identify road-related sediment problem areas in the North Fork Malheur River core area prioritizing spawning and rearing streams. Examine the ways roads capture and channel runoff, and changes in surface runoff associated with soil compaction. Stabilize roads, crossings, railroad grades, and other sources of sediment delivery; remove and vegetatively restore unneeded roads and railroad grades.
- Increase information outreach to anglers. Provide information on bull trout identification, special regulations, methods to reduce hooking mortality of bull trout caught incidentally, and the value of bull trout and their habitat.
- Investigate and implement actions to restore historic prey base by reintroducing anadromous species. Anadromous species such as steelhead and spring Chinook salmon were historically present in the North Fork Malheur River. Feasibility of restoration of spawning populations of these species to increase prey base and provide marine derived nutrients should be evaluated and implemented where feasible and biologically supportable.

Upper Malheur Core Area

1. Actions to Address Habitat Threats

1.1. Upland/Riparian Land Management

Forest Management Practices

- 1.1.1 Restore canopy and riparian cover, and native vegetation in all bull trout spawning, rearing, and migration areas. Crooked Creek, McCoy Creek, Lake Creek, Coral Basin Creek, Bosonberg Creek, Big Creek, Summit Creek, and the Malheur River downstream of Logan Valley have suppressed woody vegetation and loss of effective shade. This component is vital to restoring not only shade but also natural instream processes, hydrologic function, and thermal regimes.
- 1.1.2 Implement stream restoration projects in degraded stream reaches. Review habitat information to identify and prioritize opportunities for stream restoration, including increasing pool and gravel area, in Crooked Creek, McCoy Creek, Lake Creek, Coral Basin Creek, Bosonberg Creek, Big Creek, Summit Creek, and the Malheur River downstream of Logan Valley. Design and implement projects based on findings.
- 1.1.3 Provide a reliable source of large hardwood beaver forage. Beaver have disappeared from much of their historical range. Beavers initiate and maintain critical watershed processes important to water retention, sediment sequestration, cold water storage, and flood plain connectivity. The re-establishment of these processes in the riverscape is critical to the recovery of bull trout and their habitat. The current lack of hardwoods in riparian habitats and the necessary structure to support beaver dam construction are one of the factors limiting the recolonization of the Upper Malheur River by beaver. Grazing pressure on riparian communities is detrimental to re-establishing these critical riparian hardwoods. Implement activities to encourage riparian shrub and hardwood communities to re-establish in an effort to encourage beaver to naturally recolonize and restore the riverscape. Consider providing large support material to jump start beaver dam construction.
- 1.1.4 Evaluate and implement actions to encourage beaver recolonization. To assist in re-establishing functional riparian communities, Federal, Tribal and State resource managers should identify and implement measures to increase beaver abundance where feasible and biologically supportable. Reduction in beaver trapping pressures, increases in active releases, and

utilization of beaver control structures should be considered where effective and appropriate.

Livestock Grazing

1.1.5 Reduce grazing impacts. While recognizing no livestock grazing would likely achieve recovery of habitat and populations more rapidly, the following measures would allow livestock grazing to occur while habitat and populations recover at less than a near-natural rate of recovery. Livestock grazing within riparian areas proximate to bull trout critical habitat should be limited to light utilization and minimal bank disturbance. Based on current and best available science, threshold indicators should be monitored utilizing the Multiple Indicator Monitoring Method (Burton *et al.* 2011). Threshold indicators, when measured in early to mid-season, should not exceed:

Indicator	Foraging / Migration / Overwintering	Spawning / Rearing	Comments
Bank Alteration	Less than 20%	Less than 15%	<ul style="list-style-type: none"> • Monitor within a week of the cows coming off the pasture. • Burton <i>et al.</i> 2011 • Bengeyfield 2006
Stubble Height	6" (Early season) 8" (Late season)	8" (Early season) 10" (Late season)	<ul style="list-style-type: none"> • Goss 2013 (MS Thesis) • Clary and Webster (1989)
Browse	Light (21 to 40%)	Slight (0 to 20%)	<ul style="list-style-type: none"> • Burton <i>et al.</i> 2011

To further aid in the recovery of bull trout and minimize the potential for redd trampling, no livestock grazing should occur within sections of streams that are designated as spawning/rearing (USFWS 2010) after August 15 to the following spring. By removing livestock use after August 15 this should also aid in the recovery of woody shrubs which provide shade and stability to stream channels. These streams include: Meadow Fork of Big Creek, Big Creek, Snowshoe Creek, Lake Creek, McCoy Creek, Crooked Creek, Bosonberg Creek, and Summit Creek. McCoy Creek, Crooked Creek, Bosonberg Creek, and Summit Creek are severely degraded and are currently unoccupied during the peak of the

summer due to stream temperatures. Special emphasis should be placed on restoring these tributaries to support bull trout.

In conjunction with the above, further minimize grazing impacts with fencing, changes in timing, rest, rest rotation, off site watering, and salting. Federal land management agencies should implement PACFISH/INFISH standards and guidelines for livestock grazing, as appropriate. Priority sites within the North Fork Malheur River include the following Federal allotments: McCoy Creek allotment, Lake Creek allotment, Logan Valley allotment, Dollar Basin allotment, Star Glade Allotment, and Summit Prairie Allotment, all of which have some stream temperature, riparian habitat, and channel complexity problems.

- 1.1.6 Curtail unauthorized livestock use on U.S. Forest Service property. Implement regulations designed to reduce and eliminate violations of grazing permits and unauthorized grazing. Any cattle, sheep, goat, hog, or equine not considered wild and free roaming that is grazing without a permit is considered unauthorized by 36 CFR 222.20(b)(13).

1.2. Instream Impacts

1.3. Water Quality

Forest Management Practices and Livestock Grazing

- 1.3.1 Maintain or improve effective shade to achieve water quality objectives as outlined in the TMDL recommending no reduction of effective shade. Management activities should allow for recovery of effective shade based on site potential vegetation. The Malheur Basin TMDL and Water Quality Management Plan were completed September 2010. The Malheur National Forest is slated to complete a water quality recovery plan in 2015. Follow recommendations presented in these plans.

Agriculture Practices

- 1.3.2 Cool irrigation returns and run-off. Diversions or runoff warmer than the receiving water should be cooled when possible before allowing to it to enter the receiving system (e.g., subterranean pipes).

2. Actions to Address Demographic Threats

2.1. Connectivity Impairment

Entrainment & Fish Passage Issues

- 2.1.1 Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers. High priorities for screening include diversions on Lake Creek and Bosonberg Creek and the Drewsey Diversion. An inventory on Upper Malheur River is incomplete.
- 2.1.2 Provide passage at road-related barriers and culverts. The U.S. Forest Service is currently developing watershed action plans for the core area. These action plans include an analysis of passage issues as they relate to bull trout. Implement measures identified and prioritized in the U.S. Forest Service watershed action plans to provide passage at road related barriers including those on Corral Basin and Summit creeks.
- 2.1.3 Identify and remove barriers to juvenile and adult passage. Log weirs, culverts, legacy structures and other barriers impede juvenile and adult passage and prevent movement between spawning, rearing and overwinter habitats.

Dewatering

- 2.1.4 Improve and secure instream flows. Restore connectivity and opportunities for migration by securing instream flows and/or water rights. Improve irrigation efficiency. Lower Lake Creek in particular becomes dewatered due to management actions taken on upstream private property. In addition, implement channel restoration actions identified under Recovery Action 1. Benefits of stream channel restoration will include raising the water table and restoring natural instream flow, providing more flow during summer and late fall.

Temperature Barriers

- 2.1.5 Eliminate thermal barriers by maintaining or improving riparian vegetation communities, providing shade to streams, and increasing water quantity. Current juvenile bull trout distribution and movement is impeded by thermal barriers between spawning, rearing, and overwintering habitats. Seasonal thermal barriers exist at the mouths of Lake, McCoy, and Summit creeks as well as the upper mainstem river. Implement actions designed to cool warm water temperatures, increase flows, and improve hydrologic function as detailed under Recovery

Action 1 . Partnerships with private landowners may be necessary to eliminate thermal barriers, particularly on Lake Creek and McCoy Creek.

2.2. Fisheries Management

2.3. Small Population Size

At this time, we expect the implementation of the actions identified herein will be sufficient to increase population size and maintain gene flow among populations and will ameliorate any deleterious effects of genetic and demographic stochasticity in addition to recovering the migratory life history type. Additional measures, such as population augmentation or reintroduction within historical distribution, should be considered in the event a demographic response to these actions is not observed.

2.3.1 Investigate merits of developing a genetic management plan for the Upper Malheur core area given the extremely low population size and high abundance of brook trout.

2.4. Forage Fish Availability

3. Actions to Address Nonnative Fishes

3.1 Nonnative Fish

Competition and Hybridization

- 3.1.1 Develop and implement a comprehensive watershed-wide brook trout eradication and control strategy. In order to address the threat of brook trout long-term and over a biologically-supportable geographic scale, a watershed level plan must be developed and implemented so that eradication treatments can be evaluated in the context of overall long-term suppression, and to actively or passively encourage bull trout recolonization into treated areas. The strategy should prioritize stream reaches where success will be most likely and where threats to existing bull trout populations (*i.e.*, hybridization rates, competition, etc.) are most significant. Removal efforts should employ the use of tested and proven barriers to prevent re-invasion from adjacent reaches occupied by brook trout. Continue to work collaboratively with partner agencies to develop and implement this control and eradication strategy.
- 3.1.2 Implement actions that will ensure the expression of a migratory life history. Impacts of brook trout to bull trout populations appear to be most significant for populations of primarily resident fish. Bull trout populations containing large migratory individuals manage to maintain

despite the presence of brook trout. In addition, large fish are more fecund, have great productivity, and can out-compete smaller brook trout for food and space resources. Actions specific to fostering a migratory life history include those under Recovery Action 1 to ensure suitable overwintering habitat in the Malheur River, and those under Recovery Action 2.1 to maintain passable migratory corridors.

- 3.1.3 Develop and implement education and outreach efforts to address problems and consequences of unauthorized fish introductions. Target areas where intra-basin transfer of brook trout from adjacent systems would be most likely.

4. Research, Monitoring, and Evaluation

4.1 Habitat

Monitoring

- 4.1.1 Continue to monitor temperature, water quality, water quantity, and riparian condition to evaluate the effectiveness of habitat restoration actions and to provide the ability to detect trends in these metrics.

4.2 Demographic

Research

- 4.2.1 Further define bull trout distribution and habitat use in the core area. Research is needed to determine the extent to which bull trout express a fluvial, and potentially adfluvial, life history in the Upper Malheur River.

Monitoring

- 4.2.2 Continue ongoing population monitoring efforts within the basin. Maintain current long-term redd count datasets assessing abundance and distribution of spawning migratory bull trout. If necessary bolster the monitoring program with new protocols and methodologies consistent with other programs statewide. Continue to coordinate surveys among partner agencies.
- 4.2.3 Continue maintenance and operation of fish screens on all diversions. To prevent entrainment, consistent monitoring and maintenance is necessary to keep fish screens operating properly.

4.3 Nonnatives

Conservation Recommendations

- Promote interagency collaboration and coordination on bull trout recovery actions by supporting existing bull trout working groups or the formation of new bull trout working groups where they do not exist. While working groups may be facilitated by any interested stakeholder, most often they are organized and facilitated by the Service, a State agency, U.S. Forest Service, or a Tribal entity. Although the Service has no guidelines for format or process, existing working groups are largely informal, are organized at various scales (*e.g.*, core area, river basin, geographic region, or recovery unit), and generally meet at least annually.
- Provide long-term habitat protection through purchase of private property from willing sellers. Potential candidates include the remaining privately-held parcels in the Upper Malheur River corridor including tracts on lower Big, Summit, Lake, and Bosonberg Creeks.
- Identify and address sediment sources in Upper Malheur River basin affecting bull trout. Identify road-related sediment problem areas in the Upper Malheur River core area prioritizing spawning and rearing areas. Examine the ways roads capture and channel runoff, and changes in surface runoff associated with soil compaction. Stabilize roads, crossings, railroad grades, and other sources of sediment delivery; remove and vegetatively restore unneeded roads and railroad grades.
- Increase information outreach to anglers. Provide information on bull trout identification, special regulations, methods to reduce hooking mortality of bull trout caught incidentally, and the value of bull trout and their habitat. Education and outreach designed to assist anglers in identifying and differentiating captured brook trout from bull trout is needed to reduce unintended take of bull trout. Signage should be increased in Big Creek, Lake Creek, and access points along the main stem Upper Malheur River alerting anglers of bull trout presence in the streams.
- Investigate and implement actions to restore historic prey base by reintroducing anadromous species. Anadromous species such as steelhead and spring Chinook salmon were historically present in the Upper Malheur River. Feasibility of restoration of spawning populations of these species to increase prey base and provide marine derived nutrients should be evaluated and implemented where feasible and biologically supportable.

Jarbidge River Geographic Region

Jarbidge River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Ensure that sedimentation from road maintenance of the transportation system is minimized from entering bull trout critical habitat. Develop a road maintenance agreement with the Forest Service, Three Creeks, and local counties.
- Consider developing and implementing a vegetation management plan within the West Fork Jarbidge River.
- Consider working with Trout Unlimited to address legacy mine issues in the West Fork Jarbidge River.
- Consider development of a habitat conservation plan with private landowners in Dave Creek.

Weiser River Geographic Region

Weiser River Core Area

1. Actions to Address Habitat Threats

None

2. Actions to Address Demographic Threats

None

3. Actions to Address Nonnative Fishes

None

4. Research, Monitoring, and Evaluation

4.1 Habitat

4.2 Demographic

Monitoring

4.2.1 Continue ongoing population monitoring efforts within the basin.

Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.

4.3 Nonnatives

Conservation Recommendations

- Protect, restore, and maintain suitable habitat conditions for bull trout. Maintain or improve water quality in bull trout core areas.
- Implement brook trout removal efforts wherever feasible and biologically supported.

Implementation Schedule for the Upper Snake Recovery Unit

The Implementation Schedule that follows describes recovery action priorities, action numbers, action descriptions, duration of actions, potential or participating responsible parties, total cost estimate and estimates for the next 5 years, if available, and comments. These recovery actions, when accomplished in conjunction with implementation of recovery actions in the other bull trout recovery units, will lead to recovery of bull trout in the coterminous United States as discussed in the Revised Draft Bull Trout Recovery Plan (USFWS 2014).

Parties with authority, responsibility, or expressed interest to implement a specific recovery action are identified in the Implementation Schedule. Listing a responsible party does not imply that prior approval has been given or require that party to participate or expend any funds. However, willing participants will benefit by demonstrating that their budget submission or funding request is for a recovery action identified in an approved recovery plan, and is therefore part of a coordinated effort to recover bull trout. In addition, section 7(a)(1) of the Endangered Species Act (Act) directs all Federal agencies to use their authorities to further the purposes of the Act by implementing programs for the conservation of threatened or endangered species.

Threat Factor: Listing factor or threat category addressed by the recovery action.

- A. The Present or Threatened Destruction, Modification or Curtailment of Bull Trout Habitat or Range;
- B. Overutilization for Commercial, Recreational, Scientific, or Educational Purposes;
- C. Disease or Predation;
- D. The Inadequacy of Existing Regulatory Mechanisms; or
- E. Other Natural or Manmade Factors Affecting Its Continued Existence.

Recovery Action Priority: All priority 1 tasks are listed first, followed by priority 2 and priority 3 tasks.

Priority 1: An action that must be taken to prevent extinction or prevent the species from declining irreversibly in the foreseeable future.

Priority 2: An action that must be taken to prevent a significant decline in species population or habitat quality.

Priority 3: All other actions necessary to meet the recovery objectives.

For reference we also list additional conservation recommendations (marked Cons Rec). These actions are potentially beneficial for bull trout conservation and merit implementation, but they are not considered necessary to meet recovery objectives within a core area and so are not classified as Priority 1, 2, or 3. Conservation recommendations are not included in recovery cost estimates.

We evaluate recovery action priorities relative to the core area(s) where the action is targeted. Recovery action priorities may reflect both the severity of the threat and the expected effectiveness of the action in addressing it.

Research, monitoring and evaluation (RM&E) actions necessary for recovery are those deemed critical for developing information for planning, implementing, monitoring, and evaluating effectiveness of recovery actions addressing management of primary threats. Depending on the level of importance of this information, these RM&E actions may be classified as Priority 1, 2, or 3. Other RM&E actions, while possibly informative and potentially contributing to recovery, may not be deemed necessary and will thus be classified as conservation recommendations.

Recovery Action Number and Description: Recovery actions as numbered in the recovery outline. Refer to the Narrative for action descriptions.

Recovery Action Duration: Indicates the number of years estimated to complete the action, or other codes defined as follows:

Continual (C) – An action that will be implemented on a routine basis once begun.

Ongoing (O) – An action that is currently being implemented and will continue until no longer necessary.

To be Determined (TBD) – The action duration is not known at this time or implementation of the action is dependent on the outcome of other recovery actions.

Responsible or Participating Party: The following organizations are those with responsibility or capability to fund, authorize, or carry out the corresponding recovery tasks.

Salmon River Geographic Region

BLM	Bureau of Land Management
BPA	Bonneville Power Administration
IDFG	Idaho Department of Fish and Game

USFWS	U.S. Fish and Wildlife Service
USFS	U.S. Forest Service

Boise River, Payette River, Weiser River Geographic Regions

BLM	Bureau of Land Management
IDFG	Idaho Department of Fish and Game
USBR	U.S. Bureau of Reclamation
USFWS	U.S. Fish and Wildlife Service
USFS	U.S. Forest Service

Little Lost River Geographic Region

BLM	Bureau of Land Management
IDFG	Idaho Department of Fish and Game
USFWS	U.S. Fish and Wildlife Service
USFS	U.S. Forest Service

Malheur River Geographic Region

BLM	U.S. Bureau of Land Management
BPT	Burns Paiute Tribe
ID	irrigation districts
NRCS	Natural Resources Conservation Service
ODA	Oregon Department of Agriculture
ODEQ	Oregon Department of Environmental Quality
ODFW	Oregon Department of Fish and Wildlife
OSP	Oregon State Police
OWRD	Oregon Water Resources Department
TAC	Working Group Technical Advisory Committee
SWCD	Soil and Water Conservation District
USBR	U.S. Bureau of Reclamation
USFS	United States Forest Service
USFWS	U.S. Fish and Wildlife Service
VOID	Vale Irrigation District
WC	Watershed Councils
WSID	Warm Springs Irrigation District

Jarbidge River Geographic Region

BLM	Bureau of Land Management
EC	Elko County, Nevada
IDFG	Idaho Department of Fish and Game
Landowners	Private Landowners
NDOW	Nevada Department of Wildlife
OC	Owyhee County, Idaho
TU	Trout Unlimited

USFWS U.S. Fish and Wildlife Service
USFS U.S. Forest Service

Bolded type indicates the agency or agencies that have the lead role for task implementation and coordination, though not necessarily sole responsibility.

Cost estimates: Estimated costs assigned to each action identified in the Implementation Schedule, both for the first 5 years after release of the recovery plan and for the total estimated cost of recovery (based on time to recovery, for Continual or Ongoing actions). Cost estimates are not provided for tasks which are normal agency responsibilities under existing authorities.

An asterisk (*) in the total cost column indicates ongoing tasks that are currently being implemented as part of normal agency responsibilities under existing authorities. Because these tasks are not being done specifically or solely for bull trout conservation, they are not included in the cost estimates. Some of these efforts may be occurring at reduced funding levels and/or in only a small portion of the watershed.

Time to Recovery: Estimated time before this recovery unit could meet recovery criteria, if recovery actions are successfully implemented.

Table E-4. Upper Snake Recovery Unit Implementation Schedule

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Salmon River Geographic Region													
Little-Lower Salmon River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
South Fork Salmon River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Middle Salmon River - Chamberlin	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Middle Fork Salmon River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Middle Salmon River - Panther	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Lemhi River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Pahsimeroi River	A	1	1.2.1	Restore streams that are partially or completely dewatered. Improve instream flows.	5	BLM, IDFG, USFS		250	50	50	50	50	50
Pahsimeroi River		1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	5	BLM, IDFG, USFS		250	50	50	50	50	50

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Pahsimeroi River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Upper Salmon River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS		500	20	20	20	20	20
Opal Lake	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG		500	20	20	20	20	20
Lake Creek	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG		500	20	20	20	20	20
Estimated cost subtotal, Salmon River Geographic Region: \$5,500,000 (over 25 years, minimum estimate)													

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Boise River Geographic Region													
Anderson Ranch	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Arrowrock	A	1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	10	USFS	Ongoing. Cost covered under existing programs	*					
Arrowrock	A, E	1	2.1.2	Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.	5	USBR, IDFG, IDWR, USFWS		290	40	72	74	52	52
Arrowrock	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	25	BLM, IDFG, USFS		250	50	50	50	50	50
Arrowrock	N/A	N/A	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Estimated cost subtotal, Boise River Geographic Region: \$1,540,000 (over 25 years, minimum estimate)													

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Payette River Geographic Region													
Squaw Creek	A	1	1.1.1	Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.	25	IDFG, USFS		500	20	20	20	20	20
Squaw Creek	A	1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	10	USFS, IDFG	Ongoing. Cost covered under existing programs	*					
Squaw Creek	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	10	USFS, IDFG		250	50	50	50	50	50
Squaw Creek	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
North Fork Payette River	A	1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	10	USFS, IDFG	Ongoing. Cost covered under existing programs	*					

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
North Fork Payette River	E	1	2.3.1	Manage for demographic stochasticity by ensuring local populations contain more than 50–100 reproductive individuals and manage for environmental stochasticity with populations containing 1000-10,000 individuals, when practicable.	5	BLM, USBR, IDEQ, IDFG, IDL, USFWS, USFS		100	20	20	20	20	20
North Fork Payette River	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	10	USFS, IDFG		250	50	50	50	50	50
North Fork Payette River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Middle Fork Payette	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	10	USFS, IDFG		250	50	50	50	50	50
Deadwood River	A	1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	10	USFS, IDFG	Ongoing. Cost covered under existing programs	*					
Deadwood River	A, E	1	2.1.2	Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.	5	USBR, IDFG, IDWR, USFWS		290	40	72	74	52	52
Deadwood River	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	10	USFS, IDFG		250	50	50	50	50	50

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Deadwood River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Upper South Fork Payette River	A	1	2.1.1	Identify barriers for bull trout and implement tasks to provide passage.	10	USFS, IDFG	Ongoing. Cost covered under existing programs	*					
Upper South Fork Payette River	E	1	3.1.1	Implement brook trout removal efforts wherever feasible and biologically supported.	10	USFS, IDFG		250	50	50	50	50	50
Upper South Fork Payette River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Estimated cost subtotal, Payette River Geographic Region: \$4,140,000 (over 25 years, minimum estimate)													
Malheur River Geographic Region													
North Fork Malheur	A	1	1.1.1	Restore canopy and riparian cover, and native vegetation in all bull trout spawning, rearing and migration areas.	10	USFS, BLM, BPT, WC, landowners		250	25	25	25	25	25
North Fork Malheur	A	1	1.1.2	Implement stream restoration project in degraded stream reaches.	10	USFS, BPT, NRCS, SWCD, landowners	Ongoing. Cost covered under existing programs	*					

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)						
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19	
North Fork Malheur	A	2	1.1.3	Provide a reliable source of large hardwood beaver forage.	20	USFS, BPT, BLM, SWCD	Costs to be determined	*						
North Fork Malheur	A	2	1.1.4	Evaluate and implement actions to encourage beaver recolonization.	20	USFS, BPT, ODFW, SWCD	Costs to be determined	*						
North Fork Malheur	A	1	1.1.5	Reduce grazing impacts.	5	USFS, BPT, NRCS, SWCD, landowners		500	100	100	100	100	100	100
North Fork Malheur	A	2	1.1.6	Curtail unauthorized livestock use on USFS property.	5	USFS	Ongoing. Costs covered under existing programs.	*						
North Fork Malheur	A	2	1.2.1	Maintain a conservation pool in Beulah Reservoir to provide adequate overwinter rearing habitat for adult and sub-adult migratory bull trout.	20	USBR, USFWS, VOID, WSID	Ongoing. Costs variable depending on water year.	*						
North Fork Malheur	A	1	1.3.1	Maintain or improve effective streamside shade.	10	DEQ, WC, ODA, NRCS, USFWS		200	20	20	20	20	20	20
North Fork Malheur	A	3	1.3.2	Cool irrigation returns and run-off.	10	SWCD, ID, landowners	Costs to be determined	*						
North Fork Malheur	A	1	2.1.1	Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers.	10	ODFW, BPT, USFS, NRCS, BLM, SWCD, landowners		100	20	20	20	20	20	20
North Fork Malheur	A	2	2.1.2	Reduce occurrence of spill at Agency Valley Dam to minimize entrainment of bull trout below the dam and provide upstream passage when entrainment occurs.	20	USBR, ID, ODFW, USFWS	If spill occurs then spring time Trap and Haul is initiated.	35	7	7	7	7	7	7

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)						
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19	
North Fork Malheur	A	2	2.1.3	Identify and remove barriers to juvenile passage.	10	USFS, ODFW,		TBD						
North Fork Malheur	A,E	1	2.1.4	Improve and secure instream flows.	25	ODFW, OWRD, BPT	Partially covered by 1.1.2, 1.1.5	100	20	20	20	20	20	
North Fork Malheur	A	1	2.1.5	Eliminate thermal barriers by maintaining or improving riparian vegetation communities, providing shade to streams, and increasing instream flow.	10	USFS, BPT, NRCS, SWCD, WC, landowners	Cost covered in 1.1.1 & 1.1.2							
North Fork Malheur	E	2	3.1.1	Survey and monitor the North Fork watershed for the presence of brook trout.	25	ODFW, BPT, USFWS	Ongoing. Costs covered under existing programs.	*						
North Fork Malheur	E	2	3.1.2	Prioritize the removal of brook trout in adjacent basins.	25	ODFW, BPT, USFWS	Costs covered in Upper Malheur Plan.	*						
North Fork Malheur	A	1	3.1.3	Implement actions that will ensure the expression of a migratory life history.	25		Ongoing. Costs covered under existing programs.	*						
North Fork Malheur	E	1	3.1.4	Implement brook trout removal efforts when brook trout are detected in the North Fork Malheur core area.	25	ODFW, BPT, USFWS	Costs to be determined	*						
North Fork Malheur	E	2	3.1.5	Develop and implement an educational effort to address problems and consequences of unauthorized fish introductions.	1	ODFW, BPT, USFS, USFWS		10	10					

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)						
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19	
North Fork Malheur	A	2	4.1.1	Continue to monitor temperature, water quality, water quantity and riparian condition to evaluate the effectiveness of habitat restoration actions and to provide the ability to detect trends in these metrics.	25	USFS, BPT, BOR, ODEQ, SWCD	Ongoing. Costs covered under existing programs.	*						
North Fork Malheur	A	2	4.2.1	Further define bull trout distribution and habitat use in the core area.	25	ODFW, BPT		25	5	5	5	5	5	
North Fork Malheur	A	2	4.2.2	Continue maintenance and operation of fish screens on all diversions.	25	ODFW, BPT, USFS, USBR		50	2	2	2	2	2	
North Fork Malheur	A	2	4.2.3	Continue ongoing population monitoring efforts within the basin.	25		Ongoing. Costs covered under existing programs.	*						
North Fork Malheur	E	2	4.3.1	Implement bass and crappie monitoring efforts in Beulah Reservoir.	25	BOR	Costs to be determined	*						
Upper Malheur	A	1	1.1.1	Restore canopy and riparian cover, and native vegetation in all bull trout spawning, rearing and migration areas.	10	USFS, BLM, BPT, WC, landowners		250	25	25	25	25	25	
Upper Malheur	A	1	1.1.2	Implement stream restoration projects in degraded stream reaches.	10	USFS, USFWS	Ongoing. Costs covered under existing programs.	*						
Upper Malheur	A	2	1.1.3	Provide a reliable source of large hardwood for beaver forage.	20	USFS, BPT	Costs to be determined	*						
Upper Malheur	A	2	1.1.4	Evaluate and implement actions to encourage beaver recolonization.	20	USFS, ODFW, BPT	Costs to be determined	*						

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Upper Malheur	A	1	1.1.5	Reduce grazing impacts in all bull trout spawning areas.	5	USFS, BPT, NRCS, SWCD, landowners		500	100	100	100	100	100
Upper Malheur	A	2	1.1.6	Curtail unauthorized livestock use on USFS property.	5	USFS	Ongoing. Costs covered under existing programs.	*					
Upper Malheur	A	1	1.3.1	Maintain or improve effective shade.	10	DEQ, WC, ODA, NRCS, USFWS		250	25	25	25	25	25
Upper Malheur	A	3	1.3.2	Cool irrigation returns and run-off.	10	SWCD, ID, landowners	Costs to be determined.	*					
Upper Malheur	A	1	2.1.1	Install appropriate fish screens and passage structures around diversions and/or remove related migration barriers.	10	ODFW, BPT, USFS, NRCS, BLM, SWCD, landowners		100	20	20	20	20	20
Upper Malheur	A	2	2.1.2	Provide passage at road-related barriers and culverts.	25	USFS, ODOT, Counties	Ongoing. Project cost unknown at this time.	TBD					
Upper Malheur	A	2	2.1.3	Identify and remove barriers to juvenile and adult passage.	25	USFS, ODFW, BLM		50	10	10	10	10	10
Upper Malheur	A,E	1	2.1.4	Improve and secure instream flows.	25	ODFW, OWRD, BPT	Partially covered by 1.1.2, 1.1.5	100	20	20	20	20	20
Upper Malheur	A	1	2.1.5	Eliminate thermal barriers by maintaining or improving riparian vegetation communities and providing shade to streams.	10	USFS, BLM, BPT, WC, landowners	Cost covered in 1.1.1 & 1.1.2	250	25	25	25	25	25
Upper Malheur	A	2	2.3.1	Investigate merits of developing a genetic management plan for the Upper Malheur core area.	1	ODFW, USFWS, BPT	Costs to be determined.	*					

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)						
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19	
Upper Malheur	E	1	3.1.1	Develop and implement a comprehensive watershed-wide brook trout eradication and control strategy.	25	ODFW, BPT, USFWS, USFS	Costs to be determined.	*						
Upper Malheur	A	1	3.1.2	Implement actions that will ensure the expression of a migratory life history.	TBD		Ongoing. Costs covered under existing programs.	*						
Upper Malheur	E	2	3.1.3	Develop and implement education and outreach efforts to address problems and consequences of unauthorized fish introductions.	1	ODFW, BPT, USFS, USFWS		10	10					
Upper Malheur	A	2	4.1.1	Continue to monitor temperature, water quality, water quantity and riparian condition to evaluate the effectiveness of habitat restoration actions and to provide the ability to detect trends in these metrics.	25	USFS, BPT, BOR, ODEQ, SWCD	Ongoing. Costs covered under existing programs.	*						
Upper Malheur	A	2	4.2.1	Further define bull trout distribution and habitat use in the core area.	25	TAC, ODFW, BPT		25	5	5	5	5	5	5
Upper Malheur	A	2	4.2.2	Continue ongoing population monitoring efforts within the basin.	25	TAC, ODFW, USFW, BPT, USFWS	Ongoing. Costs covered under existing programs.	*						
Upper Malheur	A	2	4.2.3	Continue maintenance and operation of fish screens on all diversions.	25	ODFW, BPT, USFS, USBR, BLM	Ongoing. Costs covered under existing programs.	*						
Estimated cost subtotal, Malheur River Geographic Region: \$2,805,000 (over 25 years, minimum estimate)														

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Little Lost River Geographic Region													
Little Lost River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	BLM, IDFG, USFS, USFWS	Ongoing. Costs covered under existing programs.	25	1	1	1	1	1
Estimated cost subtotal, Little Lost River Geographic Region: \$25,000 (over 25 years, minimum estimate)													
Weiser River Geographic Region													
Weiser River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.	25	IDFG, USFS		500	20	20	20	20	20
Estimated cost subtotal, Weiser River Geographic Region: \$500,000 (over 25 years, minimum estimate)													

Core Area	Threat Factor	Recovery Action Priority	Recovery Action Number	Recovery Action Description	Recovery Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Jarbidge River Geographic Region													
Jarbidge River	N/A	3	4.2.1	Continue ongoing population monitoring efforts within the basin. Maintain current long term datasets assessing abundance and distribution of bull trout. Continue to coordinate surveys among partner agencies.		BLM, IDFG, NDOW, USFWS, USFS	Ongoing. Costs covered under existing programs.	25	1	1	1	1	1
Estimated cost subtotal, Jarbidge River Geographic Region: \$25,000 (over 25 years, minimum estimate)													
Estimated total cost of recovery actions within this recovery unit: \$14,535,000 (over 25 years, minimum estimate)													
Time to Recovery (estimated time required to meet recovery criteria within this recovery unit): 25 years (3-5 bull trout generations)													

Conservation Recommendations for the Upper Snake Recovery Unit

Core Area	Threat Factor	Action Priority	Action Number	Action Description	Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Little-Lower Salmon River	N/A	Cons Rec		Reduce general sediment production. Stabilize roads, road stream crossings, and other known sources of fine sediment delivery. Monitor John Day Creek and Slate Creek.									
Little-Lower Salmon River	N/A	Cons Rec		Manage for demographic stochasticity by ensuring local populations contain more than 50–100 reproductive individuals and manage for environmental stochasticity with populations containing 1000-10,000 individuals, when practicable. Focus on additional survey effort in smaller watersheds such as John Day Creek, Slate Creek, Lake-Lower Salmon, and Partridge Creek.									
Little-Lower Salmon River	N/A	Cons Rec		Implement brook trout removal efforts wherever feasible and biologically supported. Consider monitoring populations in John Day Creek and Slate Creek.									
Little-Lower Salmon River	N/A	Cons Rec		Coordinate bull trout recovery with listed anadromous fish species recovery in the Salmon River geographic area.									

Core Area	Threat Factor	Action Priority	Action Number	Action Description	Action Duration	Responsible Parties	Comments	Estimated Costs (x \$1,000)					
								Total Cost	FY 15	FY 16	FY 17	FY 18	FY 19
Anderson Ranch	N/A	Cons Rec		Implement actions necessary to accelerate recovery of riparian vegetation and streambanks and reduce negative effects from historic and current livestock grazing in identified problem areas.									
Anderson Ranch	N/A	Cons Rec		Identify barriers for bull trout and implement tasks to provide passage.									
Anderson Ranch	N/A	Cons Rec		Implement brook trout removal efforts wherever feasible and biologically supported.									
Anderson Ranch	N/A	Cons Rec		Operate reservoirs and dams to minimize negative effects on bull trout in reservoirs and habitat downstream.									
North Fork Malheur		Cons Rec		Promote interagency collaboration and coordination on bull trout recovery actions by supporting existing bull trout working groups or the formation of new bull trout working groups where they do not exist.	Ongoing	TAC							
North Fork Malheur		Cons Rec		Identify and address sediment sources in North Fork Malheur Basin affecting bull trout.	10	USFS, BLM, USBR, NRCS, SWCD							
North Fork Malheur		Cons Rec		Increase information outreach to anglers.	Ongoing	ODFW, BPT, USFS, USFWS							

References

Literature Cited

- Batt, P. E. 1996. State of Idaho Bull Trout Conservation Plan. Office of the Governor of the State of Idaho, July 1, 1996.
- Bengeyfield, P. 2006. Managing cows with streams in mind. *Rangelands* 28(1):3-6.
- Buchanan, D. V. and S. V. Gregory. 1997. Development of water temperature standards to protect and restore habitat for bull trout and other cold water species in Oregon. Pages 1-9 in: W. C. Mackay, M. K. Brewin and M. Monita, editors. Bull Trout II Conference. Proceedings. Trout Unlimited Canada, Calgary, Alberta.
- Burton, T. A., S. J. Smith, and E. R. Cowley. 2011. Riparian area management: Multiple indicator monitoring (MIM) of stream channels and streamside vegetation. Technical Reference 1737-23. BLM/OC/ST-10/003+1737+REV. U.S. Department of the Interior, Bureau of Land Management, National Operations Center, Denver, Colorado. 155 pp.
- Clary, W. P., and B. F. Webster. 1989. Managing grazing of riparian areas in the Intermountain Region. General Technical Report INT-263. USDA Forest Service, Intermountain Research Station. Ogden, Utah.
- DeHaan, P. W., M. Diggs and W. R. Ardren. 2007. Genetic population structure of bull trout in the Malheur River basin, Oregon. Report to the Burns Paiute Tribe, Burns, OR. September 28, 2007. 14 pages.
- Goss, L. 2013. Livestock disturbance protocols: understanding the relationship between bank alteration and stream conditions. Master's Thesis. Utah State University.
- [IDFG] High, B, Meyer, K., Schill, D., and E. Mamer. 2005. Bull trout status review and assessment in the State of Idaho. Grant #F-73-R-27. Idaho Department of Fish and Game. 57pp.
- [IDFG] High, B, Meyer, K., Schill, D., and E. Mamer. 2008. Distribution, abundance, and population trends of bull trout in Idaho. *North American Journal of Fisheries Management* 28:1687-1701.
- [LLRITAT] Little Lost River Interagency Technical Advisory Team. 1998. Little Lost River key watershed bull trout problem assessment. Prepared for the State of Idaho. June 29, 1998.
- [IDFG] Meyer, K., Garton, E., and D. J. Schill. 2014. Bull trout trends in abundance and probabilities of persistence in Idaho. *North American Journal of Fisheries Management*. 34:1, 202-214.
- Montana Bull Trout Restoration Team. 2000. Restoration Plan for bull trout in the Clark Fork River Basin and Kootenai River Basin, Montana. Montana Fish, Wildlife and Parks, Helena, Montana.

- Petersen, J. H., and E. E. Kofoot. 2002. Conditions for growth and survival of bull trout in Beulah Reservoir, Oregon. Annual Report for 2001. Report to the U.S. Bureau of Reclamation, Pacific Northwest Region, Boise, Idaho. 43p.
- [USFWS] U.S. Fish and Wildlife Service. 2002a. Draft recovery plan for bull trout (*Salvelinus confluentus*) in the coterminous United States: Klamath River and Columbia River Distinct Population Segments. U.S. Fish and Wildlife Service, Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2002b. Draft recovery plan for bull trout (*Salvelinus confluentus*) in the coterminous United States: St. Mary-Belly River Distinct Population Segments. U.S. Fish and Wildlife Service, Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2004a. Designation of critical habitat for the Klamath River and Columbia River populations of bull trout. Federal Register 69:59996-60075.
- [USFWS] U.S. Fish and Wildlife Service. 2004b. Draft recovery plan for the Coastal-Puget Sound distinct population segment of bull trout (*Salvelinus confluentus*). Puget Sound Management Unit, Portland, Oregon. 389 + xvii p.
- [USFWS] U.S. Fish and Wildlife Service. 2004c. Draft recovery plan for the Jarbidge River distinct population segment of bull trout (*Salvelinus confluentus*). Portland, Oregon. 132 + xiii p.
- [USFWS] U.S. Fish and Wildlife Service. 2005a. Endangered and threatened wildlife and plants; designation of critical habitat for the bull trout. September 26, 2005. 70 FR 56212-56311.
- [USFWS] U.S. Fish and Wildlife Service. 2005b. Bull trout core area templates - complete core area by core area analysis. W. Fredenberg and J. Chan, *editors*. U. S. Fish and Wildlife Service. Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2005c. Bull trout core area conservation status assessment. W. Fredenberg, J. Chan, J. Young, and G. Mayfield. U. S. Fish and Wildlife Service. Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2008. Bull trout core area templates - complete core area by core area re-analysis. W. Fredenberg and J. Chan, *editors*. U. S. Fish and Wildlife Service. Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2010. Endangered and threatened wildlife and plants; revised designation of critical habitat for bull trout in the coterminous United States; final rule. October 18, 2010. Federal Register 75:63898-64070.
- [USFWS] U.S. Fish and Wildlife Service. 2014. Revised recovery plan for the coterminous United States population of bull trout (*Salvelinus confluentus*). Portland, Oregon. xii + 151 pages.

Appendix I. Core Area Technical Meeting Summaries

Twin Falls 12/3/14

Participants: Kate Crane (BLM), Doug Megargle (IDFG), Mark Moulton (FS-SNRA), Brenda Mitchell (FS-SNF), Steve Duke (FWS), Chris Reighn (FWS), Pam Druliner (FWS), Barb Schmidt (FWS), Ben Matibag (FWS), via phone Erika Phillips (FS-SNF)

Jarbidge River: None (but check with Natural Resource Agencies in Nevada)

Upper Salmon River: brook trout (Salmon Meeting – did not identify any primary threats)

Anderson Ranch: maybe wildfire (post fire effects) (Boise Meeting – identified brook trout for resident populations)

The Service attended a meeting in Elko, Nevada on March 10, 2015 and the technical group determined that there were no primary threats present in the Jarbidge River. No trend data has been collected for the Jarbidge River core area but large areas are designated Wilderness or Wild and Scenic Rivers which offers protection for local populations and critical habitat.

The Service reviewed information regarding brook trout as a primary threat in the Upper Salmon River core area. Technical partners in Salmon did not believe that brook trout was a primary threat. Brook trout are limited in the Upper Salmon River to a few drainages and concentrated within the Valley Creek drainage and did not pose a risk to the persistence of the entire core area.

The Service also reviewed information to consider wildfire as a threat in Anderson Ranch and determined that wildfire should not be considered a primary threat. The effects of wildfire are difficult to determine on local populations. Recent wildfires have impacted known migratory corridors within this core area but have not impacted spawning and rearing areas. Primary threats are those threats that are known or likely to impact populations of bull trout and subsequently the core area. The effects of wildfire may be considered as speculative in nature.

Salmon 12/4/14

Participants: Ben Matibag (FWS), Laura Berglund (FWS), Ryan Beatty (BLM-Challis), Eric Reilend (BLM-Challis), Christine Stewart (FS-Salmon Challis), Lucy Littlejohn (BLM-Salmon), Greg Schoby (IDFG), Windy Davis (IDFG), Dan Garren (IDFG), Tricia Miller (BLM-Salmon), Mike Edmondson (OSC), Arn Berglund (BLM)

Upper Salmon River: None (Twin Falls Meeting – identified brook trout)

Lemhi River: None

Pahsimeroi River: connectivity, instream impacts, maybe small populations

(add Big Gulch as a local population)

Little Lost River: None

Middle Salmon River-Panther: None

(Delete Allison Creek as a local population)

Lake Creek: None

Opal Lake: None

Technical partners in Twin Falls believed that brook trout should be identified as a primary threat. The Service reviewed information regarding brook trout as a primary threat in the Upper Salmon River core area. Technical partners in Salmon did not believe that brook trout was should be a primary threat. Brook trout are limited in the Upper Salmon River to a few drainages and concentrated within the Valley Creek drainage and did not pose a risk to the persistence of the entire core area.

Trend data showed that populations were stable or increasing in the Upper Salmon River and Lemhi Rivercore areas. Technical partners determined that no primary threats existed in the Middle Salmon River-Panther and Little Lost River even though IDFG trend data showed either a decreasing trend or a stable-decreasing trend. Technical partners also determined that Opal Lake and Lake Creek contained no primary threats. No trend data exists for the Pahsimeroi River, Lake Creek, or Opal Lake.

Boise 12/8/14

Participants: Ben Matibag (FWS), Dave Mays (FS-BNF), Chris Reighn (FWS), Michael Morse (FWS), Dmitri Vidergar (BOR), Art Butts (IDFG), Sam Easton (OSC), Joe Kozfkay (IDFG), Dave Statler (Nez Perce Tribe), Herbert Roerick (FS-BNF), Steve Duke (FWS), Mark Robertson (FWS), via phone Paul Moroz (FWS Contractor)

Arrowrock: reservoir management, connectivity impairment, upland/riparian management (roads, grazing, mining), **nonnative fish**, **climate change**

Deadwood River: maybe entrainment, **nonnative fish**, **connectivity impairment**, **reservoir management**

Upper South Fork Payette River: climate change, future urban development, brook trout **(delete)(delete)**, **nonnative fish (delete)**, **connectivity impairment**

Squaw Creek: connectivity impairment, grazing, **nonnative fish**

North Fork Payette River: passage barriers **(delete)**, **connectivity impairment**, brook trout **(delete)**, **nonnative fish**, maybe small populations

Anderson Ranch: maybe brook trout to resident populations (Twin Falls Meeting – identified wildfire), **nonnative fish**, **connectivity impairment**, **climate change**, **riparian management (road density, grazing)**, **unmanaged water diversions**, irrigation, etc

Middle Fork Payette River: brook trout **(delete)**, **nonnative fish**

Middle Fork Salmon River: None **(delete)**, **nonnative fish**, **connectivity impairment**

South Fork Salmon River: chemical spill, brook trout **(delete)**, **nonnative fish**, **connectivity impairment**

Red edits were suggested by Boise National Forest

Blue edits were suggested by Idaho Department of Fish and Game

To attempt to be inclusive the Service has included most if not all threats recommended by the Boise National Forest and Idaho Department of Fish and Game. The Service elected to remove

climate change as a threat. The best conservation action to combat climate change is most likely improving or promoting connectivity between local populations and within core areas.

Trend data exists only for the South Fork Salmon River, Middle Fork Salmon River, and Anderson Ranch and trend data indicates that populations are increasing in both the South Fork Salmon River and Anderson Ranch. In the Middle Fork Salmon River, trend data indicates a slight decline but discussions with various technical partners indicate that there are few if any threats in the Middle Fork Salmon River. The Middle Fork Salmon River is predominately in a designated wilderness area. For these three core areas the Service made the determination that no primary threats were present. All other core areas do not have sufficient data to assess trends.

Some threats that were removed were determined by the Service to not meet the definition of a primary threat (climate change, future urban development, entrainment and chemical spills).

McCall 12/11/14

Participants: Ben Matibag (FWS), Clayton Nalder (FS-PNF), Jason Greenway (FS-PNF), Dale Allen (IDFG), Paul Janssen (IDFG), Jeff Dillon (IDFG), Chris Reighn (FWS), Caleb Zurstadt (FS-PNF), Allison Turner (FWS), via phone Paul Moroz (FWS Contractor)

Pine Creek, Indian Creek, Wildhorse River: brook trout, connectivity impairment, instream flows (these are only on Idaho portion, waiting for information on the Oregon side for Pine Creek)

Weiser River: brook trout, climate change (delete) (Grangeville Meeting – IDL suggested pond operations, brook trout)

North Fork Payette River: brook trout, small population

Little Lower Salmon River: diversions (delete), forest practices (delete) (Grangeville Meeting – identified brook trout, wildfire, and small populations predominately in Slate Creek and John Day) – No threats

(Hard Creek should not be a local population)

Middle Fork Salmon River: No threats

Middle Salmon River-Chamberlain: No threats

South Fork Salmon River: Possibly none (Grangeville Meeting – suggested to consider brook trout)

Red edits suggested by Idaho Department of Fish and Game, IDFG also suggested to not implement actions in the North Fork Payette as it was “a lost cause”

Trend information is known for the Weiser River, Little Lower Salmon River, Middle Fork Salmon River, Middle Salmon River-Chamberlain, and South Fork Salmon River core areas. For the Middle Fork Salmon, technical partners determined that there were no primary threats present even though the IDFG data indicates a declining trend. The other four core areas were determined to have either a stable or increasing trends by IDFG data and therefore no primary threats were identified.

In the Pine, Indian, Wildhorse core area we determined that brook trout was a primary threat but most other threats were in the Pine Creek drainage in Oregon.

The Little Lower Salmon River core area contained both weir data and snorkel data compiled by IDFG. The weir data at Rapid River indicated that trends were increasing while snorkel data in both John Day and Partridge indicate that trends were decreasing. The Service relied more on data from the Rapid River data due to Rapid River containing a large fluvial population in making the determination that there were no primary threats.

The Service elected to remove climate change as a threat. The best conservation action to potential address climate change is improving or promoting connectivity between local populations and within core areas. The effects of climate change are also difficult to determine, we acknowledge that climate change will have effects but are uncertain when those impacts would occur.

Grangeville 12/18/14

Participants: Ben Matibag (FWS), Katherine Thompson (FS), Craig Johnson (BLM), via phone Chris Tretter (IDL), via phone Michelle Anderson (IDL), via phone Chris Reighn (FWS), via phone Paul Moroz (FWS Contractor)

Little Lower Salmon River: brook trout, wildfire, maybe small populations (McCall – identified diversions and forest practices, and it was later suggested by IDFG that there are none)

(Hard Creek should not be a local population)

Middle Salmon River-Chamberlain: None

Discussion with IDFG indicated that they felt that since Rapid River contains a strong fluvial component that the remaining local populations located near the lower Main Salmon were not needed to ensure the persistence of the core area. FS and BLM felt the opposite in which those local populations are needed for the persistence of the core area. Trends in Rapid River collected from weir data indicate that the population is increasing while snorkel data from the smaller local populations indicate that they are decreasing. After reviewing the information the Service determined that there were no primary threats in the Little Lower Salmon River core area.

Trend data shows that populations in the Middle Salmon River – Chamberlain core area are stable and technical partners did not identify any primary threats.

Appendix II. Core Area Summaries

Summary for: Little Lower Salmon River Core Area

Geographic Description

This core area occurs in Idaho County and extends from the watersheds of the confluence of the mainstem Salmon River with the Snake River, upstream to the confluence with French Creek. In addition, the Little Salmon River watershed is included, which flows into the Salmon River at River kilometer 139 (River Mile 86.7). The western boundary is formed by Hells Canyon on the north and by the Seven Devils Mountains on the south. The eastern boundary starting from the south is the watershed crest at the headwaters of the North Fork Payette River and it continues north and crosses the Salmon River below Burgdorf Summit. This boundary continues north to the headwaters of Little Slate Creek and White Bird Creek and curves to the west around the east side of the Craig Mountains. The core area is 455,160 hectares (1,124,700 acres) and the land ownership in this core area differs from other core areas in that it contains a larger amount (approximately 38 percent) of private land. The Bureau of Land Management (6 percent) and the U.S. Forest Service (47 percent) manage the majority of lands within the core area.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least seven streams or stream complexes (*i.e.*, local populations). These local populations include Slate Creek, John Day Creek, Rapid River, Boulder Creek, Hard Creek, Lake/Lower Salmon, and Partridge Creek.

The mainstem Salmon River provides for migration, and adult and sub adult foraging, rearing and wintering habitat. The Little Salmon River also provides for foraging/adult rearing habitat and connectivity between local populations in the core area.

Current trend information for the Rapid River portion of this core area shows that the population is increasing while surveys in Slate Creek and John Day indicate that those populations are decreasing. The Service will attempt to be conservative in identifying primary threats and conservation actions.

Threats:

- Degraded Habitat
- Instream Flows
- Livestock Grazing
- Small Population

- Brook trout

Primary Threats Proposed:

- brook trout
- small population size

Summary for: South Fork Salmon River Core Area

Geographic Description

This core area occurs in Valley and Idaho Counties and enters the Mainstem Salmon River east of French Creek and extends south to its headwaters upstream of Warm Lake. The ridges that form the eastern boundary of this relatively narrow, north-south oriented area lie in the headwaters of the Middle Fork Salmon River and Big Creek. The western boundary is the divide between the upper North Fork Payette River and the South Fork Salmon River. The core area is 338,100 acres (835,000 hectares) in size.

The U.S. Forest Service manages 99 percent of the land in this core area.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 28 streams or stream complexes (*i.e.*, local populations). These local populations include Upper Lake Creek, Grouse-Flat Creek, Ruby Creek, Summit Creek, Victor Creek, Loon Creek, Lick Creek, Zena Creek, Fitsum Creek, Buckhorn Creek, Cougar Creek, Fourmile Creek, Blackmare Creek, Dollar-Six Bit Creeks, Warm Lake, Curtis Creek, Upper South Fork Salmon River, Burntlog Creek, Trapper Creek, Riordan Lake, Upper East Fork South Fork Salmon River, Sugar Creek, Tamarack Creek, Profile Creek, Quartz Creek, Elk Creek, and Pony Creek.

Both resident and fluvial populations of bull trout were documented in the mainstem South Fork Salmon River. Overwintering fluvial bull trout were observed in the lower South Fork Salmon River from the Sheep Creek confluence downstream to the mouth of the South Fork Salmon River. Bull trout also overwintered in the mainstem Salmon River from the Elkhorn Creek confluence upstream to Big Mallard Creek.

IDFG trend data indicates that this core area is increasing.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

- Connectivity Impairment
- Habitat Degradation
- Brook Trout

Proposed Primary Threats:

None

Summary for: Middle Fork Salmon River - Chamberlain Core Area

Geographic Description

This area occurs in Idaho County and includes the Salmon River from its confluence with the Middle Fork Salmon River downstream to French Creek on the western boundary. The northern boundary is comprised of the peaks that separate the Salmon River basin from the Clearwater basin. The southern boundary follows the ridges between Farrow Mountain and Mosquito Peak and then continues to the mouth of the South Fork Salmon River. The core area covers 350,700 hectares (866,600 acres) and 99 percent of this area is managed by the Federal government.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 9 streams or stream complexes (*i.e.*, local populations). These local populations include Bargamin Creek, Warren Creek, Fall Creek, California Creek, Wind River, Sheep Creek, Big Squaw Creek, Sabe Creek, and Chamberlain Creek.

The mainstem Middle Fork Salmon River provides for migration, and adult and sub adult foraging, rearing and wintering habitat. Bull trout spawning and rearing occurs in the upper reaches of the creeks, and subadult and adult rearing occurs in the remainder of the drainages.

IDFG trend data indicates that this core area is increasing.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

None

Summary for: Middle Fork Salmon River Core Area

Geographic Description

This core area includes the entire Middle Salmon River drainage which lies in Idaho County and is mostly in the Frank Church River of No Return Wilderness. The southern boundary is in the headwaters of Bear Valley Creek and the mountains to the north of Big Creek from the northern boundary. The eastern boundary follows the ridge line of the high peaks west of Panther Creek the Main Salmon River, and McElney Mountain and Twin Peaks.

This area encompasses 744,300 hectares (1,839,000 acres) and 99 percent of this area is managed by the U.S. Forest Service.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 28 streams or stream complexes (*i.e.*, local populations). These local populations include Bear Valley Creek, Marsh Creek, Upper Middle Fork Salmon River 1, Upper Middle Fork Salmon River 2, Mayfield Creek, Rapid Creek, Pistol Creek, Little Loon Creek, Warm Spring Creek, Loon Creek, Camas Creek, Lower Middle Fork Salmon River 1, Lower Middle Fork Salmon River 2, Lower Middle Fork Salmon River 3, Marble Creek, Monumental Creek, Big Raney Creek, Big Creek 1, Big Creek 2, Big Creek 3, Big Creek 4, Beaver Creek, Rush Creek, Silver Creek, Yellowjacket Creek, Wilson Creek, Indian Creek, and Sulphur Creek.

IDFG estimates that this core area contains some of the strongest bull trout local populations in the Pacific Northwest. The mainstem Middle Fork Salmon River provides for migration, and adult and sub adult foraging, rearing and wintering habitat. Bull trout spawning and rearing occurs in the upper reaches of the creeks, and subadult and adult rearing occurs in the remainder of the drainages.

IDFG trend data indicates that this core area is decreasing but technical partners determined that trends were stable.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

Brook trout

Summary for: Middle Salmon River- Panther Core Area

Geographic Description

The Middle Salmon River – Panther Creek core area is located in Lemhi and Idaho Counties. This area is bordered on the west by the mountains west of Panther Creek, the Bighorn Crags and Quartzite Mountain; the southeast boundary is the Lemhi Mountain Range; and the northeast boundary is the Bitterroot Mountain Range. The core area comprises 557,450 hectares (1,377,500 acres).

Land ownership in the Middle Salmon River - Panther basin is predominantly Federal. The U.S. Forest Service (86 percent) and the Bureau of Land Management (9 percent) manage the majority of lands within the core area. Privately owned lands make up about 5 percent of the total land in the basin.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 20 streams or stream complexes (*i.e.*, local populations). These local populations include Cow Creek, Hat Creek, McKim Creek, Iron Creek, Williams Creek, Carmen Creek, Fourth of July Creek, Jesse Creek, Twelve Mile Creek, North Fork Salmon River, Indian Creek, Squaw Creek, Spring Creek, Owl Creek, Boulder Creek, Pine Creek, Horse Creek, Panther Creek, Napias Creek, and Allison Creek. Most populations appear to exhibit resident life history expression.

They are also present in the mainstem Salmon and North Fork Salmon Rivers and in many streams of the Panther Creek drainage.

IDFG trend data indicates that this core area is decreasing but technical partners determined that trends were stable.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

- Livestock Grazing
- Mining
- Connectivity Impairment
- Brook Trout

Summary for: Lemhi River Core Area

Geographic Description

The Lemhi River core area occurs in the Boise River basin and is located in Lemhi County. This core area includes the Lemhi River and is bordered by the rugged Bitterroot Range of the Beaverhead Mountains to the north and east and the Lemhi Mountain Range to the west.

The core area is 327,260 hectares (808,670 acres) with federally-managed land divided equally between the U.S. Forest Service (40 percent) and the Bureau of Land Management (39 percent); 19 percent is privately managed.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least six streams or stream complexes (*i.e.*, local populations). These local populations include Hayden Creek, Pattee Creek, Upper Lemhi River, Geertson Creek, Kenny Creek, and Bohannon Creek.

Most bull trout are found in isolated resident populations but the mainstem Lemhi River contains fluvial bull trout. Connectivity between the tributaries and the Lemhi River is reduced because of migration barriers. Hayden Creek has year-round connectivity to the Lemhi River and contains a fluvial population.

IDFG trend data indicates that this core area is increasing.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

- Connectivity Impairment
- Habitat Degradation

Summary for: Pahsimeroi River Core Area

Geographic Description

The Pahsimeroi River core area occurs in the Boise River basin and is located in Lemhi and Custer Counties. The Pahsimeroi River watershed is located on the east side of the Salmon River, and includes the west slope of the Lemhi Mountain Range and the east slope of the Pahsimeroi Mountains in the Lost River Range. The valley floor has a low elevation of 1,418 meters (4,648 feet) and is characterized by well-developed alluvial fans that extend from the mountain fronts to near the center of the valley floor. The boulder, cobble, and gravel fans cover a large underground reservoir which provides the majority of the water that emerges as springs along the valley floor. The main Pahsimeroi River switches to subterranean flow during the late summer and winter.

Ninety-one percent of the Pahsimeroi River core area is in public ownership. The U.S. Forest Service manages 46 percent of the land area. This core area also has the highest percentage of land managed by the Bureau of Land Management (42 percent) of any of the core areas in this recovery unit. The Pahsimeroi River core area covers 217,200 hectares (536,800 acres).

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least eight streams or stream complexes (*i.e.*, local populations). These local populations include Upper Pahsimeroi River, Big Creek, Patterson Creek, Falls Creek, Morse Creek, Morgan Creek (includes the lower Pahsimeroi River), Tater Creek, and Ditch Creek.

Bull trout in the Pahsimeroi core area are found in most of the tributaries that drain the eastern, southern and southwestern portion of the area. The creeks in Upper Pahsimeroi River were considered a population stronghold in the Pahsimeroi River core area during the Subbasin Review process. The mainstem Pahsimeroi River serves as a migratory corridor for fish access to the mainstem Salmon River but lacks connectivity in multiple places on the mainstem.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Major Threats:

- Connectivity Impairment
- Instream Flows
- Instream Structures

Summary for: Upper Salmon River Core Area

Geographic Description

This core area occurs in Custer County and extends from the mouth of the Pahsimeroi River to the headwaters in the Sawtooth Mountains, including the mainstem Salmon River and tributaries. The area covers 6,242 square kilometers (2,410 square miles) and contains 5,230 kilometers (3,251 miles) of streams. Eighty-nine percent of this core area is in public ownership, and most of this public land is managed by the Federal government. The U.S. Forest Service manages 99 percent of the land in this core area.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 18 streams or stream complexes (*i.e.*, local populations). These local populations include Valley Creek, Basin Creek, Yankee Fork Creek, Thompson Creek, Squaw Creek, Challis Creek, Garden Creek, Morgan Creek, East Fork Salmon River, Slate Creek, Warms Springs Creek, Fourth of July Creek, Germania Creek, Upper Salmon River, Alturas Lake Creek, Pettit Lake, Yellowbelly Creek, and Redfish Lake Creek.

Both resident and migratory or fluvial bull trout are present in the Sawtooth Valley. The inlet of Alturas Lake has adfluvial bull trout and is one of the largest local populations in the Sawtooth Valley. Adfluvial bull trout are present in Redfish Lake. Bull trout were observed in the lower and middle reaches of Fourth of July Creek. Bull trout are found in the Valley Creek areas and are most persistent in headwater segments of several drainages. Bull trout snorkel inventories conducted by the U.S. Forest Service in the Yankee Fork of the Salmon River detected the greatest densities of fish in slow water habitat types near headwater reaches. High densities of bull trout have been documented in tributaries to the East Fork Salmon River in Big Boulder, Herd, and Warm Spring Creeks. Mainstem Challis Creek contains bull trout; however, bull trout occupancy is unknown in its tributaries. Bull trout generally move into spawning tributaries beginning in August and spawn in mid-to late-September and October within the Upper Salmon River core area. However, in the headwaters of the Salmon River, spawning has been documented in early August.

Trend information from IDFG 2014 indicates that this core area is increasing.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

- Connectivity Impairment
- Habitat Degradation
- Brook Trout

Summary for: Lake Creek Core Area

Geographic Description

This core area includes an isolated bull trout population in Williams Lake and Lake Creek. The core area is located on the west side of the Salmon River between the mouth of the Pahsimeroi and Lemhi rivers, approximately 19 kilometers (12 miles) south of Salmon, Idaho. Williams Lake was formed 8,000 to 10,000 years ago when a massive landslide dammed a creek in the steep-sided canyon and created a uniform basin. No surface outlet exists to the lake. At the base of the landslide area that created the lake, a spring-fed stream is apparently connected to the lake. The elevation of the lake is 1,601 meters (5,250 feet) and the watershed of 4,554 hectares (11,245 acres, 17.5 square miles) that surrounds the lake is 98 percent Federal land managed by the U.S. Forest Service and the Bureau of Land Management.

Current Distribution and Abundance

Bull trout are located in Williams Lake and upstream of the lake in Lake Creek. Bull trout comprise approximately 20 percent of the fish population in Williams Lake and their numbers appear to be stable but there is insufficient data to establish trend criteria for the small population in Lake Creek.

Water quality in Williams Lake in the Lake Creek core area upstream of Salmon, Idaho is impacted by recreational residential development surrounding the lake. Since this lake has no outlet, water quality has declined in recent years and may be impacting bull trout.

Summary for: Opal Lake Core Area

Geographic Description

This core area encompasses a small, isolated bull trout population in Opal Lake and upstream of the lake in Opal Creek. The area is located in the headwaters of the Panther Creek watershed and is encompassed by the Middle Salmon River-Panther core area. This natural lake has no outlet. The elevation of the lake is 2,300 meters (7,546 feet) and the watershed contains 518 hectares (1,280 acres, 1.9 square miles). The entire area is managed by the U.S. Forest Service.

Current Distribution and Abundance

Bull trout have been located by the Idaho Department of Fish and Game in Opal Lake. The lake is oligotrophic (low nutrient levels and high dissolved oxygen) and has no outlet. Good spawning habitat is located upstream of the lake; however, no positive identification of redds has been made to date.

Insufficient data is available to establish trend criteria for the small population in Opal Lake.

Summary for: Anderson Ranch Core Area

Geographic Description

Anderson Ranch core area occurs in the Boise River basin and is located in Camas and Elmore Counties. Anderson Ranch Dam on the South Fork Boise is the lower extent of the core area and presents an impassable barrier to upstream fish movement. The core area comprises approximately 257,700 hectares (636,970 acres).

Anderson Ranch Dam, on the South Fork Boise River, blocks access of bull trout residing in the lower South Fork Boise River, North Fork Boise River, and Middle Fork Boise River to the upper portion of the South Fork Boise River basin. The dam is approximately 100 meters (332 feet) tall and has no provisions for either upstream or downstream fish passage.

The Boise National Forest manages 85 percent of the watershed and private lands accounts for 11 percent.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 11 streams or stream complexes (*i.e.*, local populations). These local populations include Elk Creek, Trinity Creek (including Parks Creek), Willow Creek, Deadwood Creek, Boardman Creek (including Smokey Dome Canyon), Skeleton Creek, Bear Creek, Ross Fork Creek (including Johnson Creek and upper S.F. Boise River), Emma Creek, Big Smokey Creek (including West Fork Big Smokey), and Bluff Creek.

Migratory bull trout abundance has been estimated in Anderson Ranch Reservoir. During 1999 through 2000, abundance of adult migratory bull trout in Anderson Ranch Reservoir was estimated at 368 individuals.

The trend information based on IDFG 2014 data indicates that the core area is stable.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

None

Summary for: Arrowrock Core Area

Geographic Description

Arrowrock core area occurs in the Boise River basin and is located in Elmore and Boise Counties. Arrowrock dam on the Boise River is the lower extent of the core area and presents an impassable barrier to upstream fish movement. The core area is approximately 315,800 hectares (780,300 acres). The Boise National Forest manages 89 percent of the watershed while private lands accounts for 5 percent.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 18 streams or stream complexes (*i.e.*, local populations). These local populations include Upper Crooked River, Bear River, Bear Creek, Lodgepole Creek, Upper North Fork Boise River (including McLeod Creek, McPhearson Creek, Ballentyne Creek, and West Fork Creek), Cow Creek, Big Silver Creek, Johnson Creek, Blackwarrior Creek, Little Queens River, Queens River, Yuba River (including Trail Creek), Grouse Creek (Yuba River tributary), Decker Creek (Yuba River tributary), Buck Creek, Roaring River, Sheep Creek, and Rattlesnake Creek.

During 1996 through 1997, abundance of adult migratory bull trout (*i.e.*, fish greater than 300 millimeters (12 inches)) in Arrowrock Reservoir was estimated at 471 individuals. Current adult abundance is unknown.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Threats: (Primary Threats Identified by technical partners)

- Connectivity Impairment
- Habitat Degradation
- Water Management
- Nonnative fish

Summary for: Squaw Creek Core Area

Geographic Description

Squaw Creek core area occurs in the Payette River basin and is located in Gem, Boise, Washington, and Valley counties. The Squaw Creek drainage joins the mainstem Payette River as part of the Black Canyon Reservoir. The core area is approximately 88,300 hectares (218,200 acres). The Boise National Forest manages 47 percent of the watershed while private lands accounts for 40 percent.

Current Distribution and Abundance

Bull trout are currently known to occur in at least four streams or stream complexes (*i.e.*, local populations). These local populations include Squaw Creek, Third Fork Squaw Creek, Rammage Meadows, and Renwyck Creek.

Bull trout spawning and rearing habitat occurs only in the upper watersheds.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Major Threats: (Primary Threats Identified by technical partners)

- Connectivity Impairment
- Nonnative fish
- Livestock Grazing

Summary for: North Fork Payette River Core Area

Geographic Description

The North Fork Payette River core area is located in Valley County. The core area is approximately 159,900 hectares (395,150 acres) and is isolated upstream of Cascade Lake and a dam in the lower Gold Fork River. The U.S. Forest Service manages 47 percent of the watershed while private lands accounts for 38 percent.

Current Distribution and Abundance

Bull trout are currently known in only one stream, the Gold Fork River, in this core area.

Bull trout spawning and rearing habitat occurs only in the upper watersheds and populations appear to only be resident fish.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Threats:

- Passage barriers
- Connectivity impairment
- Small Populations

Summary for: Middle Fork Payette River Core Area

Geographic Description

The Middle Fork Payette River core area is located in both Boise and Valley counties. The South Fork Payette eventually becomes the Payette River from its' confluence with the North Fork Payette River. The core area is approximately 88,400 hectares (218,500 acres) and is predominately Federal Lands. The Forest Service manages 95 percent of the watershed.

Current Distribution and Abundance

Bull trout are currently known in only three streams or stream complexes (*i.e.*, local populations). These local populations include Upper Middle Fork Payette River (including Stoney Meadow Creek), Sixteen-to-one Creek, and Bull Creek.

Limited fluvial life history expression has been documented in this core area.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Threats: (Primary Threats Identified by technical partners)

- Nonnative fish

Summary for: Deadwood River Core Area

Geographic Description

The Deadwood River core area occurs is located in Valley County. The Deadwood River drainage eventually joins the Upper South Fork Payette River. Deadwood Dam created Deadwood Reservoir and forms an impassible barrier to fish movement. Bull trout in the upper Deadwood River and Deadwood Reservoir are isolated from fish in the lower Deadwood River and the South Fork Payette River watersheds. The core area is approximately 28,400 hectares (70,200 acres). The U.S. Forest Service manages 92 percent of the watershed.

Current Distribution and Abundance

Bull trout are currently known to in at least six streams or stream complexes (*i.e.*, local populations). These local populations include Trail Creek, North Fork Beaver Creek, South Fork Beaver Creek, Wildbuck Creek, Upper Deadwood River, and Deer Creek.

Limited fluvial life history expression has been documented in this core area.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Threats: (Primary Threats identified by technical partners)

- Connectivity Impairment
- Water management

Summary for: Upper South Fork Payette River Core Area

Geographic Description

The Upper South Fork Payette River core area is located in both Boise and Valley counties. The South Fork Payette River (SFPR) eventually becomes the Payette River from its' confluence with the North Fork Payette River. The core area is approximately 173,700 hectares (429,200 acres) and is predominately Federal Lands. The U.S. Forest Service manages 95 percent of the watershed while private lands accounts for 1 percent.

Current Distribution and Abundance

Bull trout are currently known in only 11 streams or stream complexes (*i.e.*, local populations). These local populations include Scott Creek, Warm Springs Creek (Deadwood tributary), Clear Creek, Eightmile Creek, Tenmile Creek, Chapman Creek, Warm Spring-Gates Creek (SFPR tributary), Canyon Creek, Wapiti Creek, Trail Creek, and Baron Creek.

Limited fluvial life history expression has been documented in this core area.

The trend information and total abundance for local populations in most of this core area are unknown at this time.

Technical partners also discussed both climate change and urban development. The Service considered these threats but determined that these threats are speculative in nature. The Service focused on threats that were known to occur or would likely occur. The Service elected to remove climate change as a threat. The best conservation action to potentially address climate change is improving or promoting connectivity between local populations and within core areas. The effects of climate change are also difficult to determine, We acknowledge that climate change will have effects but are uncertain when those impacts would occur.

Threats:

- Degraded Habitat
- Nonnative fish

Summary for: North Fork Malheur River Core Area

Geographic Description

This core area is located in eastern Oregon, Grant, Baker, Malheur and Harney counties. It includes the North Fork River from the headwaters and tributaries, downstream to and including Beulah Reservoir. The North Fork Malheur River is the most important of the tributaries in terms of fish habitat and bull trout abundance.

Current Distribution and Abundance

The five bull trout populations in this core area include: 1) Elk Creek, 2) Little Crane Creek, 3) Swamp Creek, 4) Sheep Creek, and 5) Horseshoe Creek. The North Fork Malheur River subpopulation was isolated by Agency Dam in 1934 (Buchanan and Gregory 1997). Buchanan and Gregory (1997) classified bull trout in the North Fork Malheur River as “of special concern”, which falls between a “low” and “moderate” risk level.

The entire occupied area of the North Fork Malheur River is essential to the conservation of bull trout in the Upper Snake Recovery Unit because it is the most western core area (along with the Upper Malheur River) in this recovery unit and is isolated from the other core areas in this unit. The five populations in this core area are spread over an isolated, large geographical area with multiple age classes, containing both resident and migratory (fluvial) fish. Bull trout were known to exist in the North Fork Malheur River watershed prior to 1992. Distribution in the North Fork Malheur River above Agency Dam has remained unchanged since the species was first documented there (Buchanan and Gregory 1997). In August 2010, two charr that looked like bull trout/brook trout hybrids were identified, through photos, in the Little Malheur River of the North Fork River.

Threats:

- brook trout
- passage barriers
- impaired stream habitat conditions
- high stream temperatures

Summary for: Upper Malheur River Core Area

Geographic Description

This core area is located in eastern Oregon in Grant and Harney Counties and includes the mainstem Malheur River from the headwaters and tributaries, downstream to the town of Drewsey.

Current Distribution and Abundance

The three local bull trout populations in this core area include: 1) Lake Creek, 2) Meadow Fork Creek, and 3) Big Creek. The Upper Malheur River subpopulation was isolated by Warm Springs Dam in 1919 (Buchanan and Gregory 1997). Buchanan and Gregory (1997) classified bull trout in the Upper Malheur River as “high risk” of extinction.

The entire occupied area of the Upper Malheur River Core area is essential to the conservation of bull trout in the Upper Snake Recovery Unit because it is the most western core area (along with the North Fork) in the Upper Snake Recovery Unit and is isolated from the other core areas in this recovery unit. The three populations in this core area are spread over a large geographical area with multiple age classes, containing both resident and fluvial fish. Recent information indicates that there is a high proportion of brook trout in the Upper Malheur River, resulting in impacts through hybridization and competition for resources. Brook trout have displaced bull trout from several historic tributaries (*i.e.*, Summit, Bosonberg, McCoy and Corral Basin creeks) and affect over 60 percent of the bull trout population. An estimate of adult abundance for the Upper Malheur River local population is not available because of the inability to distinguish between bull trout and brook trout redds when not occupied.

Threats:

- brook trout
- passage barriers
- impaired stream habitat conditions
- elevated stream temperatures

Summary for: Little Lost River Core Area

Geographic Description

The Little Lost River core area is located in Lemhi, Butte, and Custer Counties. The Little Lost River lies in a closed basin within the upper Snake River basin and encompasses an area of 252,003 hectares (622,440 acres).

Land ownership in the Little Lost River basin is mixed. The Bureau of Land Management (43 percent) and the U.S. Forest Service (43 percent) manage the majority of lands within the recovery unit (LLRITAT 1998). Privately owned lands make up about 9 percent of the total land in the basin. The Idaho Department of Lands manages small land parcels interspersed within lands administered by the Bureau of Land Management.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 10 streams or stream complexes (*i.e.*, local populations). These local populations include Badger Creek (including Bunting Canyon Creek), Williams Creek, Wet Creek (including Big Creek), Warm Creek, Squaw Creek (tributary to Sawmill Creek), Mill Creek, Iron Creek (including Hawley and Jackson creeks), Timber Creek (including Camp, Redrock and Slide creeks), Smithie Fork Creek, Upper Little Lost River (Iron Creek confluence to headwaters excluding the Timber Creek), and Smithie Fork Creek watersheds).

Abundance of bull trout (expressed as density, or the number of individuals per kilometer of stream) has declined in some areas of the Little Lost River and its tributaries. Both resident and migratory (fluvial) bull trout exist in the Little Lost River core area. Numerous connectivity projects have occurred in the main stem of the Little Lost River. Bull trout in the Little Lost River below Iron Creek road are fluvial and migrate to headwater streams to spawn. The primary spawning areas for fluvial bull trout appear to be tributary streams in Sawmill Canyon.

The trend information from IDFG in 2014 indicates that the core area is stable.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

- Connectivity Impairment
- Livestock Grazing

Summary for: Jarbidge River Core Area

Area Description

The Jarbidge River Core area is located in Elko County, Nevada and Owyhee County, Idaho. The Jarbidge River core area includes the entire Jarbidge River drainage and the portion of the Bruneau River from the confluence of the Jarbidge River to Hot Springs Idaho (Buckaroo Diversion). The core area is approximately 854,700 hectares (8,547 square kilometers) (2,112,000 acres) (3,300 square miles). Elevations range from 792 meters (2,600 feet) Bruneau River at Hot Springs, Idaho to over 3,306 meters (10,839 feet) (Matterhorn Peak) at the headwaters. Major tributaries within the core area include East and West Fork Jarbidge River, Dave Creek, Jack Creek, Pine Creek and Slide Creek.

Land managers within the core area include the Humboldt National Forest, Bureau of Land Management, State of Idaho and some small parcels of private lands. The Humboldt National Forest manages most of the watershed with the majority in the Jarbidge Wilderness. Nevada Department of Wildlife manages fisheries resources within the majority of this core area.

Current Distribution and Abundance

Bull trout are currently known to use spawning and rearing habitat in at least 6 streams or stream complexes (*i.e.*, local populations). These local populations include Dave Creek, East Fork Jarbidge River, Jack Creek, Pine Creek, Slide Creek and West Fork Jarbidge River. A tracking study documented bull trout population connectivity between many of the local populations, in particular between West Fork Jarbidge River and Pine Creek. Movement between the East and West Fork Jarbidge River was also documented. Much of the movement was in the autumn concurrent with decreasing water temperatures. The majority of bull trout that emigrated were age-2 or older fish with increased movement with age and evidence of substantial amount of fluvial life history.

Trend information and total abundance for local populations in most of this core area are unknown at this time.

Threats: (No Primary Threats – based on technical partners and IDFG 2014 trend data)

Upland/Riparian Management

Summary for: Weiser River Core Area

Geographic Description

The Weiser River core area occurs in both Adams and Washington counties. The drainage joins the Snake River as part of the Brownlee Reservoir. The core area is approximately 245,500 hectares (606,700 acres). The Forest Service manages 44 percent of the watershed while private lands accounts for 40 percent.

Current Distribution and Abundance

Bull trout are currently known to in at least 5 streams or stream complexes (*i.e.*, local populations). These local populations include Upper Hornet Creek, East Fork Weiser River, Upper Little Weiser River, Anderson Creek, and Sheep Creek

Bull trout spawning and rearing habitat occurs only in the upper watersheds on Federal and State of Idaho lands. There is no connectivity between the current three stream complexes (Hornet Creek, Anderson Creek, and East Fork Weiser River.

IDFG indicated in 2014 that trend was increasing.

Threats: (Primary Threat – none)

- Habitat Degradation – water quality