

## Aldous Browse Survey, 1992-2006

In 1992 - 20 plots were sampled.  
In 1993 - 20 plots were sampled.  
In 1994 - 15 plots were sampled.  
In 1995 - 9 plots were sampled.  
In 1996 - **NO plots were sampled (lack of personnel and time)**  
In 1997 - 19 plots were sampled.  
In 1998 - 14 plots were sampled.(Other: 5 burned, 1 flooded)  
In 1999 - 17 plots were sampled.(Other 3: burned)  
In 2000 - 18 plots were sampled.(Other 2: burned)  
In 2001 - 18 plots were sampled.(1 burned, 1 flooded)  
In 2002 - 17 plots were sampled.(Other 3: burned)  
In 2003 - 18 plots were sampled. (Other 2: burned)  
In 2004 - 13 plots were sampled. (Other 7: burned)  
In 2005 - 16 plots were sampled. (Other 4: burned)  
In 2006 - 12 plots were sampled. (Other 8: burned)  
In 2007 - 15 plots were sampled. (Other 5: burned)

### **HONEYSUCKLE BROWSE**

1992 - 44% of available honeysuckle was browsed.  
1993 - 49% of available honeysuckle was browsed.  
1994 - 49% of available honeysuckle was browsed.  
1995 - 39% of available honeysuckle was browsed.  
1996 - no record  
1997 - 44% of available honeysuckle was browsed.  
1998 - 36% of available honeysuckle was browsed.  
1999 - 37% of available honeysuckle was browsed.  
2000 - 28% of available honeysuckle was browsed.  
2001 - 50% of available honeysuckle was browsed.  
2002 - 17% of available honeysuckle was browsed.  
2003 - 45% of available honeysuckle was browsed.  
2004 - 67% of available honeysuckle was browsed.  
2005 - 70% of available honeysuckle was browsed.  
2006 - 37% of available honeysuckle was browsed.  
2007 - 18% of available honeysuckle was browsed.

### **GREENBRIER BROWSE**

1992 - 73% of available greenbrier was browsed.  
1993 - 72% of available greenbrier was browsed.  
1994 - 84% of available greenbrier was browsed.  
1995 - 79% of available greenbrier was browsed.  
1996 - no record  
1997 - 56% of available greenbrier was browsed.  
1998 - 93% of available greenbrier was browsed.  
1999 - 70% of available greenbrier was browsed.  
2000 - 71% of available greenbrier was browsed.  
2001 - 86% of available greenbrier was browsed.  
2002 - 37% of available greenbrier was browsed.  
2003 - 69% of the available greenbrier was browsed.  
2004 - 66% of the available greenbrier was browsed.  
2005 - 87% of the available greenbrier was browsed.  
2006 - 54% of the available greenbrier was browsed.  
2007 - 46% of the available greenbrier was browsed.

**Browse Comparison for Understory Vegetation, 1997-2006**  
**% of Plots that Species were browsed**  
**(where species were available)**

<b>PLANT SPECIES</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
Dogwood	18%	9%	1%	0%	-	17%	9%	0%	0%	1%	10%	6%	21%	12%	50%	24%
Hickory	50%	0%	0%	0%	-	0%	0%	0%	0%	0%	0%	0%	0%	23%	5%	1%
Black Cherry	87%	20%	1%	0%	-	0%	0%	0%	0%	16%	7%	1%	6%	54%	38%	15%
Sweetgum	64%	8%	0%	8%	-	0%	0%	0%	0%	1%	0%	2%	14%	4%	0%	0%
Vaccinium	71%	50%	3%	0%	-	0%	0%	0%	0%	0%	3%	13%	24%	22%	39%	36%
Sumac	100%	67%	35%	-	-	0%	0%	0%	25%	0%	0%	0%	0%	48%	20%	-
Blackberry	63%	50%	7%	0%	-	0%	25%	0%	8%	44%	3%	1%	62%	58%	60%	53%
Grape	33%	11%	<1%	0%	-	0%	17%	0%	0%	0%	2%	25%	7%	24%	29%	20%