

Table 1.— Preliminary estimates of passage by brood-year (BY) and run for unmarked juvenile Chinook salmon and steelhead trout captured by rotary-screw traps at Red Bluff Diversion Dam (RK391), Sacramento River, CA, for the dates listed below. Results include estimated passage, peak river discharge volume, water temperature, turbidity, and fork length (mm) range in parentheses. A dash (-) indicates that sampling was not conducted on that date.

Date	Discharge volume (cfs)	Water temperature (°C)	Water turbidity (NTU)	Estimated passage				
				BY04 Fall	BY05 Late-fall	BY04 Winter	BY04 Spring	BY05 Steelhead
4/23/05	6,050	15.0	1.6	7,477 (50 – 85)	0 (–)	0 (–)	1,467 (86 – 102)	155 (54 – 167)
4/24/05	6,470	14.2	2.1	10,103 (53 – 86)	0 (–)	88 (126 – 149)	1,377 (87 – 111)	291 (51 – 84)
4/25/05	7,570	14.1	2.0	33,785 (53 – 87)	0 (–)	0 (–)	3,084 (88 – 109)	496 (57 – 82)
4/26/05	7,530	14.7	1.5	49,281 (47 – 87)	0 (–)	0 (–)	4,546 (88 – 113)	677 (56 – 78)
4/27/05	7,400	14.6	1.9	51,073 (55 – 88)	0 (–)	0 (–)	3,678 (89 – 113)	937 (52 – 82)
4/28/05	7,330	14.6	2.7	31,407 (52 – 88)	117 (37 – 38)	0 (–)	2,178 (89 – 109)	589 (38 – 74)
4/29/05	7,420	14.8	2.0	35,451 (52 – 89)	0 (–)	0 (–)	1,690 (90 – 102)	581 (48 – 84)
4/30/05	7,420	15.3	1.8	431,578 (52 – 89)	162 (35)	0 (–)	2,282 (90 – 101)	191 (66 – 81)
5/1/05	7,640	-	-	-	-	-	-	-
5/2/05	7,530	-	-	-	-	-	-	-
5/3/05	7,690	15.7	1.7	94,402 (51 – 91)	0 (–)	0 (–)	692 (92)	605 (43 – 82)
5/4/05	7,730	14.9	2.0	77,418 (49 – 92)	0 (–)	170 (149)	1,182 (94 – 112)	927 (56 – 78)
5/5/05	8,480	-	-	-	-	-	-	-
5/6/05	12,200	14.1	61.3	437,921 (57 – 93)	887 (34)	0 (–)	1,654 (96)	764 (42 – 200)
<b>Biweekly total<sup>1</sup></b>				1,836,990	1,692	429	28,172	8,211
<b>Brood-year total</b>				10,807,182	12,376	3,275,024	414,120	16,040

<sup>1</sup> Biweekly totals may be greater than the sum of the daily estimates presented in this table if sampling was not conducted on each day of the biweekly period. A dash (-) denotes those dates. To estimate daily passage for days that were not sampled, we used a mean daily passage from the sample immediately preceding and following the un-sampled day. When consecutive days were not sampled, we calculated a mean daily passage for that period by noting the number of days not sampled and then calculating a mean daily passage using the same number of samples immediately preceding and following the un-sampled period (e.g., if three consecutive days were not sampled, we calculated a mean daily passage for each day using the three samples immediately preceding and following the un-sampled period).