

**U.S. FISH AND WILDLIFE SERVICE
SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM**

SCIENTIFIC NAME: *Hypomesus transpacificus*

COMMON NAME: Delta smelt

LEAD REGION: Region 8

DATE INFORMATION CURRENT AS OF: June 27, 2016

STATUS/ACTION

Species assessment – determined either we do not have sufficient information on threats or the information on the threats does not support a proposal to list the species and, therefore, it was not elevated to Candidate status

Listed species petitioned for uplisting for which we have made a warranted-but-precluded finding for uplisting (this is part of the annual resubmitted petition finding)

Candidate that received funding for a proposed listing determination; assessment not updated

New candidate

Continuing candidate

Listing priority number change

Former LPN: ____

New LPN: ____

Candidate removal: Former LPN: ____

A – Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

U – Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species.

F – Range is no longer a U.S. territory.

I – Insufficient information exists on biological vulnerability and threats to support listing.

M – Taxon mistakenly included in past notice of review.

N – Taxon does not meet the Act’s definition of “species.”

X – Taxon believed to be extinct.

Date when the species first became a Candidate (as currently defined):

Petition Information:

Non-petitioned

Petitioned; Date petition received: 3-8-2006

90-day substantial finding FR publication date: 7-10-2008

12-month warranted but precluded finding FR publication date: 4-7-2010

FOR PETITIONED CANDIDATE SPECIES:

- a. Is listing warranted (if yes, see summary of threats below)? Yes
- b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? Yes
- c. Why is listing precluded? Higher priority listing actions, including court-approved settlements, court-ordered and statutory deadlines for petition findings and listing determinations, emergency listing determinations, and responses to litigation, continue to preclude the proposed and final listing rules for this species. We continue to monitor populations and will change its status or implement an emergency listing if necessary. The “Progress on Revising the Lists” section of the current CNOR (<http://endangered.fws.gov/>) provides information on listing actions taken during the last 12 months.

ANIMAL/PLANT GROUP AND FAMILY: Fish, Osmeridae

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: Contra Costa, Sacramento, San Joaquin, Solano, Napa and Yolo Counties in the State of California.

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: Contra Costa, Sacramento, San Joaquin, Solano, Napa and Yolo Counties in the State of California.

LAND OWNERSHIP: This species occurs in open waters. There are no known land locked populations. The statutory Delta totals 738,000 acres including approximately 538,000 acres of agricultural land uses, 60,000 acres of open water, and 64,000 acres of urban land uses. The remainder of the region presently consists of open space and wildlife habitat.

LEAD REGION CONTACT: R8, Daniel Russell, 916-978-6191, daniel_russell@fws.gov

LEAD FIELD OFFICE CONTACT: Bay-Delta FWO, Colin Grant, 916-930-5651, colin_grant@fws.gov

In this document, we use several different terms to describe the various portions of the Delta. These terms are Bay, Delta, Bay-Delta, and San Francisco Estuary. We define these terms as follows. The “Delta” represents the legal delta encompassing all waters east of Chipps Island. The “Bay” encompasses all waters west of the Chipps Island where the legal delta ends. The “Bay-Delta” encompasses both the Bay and the Delta. The “San Francisco Estuary” encompasses all waters that have a measurable salinity within the Bay-Delta and changes seasonally with the shifting salinity zone.

BIOLOGICAL INFORMATION:

Species Description

Delta smelt are slender-bodied fish; generally about 60 to 70 millimeters (mm) (2 to 3 inches (in)) long (Figure 1), although they may reach lengths of up to 120 mm (4.7 in) (Moyle 2002, p. 227). Delta smelt are in the Osmeridae family (northern smelts) (Stanley *et al.* 1995, p. 390). Live fish are nearly translucent and have a steely blue sheen to their sides (Moyle 2002, p. 227). Delta smelt are also identifiable by their relatively large ratio of their eye diameter to head length. The eye can occupy approximately 25–30 percent of their head length (Moyle 2002, p. 227). Delta smelt have a small, translucent adipose fin located between the dorsal and caudal fins. Occasionally one chromatophore (a small dark spot) may be found between the mandibles, but most often there is none (Moyle 2002, p. 227).



Figure 1: Delta smelt

Taxonomy

We have carefully reviewed the available taxonomic information to reach the conclusion that the delta smelt (*Hypomesus transpacificus*) is a valid taxon. The delta smelt is one of six species currently recognized in the *Hypomesus* genus (Ilves and Taylor 2007, p. 8). Within the genus, delta smelt is most closely related to surf smelt (*H. pretiosus*), a species common along the western coast of North America (Ilves and Taylor 2007, p. 8). In contrast, delta smelt is a comparatively distant relation to the wakasagi (*H. nipponensis*), which was introduced into Central Valley reservoirs in 1959, and may be seasonally sympatric with delta smelt in the estuary (Trenham *et al.* 1998, p. 417). Delta smelt and Wakasagi hybrids as well as Delta smelt and longfin smelt hybrids have been observed in the Bay-Delta (California Department of Fish and Game (CDFG) 2001, p. 473). However, allozyme studies have demonstrated that wakasagi and delta smelt are genetically distinct and derived from different marine ancestors (Stanley *et al.* 1995, p. 394).

Habitat/Life History

Delta smelt are a euryhaline (tolerate a wide range of salinities) species (Moyle 2002, pp. 228–229). In captivity, some delta smelt can survive in seawater for extended periods (Komoroske *et al.* 2014, p. 6); however, in the wild they rarely occur in water with more than 10–14 salinity (about one-third seawater) (Bennett 2005, p. 11; Moyle 1992, p. 73). In the Practical Salinity Scale, salinity is defined as a pure ratio, and has no dimensions or units. Feyrer *et al.* (2007, p. 728) found that relative abundance of delta smelt was related to fall specific conductance (a surrogate for salinity) and water transparency. Delta smelt probably evolved within the naturally turbid (silt and particulate-laden) environment of the estuary's "low salinity zone" (LSZ) where the salinity ranges from approximately 0.5 to 6 (Kimmerer 1998, p.1; Moyle 2002, p. 228) and likely rely on certain levels of background turbidity at different life stages and for certain behaviors. Juvenile and sub adult delta smelt are most common within the LSZ at salinity of 1–2 (Bennett 2005, p. 10, Sommer *et al.* 2011a, p. 8).

Between December and March (Grimaldo *et al.* 2009, p.1263; Sommer *et al.* 2011, p. 12), delta smelt begin their migration back into freshwater areas where spawning is thought to occur. This migration is thought to be triggered by the first seasonal high outflow event in the Delta (Grimaldo 2009, p. 1259) although the mechanisms for this migration are still not clear. (Bennett and Burae 2014, p. 9) did find that delta smelt change their movements in response to tides, but could not distinguish the relative importance of turbidity versus changing tidal direction as cues for moving laterally or for the spawning migration. In captivity, some delta smelt can survive to age two (Bennett 2005, p. 16, Figure 9). In the wild, most delta smelt die after spawning at age one, but a small contingent of adults may survive to spawn in their second year (Bennett 2005, p. 22). Fecundity is correlated with size. In captivity, age one females spawn between 1,000 to 4,000 eggs while age two females can spawn up to 12,000 eggs (Bennett 2005, p. 15). It was once believed that delta smelt were strictly semelparous (having a single reproductive episode before death) (Moyle 2002, p.68); however, new evidence suggests that adult females can produce multiple egg batches per spawning period if water temperatures stay cool long enough and the fish get enough food to support the development of multiple clutches of eggs (Lindberg 2015, pers. comm., Fujimura 2015, pers. comm.).

Spawning is believed to occur from late January through late June or early July with most spawning occurring during April through mid-May (Bennett 2005, p. 13). Spawning likely occurs mainly at night with several males attending females that broadcast eggs onto bottom substrate (Bennett 2005, p. 13). Although preferred spawning substrate is unknown, spawning habits of close relative surf smelt, as well as preliminary studies, suggest that sandy substrate may be preferred (Bennett 2005, p. 17; Sommer *et al.* 2013, p. 13). Hatching success decreases when temperatures exceed 20⁰C (68⁰F); surveys indicate that the window of best spawning success ranges from about 14–18⁰C (57–64⁰F) (Bennett 2005, p. 17). In laboratory conditions, eggs typically hatch after 9 to 14 days and larvae begin feeding 5 to 6 days later (Mager *et al.* 2004, p. 172, Table 1). Larvae are generally most abundant in the Delta from mid-April through May (Bennett 2005, p. 13). Some delta smelt, have been observed spending their entire life cycle in freshwater within the Cache Slough region including Liberty Island (Sommer *et al.* 2011, p. 9) showing that an alternative life history strategy is possible if habitat parameters are favorable for delta smelt.

After several weeks of development, larval surveys indicate that many larvae move downstream until they reach nursery habitat in the LSZ, in part to reach cooler waters (Kimmerer 1998, p. 1; Moyle 2002, p. 228; Dege and Brown 2004, pp. 57–58). Juvenile smelt rear and grow in the LSZ and adjacent fresher water habitats for several months, where they are found in open waters (free of vegetation) (Dege and Brown 2004, pp. 56–58). By the summer, delta smelt are capable of controlling their distribution to maintain an association with suitable habitat conditions (Kimmerer 2008, 18). Growth is rapid; juvenile fish reach 40–50 mm (1.6–2 in) by early August (Erkkila *et al.* 1950; Ganssle 1966, p.78; Radtke 1966, p.118). By this time, young-of-the-year fish dominate trawl catches of delta smelt, and adults become rare. Delta smelt reach 55–70 mm (2.2 – 2.8 in) standard length in 7–9 months (Moyle 2002, p. 228). The abrupt change from a single-age, adult cohort during spawning in spring to a population dominated by juveniles in summer strongly suggests that most adults die shortly after they spawn (Radtke 1966, p. 120). Growth during September to November slows down considerably with a total length increase of only 3–9 mm (0.1 – 0.4 in over these three months (Moyle 2002, p. 228). During this time period, less food is being produced today than was produced historically at the same temperatures. Delta smelt are now 5-10 mm smaller at a given age than they were historically (Sweetnam 1999, p. 25).

Delta smelt feed primarily on small planktonic (free-floating) crustaceans, and occasionally on insect larvae (Moyle 2002, p. 228). Historically, the main prey of delta smelt was the copepod *Eurytemora affinis* and the mysid shrimp *Neomysis mercedis* (Moyle *et al.* 1992, p. 70, Table 1). The copepod *Pseudodiaptomus forbesi* has replaced *E. affinis* as a major prey source of delta smelt since its introduction into the San Francisco Bay-Delta (Baxter *et al.* 2008, p. 22). Larval smelt primarily consume the two copepods, *Eurytemora affinis* and *Pseudodiaptomus forbesi*, and in an older study, a third prey group, freshwater species of the family Cyclopidae, was also common (Nobriga 2002, p. 156; Slater and Baxter 2014, p. 8). The diversity of prey eaten by delta smelt increases as they grow; adult diets are dominated by adult copepods and somewhat larger crustaceans like amphipods, though many other invertebrates and larval fishes have been observed occasionally in stomach contents (Lott 1998, p. 19).

Water temperature also affects delta smelt distribution and this effect varies by life stage. Swanson *et al.* (2000, p. 386) reported a minimum temperature tolerance for juvenile delta smelt of 7.5°C (45.5°F) (p. 386). The approximate maximum temperature tolerances in captivity by life stage are as follows: larvae: 30°C (86°F), late larvae: 29°C (84.2°F), juvenile: 29°C (84.2°F), adult: 28.2°C (82.8°F), post spawned adult: 27.1°C (Komoroske *et al.* 2014, p. 7). Tolerance limits are typically measured at the point that delta smelt lose their equilibrium or balance. In the wild, delta smelt are seldom collected from water that approach their physiological tolerance limits (Nobriga *et al.* 2008, p. 7, Fig 4; Komoroske *et al.* 2014, p. 9, Fig 3), probably because warm water increases energetic demands (Rose *et al.* 2013a, p. 1245), which has been shown to cause behavioral impairment and lowered competitive ability in other fishes. However, at the juvenile stage, a very small fraction of the delta smelt have been collected from waters that closely represent their 29°C (84.2°F) limit tolerated in captivity (Komoroske *et al.* 2014, p. 9). This is important because these fish that tolerate higher temperatures will be more strongly selected for as waters in the delta warm due to climate change.

Currently available information indicates that delta smelt habitat is most suitable for the fish when low-salinity water is near 20°C (68°F), highly turbid, oxygen saturated, low in contaminants, and supports high densities of calanoid copepods and mysid shrimp (e.g., Moyle 2002, p. 228; Nobriga 2002, pp. 160–163, Feyrer *et al.* 2007 pp. 728–732). Almost every component listed above has been degraded over time (see five factor analysis).

POPULATION STATUS

Historical Range/Distribution

Delta smelt are endemic to the San Francisco Bay-Delta in California. Expansions and contractions to the range are discussed below.

Current Range/Distribution

Delta smelt are endemic to the upper San Francisco Bay-Delta estuary (Figure 2). The reported range of the Delta smelt extends from Berkeley in the San Francisco Bay to the City of Napa on the Napa River (Figure 8), throughout Suisun Bay and the Delta, and along the axis of the Sacramento River to Knight's Landing and along the axis of the San Joaquin River to the City of Lathrop (Merz *et al.* 2011, p. 181-182; Vincik and Julienne 2012, p. 173). At all life stages, Delta smelt distribution is strongly influenced by the position of the LSZ (Moyle *et al.* 1992, p. 72; Dege and Brown 2004, p. 56; Sommer *et al.* 2011, p. 7; Sommer and Mejia 2013, p. 8), although delta smelt commonly use tidal habitats where salinity is lower than 0.5. Delta smelt nearly always occupy a few locations throughout the year in both wet and dry years, including Honker Bay, Montezuma Slough, the Sacramento River between Sherman Lake and Rio Vista, and parts of the Liberty Island/Sacramento Deep Water Shipping Channel region (Sweetnam 1999, p. 24; Bennett 2005, p. 11; Sommer and Mejia 2013, p. 6). This portion of the habitat is located where salinity stays within a suitable range and also includes the last remaining semi-shallow and highly turbid waters (Feyrer *et al.* 2007, pp. 123–127). Delta smelt can seasonally occupy the central and eastern parts of the Delta, which can become seasonally turbid (CDFW 2016, SKT and 20 mm. surveys). During drier years, when total outflow is less than about 10,000 cfs, delta

smelt tend to aggregate and spawn in the western Delta and Cache Slough Complex where the water remains turbid and of low salinity; it is unknown how much spawning is distributed among these two regions under these occasional low flow circumstances. When outflow is high, fresh water extends farther west and delta smelt temporarily extend their range sometimes occupying San Pablo Bay, the Carquinez Strait region and the Napa River (Sweetnam 1999, p. 24).

Although a few delta smelt historically occupied the southern Delta year round (Erkkila et al. 1950, entire; Radke 1966, p. 119), presently individuals that have not been entrained typically vacate this region from July through November or December each year, suggesting a seasonal contraction of the species' range (Sommer *et al.* 2011, p. 9). During the years 1970 through 1978, delta smelt catches in the Tow Net Survey declined rapidly to zero in the Central and South Delta and have remained near zero since (Nobriga et al. 2008, p. 9). A similar shift in Fall Midwater Trawl catches occurred after 1981 (Arthur *et al.* 1996, p. 481). This portion of the Delta has had a long-term increase in water clarity during July through December, which, along with comparatively warm summer water temperatures, extensive beds of submerged aquatic vegetation, localized dissolved oxygen sags, and high risk of entrainment, has greatly reduced its suitability as delta smelt habitat (Arthur *et al.* 1996, p. 468; Feyrer *et al.* 2007, p. 730; Nobriga *et al.* 2008, p. 8).

In the north Delta, there may have been an expansion of the known seasonal range. Alternatively, this region may have always been occupied, but not routinely monitored because historically it was not a major rearing habitat for young striped bass for which surveys in the region are conducted. Routine sampling for delta smelt in the Cache Slough Complex during the summer was not initiated until 2008. Some delta smelt remain in the freshwater Cache Slough region year-round (Sommer *et al.* 2011, p. 10). This region, which includes the recently flooded Liberty Island, may be attractive due to the high diversity of habitats including tidal marshes, multiple channel sizes, and dead end sloughs (Whitley and Bollens 2014, p. 671), but the water in Liberty Island and adjacent habitats is also fairly turbid (Nobriga et al. 2005, p. 780). Other benefits of the Liberty Island/Sacramento Deep Water Shipping Channel region include channels and shoals with slow moving water that result in long residence times that support growth of phytoplankton and zooplankton (Lehman *et al.* 2010b, pp. 369–370; Nelson *et al.* 2011, p. 18).

The locations in the Delta where newly hatched larvae are present most likely indicate spawning locations. Sampling of larval delta smelt in the Bay-Delta in 1989 and 1990 suggested that spawning occurred in the Sacramento River; in Georgiana, Prospect, Beaver, Hog, and Sycamore sloughs; in the San Joaquin River adjacent to Bradford Island and Fisherman's Cut; and possibly other areas (Wang 1991, p. 11). However, in recent years, the densest concentrations of both spawners and larvae have been recorded in the Cache Slough/Sacramento Deepwater Ship Channel complex in the North Delta. Some delta smelt spawning occurs in Napa River, Suisun Bay and Suisun Marsh during wetter years (Sweetnam 1999, p. 24; Wang 1991, pp. 11–12; Hobbs *et al.* 2007, p. 522). Early stage larval delta smelt have also been recorded in Montezuma Slough near Suisun Bay (Wang 1986, p. 11). Fish spawned in the Napa River can get back to the LSZ and survive to adulthood (Hobbs *et al.* 2007, p. 523).




Figure 2: Delta Map

Population Indices

Within the Bay-Delta, delta smelt are consistently collected in the monitoring surveys that have been conducted by California Department of Fish and Wildlife, dating back to 1959. Most of the ongoing studies are currently conducted under the auspices of the Interagency Ecological Program (IEP), an entity made up of State, Federal- and non-government agencies that work collaboratively to oversee data collection and scientific analysis in the Bay-Delta. Several of the IEP's field investigations provide annual delta smelt distribution and relative abundance information, including the Spring Kodiak Trawl (SKT), the Smelt Larva Survey (SLS), the 20mm survey (20mm), the summer Towntnet Survey (TNS), and the Fall Midwater Trawl (FMWT). These surveys are presented as indices. The index numbers are not numbers of fish in the Bay-Delta, they are a relative index. The surveys tell us whether years are better or worse in relation to one another rather than estimating the population size of delta smelt.

The SKT is a surface trawl targeting spawning adult delta smelt at up to 40 stations from the Napa River landward throughout the Delta from January to May. The spring SKT has been conducted every year since 2003. The SLS and 20 mm Survey provide information on larvae and post larvae smelt. The 20-mm Survey has been conducted every year since 1995. This survey targets late-stage delta smelt larvae. Most sampling has occurred April-June. The TNS has been conducted nearly every year since 1959. This survey targets 38-mm striped bass, but collects similar-sized juvenile delta smelt. Most sampling has occurred June-August. The FMWT has been conducted nearly every year since 1967. This survey also targets age-0 striped bass, but collects delta smelt > 40 mm in length. The FMWT samples monthly, September-December. The relative abundance index data and maps of the sampling stations used in these surveys are available at CDFW - <http://www.dfg.ca.gov/delta/>.

FMWT-derived data are generally accepted as providing a reasonable basis for detecting and roughly scaling inter-annual trends in the relative abundance of delta smelt. The FMWT-derived indices have ranged from a low of 77 in 2015 to 1,673 in 1970 (Figure 3). TNS-derived indices have ranged from a low of 0.0 in 2015 to a high of 62.5 in 1978 (Figure 4), 20-mm indices have ranged from a low of 0.3 in 2015 to a high of 39.7 in 1999 (Figure 5), and SKT indices have ranged from a low of 13.8 in 2015 to a high of 147 in 2012 (Figure 6).

From 1969–1981, the mean delta smelt FMWT and TNS indices were 894 and 22.5, respectively. From 1982–1992, the mean delta smelt FMWT and TNS indices dropped to 272 and 3.2, respectively. Both indices suggest the delta smelt population declined abruptly in the early 1980s (Moyle *et al.* 1992, pp. 71–72). The population rebounded somewhat in the mid–1990s (Sweetnam 1999, p. 24); the mean FMWT and TNS indices were 529 and 7.1, respectively, during the 1993–2002 period. From 2003–2012, the FMWT and TNS index averaged 83.3 and 1.04, which is the lowest decade on record. Delta smelt numbers have trended precipitously downward since the early 2000s (Thomson *et al.* 2010, p. 1439, Figure 3). In the wet water year of 2011, the FMWT index for delta smelt increased to 343, which is the highest index recorded since 2001. It immediately declined again in 2012 to 42 and continued to decline in 2013 and 2014 when the index was 18 and 9, respectively. A new all-time low was reached in 2015 with an index of 7. Eleven of the last twelve years have seen FMWT indexes that have been the lowest ever recorded. Although, the 2016 SKT surveys have not been completed, the

first four months of 2016 have shown the lowest SKT catch on record (Figure 7). 2015-2016 results from all four of the surveys analyzed in this review have been the lowest ever recorded for the delta smelt.

Correlations exist between the 20-mm Survey and TNS indices as well as between the FMWT survey and the SKT indices. Though it is limited to data from the last 10 years, the relationship between the FMWT index of juveniles at the end of their first calendar year of life and the SKT index of spawning adults the following spring is strong and essentially linear (Nobriga 2015, no pagination). This indicates that delta smelt year-class strength is set by the end of the first calendar year of life and there is no obvious indication of a noteworthy source of overwinter mortality that decouples these indices. In contrast, the relationship between the SKTS index of adult delta smelt abundance and the 20-mm Survey index of their progeny is clearly nonlinear (Nobriga 2015, no pagination), implying there may be a carrying capacity for larval production (Nobriga 2013, p.1566).




Figure 3: Delta smelt abundance (total across year-classes) as indexed by the Fall Mid-Water Trawl of the Bay-Delta, 1967–Present. No surveys were conducted in 1974 or 1979. Eleven out of the last twelve years’ indices have been the lowest on record.




Figure 4: Summer Townnet Survey index 1959–Present: The TNS shows a decline in the population over time remaining below an index of one for ten of the last eleven years. No surveys were conducted from 1966–1968. Values of less than 3 have been labeled for better accuracy.




Figure 5: 20-mm Survey index: 1995-Present.




Figure 6: Spring Kodiak Trawl Index: 2003-Present.




Figure 7: Spring Kodiak Trawl Index: 2003-Present. Showing first four months of the year. The fifth month has not yet been tallied for 2016.

Population Estimates

There are three independent population estimates of delta smelt abundance that we are aware of: Kimmerer (2008, p. 21) used SKTS data to estimate that the abundances for 2002-2006 ranged from 100,000 to 1.5 million, while Rose et al. 2013 (p. 1250) used a combination of FMWT and SKTS data to estimate that abundances between 1995 and 2005 ranged from 500,000 to 9 million individuals. The FMWT index has declined precipitously over the last decade, indicating current abundance is likely much lower than historic abundance.

A new delta smelt abundance estimation procedure based on SKTS data was recently completed by the Service. Estimates of historical delta smelt abundances for the months of January and February based on SKTS data are provided below. The estimates (in 1000s of fish) are listed below, with 95% confidence interval estimates in parentheses. For example, the 2016 population estimate is 13,000 individuals with 95% confidence intervals ranging between 6,000 and 28,000 individuals (Newman 2016, unpublished data).

<u>Year</u>	<u>Abundance Estimates</u>
2002	597 (405,879)
2003	519 (238,1130)
2004	527 (297,934)
2005	385 (248,598)
2006	151 (105,216)

2007	235 (126,439)
2008	262 (120,576)
2009	295 (126,691)
2010	134 (83,217)
2011	234 (87,631)
2012	623 (346,1120)
2013	171 (94,312)
2014	167 (91,308)
2015	112 (54,233)
2016	13 (6,28)

Note that the 2002-2006 abundances, with confidence intervals ranging from 105,000 to 1.13 million, fall within Kimmerer’s range of 100,000 to 1.5 million. Newman, et al added the comment: “We emphasize that the 2016 SKT abundance estimate is based on only 13 fish and we have concerns about false zeros. In particular, the intensive sampling carried out in the lower San Joaquin River at Jersey Point and Prisoner’s Point (sampling 2-3 times/week and roughly 15 tows per location) as part of an Early Warning System program caught many fish while the regular SKT survey failed to catch any fish at those two locations during January and February 2016.” This considerable uncertainty is reflected in the confidence interval that ranges from 6,000 to 28,000.

These abundance estimates are an interim product of the on-going delta smelt life cycle model (DSLCLM) development work. The current version of the DSLCLM contains two life stages, adults (early February) and juveniles (June). Recent work has focused on the reproductive cycle, in particular identifying and quantifying the effects of potentially manipulable factors influencing recruitment success, e.g., the average number of juveniles produced per adult. Initial results suggest a positive association between spring outflows and recruitment success. Subsequent work is focused on (1) identifying and quantifying factors affecting the transition from juveniles to adults, namely survival rates which includes entrainment effects, (2) adding additional life stages (larvae in April and sub-adults in October-November), and (3) nesting the entire DSLCLM in a state-space model framework which simultaneously accounts for environmental variation and sampling noise.

THREATS: FIVE FACTOR ANALYSIS

Introduction of Threats

Section 4 of the Act (16 U.S.C. 1533) and implementing regulations (50 CFR part 424) set forth procedures for adding species to, removing species from, or reclassifying species on the Federal Lists of Endangered and Threatened Wildlife and Plants. Under section 4(a)(1) of the Act, a species may be determined to be endangered or threatened based on any of the following five factors:

- (A) The present or threatened destruction, modification, or curtailment of its habitat or range;
- (B) Overutilization for commercial, recreational, scientific, or educational purposes;

- (C) Disease or predation;
- (D) The inadequacy of existing regulatory mechanisms; or
- (E) Other natural or manmade factors affecting its continued existence.

In making these findings, information pertaining to each species in relation to the five factors provided in section 4(a)(1) of the Act is discussed below. The threat is significant if it drives or contributes to the risk of extinction of the species such that the species warrants listing as endangered or threatened as those terms are defined by the Act.

The primary known threats cited in the 2010 delta smelt uplisting document are: entrainment by State and Federal water export facilities (Factor E); summer and fall increases in salinity due to reductions in freshwater flow and summer and fall increases in water clarity (Factor A), and effects from introduced species, primarily the overbite clam and *Egeria densa* (Factor E). Additional threats included predation (Factor C), entrainment into power plants (Factor E), contaminants (Factor E), and small population size (Factor E). Since the 2010 warranted 12-month finding, we have identified climate change as a threat in the 2012 Candidate Notice of Review. Climate change was not analyzed in the 2010 12-month finding document. Since the 2010 up-listing, one of the two power plants within the range of the delta smelt using water for cooling has shut down and power plants are no longer thought to be a threat to the population as a whole. We have identified a number of existing regulatory mechanisms that provide protective measures that affect the stressors acting on the Delta smelt. Despite these existing regulatory mechanisms and other conservations efforts, the stressors continue to act on the species such that it is warranted for uplisting under the Act.

A. The present or threatened destruction, modification, or curtailment of its habitat or range.

Increased Salinity due to Reduced Freshwater Flow

As California's population has grown, demands for reliable water supplies and flood protection have increased. In response, local, state and federal agencies have built dams and canals, and captured water in reservoirs, to increase capacity for water storage and conveyance, resulting in one of the largest manmade water systems in the world (Nichols et al. 1986, p. 569). Operation of this system has altered the seasonal pattern of freshwater flows in the Bay-Delta. Storage in the upper watershed of peak runoff and release of the captured water for irrigation and urban needs during subsequent low flow periods result in a broader, flatter hydrograph with less seasonal variability in freshwater flows into the estuary (Kimmerer 2004, p. 15).

Two of the key hydrodynamic variables used in the resource management of the Bay-Delta are Delta inflow (from the rivers into the Delta) and Delta outflow (from the Delta into the bays). Due to high flow events, these variables are closely correlated, but they are not interchangeable. In the Bay-Delta, the location where salinity is equal to 2 is called X2. X2 is indexed as distance in kilometers from the Golden Gate Bridge. X2 is important to delta smelt because it has been shown to affect a variety of factors that contribute to delta smelt survival, making it a useful indicator of habitat conditions (Jassby *et al.* 1995, p. 282; Dege and Brown 2004, pp. 56–58). Delta outflow is the variable that most directly affects the location of X2 (Jassby *et al.* 1995, p. 284). The location of X2 is influenced by precipitation in the watershed (i.e., wetter or drier

seasonal weather patterns) and by water operations, both upstream at the dams and diversions and in the Delta at water export facilities (Jassby *et al.* 1995, entire; Kimmerer 2004, p. 18).

In addition to the system of dams and canals built throughout the Sacramento and San Joaquin River basins, the Bay-Delta is unique in having the largest water diversion system on the west coast. The State Water Project (SWP) and Central Valley Project (CVP) each operate two water export facilities in the Delta (Kimmerer and Nobriga 2008, p. 2). Project operation is dependent upon upstream water supply and export area demands, both of which are strongly affected by the interannual variability in Delta hydrology caused by variability in precipitation. From 1956 to the 1990s, water exports increased from approximately 5% of the Delta inflow to approximately 30% of the Delta inflow (Cloern and Jassby 2012, p. 7). In total, an estimated 39% of the estuary's unimpaired flow is consumed upstream or diverted from the estuary (Cloern and Jassby 2012, p. 8). Annual inflow from the watershed to the Delta is strongly correlated to unimpaired flow (runoff that would hypothetically occur if upstream dams and diversions were not in existence), mainly due to the effects of high-flow events (Kimmerer 2004, p. 15). Water operations are regulated in part by the California State Water Resources Control Board (SWRCB) according to the Water Quality Control Plan (WQCP) (SWRCB 2000, entire). The WQCP limits Delta water exports in relation to Delta inflow (the Export/Inflow, or E/I ratio). Operations are also regulated by both the Service's and NMFS's current Biological Opinions (BO's) (USFWS 2008, NMFS 2009).




Figure 8: Delta Map showing various locations of X2 as listed by kilometers from Golden Gate Bridge. Credit: Jeanne DiLeo, USGS

The close association of delta smelt with the San Francisco Estuary's LSZ and X2 has been known for many years (Stevens and Miller 1983; Moyle *et al.* 1992). There have been documented changes to the delta smelt's LSZ habitat that have led to present-day habitat conditions. Reduced Delta outflow (tidally-averaged flow at Chipps Island) causes the LSZ and X2 to move upstream which results in reduced habitat quality for the delta smelt (Bennett 2005,

pp. 11, 20). Abundance of delta smelt in the fall has been linked to fall habitat conditions (Feyrer *et al.* 2011, p. 123).

From late spring through fall and early winter, most delta smelt occur in the LSZ, which has a geographic location that varies with Delta outflow (Dege and Brown 2004, pp. 56–58; USFWS 2008, pp. 147, 150). Higher Delta outflow moves the LSZ westward into Suisun Bay and sometimes even San Pablo Bay, and lower Delta outflow allows the LSZ to encroach eastward into the Delta (Kimmerer *et al.* 2013, pp. 6–8). Delta outflow lower than ~ 11,400 cfs (amount of flow needed to keep X2 at Chipps Island) concentrates delta smelt in the Delta's channelized waterways where habitat variability is lower, the availability of shoals is lower, and entrainment risk is higher (SWRCB 1995, p. 26, Moyle 2002, p. 230, Bennett 2005, pp. 11, 20). Low outflow is associated with greater water clarity, which as discussed above can have consequences for delta smelt survival. Additionally, clam abundance in Suisun Bay tends to increase when X2 shifts upstream (See Factor E: Introduced Species for detailed analysis). The location of X2 during spring has also been shown to affect delta smelt larval abundance in recent years (Baxter *et al.* 2015, p. 47). Delta outflow varies naturally within and among years due to variation in precipitation and snowmelt. However, present-day Delta outflow is lower than historical outflows and, in some seasons, is much less variable than historical outflows due to the storage and diversion of water throughout the Sierra-Nevada and Central Valley. Recent declines in Delta outflow are closely linked to water exports at the State and Federal diversion facilities (Cloern and Jassby 2012, p. 6–8).

The seasonal distribution of the delta smelt population is strongly influenced by river flows because the quantity of fresh water flowing through the estuary changes the amount and location of suitable turbid, low-salinity, open-water habitat (Feyrer *et al.* 2007 pp. 728–732; Feyrer *et al.* 2011, p. 124, Fig 2). The FMWT data have been used to show that delta smelt relative abundance changes when the extent of habitat (low salinity and low transparency water) changes (Feyrer *et al.*, 2011, p. 124). When X2 is located downstream of the confluence at 80 km, the area of suitable habitat is increased because it encompasses the broad and relatively shallow areas of Suisun and Grizzly Bays (Feyrer *et al.* 2011, p. 125; Figure 8). Delta smelt habitat increases by approximately twofold when the location of X2 moves from 85 km to 70 km (Feyrer *et al.* 2011, p. 123).

The State of California is currently in a drought which is further decreasing freshwater flows. The Governor of California declared a State of Emergency on January 17, 2014 due to the drought and directed State officials to take all necessary actions to make water immediately available. As of October 2016, the Governor's drought declaration remains in place and the current drought conditions are comparable to the driest years on record in California. The severity of California's drought was exacerbated by record warm temperatures and below normal precipitation in 2015, resulting in a severely reduced snowpack. The Governor responded to this low precipitation by signing emergency drought relief funding for critical water infrastructure projects and emergency drought actions. During the last two years, Federal and State governments (U.S. Bureau of Reclamation [USBR] and California Department of Water Resources [DWR]) have taken actions to ensure the reduced water quality and supply does not reach a level of concern for human health and safety, while complying with biological opinions. The actions taken include the 2015 placement of a salinity rock barrier on West False River and

numerous Temporary Urgency Change Orders from the California State Water Resources Control Board to DWR and Reclamation in 2014 and 2015 to modify requirements under Decision 1641 to meet certain water quality objectives, reduction of river flows caused by low reservoir storage, and river temperature requirements. These actions have reduced fresh water outflow to the San Francisco Bay Delta Estuary. The CDFW fish surveys indicate that the relative abundance of delta smelt is currently the lowest on record. Detailed results of these surveys were presented above under population indices. The low index numbers represent the additive impact of drought to the delta smelt and its habitat.

Climate Change

Climate change is likely already impacting the delta smelt. Climate change is discussed here under Factor A because, although it may affect the delta smelt directly by creating physiological stress, the primary impacts of climate change on the species are expected to be through changes in the availability and distribution of delta smelt habitat.

Our analyses under the Act include consideration of ongoing and projected changes in climate. The terms “climate” and “climate change” are defined by the Intergovernmental Panel on Climate Change (IPCC). The term “climate” refers to the mean and variability of different types of weather conditions over time, with 30 years being a typical period for such measurements (IPCC 2013a, p. 1450). The term “climate change” thus refers to a change in the mean or variability of one or more measures of climate (for example, temperature or precipitation) that persists for an extended period, whether the change is due to natural variability or human activity (IPCC 2013a, p. 1450).

Scientific measurements spanning several decades demonstrate that changes in climate are occurring, and that the rate of change has increased since the 1950s. Examples include warming of the global climate system, and substantial increases in precipitation in some regions of the world and decreases in other regions (for these and other examples, see Solomon *et al.* 2007, pp. 35–54, 82–85; IPCC 2013b, pp. 3–29; IPCC 2014, pp. 1–32). Results of scientific analyses presented by the IPCC show that most of the observed increase in global average temperature since the mid-20th century cannot be explained by natural variability in climate and is “very likely” (defined by the IPCC as 90 percent or higher probability) due to the observed increase in greenhouse gas (GHG) concentrations in the atmosphere as a result of human activities, particularly carbon dioxide emissions from use of fossil fuels (Solomon *et al.* 2007, pp. 21–35; IPCC 2013b, pp. 11–12 and figures SPM.4 and SPM.5). Further confirmation of the role of GHGs comes from analyses by Huber and Knutti (2011, p. 4), who concluded it is extremely likely that approximately 75 percent of global warming since 1950 has been caused by human activities.

Scientists use a variety of climate models, which include consideration of natural processes and variability, as well as various scenarios of potential levels and timing of GHG emissions, to evaluate the causes of changes already observed and to project future changes in temperature and other climate conditions (Meehl *et al.* 2007, entire; Ganguly *et al.* 2009, pp. 11555, 15558; Prinn *et al.* 2011, pp. 527, 529). All combinations of models and emissions scenarios yield very similar projections of increases in the most common measure of climate change, average global surface

temperature until about 2030. Although projections of the magnitude and rate of warming differ after about 2030, the overall trajectory of all the projections is one of increasing global warming through the end of this century, even for the projections based on scenarios that assume that GHG emissions will stabilize or decline. Thus, there is strong scientific support for projections that warming will continue through the 21st century, and that the magnitude and rate of change will be influenced substantially by the extent of GHG emissions (Meehl *et al.* 2007, pp. 760–764, 797–811; Ganguly *et al.* 2009, pp. 15555–15558; Prinn *et al.* 2011, pp. 527, 529; IPCC 2013b, pp. 19–23). See IPCC 2013b (entire), for a summary of other global projections of climate-related changes, such as frequency of heat waves and changes in precipitation.

Various changes in climate may have direct or indirect effects on species. These effects may be positive, neutral, or negative, and they may change over time, depending on the species and other relevant considerations, such as threats in combination and interactions of climate with other variables (for example, habitat fragmentation) (IPCC 2014, pp. 4–11). Identifying likely effects often involves aspects of climate change vulnerability analysis. Vulnerability refers to the degree to which a species (or system) is susceptible to, and unable to cope with, adverse effects of climate change, including climate variability and extremes. Vulnerability is a function of the type, magnitude, and rate of climate change and variation to which a species is exposed, its sensitivity, and its adaptive capacity (Glick *et al.* 2011, pp. 19–22; IPCC 2014, p. 5). There is no single method for conducting such analyses that applies to all situations (Glick *et al.* 2011, p. 3). We use our expert judgment and appropriate analytical approaches to weigh relevant information, including uncertainty, in our consideration of the best scientific information available regarding various aspects of climate change.

Global climate projections are informative, and, in some cases, the only or the best scientific information available for us to use. However, projected changes in climate and related impacts can vary across and within different regions of the world (IPCC 2013b, pp. 15–16). Therefore, we use “downscaled” projections when they are available and have been developed through appropriate scientific procedures, because such projections provide higher resolution information that is more relevant to spatial scales used for analyses of a given species (see Glick *et al.* 2011, pp. 58–61, for a discussion of downscaling). With regard to our analysis for the delta smelt and overall Bay-Delta, downscaled projections are available (e.g., Dettinger 2005, p. 295–299) and they have been applied to forecasting delta smelt habitat conditions (Feyrer *et al.* 2011; Cloern *et al.* 2011; Brown *et al.* 2013).

San Francisco Bay-Delta Climate Change

The effects of climate change do not act in isolation; they are anticipated to exacerbate existing threats to delta smelt. We considered the potential effects of climate change on the delta smelt based on projections derived from various modeling scenarios. A series of publications (Feyrer *et al.* 2011; Cloern *et al.* 2011; Brown *et al.* 2013) have modeled future impacts of climate change in the delta and projected how this will affect delta smelt. These models used the B1 and A2 scenarios from the 2007 IPCC report. Each scenario included both a warmer-wetter and warmer-dryer sub scenario. Modeled predictions presented in these publications are based on current baseline conditions (no increased outflow, no breaching of levees) which may or may not change in the future. Temperature increases are likely to lead to a continued rise in sea level, further increasing salinity which will increasingly restrict delta smelt’s already limited geographic range

(Feyrer *et al.* 2011, p. 124; Cloern *et al.* 2011, p. 7; Brown *et al.* 2013, p. 761). Higher air temperatures will reduce snowpack, melt snow earlier in the winter or spring, and increase water temperatures. These changes will likely alter freshwater flows, possibly shifting and condensing the timing and location of delta smelt reproduction (Brown *et al.* 2013, p. 765).

Projections indicate that temperature and precipitation changes will diminish snowpack, changing the availability of natural water supplies (USBR 2011, p. 143). Warming may result in more precipitation falling as rain and less storage as snow. This would result in increased rain on snow events and increase winter runoff with an associated decrease in runoff for the remainder of the year (USBR 2011, p. 147). Sacramento Valley Ecoregion projections include a 27 percent decrease in annual freshwater flows and earlier snowmelts, with increased freshwater flows in January and February but reduced throughout the rest of the year (PRBO 2011, p. 27). Earlier seasonal warming increases the likelihood of rain-on-snow events, which are associated with mid-winter floods. Smaller snowpacks that melt earlier in the year may result in increased drought frequency and severity (Rieman and Isaak 2010, p. 6). Thus overall, these changes may lead to increased frequency of flood and drought cycles during the 21st century (USBR 2011, p. 149).

Sea level rise is likely to increase the frequency and range of saltwater intrusion. Salinity within the northern San Francisco Bay is projected to rise by 4.5 psu (Practical Salinity Unit) by the end of the century (Cloern *et al.* 2011, p. 7). Elevated salinity levels could push the position of X2 farther up the estuary if outflows were not increased to compensate for it. Fall X2 mean values are projected to increase by a mean of about 7 km to the area of Antioch for a distance of approximately 90 km from the Golden Gate Bridge by 2100 (Brown *et al.* 2013, p. 765). This increase in the position of X2 in the fall is expected to result in a decrease in suitable physical habitat (Brown *et al.* 2013, p. 765) if current levees and channel structures are maintained. A decrease in spring habitat due to the movement of X2 upstream due to sea level rise is also expected to result from climate change.

We expect warmer estuary temperatures to be yet another significant conservation challenge based on climate change models. Mean annual water temperatures within the Delta are expected to increase steadily during the second half of this century (Cloern *et al.* 2011, p. 7). Warmer water temperatures could reduce delta smelt growth, increase delta smelt mortality and constrict suitable habitat within the estuary during the summer months. Due to warming temperatures, delta smelt are projected to spawn an average of ten to twenty-five days earlier in the season depending on the location (Brown *et al.* 2013, p. 765). Also due to expected temperature increases, total number of high mortality days (cumulative number of days of daily average water temperature >25 °C (77°F)) is expected to increase for all IPCC climate change scenarios (Brown *et al.* 2013, p. 765). The number of stress days (cumulative number of days of daily average water temperature >20 °C (68°F)) is expected to be stable or decrease partly because many stress days will become high mortality days. This could lead to delta smelt being forced to grow under highly stressful conditions during summer and fall with less time to mature because of advanced spawning (Brown *et al.* 2013, p. 766). More recent research suggests that delta smelt will face a shorter maturation window and significant thermal habitat constriction. A shorter maturation window will likely have effects on reproduction success (Brown 2016, p. 14). Growth rates have been shown to slow as water temperatures increase, requiring delta smelt to

consume more food to reach growth rates that are normal at lower water temperatures (Rose *et al.* 2013a, p. 1252). Delta smelt are already often smaller than they used to be (Sweetnam 1999, p. 23; Bennett 2005, p. 46) and expected temperature increases due to climate change will likely further slow growth rates.

Warmer water will tend to move the spawning season earlier in the year (Brown *et al.* 2013, p. 769). This means the fish will have to grow faster to compensate for the shorter growing season to produce as many eggs as they do now. This may already be a serious limitation on their population fecundity (Rose *et al.* 2013b, p. 1268). Higher temperatures may restrict delta smelt distribution into the fall, limiting their presence in Suisun Bay for more than just salinity reasons and forcing greater inhabitation of cooler, high salinity waters (Brown *et al.* 2013, p. 769). Water temperatures are already presently above 20°C (68 °F) for most of the summer in core habitat areas, sometimes even exceeding 25 °C (77 °F) for short periods.

The delta smelt is currently at the southern limit of the inland distribution of the family Osmeridae along the Pacific coast of North America. That indicates that this region was historically already about as warm as fish in the Osmeridae family can handle. Increased temperatures associated with climate change may result in a habitat in the Bay-Delta that is outside of the delta smelt's ecological tolerance limits.

Reduced Turbidity

Turbid conditions throughout the estuary attenuate light in the water column and limit phytoplankton growth rate and thereby productivity (Cloern 1987, p. 1375–1378; Jassby 2008, p. 14). This is one reason why eutrophication has not presented a major problem here, despite high nutrient concentrations. Increasing water clarity could allow for higher productivity and eventually for eutrophication. In addition to its importance in regulating primary productivity, turbidity is an important component of fish habitat in the Bay-Delta and elsewhere and delta smelt are strongly associated with turbid water (Feyrer *et al.* 2007, p. 728; Nobriga *et al.* 2008, p. 7; Feyrer *et al.* 2011, p. 123). Turbidity has been shown to be important for successful feeding in some species (Sirois and Dodson 2000, pp. 240–243; Baskerville-Bridges *et al.* 2004, p. 223; Horppila *et al.* 2004, p. 1864) and may also help small fish to avoid predation.

First-feeding delta smelt larvae require relatively turbid waters to capture prey (Baskerville-Bridges *et al.* 2004, p. 223). Hasenbein *et al.* (2016, p. 9) placed 60 day post hatched delta smelt in waters with turbidities of 5, 12, 25, 35, 50, 80, 120 and 250 NTU to test feeding rates at these turbidities. Turbidities of 25 to 80 NTU were determined to be the optimal range in the tested conditions as evident from the highest survival, feeding and changes in gene expression compared with other treatments. Delta smelt may also use turbidity as cover from predators; this was hypothesized based on long-term monitoring of the distribution of fish in the wild (e.g., Feyrer *et al.* 2007, p. 731) and recently supported by a laboratory experiment (Ferrari *et al.* 2014, p. 87, Fig 4). From the 1950's to present, the Delta has experienced a decline in turbidity (Wright and Schoellhamer 2004, p. 12) that culminated in an estuary-wide step-decline in 1999 (Schoellhamer 2011, p. 897).

The increased water clarity in delta smelt rearing habitat in recent decades is attributed to the interruption of sediment transport by upstream dams (Arthur and Ball 1979, p. 157; Wright and

Schoellhamer 2004, pp. 7, 10) and the spread of the exotic introduced water plant *Egeria densa* (Brazilian waterweed), which traps suspended sediments (Feyrer *et al.* 2007, p. 731). The likelihood of delta smelt occurrence in trawls at a given sampling station decreases with increasing Secchi depth at the stations (Feyrer *et al.* 2007, p. 728). This is consistent with behavioral observations of captive delta smelt (Nobriga and Herbold 2008, p.11). Few daylight trawls catch delta smelt at Secchi depths over one half meter and capture probabilities of delta smelt are highest at 0.40 m depth or less. Since 1978, delta smelt have become increasingly rare in summer and fall surveys of the San Joaquin region of the Delta (Nobriga *et al.* 2008, p. 9). One reason appears to be the comparatively high water clarity in the region, although high water temperatures are also likely a contributing factor (Nobriga *et al.* 2008, pp. 8, 9).

Channel Disturbance, Dredging, Sand-mining

The placement of riprap bank protection has led to the loss of riparian habitat, large woody debris, shallow water habitat, and natural channel migration. Bank stabilization and riprapping has been shown to change natural river processes; reduce channel meandering, which reduces habitat complexity; create a smooth, hydraulically enhanced surface that is not conducive to the habitat requirements of fish including delta smelt; stop erosion, which stops woody vegetation from entering the river and reduces the long-term recruitment of large woody debris; inhibit plant growth through thick rock at the waterline, which causes vegetation to grow further from the shoreline and a subsequent reduction in outside food sources for aquatic invertebrates; and decrease near-shore roughness, which contributes to increased stream velocities and a decrease in available refuge for fish (USFWS 2000, pp. 6–12). Bank protection along the Sacramento River has contributed to habitat fragmentation. More than half of the river's banks in the lower 194 miles have been riprapped, mostly under the Army Corps of Engineers Sacramento River Bank Protection Project (SRBPP). The historical condition of the Sacramento River was free-flowing, without restrictions brought about by diversions and dams. Late summer flows were low compared to today's summer flows, and high flows during spring caused overbank flooding into areas that contained riparian forests (USFWS 2000, p. 7). Bank erosion and river meander were natural ecological processes. Today, most of the riparian forests and wetlands have been removed, and much of the historical habitat has been lost from the Sacramento River.

Ongoing maintenance dredging regularly occurs in the Sacramento Deep Water Ship Channel of the Delta. Dredging can change the light transmittance, dissolved oxygen and nutrient concentrations, salinity, temperature and pH of the water (Navy 1990, entire). Dredging will re-suspend contaminants if they are present in the surface sediments (Levine-Fricke 2004, p. 44). Dredging can result in entrainment, injury or displacement (particularly in marinas) of delta smelt (Levine-Fricke 2004, p. 67–72). These effects are localized and the plumes do not last long once dredging stops (Schoellhamer 2002, p. 491).

Sand mining is most likely to affect delta smelt at the egg and larval life stages. There are a number of measures in place to minimize the effects of sand mining in the estuary. Applicants are required to install fish screens in compliance with CDFW and NMFS criteria over sand mining vent pipes to exclude juvenile and adult fish from entrainment during sand mining events. In addition, a work window of December 1 through June 30 is in place and during this time sand mining operations are restricted to areas that are 20 ft. or greater in depth. Delta

smelt are thought to spawn in water 15 ft. or less. This will avoid spawning habitat in shallower depths for delta smelt. Sand mining volume percentages during the spawning period of delta smelt are also reduced. Because spawning substrate is not known to be limited for the species (Hobbs *et al.* 2007, entire), restrictions are in place to protect delta smelt, and sand is a dominant substrate in the estuary, sand mining is not expected to limit spawning.

Summary for Factor A

Based on a review of the best scientific and commercial information available, we find that destruction, modification, or curtailment of habitat poses a threat to delta smelt due to a suite of factors. The operation of upstream reservoirs, water exports, and cumulative water diversions has altered the magnitude, duration, and frequency of Delta outflows and the location and extent of the LSZ and has reduced habitat that the delta smelt uses. Lower turbidity reduces larval foraging efficiency and increases predation risk. Forecasted warmer water temperatures and higher salinity in the Delta due to climate change will likely further impair delta smelt habitat in the future. Channel disturbance, dredging and sand mining do not rise to the level of a threat that is currently acting on the species at the population level. Although channel modification and levee construction have altered the delta and resulted in habitat fragmentation and depletion for the delta smelt, this is a historic threat and not a current threat.

B. Overutilization for commercial, recreational, scientific, or educational purposes.

Delta smelt monitoring surveys are conducted throughout the year, including the Fall Mid-Water Trawl (FMWT), Summer Towntnet Survey (TNS), 20-mm Survey, and Spring Kodiak Trawl Survey (SKT). Other routine monitoring surveys also collect delta smelt like the Service's Beach Seine and trawl surveys targeting juvenile salmon, CDFW's San Francisco Bay Study trawling, UC Davis' Suisun Marsh fish monitoring, and Early Warning Surveys meant to notify decision makers of when delta smelt are moving towards the water project facilities. Overall take in survey collections is believed to be low compared to estimated relative abundances (Bennett 2005, p. 7); however, considering the concern for reduced abundance based on trend assessment, questions arise as to whether these and other surveys could pose a concern to the delta smelt. Because of low abundance and a high level of sampling mortality, some survey methods have been modified to limit incidental catches of delta smelt when delta smelt is not the target species.

Annual combined scientific take of adults and juveniles (> 20 mm) from the 29 surveys conducted in the last ten years has ranged from 818 individuals to 4713 individuals (Slater 2015, no pagination). Annual combined scientific take of larvae (< 20 mm) from the 29 surveys conducted in the last ten years has ranged from 412 individuals to 1769 individuals (Slater 2015, no pagination, Smith 2015, no pagination). Many of these fish are returned to the delta and survive. However, because take does include the act of catching and handling the fish, any individual that is captured is counted in the take total. Based on the low number of delta smelt collected in sampling surveys and the modified methods employed to further reduce these collections, we find that the amount of take expected to occur from sampling surveys does not reach a level substantial enough to be considered a threat. There is no evidence of use of the species for other commercial, recreational, scientific, or educational purposes.

Summary for Factor B

Based on a review of the best scientific information available, we find that overutilization for commercial, recreational, scientific or educational purposes is not a threat to the delta smelt.

C. Disease or predation.

Disease

Studies have not found evidence of significant disease infestations in wild delta smelt (Teh 2007, p. 8; Baxter *et al.* 2008, p. 14) (See contaminants discussion in Factor E for more information). Based on the best scientific and commercial information available, we conclude that disease is not a threat to the delta smelt.

Predation

There are numerous fish species that have been confirmed to be delta smelt predators either through visual analysis of their stomach contents or by detection of delta smelt DNA in their digestive tracts. It seems likely that delta smelt are also occasionally consumed by piscivorous birds that forage in their habitat but we are unaware of any data available to confirm this hypothesis. The following paragraphs focus on three predators that the local scientific community has concentrated their research efforts on for their potential to affect delta smelt viability.

The predator with the highest historical documentation of predation on delta smelt is striped bass (*Morone saxatilis*; Stevens 1963, pp. 12–28; 1966, entire; Thomas 1967, p. 58). In these studies, striped bass were confirmed to prey on both juvenile and adult delta smelt. Striped bass are widely distributed in pelagic areas of the San Francisco Bay-Delta and parts of its watershed, and thus striped bass distribution fully encompasses the distribution of delta smelt juveniles and adults (Nobriga *et al.* 2013, p. 1564). Striped bass also tend to aggregate in the vicinity of water diversion structures, where delta smelt are frequently entrained (Nobriga and Feyrer 2007, p. 9). Thus, striped bass are likely to be the most significant predator of post-larval delta smelt (Nobriga and Feyrer 2007, p. 9). No inverse correlations between the abundance of striped bass and the relative abundance of delta smelt have been found to date using a variety of statistical approaches (Mac Nally *et al.* 2010, entire; Thomson *et al.* 2010, entire; Maunder and Deriso 2011, p. 1302; Miller *et al.* 2012, p. 16; Nobriga *et al.* 2013, p. 1571). Although the relative rarity of delta smelt in the estuary food web would presumably make them an incidental prey item for striped bass, it is possible that striped bass abundance and demand for prey are always high enough to limit delta smelt population growth rate (Nobriga *et al.* 2013, p. 1574). However, focused studies of this predator-prey linkage would be required to determine whether predation by striped bass is high enough to be of concern or not.

Largemouth bass are freshwater fish that prefer clear waters along shorelines (littoral habitat) with relatively dense water plants (Nobriga and Feyrer 2007, pp. 4, 8; Brown and Michniuk 2007, p. 196; Baxter *et al.* 2008, p. 17). This is a suite of habitat characteristics that is distinctly different from those described above for delta smelt. Thus, unlike delta smelt and striped bass, delta smelt and largemouth bass have different habitat requirements (e.g., Nobriga *et al.* 2005, p.

783) and their distributions do not strongly overlap. However, there has been a major increase in the Delta's largemouth bass population since the early 1990s that is believed to have been facilitated by the spread of the introduced plant *Egeria densa*, which provides rearing habitat for the bass (Baxter *et al.* 2008, p. 17). Despite increases in largemouth bass populations and habitat, Nobriga and Feyrer (2007, p 6) did not find delta smelt as largemouth bass prey. Nor have more recent and extensive surveys of largemouth bass stomach contents (Baxter *et al.* 2015, p. 65; L. Conrad, CDWR, unpublished data). In captivity however, even young juvenile largemouth bass will attempt to consume delta smelt (Ferrari *et al.* 2014, p. 87) so they presumably represent a predation threat when the species closely co-occur in the wild. In contrast to the situation for striped bass, several researchers have found inverse correlations between the relative abundance of largemouth bass or multi-species indices that included largemouth bass and the relative abundance of delta smelt (Mac Nally *et al.* 2010, p. 1425, Fig 3b; Thomson *et al.* 2010, p. 1439, Fig 3c; Maunder and Deriso 2011, p. 1297, Table 6). At this time however, there is no way to determine whether these correlations are causative (predation by largemouth bass caused delta smelt to decline) or not (delta smelt simply use different habitats than largemouth bass and delta smelt habitat has decreased while largemouth bass habitat has increased).

Due to their size, juvenile and adult delta smelt are mainly vulnerable to larger predatory fishes. However, delta smelt eggs and larvae are very small (eggs are ~ 1 mm in diameter and larvae hatch at ~ 5–6 mm in length). Thus, these early life stages of delta smelt are potentially available prey to a much greater number of predators. One of these is the nonnative Mississippi silverside (*Menidia audens*), which like delta smelt is an annual fish with a maximum length near 100 mm (4 in.). Mississippi silversides may be both predators and competitors of delta smelt (Bennett 2005, pp. 49, 50). Mississippi silversides were first introduced to the San Francisco Bay-Delta in the mid-1970s, and have increased dramatically in numbers since the mid-1980s. They forage in schools around the shoreline habitats and tidal marsh channels of the San Francisco Bay-Delta, where they are abundant (Matern *et al.* 2002, p. 802, Table 2; Nobriga *et al.* 2005, p. 781, Table 3; Gewant and Bollens 2012, p. 479, Table 2). They readily consume delta smelt larvae in aquarium tests. Bennett (2005, p. 50) concluded that “delta smelt are at high risk if eggs or larvae co-occur with schools of foraging silversides.”

Baerwald *et al.* (2012, p. 1604) recently confirmed that delta smelt DNA was present in the guts of Mississippi silverside collected in the Sacramento Deep Water Shipping Channel. This is an area that often has high catches of delta smelt larvae (CDFW 2014, no pagination). Baerwald *et al.* (2012, p. 1606) detected delta smelt DNA in the guts of 41% of Mississippi silversides that were collected in Kodiak Trawl (N=37), but had no detections of delta smelt DNA in the guts of the much larger fraction of Mississippi silversides collected nearshore with beach seines (n=614), resulting in an overall detection rate of 2.3% (n=651). Although generally inverse relative abundance trends between delta smelt and Mississippi silverside have been recognized for several decades (Bennett and Moyle 1996, p. 530; Bennett 2005, p. 50), two recent statistical evaluations of delta smelt relative abundance trends both concluded that Mississippi silverside densities did not significantly correlate with the delta smelt trends when other factors were considered (Mac Nally *et al.* 2010, p. 1422; Thomson *et al.* 2010, p. 1441).

Two other recent studies tested for the effect of overall composite ‘predation’ variable on delta smelt trends. Maunder and Deriso (2011, p. 1300) found this variable was a good predictor in

several variations of their statistical life cycle model, while Miller *et al.* (2012, p. 14) did not include it in their “best” models. In this composite variable, Mississippi silverside relative abundance was combined with relative abundance estimates of largemouth bass and several other nonnative sunfishes and a quadratic term for average seasonal water transparency. Thus, the meaning of this variable is extremely confounded.

Summary for Factor C

Based on a review of the best scientific information available, we find that disease is not a threat to the delta smelt. Delta smelt is a rare fish and has been a rare fish (compared to other species) for at least the past several decades (Nobriga and Herbold 2008). Therefore, it has also been rare in examinations of predator stomach contents as would be expected and it is not surprising that studies are inconclusive on the extent of population level effects. Although predation is a naturally occurring mechanism, non-native fishes that have been introduced into the delta have increased the risk of predation to delta smelt. Current and historical evidence of delta smelt in stomachs of the relatively small number of non-native fishes sampled compared to overall non-native population numbers shows that predation from non-native fishes is likely having some effect on overall population numbers of delta smelt. We conclude that predation is an additional threat to delta smelt.

D. The inadequacy of existing regulatory mechanisms.

State Laws

California Endangered Species Act: The delta smelt was listed as threatened under the California Endangered Species Act (CESA) in 1993 (CDFW 2014b), and was reclassified as endangered under the CESA in 2010 (14 CCR 670.5). The CESA prohibits unpermitted possession, purchase, sale, or take of listed species. However, the CESA definition of take does not include harm, which under the Federal Endangered Species Act can include destruction of habitat that actually kills or injures wildlife by significantly impairing essential behavioral patterns (50 CFR 17.3). The CESA does require consultation between the CDFW and other State agencies to ensure that activities of State agencies will not jeopardize the continued existence of State-listed species (CDFW 2014c).

Porter Cologne Water Quality Control Act: The Porter-Cologne Water Quality Control Act (California Water Code 13000 *et seq.*) is a California State law that established the State Water Resources Control Board (SWRCB) and nine Regional Water Quality Control Boards that are responsible for the regulation of activities and factors that could degrade California water quality and for the allocation of surface water rights (California Water Code Division 7). In 1995, the SWRCB developed the Bay-Delta Water Quality Control Plan which expanded water quality objectives for the Delta. This plan is currently implemented by Water Rights Decision 1641, which imposes flow and water quality standards on the State and Federal water export facilities to assure protection of beneficial uses in the Delta (USFWS 2008, pp. 21–27). The various flow and salinity objectives can constrain export pumping and were designed, in part, to protect fisheries. These objectives include specific freshwater flow requirements throughout the year, specific water export restraints in the spring, and water export limits based on a percentage of estuary inflow throughout the year. The water quality objectives were designed to protect

agricultural, municipal, industrial, and fishery uses; they vary throughout the year and by hydrology. The SWRCB is in the process of doing a review which may potentially modify water quality standards and objectives. In addition to regulating flow requirements, the Porter Cologne Water Quality Control Act also regulates contaminants released into the delta (see Clean Water Act).

Federal Laws

National Environmental Policy Act: The National Environmental Policy Act (NEPA) (42 U.S.C. 4321 *et seq.*) requires all Federal agencies to formally document, consider, and publicly disclose the environmental impacts of major Federal actions and management decisions significantly affecting the human environment. NEPA documentation is provided in an environmental impact statement, an environmental assessment, or a categorical exclusion, and may be subject to administrative or judicial appeal. However, the Federal agency is not required to select an alternative having the least significant environmental impacts, and may select an action that will adversely affect sensitive species provided that these effects are known and identified in a NEPA document. Therefore, we do not consider the NEPA process in itself to be a regulatory mechanism that is designed to provide significant protection for the delta smelt.

Endangered Species Act: The delta smelt is currently listed as a threatened species under the Endangered Species Act of 1973, as amended (Act). The Act defines a “threatened species” as “any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range” (section 3(20) of the Act). An “endangered species” is “any species which is in danger of extinction throughout all or a significant portion of its range” (section 3(6) of the Act). Section 6 of the Act authorizes us to enter into conservation agreements with States, and to allocate funds for conservation programs to benefit threatened or endangered species. Section 7 of the act requires all Federal agencies, in consultation with and with the assistance of the Secretary (in this case the USFWS), to insure that any action authorized, funded, or carried out by such agency is not likely to jeopardize the continued existence of any endangered or threatened species or result in destruction or adverse modification of their habitat.

The CVP, operated by the Bureau of Reclamation (Reclamation), and SWP, operated by the CDWR, are currently operating under a Biological Opinion (BO) issued December 15, 2008, under section 7 of the Act (USFWS 2008, pp. 1–396). A BO sets a reasonable and prudent alternative (RPA) to protect the species from expected effects from the project if the project puts the species in jeopardy. This BO includes a RPA, according to which water export facility operations could proceed without jeopardizing the continued existence of the species or destroying or adversely modifying its designated critical habitat. It also includes an incidental take Statement (ITS) specifying reasonable and prudent measures necessary to minimize the incidental take of the species resulting from CVP and SWP operations. Reclamation has accepted the RPA provisionally, but may decide to reinstate consultation (USBR 2008, p. 1). The ITS and BO replace a previous ITS and BO issued in 2005 (USFWS 2005, p. 1), and also replace flow restrictions instituted by the District Court in the case of *NRDC v. Kempthorne* (Wanger 2007, pp. 1–11), which found the 2005 BO inadequate to conserve the species.

Central Valley Project Improvement Act: The Central Valley Project Improvement Act (CVPIA) amends the previous CVP authorizations to include fish and wildlife protection, restoration, and mitigation as project purposes having equal priority with irrigation and domestic uses, and fish and wildlife enhancement as having an equal priority with power generation. Included in CVPIA section 3406 (b)(2) was a provision to dedicate 800,000 acre-feet of Central Valley Project yield annually (referred to as “(b)(2) water”) for fish, wildlife, and habitat restoration. Since 1993, (b)(2) water has been used and in some years supplemented with acquired environmental water pursuant to CVPIA section 3406 (b)(3) to increase stream flows and reduce Central Valley Project export pumping in the Delta. These management actions were taken to contribute to the CVPIA salmonid population doubling goals and to protect delta smelt and their habitat (Guinee 2014, pers. comm.). As discussed above (under Biology and Factor A), increased freshwater flows have been shown to be beneficial to delta smelt.

Clean Water Act: The Clean Water Act (CWA) provides the basis for the National Pollutant Discharge Elimination System (NPDES). The CWA gives the EPA the authority to set effluent limits and requires any entity discharging pollutants to obtain a NPDES permit. The EPA is authorized through the CWA to delegate the authority to issue NPDES Permits to State governments. In States that have been authorized to implement CWA programs, the EPA still retains oversight responsibilities (USEPA 2014). California is one of these States to which the EPA has delegated CWA authority. The Porter-Cologne Water Quality Control Act established the California State Water Resources Control Board (SWRCB) and nine Regional Water Quality Control Boards that are now responsible for issuing these NPDES permits, including permits for the discharge of effluents such as ammonia. The SWRCB is responsible for regulating activities and factors that could degrade California water quality (California Water Code Division 7, section 13370–13389).

The release of ammonia into the estuary has been shown to have detrimental effects on the Delta ecosystem by inhibiting the production of diatoms, a kind of phytoplankton (see Factor E, below). The release of ammonia is controlled primarily by the CWA (Federal law) and secondarily through the Porter-Cologne Water Quality Control Act (State law). In 2013, the EPA updated freshwater discharge criteria that included new more stringent limits on ammonia (USEPA 2013, pp. 1–3). This was done to limit the effects of ammonia on freshwater clams and snails in the Delta. In addition, an NPDES permit for the Sacramento Regional Wastewater Treatment Plant, a major discharger, was prepared by the California Central Valley Regional Water Quality Control Board in the fall of 2010, with new ammonia limitations intended to reduce loadings to the Delta. The new ammonia limits will take effect in May 2021 (CRWQCB 2014, p. 17). Upon completion of the treatment plant modifications, ammonia discharge by the facility will be reduced by 99.4% (USFWS 2014, p. 25).

Summary for Factor D

We have identified a number of existing regulatory mechanisms that provide protective measures that affect the stressors acting on the Delta smelt. Despite these existing regulatory mechanisms and other conservations efforts, the stressors continue to act on the species such that it is warranted for uplisting under the Act.

E. Other natural or manmade factors affecting its continued existence.

Other factors affecting the continued existence of the delta smelt include entrainment (fish drawn in and transported through the flow of water) into water diversions, introduced species, contaminants, and increased vulnerabilities of small populations.

Entrainment

Water Export Facilities: The two largest pumping facilities located in the South Delta, Jones Pumping Plant (CVP) and Banks Pumping Plant (SWP) facilities exported between 4.31 and 7.74 km³ (3.49 and 6.28 million acre-feet) per year between water years (October 1–September 30) 1999 and 2008. The average annual water year export amount from 1999–2008 was 6.74 km³ (5.46 million acre-feet). In December of 2008 (water year 2009), the Service issued a BO that has actions that are designed to protect delta smelt from entrainment into the facilities. Since the BO has been in place, these two State and Federal facilities have exported between 4.29 and 8.03 km³ (3.48 and 6.51 million acre-feet) per year from 2009 through 2013. Average annual water year exports from 2009–2013 were 5.75 km³ (4.66 million acre feet) (Aasen 2014, no pagination).

Operation of water export facilities directly affects fish by entraining them into the diversion facility. Delta smelt's risk of entrainment varies with the environmental and manmade effects on Delta hydrology and the habitats occupied by delta smelt when water is being diverted (Culberson *et al.* 2004, pp. 260–262; Kimmerer and Nobriga 2008, pp. 19–20). It is important that we differentiate “entrainment” from “salvage.” Fish are considered “entrained” at the SWP when they enter the 31,000 acre-foot Clifton Court Forebay in the south Delta. Fish are considered “salvaged” at the SWP when they are actually drawn into Skinner Fish Facility within the SWP. At the CVP or Federal facility, fish are considered entrained and salvaged when they are drawn into the louvers of the pumping facility. Entrainment of delta smelt varies within and among seasons and among years. Studies of entrainment at the State and Federal export facilities found that entrainment rates increased with reverse flows in the southern Delta, which are a function of export rates and Delta inflows (Kimmerer and Nobriga 2008 p. 17, Fig 16; Kimmerer 2008, p. 20–22). The entrainment of adult delta smelt at CVP and SWP occurs mainly during their upstream spawning migrations between December and March (Grimaldo *et al.* 2009, p. 1257). The salvage of age zero delta smelt occurs from April–July with a peak in May–June (Grimaldo *et al.* 2009, p. 1257). Kimmerer (2008, p. 20, 22; 2011, p. 4) estimated that from 0 to 25 percent of the larval population and 1 to 38 percent of the adult population is entrained annually by the State and Federal export facilities with average annual losses estimated at 10% of the population (Kimmerer 2008, p. 25, 2011). Rose extrapolated from this study to find annual losses varying between 1–23% of the adult population (Rose 2013, p. 1251). The majority of entrained delta smelt do not make it to the pumps to enter the fish facility to undergo the salvage process and instead are lost through predation or other mechanisms in the Clifton Court Forebay (Castillo 2012, p. 14). Due to continuing declines in smelt numbers, very few smelt are actually making it to the salvage facility (< 30 a year).

Export of water by the CVP and SWP can limit the reproductive success of delta smelt in the San Joaquin River by entraining most larvae during downstream transport from spawning sites to rearing areas (Kimmerer and Nobriga 2008, p. 11). Winter entrainment of delta smelt represents

a loss of pre-spawning adults and their reproductive potential (Sommer *et al.* 2007, p. 275). High winter exports in the early 1980s have been associated with population decline by at least one recent study (Thomson *et al.* 2010, p. 1296), an association that was noted many years earlier by Moyle *et al.* (1992, p. 74). While there are many factors contributing to the declining trend in delta smelt abundance estimates, we consider entrainment by State and Federal water export facilities to be a significant and ongoing threat to the delta smelt.

Agricultural Diversions for Irrigation and Waterfowl Management: Water is diverted at numerous sites throughout the Bay-Delta for agriculture and waterfowl management. Herren and Kawasaki (2001, p. 343) reported over 2,200 such water diversions within the Delta and another 366 in Suisun Marsh, but CDFG (2009, p. 25) noted that number may be high because Herren and Kawasaki (2001) did not accurately distinguish intake siphons and pumps from discharge pipes. CALFED's Ecosystem Restoration Program (ERP) includes a program to screen remaining unscreened small agricultural diversions in the Delta and the Sacramento and San Joaquin Rivers. The purpose of screening fish diversions is to prevent entrainment losses; however, very little information is available on the efficacy of screening these diversions (Moyle and Israel 2005, p. 20) and the screens themselves can also harm delta smelt that come into contact with them (White *et al.* 2007, entire). Although the ERP has largely run out of funds for the program, there is money set aside in Proposition 1 funding to screen these diversions. However, all but two of the diversion screenings listed on the current priority list will take place in the upper portions of rivers and are focused on protecting salmon (CDFW 2015b, no pagination). Water diversions are primarily located on the edge of channels and along river banks. Delta smelt are generally pelagic and tend to occupy offshore habitats. The only published study that has attempted to evaluate delta smelt's vulnerability to this kind of water diversion in the field concluded they had low vulnerability to entrainment into these diversions (Nobriga *et al.* 2004, p. 293).

Power Plant Diversions: NRG Delta LLC retired the generators for one (Contra Costa Generating Station (CCGS)) of its two power stations within the range of the delta smelt in May of 2013 (NRG 2014, no pagination), although some water is still being pumped to the plant for fire suppression systems (50 acre-ft/yr) (NRG 2014, no pagination). Therefore, this analysis will only consider the other Pittsburg (PGS) power station when assessing threats. The Army Corps of Engineers (Corps) is currently in consultation with the Service for the NRG Delta Energy Project. The project includes operation and dredging activities at the PGS. The PGS is located on the shoreline of the Sacramento River and San Joaquin River confluence and utilizes once-through-cooling for its generators. As part of a settlement agreement with the Coalition for a Sustainable Delta, the Army Corps and NRG Delta (the applicant) agreed to reinstate consultation with the Service for project effects to delta smelt. Power plant operations have been substantially reduced since the late 1970s, when high entrainment and impingement were documented (CDFG 2009a, p. 24).

When the delta smelt was listed in 1993, fourteen units were generating (seven at CCGS and seven at PGS). The original cooling water design flow capacity at CCGS was 986.5 million gallons a day. For PGS, the original cooling water design flow capacity was 1,074 million gallons a day, as permitted in the 2002 Biological Opinions (NMFS 2002, USFWS 2002). From 2007-2010, total annual cooling intake pumped by both power plants ranged between 11,918 –

41,773 million gallons a year. Today there are a total of three units left generating at the PGS. Since the CCGS was retired, total volume pumped was 9,456 million gallons in 2014. The 3 remaining PGS units are expected to be retired in the next few years as the plants complete their transition to natural gas.

In summary, the operation of State and Federal export facilities constitutes an ongoing threat to delta smelt because entrained fish are lost from the reproductive population of delta smelt and this is thought to have a negative effect on the overall delta smelt population. We do not consider entrainment by agricultural diversions to be a significant threat due to their nearshore location and tendency to divert at maximum rates during the summer when many delta smelt are not in channels adjacent to them. Entrainment into power plants appears to have had a significant impact on delta smelt in the past and was cited in the 2010 12-month finding document; however, operations have been modified and the second plant is non-operational, so the effects of operations have been greatly reduced and are not thought to be significant at this time.

Introduced Species

The Bay-Delta zooplankton community has shifted in both the abundance and composition of species over the last several decades (Winder *et al.* 2011, p. 679; Kratina *et al.* 2014, p. 1070). Delta smelt feed primarily on small planktonic (free-floating) crustaceans, and occasionally on insect larvae (Moyle 2002, p. 228). The densities of these prey items depend upon a variety of factors that determine local productivity and the rate at which production from upstream sources is delivered. Historically, the main prey of delta smelt was the copepod *Eurytemora affinis* and the mysid shrimp *Neomysis mercedis*. The slightly larger copepod *Pseudodiaptomus forbesi* has replaced *E. affinis* as a major prey source of delta smelt since its introduction into the San Francisco Bay-Delta. Two other copepod species, *Limnoithona tetraspina* and *Acartiella sinenisi*, have become abundant since their introduction to the San Francisco Bay-Delta in the mid-1990s. *Limnoithona tetraspina* is now the most common copepod in the estuary and as of 2006 made up approximately 95% of the total adult copepods in the LSZ (Bouley and Kimmerer 2006, p. 219). Delta smelt eat these introduced copepods, but *P. forbesi* remains a dominant prey item (Baxter *et al.* 2010, p. 56; Slater and Baxter 2014, p. 8). It has been suggested that *L. tetraspina* may be an inferior food for pelagic fishes including delta smelt because of its small size and generally sedentary behavior (Bouley and Kimmerer 2006, pp. 220–227). Experimental studies addressing this issue have suggested that smelt larvae will attack *L. tetraspina* until they grow large enough to successfully capture larger copepods; also, growth rate of delta smelt fed *L. tetraspina* was lower than that of smelt fed the larger copepods (Sullivan *et al.*, unpublished). *L. tetraspina* is sometimes consumed in large numbers by juvenile delta smelt during late summer when this copepod is abundant in the LSZ (Slater and Baxter 2014, pp. 7–11). As mentioned previously, delta smelt are thought to require a turbid environment for efficient, successful foraging.

Copepods get most of their nutrition from phytoplankton and other direct consumers of phytoplankton. A major reason for the long-term phytoplankton reduction in the upper estuary is grazing by the introduced overbite clam (*Potamocorbula amurensis*), which became abundant by the late 1980s. The overbite clam precipitated major changes in the estuarine food web and has impaired pelagic fish production. Starting about 1987-1988, major step-declines were observed

in the abundance of phytoplankton (Alpine and Cloern 1992, p. 949), mysid shrimp (Orsi and Mecum 1986, p. 331) and the copepod *Eurytemora affinis* due to grazing by the clam (Kimmerer *et al.* 1994, p. 86). The overbite clam incidentally consumes copepod nauplii as it filters phytoplankton and other small organisms from the water (Kimmerer *et al.* 1994, p. 87); therefore, it not only reduces phytoplankton biomass but also competes directly with delta smelt for food because copepod nauplii are the primary prey for delta smelt larvae.

Pelagic primary productivity in the upper San Francisco Estuary is currently poor compared to other estuaries (Kimmerer *et al.* 2012, pp. 920–924) and low fish abundance may be expected as a consequence. Pelagic fishes responded to the food web changes brought on by the overbite clam in several ways. Northern anchovy, a marine fish, largely vacated the low-salinity zone, retreating to saltier water in San Francisco Bay (Kimmerer 2006, p. 211). Striped bass reduced their use of offshore habitats or suffered higher mortality in them giving the impression that they had moved increasingly toward inshore areas (Sommer *et al.* 2011, p. 1456). Many fish species reduced their use of mysids as prey (Feyrer *et al.* 2003, p. 281). The production per unit of flow of striped bass, longfin smelt, starry flounder, and Bay shrimp appears to have declined as a result of overbite clam grazing (Kimmerer 2002b, p. 1281; Kimmerer *et al.* 2009, no pagination). Preliminary information from studies on pelagic fish growth, condition and histology provide additional evidence for food limitation in pelagic fishes in the estuary (Armor *et al.* 2005, p. 38; Baxter *et al.* 2010, p. 50).

Declines in primary productivity and changes in zooplankton assemblages may have played a role in the decline of delta smelt (Rose *et al.* 2013a, b, entire); delta smelt growth rate may have declined (Sweetnam 1999, p. 23; Bennett 2005, p. 46). In 1999 and 2004, residual delta smelt growth was low from the Sacramento-San Joaquin confluence through Suisun Bay relative to other parts of the upper estuary (Bennett 2005, p. 48). Delta smelt collected in 2005 from the Sacramento-San Joaquin confluence and Suisun Bay also had high incidence of liver glycogen depletion, a possible indicator of food limitation, exacerbated by unusually high water temperatures (Bennett *et al.* 2008, p. 16). Similarly, during 2003 and 2004 striped bass condition factor decreased in a seaward direction from the Delta through Suisun Bay. Moreover, there has been a significant long-term decline in primary productivity in the Suisun Bay region and the lower Delta (Jassby *et al.* 2002, p. 703; Kimmerer *et al.* 2012, p. 924). Recent studies (Bennett *et al.* 2008, pp. 18–20; Winder and Jassby 2010, p. 686) suggest that summer food limitation constitutes a major stressor on delta smelt (Baxter *et al.* 2010, p. 49). Therefore, the working hypothesis is that the poor fish growth and condition in the upper estuary are due to food limitation (Baxter *et al.* 2010, p. 50).

Egeria densa and other non-native submerged aquatic vegetation (e.g., *Myriophyllum spicatum*) may affect delta smelt in direct and indirect ways. Directly, submerged aquatic vegetation can overwhelm littoral habitats (inter-tidal shoals and beaches) where delta smelt may try to spawn, making them unsuitable for spawning or increasing the risk of predation for delta smelt or their eggs. Indirectly, submerged aquatic vegetation decreases turbidity by trapping suspended sediment, which has contributed to a decrease in both juvenile and adult smelt habitat quality and a contraction of delta smelt distribution in the upper estuary (Feyrer *et al.* 2007, p. 728; Nobriga *et al.* 2008, pp. 8–9). First-feeding delta smelt larvae require relatively turbid waters to capture

prey (Baskerville-Bridges *et al.* 2004, p. 223). Clearer water may also make delta smelt more susceptible to predation (Ferrari *et al.* 2014, p. 86).

In summary, we find that the overbite clam and other introduced species have altered the Delta food web and constitute a threat to delta smelt. It is likely that impacts to delta smelt from introduced species will continue.

Contaminants

In 2014, over 21 million pounds of pesticides were applied within the five-county Bay-Delta area, and Bay-Delta waters are listed under the Clean Water Act section 303(d) as impaired for several legacy and currently used pesticides (California Department of Pesticide Regulation 2016, p. 1). Concentrations of dissolved pesticides vary in the Delta both temporally and spatially (Kuivila 1999, entire). Several areas of the Delta, particularly the San Joaquin River and its tributaries, are impaired due to elevated levels of diazinon and chlorpyrifos, which are toxic at low concentrations to some aquatic organisms (MacCoy *et al.* 1995, pp. 21–30). Several studies have demonstrated the acute and chronic toxicity of two common insecticides, diazinon and esfenvalerate, in fish species (Barry *et al.* 1995, p. 273; Goodman *et al.* 1979, p. 479; Holdway *et al.*; 1994, p. 169; Scholz *et al.* 2001, p. 1911; Tanner and Knuth 1996, p. 244). The effects to delta smelt can be direct or indirect (effects that reduce the food supply of the delta smelt).

Pyrethroid insecticides are of particular concern because of their widespread use, and their tendency to be genotoxic (DNA damaging) to fishes at low doses (in the range of micrograms per liter) (Campana *et al.* 1999, p. 159). The pyrethroid esfenvalerate is associated with delayed spawning and reduced larval survival of bluegill sunfish (*Lepomis macrochirus*) (Tanner and Knuth 1996, pp. 246–250) and increased susceptibility of juvenile Chinook salmon (*Oncorhynchus tshawytscha*) to disease (Clifford *et al.* 2005, pp. 1770–1771). In addition, pyrethroids may interfere with nerve cell function, which could eventually result in paralysis (Bradbury and Coats 1989, pp. 377–378; Shafer and Meyer 2004, pp. 304–305).

Indirect effects to delta smelt through the food web have been documented. Weston and Lydy (2010, p. 1835) found the largest source of pyrethroids flowing into the Delta to be coming from the Sacramento Regional Wastewater Treatment Plant (SRWTP), where only secondary treatment occurs. Their data not only indicate the presence of these contaminants, but the concentrations found exceeded acute toxicity thresholds for the amphipod *Hyalella azteca*. Another study of storm events in five urban creeks in Suisun Marsh in February 2014 detected concentrations of bifenthrin (a pyrethroid) and fipronil outside of the primary agricultural pesticide season. Although the concentrations of these two insecticides were not high enough to cause mortality, they did result in paralysis of either the amphipod *Hyalella azteca*, or the larval midge *Chironomus dilutus* in 70% of the samples collected (Weston *et al.* 2015, pp. 20–25). Toxicity values for estuarine and marine invertebrates are not known for these insecticides but this study raises concerns about the effects to estuarine invertebrates from urban stream insecticide loading. This is of substantial concern because the use of insecticides in the urban environment had not before been considered the primary source of insecticides flowing into the Delta. Furthermore, this was not the case for the Stockton Wastewater Treatment facility, where tertiary treatment occurs, suggesting that different treatment methods may remove or retain pyrethroids differently (Baxter *et al.* 2010, p. 33).

Ammonia loading in the Bay-Delta has increased significantly in the last 25 years (Jasby 2008, p. 15–16). Effects of elevated ammonia levels on fish range from irritation of skin, gills, and eyes to reduced swimming ability and mortality (Wicks *et al.* 2002, p. 67). Delta smelt have shown direct sensitivity to ammonia at the larval and juvenile stages (Werner *et al.* 2008, pp. 85–88). Connon *et al.* (2011, pp. 347–375) investigated the sublethal effects of ammonia exposure on the genes of juvenile delta smelt and found that ammonia altered gene transcription including specific genes related to cell membrane integrity, energy metabolism, and cellular responses to environmental stimuli. The study supports the possibility of ammonia exposure-induced cell membrane destabilization that would affect membrane permeability and thus enhance the uptake of other contaminants. Ammonia also can be toxic to several species of copepods important to larval and juvenile fishes (Werner *et al.* 2010, pp. 78–79; Teh *et al.* 2011, pp. 25–27).

In addition to direct effects on fish, ammonia in the form of ammonium is thought to reduce primary production by diatoms, an important kind of phytoplankton, because ammonium inhibits the ability of diatoms to take nitrate out of the water. Diatoms grow faster using nitrate than ammonium. Ammonium in the estuary has been shown to suppress spring phytoplankton blooms in Suisun and Grizzly Bays by slowing down the growth rates of diatoms (Dugdale *et al.* 2007, pp. 26–28; Parker *et al.* 2012, pp. 6–8). However, Kimmerer *et al.* 2014 (p. 1214) found no direct evidence of ammonium effects on phytoplankton production within the LSZ from 2006–2008. A recent study conducted found no evidence that ammonium was inhibiting diatom production in Pacheco Slough in the fall (Esparza *et al.* 2014, p. 198). There were several differences between this recent research and past research. The Pacheco Slough study was conducted from August to October as opposed to the Wilkerson study that started in November, it measured biomass and not growth rates, and it took place in a slough rather than in open waters.

The role of ammonium nitrogen uptake inhibition in Sacramento River primary production is not fully understood. Parker *et al.* (2012, pp. 577–580) observed primary production in the Sacramento River decreased in the SRWTP region as compared to the upper river region during the months of March and April. However, a previous study found that declines in phytoplankton density above the SRWTP between the Tower Bridge in Sacramento and Garcia Bend are a possible cause of this decline in productivity (Foe *et al.* 2010, p. 13). The application of general ecological principles would lead us to believe that decreased primary productivity, wherever it occurs in delta smelt habitat, is likely to lead to a decrease in copepods and other zooplankton that delta smelt rely upon for food. A link between primary productivity and productivity in higher trophic levels has been documented in various pelagic food webs (Nixon 1988, p. 1019; Sobczak *et al.* 2005, p. 133). At this time, we conclude that more science is needed to determine the role of ammonium in the food web. However, because ammonium may be affecting the food web as shown in research described above, we support future actions in the Bay-Delta that would reduce ammonium outputs.

Selenium, introduced into the estuary primarily from agricultural irrigation runoff via the San Joaquin River drainage and oil refineries in San Francisco Bay, has been implicated in toxic and reproductive effects in fish and wildlife (Linville *et al.* 2002, p. 52). Selenium exposure has been shown to have effects on some benthic foraging species; deformities typical of selenium exposure including lordosis (spinal deformities) have been observed in splittail collected from Suisun Bay (Stewart *et al.* 2004, p. 4524); however, there is no evidence that selenium exposure

is contributing to the decline of delta smelt or other pelagic species in the Bay-Delta (Baxter *et al.* 2010, p. 28).

Complex mixtures of contaminants spanning many different classes can be common in regions heavily influenced by agricultural or urban environments. To date, a variety of studies have documented the impacts of complex chemical mixtures on aquatic organisms. Laetz *et al.* (2009, p. 351) exposed juvenile Coho salmon (*Oncorhynchus ktsutch*) to a sub-lethal concentrations of five current-use pesticides and found the compounds were acting as synergists with each other. Nørgaard and Cedergreen (2010, p. 962) found that a mixture of fungicides and pyrethroids could produce a 12-fold increase in toxicity over what was expected using an additive model when looking at impacts to *Daphnia magna*. LeBlanc *et al.* (2012, p. 383) examined the sub-lethal effects of three pesticides: chlorpyrifos, dimethoate (both organophosphate) and imidacloprid (neonicotinoid), and found a synergistic interaction when aquatic invertebrates (*Chironomus dilutus* larvae) were exposed to all three at once. Carvalho *et al.* (2014, pp. 225–228) produced two mixtures of 14 and 19 different compounds of concern, including metals, pesticides, pharmaceuticals, and hydrocarbons, all at concentrations below the Environmental Quality Standards. A host of sub-lethal impacts to many different aquatic organisms were detected including fish embryo toxicity, increased oxidative stress, and decreased invertebrate mobility.

Contaminants are suspected to be a stressor on delta smelt despite little direct evidence (Kuivila and Moon 2004, pp. 237–241; Brooks *et al.* 2012, pp. 611–614). A study of juvenile delta smelt in five different regions encompassing their range examined fish for signs of contaminants and food limitation. The histopathological analysis of 244 fish sampled in 2012 and 2013 found an 11-fold increase in gill and liver lesion scores in Cache Slough as compared to Suisun Marsh. Higher lesion scores indicate less healthy tissues and are indicative of contaminant-related stress (Hammock *et al.* 2015, p. 320).

Large blooms of toxic *Microcystis aeruginosa* (a species of cyanobacteria) were first documented in the Bay-Delta during the summer of 1999 (Lehman *et al.* 2005, p. 87). *M. aeruginosa* forms large colonies throughout most of the Delta and increasingly down into Suisun Bay (Lehman *et al.* 2005, p. 92; 2013, p. 150). Blooms typically occur when water temperatures are above 20 °C (68 °F) (Lehman *et al.* 2010a, p. 238). It is unclear whether microcystins and other toxins produced by local blooms are acutely toxic to fishes at current concentrations; however, the toxins accumulate in fish and their prey. During summer 2005, Age-0 striped bass and Mississippi silversides that were co-occurring with the *Microcystis* bloom showed various forms of liver damage (Lehman *et al.* 2010a, p. 241). When ingested with food, microcystins have been experimentally shown to cause substantial impairment of health in threadfin shad (Acuña *et al.* 2012, p. 1195). In addition, the copepods that delta smelt eat are particularly susceptible to these toxins (Ger 2008, pp. 12, 13; Ger *et al.* 2010, p. 1554). An investigation of food web effects and fish toxicity concluded that even at low abundances, *M. aeruginosa* may impact estuarine fish productivity through both toxicity and food web impacts (Lehman *et al.* 2010, p. 241–245). *M. aeruginosa* is most likely to affect juvenile delta smelt during summer blooms.

Vulnerability of Small Populations

Delta smelt are relatively concentrated in their rearing habitat during the fall, making them vulnerable to environmental conditions such as droughts, contaminant spills, and predation. Small, isolated populations are more likely to lose genetic variability due to genetic drift, and to suffer inbreeding depression due to the fixation of deleterious alleles (gene variants) (Lande 1998, pp. 11–17). Populations at low densities are often subject to Allee effects, which involve decreases in the ratio of offspring to adults as the population density decreases (Dennis 2002, p. 389). It is unknown if small population size has contributed to delta smelt's decline. It was recently documented that delta smelt's genetic effective population size declined between 2003 and 2009 as the population declined (Fisch *et al.* 2011, p.7, Fig 2). This study estimated that the effective population size declined from about 4,000–12,000 fish in 2003 to 1,000–2,000 fish during 2007–2009. The effective population size is an estimate of the number of adult fish that produced offspring that survived to adulthood in any given year. It is thought that the population experienced a genetic bottleneck during the 2007-2009 time period and may have experienced subsequent bottlenecks since that time. It is likely that the population is currently experiencing a genetic bottleneck, with a population estimate of 13,000 fish for 2016. This loss of genetic diversity in the population reduces the ability for fish to respond to changes and evolve, and acts as a threat to the delta smelt population.

Summary for Factor E

Based on a review of the best scientific and commercial information available, we find that the following natural or manmade factors pose primary ongoing threats to the delta smelt: entrainment by the State and Federal water export facilities, and introduced species, primarily the overbite clam and *Egeria densa*. Additional threats include contaminants and small population size. Ammonia in the form of ammonium may also constitute a threat to the delta smelt.

CONSERVATION MEASURES PLANNED OR IMPLEMENTED

A variety of conservation measures have been proposed and/or undertaken in the estuary that benefit the delta smelt. The following list is not exhaustive.

Suisun Marsh Habitat Management, Preservation and Restoration Plan: The Suisun Marsh Plan, signed in 2014, was developed to balance the goals and objectives of the Bay-Delta Program, Suisun Marsh Preservation Agreement and other management and restoration programs within Suisun Marsh. The Plan provides for simultaneous protection and enhancement of Pacific Flyway and existing wildlife values in managed wetlands, endangered species recovery, and water quality. The Plan addresses water quality, fisheries, wildlife, vegetation, special-status species, land use, land use development patterns, population, housing, economics, and public services (fire protection, vector control), cultural resources, air quality, noise, recreation, energy, visual impacts, and socioeconomic condition. (Suisun Marsh Habitat Restoration Plan 2011, entire). The Suisun Marsh Plan will benefit the delta smelt by restoring areas that are key to the species habitat requirements.

Ecosystem Restoration Program (ERP): The goal of the ERP is to improve and increase aquatic and terrestrial habitats and improve ecological functions in the Bay-Delta to support sustainable populations of diverse and valuable plant and animal species. ERP began as a CALFED program, but is now housed within the CDFW. The ERP has operated mainly upstream of the

Delta to protect, restore and enhance aquatic and terrestrial habitat and improve fish passage. The Delta Regional Ecosystem Restoration Implementation Plan (DRERIP) is one of four regional plans intended to implement the ERP. The DRERIP has produced a series of models that have informed agencies about processes, habitats, species, and stressors of the Bay-Delta. The ERP issued a revised draft Conservation Strategy for the Restoration of the Sacramento-San Joaquin Delta and the Sacramento and San Joaquin Valley Regions in February of 2013. The CDFW and the CDWR are continuing to implement and plan for ecosystem restoration projects begun under the CALFED Bay-Delta Program located in Suisun Marsh, at Dutch Slough, at Cache Slough, in the Yolo Bypass, and at the Cosumnes Preserve's North Delta project. These ecosystem restoration projects are expected to benefit the delta smelt.

Recovery Plan for Tidal Marsh Ecosystems of Northern and Central California: This multi-species recovery plan, published in 2014, addresses conservation needs for the San Francisco Estuary, with a focus on the following listed plant and terrestrial species: *Cirsium hydrophilum* var. *hydrophilum* (Suisun thistle), *Cordylanthus mollis* ssp. *Mollis* (soft bird's beak), *Suaeda californica* (California sea-blite), California clapper rail (*Rallus longirostris obsoletus*), and salt marsh harvest mouse (*Reithrodontomys raviventris*). Restoration efforts from this plan are identified in the implementation table of this recovery plan. One of the actions in the plan is to restore 5,000 acres of high quality marsh habitat within the Suisun Bay Recovery Unit. Restoration of this area is expected to help the delta smelt by increasing food web productivity and generally benefiting the habitat of the delta smelt.

Interagency Ecological Program: The mission of the IEP is, in collaboration with others, to provide ecological information and scientific leadership for use in management of the San Francisco Estuary. The goals of IEP are to: describe the status and trends of aquatic ecological factors of interest in the estuary; develop an understanding of environmental factors that influence observed aquatic ecological status and trends; use knowledge of the above information in a collaboration process to support natural resource planning, management, and regulatory activities in the estuary; continually reassess and enhance long-term monitoring and research activities that demonstrate scientific excellence; and to provide scientific information about the estuary that is accurate, accessible, reliable, and timely. The IEP studies benefit the delta smelt by providing information that will help with the conservation of the species.

Summary of Threats

This status review identified threats to the Bay-Delta DPS of delta smelt attributable to Factors A, D, and E, as well as interactions between these threats. The primary known threats cited in the 2010 delta smelt uplisting document are: entrainment by State and Federal water export facilities (Factor E); summer and fall increases in salinity due to reductions in freshwater flow and summer and fall increases in water clarity (Factor A), and effects from introduced species, primarily the overbite clam and *Egeria densa* (Factor E). Additional threats included predation (Factor C), entrainment into power plants (Factor E), contaminants (Factor E), and small population size (Factor E). Since the 2010 proposed up-listing, we have identified climate change as a threat in the 2012 Candidate Notice of Review. Climate change was not analyzed in the 2010 up-listing document. Since the 2010 up-listing, one of the two power plants within the range of the delta smelt using water for cooling has shut down and power plants are no longer thought to be a threat to the population as a whole. We have identified a number of existing

regulatory mechanisms that provide protective measures that affect the stressors acting on the Delta smelt. Despite these existing regulatory mechanisms and other conservations efforts, the stressors continue to act on the species such that it is warranted for uplisting under the Act.

Upstream dams and water storage, exacerbated by water diversions, especially from the SWP and CVP water export facilities, result in reduced freshwater flows within the estuary, and these reductions in freshwater flows result in reduced habitat suitability for delta smelt by moving the position of X2 in the estuary. First-feeding delta smelt larvae require relatively turbid waters to capture prey and delta smelt may also use turbidity as cover from predators. The increased water clarity in delta smelt rearing habitat in recent decades is attributed to the interruption of sediment transport by upstream dams and the spread of the exotic introduced water plant *Egeria densa* (Brazilian waterweed), which traps suspended sediments. Increased water clarity is therefore a threat to delta smelt. Models indicate a steady log-linear decline in abundance of delta smelt since about the time of the invasion of the nonnative overbite clam in 1987 (Thomson et al. 2010, p. 1442; see Factor E: Introduced Species) in the Bay-Delta. The long-term decline in abundance of delta smelt in the Bay-Delta has been partially attributed to reductions in food availability caused by establishment of the nonnative overbite clam in 1987 (Factor E) and possibly by ammonium concentrations (Factor E) and water diversions (Factor A).

Delta smelt is a rare fish and has been a rare fish (compared to other species) for at least the past several decades (Nobriga and Herbold 2008). Therefore, it has also been rare in examinations of predator stomach contents as would be expected and it is not surprising that studies are inconclusive on the extent of population level effects. Current and historical evidence of delta smelt in stomachs of the relatively small number of non-native fishes sampled compared to overall non-native population numbers shows that predation from non-native fishes is likely having some effect on overall population numbers of delta smelt. It is likely that the population is currently experiencing a genetic bottleneck with a population estimate of 13,000 fish for 2016. This loss of genetic diversity in the population reduces the ability for fish to respond to changes and evolve and acts a threat to the delta smelt population.

The threats identified are likely acting together to contribute to the decline of the population (Baxter et al. 2010, p. 69). Reduced freshwater flows result in effects to delta smelt habitat suitability, at the same time that the food web has been altered by introduced species and ammonium concentrations. Climate change will likely exacerbate these threats. The combined effects of reduced freshwater flows, the invasive overbite clam (reduced levels of phytoplankton and zooplankton that are important to the Bay-Delta food web), entrainment, predation, small population size, and contaminants act to significantly degrade conditions for delta smelt.

The best scientific and commercial information available indicates that the threats facing the delta smelt are of sufficient imminence, intensity, and magnitude to endanger the continued existence of the species.

In 2010, we completed a 12-month finding for delta smelt in which we determined a change in status from threatened to endangered was warranted. The continuing downward trend in delta smelt abundance indices supports that finding. Eleven of the last twelve years have seen FMWT indexes that have been the lowest ever recorded. 2015-2016 results from all four of the surveys analyzed in this review have been the lowest ever recorded for the delta smelt. A new 2016

population estimate of 13,000 individuals with 95% confidence intervals ranging between 6,000 and 28,000 individuals (Newman 2016, unpublished data) is also the lowest ever estimated declining sharply from the 2015 estimate of 112,000 individuals. Although conservation measures are in place to protect the species including the 2008 Biological Opinion, these measures have not been sufficient to halt the decline of the species. Therefore, based on a review of the best scientific and commercial information available, we find that the delta smelt still meets the definition of an endangered species under the Act, and that it warrants reclassification from threatened to endangered. However, at this time, the promulgation of a formal rulemaking to reclassify delta smelt is precluded by higher priority actions.

Recommended Conservation Measures

Increasing Delta outflows so that they more closely approximate unimpaired flows in the watershed would address several needs of the delta smelt, likely improving its habitat quality and quantity and by extension its reproduction and survival. Furthermore, increased winter and spring flows may reduce water clarity, which would increase habitat quality for delta smelt. Contaminant reduction within the Bay-Delta could improve primary and secondary productivity while at the same time limiting toxicity exposure to delta smelt. The reduction of pesticides entering the Delta could also improve habitat conditions and fish health. Reducing ammonia loads may also help to increase primary productivity within the Bay-Delta, contributing to better growth and survival. Therefore, the FWS recommends higher outflows and the reduction of contaminants entering the estuary.

LISTING PRIORITY

As a result of our analysis of the best available scientific and commercial information, we have assigned the delta smelt a Listing Priority Number of 2, based on high magnitude and immediacy of threats. While we conclude that reclassifying the species as endangered is still warranted, an immediate proposal to reclassify this species is precluded by other higher priority actions.

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/Population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/Population	6
Moderate to Low	Imminent	Monotype genus	7

		Species	8
		Subspecies/Population	9
	Non-Imminent	Monotype genus	10
		Species	11
		Subspecies/Population	12

Rationale for listing priority number:

Magnitude:

The magnitude of threats is high due to the number and severity of ongoing threats. These threats include turbidity changes, entrainment and invasive species. Ammonium may also be a threat to the survival of the delta smelt. The ecology and biology of the San Francisco Bay-Delta has changed drastically over the last 160 years. Although a number of conservation measures have been put in place to protect the delta smelt and its habitat, the population continues to decline. Turbidity changes due to levees and dams and increased ammonium concentrations have taken place throughout the range of the delta smelt. Changes in the position of the LSZ in the Bay-Delta have altered foraging and breeding habitat. Although this threat does not extend throughout the range of the delta smelt, it does encompass areas that are key to the delta smelt's survival, including Suisun Marsh and Suisun Bay. Delta smelt numbers have dwindled to a small fraction of what they were before these changes took place. Stress from water pumping operations and invasive species is expected to continue into the future as water demands for the growing population in California continue to increase.

Imminence

The threats discussed above are ongoing and likely to continue into the future. We therefore consider threats to be imminent.

Have you promptly reviewed all of the information received regarding the species for the purpose of determining whether emergency listing is needed? Yes

Is Emergency Listing Warranted? No

DESCRIPTION OF MONITORING

The Interagency Ecological Program (IEP) was created by State and Federal resource agencies to focus scientific inquiry on the effects of CVP and SWP water operations on the aquatic resources of the Bay-Delta. The IEP initiated the Pelagic Organism Decline (POD) study effort in 2005 to focus scientific effort and resources on the most recent and precipitous declines in abundance of

several species, including delta smelt. Delta smelt are regularly captured in the IEP’s monitoring surveys. IEP monitoring and research includes effects of contaminants, invasive species, export pump entrainment and freshwater outflow on delta smelt biology. Species experts from universities, state and federal governments, and non-profits work cooperatively to yield crucial information on the status of the species through the IEP program. The primary focus of the IEP is on monitoring status and trends within the estuary and on water project effects on the estuary’s ecology. The Service funds a delta smelt research biologist position at the Stockton Fish and Wildlife Office to investigate potential threats to the species. Existing funding comes from the Service and from CalFed agencies and through grants from the Delta Science Program. The Service is presently cooperating with scientists from the University of California-Davis and the California Department of Water Resources to develop two genetic refuge populations of delta smelt and a delta smelt culture facility. In addition to its routine monitoring programs, the IEP is currently funding or conducting research into delta smelt vitality rates, population genetics, aquaculture and tagging/marking techniques, vulnerability to sampling gear, and large conceptual and mathematical summaries of delta smelt abundance trends and population dynamics.

Figure 1: Monitoring surveys from which Delta smelt abundance trends are derived.

Survey	Agency Lead	Target Species	Season of Sample	Frequency	Application of Data
Spring Kodiak Trawl (SKT)	CDFW	Delta smelt	Jan–May	Monthly	Adult spawning survey, distribution and relative abundance
Smelt Larva Survey (SLS)	CDFW	Longfin Smelt*	Jan–March	Biweekly	Distribution and relative abundance of smelt larvae
20mm Survey	CDFW	Delta smelt	March–July	Biweekly	Larval–juvenile survey, distribution and relative abundance
Summer Townet Survey (TNS)	CDFW	Striped Bass**	June–Aug	Biweekly	Distribution and relative abundance
Fall Midwater Trawl (FMWT)	CDFW	Striped Bass**	Sept–Dec	Monthly	Annual delta smelt abundance index calculation

* The SLS was originally a delta smelt-targeted pilot study, with a study design that changed from year to year. In 2009, the SLS was redesigned to collect longfin smelt distribution data for the purpose of providing recommendations for water operations in the Delta.

** The FMWT and TNS were originally implemented to monitor distribution and abundance of striped bass. Because these surveys also collected information on the distribution and abundance of delta smelt, they are both mandated by USFWS in its State and Federal water operations BO.

COORDINATION WITH STATES

The Delta smelt is known only from California. Therefore, coordination is done with the State of California. Much of the coordination is done through the IEP and includes research and abundance surveys (See Description of Monitoring above).

LITERATURE CITED

Aasen, G. 2014. CVP and SWP exports 1981-2013. Powerpoint slide.

Acuña, S, D. Baxa, S. Teh. 2012. Sublethal dietary effects of microcystin producing *Microcystis* on threadfin shad, *Dorosoma petenense*. *Toxicon* 60: 1991-1202.

Alpine, A.E. and J.E. Cloern. 1992. Trophic interactions and direct physical effects control phytoplankton biomass and production in an estuary. *Limnol. Oceanogr.* 37: 946-955

Armor, C., R. Baxter, B. Bennett, R. Breuer, M. Chotkowski, P. Coulston, D. Denton, B. Herbold, W. Kimmerer, K. Larsen, M.L. Nobriga, K. Rose, T. Sommer, and M.T. Stacey. 2005. Interagency Ecological Program synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Draft Report. Prepared for U.S. Bureau of Reclamation. 55 pages. IEP.

Arthur, J.F., M.D. Ball. 1979. Factors Influencing the Entrapment of Suspended Material in the San Francisco Bay-Delta Estuary. USBR, 2800 Cottage Way, Sacramento CA, 95825. 33 pp.

Arthur, J.F., M.D. Ball and S.Y. Baughman. 1996. Summary of federal and state water project impacts in the San Francisco Bay-Delta estuary, California. pp 445-495 *in* San Francisco Bay: the ecosystem, J.T. Hollibaugh, ed., AAAS, San Francisco, CA, USA.

Barry, M.J., D.C. Logan, J.T. Ahokas, and D.A. Holdway. 1995. Effect of Algal food concentration on toxicity of two agricultural pesticides to *Daphnia carinata*. *Ecotoxicology and Environmental Safety* 32:273-279.

Baskerville-Bridges, B., J.C. Lindberg and S.I. Doroshov. 2004. The effect of light intensity, alga concentration, and prey density on the feeding behavior of delta smelt larvae. pp 219-228 *in* Early life history of fishes in the San Francisco Estuary and watershed, F. Feyrer, L. Brown, R. Brown and J. Orsi, eds. American Fisheries Society Symposium 39, Bethesda, MD.

Baxter, R., R. Breuer, L. Brown, L. Conrad, F. Feyrer, S. Fong, K. Gehrts, L. Grimaldo, B. Herbold, P. Hrodey, A. Mueller-Solger, T. Sommer and K. Souza. 2010. Interagency ecological program pelagic organism decline work plan and synthesis of results. Interagency Ecological Program, Sacramento, CA 259 pp
<http://www.water.ca.gov/iep/docs/FinalPOD2010Workplan12610.pdf>

Baxter, R., L.R. Brown, G. Castillo, L. Conrad, S. Culberson, M. Dekar, M. Dekar, F. Feyrer, T. Hunt, K. Jones, J. Kirsh, A. Mueller-Solger, M. Nobriga, S.B. Slater, T. Sommer, K. Souza, 2015. An updated conceptual model of Delta Smelt biology: our evolving understanding of an

estuarine fish. Interagency Ecological Program. Management, Analysis and Synthesis Team. Technical Report 90. January, 2015.

Baerwald, M.R., B.M. Schreier, G. Schumer and B. May. 2012. Detection of threatened delta smelt in the gut contents of the invasive Mississippi silverside in the San Francisco estuary using TaqMan assays. *Transactions of American Fisheries Society* 141: 1600-1607.

Bennett, W.A. 2005. Critical assessment of the delta smelt population in the San Francisco Estuary, California. *San Francisco Estuary and Watershed Science* Vol. 3, Issue 2 (September 2005), Article 1. <http://repositories.cdlib.org/jmie/sfewsvol3/iss2/art1>

Bennett, W.A., Burau, J.R. 2014. Riders on the Storm: Selective Tidal Movements Facilitate the Spawning Migration of Threatened Delta Smelt in the San Francisco Estuary. *Estuaries and Coasts*. DOI 10.1007/s12237-014-9877-3.

Bennett, W.A., and P.B. Moyle. J.T. Hollibaugh, editor. 1996. Where have all the fishes gone? Interactive factors producing fish declines in the Sacramento-San Joaquin Estuary. *San Francisco Bay: The ecosystem: further investigations into the natural history of San Francisco Bay and Delta with reference to the influence of man*. 26 pp. American Association for the Advancement of Science, Romsberg Tiburon Center, San Francisco State University.

Bennett, W.A., J.A. Hobbs, and S.J. Teh. 2008. Interplay of environmental forcing and growth-selective mortality in the poor year-class success of delta smelt in 2005. Final Report. Prepared for The Pelagic Organism Decline Management Team. 30 pages.

Bouley, P., and W. Kimmerer. 2006. Ecology of a highly abundance, introduced cyclopoid copepod in a temperate estuary. *Marine Ecology Progress Series* 324:219-228.

Bradbury, S.P. and J.R. Coats. 1989. Toxicokinetics and toxicodynamics of pyrethroid insecticides in fish. *Environmental Toxicology and Chemistry* 8:373-380.

Brooks, M. L., E. Fleishman, L.R. Brown, P.W. Lehman, I. Werner, N. Scholz, S. van Drunick, J. Drever, D. Stoms, A. Parker, R. Dugdale. 2012. Life histories, salinity zones, and sublethal contributions of contaminants to pelagic fish declines illustrated with a case study of San Francisco Estuary, California, USA. *Estuaries and Coasts* 35(2): 603-621.

Brown, L. R., and D. Michniuk. 2007. Littoral fish assemblages of the alien-dominated Sacramento-San Joaquin Delta, California 1980-1983 and 2001-2003. *Estuaries and Coasts* 30(1):186-200.

Brown, L. R., W. A. Bennett, R. W. Wagner, T. Morgan-King, N. Knowles, F. Feyrer, D. H. Schoellhamer, M. T. Stacey, and M. Dettinger. 2013. Implications for future survival of delta smelt from four climate change scenarios for the Sacramento-San Joaquin Delta, California. *Estuaries and Coasts* 36 (4):754-774.

Brown, L.R., L.M. Komoroske, R.W. Wagner, T. Morgan-King, J.T. May, R.E. Connon. 2016.

Coupled Downscaled Climate Models and Ecophysiological Metrics Forecast Habitat Compression for an Endangered Estuarine Fish. PLoS ONE 11(1): e0146724. doi:10.1371/journal.pone.0146724

[CDFG] California Department of Fish and Game. 2001. True Smelts. In: California's Living Marine Resources: A Status Report (eds.) W.S. Leet, C.M. Dewees, R. Klingbiel, & E.J. Larson. pp. 472-479.

California Department of Fish and Wildlife. 2016. Survey results from webpage. <https://www.wildlife.ca.gov/Regions/3> <Accessed on December 24, 2014>.

California Department of Pesticide Regulation. 2016. CDPR website accessed on 04/12/16: http://www.cdpr.ca.gov/docs/pur/pur14rep/lbsby_co_14.pdf

[CRWQCB] California Regional Water Quality Control Board. 2014. Waste Water Discharge Requirements for the Sacramento Regional County Sanitation District Sacramento Regional Wastewater Treatment Plant. 221 pp.

Campana, M.A., A.M. Panzeri, V.J. Moreno and F.N. Dulout. 1999. Genotoxic evaluation on the pyrethroid lambda-cyhalothrin using the micronucleus test in erythrocytes of the fish *Cheirodon interruptus interruptus*. Mutation Research 438:155-161.

Castillo, G., J. Morinaka, J. Lindberg, R. Fujimura, B. Baskerville-Bridges, J. Hobbs, G. Tigan, L. Ellison. 2012. Pre-Screen Loss and Fish Facility Efficiency for Delta Smelt at the South Delta's State Water Project, California. San Francisco Estuary and Watershed Science. 25 pp.

Clifford, M.A., K.J. Eder, I. Werner and R. Hedrick. 2005. Synergistic effects of esfenvalerate and infectious hematopoietic necrosis virus on juvenile Chinook salmon mortality. Environmental Toxicology and Chemistry 24(7):1766-1772.

Cloern, J.E. 1987. Turbidity as a Control on Phytoplankton Biomass and Productivity in Estuaries. Continental Shelf Research 7: 1367-1381.

Cloern, J. E., and A. D. Jassby. 2012. Drivers of change in estuarine-coastal ecosystems: Discoveries from four decades of study in San Francisco Bay. Reviews of Geophysics 50(4):1-33.

Cloern, J. E., N. Knowles, L. R. Brown, D. Cayan, M. D. Dettinger, T. L. Morgan, D. H. Schoellhamer, M. T. Stacey, M. van der Wegen, R. W. Wagner, and A. D. Jassby. 2011. Projected evolution of California's San Francisco Bay-Delta-river system in a century of climate change. PLoS One 6(9):e24465.

Connon, R. E., L. A. Deanovic, E. B. Fritsch, L. S. D'Abronzio, and I. Werner. 2011. Sublethal responses to ammonia exposure in the endangered delta smelt; *Hypomesus transpacificus* (Fam. Osmeridae). Aquat Toxicol 105(3-4):369-377.

- Culberson, S.D., C.B. Harrison, C. Enright, and M.L. Nobriga. 2004. Sensitivity of larval fish transport to location, timing, and behavior using a particle tracking model in Suisun Marsh, California. *American Fisheries Society Symposium* 39:257-267.
- Dege, M. and L.R. Brown. 2004. Effect of outflow on spring and summertime distribution and abundance of larval and juvenile fishes in the upper San Francisco Estuary. pp *in* Early life history of fishes in the San Francisco Estuary and watershed, F. Feyrer, L. Brown, R. Brown and J. Orsi, eds. *American Fisheries Society Symposium* 39, Bethesda, MD. F. Feyrer, L. Brown, R. Brown and J. Orsi, eds. *American Fisheries Society Symposium* 39, Bethesda, MD.
- Dennis, B. 2002. Allee effects in stochastic populations. *Oikos* 96(3):389-401.
- Dettinger, M. 2005. From Climate-Change Spaghetti to Climate-Change Distributions for 21st Century California. *San Francisco Estuary and Watershed Science* 3(1):Article 4. 14.
- Dugdale, R.C., F.P. Wilkerson, V.E. Hogue, and A. Marchi. 2007. The role of ammonia and nitrate in spring bloom development in San Francisco Bay. *Estuarine, Coastal and Shelf Science* 73:17-29.
- Erkkila, L.F., J.F. Moffett, O.B. Cope, B.R. Smith and R.S. Nelson. 1950. Sacramento-San Joaquin Delta fishery resources: effects of Tracy pumping plant and delta cross channel. U.S. Fish and Wildl. Svc. Spec. Sci. Rpt. Fisheries 56. 109 pp
- Esparza, M. L., A. E. Farrell, D. J. Craig, C. Swanson, B. S. Dhaliwal, and G. M. Berg. 2014. Impact of atypical ammonium concentrations on phytoplankton abundance and composition in fresh versus estuarine waters. *Aquatic Biology* 21(3):191-204.
- Ferrari, M.C.O., L. Ranaker, K.L. Weinersmith, M.J. Young, A. Sih, J.L. Conrad. 2014. Effects of turbidity and an invasive waterweed on predation by introduced largemouth bass. *Environ Biol Fish* 97: 79–90.
- Feyrer, F., B. Herbold, S.A. Matern and P.B. Moyle. 2003. Dietary shifts in a stressed fish assemblage: consequences of a bivalve invasion in the San Francisco Estuary. *Environmental Biology of Fishes* 67:277-288.
- Feyrer, F., M.L. Nobriga and T.R Sommer. 2007. Multidecadal trends for three declining fish species: habitat patterns and mechanisms in the San Francisco Estuary, California, USA. *Canadian Journal of Fisheries and Aquatic Science* 64:723-734.
- Feyrer, F., K. Newman, and M.L. Nobriga. 2011. Modeling the Effects of Future Outflow on the Abiotic Habitat of an Imperiled Estuarine Fish. *Estuaries and Coasts* 34(1):120-128.
- Fisch, K. M., J. M. Henderson, R. S. Burton, and B. May. 2011. Population genetics and conservation implications for the endangered delta smelt in the San Francisco Bay-Delta. *Conservation Genetics* 12(6):1421-1434.

Foe, C., A. Ballard, and S. Fong. 2010. Nutrient concentrations and biological effects in the Sacramento-San Joaquin Delta. 90 pages. California Regional Water Quality Control Board, Central Valley Region.

Ganguly, A.R., K. Steinhaeuser, D.J. Erickson III, M. Branstetter, E.S. Parish, N. Singh, J.B. Drake, and L. Buja. 2009. Higher trends but larger uncertainty and geographic variability in 21st century temperature and heat waves. Proceedings of the National Academy of Sciences. September 15, 2009, Vol. 106 no. 37:15555–15559.

Ganssle, D. 1966. Fishes and decapods of San Pablo and Suisun Bay. Ecological studies of the Sacramento-San Joaquin Estuary, Pt. 1, D.W. Kelley, ed., Calif. Dept. Fish Game Bull. 133: 64-94.

<http://content.cdlib.org/view?docId=kt4j49n6r5;NAAN=13030&doc.view=frames&chunk.id=d0e2097&toc.depth=1&toc.id=d0e2097&brand=calisphere>

Ger, K.A. 2008. Extent of acute, chronic and nutritional impacts of *Microcystis aeruginosa* blooms on the calanoid copepods of the upper San Francisco Estuary. Dissertation. University of California, Davis. 77 pp.

Ger, K.A., S.J. Teh, D.V. Baxa, S. Lesmeister, and C.R. Goldman. 2010. The effects of dietary *Microcystis aeruginosa* and microcystin on the copepods of the upper San Francisco Estuary. Freshwater Biology 55(7):1548–1559.

Gewant, D., S.M. Bollens. 2012. Fish Assemblages of Interior Tidal Marsh Channels in Relation to Environmental Variables in the Upper San Francisco Estuary. Environmental Biology Fish 94: 483-499.

Goodman, L.R., D.J. Hansen, D.L. Coppage, J.C. Moore and E. Matthews. 1979. Diazinon: Chronic toxicity to and brain acetylcholinesterase inhibition in, the sheepshead minnow, *Cyprinodon variegatus*. Transaction of the American Fisheries Society 108:479-488.

Grimaldo, L.F., T. Sommer, N. Van Ark, G. Jones, E. Holland, P. Moyle and P. Smith. 2009. Factors affecting fish entrainment into massive water diversions in a tidal freshwater estuary: can fish losses be managed? North American Journal of Fisheries Management 29: 1253-1270

Guinee, R. 2014. Personal communication between Roger Guinee of the U. S. Fish and Wildlife Service and Colin Grant of the U.S. Fish and Wildlife Service.

Hammock, B.G. J. A. Hobbs, S.B. Slater, S. Acuna and S.J. Teh. 2015. Contaminants and food limitation stress in an endangered estuarine fish. Science of the Total Environment 532: 316-326.

Hasenbein, M., L.M. Komoroske, R.E. Connon, J. Geist and N.A. Fangué. 2013. Turbidity and Salinity Affect Feeding Performance and Physiological Stress in the Endangered Delta Smelt. Integrative and Comparative Biology. August 14, 2013 edition. pp. 1-15.

Hasenbein, M., Fangué N.A., Geist J., Komoroske L.M., Truong J., McPherson R., Connon R.E. 2016. Assessments at multiple levels of biological organization allow for an integrative

determination of physiological tolerances to turbidity in an endangered fish species. *Conserv Physiol* 4: doi:10.1093/conphys/cow004.

Herren, J.R. and S. Kawasaki. 2001. Inventory of water diversions in four geographic areas in California's Central Valley. Contributions to the biology of Central Valley salmonids, Volume 2. R.L. Brown, Department of Fish and Game Fish Bulletin. 179: 343-355

Hobbs, J.A., W.A. Bennett, J. Burton. 2007. Classification of Larval and Delta Smelt to Nursery Areas by Use of Trace Elemental Fingerprinting. Transactions of the American Fisheries Society 136:518-527.

Holdway, D.A., M.J. Barry, D.C. Logan, D. Robertson, V. Young, J.T. Ahokas. 1994. Toxicity of pulse-exposed fenvalerate and esfenvalerate to larval Australian crimson-spotted rainbow fish (*Melanotaenia fluviatilis*). Aquatic Toxicology 28:169-187.

Horppila, J., A. Liljendahl-Nurminen, and T. Malinen. 2004. Effects of clay turbidity and light on the predator prey interaction between smelts and chaoborids. Canadian Journal of Fisheries and Aquatic Sciences 61(10):1862-1870.

Huber, M., and R. Knutti. 2011. Anthropogenic and natural warming inferred from changes in Earth's energy balance. Nature Geoscience Letters. Published online: 4 December 2011, DOI:10.1038/NCEO1327 (*Nature Geoscience* 5, 31-36 (2012)).

Ilves, K.L., E.B. Taylor. 2008. Evolutionary and Biogeographical patterns within the smelt genus *hypomesus* in the North Pacific Ocean. Journal of Biogeography. 35: 48-64.

Intergovernmental Panel on Climate Change [IPCC]. 2013a. Annex III: Glossary [Planton, S. (ed.)]. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. pg. 1450.

Intergovernmental Panel on Climate Change [IPCC]. 2013b. Summary for Policymakers. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. Pp. 3-29;

Intergovernmental Panel on Climate Change [IPCC]. 2014. Summary for policymakers. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, Pp. 1-32.

Jassby, A.D. 2008. Phytoplankton in the upper San Francisco Estuary: Recent biomass trends, their causes and their trophic significance. *San Francisco Estuary and Watershed Science* 6(1):Article 2: 24 pages.

Jassby, A.D., W.J. Kimmerer, S.G. Monismith, C. Armor, J.E. Cloern, T.M. Powell, J.R. Schubel and T.J. Vendlinski. 1995. Isohaline position as a habitat indicator for estuarine populations. *Ecol. Appl.* 5: 272-289.

Jassby, A.D., J.E. Cloern, and B.E. Cole. 2002. Annual primary production: Patterns and mechanisms of change in a nutrient-rich tidal ecosystem. *Limnology and Oceanography* 47(3):698-712.

Kimmerer, W.J., 1998. Report of the 1994 Entrapment Zone Study No. Technical Report 56, Interagency Ecological Program for the San Francisco Bay/Delta Estuary, Sacramento.

Kimmerer, W.J. 2002b. Physical, biological, and management responses to variable freshwater flow into the San Francisco Estuary. *Estuaries* 25:1275-1290

Kimmerer, W. 2004. Open water processes of the San Francisco Estuary: from physical forcing to biological responses. *SFEWS* 2(1) (February 2004), Article 1.
<http://repositories.cdlib.org/jmie/sfews/vol2/iss1/art1>

Kimmerer, W. 2006. Response of anchovies dampens effects of the invasive bivalve *Corbula amurensis* on the San Francisco estuary foodweb. *Marine Ecology Progress Series* 324:207-218.

Kimmerer, W. 2008. Losses of Sacramento River Chinook salmon and delta smelt to entrainment in water diversions in the Sacramento-San Joaquin Delta. *San Francisco Estuary and Watershed Science* 6(2): Article 2, 28 pages.

Kimmerer, W. 2011. Modeling delta smelt losses at the South Delta export facilities. *San Francisco Estuary and Watershed Science* 9(1):9.

Kimmerer, W.J. and M.L. Nobriga. 2008. Investigating particle transport and fate in the Sacramento-San Joaquin Delta using a particle tracking model. *SFEWS* 6(1) (February 2008), Article 4 <http://repositories.cdlib.org/jmie/sfews/vol6/iss1/art4>

Kimmerer, W.J. and J.K. Thompson. 2014. Phytoplankton Growth Balanced by Clam and Zooplankton Grazing and Net Transport into the Low-Salinity Zone of the San Francisco Estuary. *Estuaries and Coasts* 37:1202–1218. DOI 10.1007/s12237-013-9753-6.

Kimmerer, W.J., E. Gartside and J.J. Orsi. 1994. Predation by an introduced clam as the likely cause of substantial declines in zooplankton of San Francisco Bay. *Mar. Ecol. Progr. Ser.* 113: 81-93.

Kimmerer, W.J., E.S. Gross and M.L. MacWilliams. 2009. Is the response of estuarine nekton to freshwater flow in the San Francisco Estuary explained by variation in habitat volume? *Estuaries and Coasts* 32: 375-389.

Kimmerer, W.J., A.E. Parker, U.E. Lidström, and E.J. Carpenter. 2012. Short-Term and interannual variability in primary production in the low-salinity zone of the San Francisco Estuary. *Estuaries and Coasts* 35(4):913-929.

Kimmerer, W.J., M.L. MacWilliams, and Gross, E.S.. 2013. Variation of Fish Habitat and Extent of the Low-Salinity Zone with Freshwater Flow in the San Francisco Estuary. *San Francisco Estuary and Watershed Science*. Dec 2013. 18 pp.

Komoroske, L.M., R.E. Connon, J. Lindberg , B.S. Cheng, G. Castillo, M. Hasenbein, N.A. Fangue. 2014. Ontogeny influences sensitivity to climate change stressors in an endangered fish. *Conserv Physiol* 2: doi:10.1093/conphys/cou008.

Kratina, P., R. Mac Nally, W.J. Kimmerer, J.R. Thompson, M. Winder. 2014. Human-Induced Biotic Invasions and Changes in Plankton Interaction Networks. *Journal of Applied Ecology* 51:1066-1074.

Kuivila, K.M. 1999. Studies Relating Pesticide Concentrations to Potential Effects on Aquatic Organisms in the San Francisco Bay-Estuary, California. U.S. Geological Survey Toxic Substances Hydrology Program. Proceedings of the Technical Meeting, Charleston, South Carolina, March 8-12, 1999, v. 2. Contamination of hydrologic Systems and Related Ecosystems, 1999, U.S. Geological Survey Water-Resources Investigations Report 99-4018 B.

Kuivila, K, and G.E. Moon. 2004. Potential exposure of larval and juvenile delta smelt to dissolved pesticides in the Sacramento-San Joaquin Delta, California. Pages 229-242 *In American Fisheries Society Symposium*, American Fisheries Society.

Lande, R. 1998. Demographic stochasticity and allee effect on a scale with isotropic noise. *Oikos* 83(2):353-358.

Lehman, P.W. G. Boyer, C. Hall, S. Walker and K. Gehrts. 2005. Distribution and toxicity of a new colonial *Microcystis aeruginosa* bloom in the San Francisco Bay Estuary, California. *Hydrobiologia* 541:87-99.

Lehman, P.W., S.J. Teh, G.L. Boyer, M.L. Nobriga, E. Bass, C. Hogle. 2010a. Initial impacts of *Microcystis aeruginosa* blooms on the aquatic food web in the San Francisco Estuary. *Hydrobiologia* 637:229–248.

Lehman, P.W., S. Mayr, L. Mecum, C. Enright. 2010b. The freshwater tidal wetland Liberty Island, CA was both a source and sink of inorganic and organic material to the San Francisco Estuary. *Aquat Ecology* 44:359–372.

Lehman, P.W., K. Marr, G.L. Boyer, S. Acuna, and S.J. Teh. 2013. Long-term trends and causal factors associated with *Microcystis* abundance and toxicity in San Francisco Estuary and implications for climate change impacts. *Hydrobiologia* 718(1):141-158.

LFR Levine-Fricke. 2004. Framework for assessment of potential effects of dredging on sensitive fish species in San Francisco Bay. Final report. Prepared for U.S. Army Corps of Engineers. 141 pp. LFR Levine-Fricke, Emeryville, California.

Linville, R., G.S. Luoma, L. Cutter and G. Cutter. 2002. Increased selenium threat as a result of invasion of the exotic bivalve *Potamocorbula amurensis* into the San Francisco Bay-Delta. *Aquatic Toxicology* 57:51-64.

Lott J. 1998. Feeding habits of juvenile and adult delta smelt from the Sacramento–San Joaquin River Estuary. Interagency Ecological Program for the Sacramento–San Joaquin Estuary. IEP newsletter 11(1):14–19. Sacramento (CA): California Department of Water Resources. Available from: <http://www.water.ca.gov/iep/newsletters/1998/IEP-winter-1998>. *Cfm*

MacCoy, D., K.L. Crepeau, K.M. Kuivila. 1995. Dissolved pesticide data for the San Joaquin River at Vernalis and the Sacramento River at Sacramento, California, 1991-94. US Geological Survey, Earth Science Information Center Open-File Reports Section, Denver, CO. 27 pp.

Mac Nally, R., J.R. Thomson, W. Kimmerer, F. Feyrer, K. Newman, A. Sih, W.A. Bennett, L.R. Brown, E. Fleishman, S.D. Culberson, and G. Castillo. 2010. Analysis of pelagic species decline in the upper San Francisco Estuary using multivariate autoregressive modeling (MAR). *Ecological Application* 20(5):1417-1430.

Mager, R.C., S.I. Doroshov, J.P. Van Eenennaam and R.L. Brown. 2004. Early life stages of delta smelt *in* Early life history of fishes in the San Francisco Estuary and watershed, F. Feyrer, L. Brown, R. Brown and J. Orsi, eds. American Fisheries Society Symposium 39: 169-180, F. Feyrer, L.R. Brown, R. Brown and J.J. Orsi, eds. Bethesda, MD.

Matern, S.A., P.B. Moyle, L.C. Pierce. 2002. Native and Alien Fishes in a California Estuarine Marsh: Twenty-One Years of Changing Assemblages. *Transactions of American Fisheries Society* 131: 797-816.

Maunder, M.N., and R.B. Deriso. 2011. A state–space multistage life cycle model to evaluate population impacts in the presence of density dependence: illustrated with application to delta smelt (*Hyposmesus transpacificus*). *Canadian Journal of Fisheries and Aquatic Sciences* 68(7):1285-1306.

Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao, 2007: Global Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

- Merz, J.E., S. Hamilton, P.S. Bergman, B. Cavallo. Spatial perspective for delta smelt: a summary of contemporary survey data. *California Fish and Game* 97(4):164-189.
- Miller, D.J., and R.N. Lea. 1972. Smelt family Osmeridae. *Fish Bulletin* 157: Guide to the coastal marine fishes of California. 7 pp. California Department of Fish and Game.
- Miller, W.J., B.F. J. Manly, D.D. Murphy, D. Fullerton, and R.R. Ramey. 2012. An investigation of factors affecting the decline of delta smelt (*Hypomesus transpacificus*) in the Sacramento-San Joaquin Estuary. *Reviews in Fisheries Science* 20(1):1-19.
- Moyle, P.B. 2002. *Inland fishes of California*. Revised and expanded. University of California Press, Berkeley, CA.
- Moyle, P.B., and J. A. Israel. 2005. Untested assumptions: effectiveness of screening diversions for conservation of fish populations. *Fisheries* 30(5):20-28.
- Moyle, P.B., B. Herbold, D.E. Stevens and L.W. Miller. 1992. Life history and status of delta smelt in the Sacramento-San Joaquin estuary, California. *T. Am. Fish. Soc.* 121: 67-77
- Nelson, C., G. Benigno, L. Conrad. 2011. Mysid Abundance in a Restored Tidal Freshwater Wetland. *IEP newsletter*. Volume 24. 3: 16-22.
- Newman, K., L. Polansky, L. Mitchell. 2016. Unpublished Modelling Data on Delta Smelt Population Estimates.
- [NRC] National Research Council. 2012. Sustainable water and environmental management in the California Bay-Delta. National Academies Press, Washington, D.C. Available at http://www.nap.edu/catalog.php?record_id=13394.
- Nichols, F.H., J.E. Cloern, S.N. Luoma, and D.H. Peterson. 1986. The modification of an Estuary. *Science* 231:567-573.
- Nixon, S.W. 1988. Physical energy inputs and the comparative ecology of lake and marine ecosystems. *Limnology and Oceanography* 33(4, part 2):1005-1025.
- [NMFS] National Marine Fisheries Services [NOAA] National Oceanographic and Atmospheric Administration. 2009. Biological and Conference Opinion on the Long-Term Operations of the Central Valley Project and State Water Project. Available online. Accessed on December 22, 2014.
http://www.westcoast.fisheries.noaa.gov/publications/Central_Valley/Water%20Operations/Operations,%20Criteria%20and%20Plan/nmfs_biological_and_conference_opinion_on_the_long-term_operations_of_the_cvp_and_swp.pdf
- Nobriga, M.L. 2002. Larval delta smelt diet composition and feeding incidence: environmental and ontogenetic influences. *California Fish and Game* 88(4):149-164.

- Nobriga, M.L. 2015. Unpublished Data. Survey data analysis.
- Nobriga, M.L., F. Feyrer. 2007. Shallow-Water Piscivore-Prey Dynamics in California's Sacramento-San Joaquin Delta. *San Francisco Estuary and Watershed Science*. Vol. 5, Issue 2. Article 4. 13 pp.
- Nobriga, M.L., and B. Herbold. 2008. Conceptual model for delta smelt (*Hypomesus transpacificus*) for the Delta Regional Ecosystem Restoration and Implementation Plan (DRERIP). 47 pp.
- Nobriga, M.L., Z. Matica, and Z.P. Hymanson. 2004. Evaluating entrainment vulnerability to agricultural irrigation diversions: A comparison among open-water fishes. *American Fisheries Society Symposium* 39:281-295.
- Nobriga, M.L., F. Feyrer, R. Baxter, and M. Chotkowski. 2005. Fish community ecology in an altered river delta: Spatial patterns in species composition, life history strategies, and biomass. *Estuaries* 28(5):776-785.
- Nobriga, M.L., T.R. Sommer, F. Feyrer and K. Fleming. 2008. Long-term trends in summertime habitat suitability for delta smelt (*Hypomesus transpacificus*). *San Francisco Estuary and Watershed Science* Vol. 6, Issue 1 (February 2008), Article 1.
<http://repositories.cdlib.org/jmie/sfews/vol6/iss1/art1>
- Nobriga, M.L., E. Loboschfsky, F. Feyrer. 2013. Common Predator, Rare Prey: Exploring Juvenile Striped Bass Predation on Delta Smelt in California's San Francisco Estuary. *Transactions of the American Fisheries Society*. 142:1563-1575.
- NRG Delta. 2014. 2013 Annual Biological Opinions Letter Report. NGG Delta, LLC-Contra Costa and Pittsburg Generating Stations. 5 pp.
- Orsi, J.J, W.L. Mecum. 1986. Zooplankton Distribution and Abundance in the Sacramento-San Joaquin Delta in Relation to Certain Environmental Factors. *Estuaries* 9 (4B): 326-339.
- Parker, A.E., R.C. Dugdale, and F.P. Wilkerson. 2012. Elevated ammonium concentrations from wastewater discharge depress primary productivity in the Sacramento River and the Northern San Francisco Estuary. *Mar Pollut Bull* 64(3):574-586.
- PRBO Conservation Science. 2011. Projected Effects of Climate Change in California: Ecoregional. Summaries Emphasizing Consequences for Wildlife. Version 1.0.
<http://data.prbo.org/apps/bssc/uploads/Ecoregional021011.pdf> (Accessed December 22, 2014).
- Prinn, R., S. Paltsev, A. Sokolov, M. Sarofim, J. Reilly, H. Jacoby. 2011. Scenarios with MIT integrated global systems model: significant global warming regardless of different approaches. *Climatic Change* (2011) 104:515-537.

Radtke, L.D. 1966. Distribution of smelt, juvenile sturgeon and starry flounder in the Sacramento-San Joaquin Delta. pp 115-119 in Ecological studies of the Sacramento-San Joaquin estuary, Pt. 2, S.L. Turner and D.W. Kelley, eds., Calif. Dept. Fish Game Bull. 136
<http://content.cdlib.org/view?docId=kt8h4nb2t8;NAAN=13030&doc.view=frames&chunk.id=d0e3269&toc.depth=1&toc.id=d0e3269&brand=calisphere>

Rieman, B.E., and D.J. Isaak. 2010. Climate change, aquatic ecosystems, and fishes in the Rocky Mountain west: Implications and alternatives for management. General technical report RMRS-GTR-250. 46 pages. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO.

Rose, K.A., W.J. Kimmerer, K.P. Edwards and W.A. Bennett. 2013a. Individual-Based Modeling of Delta Smelt Population Dynamics in the Upper San Francisco Estuary: I. Model Description and Baseline Results, Transactions of the American Fisheries Society, 142:5, 1238-1259, DOI: [10.1080/00028487.2013.799518](https://doi.org/10.1080/00028487.2013.799518)

Rose, K.A., W.J. Kimmerer, K.P. Edwards and W.A. Bennett. 2013b. Individual-Based Modeling of Delta Smelt Population Dynamics in the Upper San Francisco Estuary: II. Alternative Baselines and Good versus Bad Years, Transactions of the American Fisheries Society, 142:5, 1260-1272, DOI: [10.1080/00028487.2013.799519](https://doi.org/10.1080/00028487.2013.799519)

Schoellhamer, D.H. 2002. Comparison of the Basin-scale Effect of Dredging Operations and Natural Estuarine Processes on Suspended Sediment Concentration. Estuaries Vol. 25, No. 3, p. 488–495.

Schoellhamer, D.H. 2011. Sudden Clearing of Estuarine Waters upon Crossing the Threshold from Transport to Supply Regulation of Sediment Transport as an Erodible Sediment Pool is Depleted: San Francisco Bay, 1999. Estuaries and Coasts 34: 885–899.

Scholz, N.L., N.K. Truelove, B.L. French, B.A. Berejikian, T.P. Quinn, E. Casillas, T.K. Collier. 2000. Diazinon disrupts antipredator and homing behaviors in Chinook salmon (*Oncorhynchus tshawytscha*). Canadian Journal of Fisheries and Aquatic Science 57:1911-1918.

Shafer, T.J. and D.A. Meyer. 2004. Effects of pyrethroids on voltage-sensitive calcium channels: a critical evaluation of strengths, weaknesses, data needs, and relationship to assessment of cumulative neurotoxicity. Toxicology and Applied Pharmacology 196:303-318.

Sirois, P., and J.J. Dodson. 2000. Critical periods and growth-dependent survival of larvae of an estuarine fish, the rainbow smelt *Osmerus mordax*. Marine Ecology Progress Series 203:233-245.

Slater, S.B., R.D. Baxter. 2014. Diet, Prey Selection, and Body Condition of Age-0 Delta Smelt, *Hypomesus transpacificus*, in the Upper San Francisco Estuary. San Francisco Estuary and Watershed Science. 26 pp.

Slater, S. 2015. Interagency Ecological Program scientific take. Unpublished data. 2005-2015.

Sobczak, W.V., J.E. Cloern, A.D. Jassby, and A.B. Muller-Solger. 2002. Bioavailability of organic matter in a highly disturbed estuary: the role of detrital and algal resources. *Proceedings of the National Academy of Sciences* 99(12):8101-8105.

Solomon, S., D. Qin, M. Manning, R.B. Alley, T. Berntsen, N.L. Bindoff, Z. Chen, A. Chidthaisong, J.M. Gregory, G.C. Hegerl, M. Heimann, B. Hewitson, B.J. Hoskins, F. Joos, J. Jouzel, V. Kattsov, U. Lohmann, T. Matsuno, M. Molina, N. Nicholls, J. Overpeck, G. Raga, V. Ramaswamy, J. Ren, M. Rusticucci, R. Somerville, T.F. Stocker, P. Whetton, R.A. Wood, and D. Wratt. 2007. Technical Summary. *In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Sommer, T., C. Armor, R. Breuer, L. R. Brown, M. Chotkowski, S. D. Culberson, F. Feyrer, M. Gingras, B. Herbold, W. Kimmerer, A. Mueller-Solger, M. L. Nobriga, and K. Souza. 2007. The collapse of pelagic fishes in the upper San Francisco Estuary. *Fisheries* 32(6):270-277.

Sommer, T.R., F. Heija, M. Nobriga, F. Feyrer, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science*. June 2011. 17 pp.

Sommer, T., F. Mejia. 2013. A Place to Call Home: A Synthesis of Delta Smelt Habitat in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science*. June 2013. 25 pp.

Stanley, S., P.B. Moyle and H.B. Shaffer. 1995. Electrophoretic analysis of delta smelt, *Hypomesus transpacificus*, and longfin smelt, *Spirinchus thaleichthys*, in the Sacramento-San Joaquin estuary, California. Final report to the Department of Water Resources, Sacramento, CA 20 pp.

[SWRCB] State Water Resources Control Board. 1995. Water Quality Control Plan for the San Francisco Bay/Sacramento-San Joaquin Delta Estuary. Resolution no. 95-24. California Environmental Protection Agency. 55 pp.

[SWRCB] State Water Resources Control Board. 2000. Revised Water Right Decision 1641. Implementation of Water Quality Objectives for the San Francisco Bay/Sacramento-San Joaquin Delta Estuary.

Stevens, D.E. 1963. Food habits of striped bass, *Roccus saxatilis*, in the Sacramento-Rio Vista Area of the Sacramento delta. Masters Thesis. University of California. 57 pp.

Stevens, D.E. 1966. Food habits of striped bass, *Roccus saxatilis*, in the Sacramento-San Joaquin Delta. Pages 68-96 in Turner JL, Kelley DW (eds). *Ecological studies of the Sacramento-San Joaquin Delta, part II, fishes of the Delta*. California Department of Fish and Game Fish Bulletin 136.

Stevens, D.E., and L.W. Miller. 1983. Effects of river flow on abundance of young Chinook salmon, American shad, longfin smelt, and delta smelt in the Sacramento-San Joaquin River system. *North American Journal of Fisheries Management* 3:425-437.

Stewart, A.R., S.N. Luoma, C.E. Schlekot, M.A. Doblin, K.A. Heib. 2004. Food Web Pathway Determines How Selenium Affects Aquatic Ecosystems: A San Francisco Bay Case Study. *Environmental Science Technology* 38: 4519-4526.

Suisun Marsh Habitat Restoration Plan. 2011. Accessed on January 13, 2015. Available at \\ifw8delt-fs4\DFSRoots\Data\ESA -REGULATORY\Section 4\Recovery plans\2014 Delta Native Fishes Plan\Citations\Plans\Suisun Marsh Plan_Final EIS.mht

Swanson, C., T. Reid, P.S. Young and J.J. Cech. 2000. Comparative environmental tolerances of threatened delta smelt (*Hypomesus transpacificus*) and introduced wakasagi (*H. nipponensis*) in an altered estuary. *Oecologia* 123:384-390.

Sweetnam, D.A. 1999. Status of delta smelt in the Sacramento-San Joaquin Estuary. *California Fish and Game* 85:22-27.

Taplin, J., and C. H. Hanson. 2006. Results of 2006 investigation of juvenile Chinook salmon and steelhead entrainment into a hydraulic suction dredge during sand mining within the Sacramento-San Joaquin Bay-Delta Estuary. Prepared for: Hanson Marine Operations, Inc., Cemex Construction Materials LTD, Jerico Products Inc./Morris Tug and Barge. 37 pp.

Tanner, D.K. and M.L. Knuth. 1996. Effects of esfenvalerate on the reproductive success of the bluegill sunfish, *Lepomis macrochirus* in littoral enclosures. *Archives of Environmental Contaminants and Toxicology* 31:244-251.

Teh, S. J. 2007. Pilot study of the health status of the 2005 adult delta smelt in the upper San Francisco Estuary. Final Report of Histopathological Evaluation of Starvation and/or Toxic Effects on Pelagic Fishes. 28 pp.

Teh, S., I. Flores, M. Kawaguchi, S. Lesmeister and C. Teh. 2011. Full life-cycle bioassay approach to assess chronic exposure of *Pseudodiaptomus forbesi* to ammonia/ammonium. Final report. Submitted to the State Water Board on March 4, 2011. UC Davis, Aquatic Toxicology Program. Davis, CA.

Thomas, J.L..1967. The Diet of Juvenile and Adult Striped Bass, *Roccus Saxatilis*, In the Sacramento-San Joaquin River System. *California Fish and Game Journal* 53 (1) : 49-62.

Thomson, J.R., W.J. Kimmerer, L.R. Brown, K.B. Newman, R. MacNally, W.A. Bennett, F. Feyrer and E. Fleishman. 2010. Bayesian change point analysis of abundance trends for pelagic fishes in the upper San Francisco estuary. *Ecol. Appl.* 20: 1431-1448.

Trenham, P.C., H.B. Shaffer, P.B. Moyle. Biochemical Identification and Assessment of Population Subdivision in Morphologically Similar Native and Invading Smelt Species

(*Hypomesus*) in the Sacramento–San Joaquin Estuary, California. Transactions of the American Fisheries Society 127: 417-424.

[USBR] U.S. Bureau of Reclamation. 2008. Central Valley Project and State Water Project Operations Criteria and Plan biological assessment. 1016 pages.

[USBR] U.S. Bureau of Reclamation. 2011. SECURE water act section 9503(c) – Reclamation climate change and water 2011. U.S. Department of the Interior, Denver, CO. 206 pp.

[USEPA] U.S. Environmental Protection Agency. 2013. Aquatic Life Ambient Water Quality Criteria for Ammonia-Freshwater. Office of Water. August 2013. 3 pp.

[USEPA] U.S. Environmental Protection Agency. 2014. National Pollution Discharge Elimination System website. <http://water.epa.gov/polwaste/npdes/> <Accessed on December 29, 2014>.

[USFWS] U.S. Fish and Wildlife Service. 2000. Impacts of riprapping to ecosystem functioning, Lower Sacramento River, California. Prepared for U.S. Army Corps of Engineers, Sacramento District. 40 pp. Sacramento Fish and Wildlife Office, Sacramento, California.

[USFWS] U.S. Fish and Wildlife Service. 2005. Operational Criteria and Plan Biological Opinion. 238 pages, Sacramento Field Office, Sacramento, California.

[USFWS] U.S. Fish and Wildlife Service. 2008. Formal endangered species act consultation on the proposed coordinated operations of the Central Valley Project (CVP) and State Water Project (SWP). Sacramento, CA 396 pp

[USFWS] U.S. Fish and Wildlife Service. 2010. Delta smelt 5-year review. 19 pp.

U.S. Navy. 1990.

Vincik, R. F., J. M. Julienne. 2012. Occurrence of delta smelt (*Hypomesus transpacificus*) in the lower Sacramento River near Knights Landing, California. California Fish and Game 98(3):171-174.

Wang, J.C.S. 1986. Fishes of the Sacramento-San Joaquin estuary and adjacent waters, California: a guide to early life histories. Interagency Ecological Studies Program Tech. Rpt. 9

Wang, J.C.S. 1991. Early life stages and early life history of the delta smelt, *Hypomesus transpacificus*, in the Sacramento-San Joaquin estuary with comparison of early life stages of the longfin smelt, *Spirinchus thaleichthys*. Interagency Ecological Studies Program Tech. Rpt. 28

Werner, I., L. Deanovic, D. Markiewicz, M. Stillway, N. Offer, R. Connon, S. Brander. 2008. Pelagic Organism Decline (POD): Acute and chronic invertebrate and fish toxicity testing in the Sacramento-San Joaquin Delta, 2006-2007. Final Report, 30 April, 2008. Aquatic Toxicology Laboratory, School of Veterinary Medicine, University of California, Davis.

Werner, I., D. Markiewicz, L. Deanovic, R. Connon, S. Beggel, S. Teh, M. Stillway and C. Reece. 2010. Pelagic Organism Decline (POD): Acute and chronic invertebrate and fish toxicity testing in the Sacramento-San Joaquin Delta, 2008-2010. Final Report submitted to the California Department of Water Resources, July 24, 2010. Aquatic Toxicology Laboratory, School of Veterinary Medicine, University of California, Davis.

Weston, D.P. and M.J. Lydy. 2010. Urban and agricultural sources of pyrethroid insecticides to the Sacramento-San Joaquin Delta of California. *Environmental Science and Technology* 44:1833-1840.

Weston, D.P., D. Chen, and M.J. Lydy. 2015. Stormwater-related transport of the insecticides bifenthrin, fipronil, imidacloprid, and chlorpyrifos into a tidal wetland, San Francisco Bay, California. *Science of the Total Environment* 527:18-25.

White, D.K., C. Swanson, P.S. Young, J.J. Cech, Z.Q. Chen, and M.L. Kavvas. 2007. Close encounters with a fish screen II: delta smelt behavior before and during screen contact. *Transactions of the American Fisheries Society* 136 (2):528-538.

Wicks, B.J., R. Joensen, Q. Tang, and D.J. Randall. 2002. Swimming and ammonia toxicity in salmonids: the effect of sub lethal ammonia exposures on the swimming performance of coho salmon and the acute toxicity of ammonia in swimming in rainbow trout. *Aquatic Toxicology* 59:55-69.

Winder, M., A.D. Jassby, and R. Mac Nally. 2011. Synergies between climate anomalies and hydrological modifications facilitate estuarine biotic invasions. *Ecol Lett* 14(8):749-757.

Wright, S.A. and D.H. Schoellhamer. 2004. Trends in the sediment yield of the Sacramento River, California, 1957-2001. *San Francisco Estuary and Watershed Science* Vol. 2, Issue 2 (May 2004), Article 2 <http://repositories.cdlib.org/jmie/sfews/vol2/iss2/art2>

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve: _____
Regional Director, Fish and Wildlife Service Date

Concur: _____
Director, Fish and Wildlife Service Date

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks: