


U.S. Fish & Wildlife Service

Arcata Fisheries Data Series Report DS 2019-62

Mainstem Trinity River Chinook Salmon Spawning Survey, 2017

Stephen A. Gough¹, Nicholas A. Som¹, Shane Quinn², William C. Matilton³,
Andrew M. Hill⁴, and William Brock⁵



U.S. Fish and Wildlife Service¹
Arcata Fish and Wildlife Office
1655 Heindon Road, Arcata, California 95521
(707) 822-7201

Yurok Tribal Fisheries Program²
Willow Creek, California 95573
(530) 629-3333

Hoopa Valley Tribal Fisheries Department³
Hoopa, California 95546
(530) 625-4267

California Department of Fish and Wildlife⁴
Weaverville, California 96093
(530) 623-5309

U.S. Forest Service⁵
Shasta-Trinity National Forest
Weaverville Ranger District
Weaverville, California 96093
(530) 623-2121



July 2019



Arcata Fish and Wildlife Office participation in this study was funded by the U.S. Fish and Wildlife Service. Participation of the Yurok Tribal Fisheries Program, Hoopa Valley Tribal Fisheries Department, and California Department of Fish and Wildlife was funded by the U.S. Bureau of Reclamation and the U.S. Fish and Wildlife Service. Shasta–Trinity National Forest participation was funded by the U.S. Forest Service.

Disclaimer: The mention of trade names or commercial products in this report does not constitute endorsement or recommendation for use by the Federal Government.

The Fish and Aquatic Conservation Program of the Arcata Fish and Wildlife Office reports its study findings through two publication series. The **Arcata Fisheries Data Series** was established to provide timely dissemination of data to managers and for inclusion in agency databases. **Arcata Fisheries Technical Reports** publish scientific findings from single and multi-year studies that have undergone more extensive peer review and statistical testing. Additionally, results of Program studies may be published in a variety of peer-reviewed technical journals. To ensure consistency with Service policy, Arcata Fisheries Data Series and Technical Reports are distributed electronically and made available in the public domain. Paper copies are no longer produced and circulated.

key words: carcass, Chinook Salmon, escapement, hatchery, pre-spawn mortality, redd, restoration, Trinity River

The correct citation for this report is:

Gough, S. A., N. A. Som, N. S. Quinn, W. C. Matilton, A. M. Hill, and W. Brock. 2019. Mainstem Trinity River Chinook Salmon spawning survey, 2017. U.S. Fish and Wildlife Service. Arcata Fish and Wildlife Office, Arcata Fisheries Data Series Report Number DS 2019–62, Arcata, California.

Table of Contents

| | Page |
|--|-----------|
| List of Tables..... | <i>iv</i> |
| List of Figures | <i>v</i> |
| List of Appendices..... | <i>vi</i> |
| Introduction | 2 |
| Methods..... | 4 |
| Survey Area and Timing..... | 4 |
| Redd Identification..... | 4 |
| Carcasses Estimation..... | 4 |
| Pre-Spawn Mortality | 5 |
| Redd–Carcass Relationship | 7 |
| Trends in Redd Abundance and Distribution | 7 |
| Results..... | 11 |
| Survey Success and Conditions | 11 |
| Salmon Carcasses | 11 |
| <i>Carcass Estimates</i> | 14 |
| <i>Pre-spawn Mortality</i> | 14 |
| Salmon Redds..... | 16 |
| <i>Redd–Carcass Relationship</i> | 19 |
| <i>Redd Abundance and Distribution: System Scale</i> | 19 |
| <i>Redd Abundance and Distribution: Reach Scale</i> | 22 |
| <i>Redd Abundance and Distribution: Site Scale</i> | 22 |
| Discussion | 28 |
| Acknowledgements..... | 30 |
| Literature Cited | 31 |
| Appendices | 33 |

List of Tables

| | Page |
|--|------|
| Table 1. Reach boundaries [and river kilometer (rkm)] for the mainstem Trinity River, California, salmon spawning surveys. | 6 |
| Table 2. The reach- and site-scale sections used for redd abundance and distribution analysis within the restoration reach. | 9 |
| Table 3. River sections [with river kilometer (rkm)] used for the reach-scale analysis of redd abundance..... | 10 |
| Table 4. Summary of fresh (conditions 1 and 2) Chinook Salmon carcass data by survey reach, 2017 Trinity River surveys. | 12 |
| Table 5. Coded-wire tag (CWT) information retrieved from fresh adipose fin-clipped Chinook Salmon carcasses, 2017 Trinity River surveys..... | 12 |
| Table 6. Summary of fresh (conditions 1 and 2) Coho Salmon carcass data by survey reach, 2017 Trinity River surveys..... | 13 |
| Table 7. Pre-spawn mortality rates of Chinook Salmon in the Trinity River below Lewiston Dam (Reaches 1–14) and in the restoration reach (Reaches 1–7), 2009–2017 surveys..... | 14 |
| Table 8. Pre-spawn mortality rates of natural- and hatchery-origin Coho Salmon, Trinity River surveys, 2009–2017..... | 15 |
| Table 9. Redd counts (before species differentiation) by week and reach, Trinity River surveys 2017. | 16 |
| Table 10. Estimated numbers and bootstrap-generated 95% confidence intervals of salmon redds by species and origin observed in the mainstem Trinity River, 2017..... | 17 |

List of Figures

| | Page |
|---|------|
| Figure 1. Survey Reaches 1–14 (Lewiston Dam to Weitchpec) on the mainstem Trinity River, California. | 5 |
| Figure 2. The ten sections of the mainstem Trinity River used for reach-scale analyses of Chinook Salmon redd distribution. | 10 |
| Figure 3. Distribution of coded-wire-tagged (CWT) spawned female Chinook Salmon carcasses by brood type (spring and fall) located in the mainstem Trinity River downstream of Lewiston Dam in 2017. | 13 |
| Figure 4. Weekly pre-spawn mortality from fresh (conditions 1 and 2) female Chinook Salmon carcasses, Trinity River surveys 2017. | 15 |
| Figure 5. Spatiotemporal distribution of mainstem Trinity River salmon redds from Lewiston Dam to Weitchpec, 2017. | 18 |
| Figure 6. Relationship between counts of <i>ln</i> -transformed Chinook Salmon redds and <i>ln</i> -transformed estimates of spawned female Chinook Salmon carcasses in Survey Reaches 1 and 2 (solid line), 2012–2017. | 19 |
| Figure 7. Estimated number of redds constructed in the entire mainstem Trinity River (left), within the restoration reach (center), and downstream (DS) of the restoration reach (right) by all Chinook Salmon (top), natural-origin Chinook Salmon (middle), and hatchery-origin Chinook Salmon (bottom) from 2002 to 2017. | 20 |
| Figure 8. Mean distance from Lewiston Dam of redds constructed by natural- (left) and hatchery-origin (right) Chinook Salmon females between Lewiston Dam and Cedar Flat (0–102.8 km from Lewiston Dam; Reaches 1–10) on the mainstem Trinity River, 2002–2017. | 21 |
| Figure 9. Estimated number of mainstem Trinity River natural-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. | 24 |
| Figure 10. Proportions of mainstem Trinity River natural-origin Chinook Salmon redds relative to the total mainstem count of natural-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. | 25 |
| Figure 11. Estimated number of mainstem Trinity River hatchery-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. | 26 |
| Figure 12. Proportions of mainstem Trinity River hatchery-origin Chinook Salmon redds relative to the total mainstem count of hatchery-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. | 27 |

List of Appendices

| | Page |
|--|------|
| Appendix A. Trinity River water visibility by week and reach throughout the 2017 survey period. | 33 |
| Appendix B. Trinity River mean daily discharge at Lewiston (USGS Gage 11525500) and Hoopa, California (USGS Gage 11530000) during the 2017 survey season. | 34 |
| Appendix C. Pre-spawn mortality numbers by week and reach of unmarked and ad-clipped fresh (conditions 1 and 2) female Chinook Salmon carcasses, mainstem Trinity River surveys 2017. | 35 |
| Appendix D. Proportion of TRRP restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. | 36 |
| Appendix E. Proportion of natural-origin Chinook Salmon redds within site-scale sections in the TRRP restoration reach that encompass mechanical channel rehabilitation locations, 2002–2017. | 42 |
| Appendix F. Natural-origin Chinook Salmon redd counts versus estimates of pre-spawn mortality from Lewiston Dam to the North Fork confluence, Trinity River surveys, 2009–2017. | 45 |

Mainstem Trinity River Chinook Salmon Spawning Survey, 2017

Stephen A. Gough¹, Nicholas A. Som¹, Shane Quinn², Billy C. Matilton³,
Andrew M. Hill⁴, and William Brock⁵

¹ U.S. Fish and Wildlife Service, Arcata Fish and Wildlife Office
1655 Heindon Road, Arcata, California 95521
steve_gough@fws.gov
nicholas_som@fws.gov

² Yurok Tribal Fisheries Program
Highway 96 Box 36, Willow Creek, California 95573
squinn@yuroktribe.nsn.us

³ Hoopa Valley Tribal Fisheries Department
P.O. Box 417, Hoopa, California 95546
bmatilton@hoopa-nsn.gov

⁴ California Department of Fish and Wildlife
80 Nugget Lane, Suite A, Weaverville, California 96093
andrew.hill@wildlife.ca.gov

⁵ U.S. Forest Service, Shasta-Trinity National Forest
360 West Main Street, Weaverville, California 96093
wbrock@fs.fed.us

Abstract.— Salmon redds and carcasses were surveyed on the mainstem Trinity River, California from Lewiston Dam to the confluence with the Klamath River, during the 2017 spawning season to map spawning abundance and distribution, evaluate pre-spawn mortality, and characterize redds by species and spawner origin. The total redd count in 2017 was 1,982. We applied generalized additive models to the spatiotemporal distribution of unmarked and hatchery-marked spawned female salmon carcasses to apportion redd counts by natural- and hatchery-origin Chinook Salmon *Oncorhynchus tshawytscha* and Coho Salmon *O. kisutch*. This methodology only allows for the partitioning of redds constructed by hatchery- and natural-produced females and does not account for the origin of the male spawners. We estimated that 1,600 (95% c.i.: 1,435–1,762) redds were constructed by natural-origin Chinook Salmon, 348 (95% c.i.: 186–513) by hatchery-origin Chinook Salmon, and the remaining 34 were attributed to Coho Salmon. Natural-origin Chinook Salmon spawned throughout the mainstem river while the distribution of redds constructed by hatchery-origin Chinook Salmon was highly skewed toward Lewiston Dam and the Trinity River Hatchery (about 59% were within 10 km of the

dam). Pre-spawn mortality of female Chinook Salmon was 1.8% for carcasses observed in all reaches and 2.0% within an intensively managed ‘restoration reach’, which is a focal area for habitat restoration improvements being implemented by the Trinity River Restoration Program (TRRP). Long-term trend analyses from 2002 to 2017 showed no significant change in the abundance of natural-origin Chinook Salmon redds constructed in the mainstem Trinity River, while the number of hatchery-origin Chinook Salmon redds decreased. The proportion of total annual natural-origin Chinook Salmon redds decreased in the reaches nearest to Lewiston Dam and increased in reaches farther downstream from 2002 to 2017, while the annual component of hatchery-origin Chinook Salmon redds remained almost completely within the two reaches nearest to Lewiston Dam.

Introduction

The Trinity River, California, once supported large populations of naturally produced anadromous salmonids, including spring- and fall-run Chinook Salmon *Oncorhynchus tshawytscha* (USFWS and HVT 1999). Prior to the construction of Trinity and Lewiston dams, the spawning of spring- and fall-run Chinook Salmon was separated temporally and spatially due to the timing of adult upstream migration of each race and the hydrology of the river. In 1940s, Moffett and Smith (1950) noted that “almost without exception, Trinity River salmon migrating above the South Fork spawn in the 72 miles of river between the North Fork and Ramshorn Creek.”

Following construction of Lewiston Dam [river kilometer (rkm 182.2)], spring- and fall-run Chinook Salmon spawning in the mainstem Trinity River exhibited considerable spatial and temporal overlap due to lack of access to historic spawning areas for the spring-run. High redd densities became frequent within the upper-most portions of the river below the dam, where presumably hatchery-origin salmon and their progeny comingled and spawned with naturally produced fish. Trinity River Hatchery (TRH), located at the base of Lewiston Dam, is operated to mitigate for the loss of Chinook Salmon, Coho Salmon *O. kisutch*, and steelhead *O. mykiss* production upstream of the dam. Rogers (1972) documented that in 1970 more than 50% of Chinook Salmon spawned in the two miles (3.2 km) below Lewiston Dam and 80% spawned above Douglas City (around rkm 150.1). Redd surveys in the 1980s and 1990s between North Fork Trinity River (rkm 118.2) and Cedar Flat (rkm 79.1) documented variable spawning use in these reaches, with redd counts ranging from a low of 187 in 1998 to a high of 928 redds in 1997 (USFWS 1986, 1987; Quihillalt 1999). Chamberlain et al. (2012) noted that the mean distance from Lewiston Dam of natural-origin Chinook Salmon redds upstream of Cedar Flat increased from 2002 to 2011. Rupert et al. (2017a) noted that when the mainstem Trinity River was divided into reach-scale sections, natural-origin Chinook Salmon spawning activity decreased near Lewiston Dam and increased in sections of the river farther downstream.

In an effort to restore the fishery resources of the Trinity River, the Secretary of the Interior signed the Trinity River Mainstem Fishery Restoration Record of Decision (ROD) in 2000 (USDOI 2000) and the Trinity River Restoration Program (TRRP) was established. The goal of the TRRP is to:

“...restore and sustain natural production of anadromous fish populations downstream of Lewiston Dam to pre-dam levels, to facilitate dependent tribal, commercial, and sport fisheries’ full participation in the benefits of restoration via enhanced harvest opportunities” (TRRP and ESSA 2009).

To achieve this goal, the TRRP implements a suite of actions (flow management, mechanical channel rehabilitation, coarse sediment augmentation, and watershed restoration) to restore riverine habitats and restore habitat-creating alluvial processes (USFWS and HVT 1999; USDOJ 2000). Collectively, these actions are intended to increase and maintain salmonid habitats in the 64-km section of the Trinity River from Lewiston Dam downstream to the North Fork Trinity River (restoration reach), which was severely degraded due the operation of the Trinity River Division (TRD) of the Central Valley Project. Downstream of the North Fork confluence, the Trinity River valley narrows and accretions of flow and sediment from tributaries attenuate many of the morphological impacts that have occurred in the restoration reach (USFWS and HVT 1999).

The Integrated Assessment Plan (IAP; TRRP and ESSA 2009) sets forth a list of objectives to evaluate the effectiveness of TRRP restoration actions. Salmon spawning surveys are preformed to provide data to address Objective 3, specifically sub-objectives 3.1 and 3.3:

Objective 3: Restore and maintain natural production of anadromous fish populations.

Sub-objective 3.1: Increase spawning, incubation, and emergence success of anadromous spawners.

Sub-objective 3.3: Minimize impacts of predation and genetic interactions between and among hatchery and natural anadromous fish.

The IAP proposes assessing spawning at three spatial scales: system, reach, and site scales. Each of these spatial scales evaluates the effects of restoration efforts on Chinook Salmon spawning at different resolutions. System-scale analysis evaluates the response to all restoration activities combined over time. Reach-scale analysis evaluates the response to management actions within sections of the river that have unique hydrology and sediment supplies. Finally, site-scale analysis provides insight on changes in spawning distribution/abundance within restoration sites and the localized effects of mechanical channel rehabilitation. The IAP also states that “increased spawner success will likely occur within 3–4 brood cycles following completion of channel rehabilitation and subsequent fluvial and geomorphic evolution.”

This report details the results from salmon spawning survey data collected in 2017 on the mainstem Trinity River. Surveying salmon carcasses provides pre-spawn mortality data and carcass estimates and reflect the species and origin composition of spawned salmon. Surveying salmon redds provides the location and spawn timing of individual redds. When analyzed together, each year’s data produces a spatially and temporally explicit set of observed redd locations with each redd having an associated probability of construction by female natural-origin Chinook Salmon, hatchery-origin Chinook Salmon, natural-origin Coho Salmon, and hatchery-origin Coho Salmon. We define ‘hatchery-origin’ as fish produced and released from Trinity River Hatchery (TRH), and ‘natural-origin’ as fish that emerge from a redd, regardless of parental origin. These data sets facilitate an array of

analyses over a range of spatial and temporal scales, which we use to investigate spawning distribution and abundance. Where applicable, we use the performance measures set forth by the IAP to evaluate changes in spawning as responses to the restoration actions of the TRRP.

Methods

Survey Area and Timing

The Trinity River from Lewiston Dam to its confluence with the Klamath River was delineated into 14 survey reaches ranging in length from 3.3 to 21.3 km (Figure 1, Table 1). Reach breaks were based on river access locations and channel distances that could be surveyed in a day. Two whitewater sections were not surveyed: the 9.7-km Pigeon Point run (Reach 8) and the 15.6-km section that includes the Burnt Ranch Gorge (Reach 11). In 2016, the boundary separating Reaches 5 and 6 was moved from Roundhouse (rkm 135.7) to Evan's Bar (rkm 137.4) because of a change in private landowner permission to use their river access.

Reaches 1–7 were surveyed weekly and Reaches 9–14 (excluding Reach 11) were surveyed every other week, as conditions permitted, for salmon carcasses and redds as described in Rupert et al. (2017a). Surveys in 2017 began August 30 and concluded December 20. This period was intended to encompass the majority of Chinook Salmon spawning activity.

Redd Identification

Chinook and Coho salmon spawning periods temporally overlap and natural- and hatchery-origin salmon spawn in the same areas in the mainstem Trinity River. Given that redds are not visually distinguishable by these species and origin types, the estimated proportion and spatial distribution of fresh female carcasses of hatchery- and natural-origin Chinook and Coho salmon were used to infer the probability of redd construction by species and origin. Since only female carcasses are used in the hatchery–natural analysis, the estimates of redds constructed by natural-origin females do not account for hatchery-produced males spawning with naturally produced females. Therefore natural-origin spawning estimates should be considered maximum values given that estimates were not adjusted downward to account for hatchery–natural mating pairs. Generalized Additive Models (GAM) were used with the spatiotemporal distribution of carcasses to estimate the longitudinal gradient in proportional distribution of spawned females by species (Chinook or Coho salmon) and origin (hatchery or natural) along the river channel and over time (Rupert et al. 2017a). Cumulative redd counts were arranged by survey day within reach boundaries and season total estimates of redds by species and origin were calculated by summing predicted probabilities of construction for each species–origin category (Rupert et al. 2017a).

Carcasses Estimation

Carcass abundance estimates for Reaches 1 and 2 were generated via a hierarchical latent variables model as described in Rupert et al. (2017a). This model assumes a latent (unobservable) ecological process interacts with a detection process to produce the observed counts of carcasses (Kery and Schaub 2012). For this survey, the latent process is the true




Figure 1. Survey Reaches 1–14 (Lewiston Dam to Weitchpec) on the mainstem Trinity River, California. Dangerous whitewater conditions precluded surveys in Reaches 8 and 11.

abundance of carcasses. As not all carcasses are observed (imperfect detection), a separate observation process links the unobserved latent process to the observed data. In essence, annual carcass estimates were generated by first estimating weekly detection probabilities. Next, weekly counts of fresh carcasses (those arriving since the prior survey) were assumed to arise from a binomial process, which allows the estimation of weekly abundances. Finally, weekly estimates were summed to create an annual abundance estimate as a derived parameter.

Pre-Spawn Mortality

Fresh carcasses were described as spawned ($\leq 1/3$ eggs retained), partially spawned ($1/3$ – $2/3$ eggs retained), or unspawned ($\geq 2/3$ eggs retained). These spawning condition data were used to assess levels of pre-spawn mortality. Female carcasses designated as ‘spawned’ and ‘partially spawned’ were considered successful spawners. Unspawned carcasses were considered pre-spawn mortalities. Measurement of pre-spawn mortality is limited to occurrence within the time and space of the surveys. Therefore, pre-spawn mortality in the lower Klamath River of Trinity River-bound fish and pre-spawn mortality of spring-run Chinook Salmon prior to the first survey are not reflected in our data and analyses.

Table 1. Reach boundaries [and river kilometer (rkm)] for the mainstem Trinity River, California, salmon spawning surveys. Agencies involved in data collection include California Department of Fish and Wildlife (CDFW), Shasta–Trinity National Forest (USFS), U.S. Fish and Wildlife Service (USFWS), Yurok Tribal Fisheries Program (YTFP), and Hoopa Valley Tribal Fisheries Department (HVT).

| Reach | Boundaries | | Surveying agency |
|-------|---------------------------------------|--|------------------|
| | Upstream | Downstream (rkm) | |
| 1 | Lewiston Dam (rkm 182.2) ^a | Old Lewiston Bridge (178.7) | USFS, YTFP, CDFW |
| 2 | Old Lewiston Bridge | Bucktail River Access (171.6) | CDFW, YTFP |
| 3 | Bucktail River Access | Steel Bridge River Access (160.7) | CDFW, YTFP |
| 4 | Steel Bridge River Access | Douglas City Campground (150.1) | CDFW, YTFP |
| 5 | Douglas City Campground | Evan's Bar (137.4) ^b | CDFW, YTFP |
| 6 | Evan's Bar ^b | Junction City Campground (127.1) | USFWS, HVT |
| 7 | Junction City Campground | Pigeon Point Campground ^c (117.4) | USFWS, HVT |
| 8 | Pigeon Point Campground ^c | Big Flat River Access (107.6) | NOT SURVEYED |
| 9 | Big Flat River Access | Del Loma River Access (93.8) | USFWS, HVT |
| 10 | Del Loma River Access | Cedar Flat River Access (79.1) | USFWS, HVT |
| 11 | Cedar Flat River Access | Hawkins Bar (63.4) | NOT SURVEYED |
| 12 | Hawkins Bar | Camp Kimtu in Willow Creek (42.6) | USFWS, HVT |
| 13 | Camp Kimtu in Willow Creek | Roland's Bar in Hoopa Valley (21.3) | USFWS, HVT |
| 14 | Roland's Bar in Hoopa Valley | Weitchpec (Trinity mouth; 0.0) | USFWS, HVT |

^a The spillway and pool directly downstream of Lewiston Dam were not surveyed and presumed to have no redds.

^b In 2015 and earlier the river access separating Reaches 5 and 6 was at Roundhouse (rkm 135.7).

^c Pigeon Point Campground access is 0.8 km downstream of the North Fork Trinity River confluence (rkm 118.2). The primary area where Trinity River Restoration Program actively manages to improve channel morphology and salmon habitat is in Reaches 1–7.

Redd–Carcass Relationship

Spawning density was hypothesized to affect the crews' ability to observe redds and carcasses with equal efficiency, especially in the high spawning density areas of Reaches 1 and 2 (Bradford and Hankin 2012). This hypothesis would be supported if the number of redds surveyed in an area was not proportional to the number of spawned female carcasses found in that same area. To determine if this occurred, the estimates of spawned female Chinook Salmon carcasses were compared with corresponding counts of Chinook Salmon redds from Reaches 1 and 2. These values were log-transformed and analyzed using linear regression. These two variables would be considered proportional if the slope of their linear relationship was not significantly different than '1'. A slope that is significantly different than '1' would indicate that these variables are not proportional and some density-dependent observer error could be inferred.

Trends in Redd Abundance and Distribution

Data from 2017 were combined with the preceding fifteen years (2002–2016) of mainstem Trinity River redd data from Chamberlain et al. (2012) and Rupert et al. (2017a, 2017b) for long-term analyses of redd abundance and distribution. Past years' data availability was sometimes limited since not all variables analyzed were previously collected (i.e., spatially explicit redd data are not available for Reaches 12–14 prior to 2007). Redd abundance and distribution were analyzed at three spatial scales: the system (~50–100 km sections), reach (~10–20 km sections), and site (~1–2 km sections) scales. The 2017 data were examined and, when applicable, included with previous years' data for multi-year trend analyses.

For spatial analyses, the river was partitioned into individual segments based on morphology and referred to as 'riffle units' (Rupert et al. 2017b). A riffle unit is defined as a section of river that corresponds to a singular pool–riffle–pool sequence that typically ranges between 0.1 and 0.5 km in length. These units were delineated by this sequence for redd abundance analyses because Chinook Salmon typically build redds in patches proximate to riffle crests. Therefore, riffle units generally contain an undivided group of redds. Riffle unit designations were based on the 'morphological units' delineated by Gaeuman et al. (2016). Where Gaeuman et al. (2016) used hydraulic controls (i.e., riffles) to delineate morphological units, the deepest locations (i.e., pools) between these hydraulic controls were used to split riffle units. As a result, the morphological units from Gaeuman et al. (2016) were shifted slightly upstream. Aerial photography was used to construct riffle units downstream of the restoration reach (excluding Reaches 8 and 11) because the morphological units developed by Gaeuman et al. (2016) were limited to the restoration reach. In total, the mainstem Trinity River was divided into 482 riffle units.

The riffle unit method described in this report refers to the method used for partitioning the river in Rupert et al. (2017b). In Rupert et al. (2017a), the smallest spatial units were based on contiguous 400-m (and occasionally 200-m) sections of the Science Advisory Board dataframe (SAB units; Buffington et al. 2014). This change in methodology is an improvement over that used in Rupert et al. 2017a because redd groupings are no longer split and the three spatial scale sections better reflect local spawning habitat and TRRP channel rehabilitation sites or suites of sites. The upstream and downstream site-, reach-, and system-scale section boundaries changed slightly as a result to reflect the newer riffle

unit divisions. The complete 2002–2017 data set was analyzed using the newer riffle unit-based divisions at each spatial scale.

Contiguous groups of riffle units were combined to create the sections used for the site-scale analysis (Table 2). These site designations were generally based on the TRRP site designations of the Science Advisory Board dataframe (Buffington et al. 2014). However, the total count of site-scale units was reduced from 57 to 44 by merging the smallest site-scale sections of the SAB dataframe into the most appropriate adjacent site-scale sections. This spatial scale was used to evaluate changes in natural- and hatchery-origin Chinook Salmon redd abundance at a scale similar to TRRP restoration sites or suites of sites. Changes in spawning abundance within these sites was analyzed using linear regression of the annual proportion (number of redds in the site / sum of redds in the restoration reach) of redds.

Ten reach-scale sections were also used to evaluate long-term trends in natural- and hatchery-origin Chinook Salmon redd abundance (Figure 2, Table 3). These reaches consisted of groups of sites and were intended to evaluate redd abundance at a spatial scale that was an intermediate between the system and site scales. Our reach-scale designations closely resemble those defined by HVT et al. (2011), who partitioned the restoration reach into five ‘rehabilitation reaches’ that were delineated by differences in hydrology and sediment supply characteristics. Boundaries of the other five river sections downstream of the restoration reach were set similarly. Changes in spawning abundance within these reaches were analyzed using linear regression analyses of both the annual number and proportion (number of redds in reach / sum of redds in all reaches) of natural- and hatchery-origin Chinook Salmon redds.

Changes in redd abundance and distribution at the system scale were evaluated over the entire mainstem and also separately for the restoration reach (Reaches 1–7) and remaining surveyed river downstream of the restoration reach (Reaches 9–10 and 12–14). Linear models were used to detect trends in redd abundance. Mean distance from Lewiston Dam of natural- and hatchery-origin Chinook Salmon redds built upstream of Cedar Flat were evaluated using linear regression models.

Table 2. The reach- and site-scale sections used for redd abundance and distribution analysis within the restoration reach. Sites are listed with the approximate location of their upstream boundary, shown as distance from the Klamath River confluence (rkm).

| Reach | Site (rkm) | TRRP Rehabilitation | Length (km) |
|---------------|----------------------------------|---------------------|-------------|
| Lewiston | Hatchery (182.20) | 2006 | 0.69 |
| | Sven Olbertson (181.51) | 2008 | 1.28 |
| | Old Bridge (180.22) | 2008 | 1.75 |
| | Sawmill (178.47) | 2009 | 1.60 |
| | Upper Rush Creek (176.87) | | 1.46 |
| Limekiln | Lower Rush Creek (175.41) | | 1.33 |
| | Dark Gulch (174.08) | 2008 | 2.81 |
| | Lowden Ranch (171.27) | 2010 | 1.73 |
| | Trinity House Gulch (169.54) | 2010 | 0.72 |
| | Tom Lang Gulch (168.82) | | 1.48 |
| | Poker Bar (167.34) | | 2.30 |
| | China Gulch (165.05) | | 1.47 |
| | Limekiln Gulch (163.57) | 2015 | 2.38 |
| | Steel Bridge (161.20) | | 1.67 |
| | McIntyre Gulch (159.53) | | 1.53 |
| | Vitzthum Gulch (158.00) | 2007 | 2.02 |
| | Upper Indian Creek (155.98) | 2007 | 0.56 |
| Douglas City | Lower Indian Creek (155.42) | 2007 | 1.52 |
| | Upper Douglas City (153.90) | 2007, 2015 | 0.83 |
| | Douglas City (153.07) | 2013 | 1.30 |
| | Reading Creek (151.77) | 2010 | 1.77 |
| | Upper Steiner Flat (150.00) | | 1.26 |
| | Lower Steiner Flat (148.74) | 2012 | 1.90 |
| | Lorenz Gulch (146.83) | 2013 | 1.49 |
| | The Canyon (upstream) (145.34) | | 2.17 |
| Junction City | The Canyon (downstream) (143.18) | | 2.23 |
| | Dutch Creek (140.95) | | 2.56 |
| | Evan's Bar (138.38) | | 1.28 |
| | Soldier Creek (137.11) | | 0.89 |
| | Chapman Ranch (136.22) | | 1.10 |
| | Deep Gulch (135.13) | | 1.11 |
| | Sheridan Creek (134.02) | | 1.15 |
| | Oregon Gulch (132.87) | | 0.76 |
| | Sky Ranch (132.12) | | 1.20 |
| | Upper Junction City (130.91) | 2012 | 0.89 |
| | Lower Junction City (130.01) | 2014 | 0.67 |
| North Fork | Hocker Flat (129.34) | 2005 | 1.88 |
| | Upper Conner Creek (127.46) | | 1.12 |
| | Conner Creek (126.34) | 2006 | 1.71 |
| | Wheel Gulch (124.63) | 2011 | 1.05 |
| | Valdor Gulch (123.58) | 2006 | 1.84 |
| | Elkhorn (121.74) | 2006 | 1.50 |
| | Pear Tree Gulch (120.24) | 2006 | 1.33 |
| | Bagdad (118.92) ^a | | 1.52 |

^a the downstream boundary of the Bagdad site was at rkm 117.4




Figure 2. The ten sections of the mainstem Trinity River used for reach-scale analyses of Chinook Salmon redd distribution.

Table 3. River sections [with river kilometer (rkm)] used for the reach-scale analysis of redd abundance.

| Section | Boundaries | | Length (km) |
|---------------------|--------------------------------------|---------------------------------------|-------------|
| | Upstream (rkm) | Downstream (rkm) | |
| Lewiston Rehab | Lewiston Dam (182.20) | Rush Creek (175.41) | 6.79 |
| Limekiln Rehab | Rush Creek | Indian Creek (155.42) | 19.99 |
| Douglas City Rehab | Indian Creek | Browns Creek (143.18) | 12.25 |
| Junction City Rehab | Browns Creek | Canyon Creek (129.34) | 13.84 |
| North Fork Rehab | Canyon Creek | North Fork Trinity River (117.40) | 11.94 |
| Big Bar | Big Flat access riffle unit (107.82) | Del Loma access riffle unit (94.03) | 13.79 |
| Del Loma | Del Loma access riffle unit | Cedar Flat access riffle unit (79.31) | 14.72 |
| Salyer Gorge | Hawkins Bar river access (63.76) | South Fork Trinity River (50.33) | 13.41 |
| Willow Creek Valley | South Fork Trinity River | Tish Tang a Tang Creek (26.95) | 23.40 |
| Hoopa Valley | Tish Tang a Tang Creek | Weitchpec (Trinity River mouth; 0.0) | 26.95 |

Results

Survey Success and Conditions

Crews were able to complete 86% of the originally scheduled surveys in 2017, including missed surveys that were rescheduled for the following week (Appendix A). The first scheduled surveys on Reaches 4–7 were cancelled due to wildfires causing smoky air conditions and road and river access closures. Other missed surveys, which were mostly for Reach 6 and downstream from mid-November to early December, were usually cancelled due to rain events causing increased turbidity and poor visibility. Additionally, surveys on Reaches 1–4 and 13 were completed the week of December 17, which was one week more than initially scheduled.

Trinity River discharge at Lewiston, California, was about 13.1 m³/s during the first half of the survey season before dropping to about 9.0 m³/s in mid-October, at which it remained for the remainder of season (Appendix B). At Hoopa, California, mean daily flows on the mainstem Trinity River ranged between 18.2 and 31.4 m³/s from the start of the survey season to early November before rain events caused flows to increase in mid-November. Mean daily flow peaked at 277.5 m³/s on November 21 before coming back down to about 36.0 m³/s by mid-December.

Crews reported water visibility between 1.5 and 3.0 m during most of the surveys in 2017 (Appendix A). Visibility was occasionally higher (>3.0), particularly in the lower reaches. Visibility was lower (0.9–1.5 m) during some early season surveys and less than 0.9 m once in Reach 9 in early September after a project in Sheridan Creek temporarily increased turbidity.

Salmon Carcasses

During the 2017 surveys, 527 fresh (conditions 1 and 2 as described in Rupert et al. 2017a) Chinook Salmon carcasses were examined (Table 4). Of these fresh carcasses, 333 (63.4%) were females, 39 (7.4%) were adipose fin-clipped ('ad-clip'), and 32 (6.1%) had been marked with a spaghetti tag at the Willow Creek or Junction City weir operated by the California Department of Fish and Wildlife. Chinook Salmon released from the TRH are batch-marked with coded-wire tags (CWT) and externally marked using an ad-clip at a constant fractional mark rate of about 25%. From the 39 ad-clipped fresh Chinook Salmon carcasses observed, 31 head samples were collected (Table 5). Data from CWT recoveries yielded an average annual production multiplier (i.e., tagging rate) of 0.240 in 2017.

Of the 333 fresh female Chinook Salmon carcasses recovered, 25 (7.5%) were ad-clipped, and of these, 20 heads were collected. CWTs were recovered and read from all 20 (100%) of these heads. Of the spawned female hatchery-origin Chinook Salmon carcasses (spring and fall broods combined) with associated CWT data, 90% (18 of 20) were recovered within 10 km of Lewiston Dam (Figure 3).

Relatively few (six) Coho Salmon carcasses were recovered during the 2017 surveys (Table 6). Of these, three were fresh and of these, none (0%) were right maxillary-clipped, which would indicate hatchery origin. Only one of the Coho Salmon carcasses was a fresh spawned female. The limited number of spawned female Coho Salmon carcasses recovered inhibited the ability to differentiate Coho Salmon redds by origin in 2017.

Table 4. Summary of fresh (conditions 1 and 2) Chinook Salmon carcass data by survey reach, 2017 Trinity River surveys.

| Reach | Total | Males | Females | Female proportion | Ad-clipped | Weir-tagged |
|-------|------------------|-------|---------|-------------------|-----------------|-------------|
| 1 | 120 ^a | 33 | 85 | 72.0% | 17 | 9 |
| 2 | 119 | 43 | 76 | 63.9% | 13 | 9 |
| 3 | 76 | 38 | 38 | 50.0% | 3 | 4 |
| 4 | 38 | 18 | 20 | 52.6% | 1 | 1 |
| 5 | 53 | 24 | 29 | 54.7% | 4 | 3 |
| 6 | 62 | 18 | 44 | 71.0% | 1 | 1 |
| 7 | 20 | 6 | 14 | 70.0% | 0 | 0 |
| 9 | 25 | 9 | 16 | 64.0% | 0 | 3 |
| 10 | 13 | 3 | 10 | 76.9% | 0 | 2 |
| 12 | 0 | 0 | 0 | - | 0 | 0 |
| 13 | 1 | 0 | 1 | 100.0% | 0 | 0 |
| 14 | 0 | 0 | 0 | - | 0 | 0 |
| Total | 527 ^a | 192 | 333 | 63.4% | 39 ^b | 32 |

^a includes two carcasses of unknown sex^b head samples were collected from 31 of the 39 fresh ad-clipped Chinook Salmon carcasses

Table 5. Coded-wire tag (CWT) information retrieved from fresh adipose fin-clipped Chinook Salmon carcasses, 2017 Trinity River surveys.

| Carcasses | CWT | Brood Year | Run type | Release type | Production multiplier | Production multiplier |
|-----------|------------------------|------------|----------|---------------------|-----------------------|-----------------------|
| 1 | 060605 | 2013 | Spring | Advanced fingerling | 4.24 | 0.236 |
| 1 | 060606 | 2013 | Spring | Advanced fingerling | 4.15 | 0.241 |
| 1 | 060609 | 2013 | Fall | Fingerling | 4.12 | 0.243 |
| 2 | 060612 | 2013 | Spring | Yearling | 4.22 | 0.237 |
| 1 | 060615 | 2014 | Fall | Fingerling | 4.13 | 0.242 |
| 2 | 060689 | 2014 | Spring | Advanced fingerling | 4.27 | 0.234 |
| 1 | 060691 | 2014 | Spring | Advanced fingerling | 4.14 | 0.242 |
| 6 | 060692 | 2014 | Fall | Advanced fingerling | 4.09 | 0.244 |
| 5 | 060693 | 2014 | Fall | Advanced fingerling | 4.08 | 0.245 |
| 1 | 060694 | 2014 | Fall | Fingerling | 4.28 | 0.233 |
| 1 | 060696 | 2014 | Spring | Yearling | 4.27 | 0.234 |
| 2 | 060697 | 2014 | Fall | Yearling | 4.18 | 0.239 |
| 1 | 060775 | 2015 | Fall | Fingerling | 4.27 | 0.234 |
| 4 | 060780 | 2015 | Fall | Yearling | 4.25 | 0.236 |
| 1 | 068849 | 2013 | Spring | Fingerling | 4.18 | 0.239 |
| 1 | -- Missing CWT/head -- | | | | NA | NA |
| | | | | | Mean = 4.17 | Mean = 0.240 |




Figure 3. Distribution of coded-wire-tagged (CWT) spawned female Chinook Salmon carcasses by brood type (spring and fall) located in the mainstem Trinity River downstream of Lewiston Dam in 2017.

Table 6. Summary of fresh (conditions 1 and 2) Coho Salmon carcass data by survey reach, 2017 Trinity River surveys.

| Reach | Total | Males | Females | Female proportion | Maxillary-clipped | Weir-tagged |
|-------|-------|-------|---------|-------------------|-------------------|-------------|
| 1 | 1 | 1 | 0 | 0.0% | 0 | 0 |
| 2 | 0 | 0 | 0 | - | 0 | 0 |
| 3 | 2 | 1 | 1 | 50.0% | 0 | 0 |
| 4 | 0 | 0 | 0 | - | 0 | 0 |
| 5 | 0 | 0 | 0 | - | 0 | 0 |
| 6 | 0 | 0 | 0 | - | 0 | 0 |
| 7 | 0 | 0 | 0 | - | 0 | 0 |
| 9 | 0 | 0 | 0 | - | 0 | 0 |
| 10 | 0 | 0 | 0 | - | 0 | 0 |
| 12 | 0 | 0 | 0 | - | 0 | 0 |
| 13 | 0 | 0 | 0 | - | 0 | 0 |
| 14 | 0 | 0 | 0 | - | 0 | 0 |
| Total | 3 | 2 | 1 | 33.3% | 0 | 0 |

Carcass Estimates

The hierarchical latent variables model estimated 366 (95% CI: 277–499) Chinook Salmon carcasses in Reach 1 and 498 (95% CI: 356–735) in Reach 2 in 2017. Estimates of spawned female Chinook Salmon carcasses were 250 (95% CI: 186–353) in Reach 1 and 316 (95% CI: 218–475) in Reach 2.

Pre-spawn Mortality

Six fresh unspawned female Chinook Salmon carcasses were found in 2017, all without a hatchery mark, which yielded a pre-spawn mortality rate among female Chinook Salmon throughout the mainstem Trinity River of 1.8% (Table 7). Weekly pre-spawn mortality rates ranged from 0.0% to 8.0% (the first six survey weeks were combined, as were the final three, due to small sample sizes; Figure 4). Annual pre-spawn mortality of female Chinook Salmon in the Trinity River restoration reach was 2.0% in 2017.

The lone (one) fresh female Coho Salmon carcass encountered in 2017 was of natural-origin and had spawned (Table 8). Note that pre-spawn mortality rates were based on data collected through late December, while Coho Salmon are still spawning.

Table 7. Pre-spawn mortality rates of Chinook Salmon in the Trinity River below Lewiston Dam (Reaches 1–14) and in the restoration reach (Reaches 1–7), 2009–2017 surveys. Pre-spawn mortalities by week and reach for unmarked and ad-clipped Chinook Salmon are presented in Appendix C.

| Year | Reaches 1-14 | Reaches 1-7 |
|------|---------------------------------|------------------------------|
| | (Lewiston Dam to Klamath River) | (Lewiston Dam to North Fork) |
| 2009 | 7.9% | 6.8% |
| 2010 | 10.2% | 9.5% |
| 2011 | 4.6% | 4.6% |
| 2012 | 2.4% | 2.4% |
| 2013 | 5.1% | 6.1% |
| 2014 | 11.5% | 9.1% |
| 2015 | 0.8% | 0.0% |
| 2016 | 0.7% | 0.8% |
| 2017 | 1.8% | 2.0% |




Figure 4. Weekly pre-spawn mortality from fresh (conditions 1 and 2) female Chinook Salmon carcasses, Trinity River surveys 2017. Calendar weeks 36–40 and 48–51 were combined because sample sizes were low in at least one of those weeks.

Table 8. Pre-spawn mortality rates of natural- and hatchery-origin Coho Salmon, Trinity River surveys, 2009–2017. Note that these pre-spawn mortality rates were based on data only collected through late December. Spawning success often varies, typically improving over time, and our surveys did not extend over the entire Coho Salmon spawning period.

| Year | Natural-origin | Hatchery-origin | Combined |
|------|--------------------|--------------------|--------------------|
| 2009 | 7.1% | 20.3% | 16.1% |
| 2010 | 21.9% | 16.2% | 17.0% |
| 2011 | 6.1% | 15.1% | 11.6% |
| 2012 | 3.6% | 11.8% | 10.4% |
| 2013 | 10.7% | 6.1% | 6.6% |
| 2014 | 35.1% | 28.5% | 29.8% |
| 2015 | 33.3% ^a | 50.0% ^a | 40.0% ^a |
| 2016 | 0.0% ^b | 0.0% ^b | 0.0% ^b |
| 2017 | 0.0% ^c | - | 0.0% ^c |

^a the sample size for Coho Salmon was only five carcasses in 2015

^b the sample size for Coho Salmon was only two carcasses in 2016

^c the sample size for Coho Salmon was only one carcass in 2017

Salmon Redds

During the 2017 surveys, 1,982 salmon redds were identified (Table 9). A majority of the redds (1,600; 80.7%) were estimated to have been constructed by natural-origin female Chinook Salmon, while hatchery-origin female Chinook Salmon accounted for 348 (17.6%) of the total redd count (Table 10). Coho Salmon redds accounted for 34 (1.7%) of the surveyed redds. The low numbers of spawned female Coho Salmon carcasses collected in 2017 precluded the differentiation of hatchery- and natural-origin Coho Salmon redds. Note that Coho Salmon spawning continued beyond our survey season, and our estimates of Coho Salmon redds are included only to differentiate them from Chinook Salmon redds.

Natural-origin Chinook Salmon redds were constructed throughout most of the mainstem Trinity River in 2017, though the lowest numbers were in the downstream-most reaches (Figure 5). Hatchery-origin Chinook and Coho (both origin types) salmon redds were consistently skewed toward Lewiston Dam. Little to no spawning by hatchery-origin Chinook Salmon or Coho Salmon was detected downstream of Reach 7.

Table 9. Redd counts (before species differentiation) by week and reach, Trinity River surveys 2017. NS = No Survey for scheduled surveys that were missed. Dashes (-) represent days when surveys were not scheduled.

| Week start | Reach | | | | | | | | | | | | |
|---------------|-------|-----|-----|-----|-----|-----|-----|-----------------|-----------------|-----------------|----|----|-------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 | 10 | 12 | 13 | 14 | Total |
| Aug. 27 | 0 | - | - | - | - | - | - | - | - | - | - | - | - |
| Sep. 3 | 1 | 0 | 0 | NS | NS | NS | NS | - | - | - | - | - | 1 |
| Sep. 10 | 6 | 3 | 1 | 1 | 2 | 1 | 0 | 0 | 0 | - | - | - | 14 |
| Sep. 17 | 13 | 13 | 3 | 10 | 15 | 2 | 1 | - | - | - | - | - | 57 |
| Sep. 24 | 8 | 13 | 16 | 34 | 32 | 18 | NS | 3 | 3 | - | - | - | 127 |
| Oct. 1 | 22 | 52 | 21 | 29 | 24 | 44 | NS | - | - | 0 | 0 | 0 | 192 |
| Oct. 8 | 6 | 14 | 26 | 21 | 41 | 53 | 60 | 122 | 3 | - | - | - | 346 |
| Oct. 15 | 16 | 15 | 21 | 17 | 25 | 37 | 17 | - | - | 16 | 2 | NS | 166 |
| Oct. 22 | 8 | 5 | 4 | 17 | 31 | 21 | 54 | 78 | NS | - | - | - | 218 |
| Oct. 29 | 8 | 6 | 13 | 5 | 43 | 15 | 26 | - | - | 16 | 32 | 17 | 181 |
| Nov. 5 | 16 | 8 | 19 | 10 | 15 | 3 | 22 | 111 | 96 | - | - | - | 300 |
| Nov. 12 | 21 | 25 | 14 | 7 | 8 | 3 | 7 | - | - | NS | NS | NS | 85 |
| Nov. 19 | 51 | 18 | 16 | NS | NS | 1 | NS | NS ^a | NS ^a | - | - | - | 86 |
| Nov. 26 | 21 | 19 | 17 | 10 | 10 | 4 | 2 | 44 | 27 | NS ^a | NS | NS | 154 |
| Dec. 3 | 8 | 8 | 3 | 0 | 5 | 2 | 0 | NS ^a | NS ^a | 6 | - | - | 32 |
| Dec. 10 | 5 | 4 | 0 | 0 | 0 | NS | NS | 1 | 6 | NS ^a | 3 | 2 | 21 |
| Dec. 17 | 0 | 0 | 1 | 0 | - | - | - | - | - | 1 | 0 | - | 2 |
| Total | 210 | 203 | 175 | 161 | 251 | 204 | 189 | 359 | 135 | 39 | 37 | 19 | 1,982 |

^a missed survey rescheduled for the following week

Table 10. Estimated numbers and bootstrap-generated 95% confidence intervals of salmon redds by species and origin observed in the mainstem Trinity River, 2017. Natural- and hatchery-origin estimates are for the maternal first generation only.

| Species | Origin | Redd estimate | 95% confidence limits | |
|--------------------------|----------|--------------------|-----------------------|-------|
| | | | Lower | Upper |
| Chinook Salmon | All | 1,948 ^b | - | - |
| | Natural | 1,600 | 1,435 | 1,762 |
| | Hatchery | 348 | 186 | 348 |
| Coho Salmon ^a | All | 34 ^b | - | - |
| | Natural | NA ^c | - | - |
| | Hatchery | NA ^c | - | - |

^a The survey season only partially covers the Coho Salmon spawning period

^b Confidence intervals are generated with both Chinook and Coho salmon data. Not enough female Coho Salmon carcasses were found in 2017 to calculate a confidence interval.

^c Not enough Coho Salmon carcasses were observed in 2017 to calculate separate estimates for natural- and hatchery-origin Coho Salmon redds.




Figure 5. Spatiotemporal distribution of mainstem Trinity River salmon redds from Lewiston Dam to Weitchpec, 2017. Surveys were not conducted in Reaches 8 (rkm 107.6–117.4) and 11 (rkm 63.4–79.1). The Coho Salmon carcass data precluded the differentiation of hatchery- and natural-origin groups. Survey day 1 = September 1.

Redd–Carcass Relationship

Chinook Salmon redds [natural log-(\ln -) transformed] and fresh spawned female Chinook Salmon carcasses (\ln -transformed) in Reaches 1 and 2 from 2012 to 2017 had a positive linear correlation ($R^2 = 0.8387$, $p < 0.001$; Figure 6). A significant difference was detected between a slope of ‘1’ and the slope of the linear regression between log-transformed Chinook Salmon redd estimates and Chinook Salmon carcass estimates (slope = 0.637, 95% CI: 0.465–0.809).

Redd Abundance and Distribution: System Scale

From 2002 to 2017, the number of mainstem salmon redds ranged between 1,671 and 7,588 redds and generally decreased over time ($R^2 = 0.2984$, $p = 0.03$; Figure 7). The number of redds constructed by natural-origin Chinook Salmon in the mainstem Trinity River also generally decreased over time, but with no significant trend ($R^2 = 0.0488$, $p = 0.4$), while the number of redds constructed by hatchery-origin Chinook Salmon trended downward ($R^2 = 0.5175$, $p < 0.001$) over this time frame.




Figure 6. Relationship between counts of \ln -transformed Chinook Salmon redds and \ln -transformed estimates of spawned female Chinook Salmon carcasses in Survey Reaches 1 and 2 (solid line), 2012–2017. The dashed line is included to represent a slope of ‘1’, which would be the slope of two perfectly proportional variables. Dotted lines represent 95% confidence limits of the linear model.




Figure 7. Estimated number of redds constructed in the entire mainstem Trinity River (left), within the restoration reach (center), and downstream (DS) of the restoration reach (right) by all Chinook Salmon (top), natural-origin Chinook Salmon (middle), and hatchery-origin Chinook Salmon (bottom) from 2002 to 2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines).

The trends in redd abundance within the restoration reach were similar to the mainstem-wide data (Figure 7). From 2002 to 2017, the number of redds constructed annually by natural- and hatchery-origin Chinook Salmon in the restoration reach were variable but trended downward ($R^2 = 0.2562$, $p < 0.05$ and $R^2 = 0.5528$, $p < 0.001$, respectively).

Downstream of the restoration reach the number of natural-origin Chinook Salmon redds constructed from 2002 to 2017 generally increased but with no significant trend ($R^2 = 0.1979$, $p = 0.07$; Figure 7). A significant decrease in hatchery-origin Chinook Salmon redds was detected downstream of the restoration reach ($R^2 = 0.4773$, $p = 0.005$), but relatively few to no redds were constructed by hatchery-origin Chinook Salmon in this section of river. From 2002 to 2006 between 33 and 72 redds per year were estimated to be constructed by hatchery-origin Chinook Salmon downstream of the restoration reach except for 2004 when none were estimated. From 2007 to 2017 between 0 and 14 redds per year were estimated to be constructed by hatchery-origin Chinook Salmon downstream of the restoration reach and only zero or one redd was estimated in 8 of those 11 years.

In the section of river from Lewiston Dam to Cedar Flat (Reaches 1–10), the mean distance from the dam of redds constructed by natural- (49.2 km) and hatchery-origin (14.2 km) Chinook Salmon were both the highest in the 16-year history of this project. From 2002 to 2016, the mean distance of redds from the dam ranged between 15.3 and 48.9 km for natural-origin and between 2.1 and 7.5 km for hatchery-origin Chinook Salmon. In this section of river, the mean distance from Lewiston Dam of natural-origin Chinook Salmon redds shifted downstream from 2002 to 2017 ($R^2 = 0.7697$, $p < 0.001$; Figure 8). This trend, to a lesser degree, was also evident for redds constructed by hatchery-origin Chinook Salmon ($R^2 = 0.2508$, $p < 0.05$), which also consistently spawned near Lewiston Dam.




Figure 8. Mean distance from Lewiston Dam of redds constructed by natural- (left) and hatchery-origin (right) Chinook Salmon females between Lewiston Dam and Cedar Flat (0–102.8 km from Lewiston Dam; Reaches 1–10) on the mainstem Trinity River, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an ‘*’ if < 0.05), and 95% confidence limits (dotted lines).

Redd Abundance and Distribution: Reach Scale

Long-term changes in natural-origin Chinook Salmon redd distribution were detected at the reach scale (~10–20 km). Redds by natural-origin Chinook Salmon most drastically trended downward in the Lewiston ($R^2 = 0.5252$, $p = 0.002$) and Limekiln ($R^2 = 0.3047$, $p = 0.03$) reaches and generally decreased, although not significantly, in the Douglas City reach from 2002 to 2017 (Figure 9). The number of redds between the Junction City and Del Loma reaches generally increased over this time period and generally decreased, although not significantly, in the Salyer Gorge, Willow Creek Valley, and Hoopa Valley reaches over the shorter time period from 2007 to 2017. To account for annual variation in run size, the proportions of natural-origin Chinook Salmon redds within each of the ten reach-scale segments relative to the annual total in the entire mainstem river were compared (Figure 10). This analysis revealed a shift in spawning distribution, where natural-origin Chinook Salmon redds decreased in the two upstream-most reaches [Lewiston ($R^2 = 0.8034$, $p < 0.001$) and Limekiln ($R^2 = 0.4771$, $p = 0.003$)], did not significantly change in the Douglas City reach, and increased in the mid-river reaches [Junction City ($R^2 = 0.5326$, $p = 0.001$), North Fork ($R^2 = 0.5184$, $p = 0.002$), Big Bar ($R^2 = 0.6798$, $p < 0.001$), and Del Loma ($R^2 = 0.7897$, $p < 0.001$) reaches]. The proportion of redds in the downstream-most reaches (Salyer Gorge, Willow Creek Valley, and Hoopa Valley) have not changed significantly.

Most hatchery-origin Chinook Salmon redds were constructed in the Lewiston rehabilitation reach (range = 72–1,888 redds/year, mean = 770 redds/year) and, to a lesser degree, in the Limekiln rehabilitation reach (range = 19–236 redds/year, mean = 84 redds/year) from 2002 to 2017. Over this time frame, the abundance of hatchery-origin Chinook Salmon redds significantly decreased in the Lewiston reach ($R^2 = 0.5648$, $p < 0.001$) and generally decreased in the Limekiln reach (Figure 11). Fewer hatchery-origin Chinook Salmon redds were found downstream of the Limekiln reach to the Del Loma reach where their redd numbers averaged between 7 and 18 per year in each reach and only changed significantly in the Del Loma reach ($R^2 = 0.2753$, $p = 0.04$). No redds were predicted to be associated with hatchery-origin Chinook Salmon downstream of the Del Loma reach.

To account for annual variation in run size, the proportions of hatchery-origin Chinook Salmon redds within each of the reaches were compared to the annual total in the entire mainstem river (Figure 12). The majority of hatchery-origin Chinook Salmon redds were consistently observed in the Lewiston reach (range = 51.7%–95.4%, mean = 82.3%) and, to a smaller degree, in the Limekiln reach (range = 3.5%–30.2%, mean = 11.5%) from 2002 to 2017. The proportion of hatchery-origin Chinook Salmon redds in the Lewiston reach generally decreased while the proportion of redds in the Limekiln reach significantly increased ($R^2 = 0.4229$, $p = 0.006$) over this time period. The mean proportion of hatchery-origin Chinook Salmon redds in each reach downstream of the Limekiln reach ranged between 0.0% and 2.2% and did not change significantly in any of the reaches (Figure 12).

Redd Abundance and Distribution: Site Scale

The proportional abundance of natural-origin Chinook Salmon within the 44 site-scale river sections show a range of long-term (2002–2017) trends. Most sites (21) did not show a significant change, 17 sites showed an increasing trend, and 6 sites showed a decreasing trend (Appendix D). The three upstream-most sites (Lewiston Hatchery, Sven Olbertson, and Old Bridge sites) underwent significant decreases in the proportion of natural-origin

Chinook Salmon redds, followed by a less drastic general decrease at the Sawmill site and significant decrease at the Upper Rush Creek site. Most sections from the Lower Rush Creek site to the Douglas City site did not significantly change. At each site downstream of the Douglas City site, from the Reading Creek site to the Bagdad site, the proportion of natural-origin Chinook Salmon redds either generally or significantly increased.

Of the 22 mechanical channel rehabilitation sites with at least five years of post-construction data, the proportional abundance of natural-origin Chinook Salmon redds trended upward at 7 sites, trended downward at 2 sites, and displayed no significant change at 13 sites (Appendix E). Similar to the long-term trends, the proportional abundance of natural-origin Chinook Salmon redds generally or significantly decreased in the upstream-most sites (Lewiston Hatchery to Sawmill sites), did not change in the middle sites (Dark Gulch to Upper Douglas City sites), and generally or significantly increased in most of the downstream-most sites (Douglas City to Pear Tree Bar sites).

Hatchery-origin Chinook Salmon redds were not distributed throughout the restoration sites and were too few or absent to merit statistical analysis at the site scale. Like at the reach scale, the proportion of hatchery-origin fish were at or close to zero at most sites below the Limekiln reach from 2002 to 2017.




Figure 9. Estimated number of mainstem Trinity River natural-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if <0.05), and 95% confidence limits (dotted lines).




Figure 10. Proportions of mainstem Trinity River natural-origin Chinook Salmon redds relative to the total mainstem count of natural-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if <0.05), and 95% confidence limits (dotted lines).




Figure 11. Estimated number of mainstem Trinity River hatchery-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if <0.05), and 95% confidence limits (dotted lines).




Figure 12. Proportions of mainstem Trinity River hatchery-origin Chinook Salmon redds relative to the total mainstem count of hatchery-origin Chinook Salmon redds within ten reach-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if <0.05), and 95% confidence limits (dotted lines).

Discussion

Redd counts from the 2017 spawning season were the second lowest since this survey's inception in 2002 and salmon carcass estimates were the third lowest. Our 2017 results are consistent with the California Department of Fish and Wildlife Chinook Salmon natural spawner escapement estimates for the Trinity River Basin, which estimated the third lowest numbers of both spring- and fall-run Chinook Salmon since 2002 (CDFW 2018a, 2018b).

Flows were generally stable throughout the survey period in the upper reaches and most of the survey period in the lower reaches. Rain events elevated water turbidity and the reduced visibility impaired the ability to detect redds and carcasses in the lower reaches from mid- to late November. Though scheduled lower river (Reaches 12–14) surveys in mid- to late November were cancelled due to high flow and poor visibility, spawning is typically sparse in these reaches and any missed redds from this section would likely have only been a minor contribution to the total redd count.

The analyses of long-term data from our spawning surveys provide insight into the dynamics of Chinook Salmon spawning activity on the Trinity River. The main themes that emerge are 1) the overall abundance of natural-origin Chinook Salmon redds did not change significantly from 2002 to 2017, 2) straying and spawning of hatchery-origin salmon is generally confined to areas near the hatchery below Lewiston Dam, 3) the spatial distribution of natural-origin Chinook Salmon spawning continues to change, and 4) pre-spawn mortality has been relatively low in recent years.

The annual natural-origin Chinook Salmon redd count from 2002 to 2017 ranged between 1,516 (in 2016) and 6,170 (in 2012). Spawner abundance was hypothesized to increase following restoration actions (TRRP and ESSA 2009), but the abundance of natural-origin Chinook Salmon redds in the mainstem Trinity River from 2002 to 2017 did not significantly change (Figure 7). Other factors (e.g., harvest, ocean conditions, in-river conditions, etc.) that influence in-river escapement may have masked any responses in spawning activity to river restoration. Shifts in abundance are common to Chinook Salmon populations (Mantua et al. 1997; Brown 2002) and are evident in the Klamath Basin (CDFW 2018a, 2018b). The estimates of Trinity River natural-spawner adult escapement (2,532 spring-run and 6,072 fall-run; CDFW 2018a, 2018b) in 2017 were notably below the TRRP annual river escapement goal of 68,000 natural-origin Chinook Salmon spawners (6,000 spring-run adults and 62,000 fall-run adults).

Although the abundance of natural-origin Chinook Salmon redds did not show a significant trend from 2002 to 2017, the spatial distribution of redds shifted downstream. The increase in mean distance from Lewiston Dam of natural-origin Chinook Salmon redds was previously documented (Chamberlain et al. 2012; Rupert et al. 2017a, 2017b) and data collected in 2017 continue to follow this trend. This shift is consistent with the IAP's suggestion that changes in longitudinal redd distribution would happen within three to four brood cycles following restoration activities (TRRP and ESSA 2009).

The abundance of hatchery-origin Chinook Salmon redds (redds constructed by hatchery-produced females regardless of male origin) decreased significantly from 2002 to 2017, as evident in the Lewiston Reach where the majority of hatchery-origin Chinook Salmon

spawn (Figure 11). Also, even though the distribution of hatchery-origin Chinook Salmon redds has remained skewed towards the TRH (Figure 5), the proportion of hatchery-origin Chinook Salmon redds has generally decreased in the Lewiston Reach and increased in the Limekiln Reach (Figure 12). The number and release timing of hatchery-reared juvenile Chinook Salmon has remained relatively constant over these years, so the reason for the decrease in abundance of hatchery-origin Chinook Salmon redds is unclear. While IAP objectives advocate limiting the genetic interaction of hatchery- and natural-origin Chinook Salmon, and having fewer hatchery-origin Chinook Salmon redds on the spawning grounds does support these objectives, further investigations are suggested to examine the causes for this decrease in hatchery-origin Chinook Salmon redds.

Reach-scale analyses revealed the clearest resolution for analyzing spawning distribution shifts of natural-origin Chinook Salmon. The proportion of natural-origin Chinook Salmon that spawned near TRH and Lewiston Dam (Lewiston and Limekiln reaches) decreased from 2002 to 2017 and more spawned in the mid-river sections (Junction City–Del Loma reaches; Figure 10). This shift is contrary to the IAP hypothesis that redd abundance in the reaches below the North Fork Trinity River would not increase until escapement began to approach restoration goals (TRRP and ESSA 2009). TRRP restoration actions may therefore be influencing a larger portion of the Trinity River than expected. Presumably, flow management is the primary factor for the spawning distribution shift of natural-origin Chinook Salmon since the effects of flow extend downstream much further than the generally localized effects of mechanical channel rehabilitation, coarse sediment augmentation, and watershed (tributaries) restoration.

Changes in redd abundance at the site scale was specifically used to evaluate the effect of TRRP channel rehabilitation activities. Our analysis revealed no clear post-construction response at rehabilitation sites. As reported in Rupert et al. (2017a), despite being the smallest scale used in our analyses, the site scale may still be too spatially broad and too few years have passed since construction to detect responses to restoration. A positive response in the abundance of Chinook Salmon redds to channel rehabilitation may take many generations that encompass several years of geomorphic change and restoration site maturation. TRRP channel rehabilitation sites only secondarily affect spawning habitat since many constructed features are intended to increase and diversify juvenile rearing habitats and/or change the geomorphology of the site. The long-term effects of flow management, however, are intended to increase spawning habitat, though this would presumably affect all sites regardless of channel rehabilitation treatments (TRRP and ESSA 2009).

The relationship between redd counts and the estimated number of spawned female Chinook Salmon in Reaches 1 and 2 using the 2012–2017 data set indicate a density-dependent redd observation bias (Figure 6). This is contrary to the result that Rupert et al. (2017a) found with just the 2012–2014 data set. The Reach 2 data point from 2012, the largest run year, appears to have a negative influence on the slope of the regression line. Large spawning runs in the future may help validate or refute the density-dependent observation bias within this section of the river.

The importance of describing pre-spawn mortality has increased in recent years with ongoing drought conditions and associated higher risks of epizootic events. Aguilar et al. (1996) reported that pre-spawn mortality for Chinook Salmon ranged between 1.1% and 44.9% in the mainstem Trinity River above the North Fork confluence from 1978 to 1982

and 1987 to 1995. In comparison, pre-spawn mortality rates that we measured were relatively low (between 0.0% and 9.5% from 2009 to 2016 and 2.0% in 2017) in this section of the river. Salmon pre-spawn mortality rates are typically highest at the beginning of the spawning season and decrease as the season advances (Aguilar et al. 1996; Gough and Williamson 2012). Too few pre-spawn mortality Chinook Salmon carcasses (six) were observed in 2017 to conduct a temporal analysis. Aguilar et al. (1996) also reported a positive correlation between pre-spawn mortality and run size for Trinity River Chinook Salmon from 1978 to 1995. After adding the data from 2017, which had the second lowest redd count and third lowest pre-spawn mortality rate since 2009, to the data from 2009 to 2016, no correlation was detected between these two parameters in the restoration reach (Appendix F). The lack of correlation suggests that other factors beyond run size (i.e., river conditions, run timing, etc.) may be influencing pre-spawn mortality rates. The 2017 Coho Salmon run size was notably small and the carcasses sample size ($n = 1$ fresh female) was inadequate to assess pre-spawn mortality for this species. Interpretation of results pertaining to spawning success should take into account that pre-spawn mortality occurs outside of the temporal and spatial extent of the surveys. Pre-spawn mortality fish are available to our carcass survey because they expired prior to spawning. The spatiotemporal location of carcass recovery is unlikely to be an accurate depiction of when and where fish were destined to spawn had they survived. For instance, pre-spawn mortality occurring in the Lower Klamath River for Trinity River-bound fish were not detectable during our Trinity River spawn surveys. Likewise, spring-run Chinook Salmon that expired well before the first surveys in September were also undetectable.

Acknowledgements

We extend special gratitude to Charlie Chamberlain (USFWS) for producing the modeling scripts that provide the foundation for this project's analyses and to Derek Rupert [U.S. Bureau of Reclamation (BOR)] for his technical assistance and for the cover art on this report. Thanks to the field personnel that collected the data for this survey, including: Anne Beulke (USFS), Mike Bradford (CDFW), Todd Buxton (USFWS), Alfredo Billy Colegrove (HVT), Axel Erickson IV (YTFP), Amelia Fleitz (USFS), Sterling Fulford (USFWS), Brandt Gutermuth (BOR), Kevin Held (BOR), Kyle Hopkins (YTFP), Dionisio Hurtado (HVT), Yadao Inong (YTFP), Erik Kenas (USFWS), Chris Laskodi (YTFP), Michael Macon (USFWS), Thomas Masten (HVT), Hawk Mattz (YTFP), Albert Perry (HVT), James Rickaby (HVT), David Ruiz (HVT), Derek Rupert (USFWS), Ron Smith (CDFW), Brianna Walsh (USFWS), and William Willson (YTFP).

Literature Cited

- Aguilar, B., L. D. Davis, B. W. Collins, L. Hanson, W. Sinnen, M. Zuspan, and M. Dean. 1996. Annual Report: Trinity River Basin salmon and steelhead monitoring project, 1994–1995 season. R. M. Kano, editor. California Department of Fish and Game, Inland Fisheries Division, Sacramento, California. 223 p.
- Bradford, M., and D. Hankin. 2012. Trinity River Restoration Program adult salmonid monitoring evaluation. Prepared for TRRP. Weaverville, California. 47 p.
- Brown, E. M. 2002. 2000 salmon spawning ground surveys. Pacific Salmon Treaty Program, 538 Award Number NA77FP0445. Idaho Fish Game Report 02-33, Boise, Idaho.
- Buffington, J., C. Jordan, M. Merigliano, J. Peterson, and C. Stalnaker. 2014. Review of the Trinity River Restoration Program following Phase 1, with emphasis on the Program's channel rehabilitation strategy. Prepared by the Trinity River Restoration Program's Science Advisory Board for the Trinity River Restoration Program with assistance from Anchor QEA, LLC, Stillwater Sciences, BioAnalysts, Inc., and Hinrichsen Environmental Services.
- CDFW (California Department of Fish and Wildlife). 2018a. Klamath River Basin fall Chinook Salmon spawner escapement, in-river harvest and run-size estimates, 1978–2017. California Department of Fish and Wildlife, Arcata, Californian.
- CDFW (California Department of Fish and Wildlife). 2018b. Klamath River Basin spring Chinook Salmon spawner escapement, river harvest and run-size estimates, 1980–2017. California Department of Fish and Wildlife, Arcata, Californian.
- Chamberlain, C. D., S. Quinn, and W. Matilton. 2012. Distribution and abundance of Chinook Salmon redds in the mainstem Trinity River 2002 to 2011. U.S. Fish and Wildlife Service. Arcata Fish and Wildlife Office, Arcata Fisheries Technical Report TR 2012-16, Arcata, California.
- Gaeuman, D., R. Stewart, and T. Buxton. 2016. First steps towards a river corridor management strategy. Technical report: TR-TRRP-2016-1. Trinity River Restoration Project, Weaverville, California.
- Gough, S. A., and S. C. Williamson. 2012. Fall Chinook Salmon run characteristics and escapement for the main-stem Klamath River, 2001–2010. U.S. Fish and Wildlife Service, Arcata Fish and Wildlife Office, Arcata Fisheries Technical Report Number TR 2012–14, Arcata, California.
- HVT (Hoopa Valley Tribe), McBain and Trush, and Northern Hydrology and Engineering. 2011. Channel rehabilitation design guidelines for the mainstem Trinity River. Prepared for the Trinity River Restoration Program, Hoopa, California.
- Kery, M., and M. Schaub. 2012. Bayesian population analysis using WinBUGS. Academic Press, Oxford, United Kingdom.

- Mantua, N. J., S. R. Hare, Y. Zhang, J. M. Wallace, R. C. Francis. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society* 78(6):1069–1079.
- Moffett, J. W., and S. H. Smith. 1950. Biological investigations of the fishery resources of the Trinity River, California. Special Scientific Report No. 12, U.S. Fish and Wildlife Service. 71 p.
- Quihillalt, R. R. 1999. Mainstem Trinity River fall Chinook Salmon spawning redd survey, 1996 through 1998. U.S. Fish and Wildlife Service, Arcata, California.
- Rogers, D. W. 1972. King Salmon (*Oncorhynchus tshawytscha*) and Silver Salmon (*Oncorhynchus kisutch*) spawning escapement and spawning habitat in the upper Trinity River, 1970. California Fish and Game, Region 1, Anadromous Fisheries Branch Administrative Report Number 73-10.
- Rupert, D. L., C. D. Chamberlain, S. A. Gough, N. A. Som, N. J. Davids, W. C. Matilton, A. M. Hill, and E. R. Wiseman. 2017a. Mainstem Trinity River Chinook Salmon Spawning Survey 2012-2014. U.S. Fish and Wildlife Service. Arcata Fish and Wildlife Office, Arcata Fisheries Technical Report Number TR 2017–52, Arcata, California.
- Rupert, D. L., S. A. Gough, N. A. Som, N. J. Davids, W. C. Matilton, A. M. Hill, and J. L. Pabich. 2017b. Mainstem Trinity River Chinook Salmon Spawning Distribution, 2015 and 2016. U.S. Fish and Wildlife Service. Arcata Fish and Wildlife Office, Arcata Fisheries Technical Report Number TR 2017–56, Arcata, California.
- TRRP and ESSA (Trinity River Restoration Program and ESSA Technologies Ltd.). 2009. Integrated Assessment Plan, Version 1.0 – September 2009. Draft report prepared for the Trinity River Restoration Project, Weaverville, California. 285 p.
- USDOI (U.S. Department of Interior). 2000. Record of Decision: Trinity River mainstem fishery restoration final Environmental Impact Statement / Environmental Report.
- USFWS (U.S. Fish and Wildlife Service). 1986, 1987. Trinity River flow evaluation study, 2 reports. Annual Report, U.S. Fish and Wildlife Service, Sacramento, California.
- USFWS (U.S. Fish and Wildlife Service) and HVT (Hoopa Valley Tribe). 1999. Trinity River flow evaluation, Final Report. U.S. Fish and Wildlife Service, Arcata Fish and Wildlife Office, Arcata, California and Hoopa Valley Tribe, Hoopa, California.

Appendices



Appendix A. Trinity River water visibility by week and reach throughout the 2017 survey period. Grey boxes represent surveys with sub-optimal visibility. NS = No Survey for scheduled surveys that were missed. Dashes (-) represent days when surveys were not scheduled or performed.

| Week start | Reach | | | | | | | | | | | |
|---------------|---------|---------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|-----------------|-----------------|------|---------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 | 10 | 12 | 13 | 14 |
| Aug. 27 | 1.5-3.0 | - | - | - | - | - | - | - | - | - | - | - |
| Sep. 3 | 0.9-1.5 | 0.9-1.5 | 0.9-1.5 | NS | NS | NS | NS | - | - | - | - | - |
| Sep. 10 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 0.9-1.5 | 0.9-1.5 | <0.9 | 0.9-1.5 | - | - | - |
| Sep. 17 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 0.9-1.5 ^b | 0.9-1.5 ^b | - | - | - | - | - |
| Sep. 24 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 0.9-1.5 | NS | 0.9-1.5 | 0.9-1.5 | - | - | - |
| Oct. 1 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^a | NS | - | - | 1.5-3.0 | >3.0 | >3.0 |
| Oct. 8 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | >3.0 | >3.0 | 1.5-3.0 ^b | 0.9-1.5 | - | - | - |
| Oct. 15 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^a | 0.9-1.5 | 1.5-3.0 ^a | >3.0 | >3.0 | - | - | >3.0 | >3.0 | NS |
| Oct. 22 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | 1.5-3.0 ^b | 1.5-3.0 ^b | NS | - | - | - |
| Oct. 29 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | 1.5-3.0 ^b | - | - | >3.0 | >3.0 | >3.0 |
| Nov. 5 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | >3.0 | >3.0 | >3.0 | >3.0 | - | - | - |
| Nov. 12 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^a | 1.5-3.0 | 0.9-1.5 | 0.9-1.5 | - | - | NS | NS | NS |
| Nov. 19 | 1.5-3.0 | 1.5-3.0 | 0.9-1.5 | NS | NS | 0.9-1.5 | NS | NS ^c | NS ^c | - | - | - |
| Nov. 26 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | 1.5-3.0 ^b | 1.5-3.0 ^b | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 | >3.0 | NS ^c | NS | NS |
| Dec. 3 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | 1.5-3.0 ^b | 1.5-3.0 ^b | >3.0 | >3.0 | NS ^c | NS ^c | 1.5-3.0 | - | - |
| Dec. 10 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | 1.5-3.0 ^b | 1.5-3.0 | NS | NS | 1.5-3.0 ^b | >3.0 | NS ^c | >3.0 | 1.5-3.0 |
| Dec. 17 | 1.5-3.0 | 1.5-3.0 | 1.5-3.0 ^b | >3.0 | - | - | - | - | - | >3.0 | >3.0 | - |

^a this is the higher visibility reported by the two crews. The other crew reported visibility 0.9-1.5 m

^b this is the lesser visibility reported by the two crews. The other crew reported visibility >3.0 m

^c missed survey rescheduled for following week




Appendix B. Trinity River mean daily discharge at Lewiston (USGS Gage 11525500) and Hoopa, California (USGS Gage 11530000) during the 2017 survey season.


Appendix C. Pre-spawn mortality numbers by week and reach of unmarked and ad-clipped fresh (conditions 1 and 2) female Chinook Salmon carcasses, mainstem Trinity River surveys 2017. Also included are weekly pre-spawn mortality proportions among like mark-type carcasses. Ad-clipped carcass numbers were not expanded by CWT-specific production multipliers and are therefore about 25% of hatchery-origin carcass numbers. Likewise, unmarked carcass numbers include hatchery-origin carcasses that were not ad-clipped. ‘NS’ = no survey and dashes (-) represent a sample size of zero.

| Unmarked | | | | | | | | | | | | | | | | |
|-----------|------------------|-------|----|----|----|----|----|----|----|----|----|----|----|---|-------------|--|
| Calendar | | Reach | | | | | | | | | | | | | All reaches | |
| week | Dates | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 | 10 | 12 | 13 | 14 | n | Pct. | |
| 35 | Aug. 27 - Sep. 2 | - | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | - | - | |
| 36 | Sep. 3 - 9 | - | - | - | - | - | - | NS | NS | NS | NS | NS | NS | - | - | |
| 37 | Sep. 10 - 16 | - | - | - | - | - | - | - | - | - | NS | NS | NS | - | - | |
| 38 | Sep. 17 - 23 | 1 | 0 | - | - | 1 | - | - | NS | NS | NS | NS | NS | 2 | 40.0% | |
| 39 | Sep. 24 - 30 | - | 0 | - | - | - | - | NS | - | - | NS | NS | NS | 0 | 0.0% | |
| 40 | Oct. 1 - 7 | 0 | 0 | 0 | 0 | - | 0 | NS | NS | NS | - | - | - | 0 | 0.0% | |
| 41 | Oct. 8 - 14 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | - | 0 | NS | NS | NS | 1 | 4.2% | |
| 42 | Oct. 15 - 21 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | NS | NS | - | - | NS | 0 | 0.0% | |
| 43 | Oct. 22 - 28 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | NS | NS | NS | 1 | 1.9% | |
| 44 | Oct. 29 - Nov. 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | NS | NS | - | 0 | - | 0 | 0.0% | |
| 45 | Nov. 5 - 11 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | NS | NS | NS | 0 | 0.0% | |
| 46 | Nov. 12 - 18 | 0 | 0 | 0 | 0 | - | - | - | NS | NS | NS | NS | NS | 0 | 0.0% | |
| 47 | Nov. 19 - 25 | 1 | 0 | 0 | NS | NS | 0 | NS | NS | NS | NS | NS | NS | 1 | 3.7% | |
| 48 | Nov. 26 - Dec. 2 | 1 | 0 | 0 | 0 | - | 0 | - | - | 0 | NS | NS | NS | 1 | 3.2% | |
| 49 | Dec. 3 - 9 | 0 | 0 | 0 | - | - | - | - | NS | NS | - | NS | NS | 0 | 0.0% | |
| 50 | Dec. 10 - 16 | 0 | 0 | - | - | - | - | NS | - | - | NS | - | - | 0 | 0.0% | |
| 51 | Dec. 17 - 23 | - | 0 | - | - | - | NS | NS | - | - | - | - | - | 0 | 0.0% | |
| All weeks | | 3 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | - | - | - | 6 | 2.0% | |


| Ad-clipped | | | | | | | | | | | | | | | | |
|------------|------------------|-------|----|----|----|----|----|----|----|----|----|----|----|---|-------------|--|
| Calendar | | Reach | | | | | | | | | | | | | All reaches | |
| week | Dates | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 | 10 | 12 | 13 | 14 | n | Pct. | |
| 35 | Aug. 27 - Sep. 2 | - | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | - | - | |
| 36 | Sep. 3 - 9 | - | - | - | - | - | - | NS | NS | NS | NS | NS | NS | - | - | |
| 37 | Sep. 10 - 16 | - | - | - | - | - | - | - | - | - | NS | NS | NS | - | - | |
| 38 | Sep. 17 - 23 | - | - | - | - | - | - | - | NS | NS | NS | NS | NS | - | - | |
| 39 | Sep. 24 - 30 | - | - | - | - | - | 0 | NS | - | - | NS | NS | NS | 0 | 0.0% | |
| 40 | Oct. 1 - 7 | - | 0 | 0 | - | - | - | NS | NS | NS | - | - | - | 0 | 0.0% | |
| 41 | Oct. 8 - 14 | 0 | - | - | - | 0 | - | - | - | - | NS | NS | NS | 0 | 0.0% | |
| 42 | Oct. 15 - 21 | - | 0 | - | - | - | - | - | NS | NS | - | - | NS | 0 | 0.0% | |
| 43 | Oct. 22 - 28 | 0 | - | - | - | - | - | - | - | - | NS | NS | NS | 0 | 0.0% | |
| 44 | Oct. 29 - Nov. 4 | 0 | 0 | - | - | 0 | - | - | NS | NS | - | - | - | 0 | 0.0% | |
| 45 | Nov. 5 - 11 | - | - | - | - | - | - | - | - | - | NS | NS | NS | - | - | |
| 46 | Nov. 12 - 18 | 0 | - | - | - | - | - | - | NS | NS | NS | NS | NS | 0 | 0.0% | |
| 47 | Nov. 19 - 25 | 0 | 0 | - | NS | NS | - | NS | NS | NS | NS | NS | NS | 0 | 0.0% | |
| 48 | Nov. 26 - Dec. 2 | 0 | 0 | - | - | - | - | - | - | - | NS | NS | NS | 0 | 0.0% | |
| 49 | Dec. 3 - 9 | 0 | - | - | - | - | - | - | NS | NS | - | NS | NS | 0 | 0.0% | |
| 50 | Dec. 10 - 16 | - | - | - | - | - | - | NS | - | - | NS | - | - | - | - | |
| 51 | Dec. 17 - 23 | - | - | - | - | - | NS | NS | - | - | - | - | - | - | - | |
| All weeks | | 0 | 0 | 0 | - | 0 | 0 | - | - | - | - | - | - | 0 | 0.0% | |




Appendix D. Proportion of TRRP restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars. Note the change in y-axis scale in the Sven Olbertson site.




Appendix D (continued). Proportion of restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.




Appendix D (continued). Proportion of restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an '*' if <0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.




Appendix D (continued). Proportion of restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an ‘*’ if < 0.05), and 95% confidence limits (dotted lines).




Appendix D (continued). Proportion of restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an ‘*’ if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.




Appendix D (continued). Proportion of restoration reach natural-origin Chinook Salmon redds within site-scale sections, 2002–2017. Each plot includes a linear model with the R^2 value, p -value (noted with an ‘*’ if <0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.




Appendix E. Proportion of natural-origin Chinook Salmon redds within site-scale sections in the TRRP restoration reach that encompass mechanical channel rehabilitation locations, 2002–2017. Each plot includes a post-construction linear model with the R^2 value, p-value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.



Appendix E (continued). Proportion of natural-origin Chinook Salmon redds within site-scale sections in the TRRP restoration reach that encompass mechanical channel rehabilitation locations, 2002–2017. Each plot includes a post-construction linear model with the R^2 value, p-value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.



Appendix E (continued). Proportion of natural-origin Chinook Salmon redds within site-scale sections in the TRRP restoration reach that encompass mechanical channel rehabilitation locations, 2002–2017. Each plot includes a post-construction linear model with the R^2 value, p-value (noted with an '*' if < 0.05), and 95% confidence limits (dotted lines). The time mechanical channel rehabilitation was initiated is shown as black vertical bars.



Appendix F. Natural-origin Chinook Salmon redd counts versus estimates of pre-spawn mortality from Lewiston Dam to the North Fork confluence, Trinity River surveys, 2009–2017.