

**RANGE-WIDE MONITORING OF
THE MOJAVE DESERT
TORTOISE (*GOPHERUS
AGASSIZII*):
2019
ANNUAL REPORTING**

**PREPARED BY LINDA ALLISON
DESERT TORTOISE MONITORING COORDINATOR
U.S. FISH AND WILDLIFE SERVICE**

FEBRUARY 2020

Recommended Citation: U.S. Fish and Wildlife Service. 2020. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2019 Annual Reporting DRAFT. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

TABLE OF CONTENTS

Executive Summary	1
Introduction.....	2
Methods.....	2
Study areas and transect locations	2
Distance sampling transect completion	5
Proportion of tortoises available for detection by line distance sampling, G_0	6
Field observer training	7
Telemetry training.....	7
Distance sampling training	8
Data management, quality assurance, and quality control.....	12
Tortoise encounter rate and development of detection functions	14
Proportion of available tortoises detected on the transect centerline, $g(0)$	15
Estimates of tortoise density	16
Results.....	17
Field observer training	17
Proportion of tortoises detected at distances from the transect centerline.....	17
Quality assurance and quality control.....	21
Transect completion.....	22
Proportion of tortoises available for detection by line distance sampling, G_0	27
Tortoise encounter rates and detection functions.....	27
Proportion of available tortoises detected on the transect centerline, $g(0)$	29
Estimates of tortoise density	30
Discussion	32
Literature Cited	34

LIST OF TABLES

Table 1. Training schedule for 2019 for a) Kiva transect crew, b) GBI transect crews, and c) GBI telemetry trainees.	8
Table 2. Proportion of tortoise models detected in 2019 by crews within 1-, 2-, or 5-m of the transect centerline. Values that scored below the target of 0.90 at 1- and 2-m are highlighted. Crews 1-12 surveyed for Kiva Biological; the remaining crews surveyed for Great Basin Institute.	18
Table 3. Diagnostics for individual crews after training in 2019.	19
Table 4. Number and completion of transects in each stratum in 2019.	22
Table 5. Availability of tortoises (G_{θ}) when transects were walked in 2019 in the same or in neighboring strata.	27
Table 6. Stratum-level encounters and densities in 2019 for tortoises of $MCL \geq 180$ mm.	31

LIST OF FIGURES

Figure 1. Long-term monitoring strata (n=17) corresponding to tortoise conservation areas (USFWS, 2011) in each recovery unit.	4
Figure 2. Data flow from collection through final products.	13
Figure 3. Relationship between single-observer detections (by the leader, p) and dual-observer (team) detections, $g(\theta)$	16
Figure 4. Process for developing density estimates in 2019. For estimate (one per column), the full set of data was factored as indicated by divisions within the columns.	17
Figure 5. Detection curves for each of the 2019 Kiva crews during training. Each curve is based on a 16 km trial with approximately 100 detections.	20
Figure 6. Detection curves for each of the 2019 GBI crews during training. Each curve is based on a 16 km trial with approximately 100 detections.	21
Figure 7. Distribution of distance sampling transects and live tortoise observations in 2019 in Chocolate Mountain Aerial Gunnery Range, Joshua Tree, Pinto Mountains, and Chuckwalla in the southern part of the Colorado Desert Recovery Unit.	23
Figure 8. Distribution of distance sampling transects and live tortoise observations in 2019 in the Ivanpah Valley and Eldorado Valley strata of the Eastern Mojave Recovery Unit and in the Fenner stratum of the Colorado Desert Recovery Unit.	24
Figure 9. Distribution of transects and live tortoise observations in 2019 in the Ord-Rodman, Fremont-Kramer, and Superior-Cronese strata of the Western Mojave Recovery Unit.	25
Figure 10. Distribution of transects and live tortoise observations in 2019 in the Coyote Springs Valley and Beaver Dam Slope strata of the Northeastern Mojave Recovery Unit.	26
Figure 11. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180$ mm found by Kiva in 2019.	28

Figure 12. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180\text{mm}$ found by GBI in 2019..... 29

Figure 13. Detection pattern for the leader (p) and by the team ($g(\theta)$) based on all observations out to a given distance (x) from the centerline in 2019. Note convergence of $g(\theta)$ on 1.0 as x goes to 0. 30

ACKNOWLEDGEMENTS

Funding was provided to this collaborative project in 2019 by Arizona State BLM; the BLM California Desert District; the National Training Center, Ft. Irwin; the Clark County Desert Conservation Program; the Marine Corps Air Station, Yuma; and the Marine Corps Air Ground Command Center, Twentynine Palms.

The original design for this project and considerations for optimizing it based on new information and experience were first set out in Anderson and Burnham (1996) and Anderson et al. (2001).

Personnel from Kiva Biological Consulting (California) led by M. Bassett and K. Hayes, and from the Great Basin Institute (Nevada and the Beaver Dam Slope of Utah and Arizona) led by T. Christopher and M. Spangler conducted the field surveys. The survey in Chocolate Mountains Aerial Gunnery Range was conducted by Vernadero Group Incorporated. The field monitors from these teams who did the hard work of collecting and verifying the data were:

K. Anderson, R. Arney, M. Bassett, C. Benson, C. Boulden, V. Chaidez, T. Chizinski, S. Clegg, J. Curiak, R. DePond, A. d'Eprenesnil, M. Dipane, A. Drummer, C. Farchaus, S. Hanner, K. Hayes, L. Henzler, T. Hobbs, N. Holmes, K. Hotchkiss, S. Jones, A. Kalyn, T. Kneppers, J. Knight, K. Koller, A. LaBarre, C. Lillard, K. Lund, L. McDiffitt, S. Nelson, D. Nilsson, B. O'Brien, K. Patronik, A. Pawlicki, J. Perez-Jimenez, M. Ress, K. Richardson, B. Role, B. Scavone, B. Semone, E. Smith, T. Sparrow, S. Till, S. Treu, C. Veety, A. Wilson, T. Winqvist, C. Wooster.

B. Sparks (GBI) provided specialized training instruction for field crews. R. Patil (GBI) updated the electronic data-collection forms and procedures. M. Spangler (GBI) ran first-level quality assurance/quality control of data submitted by both field groups. M. Brenneman (Topoworks) provided independent review and post-processing of data and developed the final databases.

EXECUTIVE SUMMARY

The recovery program for Mojave desert tortoises (*Gopherus agassizii*) throughout their range in the Mojave and Colorado deserts (USFWS, 2011) requires range-wide, long-term monitoring to determine whether recovery goals are met. Specifically, will population trends within recovery units increase for a period of 25 years? In 1999, the Desert Tortoise Management Oversight Group endorsed the use of line distance sampling (Buckland et al., 2001) for estimating range-wide desert tortoise density. From 2001 to 2005 and 2007 to 2019, the USFWS has coordinated the distance sampling monitoring program for desert tortoises in 4 of the 5 recovery units. (The Upper Virgin River Recovery Unit is monitored by Utah Division of Wildlife Resources (UDWR; McLuckie et al., 2018) and will not be further addressed herein.)

This report describes quality assurance steps and final results for the 2019 monitoring effort. During the first years of the project, survey effort was directed annually at all 16 long-term monitoring strata. After agency funding was severely curtailed in 2012, the decision was made to survey only in well-funded strata to generate robust estimates rather than attempting to cover more strata in a less satisfactory manner, and this approach continued again in 2019, when crews completed 793 transects (8559.4 km) in 12 strata between 9 March and 15 May. In the course of these surveys, they reported 471 live tortoises, 400 of which were at least 180 mm midline carapace length (MCL) and used to generate density estimates.

In 2019, we surveyed 12 of the 16 strata outside Upper Virgin River. Chuckwalla, Pinto Mountains, and Superior-Cronese were the 3 strata with estimated densities less than 2.0 adult tortoises/km². Although the southern portion of the Chocolate Mountain Aerial Gunnery Range had densities similar to those in other strata (3.0 adults/km²), the northern portion had much higher densities (14.2 adults/km²), a pattern that was also seen in past years of these surveys. Other strata surveyed in 2019 (and the estimated density of adults/km²) were Beaver Dam Slope (2.0), Coyote Springs Valley (3.2), Eldorado Valley (2.3), Fenner (2.8), Fremont-Kramer (2.7), Ivanpah Valley (2.6), JoshuaTree (3.1), and Ord-Rodman (2.5). Over all strata, the encounter rate averaged 22.8 km for each adult tortoise that was observed.

These surveys are reported annually, corresponding to the reporting requirements for annual funding. However, the survey effort is not planned specifically for precise and accurate annual density estimates; it is directed at accurately describing population trends by using multiple years of density estimates in each monitored stratum. Based on these population trends, we can provide an estimate of the density in any one year that is more accurate than the density estimate based only on the data from that one year, such as the estimates in this report. The most accurate existing density estimates for each stratum are therefore those based on trend estimates through 2014 (USFWS 2015); an updated analysis with more recent years of data is anticipated in conjunction with the 2020 report.

RANGE-WIDE MONITORING OF THE MOJAVE DESERT TORTOISE 2019

INTRODUCTION

The Mojave Desert population of the desert tortoise was listed as threatened under the Endangered Species Act in 1990. This group of desert tortoises north and west of the Colorado River are now recognized as the species *Gopherus agassizii*, separate from *G. morafkai* south and east of the Colorado River (Murphy et al., 2011). The revised recovery plan (USFWS, 2011) designates five recovery units to which decisions about continued listing status should be applied. The recovery plan specifies that consideration of delisting should only proceed when populations in each recovery unit have increased for at least one tortoise generation (25 years), as determined through a rigorous program of long-term monitoring. This report describes implementation of monitoring and presents the analysis of desert tortoise density in 2019. A more thorough description of the background of the monitoring program is provided in USFWS (2015), and use of annual density estimates to describe population trends from 2004-2014, is provided in Allison and McLuckie (2018).

METHODS

Study areas and transect locations

Long-term monitoring strata (Figure 1) will be used over the life of the project to describe population trends in areas where tortoise recovery will be evaluated. These areas are called “tortoise conservation areas” (TCAs) in the recovery plan to describe designated critical habitat as well as contiguous areas with potential tortoise habitat and compatible management. The area associated with each critical habitat unit (CHU) is generally treated as one monitoring stratum, although the portion of Mormon Mesa CHU that is associated with Coyote Springs Valley is treated as a separate stratum. Chuckwalla CHU is also treated as dual monitoring strata, with potentially unequal sampling effort in the areas managed by the Department of Defense (Chocolate Mountain Aerial Gunnery Range, CMAGR) and by the Bureau of Land Management (BLM). New recovery units were established under the revised recovery plan (USFWS, 2011), which led to separating the Piute and Eldorado Valleys into 2 distinct strata which are in different recovery units. Fenner Valley is in the same recovery unit but is a distinct stratum from Piute Valley to simplify reporting by state. The Joshua Tree stratum does not encompass all suitable habitat for desert tortoises in Joshua Tree National Park (JTNP). The national park designation and current boundaries just post-date the designation of CHUs, so some of the Pinto Mountains and Chuckwalla CHUs (and monitoring strata) are in the current JTNP.

In 2019, surveys were conducted in California in AG, CK, FE, FK, IV, JT, OR, PT, and SC strata; and in BD, CS, and EV in Nevada, Utah, and Arizona. The optimal number of transects in

a monitoring stratum was determined by evaluating how these samples would contribute to the precision of the annual density estimate for a given stratum (Anderson and Burnham, 1996; Buckland et al., 2001). Power to detect an increasing population size is a function of 1) the magnitude of the increasing trend, 2) the “background noise” against which the trend operates, and 3) the length of time the trend is followed (even a small annual population increase will result in a noticeably larger population size if the increase continues for many years).

Anderson and Burnham (1996) recommended that transect number and length be chosen to target precision reflected in a coefficient of variation (CV) of 10-15% for the estimate of density in each recovery unit. The CV describes the standard deviation (a measure of variability) as a proportion of the mean and is often converted to a percentage. The target CV is achieved based on the number of tortoises that might be encountered there (some strata have higher densities than others). Operationally for this species, this typically entails surveying sufficient kilometers to encounter approximately 30 tortoises in each stratum.

The actual number of transects assigned in each stratum was a function of the optimal numbers described above, as well as on available funding. Transects were selected from among a set of potential transects laid out systematically across strata, with a random origin that was established in 2007 for the lattice of transects. Systematic placement provides more even coverage of the entire stratum, something that may not occur when strictly random placement of transects is used. Once the number of transects to survey in each stratum was determined, these were selected using randomization procedures; since 2013 R software has been used to implement the Generalized Random Tessellated Stratified (GRTS) spatially balanced survey design procedure (R Core Team, 2018; Kincaid and Olsen, 2017). The US Environmental Protection Agency developed GRTS as a means to generate a spatially balanced, random sample (Stevens and Olsen, 2004). Each year GRTS was used to select planned transects with these qualities and to select a set of alternative transects that would contribute to the final sample having the same spatially representative and random properties if any planned transects were replaced due to field logistics. Because the same set of potential transects has been used since 2007, some transects are repeated between years but others may not have been selected in the past.




Figure 1. Long-term monitoring strata (n=17) corresponding to tortoise conservation areas (USFWS, 2011) in each recovery unit. Stratum abbreviations are given in Table 6. Potential habitat (Nussear et al., 2009) is overlain on the southwestern United States in the extent indicator.

Distance sampling transect completion

One adaptation that tortoises have for living in the desert is to restrict surface activity to fairly narrow windows of time during the year. In general, tortoises emerge from deep within shelters (burrows) from mid-March through mid-May and then again (less predictably) in the fall. These periods coincide with flowering of their preferred food plants (in spring) and with annual mating cycles (in fall). The annual range-wide monitoring effort is scheduled to match the spring activity period for tortoises.

During this season, not all tortoises are above ground or visible in burrows. To encounter as many tortoises as possible, monitoring is scheduled for early in the day and to be completed before the hottest time of day. Because tortoises are located visually, monitoring is restricted to daylight hours. Based on past experience, we expect tortoises to become most active after 8am during March (it is usually too cool before this time), but to emerge earlier and earlier until their optimal activity period includes sunrise by the beginning of May. In May, we also expect daytime temperatures to limit tortoise above-ground activity as the morning progresses to afternoon.

Field crews completed transects during this optimal period each day. Start times were decided a week in advance, so crews arrived at transects at similar times on a given morning. However, completion times will be more variable, as a consequence of terrain, number of tortoises encountered, etc. Under normal conditions, each team walked one 12-km square transect each day. Teams were comprised of two field personnel who switched lead and follow positions at each corner of each transect, so they each spent an equal amount of time in the leader and follower positions. The leader walked on the designated compass bearing while pulling a 25 m length of durable cord; the walked path is also the transect centerline and was indicated by the location of the cord. The length of cord also spaced the two observers, guiding the path of the follower; when the cord was placed on the ground after a tortoise or carcass was detected, it facilitated measurement of the local transect bearing. The walked length of each transect was calculated as the straight-line distance between GPS point coordinates that were recorded at approximate 500 m intervals (waypoints) along the transect and/or whenever the transect bearing changed. Leader and follower each scanned for tortoises independently without leaving the centerline, and the role of the crew member finding each tortoise was recorded in the data. Although the leader saw most of the tortoises, the role of the follower was to see any remaining tortoises near the centerline, crucial to unbiased estimation of tortoise densities.

Distance sampling requires that distance from the transect centerline to tortoises is measured accurately. When a tortoise was observed, crews 1) used a compass to determine the local transect bearing based on the orientation of the 25 m centerline, 2) used a compass to determine the bearing from the point of observation to the tortoise, and 3) used a measuring tape to determine the distance from the observer to the tortoise. These data are sufficient to calculate the

perpendicular distance from the observed tortoise to the local transect line. If the tortoise was outside of a burrow, it was handled enough to measure midline carapace length (MCL), to determine its sex, assess its body condition (USFWS, 2012a), and to apply a small numbered tag to one scute. If a tortoise could not be measured because it was in a burrow, because temperatures precluded handling, or for any other reason, crews attempted to establish by other means whether the animal was at least 180 mm MCL, the criterion for including animals in density estimates.

Because transects are 3 km on one side, it is not unusual for that path to cross through varied terrain or even be blocked by an obstacle such as an interstate highway. In the first years of this program, smaller transects in inconvenient locations were shifted or replaced, but this compromised the representative nature of the sample. Since 2007, the basic rules for modifying transects involve 1) reflecting transects to avoid obstacles associated with human infrastructure or jurisdictions (large roads, private inholdings, administrative boundaries, etc.), or 2) shortening transects in rugged terrain (USFWS, 2012b). Substrate and access to transects can also make it difficult to complete transects during the optimal period of times, so 3) transects could be shortened to enable completion before 4pm each day.

If it was anticipated that fewer than 6 km could be walked due to difficult terrain, the transect was replaced with a transect from the alternate list that were also selected using the GRTS procedure. It was assumed that the proportion of the area that was unwalkable was the same as the proportion of total planned kilometers (12 X number of planned transects) that were unwalkable. Specifics of how transect paths were to be modified for rugged terrain (shortened) or for administrative boundaries (reflected) can be found online in the current version of the handbook (USFWS, 2017a).

Proportion of tortoises available for detection by line distance sampling, G_0

Basing density estimates only on the tortoises that are visible will result in density estimates that are consistently underestimated (biased low). Instead, we use telemetry to estimate the proportion of tortoises available for sampling, G_0 (“gee-sub-zero”), which was incorporated in estimate of adult tortoise density to correct this bias.

We used telemetry to locate radio-equipped tortoises that were visible as well as those that were otherwise undetectable in deep burrows or well-hidden in dense vegetation. To quantify the proportion that were available for detection (visible), telemetry technicians used a VHF radio receiver and directional antenna to locate 9-16 radio-equipped G_0 tortoises in each of 6 of the full set of focal sites throughout the Mojave and Colorado deserts (Fig. 1).

Each time a transmittered tortoise was located, the observer determined whether the tortoise was visible (*yes* or *no*). Through careful coordination, observers at telemetry sites monitored visibility during the same daily time period when field crews were walking transects in the same region of

the desert. Observers completed a survey circuit of all focal animals as many times as possible during the allotted time, recording visibility each time. Bootstrapped estimates of G_0 started by selecting one visibility record at random for each tortoise on each day it was located. The average visibility of all tortoise observations at a site on a given day was calculated and used to estimate the mean and variance of G_0 at that site. One thousand bootstrap samples were generated in Microsoft Excel to estimate G_0 and its standard error.

Field observer training

Training for careful data collection and consistency between crews is fundamental part of quality assurance for this project. This training includes instruction as well as required practice time on skills such as tortoise handling, walking practice transects, and developing detection and distance-measuring techniques on a training course with tortoise models in measured locations. Chapters of the monitoring handbook are updated as needed and posted to the Desert Tortoise Recovery Office website (http://www.fws.gov/nevada/desert_tortoise/reports).

Kiva Biological (Kiva) supplied crews for monitoring in California strata. Great Basin Institute (GBI) supplied crews for monitoring in strata in Nevada, Utah, and Arizona. Fourteen of the 24 personnel for Kiva had previous transect experience with this monitoring program, as did five of 22 surveyors for GBI. The two teams were trained separately by the same USFWS instructor for consistency. To accommodate logistics on Chocolate Mountain Aerial Gunnery Range, California surveys started approximately one month earlier than those in Nevada, so it was not practical to overlap the training schedules (Table 1).

Telemetry training

The primary goals of G_0 training include correct use of telemetry equipment, understanding G_0 data collection fields, observation of as many radio-equipped tortoises as possible during the day, and covering a window of observation that overlaps the transect observation period for each sampling area on that day. Although all telemetry crews had some prior telemetry experience, performance on this project differs from others that do not require confirmation of the exact location of the tortoise. Unless the exact location is determined, visibility of the tortoise cannot be accurately recorded. Beyond instruction and testing on use of the equipment in desert terrain, several days of practice were compulsory to be able to troubleshoot locating the tortoise and confirming the location when it could not be seen. In addition, some instruction for telemetry and transect crews overlapped to help each group better understand the purpose their data serve and how separate data types are related to the final density estimate.

Distance sampling training

Transect walkers were given classroom instruction, skills training, field demonstrations, and practice transects to complete (Table 1). Ultimately each team was evaluated based on performance on a field arena outfitted with polystyrene tortoise models placed in measured locations (Anderson et al., 2001), as well as on performance meeting protocol requirements on full-day staged transects.

Polystyrene desert tortoise models were set out on the training course each year using placement instructions (vegetation or open placement, tape-measured distance along training line, and tape-measured distance perpendicular from training line). This course was used to determine whether 1) individual teams are able to detect all models on the transect centerline, 2) whether their survey techniques yield useful detection functions, and 3) whether they can accurately report the distance of each model from the transect centerline. For each purpose, many opportunities must be provided, so the course is populated at a very high density of models (410/km²).

Crews were sent on transects and training lines as paired, independent observers. That is, the follower was 25 m behind the leader, with the opportunity to detect models not found by the leader. If the leader detected 80% of all tortoises that are found, the assumption was that the follower detected 80% of the tortoises that were missed by the leader. In this example, the pair together would detect $0.80 + (0.80 \times (1 - 0.80)) = 0.96$ of all tortoises on the centerline. These data on models were used to evaluate and correct crew performance before the field season, but were not used in any way to estimate densities of live tortoises once range-wide field surveys began.

Table 1. Training schedule for 2019 for a) Kiva transect crew, b) GBI transect crews, and c) GBI telemetry trainees.

1a. Training schedule for 2019 for Kiva transect crews

Date	Activity	Location	Instructors
Saturday, 2 March	Transect methods overview	GBI Field Station	Allison
	Phones – Training database	GBI Field Station	Spangler
	Review protocol and goals on training lines	GBI Field Station	Allison
	Compass work	GBI Field Station	Allison

Range-wide Monitoring of the Mojave Desert Tortoise: 2019

Date	Activity	Location	Instructors
	How to collect data in the styrotort arena and estimate 2-, 5-, and 15-m distances	GBI Field Station	Hayes/ Bassett
	Practice epoxy for tag attachment	GBI Field Station	Woodman
3 March	Training Lines I (8km)	BLM Desert Tortoise Mgmt Area (DTMA)	Allison
4 March	Review training line I results	GBI Field Station	Allison / Spangler
	Standard protocol and nonstandard transects, start point and reflection exercise	GBI Field Station	Allison
	Phones – Transect database	GBI Field Station	Allison
	GPS and compass use for tortoise monitoring	GBI Field Station	Allison
	Tortoise handling	GBI Field Station	Bassett/ Hayes
	Tortoise visibility examples	GBI Field Station	Allison
5 March	Training Lines II (8km)	BLM DTMA	Allison
6 March	Full transects (12km)	Large Scale Translocation Study Area (LSTS)	Hayes/ Bassett
7 March	Review training line II results	GBI Field Station	Allison / Spangler
	Review LSTS practice transect results	GBI Field Station	Allison / Spangler
	Wrap up discussion	GBI Field Station	Allison

Table 1b. Training schedule for GBI transect crews.

Date	Activity	Location	Instructors
Monday, 18 March	Transect methods overview	GBI Field Station	Allison
	Compass work	GBI Field Station	Christopher
	Training line protocol and objectives	GBI Field Station	Allison
19 March	Training Lines I (8km)	BLM DTMA	Allison
20 March	Review training line I results	GBI Field Station	Allison / Spangler
	Monitoring on Public Lands	GBI Field Station	Allison
	Standard protocol and nonstandard transects, start point and reflection exercise	GBI Field Station	Allison
	GPS and compass use	GBI Field Station	Allison
	Tortoise visibility examples	GBI Field Station	Allison
21 March	Tortoise handling	GBI Field Station	Dr. Johnson
22 March	Full 12km transect with interruption for terrain	Large Scale Translocation Site	Christopher
25 March	Search image for tortoises	River Mtns, NV	Christopher/Sparks
26 March	Training Lines II (16km)	BLM DTMA	Christopher
27 March	Training Lines II (continued)	BLM DTMA	Christopher
28 March	Review LSTS I	GBI Field Station	Allison

Range-wide Monitoring of the Mojave Desert Tortoise: 2019

Date	Activity	Location	Instructors
	Read a bearing from a map	GBI Field Station	Allison
	Plan reflected transects for LSTS 2	GBI Field Station	Allison
	Training line results – Trial II	GBI Field Station	Allison
	Handling 2	GBI Field Station	Christopher
29 March	Full transects (12km) reflected for highway	LSTS	Allison
1 April	Review LSTS transects	GBI Field Station	Allison
	Handling 3	GBI Field Station	Christopher
	Wrap up discussion	GBI Field Station	Allison

Table 2c. Training schedule for GBI telemetry technicians.

Date	Activity	Location	Instructors
5 March	Introduction to distance sampling	GBI Field Station	Allison
	Visibility descriptions	GBI Field Station	Allison
7 March	Introduction to tortoise telemetry	Boulder City Conservation Easement (BCCE)	Sparks
8 March	Telemetry practice	Halfway Wash	Sparks
9 March	Telemetry practice	BCCE	Sparks
14 March	Telemetry practice	Gold Butte focal site	Sparks
19 March	Telemetry practice	River Mountains, Nevada	Sparks
20 March	Telemetry practice	River Mountains, Nevada	Sparks
21 March			

Date	Activity	Location	Instructors
	Tortoise handling	GBI Field Station	Dr. Johnson
22 March	Telemetry practice	BCCE	Sparks
23 March	Telemetry practice	River Mountains, Nevada	Sparks
26 March	Surveyor search image for tortoises	River Mountains, Nevada	Christopher/Sparks
28 March	Handling 2	GBI Field Station	Christopher
29 March	Telemetry to derive start times	Piute-Mid focal site	Sparks
30 March	Telemetry to derive start times	Piute-Mid focal site	Sparks

Data management, quality assurance, and quality control

Two sets of data tables were maintained through the field season, organizing data collected on transects and at the G_0 focal sites. Collection data forms, paper datasheets, and databases were designed to minimize data entry errors and facilitate data verification and validation. Data were collected in both electronic and paper formats by the separate survey organizations, then combined into a single database by a single data manager provided by GBI. Data were submitted to the USFWS for evaluation at 7-14-day intervals over the course of surveys. Data were evaluated for completeness and correctness but also for consistency among crews and between field teams. Written review of the datasets was provided by USFWS to the field teams, who worked with the Phase I data manager to address and/or clarify any identified inconsistencies in the data and to ensure all crews applied the field protocols consistently.

Data quality assurance and quality control (data QA/QC, also known as verification and validation) was performed during the data collection (Phase I, described above), data integration, and data finalization phases. In each phase, processing steps were also implemented. For instance, in Phase I, datasheets were scanned and named to be easily associated with their electronic records. During the data integration phase (II), additional attribute fields were added to enable data from different UTM zones to be utilized simultaneously, and all fields were formatted for final processing. The third phase, data finalization (III), involved generation of final spatial and non-spatial data products used for analysis. Because processing steps can introduce errors, each phase of QA/QC included checks of collection but also of processing information. Figure 2 describes the overall data flow.




Figure 2. Data flow from collection through final products.

Tortoise encounter rate and development of detection functions

The number of tortoises seen in each stratum and their distances from the line were used to estimate the encounter rate (tortoises seen per kilometer walked) and the detection rate (proportion of available tortoises that are detected out to a certain distance from the transect centerline). Detection function estimation is “pooling robust” under most conditions (Buckland et al., 2001). This property holds as long as factors that cause variability in the curve shape are represented proportionately (Marques et al., 2007). Factors that can affect curve shape include vegetation that differentially obscures vision with distance and different detection protocols used by individual crews (pairs). I expected to develop one detection curve for each field team each year because each of the pairs on a team contributes the same number of transects to the effort, and because each team works in geographically different sites. The encounter rate is less sensitive to small sample sizes, so it was estimated for each stratum separately.

Program DISTANCE, Version 7, Release 3 (Thomas et al., 2010) was used to fit appropriate detection functions, to estimate the encounter rate of tortoises in each stratum, and to calculate the associated variances. Analysis was applied to all live tortoises at least 180 mm MCL. Transects were packaged into monitoring strata (“regions” in Program DISTANCE).

Observations were truncated to improve model fit as judged by the simplicity (reasonableness) of the resulting detection function estimate (Buckland et al., 2001:15-16) as well as fit diagnostics near the transect centerline. Any observations that were not used to estimate detection functions were also not used to estimate the encounter rate (tortoises detected per kilometer walked). In distance sampling applications for many other species, encounter rate can be estimated with relatively high precision, but tortoise encounter rates are low enough that truncation was applied conservatively to maximize the number of observations per stratum. Using truncated data, I considered the Akaike Information Criterion (AIC) to compare detection-function models (uniform, half normal, and hazard-rate) and key function/series expansions (none, cosine, simple polynomial, hermite polynomial) recommended in Buckland et al. (2001). To determine whether a single detection curve might be used for both survey teams, AIC was also used to compare separate models to a single one that included a factor for field team to modify the shape of the curve.

In April 2017, 366 tortoises that were at least 180 mm median carapace length (“adults”) were marked and then translocated into Ord-Rodman TCA (OR) in the Lucerne Valley just south of the Ord Mountains. This translocation was done in conjunction with the base expansion at 29 Palms Marine Corps Air Gunnery Command Center (MCAGCC). Another 2 were translocated to OR in the fall of 2017, 38 more in the spring of 2018, and 19 more in the spring of 2019 before the surveys reported here. A second recipient site, Rodman-Sunshine Peak North, is just adjacent to OR and received 301 adult translocatees through spring 2018. For this report, encounter rates were calculated in OR with all encountered animals and then again separately to report on only

resident animals. The latter density captures the population status before translocations. From experience with previous translocations, it is expected that translocatees will be much more active than resident tortoises for the first 2 years, so analyses will separate residents from translocatees through 2021.

Proportion of available tortoises detected on the transect centerline, $g(0)$

Transects were conducted by two-person crews using the method adopted beginning in 2004 (USFWS, 2006). Transects were walked in a continuous fashion, with the lead crew member walking a straight line on a specified compass bearing, trailing about 25 m of line, and the second crew member following at the end of the line. This technique involves little lateral movement off the transect centerline, where attention is focused. Use of two observers allows estimation of the proportion of tortoises detected on the line; and thereby provides a test of the assumption that all tortoises on the transect centerline are recorded ($g(0) = 1$). The capture probability (p) for tortoises within increasing distances from the transect centerline was estimated as for a two-pass removal or double-observer estimator (White et al., 1982): $p = (\text{lead} - \text{follow}) / \text{lead}$, where lead = the number of tortoises first seen by the observer in the leading position and follow = the number of tortoises seen by the observer in the follower position. The corresponding proportion detected near the line by two observers was estimated by $g = 1 - q^2$, where $q = 1 - p$. Figure 4 graphs the relationship between the single-observer detection rate (p) and the corresponding dual-observer detection rate ($g(0)$; “*gee at zero*”). The actual proportion detected can be estimated, but to avoid the necessity of compensating for imperfect detection, during training field crews (pairs) are expected to detect 96% of all models within 1 m of the transect centerline. This corresponds to the leader being responsible for at least 80% of the team’s detections near on the centerline in order to meet this standard and is the basis for one of the training metrics.

Few or no tortoises are located exactly on the line, and even examining a small interval (such as 1 m on each side of the transect line) results in few observations to precisely estimate $g(0)$. Instead, my test of the assumption involves examination of the lead and follow proportions starting with counts of tortoises in larger intervals from the line, moving to smaller intervals centered on the transect centerline. As the intervals get smaller the sample sizes also get smaller, but the estimates are more relevant to the area right at the transect centerline. The expectation is that the estimates should converge on $g(0) = 1.0$.

If the test does not indicate that all tortoises were seen on the transect centerline, the variance of p can be estimated as the binomial variance = $q(1 + q)/np$ (White et al., 1982), where n = the estimated number of tortoises within 1 m of the transect centerline, and the variance of $g(0)$ is estimated as twice the variance of p .




Figure 3. Relationship between single-observer detections (by the leader, p) and dual-observer (team) detections, $g(0)$.

Estimates of tortoise density

Each year, the density of tortoises is estimated at the level of the stratum. The calculation of these densities starts with estimates of the density of tortoises in each stratum from Program DISTANCE, as well as their variance estimates:

$$D = \frac{n}{2wLP_a G_0 g(0)},$$

where L is the total length of kilometers walked in each stratum and w is the distance to which observations are truncated, so $2wL$ is the area searched in each stratum. This is a known quantity (not estimated). P_a is the proportion of desert tortoises detected within w meters of the transect centerline and was estimated using distance assumptions in Program DISTANCE. The encounter rate (n/L) and its variance were estimated in Program DISTANCE for each stratum. Calculation of D required estimation of n/L , P_a , G_0 , and $g(0)$, so the variance of D depended on the variance of these quantities as well.

Proportion of available tortoises was estimated for all strata near each G_0 site and the proportion of available tortoises detected on the transect centerline ($g(0)$) was estimated jointly for all strata. The detection function, which comes into the above equation as P_a , may be estimated jointly or separately for each team, depending on the number and quality of observations. In 2019, the two

teams had very different detection patterns, so separate curves were developed for each. A schematic of the process leading to density estimates is given in Figure 4. Each of the four left-hand columns represent one estimate that contributed to the final density estimates, and the rows in each column show the subsets of the data on which they are based. These estimates combined from left to right to generate stratum and recovery unit density estimates.

	Tortoise encounter rate	Proportion that are visible, G_0	Detection rate, P_a	Proportion seen on the line, $g(0)$	Density	
<i>Recovery unit</i>	<i>Stratum</i>	<i>Neighboring G_0 sites</i>	<i>Data collection group</i>	<i>Overall</i>	<i>Stratum</i>	
West Mojave	FK	SC	Kiva	All data	FK	
	SC				SC	
	OR	OR				
Colorado Desert	AG	CK			All data	AG
	CK					CK
	JT					JT
	PT					PT
	FE	FE				
Eastern Mojave	IV	IV			All data	IV
	EV	PV				EV
Northeastern Mojave	CS	HW	GBI	All data	CS	
	BD				BD	

Figure 4. Process for developing density estimates in 2019. For estimate (one per column), the full set of data was factored as indicated by divisions within the columns.

RESULTS

Field observer training

Training in 2019 lasted from 2 March – 1 April (Table 1). Tests of field detection abilities occurred toward the end of each period, as indicated in the schedules.

Proportion of tortoises detected at distances from the transect centerline

Table 2 reports the proportion of models that were available and were detected over 16 km of transects by each team at 1-, 2-, and 5-m from the transect centerline. Teams were tested after a trial run on the detection lines or after returning crews walked practice transects to refresh the search pattern. The target for detection on the centerline is 100%, and nine of the 22 crews achieved this.

Table 2. Proportion of tortoise models detected in 2019 by crews within 1-, 2-, or 5-m of the transect centerline. Values that scored below the target of 0.90 at 1- and 2-m are highlighted. Crews 1-12 surveyed for Kiva Biological; the remaining crews surveyed for Great Basin Institute.

Crew Number	1m	2m	5m
1	1.00	0.96	0.94
2	1.00	1.00	0.94
3	1.00	1.00	0.94
4	0.93	0.96	0.93
5	0.86	0.92	0.92
6	0.92	0.96	0.92
7	0.93	0.96	0.90
8	0.86	0.89	0.88
9	1.00	0.96	0.90
10	0.93	0.89	0.88
11	0.93	0.92	0.92
12	1.00	0.96	0.93
13	1.00	1.00	0.88
14	1.00	0.96	0.87
15	0.93	0.96	0.91
16	0.93	0.96	0.87
17	0.87	0.92	0.91
18	1.00	0.96	0.90
19	0.81	0.84	0.80
20	0.87	0.92	0.90
21	1.00	1.00	0.90
22	0.87	0.92	0.91
Kiva	0.948	0.949	0.917
GBI	0.926	0.947	0.885
Overall	0.938	0.948	0.902

Table 3 gives the average [absolute] difference between the expected and measured perpendicular distances from the model to the walked line. All measurements for all models during the 2-day trial were used for this estimate and capture two different sources of inaccuracies: 1) using a compass and measuring tape to record distances to the models, plus 2) inaccurately following the trajectory of the transect. The latter source of error does not occur on monitoring transects, because the walked transect is the true transect. On training lines, measurement error increased if crew path diverged from the measured line used to place the models. The “Available Models Detected by Leader” column reports the proportion of all models

that were found first by the leader. During training, this number was used to identify crews in which the leader was not finding at least 80% of all detected. With an 80% detection rate for the leader, a 96% detection rate was expected for the team.

Table 3. Diagnostics for individual crews after training in 2019.

Team	Proportion available models within 2m of centerline by leader	Proportion available models within 2m of centerline by team	Measured versus exact distance (m)	Estimated abundance	95% CI Lower limit	95% CI Upper limit
1	0.85	0.96	0.70	381	321.0	452.2
2	0.93	1.00	0.83	418	373.9	466.8
3	0.96	1.00	0.72	410	298.3	563.6
4	0.88	0.96	0.94	424	367.1	489.2
5	0.88	0.92	0.88	427	393.6	462.6
6	0.96	0.96	0.78	364	319.8	414.1
7	0.88	0.96	0.80	436	321.9	590.2
8	0.82	0.89	0.77	383	306.2	479.8
9	0.89	0.96	0.74	416	347.4	498.6
10	0.89	0.89	0.85	457	375.0	557.9
11	0.92	0.92	0.71	399	348.6	456.5
12	0.84	0.96	0.98	400	288.8	553.9
13	0.89	1.00	0.86	374	308.7	453.1
14	0.92	0.96	1.02	411	305.2	554.3
15	0.92	0.96	0.69	487	391.6	606.6
16	0.85	0.96	1.00	348	298.4	405.9
17	0.73	0.92	0.82	414	329.3	521.6
18	0.96	0.96	0.79	432	367.0	509.1
19	0.80	0.84	0.70	339	261.9	437.8
20	0.88	0.92	0.79	352	312.6	395.3
21	1.00	1.00	0.92	360	311.9	415.4
22	0.81	0.92	0.78	419	313.5	560.4
Kiva	0.892	0.949	0.808	424	376.6	476.2
GBI	0.879	0.947	0.896	398	360.9	438.5
Overall	0.886	0.948	0.821	402	330.1	492.9

Although some individual metrics were below-target (gray cells in Tables 2 and 3), all teams performed well overall so after corrective instruction to fine tune search techniques of specific crews, no pairs were rebuilt. During training, detection curves were fit to each crew’s set of tortoise model observations. In no case was the best-fitting model one without a “shoulder” describing detections near the centerline. The best-fitting detection curves for each team are

plotted in. Figure 5 and 6 and were used to generate density estimates in Table 3. Crews were not evaluated on their ability to match curves of teammates; however, such overlays were used to focus field personnel on an additional level of conformity they could work toward. Distance sampling and development of a single detection curve from many observers is robust to the effects of pooling across observations from crews with variable search patterns, when observers contribute proportionally to the overall pattern (Marques et al., 2007).

In 2019, 10 of the 24 Kiva surveyors were returnees to the project. Five of the 20 GBI surveyors had previous experience with this project, and a sixth one had surveyed for tortoises without using distance sampling. The statistics for the relatively inexperienced GBI team were comparable to those for the experienced Kiva surveyors by the end of training.




Figure 5. Detection curves for each of the 2019 Kiva crews during training. Each curve is based on a 16 km trial with approximately 100 detections.




Figure 6. Detection curves for each of the 2019 GBI crews during training. Each curve is based on a 16 km trial with approximately 100 detections.

Quality assurance and quality control

There were 21,789 transect records and 2232 G_0 records associated with the monitoring effort in 2019. The first data specialist worked with the field teams to resolve 701 cases with fields that were inconsistent with constraints and expectations. After this phase of QA/QC had finished verifying and validating the information in these databases, Phase II provided independent review, repackaged tables into their final configuration, and added some spatial information. An additional 193 issues remained or were discovered in the third (final) phase of QA/QC. Only 117 were errors created by the field crews (sometimes faulty equipment or crews otherwise entering electronic data after the transect was completed, other times data entry error), of which all but 34 were corrected with recourse to paper datasheets. The remaining errors in 2019 indicated a failure to comply with protocols (e.g., first timestamps indicating the transect record was initiated the night before the survey), not because the data were erroneous.

Data for these and previous years can be requested from the author at Linda_Allison@fws.gov.

Transect completion

Table 4 reports the number of assigned and completed transects in each stratum in 2019. Table 4 also indicates the number of assigned transects that could be completed as standard square 12-km transects or by reflecting around property boundaries and infrastructure (column 4). An additional number (column 5) were shortened and represent more rugged terrain. Finally, some transects were considered unwalkable (column 6). Figures 7 to 10 show locations of transects and observations of live tortoises.

Table 4. Number and completion of transects in each stratum in 2019.

Stratum	Assigned transects	Assigned and alternate transects completed	Assigned, completed 12k	Assigned, completed shortened	Assigned, judged unwalkable*
BD	70	70	38	29	3
CS	72	72	35	29	8
EV	80	80	45	21	14
GBI	222	222	118	79	25
AG	35	35	18	14	3
CK	100	100	41	38	21
FE	40	40	37	3	0
FK	59	59	53	4	2
IV	74	74	62	12	0
JT	60	60	14	22	24
OR	56	56	28	14	14
PT	57	57	9	31	17
SC	90	90	55	19	16
Kiva	571	571	317	157	97
Total	793	793	435	236	122

*Assigned transects that were not walked were to be replaced by alternates. In addition to transects that were unwalkable due to terrain and counted in the far right column above, 7 were replaced due to time constraints of accessing through wilderness, and 1 was replaced after consulting with EOD crews on Chocolate Mountain AGR. Four walkable assigned transects in CS could not be accessed through a locked gate, 11 were not completed because China Lake NAWS could not provide escort, and 3 were replaced for logistical considerations pairing teams or complying with military installation requirements. Three were replaced inadvertently.




Figure 7. Distribution of distance sampling transects and live tortoise observations in 2019 in Chocolate Mountain Aerial Gunnery Range, Joshua Tree, Pinto Mountains, and Chuckwalla in the southern part of the Colorado Desert Recovery Unit.




Figure 8. Distribution of distance sampling transects and live tortoise observations in 2019 in the Ivanpah Valley and Eldorado Valley strata of the Eastern Mojave Recovery Unit and in the Fenner stratum of the Colorado Desert Recovery Unit.




Figure 9. Distribution of transects and live tortoise observations in 2019 in the Ord-Rodman, Fremont-Kramer, and Superior-Cronese strata of the Western Mojave Recovery Unit.

Proportion of tortoises available for detection by line distance sampling, G_0

In general, telemetry sites and associated strata were completed sequentially, from south to north. This pattern corresponds to the expected timing of tortoise activity; peaking first in the south, later in the north. Unlike previous years, visibility was very high in all areas for extended periods of the spring in 2019 (Table 5). This is undoubtedly related to the widespread and long-lasting flowering period this year. Temperatures were relatively cool which resulted in extended blooming periods and an extended period before tortoises would experience heat stress. Only in the easternmost sites (Halfway Wash and Piute-Mid) were G_0 estimates below 0.94, yet the estimated visibility was still higher than average for these sites.

Table 5. Availability of tortoises (G_0) when transects were walked in 2019 in the same or in neighboring strata.

G_0 site	Stratum	Dates	Days	G_0 (Std Error)
Chuckwalla	Chocolate Mtn south	10-Mar - 13-Mar	4	0.94 (0.265)
Chuckwalla	Chocolate Mtn north	15-Mar - 17-Mar	3	0.97 (0.182)
Chuckwalla	Chuckwalla	09-Mar - 19-Mar	11	0.96 (0.209)
Chuckwalla	Joshua Tree	22-Mar - 27-Mar	6	1.00 (0.000)
Chuckwalla	Pinto Mountains	27-Mar - 31-Mar	5	0.97 (0.188)
Ord-Rodman	Ord-Rodman	12-Apr - 15-Apr	4	1.00 (0.000)
Superior-Cronese	Fremont-Kramer	07-Apr - 10-Apr	4	1.00 (0.000)
Superior-Cronese	Superior-Cronese	18-Apr - 27-Apr	10	1.00 (0.000)
Ivanpah	Fenner	03-Apr - 06-Apr	4	0.97 (0.176)
Ivanpah	Ivanpah	28-Apr - 03-May	6	0.97 (0.190)
Piute-Mid	Eldorado Valley	02-Apr - 16-Apr	15	0.83 (0.394)
Halfway Wash	Coyote Springs Valley	17-Apr - 30-Apr	14	0.75 (0.459)
Halfway Wash	Beaver Dam Slope	30-Apr - 15-May	16	0.82 (0.405)

Tortoise encounter rates and detection functions

All survey pairs worked together from the beginning to the end of the season. Each Kiva crew surveyed a median 48 transects (no fewer than 45) and overall they detected 321 tortoises larger than 180 mm MCL (“adults”). GBI surveyors walked a median 23 transects each excluding one team that ended surveys after 12 transects due to injury. GBI teams reported 79 adult tortoises. Because GBI did not have a large number of observations on which to base their detection curve, a single detection curve was tested against separate curves for each group, but at several truncation distances, separate curves for each team were most strongly supported. Kiva’s detection pattern best fit a uniform curve with second-order cosine adjustment and using all observations up to 24 m from the centerline. GBI best fit a hazard rate curve and also using observations as far as 20 m from the centerline. Figure 11 and 12 are histograms of the observed number of tortoises seen at increasing distance from the transect centerline. Truncation distance for Kiva removed 6% of the most distant observations resulted in good fit overall and near the

centerline. All but one stratum surveyed by Kiva had at least 25 observations (n=16 for Pinto Mountains). Truncation distance for GBI removed only 5 of the observations and also had a simple shape (no adjustments). All three strata surveyed by GBI retained at least 20 observations within the 20 m truncation distance. The detection rate for Kiva crews within 24 m of the transect centerline was 40.5% (Kiva; CV=0.041) and for GBI crews it was 38.1% (CV=0.149).




Figure 11. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with MCL \geq 180mm found by Kiva in 2019.

This curve uses only the n=302 observations found within 24 m of the line.




Figure 12. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180\text{mm}$ found by GBI in 2019.

This curve uses only the $n=74$ observations found within 20 m of the line.

Proportion of available tortoises detected on the transect centerline, $g(0)$

Because they are cryptic, even tortoises that are visible (not covered by dense vegetation or out of sight in a burrow) and close to the surveyor may not be detected. In 2019, for 112 detections of tortoises within 5 m of the transect centerline, 94 were found by the observer in the lead position and 18 by the follower, so that the probability of detection by single observer, $p = 0.808$, and the proportion detected using the dual observer method, $g(0 \text{ to } 5 \text{ m}) = 0.963$ (SE = 0.071). Figure 13 shows that $g(0)$ was converging on 1.0 in 2019 although this pattern weakens within a meter of the line, where estimates are based on a diminishing number of observations. The curves since dual observers were first used in 2004 have all supported the premise that complete detection on the transect line was achieved for years in which the dual-observer method was used (USFWS 2009, 2012b, 2012c, 2013, 2014, 2015, 2016, 2018, 2019). Previous years of data and the pattern in Fig. 13 indicate the assumption of perfect detection on the centerline was met; consequently, no adjustment was made to the final density estimate.




Figure 13. Detection pattern for the leader (p) and by the team ($g(\theta)$) based on all observations out to a given distance (x) from the centerline in 2019. Note convergence of $g(\theta)$ on 1.0 as x goes to 0.

Estimates of tortoise density

Density estimates were generated separately for each monitoring stratum, and for OR they were generated based only on resident tortoises and then again after including marked animals that had been translocated (Table 6). The reported densities for OR represent an increase of 366 (SE=77.3) tortoises, an estimate which may not stabilize until a couple years after the last translocations in spring 2019 when the movements of these animals should settle into more typical home ranges (Nussear et al. 2012, Farnsworth et al. 2015).

Table 6. Stratum-level encounters and densities in 2019 for tortoises of MCL \geq 180 mm.
Coefficients of variation expressed as percentages.

Recovery Unit/ Stratum		Area (km ²)	# Transects	Transect length (km)	Begin date	End date	<i>n</i> (torts observed)	CV(<i>n</i>)	Density (/km ²)	CV(Density)
Western Mojave		6873	205	2308	16-Mar	26-Apr	103			
Fremont-Kramer	FK	2417	59	692	9-Apr	9-Apr	36	23.7	2.7	24.00
Ord-Rodman	OR	1124	56	601	16-Mar	16-Mar	29	19.9	2.5	20.33
OR – residents only	ORr	1124	75	601	16-Mar	16-Mar	24	21.6	2.1	21.96
Superior-Cronese	SC	3332	90	1016	26-Apr	26-Apr	38	23.3	1.9	23.65
Colorado Desert		8422	292	2946	9-Mar	20-Apr	157			
Chocolate Mtn north	AGN	351	14	126	12-Mar	14-Mar	34	15.7	14.2	24.73
Chocolate Mtn south	AGS	403	21	237	9-Mar	11-Mar	13	32.4	3.0	43.04
Chocolate Mtn	AG	755	35	363	9-Mar	14-Mar	47	19.6	7.0	29.51
Chuckwalla	CK	3509	100	1032	22-Mar	7-Apr	34	18.5	1.8	28.83
Fenner	FE	1841	40	473	9-Apr	16-Apr	25	22.7	2.8	29.27
Joshua Tree	JT	1567	60	578	16-Mar	20-Apr	35	19.7	3.1	20.15
Pinto Mountains	PT	751	57	500	16-Mar	20-Apr	16	24.9	1.7	31.81
Eastern Mojave		3720	154	1755	16-Apr	30-Apr	68			
Eldorado Valley	EV	1153	80	890	17-Apr	26-Apr	26	21.6	2.3	54.35
Ivanpah Valley	IV	2567	74	866	16-Apr	30-Apr	42	14.8	2.6	24.92
Northeastern Mojave		1853	142	1549	3-Apr	9-May	48			
Beaver Dam Slope	BD	828	70	780	26-Apr	9-May	20	25.7	2.0	57.54
Coyote Springs	CS	1025	72	769	3-Apr	17-Apr	28	20.7	3.2	65.99

DISCUSSION

Surveyors in 2019 had a spectacular experience – there had been heavy winter precipitation throughout the desert so that the spring flowering period was dramatic and widespread. Also, temperatures were cooler than average through the spring, probably prolonging the blooming period for most annual plants. This long period of food availability and moderate temperatures also probably explains the large number of tortoises seen on transects. However, this did not cause our density estimates to increase dramatically because at the same time, a larger proportion of tortoises was aboveground at telemetry sites, so the correction factor for tortoises that were below ground was minimal compared to other years.

Base expansion of 29 Palms MCAGCC affected many tortoises, and starting in 2017, 727 adult tortoises were translocated to two areas either inside (n=426) or at the boundary of the Ord-Rodman critical habitat unit (n=301) before the surveys reported here. Although this expansion negatively impacts tortoises and their habitat (USFWS 2017b) elsewhere, augmenting the resident population in OR with reproductive adults may result in local population growth by increasing the number of juveniles produced each year. In conjunction with fencing, law enforcement, and other mitigation that is implemented in OR, population augmentation is a strategy that may accelerate the process of stabilizing this population (USFWS 2011). Other monitoring is in place to assess the success of the translocations measured by survivorship, for instance, but the ongoing range-wide monitoring program reported here will provide a composite view of the success of the suite of recovery activities that are now occurring in OR. This year, the program estimated an increase of 366 adult tortoises due to translocations (Table 6). While this estimate is lower than the 727 that were actually translocated in and nearby, we do not have a specific prediction for the number of translocatees that would settle inside or outside the boundaries of the surveyed critical habitat unit. Instead, these surveys will document whether there is an overall increase in the number of tortoises over time. The northwestern part of Ord-Rodman did not receive translocatees. In Figure 9, the relatively high number of dead tortoises found there this year is striking. At least four of the tortoises had died in the last 1-2 years and were found along a barbed-wire fence on the east end of Stoddard Valley.

Although these surveys do not search for tortoises directly on MCAGCC, areas of other installations such as Chocolate Mountains AGR and NAWS China Lake are in critical habitat and are surveyed. In 2019, access protocols for China Lake had changed, necessitating 2 separate visits for each surveyor seeking security clearance. This represents 2 days fewer transects but can be accommodated. However, the base also requires escorts on the transects and did not have sufficient escorts to ensure any would be available if we completed the process for security clearances. We did not send surveyors for the clearance process and transects were not completed on China Lake.

Access to Chocolate Mountain is limited to those times when it is closed to military training, and this period has shifted steadily earlier in the year. Although tortoises may be active earlier than in past decades, the shift in the military training schedule has moved at a faster rate that is not based on tortoise activity and may necessitate renegotiating the surveys; surveying for tortoises too early and before they have emerged from their winter burrows would compromise density estimation. Separate from scheduling needs on the military installations, one priority for the next years will be to determine whether there is a pattern of tortoise activity moving earlier in the season in any parts of the range. This will inform the optimal timing of surveys but of course more importantly would reflect a response to the changing climate in the Mojave and Colorado deserts. To describe annual tortoise activity patterns, we could use the currently transmittered tortoises for observation, but would have to expand the monitoring period beyond the transect surveys to capture changes over the season.

In 2018, annual density estimates were used to describe population trends between 2004 and 2014 in each of the monitoring strata (Allison and McLuckie 2018). These trend estimates will be updated based on more recent information only after there have been at least three new annual density estimates for each monitoring stratum; probably about every six years. The next evaluation of population trends may be possible after the 2020 field season depending on sufficient funding to cover the last strata for a third time.

LITERATURE CITED

- Allison, L.J. and A. McLuckie. 2018. Population trends in Mojave Desert Tortoises (*Gopherus agassizii*). *Herpetological Conservation and Biology* 13(2):433-452.
- Anderson, D.R., and K.P. Burnham. 1996. A monitoring program for the desert tortoise. Report to the Desert Tortoise Management Oversight Group.
- Anderson, D.R., K.P. Burnham, B.C. Lubow, L. Thomas, P.S. Corn, P.A. Medica, and R.W. Marlow. 2001. Field trials of line transect methods applied to estimation of desert tortoise abundance. *Journal of Wildlife Management* 65:583-597.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers, and L. Thomas. 2001. *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford Univ. Press, Oxford. 432 pp.
- Darst C.R., P.J. Murphy, N.W. Strout, S.P. Campbell, K.J. Field, L.Allison, and R.C. Averill-Murray. 2013. A strategy for prioritizing threats and recovery actions for at-risk species. *Environmental Management* 51:786–800.
- Farnsworth, M.L.; B.G. Dickson; L.J. Zachmann; E.E. Hegeman; A.R. Cangelosi; T.G. Jackson, Jr.; and A.F. Scheib. 2015. Short-term space-use patterns of translocated Mojave Desert Tortoise in southern California.
- Kincaid, T.M. and A.R. Olsen. 2017. spsurvey: Spatial Survey Design and Analysis. R package version 3.4. URL: <https://CRAN.R-project.org/package=spsurvey>.
- Lindenmayer, D.B., G.E. Likens, A. Haywood, and L. Miezi. 2010. Adaptive monitoring in the real world: proof of concept. *Trends in Ecology and Evolution* 26:641–646.
- Lyons, J.E., M.C. Runge, H.P. Laskowski, and W.L.Kendall. 2008. Monitoring in the context of structured decision-making and adaptive management. *Journal of Wildlife Management* 72:1683–1692.
- Marques, T.A., L. Thomas, S.G. Fancy, and S. T. Buckland. 2007. Improving estimates of bird density using multiple-covariate distance sampling. *The Auk* 124(4) 1229-1243.
- Murphy, R.W., K.H. Berry, T. Edwards, A.E. Leviton, A. Lathrop, J.D. Riedle, 2011. The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation. *ZooKeys* 113: 33-71. doi: 10.3897/zookeys.113.1353.

- Nichols, J.D., and B.K. Williams. 2006. Monitoring for conservation. *Trends in Ecology and Evolution* 21:668–673.
- Nussear, K.E., T.C. Esque, R.D. Inman, L. Gass, K.A. Thomas, C.S.A. Wallace, J.B. Blainey, D.M. Miller, and R.H. Webb. 2009. Modeling habitat of the desert tortoise (*Gopherus agassizii*) in the Mojave and parts of the Sonoran deserts of California, Nevada, Utah, and Arizona. U.S. Geological Survey Open-file Report 2009-1102.
- Nussear, K.E., C.R. Tracy, P.A. Medica, D.S. Wilson, R.W. Marlow, and P.S. Corn. 2012. Translocation as a conservation tool for Agassiz's Desert Tortoises: Survivorship, reproduction, and movements.
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Stevens, D.L., Jr. and A.R. Olsen. 2004. Spatially-balanced sampling of natural resources. *Journal of American Statistical Association* 99(465): 262-278.
- Thomas, L, S.T. Buckland, E.A. Rexstad, J.L. Laake, S. Strindberg, S.L. Hedley, J.R.B. Bishop, T.A. Marques, and K.P. Burnham. 2010. Distance software: design and analysis of distance sampling surveys for estimating population size. *J. of Applied Ecology* 47:5-14.
- [USFWS] U.S. Fish and Wildlife Service. 2006. Range-wide Monitoring of the Mojave Population of the Desert Tortoise: 2001-2005 Summary Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- [USFWS] U.S. Fish and Wildlife Service. 2009. Range-wide Monitoring of the Mojave Population of the Desert Tortoise: 2007 Annual Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- [USFWS] U.S. Fish and Wildlife Service. 2011. Revised recovery plan for the Mojave Population of the desert tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California. 222 pp.
- [USFWS] U.S. Fish and Wildlife Service. 2012a. Health Assessment Procedures for the Mojave Desert Tortoise (*Gopherus agassizii*): A Handbook Pertinent to Translocation. Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. Accessible through: http://www.fws.gov/nevada/desert_tortoise/index.html
- [USFWS] U.S. Fish and Wildlife Service. 2012b. Range-wide Monitoring of the Mojave Desert Tortoise: 2008 and 2009 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2012c. Range-wide Monitoring of the Mojave Desert Tortoise: 2010 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2013. Range-wide Monitoring of the Mojave Desert Tortoise: 2011 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2014. Range-wide Monitoring of the Mojave Desert Tortoise: 2012 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2015. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2013 and 2014 Annual Reports. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2016. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2015 and 2016 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2017a. Desert Tortoise Monitoring Handbook. Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. Version: 28 February 2017. http://www.fws.gov/nevada/desert_tortoise/reports.

[USFWS] U.S. Fish and Wildlife Service. 2017b. Biological Opinion for Land Acquisition and Airspace Establishment, Twentynine Palms, California (8-8-11-F-65R). Letter to Lt. Col T.B. Pochop, Marine Corps Air Ground Combat Center, Marine Air Ground Task Force Training Command, Natural Resources and Environmental Affairs Division, Twentynine Palms, California. Dated January 31. From G. Mendel Stewart, Field Supervisor, Carlsbad Fish and Wildlife Office, Carlsbad, California.

[USFWS] U.S. Fish and Wildlife Service. 2018. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2017 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2019. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2018 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

White, G.C., D.R. Anderson, K.P. Burnham, and D.L. Otis. 1982. Capture-recapture and removal methods for sampling closed populations. LA-87-87-NERP. Los Alamos National Laboratory, Los Alamos, NM. 235pp.