


# RELATIONSHIPS BETWEEN PALUSTRINE WETLANDS OF FORESTED RIPARIAN FLOODPLAINS AND FISHERY RESOURCES: A REVIEW



Fish and Wildlife Service

U.S. Department of the Interior

Biological Report 88(32)  
August 1988

RELATIONSHIPS BETWEEN PALUSTRINE WETLANDS OF FORESTED  
RIPARIAN FLOODPLAINS AND FISHERY RESOURCES: A REVIEW

by

Johnie H. Crance  
U.S. Fish and Wildlife Service  
National Ecology Research Center  
Creekside One Building  
2627 Redwing Road  
Fort Collins, CO 80526-2899

U.S. Department of the Interior  
Fish and Wildlife Service  
Research and Development  
Washington, DC 20240

Suggested citation:

Crance, J.H. 1988. Relationships between palustrine wetlands of forested riparian floodplains and fishery resources: a review. U.S. Fish Wildl. Serv. Biol. Rep. 88(32). 27 pp.

## PREFACE

This report stemmed from a workshop held at Annapolis, Maryland, February 9-10, 1987, to identify priority information needs for nontidal, freshwater palustrine and riverine wetlands (exclusive of isolated wetlands). Participants in the workshop were primarily representatives of the U.S. Fish and Wildlife Service's Ecological Services Field Offices located at Annapolis, Maryland, and State College, Pennsylvania. A high-priority information need identified by the workshop participants was documentation of the values to fish of freshwater palustrine wetlands that are generally tree- and shrub-dominated floodplains of streams that flow to coastal areas of the eastern portion of the United States. The workshop participants concluded that a synthesis of the relevant literature should be the first step in addressing this information need. The participants also recognized that long-term field research will be required to determine more clearly the relationships between these selected palustrine wetlands and associated fisheries resources. Soon after a review of the relevant literature was initiated it became evident that there is sufficient evidence to support the hypothesis that palustrine wetlands of forested riparian floodplains are important to fishery resources. It also became evident that much additional research on this subject is needed for use in management and protection of these vital wetland resources and their associated fishery resources.

Wetlands are of national interest because of the many benefits they provide for society. Palustrine wetlands of forested riparian floodplains are notable because of the large areas they cover and because of the rapid rate at which they are being converted for timber production, agriculture, and human settlement. These wetlands are inextricably linked, directly or indirectly, to streams, estuaries, and upland watersheds. This report focuses on linkages between these wetlands, which flank many of the major streams that flow through the Piedmont and Coastal Plain to the sea, and their relationship to fishery resources. The draft manuscript was reviewed by six wetland ecologists for technical content and soundness of concepts. Some major changes in format and content of the draft manuscript along with other comments and recommendations by these six reviewers were incorporated where appropriate in the final manuscript.

Suggestions or comments relating to this report should be sent to the author at the noted address.

## CONTENTS

	<u>Page</u>
PREFACE .....	iii
ACKNOWLEDGMENTS .....	vi
INTRODUCTION .....	1
Focus, Scope, and Purpose of This Report .....	1
LITERATURE REVIEW .....	3
Use of Floodplains by Fish .....	3
Structure, Function, and Hydroperiod of Floodplains .....	7
SUMMARY AND CONCLUSION .....	16
LITERATURE CITED .....	18

## ACKNOWLEDGMENTS

Review of the draft manuscript and helpful suggestions by the following individuals are gratefully acknowledged: M.M. Brinson, D.L. Correll, J.W. Day, Jr., K.W. Cummins, G. Kinser, V.W. Lambou, L.C. Lee, and M.L. Scott.

Sincere thanks are extended to C.F. Bryant, L. Ischinger, and C.B. Stalnaker for reviewing the final manuscript and providing helpful suggestions.

The usual excellent editorial service provided by J.R. Zuboy is acknowledged and appreciated. I extend special thanks to Jennifer Shoemaker for designing the cover, and Ann Crance and Jennifer Shoemaker for preparing drawings, Patricia Gillis and Dora Ibarra for providing word processing support, and Kay Lindgren for locating the many articles that I reviewed.

## INTRODUCTION

At the functional level, wetlands moderate the effects of flooding, maintain and improve water quality, provide fish and wildlife habitat, support food chains, and have aesthetic and heritage values (Table 1). Wetlands also contribute to the stability of levels of atmospheric nitrogen, sulfur, carbon dioxide, and methane (Mitsch and Gosselink 1986).

Numerous laws and regulations have been enacted to protect and manage wetlands (Zinn and Copeland 1982; Kusler 1983; Mitsch and Gosselink 1986), but the area of wetlands in the lower conterminous United States declined from about 215 million acres in presettlement times (Roe and Ayres 1954) to about 99 million acres in the mid-1970's (Frayer et al. 1983). During the past 40 years, there have been significant losses of palustrine wetlands of forested riparian floodplains in the southeastern portion of the United States. A net loss of 4.5 million acres of palustrine forested wetlands occurred in the Mississippi Flyway alone during the 1950's-1970's (Frayer et al. 1983).

Until the second half of the twentieth century, the only functional value widely attributed to wetlands was that of providing habitat for wildlife, particularly waterfowl. More recently, wetlands have attracted significant attention related to other functions and values (Good et al. 1978; Greenson et al. 1979; Brinson et al. 1981a,b; Richardson 1981; Sather and Smith 1984; Strickland 1986; Mitch and Gosselink 1986).

At the population level, wetland-dependent fish, shellfish, furbearers, and waterfowl provide a valuable harvest and millions of man-days of recreational fishing and hunting. Commercial landings of wetland-dependent fish species had a landing value exceeding \$700 million in 1976; and in 1975, recreational fishermen spent \$31.1 billion pursuing wetland-associated fishes (Peters et al. 1979). About one-third of the North American bird species use wetlands (Kroodsman 1979), as do numerous aquatic invertebrates, amphibians, reptiles, fish, and mammals (Lambou 1965, 1984; Ziser 1978; Clark 1979; Fredrickson 1979; Schitoskey and Linder 1979; Brinson et al. 1981a,b; Frizell 1988). About 50% of the threatened or endangered animal species and 28% of the plant species are associated with wetlands, including 22 of the 41 species and subspecies of United States fishes listed as endangered or threatened (Williams and Dodd 1979; Niering 1988). The timber value of southern wetland forests was estimated to be \$8 billion (Johnson 1979).

### FOCUS, SCOPE, AND PURPOSE OF THIS REPORT

There is no universal, correct, indisputable, ecologically sound definition for wetlands, primarily because of the diversity of wetlands and

Table 1. Summary of representative examples of functions of forested palustrine wetlands (adapted from Roelle et al. 1987).

Major functional categories	Specific functions
Hydrology	Flood storage Velocity reduction Groundwater discharge modification
Water quality	Sediment retention Erosion control Nutrient retention and transformation Contaminant retention and transformation
Fisheries and wildlife	Habitat for finfish Habitat for shellfish Habitat for wood ducks Habitat for wintering dabbling ducks Habitat for black bear Habitat for white-tailed deer Habitat for migrating passerine birds
Ecosystem processes	Maintenance of natural biotic diversity, cumulative level Maintenance of natural biotic diversity, site-specific level Food chain support Streamflow mediation Transforming and filtering (water quality)
Cultural/recreational/economic	Sensory experience Recreation experience Information storage Renewable harvesting

because demarcation between wet and dry environments lies along a continuum. However, several wetland classification systems impose boundaries on wetlands for the purposes of inventory, evaluation, and management (e.g., Martin et al. 1953; Shaw and Fredine 1956; Stewart and Kantrud 1971; Golet and Larson 1974; Zoltai et al. 1975; Millar 1976; Cowardin et al. 1979). Definitions of wetlands often depend on the objectives and field of interest of the user. I have chosen palustrine wetlands of forested riparian floodplains for the focus of this report. These wetlands are primarily forested and scrub/shrub classes

of the palustrine wetlands system as classified by Cowardin et al. (1979). As used in this report, palustrine wetlands of forested riparian floodplains include only wetlands that are directly coupled to upland watersheds and to stream systems (generally higher order streams) which flow to estuaries along the mid- and south-Atlantic and the northern Gulf of Mexico coastal areas of the United States. Upslope, these wetlands are bounded by terrestrial habitat. They are coupled to riverine habitat via a "water bridge" during flooding (Figure 1). The water bridge is essential from a fishery functional value standpoint because during flooding it allows fish access to productive floodplain areas used for spawning, nursery, feeding, and cover. The water bridge also provides passage for fish to return to the stream channel during receding floodwaters. A stream corridor with palustrine forested wetlands located in its floodplain may also contain wetlands classified as Estuarine Intertidal Forested Wetlands, Riverine Lower Perennial Emergent Wetlands and Aquatic Beds, and Palustrine Aquatic Beds (Cowardin et al. 1979). These wetland habitats are also functionally important to fish that use the stream corridor and associated wetlands. However, such wetlands are not the focus of this report. Terms other authors have used to describe palustrine wetlands of special interest and which I consider to be generally synonymous to wetlands that are the focus of this report are listed in Table 2.

A recent survey by Abernethy and Turner (1987) showed that 57% of the forested wetland area in the United States is located in Alabama, Arkansas, Georgia, Florida, Kentucky, Louisiana, Mississippi, Missouri, North Carolina, South Carolina, Tennessee, and Virginia. In this area of the United States, palustrine wetlands of forested riparian floodplains are commonly referred to as bottomland hardwoods. However, this term is used very loosely from an anatomical standpoint, since both hardwood and softwood tree species occur on floodplains of streams in these States.

## LITERATURE REVIEW

### USE OF FLOODPLAINS BY FISH

It is generally known that palustrine wetlands of forested riparian floodplains provide cover, spawning, and nursery habitat in situ for numerous fish species (e.g., Hinchey 1977; Hastings 1979; Welcomme 1979; Wharton and Brinson 1979; Chambers 1980; Wharton et al. 1981). It is generally known or assumed that such wetlands usually import, produce, store, recycle, and export biotic and abiotic materials that are used in food chains by fish in situ or at sites downstream (e.g., Day et al. 1977, 1980; Livingston and Loucks 1979; Conner and Day 1982; Taylor et al. 1984).

Most of the available information on the occurrence, population, and production of fish in forested floodplains has been reviewed and summarized by Wharton et al. (1981, 1982). It has been shown that 90 fish species use the wooded floodplains of the Atchafalaya River, Louisiana (Lambou 1965, 1984; Bryan et al. 1975, 1976). Of the 90 species, 51 used the overflow wooded areas for spawning or for rearing of young, while 53 species used these areas




Figure 1. Simplified schema of a cross section of a stream corridor showing the proximity of a palustrine wetland in a forested riparian floodplain to upland, riverine, and some other palustrine habitats (adapted from Cowardin et al. 1979).

for feeding (Table 3). Bottomland hardwood areas of the Henderson Lake area in the Atchafalaya floodplain are used by at least 37 fish species for spawning and nursery habitat (Pollard et al. 1983; Lambou 1984). The number of fish species that have been collected from other floodplains areas include: 13 in a cypress swamp in Florida (Carlson and Duever 1977), 16 from a blackwater creek floodplain in North Carolina (Walker 1980), and 20 from floodplain areas along the Savannah River in Georgia (Patrick et al. 1967). Numerous fish species use the floodplain along the Suwanee River in Georgia for spawning and nursery habitat (Wyatt and Holder 1969; Holder 1970; Holder et al. 1970, 1971; Germann 1973). Forested floodplains are used as spawning sites by hickory shad (*Alosa mediocris*) and blueback herring (*A. sapidissima*) in New Jersey

Table 2. Some terms used by various authors in discussing Palustrine Forested Wetlands and Scrub/Shrub Wetlands (Cowardin et al. 1979) and that are considered to be generally synonymous to palustrine wetlands of forested riparian floodplains.

Term	Author
Floodplain Floodplains of rivers	Welcomme (1979, 1985)
Forested wetlands Swamp forests	Conner and Day (1982)
Wetlands of bottomland hardwood forests	Clark and Benforado (1981), Roelle et al. (1987), Wilkinson et al. (1987), Lea (1988)
Floodplain swamps	Kuenzler et al. (1980)
Riparian ecosystems Riparian wetlands	Ewel (1978), Brinson et al. (1981b), Mitsch and Gosselink (1986)
Bottomland forests	Day et al. (1981), McKnight et al. (1981)
Bottomland hardwood-cypress forests	Langdon et al. (1981)
Lowland hardwood wetlands	Fredrickson (1979a,b)
Pine barrens Cedar swamps	Patrick et al. (1979)

(unpublished report cited by Hastings 1979). The blue crab (*Callinectes sapidus*), shrimp (*Panaeus* sp.), and several species of estuarine finfish were found to use freshwater forested floodplains in Louisiana (Hinchee 1977; Chambers 1980; Lambou 1984). Crayfish form a large percentage of the aquatic animal biomass on some floodplains (Holder 1971; Konikoff 1977) and are an important food source for the large predator fishes found on floodplains (Penn 1950; Holder 1971; Wharton 1977). Several species of freshwater mussels dominate the animal biomass of some floodplains (Parson and Wharton 1978). The glochidian stage of most freshwater mussels cannot survive unless they

Table 3. Summary of fish use of habitat types for a particular purpose in the leveed Atchafalaya Basin between Morgan City and Simmesport, Louisiana (adapted from Lambou 1984).

Habitat type and use	Number of fish species	Percent of total number (n = 90) of species
Used overflow wooded areas for spawning or rearing of young	51	53.7
Used overflow wooded areas for feeding	53	55.8
Spawned in marine waters and adults and subadults used overflow wooded and permanent-water areas for feeding	2	2.1
How extensively overflow wooded areas were used is not known	31	32.6
Used permanent-water areas with current, how extensive overflow wooded areas were used is not known	5	5.3
Mainly inhabited marine or estuarine waters, a few individuals invaded area during low-water periods	4	4.2

become attached to their host-specific species of fish. The average annual yield of finfish and shellfish from floodplains to man is largely dependent on the maximum area flooded (Bryan and Sabins 1979) and the amount of fishing effort. Yields may range from about 4,000 to 6,000 kg/km<sup>2</sup> of finfish (Welcomme 1979) to as high as 8,797 kg/km<sup>2</sup> of finfish and shellfish combined (Lambou 1984). Fish density and biomass in a cypress (*Taxodium distichum*) swamp in Florida ranged from 16 to 43 fish/m<sup>2</sup> and 1 to 3 g/m<sup>2</sup> respectively (Carlson and Duever 1977). The area of bottomland hardwood wetlands in the Mississippi River basin is likely an important determinant of the overall level of fishery productivity in that river basin (Risotto and Turner 1985).

## STRUCTURE, FUNCTION, AND HYDROPERIOD OF FLOODPLAINS

A floodplain may be considered the part of a stream channel that is used to accommodate high flows (Brinson et al. 1981). Floodplains result from the combination of the deposition of alluvial materials (aggradation) and down-cutting of surface material over a period of many years. In a review of the structure and function of riparian floodplains Mitsch and Gosselink (1986) pointed out that the importance of the stream to the floodplain and the floodplain to the stream cannot be overemphasized. If the stream or the floodplain is altered the other will surely change in time because streams and their floodplains are in continual dynamic balance between the building and removal of structure. Generally, there are four hydrological phases of a stream in relation to its floodplain (Figure 2).

The hydroperiod (Figure 3) defines the pattern of rise and fall of the floodplain's surface and subsurface water. The hydroperiod is unique to each stream and each floodplain and is the driving force which determines the floodplain's structure and function. Changes in the frequency, duration, and timing of the hydroperiod may prevent passage of fish between the stream and its floodplain, alter the age structure and species composition of the floodplain flora and fauna, and interfere with ecological processes that support food chains in the wetland and in downstream systems (Bedinger 1981; Kadlec 1987; Klimas 1988). The timing of flooding is particularly important because flooding during the growing season has a greater effect on floodplain productivity than does an equal amount of flooding during the nongrowing season (Mitsch and Gosselink 1986). Floodplain vegetation is dependent on inundation and drainage patterns that differentially influence the ability of various plant species to become established and compete on a given site (Klimas 1988). Timing and magnitude of floods are especially critical to fish species that use the floodplain for spawning. Migration routes may be blocked if flood levels are too low.

### The Role of Hydrology in Floodplains

The role of hydrology in maintenance of floodplain wetlands has been discussed by numerous authors (e.g., Conner and Day 1976; Gosselink and Turner 1978; Bedinger 1979, 1981; Carter et al. 1979; Wharton 1980; Brinson et al. 1981b; Gosselink et al. 1981; Klimas 1988). Gosselink and Turner (1978) developed a conceptual model of the role of hydrology in freshwater wetlands. Their model was adapted (Figure 4) to represent the role of hydrology in palustrine wetlands of forested riparian floodplains discussed in this report. This model shows that some climatic factors may have an overriding influence on the types of species that develop in wetlands of forested riparian floodplains. However, certain attributes of the hydrologic regime (i.e., water source, velocity, renewal rate, and timing) are of most importance to the biota, and especially to fish. The source of water strongly influences its ionic composition, oxygen saturation, and toxin load, each of which may influence the water quality suitability for fish. Water velocity affects turbulence and sediment transport and distribution, each of which may be determining factors for fish spawning and feeding habitat. The renewal rate or frequency of replacement of the water is dependent on volume and frequency of inundation and velocity. The renewal rate and the timing of frequency of




Figure 2. Schematic showing four of the hydrological phases of a stream in relation to its forested riparian floodplain: a, drought phase with the water level on the floodplain too low to support fish; b, flood phase when overbank flow from the stream channel contributes water to the floodplain allowing fish passage; c, post-flood phase with surface drainage from the floodplain to the stream, allowing fish passage; and d, normal channel-contained streamflow which originates from floodplain seepage, precipitation, or lateral runoff from adjacent uplands. (Adapted from M.M. Brinson, H.D. Bradshaw, and R.N. Holmes, "Significance of floodplains in nutrient exchange between a stream and its floodplain," in T.D. Fontaine and S.M. Bartell, *Dynamics of Lotic Ecosystems*, pp. 199-221, with permission from Butterworth Publishers, Stoneham, MA.)




Figure 3. Example of hydrographs from the Apalachicola River, Florida (solid line) at river mile 86, and from two nearby floodplain sites along the river during the water year 1980. Floodplain site 1, (dashed line) was located one mile east of the river in a large tupelo-cypress swamp that was permanently ponded with 1-2 feet of water. Floodplain site 2 (dotted line) was located about 0.5 mile west of the river in a floodplain with high flow velocities during floods, but which was ponded during low flows (adapted from Leitman et al. 1984).

inundation and its regularity may limit the availability and use of the floodplain by fish and wildlife species. Water nutrients, toxins, and oxygen availability are key chemophysical properties of the floodplain's substrate and are strongly influenced by the hydrologic regime. Detailed discussions on the influence of the hydrologic regime on floodplain flora are found in Gosselink and Turner (1978).

Openness or coupling and primary productivity, secondary productivity and food chain support, and water quality characteristics of wetlands are known to respond to hydrology, which in turn determines the fisheries value of palustrine wetlands of forested riparian floodplains. These characteristics, their responses to hydrology, and their relationship to fishery resources are discussed in the following sections of this paper.

## Local Climatic Influences

Solar radiation →  
Temperature →  
Precipitation →  
Relative humidity →  
Cyclic regularity →




Figure 4. A conceptual model of the role of hydrology in palustrine wetlands of forested riparian floodplains. (Modified from Gosselink and Turner (1978). Original copyright 1978 by Academic Press, Inc. Modified and reproduced with permission from the publishers.)

## Floodplain Coupling and Primary Productivity

Portions of the length of a stream corridor, including tributaries, have vital hydrologic and biologic attributes that couple a stream upstream and downstream, and laterally with its floodplain (Wharton and Brinson 1979). Palustrine wetlands of forested riparian floodplains are coupled directly or indirectly to upland watersheds, streams, and estuaries. By virtue of this coupling, gravity, the solubility and transport properties of water, and the mobility of various aquatic organisms, including fish, these wetlands are open systems accounting for exchange of materials among various components of the stream corridor (Figure 5). The functional aspect of this coupling depends on pulses of water flow which are vital to primary and secondary productivity in the floodplain and at downstream sites (Wharton and Brinson 1979).

Studies have shown that primary productivity of forested floodplains depends on hydrologic and nutrient conditions (see summary by Mitsch and Gosselink 1986). Highest productivity results from wetlands that are neither too wet nor too dry but that have seasonal hydrologic pulsing (Brown et al. 1979; Brinson et al. 1981b; Conner and Day 1982) (Figure 6). Floodplain forests are among the highest in primary productivity of any ecosystem in the southeastern United States (Boyd 1976; Conner and Day 1976; Brown et al. 1979). Nutrient supplies from the watershed deposited during stream flooding provide resources necessary to sustain high rates of productivity. Trees are the floodplain's primary producers and the source of much of the detritus on the floodplain and in the associated stream (Wharton and Brinson 1979). Aboveground biomass varies widely, ranging from 10 kg/m<sup>2</sup> to 119 kg/m<sup>2</sup> (Brinson et al. 1981b). Some tree species commonly found on forested floodplains in the Southeast are bald cypress (*Taxodium distichum*), water tupelo (*Nyssa aquatica*), swamp tupelo (*N. sylvatica*), black gum (*N. biflora*), water hickory (*Cara aquatica*), overcup oak (*Quercus lyrata*), red maple (*Acer rubrum*), Atlantic white cedar (*Chamaecyparis thyoides*), and water ash (*Fraxinus caroliniana*). However, tree species composition may vary greatly depending on hydrologic regimes, changes in floodplain elevation, and other factors (Conner and Day 1976; Wharton and Brinson 1979; Brinson et al. 1981; Larson et al. 1981; McKnight et al. 1981; Abernethy and Turner 1987).

One of the fundamental functions of primary productivity in forested floodplains is that of providing energy for detritus-based food webs, including fish food webs, in situ and at downstream sites. Detritus-based food webs add stability to ecosystems by making the energy that is seasonally fixed by primary production available to consumers over longer periods of time.

## Secondary Productivity and Food Chain Support Related to the Floodplain

A review of the secondary production values of wetlands by Crow and Macdonald (1979) suggests that three distinct and significant wetland contributions to secondary productivity stand above all others. Two of these contributions relate to wetland trophic resources and one relates to nontrophic resources. Generally, food energy use in wetlands is of two types: direct and indirect consumption. Direct consumption is concerned with energy and matter transferred by means of grazing by consumers, including insects (Wallace and O'Hop 1985; Scott and Haskins 1987), waterfowl (Chamberlain 1959; Glasgow

## TERRESTRIAL WATERSHED




Figure 5. Schema showing the potential for transfer of materials between a palustrine wetland of a forested riparian floodplain and other systems linked directly or indirectly to the wetland. For each system (i.e., terrestrial watershed, riverine, forested riparian wetland, estuarine, and marine), examples are given of major sources of energy for the system (E), some major primary producers for the system (P), and some major consumers for the system (C). Arrows with broken line indicate direct linkage and direction of transfer of water, detritus, nutrients, and sediments. Arrows with solid line indicate direct linkage and direction of movement of fish and other primary consumers.





phytoplankton → zooplankton → insects  
↓  
→ small fish → larger fish → man

### Water Quality and the Floodplain

The role of forested wetlands in the southeastern United States in maintaining water quality was reviewed by Winger (1986). Good water quality enhances the quality of fish habitat. The quality of water entering or leaving the floodplain and exported downstream may be affected by the nutrient content of surface water or ground water from the watershed. In the past two decades, concerns have increased that the use of agricultural fertilizers and pesticides may result in contamination of surface and ground water and increased nutrient loading to streams (Kemp 1949; Scheuler and Kemp 1979; Kuenzler and Craig 1986). Gilliam et al. (1974) found ground water nitrate concentrations to be higher under cultivated and fertilized fields than from unfertilized woods. A study in four watersheds of the Little River in the Georgia Coastal Plain (Lowrance et al. 1985) showed that two watersheds with more agricultural land had consistently higher loads of N, K, Ca, Mg, and Cl in streamflow, and had NO<sub>3</sub>-N loads 1.5 to 4.4 times higher than loads from two less-agricultural watersheds. Agricultural nonpoint sources are a major contributor to nutrient loading to some stream systems (Kemp 1949; Scheuler and Kemp 1979; Kuenzler and Craig 1986).

Forested floodplains can be effective filters for nutrient materials in lateral runoff and ground water. A 50-m-wide riparian forest in an agricultural watershed near the Chesapeake Bay in Maryland removed about 89% of the nitrogen and 80% of the phosphorus that entered the forest from upland runoff, ground water, and bulk precipitation (Peterjohn and Correll 1984). Correll (1987) concluded that protection and proper management of riparian forest buffer strips between cropland and urban areas along the first order streams in the watershed of Chesapeake Bay would prevent most of the nitrate from these watershed sources from being discharged into the bay. Kuenzler et al. (1980) reported a significant retention of phosphorous by a coastal plain floodplain swamp in North Carolina. Other studies have provided evidence that forested riparian floodplains reduce sediment and phosphorous loads in adjacent streams (e.g., McColl 1978; Schlosser and Karr 1981a,b). The same general conclusion that forested floodplains effectively reduce levels of nutrients from agricultural lands to receiving waters has been reached as a result of numerous studies (e.g., Lowrance et al. 1983, 1984a,b,c; Correll et al. 1984, 1986; Cooper et al. 1986; Peterjohn and Correll 1986; Schnabel 1986; Gilliam et al. 1988; Correll and Weller, in press).

## SUMMARY AND CONCLUSION

Palustrine wetlands of forested riparian floodplains were defined as freshwater wetlands that are coupled to upland watersheds and to adjacent streams. Coupling between the floodplain and the watershed forms a continuum. Coupling to the associated stream is via a "water bridge," at least during flooding and generally during the growing season. The associated stream forms a continuum to downstream aquatic systems. This openness of a stream corridor ecosystem strongly suggests that a systems approach be taken in the examination of the relationship between wetlands of forested riparian floodplains and fishery resources. Numerous studies show that many species of fish and several species of shellfish use palustrine wetlands of forested riparian floodplains, but few studies have been made to quantify the value of these wetlands to fisheries production. Fish yields to man by many of the major river floodplains of the world have been estimated to be 4,000 to 6,000 kg/km<sup>2</sup>/year/maximum area flooded (Welcomme 1979). In the United States, few if any floodplain fisheries are fully exploited, and fish production on floodplains has not been adequately quantified. The average annual harvest of fish and shellfish from floodplains in Louisiana has been estimated to be as high as 8,797 kg/km<sup>2</sup> (Lambou 1984).

It is generally known, but not sufficiently quantified or substantiated, that palustrine wetlands of forested riparian floodplains import, store, produce, and recycle materials used in food chains in situ by numerous organisms, including many species of fish. In most, if not all of these wetlands, some residual materials are exported to downstream aquatic systems where the materials are available for use in food chains of fish. Thus, palustrine wetlands of forested riparian floodplains may be used by fish for feeding, spawning, nursery and rearing habitat, or as a source of materials used in food chains at downstream sites. This openness emphasizes that one of the fundamental functions of wetland primary production is providing energy flow to detrital food webs. Detritus-based food webs add stability to aquatic systems by making the seasonally fixed photosynthetic energy available to fish, shellfish, and other consumers over longer periods of time and at sites remote from where the detritus was produced.

All wetlands exist as a result of hydrologic regimes. Food chain support, openness, productivity, species diversity and structure, and nutrient cycling are some of the important characteristics of forested floodplains that respond to hydrologic regimes. Buffer strips of forested riparian wetlands effectively reduce levels of nutrients and possibly levels of pesticides from agricultural lands reaching receiving waters. The importance of the relationship between the stream hydroperiod and floodplain is undeniable. If either is altered the other will surely change. For example, stream channelization or stream impoundment may or may not reduce wetland area on the adjacent floodplain, but either activity may drastically alter the quality and quantity of the wetland by altering its hydroperiod. Timing, magnitude, and duration of flooding are the primary determinants of floodplain structure and function. We have gained

a considerable amount of insight on how palustrine wetlands of forested riparian floodplains function. However, more information is required to help ensure that these wetlands are managed in a manner that provides the greatest benefits to society. Three research topics that need immediate attention are (1) trophic relationships between fish communities of the wetland and associated streams and estuaries, (2) relationships between watershed land use and the wetlands, and (3) the effects of hydrology (i.e., timing, extent, duration, and frequency of flooding) on ecological processes (i.e., productivity, nutrient cycling, food chain support, and species diversity and structure) in the wetland. Research in each of these areas should lead to better guidelines for protecting and managing palustrine wetlands of forested riparian floodplains and the fishery resources that these wetlands sustain.

## LITERATURE CITED

- Abernethy, Y., and R.E. Turner. 1987. U.S. forested wetlands: 1940-1980. *BioScience* 37:721-727.
- Bedinger, M.S. 1979. Relation between forest species and flooding. Pages 427-435 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. *Wetland function and values: the state of our understanding*. American Water Resources Assoc., Minneapolis, MN.
- Bedinger, M.S. 1981. Hydrology of bottomland hardwood forests of the Mississippi embayment. Pages 161-176 in J.R. Clark and J. Benforado, eds. *Wetlands of bottomland hardwood forests*. Elsevier Scientific Pub. Co., New York.
- Boyd, H.E. 1976. Biological productivity in two Georgian river swamps. Ph.D. Dissertation. University of Tennessee, Knoxville. 110 pp.
- Brinson, M.M., A.E. Lugo, and S. Brown. 1981a. Primary productivity, decomposition, and consumer activity in freshwater wetlands. *Annu. Rev. Ecol. Syst.* 12:123-161.
- Brinson, M.M., B.L. Swift, R.C. Plantico, and J.S. Barclay. 1981b. Riparian ecosystems: their ecology and status. U.S. Fish Wildl. Serv. FWS/OBS-81/17. 155 pp.
- Brinson, M.M., H.D. Bradshaw, and R.N. Holmes. 1983. Significance of floodplain sediments in nutrient exchange between a stream and its floodplain. Pages 199-219 in T.D. Fontaine, III, and S.M. Bartell, eds. *Dynamics of lotic ecosystems*. Ann Arbor Science, Ann Arbor, MI.
- Brown, S., M.S. Brinson, and A.E. Lugo. 1979. Structure and function of riparian wetlands. Pages 17-31 in R.R. Johnson and J.F. McCormick, technical coordinators. *Strategies for protection and management of floodplain wetlands and other riparian ecosystems*. U.S. Forest Service, Washington, DC Pub. GTR-WO-12. 410 pp.
- Bryan, C.F., F.M. Truesdale, and D.S. Sabins. 1975. Annual Report: a limnological survey of the Atchafalaya Basin. Louisiana Cooperative Fish and Wildlife Research Unit, Louisiana State University, Baton Rouge. 203 pp.
- Bryan, C.F., D.J. DeMont, D.S. Sabins, and J.P. Newman, Jr. 1976. Annual report: a limnological survey of the Atchafalaya Basin. Louisiana Cooperative Fish and Wildlife Research Unit, School of Forestry and Wildlife Management. Louisiana State University, Baton Rouge. 285 pp.

- Bryan, C.F., and D.S. Sabins. 1979. Management implications in water quality and fish standing stock information in the Atchafalaya River Basin, Louisiana. Pages 293-316 in J.W. Day, Jr., D.D. Culley, Jr., R.E. Turner, and A.J. Mumphrey, Jr., eds. Third coastal marsh and estuary symposium. Louisiana State University, Division of Continuing Education, Baton Rouge.
- Carlson, J.E., and M.J. Duever. 1977. Seasonal fish population fluctuations in a South Florida Swamp. Proc. Annu. Conf. Southeast Assoc. Fish Wildl. Agencies. 31:603-611.
- Carter, V., M.S. Bedinger, R.P. Novitzki, and W.O. Wilen. 1979. Water resources and wetlands. Pages 349-376 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. Wetland functions and values: the state of our understanding. American Water Resources Assoc., Minneapolis, MN.
- Chambers, D.G. 1980. An analysis of nekton communities in the upper Barataria Basin, Louisiana. M.S. Thesis. Louisiana State University, Baton Rouge. 286 pp.
- Clark, J.E. 1979. Freshwater wetlands: habitat for aquatic invertebrates, amphibians, reptiles, and fish. Pages 330-343 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. Wetland functions and values: the state of our understanding. American Water Resources Assoc., Minneapolis, MN.
- Clark, J.R., and J.E. Benforado, eds. 1981. Wetlands of bottomland hardwood forests. Elsevier Scientific Pub. Co., Amsterdam. 401 pp.
- Conner, W.H., and J.W. Day. 1976. Productivity and composition of a bald cypress-water tupelo site and a bottomland hardwood site in a Louisiana swamp. Am. J. Bot. 63:1354-1364.
- Conner, W.H., and J.W. Day, Jr. 1982. The ecology of forested wetlands in the southeastern United States. Pages 69-87 in B. Gopal, R.E. Turner, R.G. Wetzel, and D.F. Wigham, eds. Wetlands: ecology and management. National Institute of Ecology and International Scientific Publications, Jaipur, India.
- Cooper, J.R., J.W. Gilliam, and T.C. Jacobs. 1986. Riparian areas as a control of nonpoint pollutants. Pages 166-192 in D.L. Correll, ed. Watershed research perspectives. Smithsonian Institute Press, Washington, DC.
- Correll, D.L. 1987. Nutrients in Chesapeake Bay. Pages 298-320 in S.K. Majumber, L.W. Hall, Jr., and H.M. Austin, eds. Contaminant problems and management of living Chesapeake Bay resources. Pennsylvania Academy of Science, Easton, PA.
- Correll, D.L., N.M. Goff, and W.T. Peterjohn. 1984. Ion balances between precipitation inputs and Rhode River watershed discharges. Pages 77-111 in O.P. Bricker, ed. Geological aspects of acid deposition. Butterworth Publications, Boston, MA.

- Correll, D.L., J.J. Miklas, A.H. Hines, and J.J. Schafer. 1986. Chemical and biological trends associated with acidic atmospheric deposition in the Rhode River watershed and estuary. *Water, Air, Soil Pollution* 35:63-86.
- Correll, D.L., and D.E. Weller. Factors limiting processes in freshwater wetlands: an agricultural primary stream riparian forest. *Freshwater Wetlands and Wildlife Ninth Symposium*, Savannah River Ecology Laboratory. In press.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish Wildl. Serv. FWS/OBS-79/31. 131 pp.
- Crow, J.H., and K.B. Macdonald. 1979. Wetland values: secondary production. Pages 146-161 in P. Greeson, J.R. Clark, and J.E. Clark, eds. *Wetland functions and values: the state of our understanding*. American Water Resources Assoc., Minneapolis, MN.
- Day, J.W., Jr., T.J. Butler, and W.H. Conner. 1977. Productivity and nutrient export studies in a cypress swamp and lake system in Louisiana. Pages 255-269 in M. Wiled, ed. *Estuarine processes*, vol. 2. Academic Press, New York.
- Day, J.W., Jr., W.H. Conner, and G.P. Kemp. 1980. Contribution of wooded swamps and bottomland forests to estuarine productivity. Pages 33-65 in P.L. Fore and R.D. Peterson, eds. *Proceedings of the Gulf of Mexico coastal ecosystems workshop*. U.S. Fish Wildl. Serv. FWS/OBS-80/30.
- Day, J.W., Jr., W.H. Conner, G.P. Kemp, and D.G. Chambers. 1981. The relationship of estuarine productivity to wooded swamps and bottomland hardwood forests in the southeastern U.S. Pages 193-213 in R.C. Carey, P.S. Markevits, and J.B. Kirkwood, eds. *Proceedings of the workshop on coastal ecosystems of the southeastern United States*. U.S. Fish Wildl. Serv. FWS/OBS-80/59.
- Ewel, K.C. 1978. Riparian ecosystems: conservation of their unique characteristics. Pages 56-62 in *Strategies for protection and management of floodplain wetlands and other riparian ecosystems*. General Tech. Rep. WO-12. U.S. Dept. Agric., Forest Service, Washington, DC.
- Framer, W.E., T.J. Monahan, D.C. Bowden, and F.A. Graybill. 1983. Status and trends of wetlands and deepwater habitats in the conterminous United States, 1950's to 1970's. Dept. Forest and Wood Sciences, Colorado State University, Fort Collins. 32 pp.
- Fredrickson, L.H. 1979a. Lowland hardwood wetlands: current status and habitat values for wildlife. Pages 296-306 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. *Wetland functions and values: the state of our understanding*. American Water Resources Assoc., Minneapolis, MN.

- Fredrickson, L.H. 1979b. Floral and faunal changes in lowland hardwood forests in Missouri resulting from channelization, drainage, and impoundment. U.S. Fish Wildl. Serv. FWS/OBS-78/91. 130 pp.
- Fritzell, E.K. 1988. Mammals and wetlands. Pages 213-226 in D.D. Hook et al., eds. The ecology and management of wetlands, vol. I: ecology of wetlands. Timber Press, Portland, OR.
- Germann, J.F. 1973. Population studies of streams. Suwannee River, Annu. Progress Rep., Statewide Fisheries Investigation F-21-4, Study XVI, Job 2. Game and Fish Comm., Georgia Dept. Natural Resources, Atlanta. 13 pp.
- Gilliam, J.W., R.B. Daniels, and J.F. Lutz. 1974. Nitrogen contents of shallow groundwater in the North Carolina Coastal Plain. J. Environ. Qual. 3:147-151.
- Gilliam, J.W., G.W. Cheschire, and R.W. Skaggs. 1988. Effects of pumped agricultural drainage water on wetland water quality. Pages 275-283 in D.O. Hook et al., eds. The ecology and management of wetlands, vol. 2: management, use, and value of wetlands. Timber Press, Portland, OR.
- Golet, F.C., and J.S. Larson. 1974. Classification of freshwater wetlands in the glaciated northeast. U.S. Fish Wildl. Serv. Resour. Publ. 16. 56 pp.
- Good, R.E., D.F. Whigham, and R.L. Simpson, eds. 1978. Freshwater wetlands: ecological processes and management potential. Academic Press, New York. 378 pp.
- Gosselink, J.G., and R.E. Turner. 1978. The role of hydrology in freshwater wetland ecosystems. Pages 63-77 in R.E. Good, D.F. Wigham, and R.L. Simpson, eds. Freshwater wetlands: ecological processes and management potential. Academic Press, New York.
- Gosselink, J.G., W.H. Conner, Jr., and R.E. Turner. 1981. Classification of wetland resources: land, timber, and ecology. Pages 28-48 in B.D. Jackson and J.L. Chambers, eds. Timber harvesting in wetlands. Louisiana State University, Division of Continuing Education, Baton Rouge.
- Greeson, P.E., J.R. Clark, and J.E. Clark, eds. 1979. Wetland functions and values: the state of our understanding. Proc. Nat. Symp. Wetlands, Lake Buena Vista, FL. Am. Water Resour. Assoc., Tech. Publ. TPS-79-2, Minneapolis, MN. 674 pp.
- Hastings, R.W. 1979. Fish of the Pine Barrens. Pages 489-504 in R.T.T. Forman, ed. Pine Barrens: ecosystems and landscape. Academic Press, New York.
- Hinchee, R.E. 1977. Selected aspects of the biology of Lake Pontchartrain, Louisiana. M.S. Thesis. Louisiana State University, Baton Rouge.

- Holder, D.R. 1970. A study of fish movements from the Okefenokee Swamp into the Suwannee River. Proc. Annu. Conf. Southeast Assoc. Fish. Wildl. Agencies. 24:591-608.
- Holder, D.R. 1971. Benthos studies in warmwater streams. Statewide Fisheries Investigation, Annu. Prog. Rep. Project F-21-2, Georgia Game and Fish Comm., Dept. of Natural Resources, Atlanta. 17 pp.
- Holder, D.R., L. McSwain, W.D. Hill, Jr., W. King, and C. Sweet. 1970. Population studies - streams. Statewide Fisheries Investigation, Annu. Prog. Rep. F-21-2, Study XVI, Job 2. Game and Fish Comm., Georgia Dept. Natural Resources, Atlanta. 66 pp.
- Holder, D.R., J. Sandow, L. McSwain, W.D. Hill, Jr., W. King, and C. Sweet. 1971. Population studies - streams. Statewide Fisheries Investigation, Annu. Prog. Rep. F-21-3, Study XVI, Job 2. Game and Fish Comm., Georgia Dept. Nat. Resources, Atlanta. 42 pp.
- Johnson, R.L. 1979. Timber harvests from wetlands. Pages 598-605 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. Wetland function and values: the state of our understanding. American Water Resource Assoc., Minneapolis, MN.
- Kadlec, J.A. 1987. Nutrient dynamics in wetlands. Pages 393-419 in K.R. Reddy and W.H. Smith, eds. Aquatic plants for water treatment and resource recovery. Magnolia Pub. Inc., Orlando, FL.
- Kaushik, N.K., and H.B.N. Hynes. 1971. The fate of the dead leaves that fall into streams. Arch. Hydrobiol. 68:465-515.
- Kemp, H. 1949. Soil pollution in the Potomac River basin. Interstate Commission on the Potomac River Basin, Washington, DC. 21 pp.
- Klimas, C.V. 1988. River regulation effects on floodplains hydrology and ecology. Pages 40-49 in D.D. Hook et al., eds. The ecology and management of wetlands, vol 1. Timber Press, Portland, OR.
- Konikoff, M. 1977. Studies of the life history and ecology of the red swamp crawfish, Procambarus clarkii, in the lower Atchafalaya basin floodway. Final Report to U.S. Fish Wildl. Serv. Dept. Biology, University of Southeastern Louisiana, Lafayette. 81 pp.
- Kroodsman, D.E. 1979. Habitat values for non-game wetland birds. Pages 320-329 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. Wetland functions and values: the state of our understanding. American Water Resources Assoc., Minneapolis, MN.
- Kuenzler, E.J., P.J. Mulholland, L.A. Yarbrow, and L.A. Smock. 1980. Distributions and budgets of carbon, phosphorous, iron, and manganese in a floodplain swamp ecosystem. Water Resources Research Institute Report 15-7. University North Carolina, Chapel Hill. 234 pp.

- Kuenzler, E.J., and N.J. Craig. 1986. Land use and nutrient yields of the Chowan River watershed. Pages 77-107 in D.L. Correll, ed. Watershed research perspective. Smithsonian Institution Press, Washington, DC.
- Kusler, J.A. 1983. Our national wetland heritage: a protection guidebook. Environmental Law Institute, Washington, DC. 167 pp.
- Lambou, V.W. 1965. The commercial and sport fisheries of the Atchafalaya Basin Floodway. Proc. 17th Annu. Conf. Southeast Assoc. Game Fish Comm. 17:256-281.
- Lambou, V.W. 1984. Importance of bottomland hardwood forest zones to fish and fisheries: the Atchafalaya Basin a case history. Paper presented at workshop concerning ecological zonation in bottomland hardwoods, St. Francisville, LA, December 3-7, 1984.
- Langdon, O.G., J.P. McClure, D.D. Hook, J.M. Crockett, and R. Hunt. 1981. Extent, condition, management, and research needs of bottomland hardwoods-cypress forests in the southeastern United States. Pages 71-80 in J.R. Clark and J. Benforado, eds. Wetlands of bottomland hardwood forests. Elsevier Scientific Pub. Co., New York.
- Larson, J.S., M.S. Bedinger, C.F. Bryan, S. Brown, R.T. Huffman, E.L. Miller, D.G. Rhodes, and B.A. Touchet. 1981. Transition from wetlands to uplands in southeastern bottomland hardwood forests. Pages 225-273 in Wetlands of bottomland hardwood forests. Elsevier Scientific Pub. Co., New York.
- Lea, R. 1988. Management of eastern United States bottomland hardwood forests. Pages 185-194 in D.O. Hook et al., eds. The ecology and management of wetlands, vol. 2: management, use, and value of wetlands. Timber Press, Portland, OR.
- Leitman, H.M., J.E. Sohm, and M.A. Franklin. 1984. Wetland hydrology and tree distribution of the Apalachicola River flood plain. Water Supply Paper 2196-A, U.S. Geological Survey, Washington, DC. 52 pp.
- Livingston, R.J., and O.L. Loucks. 1979. Productivity, trophic interactions, and food-web relationships in wetlands and associated systems (theme paper). Pages 101-119 in P. Greeson, J.R. Clark, and J.E. Clark, eds. Wetland functions and values: the state of our understanding. American Water Resources Assoc., Minneapolis, MN.
- Lowrance, R.R., R.L. Todd, and L.E. Asmussen. 1983. Waterborne nutrient budgets for the riparian zone of an agricultural watershed. Agriculture, ecosystems, and environment 10:371-384. Elsevier Science Pub. Co., Amsterdam.
- Lowrance, R.R., R.L. Todd, and L.E. Asmussen. 1984a. Nutrient cycling in an agricultural watershed: I. phreatic movement. J. Environ. Qual. 13:22-27.


- Lowrance, R.R., R.L. Todd, and L.E. Asmussen. 1984b. Nutrient cycling in an agricultural watershed: II. streamflow and artificial drainage. *J. Environ. Qual.* 13:27-32.
- Lowrance, R., R. Todd, J. Fail, Jr., O. Hendrickson, Jr., R. Leonard, and L. Asmussen. 1984c. Riparian forests as nutrient filters in agricultural watersheds. *BioScience* 34:374-377.
- Lowrance, R.R., R.A. Leonard, L.E. Asmussen, and R.L. Todd. 1985. Nutrient budgets for agricultural watersheds in the southeastern coastal plain. *Ecology* 66:287-296.
- Martin, A.C., N. Hutchkiss, F.M. Uhler, and W.S. Bourn. 1953. Classification of wetlands of the United States. U.S. Fish Wildl. Serv. Spec. Sci. Rep. Wildlife 20. 14 pp.
- McColl, R.H.S. 1978. Chemical runoff from pastures: the influence of fertilizer and riparian zones. *New Zealand Jour. Marine and Freshwater Research* 12:371-380.
- McKnight, J.S., D.D. Hook, O.G. Langdon, and R.L. Johnson. 1981. Flood tolerance and related characteristics of trees of the bottomland forests of the southern United States. Pages 29-69 in J.R. Clark and J. Benforado, eds. *Wetlands of bottomland hardwood forests*. Elsevier Scientific Pub. Co., New York.
- Melchiorri-Santolini, V., and J.W. Hampton, eds. 1972. Detritus and its ecological role in aquatic systems. Proc. IBP-UNESCO Symp. Mem. Inst. Ital. Idrobiol., Pallanza, Italy.
- Millar, J.B. 1976. Wetland classification in western Canada: a guide to marshes and shallow open water wetlands in the grasslands and parklands of the Prairie Provinces. *Can. Wildl. Serv. Rep. Ser.* 37. 38 pp.
- Mitsch, W.J., and J.G. Gosselink. 1986. *Wetlands*. Van Nostrand Reinhold Co., New York. 539 pp.
- Niering, W.A. 1988. Endangered, threatened, and rare wetland plants and animals of the continental United States. Pages 227-238 in D.D. Hook et al., eds. *The ecology and management of wetlands, vol. I: ecology of wetlands*. Timber Press, Portland, OR.
- Parsons, K., and C.H. Wharton. 1978. Macroinvertebrates of pools on a Piedmont river floodplain. *Georgia J. Sci.* 36:25-33.
- Patrick, R., J. Cairns, Jr., and S.S. Roback. 1967. An ecosystematic study of the fauna and flora of the Savannah River. *Proc. Acad. Nat. Sci. Phila.* 118(5):109-407.
- Patrick, R., B. Matson, and L. Anderson. 1979. Streams and lakes in the Pine Barrens. Pages 169-193 in R.T.T. Forman, ed. *Pine Barrens: ecosystem and landscape*. Academic Press, New York.

- Penn, G.H. 1950. Utilization of crawfishes by cold-blooded vertebrates in the eastern U.S. *Am. Midl. Nat.* 44:643-658.
- Peterjohn, W.T., and D.L. Correll. 1984. Nutrient dynamics in an agricultural watershed: observations on the role of a riparian forest. *Ecology* 65:1466-1475.
- Peterjohn, W.T., and D.L. Correll. 1986. The effect of riparian forest on the volume and chemical composition of baseflow in an agricultural watershed. Pages 244-262 in D.L. Correll, ed. *Watershed research perspectives*. Smithsonian Institution Press, Washington, DC.
- Peters, D.S., D.W. Ahrenholz, and T.R. Rice. 1979. Harvest and value of wetland associated fish and shellfish. Pages 606-617 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. *Wetland functions and values: the state of our understanding*. American Water Resources Assoc., Minneapolis, MN.
- Pollard, J.E., S.M. Melancon, and L.S. Blakey. 1983. Importance of bottomland hardwoods to crayfish and fish in the Henderson Lake area. *Atchafalaya Basin, Louisiana. Wetlands* 3:1-25.
- Richardson, C.J., ed. 1981. *Pocosin wetlands: an integrated analysis of coastal plain freshwater bogs in North Carolina*. Hutchinson Ross, Stroudsbury, PA. 364 pp.
- Roe, H.B., and Q.C. Ayers. 1954. *Engineering for agricultural drainage*. McGraw Hill, New York. 501 pp.
- Roele, J.E., G.T. Auble, D.B. Hamilton, G.C. Horak, R.L. Johnson, and C.A. Segelquist, eds. 1987. *Results of a workshop concerning impacts of various activities on the function of bottomland hardwoods*. U.S. Fish. Wildl. Serv., National Ecology Research Center, Fort Collins, CO. NEC-87/15. 171 pp.
- Risotto, S.P., and R.E. Turner. 1985. Annual fluctuation in abundance of the commercial fisheries of the Mississippi River and tributaries. *N. Am. J. Fish. Manage.* 5:557-574.
- Sather, J.H., and R.D. Smith. 1984. An overview of major wetland functions and values. U.S. Fish Wildl. Serv. FWS/OBS-84/18. 68 pp.
- Scheuler, T.R., and W.M. Kemp. 1979. An agricultural mass balance model of nitrogen movement in the Choptank River basin. Pages 80-86 in W.M. Kemp, W. Boynton, and J.W. Stevenson, eds. *Submerged aquatic vegetation in the Chesapeake Bay*. Annu. Rep. EPA/CBF, Vol. 1. UMCEES Ref. No. 80-68-HPEL.
- Schitoskey, F., Jr. 1979. Use of wetlands by upland game. Pages 307-319 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. *Wetland functions and values: the state of our understanding*. American Water Resources Assoc., Minneapolis, MN.

- Schlosser, I.J., and J.R. Karr. 1981a. Water quality in agricultural watersheds: impact of riparian vegetation during base flow. *Water Resour. Bull.* 17:233-240.
- Schlosser, I.J., and J.R. Karr. 1981b. Riparian vegetation and channel morphology impact on spatial patterns of water quality in agricultural watersheds. *Environ. Manage.* 5:233-243.
- Schnabel, R.R. 1986. Nitrate concentrations in a small stream as affected by chemical and hydrologic interactions in the riparian zone. Pages 263-282 in D.L. Correll, ed. *Watershed research perspectives*. Smithsonian Institution Press, Washington, DC.
- Scott, M.L., and J.L. Haskins. 1987. Effects of grazing by chrysomelid beetles on two wetland herbaceous species. *Bull. Torrey Botanical Club* 114:13-17.
- Shaw, S.P., and C.G. Fredine. 1956. Wetlands of the United States: their extent, and their value for waterfowl and other wildlife. U.S. Fish Wildl. Serv. Circular 39. 67 pp.
- Stewart, R.E., and H.A. Kantrud. 1971. Classification of natural ponds and lakes in the glaciated prairie region. U.S. Fish Wildl. Serv. Res. Publ. 92. 57 pp.
- Strickland, R., ed. 1986. Wetland functions, rehabilitation, and creation in the Pacific Northwest: the state of our understanding. Proc. April 30-May 2, 1986 conf. Washington State University, Dept. Ecology, Olympia. 184 pp.
- Taylor, J.R., M.A. Cardamone, and M.J. Mitsch. 1984. Bottomland hardwood forests: their functions and values. Paper presented at Bottomlands Hardwoods Workshop, St. Francisville, Louisiana, December 1984. Mitch and Associates, Louisville, KY. 87 pp.
- Walker, M. 1980. Utilization by fishes of a blackwater creek floodplain in North Carolina. M.S. Thesis. E. Carolina University, Greenville, NC.
- Wallace, J.E., and J. O'Hop. 1985. Life on a fast pad: waterlily leaf beetle impact on waterlilies. *Ecology* 66:1534-1544.
- Welcomme, R.L. 1979. Fisheries ecology of floodplain rivers. Longman, New York. 317 pp.
- Welcomme, R.L. 1985. River floodplains. FAO Fisheries Tech. Paper 262. FAO, Rome, Italy. 330 pp.
- Wharton, C.H. 1977. The natural environments of Georgia. Georgia Geological Survey, Dept. of Natural Resources, Atlanta. 227 pp.
- Wharton, C.H. 1980. Values and functions of bottomland hardwoods. Trans. N. Am. Wildl. Nat. Resour. Conf. 45:341-353.

- Wharton, C.H., and M.M. Brinson. 1979. Characteristics of southeastern river systems. Pages 32-40 in R.R. Johnson and J.F. McCormick, technical coordinators. Strategies for protection and management of floodplain wetlands and other riparian ecosystems, U.S. Forest Serv. Publ. GTR-WO-12.
- Wharton, C.H., V.W. Lambou, J. Newsom, P.V. Winger, L.L. Gaddy, and R. Mancks. 1981. The fauna of bottomland hardwoods in southeastern United States. Pages 87-160 in J.R. Clark and J. Benforado, eds. Wetlands of bottomland hardwood forests. Elsevier Scientific, New York.
- Wharton, C.H., W.H. Kitchens, E.C. Pendleton, and T.W. Snipe. 1982. The ecology of bottomland hardwood swamps of the southeast: a community profile. FWS/OBS-81/37. 133 pp.
- Wilkinson, D.L., K. Schneller-McDonald, R.W. Olson, and G.T. Auble. 1987. Synopsis of wetland functions and values: bottomland hardwoods with special emphasis on eastern Texas and Oklahoma. U.S. Fish Wildl. Serv. Biol. Rep. 87(12). 132 pp.
- Williams, J.D., and C.K. Dodd, Jr. 1979. Importance of wetlands to endangered and threatened species. Pages 565-575 in P.E. Greeson, J.R. Clark, and J.E. Clark, eds. Wetlands functions and values: the state of our understanding. American Water Resources Assoc., Minneapolis, MN.
- Winger, P.V. 1986. Forested wetlands of the southeast: review of major characteristics and role in maintaining water quality. U.S. Fish Wildl. Serv. Res. Publ. 163. 16 pp.
- Wyatt, H., and D.R. Holder. 1969. Life history studies. Annu. Rep. Federal Aid to Fish Restoration F-19-R-3 (July 1, 1967 - June 30, 1968). Georgia Dept. Natural Resources, Game and Fish Commission, Atlanta.
- Zinn, J.A., and C. Copeland. 1982. Wetland management. Congr. Res. Serv., Library of Congress, Washington, DC. 149 pp.
- Ziser, S.W. 1978. Seasonal variations in water chemistry and diversity of the phytaphilia macroinvertebrates of three swamp communities in southeastern Louisiana. Southwest Nat. 23:545-562.
- Zoltai, S.C., F.C. Pollett, J.K. Jeglum, and G.D. Adams. 1975. Developing a wetland classification for Canada. Proc. N. Am. Soils Conf. 4:497-511.

<b>REPORT DOCUMENTATION PAGE</b>		<b>1. REPORT NO.</b> Biological Report 88(32)	<b>2.</b>	<b>3. Recipient's Accession No.</b>
<b>4. Title and Subtitle</b> Relationships Between Palustrine Forested Wetlands of Forested Riparian Floodplains and Fishery Resources: a review			<b>5. Report Date</b> August 1988	
<b>7. Author(s)</b> J. H. Crance			<b>6.</b>	
<b>9. Performing Organization Name and Address</b> National Ecology Research Center U.S. Fish and Wildlife Service Creekside One Bldg., 2627 Redwing Rd. Fort Collins, CO 80526-2899			<b>8. Performing Organization Rept. No.</b>	
<b>12. Sponsoring Organization Name and Address</b> U.S. Department of the Interior Fish and Wildlife Service Research and Development Washington, DC 20240			<b>10. Project/Task/Work Unit No.</b>	
			<b>11. Contract(C) or Grant(G) No.</b> (C) (G)	
			<b>13. Type of Report &amp; Period Covered</b>	
<b>15. Supplementary Notes</b>			<b>14.</b>	
<b>16. Abstract (Limit: 200 words)</b> <p>This literature review should help fishery, wetland, and water resource managers focus on management approaches and research for the identification and mitigation of wetland impacts and on the importance of wetlands to fisheries. Wetlands and their attendant fisheries exist as a result of hydrology. Productivity, species diversity and structure, and nutrient cycling are wetland characteristics that respond to hydrologic regimes. Anthropogenic changes often appear subtle but can result in ecological catastrophes. Deterioration of hydrologic regimes of these wetlands is likely to reduce habitat quality or quantity and may block fish access. Wetland-related fishery resources should be given high priority when regulated stream flow, channelization, diking, and watershed practices are being evaluated.</p>				
<b>17. Document Analysis a. Descriptors</b> Riparian ecosystems				
<b>b. Identifiers/Open-Ended Terms</b> Floodplain wetlands				
<b>c. COSATI Field/Group</b>				
<b>18. Availability Statement</b> Release unlimited		<b>19. Security Class (This Report)</b> Unclassified		<b>21. No. of Pages</b> 27
		<b>20. Security Class (This Page)</b> Unclassified		<b>22. Price</b>



**REGION 1**

Regional Director  
 U.S. Fish and Wildlife Service  
 Lloyd Five Hundred Building, Suite 1692  
 500 N.E. Multnomah Street  
 Portland, Oregon 97232

**REGION 2**

Regional Director  
 U.S. Fish and Wildlife Service  
 P.O. Box 1306  
 Albuquerque, New Mexico 87103

**REGION 3**

Regional Director  
 U.S. Fish and Wildlife Service  
 Federal Building, Fort Snelling  
 Twin Cities, Minnesota 55111

**REGION 4**

Regional Director  
 U.S. Fish and Wildlife Service  
 Richard B. Russell Building  
 75 Spring Street, S.W.  
 Atlanta, Georgia 30303

**REGION 5**

Regional Director  
 U.S. Fish and Wildlife Service  
 One Gateway Center  
 Newton Corner, Massachusetts 02158

**REGION 6**

Regional Director  
 U.S. Fish and Wildlife Service  
 P.O. Box 25486  
 Denver Federal Center  
 Denver, Colorado 80225

**REGION 7**

Regional Director  
 U.S. Fish and Wildlife Service  
 1011 E. Tudor Road  
 Anchorage, Alaska 99503



*Preserve Our Natural Resources*



**DEPARTMENT OF THE INTERIOR**  
**U.S. FISH AND WILDLIFE SERVICE**



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.