


FEDERAL AND STATE ENDANGERED AND THREATENED SPECIES EXPENDITURES



Fiscal Year 2022

TABLE OF CONTENTS

ENDANGERED AND THREATENED SPECIES EXPENDITURES FISCAL YEAR 2022	
PURPOSE	3
BACKGROUND	4
SECTION 18(1), FEDERAL EXPENDITURES	5
SECTION 18(2), STATE EXPENDITURES	5
SUMMARY OF DATA	5
COMPARISON TO PREVIOUS REPORTS	6
Table 1. FY 2022 Reported Expenditures for Endangered and Threatened Species	7
Table 2. Species Ranked in Descending Order of Total FY 2022 Reported Expenditures	114
Table 3. FY 2022 Reported Land Acquisition Expenditures for Endangered and Threatened Species	169
Table 4. FY 2022 Expenditures Reported by Individual Federal Agencies and the States	179
Appendix A. Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species	181
Appendix B. Reports Provided by Other Federal Agencies	185
Appendix C. Expenditures Reported by the States	262

ENDANGERED SPECIES EXPENDITURES FISCAL YEAR 2022

PURPOSE

The Endangered Species Act of 1973 (16 U.S.C. 1531 *et seq.*) (Act) was reauthorized and amended on October 7, 1988, under Public Law 100-478. A new section 18 was added to the Act that requires the U.S. Fish and Wildlife Service (Service) on behalf of the Secretary of the Interior to report annually on certain expenditures on a species-by-species basis. The section reads:

SEC. 18. On or before January 15, 1990, and each January 15 thereafter, the Secretary of the Interior, acting through the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain

(1) the accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act; and

(2) an accounting on a species-by-species basis of all reasonably identifiable expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act by States receiving grants under section 6.

This document constitutes the 34th report by the Service pursuant to the requirements of section 18 of the Act. The report covers fiscal year 2022 (October 1, 2021 - September 30, 2022), although some States may be reporting for a different fiscal year period (without duplicating a previous year's reported data). To assist Federal and State agencies with more accurate accounting of expenditures, a multi-species reporting category was added to the FY12 and subsequent reports. This option gives agencies another way to report expenditures when it is not possible to determine individual species costs for those species that are part of a multi-species conservation activity.

The term "State" as used in this report is defined in section 3(17) of the Act to mean any of the several States, the District of Columbia, the Commonwealth of Puerto Rico, American Samoa, the Virgin Islands, Guam, and the Trust Territory of the Pacific Islands.

The term "State agency" as used in this report is defined in section 3(18) of the Act to mean any State agency, department, board, commission, or other governmental entity which is responsible for the management and conservation of fish, plant, or wildlife resources within a State.

The term "conservation," as defined in section 3(3) of the Act, means the use of all methods and procedures necessary to bring any endangered or threatened species to the point where the measures provided for by the Act are no longer necessary--including, but not limited to, research, census, law enforcement, habitat acquisition and maintenance, propagation, live trapping, and transplantation. For the purposes of this report, the definition of conservation is further expanded to include any and all actions taken by Federal and State agencies on behalf of threatened or endangered species listed pursuant to the Act.

BACKGROUND

At the time the reporting requirement was enacted by Congress, the Service determined that a good faith effort was expected to develop methods to track and obtain data that are "reasonably identifiable." No extensive or extraordinary measures to develop exceptionally precise data were sought, nor were the reporting agencies expected to prorate staff salaries and other normal operational and maintenance costs not directed toward a particular species. The expenditures reported in this and the preceding reports do not necessarily constitute all expenditures made for each species.

This report represents a compilation of reasonably identifiable expenditures for the conservation of listed species reported independently by Federal and State agencies. Expenditures for FY 2022 are reported for 1,625 domestic threatened and endangered species of the 1,850 domestic species that were listed under United States jurisdiction in 50 CFR Part 17 during FY 2022 up through September 30, 2022 (Tables 1 and 3).

Expenditures are identified by listed entity, which may be a species¹, subspecies², Distinct Population Segment (DPS)³, or Evolutionarily Significant Unit (ESU)⁴, as identified in the Act's implementing regulations at 50 CFR 17.11 and 17.12.

These expenditures are reported separately for the Service and in aggregate for all other Federal agencies and the States (Tables 1 and 3).

For these annual reports, the Service requests that all reporting agencies submit their data on the species spreadsheet provided with the data call. At the time of submission, the reporting agencies certify that their information is accurate. The information provided in this report, pursuant to section 18 of the Endangered Species Act, is presented as it was submitted by these agencies. We do not verify an agency's data after the data are submitted to us.

Expenditures for candidate species or other species not listed in 50 CFR Part 17 as of the end of each fiscal year are not included in this report. In addition, other types of expenses that are not incurred primarily for the conservation of listed species, such as those for litigation, are not reported.

¹ **Species** For purposes of the ESA, this term includes any species or subspecies of fish or wildlife or plants, and any Distinct Population Segment of any species of vertebrate fish or wildlife that interbreeds when mature.

² **Subspecies** A taxonomic rank below that of species, usually recognizing individuals that have certain heritable characteristics distinct from other subspecies of a species.

³ **Distinct Population Segment** A subdivision of a vertebrate species that is treated as a species for purposes of listing under the Endangered Species Act.

⁴ **Evolutionarily Significant Unit** A Pacific salmonid stock that is substantially reproductively isolated from other stocks of the same species and which represents an important part of the evolutionary legacy of the species; functionally equivalent to DPS for the purpose of listing under ESA.

SECTION 18(1), FEDERAL EXPENDITURES

Expenditures Reported by the Fish and Wildlife Service

The Service's operational categories of expenditures include fisheries, refuges, land acquisition, law enforcement, research, international conservation, and regional and field office operations for listing, recovery, consultation, migratory birds, and grants. All reasonably identifiable expenditures that could be attributed to individual species within these activities have been included.

The Federal assistance portion under section 6 of the Act for each species is included in the FWS Total in Table 1. The State share for each species for which a Federal contribution was made is included in the States Total column of Table 1.

A portion of the Service's funds for consultation, listing, and recovery are incorporated into salaries, operations, and maintenance costs. If these costs contributed to the conservation of listed species and are reasonably identifiable to specific species, they are included in this report.

Expenditures Reported by other Federal Agencies

The Director of the Service sends a request to other Federal agencies identified as likely to have species-specific expenditures under the Act to report for each fiscal year. The Service provides guidance identifying what constitutes reasonably identifiable expenditures and how these are to be reported (Appendix A). For FY 2022, the Service and 11 other Federal agencies identified expenditures for the conservation of individual threatened and endangered species. Results from all responding Federal agencies are included in Appendix B. The data identified by those responding Federal agencies for individual species and multi-species conservation activities are tabulated in the “Other Fed” columns of this report (Tables 1 and 3).

SECTION 18(2), STATE EXPENDITURES

The assistance of the Association of Fish and Wildlife Agencies is sought by the Service to help coordinate the reporting of reasonably identifiable expenditures made by the States. States responding to this data call do so by reporting their data by species directly to the Service via our Web-based system. Collective results from all responding State agencies are included in Appendix C. For FY 2022, 26 State agencies identified expenditures for the conservation of individual threatened and endangered species.

SUMMARY OF DATA

Table 1 reports the activity-related expenditures that are reasonably identifiable to specific individual species as well as multi-species activity costs by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land acquisition (Table 3) are not included in this tabulation. Species are grouped in the same general order as they appear in 50 CFR Part 17. Table 1 also provides the rank for each species from Table 2.

Table 2 ranks each species in descending order of the species total from Table 1. A cumulative total of the activity-related, species-specific expenditures is also provided.

Table 3 displays land acquisition expenditures, reasonably identifiable to specific individual species, by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land purchased for multi-species conservation activities are listed at the end of the table. Table 3 also provides the rank for land expenditures by species.

Table 4 shows the combined total expenditures for species and land acquisition by reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species and multi-species conservation activities are included.

Total expenditures reported for domestic and foreign species in **FY 2022** were **\$ 834,772,899**, of which **\$ 801,427,923** was reported by Federal agencies and **\$ 33,344,976** was reported by the States (Table 4).

COMPARISON TO PREVIOUS REPORTS

Over the years, these reports have been refined as our experience in planning and coordinating with the reporting agencies to track such expenditures has improved. A good faith effort is made to develop species-by-species expenditures for these reports, but due to differences in reporting methods, this report cannot be easily compared to prior expenditures reports. These year-to-year differences include changes in the number of Federal and State agencies reporting data, changes in how expenditures are calculated, changes in the agencies' abilities to track expenditures, and changes in the number of listed species. Therefore, real dollar increases or decreases cannot accurately be calculated or inferred from the overall or individual totals.

TABLE 1. FY 2022 REPORTED EXPENDITURES FOR ENDANGERED AND THREATENED SPECIES

The list of species in Table 1 essentially follows the taxonomic and nomenclatural conventions used in 50 CFR Part 17 for the identification and sequence of species' names. For ease of reference, the alphabetized, inverted common name is provided within each group (usually Class) of animals and plants. Thus, the gray bat is found with other bats under "Bat, gray" within mammals, and Tar River spiny mussel is found under "Spiny mussel, Tar River" within the clams. Groups of animals follow the sequence: Amphibians, Birds, Clams, Conifers and Cycads, Corals, Crustaceans, Ferns and Allies, Fishes, Flowering Plants, Insects, Lichens, Mammals, Reptiles, and Snails.

For additional information about the names, historical range, or status of an endangered or threatened species, the species can be found in the Code of Federal Regulations at [50 CFR 17.11 \(list of wildlife\)](#) or [17.12 \(list of plants\)](#), or through our Web site at <http://www.fws.gov/endangered>. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

Fish and Wildlife Service expenditures (FWS Total) and other Federal agency expenditures (Other Fed) are totaled in the "Federal Total" column. The States Total is the amount reported to the Service directly by the States. None of these reported totals includes costs for land acquisition. (See Table 3 for reported land acquisition expenditures.)

The RANK is from Table 2 and indicates each species' rank by identifiable expenditures, where #1 is the highest expenditure.

Multi-species conservation activity-related expenditures are tabulated at the end of Table 1.

TABLE 1. REPORTED EXPENDITURES FOR FY 2022 FOR ENDANGERED AND THREATENED SPECIES, NOT INCLUDING LAND ACQUISITION COSTS

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
Amphibians	1416	Coqui, golden (<i>Eleutherodactylus jasperi</i>) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	708	Coqui, Llanero (<i>Eleutherodactylus</i> <i>juanariveroi</i>) -Wherever found	E	\$28,500	\$0	\$28,500	\$0	\$28,500
	94	Frog, California red-legged (<i>Rana draytonii</i>) -Wherever found	T	\$929,960	\$156,196	\$1,086,156	\$0	\$1,086,156
	456	Frog, Chiricahua leopard (<i>Rana</i> <i>chiricahuensis</i>) -Wherever found	T	\$0	\$67,565	\$67,565	\$0	\$67,565
	216	Frog, dusky gopher (<i>Rana sevosa</i>) - Wherever found	E	\$241,500	\$0	\$241,500	\$106,618	\$348,118
	300	Frog, mountain yellow-legged (<i>Rana</i> <i>muscosa</i>) -U.S.A., northern California	E	\$1,300	\$150,907	\$152,207	\$0	\$152,207
	299	Frog, mountain yellow-legged (<i>Rana</i> <i>muscosa</i>) -U.S.A., southern California	E	\$152,600	\$0	\$152,600	\$0	\$152,600
	45	Frog, Oregon spotted (<i>Rana pretiosa</i>) - Wherever found	T	\$3,617,872	\$13,815	\$3,631,687	\$0	\$3,631,687
	184	Frog, Sierra Nevada Yellow-legged (<i>Rana</i> <i>sierrae</i>) -Wherever found	E	\$109,700	\$325,907	\$435,607	\$0	\$435,607
	878	Guajon (<i>Eleutherodactylus cooki</i>) - Wherever found	T	\$17,500	\$0	\$17,500	\$0	\$17,500
	463	Hellbender, Eastern (<i>Cryptobranchus</i> <i>alleganiensis alleganiensis</i>) -Missouri DPS	E	\$39,000	\$0	\$39,000	\$25,500	\$64,500
	374	Hellbender, Ozark (<i>Cryptobranchus</i> <i>alleganiensis bishopi</i>) -Wherever found	E	\$50,500	\$15,552	\$66,052	\$40,500	\$106,552
	311	Salamander, California tiger (<i>Ambystoma</i> <i>californiense</i>) -U.S.A. (CA - Central California)	T	\$142,700	\$0	\$142,700	\$0	\$142,700
	123	Salamander, California tiger (<i>Ambystoma</i> <i>californiense</i>) -U.S.A. (CA - Santa Barbara County)	E	\$755,300	\$8,837	\$764,137	\$0	\$764,137
	42	Salamander, California tiger (<i>Ambystoma</i> <i>californiense</i>) -U.S.A. (CA - Sonoma County)	E	\$4,139,700	\$0	\$4,139,700	\$0	\$4,139,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	310	Salamander, Cheat Mountain (<i>Plethodon nettingi</i>) -Wherever found	T	\$140,500	\$0	\$140,500	\$3,137	\$143,637
	219	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) -Wherever found	T	\$295,000	\$0	\$295,000	\$40,531	\$335,531
	1507	Salamander, Georgetown (<i>Eurycea naufragia</i>) -Wherever found	T	\$0	\$0	\$0	\$4,000	\$4,000
	391	Salamander, Jemez Mountains (<i>Plethodon neomexicanus</i>) -Wherever found	E	\$0	\$96,000	\$96,000	\$0	\$96,000
	1507	Salamander, Jollyville Plateau (<i>Eurycea tonkawae</i>) -Wherever found	T	\$0	\$0	\$0	\$4,000	\$4,000
	661	Salamander, Red Hills (<i>Phaeognathus hubrichti</i>) -Wherever found	T	\$32,000	\$0	\$32,000	\$0	\$32,000
	306	salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) -Wherever found	E	\$109,000	\$0	\$109,000	\$40,531	\$149,531
	1338	Salamander, Salado (<i>Eurycea chisholmensis</i>) -Wherever found	T	\$0	\$0	\$0	\$8,000	\$8,000
	365	Salamander, Santa Cruz long-toed (<i>Ambystoma macrodactylum croceum</i>) -Wherever found	E	\$110,000	\$0	\$110,000	\$0	\$110,000
	466	Salamander, Shenandoah (<i>Plethodon shenandoah</i>) -Wherever found	E	\$3,000	\$60,000	\$63,000	\$0	\$63,000
	468	Toad, arroyo (=arroyo southwestern) (<i>Anaxyrus californicus</i>) -Wherever found	E	\$62,600	\$0	\$62,600	\$0	\$62,600
	410	Toad, Houston (<i>Bufo houstonensis</i>) -Wherever found	E	\$25,000	\$0	\$25,000	\$61,000	\$86,000
	987	Toad, Puerto Rican crested (<i>Peltophryne lemur</i>) -Wherever found	T	\$13,500	\$0	\$13,500	\$0	\$13,500
	234	Toad, Wyoming (<i>Anaxyrus baxteri</i>) -Wherever found	E	\$165,000	\$0	\$165,000	\$105,370	\$270,370
	773	Waterdog, black warrior (=Sipsey Fork) (<i>Necturus alabamensis</i>) -Wherever found	E	\$23,500	\$350	\$23,850	\$0	\$23,850
	663	Waterdog, Neuse River (<i>Necturus lewisi</i>) -Wherever found	T	\$22,000	\$3,200	\$25,200	\$6,500	\$31,700
	249	Yosemite Toad (<i>Anaxyrus canorus</i>) -Wherever found	T	\$161,900	\$75,591	\$237,491	\$0	\$237,491
	284	Toad, Dixie Valley (<i>Anaxyrus williamsi</i>) -	E	\$177,000	\$0	\$177,000	\$0	\$177,000
Amphibians subtotal				\$11,572,632	\$973,920	\$12,546,552	\$445,687	\$12,992,239

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
Arachnids	760	Spider, Kauai cave wolf or pe'e pe'e maka 'ole (Adelocosa anops) -Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	652	Spider, spruce-fir moss (Microhexura montivaga) -Wherever found	E	\$33,500	\$0	\$33,500	\$0	\$33,500
	1507	Spider, Tooth Cave (Tayshaneta myopica) -Wherever found	E	\$0	\$0	\$0	\$4,000	\$4,000
	1667	Tarantula, Sri Lankan Ornamental (Poecilotheria fasciata) -Wherever found	E	\$521	\$0	\$521	\$0	\$521
Arachnids subtotal				\$58,521	\$0	\$58,521	\$4,000	\$62,521
Birds	854	(honeyeater), Mao (= maomao) (Gymnomyza samoensis) -Wherever found	E	\$18,430	\$500	\$18,930	\$0	\$18,930
	50	'I'iwi (Drepanis coccinea) -Wherever found	T	\$414,123	\$2,409,875	\$2,823,998	\$0	\$2,823,998
	987	'O'o, Kauai (honeyeater) (Moho braccatus) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	895	'O'u (honeycreeper) (Psittirostra psittacea) -Wherever found	E	\$16,791	\$0	\$16,791	\$0	\$16,791
	221	Akekee (Loxops caeruleirostris) -Wherever found	E	\$329,895	\$0	\$329,895	\$0	\$329,895
	323	akepa, Hawaii (Loxops coccineus) -Wherever found	E	\$130,177	\$0	\$130,177	\$0	\$130,177
	1002	akepa, Maui (Loxops ochraceus) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	952	akialoa, Kauai (honeycreeper) (Akialoa stejnegeri) -Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	324	akiapolaau (Hemignathus wilsoni) -Wherever found	E	\$129,177	\$0	\$129,177	\$0	\$129,177
	192	Akikiki (Oreomystis bairdi) -Wherever found	E	\$398,560	\$0	\$398,560	\$0	\$398,560
	58	Akohekohe (Palmeria dolei) -Wherever found	E	\$39,500	\$2,474,560	\$2,514,060	\$0	\$2,514,060
	1002	alauahio, Oahu (Paroreomyza maculata) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	223	Albatross, short-tailed (Phoebastria (=Diomedea) albatrus) -Wherever found	E	\$328,491	\$0	\$328,491	\$0	\$328,491
	896	Blackbird, yellow-shouldered (Agelaius xanthomus) -Wherever found	E	\$16,500	\$0	\$16,500	\$0	\$16,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	117	Caracara, Audubon's crested (<i>Polyborus plancus audubonii</i>) -U.S.A. (FL)	T	\$809,500	\$0	\$809,500	\$0	\$809,500
	505	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. (Oregon, and specific portions of northern California and northwest Nevada)	EXPN	\$53,761	\$1,500	\$55,261	\$0	\$55,261
	399	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. (specific portions of Arizona, Nevada, and Utah)-see 17.84(j)	EXPN	\$500	\$92,300	\$92,800	\$0	\$92,800
	119	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. only, except where listed as an experimental population	E	\$281,600	\$512,890	\$794,490	\$0	\$794,490
	241	coot, Hawaiian (<i>Fulica americana alai</i>) - Wherever found	E	\$228,961	\$31,100	\$260,061	\$0	\$260,061
	133	Crane, Mississippi sandhill (<i>Grus canadensis pulla</i>) -Wherever found	E	\$708,000	\$0	\$708,000	\$0	\$708,000
	226	Crane, whooping (<i>Grus americana</i>) -U.S.A (Southwestern Louisiana)	EXPN	\$308,000	\$0	\$308,000	\$0	\$308,000
	1064	Crane, whooping (<i>Grus americana</i>) -U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$11,500	\$0	\$11,500	\$0	\$11,500
	238	Crane, whooping (<i>Grus americana</i>) -U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$261,900	\$0	\$261,900	\$0	\$261,900
	160	Crane, whooping (<i>Grus americana</i>) - Wherever found, except where listed as an experimental population	E	\$214,900	\$156,468	\$371,368	\$158,851	\$530,219
	322	Creeper, Hawaii (<i>Oreomystis mana</i>) - Wherever found	E	\$131,177	\$0	\$131,177	\$0	\$131,177
	1002	Creeper, Molokai (<i>Paroreomyza flammea</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	115	Crow, Hawaiian (=alala) (<i>Corvus hawaiiensis</i>) -Wherever found	E	\$827,695	\$0	\$827,695	\$0	\$827,695
	112	Crow, Mariana (=aga) (<i>Corvus kubaryi</i>) - Wherever found	E	\$845,897	\$0	\$845,897	\$0	\$845,897

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1464	Crow, white-necked (<i>Corvus leucognaphalus</i>) -Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	69	Cuckoo, yellow-billed (<i>Coccyzus americanus</i>) -Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	T	\$500,599	\$1,349,098	\$1,849,697	\$40,000	\$1,889,697
	1039	Curassow, blue-billed (<i>Crax alberti</i>) -Wherever found	E	\$12,062	\$0	\$12,062	\$0	\$12,062
	278	Duck, Hawaiian (=koloa) (<i>Anas wyvilliana</i>) -Wherever found	E	\$182,841	\$2,400	\$185,241	\$0	\$185,241
	351	Duck, Laysan (<i>Anas laysanensis</i>) -Wherever found	E	\$114,251	\$0	\$114,251	\$0	\$114,251
	756	Eagle, harpy (<i>Harpia harpyja</i>) -Wherever found	E	\$24,818	\$0	\$24,818	\$0	\$24,818
	124	Eider, spectacled (<i>Somateria fischeri</i>) -Wherever found	T	\$666,000	\$77,500	\$743,500	\$16,983	\$760,483
	151	Eider, Steller's (<i>Polysticta stelleri</i>) -U.S. Alaska breeding population only	T	\$505,600	\$65,000	\$570,600	\$0	\$570,600
	469	elepaio, Oahu (<i>Chasiempis ibidis</i>) -Wherever found	E	\$62,000	\$0	\$62,000	\$0	\$62,000
	581	Falcon, Eurasian peregrine (<i>Falco peregrinus peregrinus</i>) -Wherever found	E	\$0	\$0	\$0	\$41,559	\$41,559
	699	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) -U.S.A (AZ, NM)	EXPN	\$0	\$29,400	\$29,400	\$0	\$29,400
	523	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) -Wherever found, except where listed as an experimental population	E	\$51,000	\$450	\$51,450	\$0	\$51,450
	486	Finch, Laysan (honeycreeper) (<i>Telespyza cantans</i>) -Wherever found	E	\$58,000	\$0	\$58,000	\$0	\$58,000
	460	Finch, Nihoa (honeycreeper) (<i>Telespyza ultima</i>) -Wherever found	E	\$65,500	\$0	\$65,500	\$0	\$65,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	100	Flycatcher, southwestern willow (Empidonax traillii extimus) -Wherever found	E	\$148,900	\$861,458	\$1,010,358	\$0	\$1,010,358
	250	gallinule, Hawaiian common (Gallinula galeata sandvicensis) -Wherever found	E	\$231,740	\$2,400	\$234,140	\$0	\$234,140
	227	Gnatcatcher, coastal California (Polioptila californica californica) -Wherever found	T	\$304,500	\$0	\$304,500	\$0	\$304,500
	104	Goose, Hawaiian (Branta (=Nesochen) sandvicensis) -Wherever found	T	\$647,671	\$269,639	\$917,310	\$0	\$917,310
	798	Ground-Dove, Friendly (Gallicolumba stairi) -American Samoa DPS	E	\$21,430	\$500	\$21,930	\$0	\$21,930
	809	Hawk, Puerto Rican broad-winged (Buteo platypterus brunescens) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	365	Hawk, Puerto Rican sharp-shinned (Accipiter striatus venator) -Wherever found	E	\$110,000	\$0	\$110,000	\$0	\$110,000
	162	Horned lark, streaked (Eremophila alpestris strigata) -Wherever found	T	\$509,370	\$1,863	\$511,233	\$0	\$511,233
	1663	Ibis, northern bald (Geronticus eremita) -Wherever found	E	\$0	\$600	\$600	\$0	\$600
	231	Kingfisher, Guam Micronesian (Halcyon cinnamomina cinnamomina) -Wherever found	E	\$255,708	\$36,800	\$292,508	\$0	\$292,508
	164	Kite, Everglade snail (Rostrhamus sociabilis plumbeus) -Wherever found	E	\$367,000	\$5,000	\$372,000	\$125,743	\$497,743
	130	Knot, red (Calidris canutus rufa) -Wherever found	T	\$649,400	\$78,602	\$728,002	\$10,438	\$738,440
	1639	Macaw, military (Ara militaris) -Wherever found	E	\$0	\$1,000	\$1,000	\$0	\$1,000
	1539	macaw, Scarlet (Ara macao ssp. cyanopterus) -Wherever found	E	\$3,132	\$0	\$3,132	\$0	\$3,132
	415	Megapode, Micronesian (Megapodius laperouse) -Wherever found	E	\$84,500	\$0	\$84,500	\$0	\$84,500
	441	Millerbird, Nihoa (old world warbler) (Acrocephalus familiaris kingi) -Wherever found	E	\$75,000	\$0	\$75,000	\$0	\$75,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	401	Moorhen, Mariana common (<i>Gallinula chloropus guami</i>) -Wherever found	E	\$91,000	\$0	\$91,000	\$0	\$91,000
	48	Murrelet, marbled (<i>Brachyramphus marmoratus</i>) -U.S.A. (CA, OR, WA)	T	\$1,773,996	\$1,503,126	\$3,277,122	\$0	\$3,277,122
	1292	Nightjar, Puerto Rican (<i>Caprimulgus noctitherus</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1002	nukupuu, Kauai (<i>Hemignathus hanapepe</i>) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	792	nukupuu, Maui (<i>Hemignathus affinis</i>) -Wherever found	E	\$22,000	\$0	\$22,000	\$0	\$22,000
	181	Owl, Mexican spotted (<i>Strix occidentalis lucida</i>) -Wherever found	T	\$19,000	\$421,513	\$440,513	\$0	\$440,513
	19	Owl, northern spotted (<i>Strix occidentalis caurina</i>) -Wherever found	T	\$4,735,423	\$4,326,999	\$9,062,422	\$0	\$9,062,422
	269	Palila (honeycreeper) (<i>Loxioides bailleui</i>) -Wherever found	E	\$200,341	\$0	\$200,341	\$0	\$200,341
	79	Parrot, Puerto Rican (<i>Amazona vittata</i>) -Wherever found	E	\$1,430,000	\$0	\$1,430,000	\$0	\$1,430,000
	55	parrotbill, Maui (<i>Kiwikiu</i>) (<i>Pseudonestor xanthophrys</i>) -Wherever found	E	\$218,975	\$2,474,560	\$2,693,535	\$0	\$2,693,535
	1653	penguin, African (<i>Spheniscus demersus</i>) -Wherever found	E	\$0	\$700	\$700	\$0	\$700
	839	penguin, emperor (<i>Aptenodytes forsteri</i>) -Wherever found	T	\$19,663	\$0	\$19,663	\$0	\$19,663
	1541	petrel, Bermuda (<i>Pterodroma cahow</i>) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	98	petrel, Hawaiian (<i>Pterodroma sandwichensis</i>) -Wherever found	E	\$674,887	\$370,902	\$1,045,789	\$0	\$1,045,789
	610	Pigeon, Puerto Rican plain (<i>Columba inornata wetmorei</i>) -Wherever found	E	\$38,000	\$0	\$38,000	\$0	\$38,000
	66	Plover, piping (<i>Charadrius melodus</i>) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$891,100	\$964,828	\$1,855,928	\$59,176	\$1,915,104

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	93	Plover, piping (Charadrius melodus) -[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$948,500	\$55,308	\$1,003,808	\$115,325	\$1,119,133
	78	Plover, western snowy (Charadrius nivosus nivosus) -Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$1,318,901	\$189,600	\$1,508,501	\$0	\$1,508,501
	792	Po`ouli (honeycreeper) (Melamprosops phaeosoma) -Wherever found	E	\$22,000	\$0	\$22,000	\$0	\$22,000
	848	Prairie-chicken, Attwater's greater (Tympanuchus cupido attwateri) -Wherever found	E	\$19,000	\$0	\$19,000	\$0	\$19,000
	170	Rail, California clapper (Rallus longirostris obsoletus) -Wherever found	E	\$484,400	\$5,702	\$490,102	\$0	\$490,102
	101	rail, Eastern Black (Laterallus jamaicensis ssp. jamaicensis) -Wherever found	T	\$901,500	\$48,862	\$950,362	\$48,600	\$998,962
	724	Rail, Guam (Rallus owstoni) -Rota	EXPN	\$27,000	\$0	\$27,000	\$0	\$27,000
	247	Rail, Guam (Rallus owstoni) -Wherever found, except where listed as an experimental population	E	\$223,779	\$15,900	\$239,679	\$0	\$239,679
	295	Rail, light-footed clapper (Rallus longirostris levipes) -Wherever found	E	\$156,600	\$0	\$156,600	\$0	\$156,600
	176	Rail, Yuma Ridgway's (Rallus obsoletus yumanensis) -Wherever found	E	\$168,900	\$291,431	\$460,331	\$0	\$460,331
	83	sage-grouse, Gunnison (Centrocercus minimus) -Wherever found	T	\$103,000	\$768,661	\$871,661	\$427,782	\$1,299,443
	169	scrub-jay, Florida (Aphelocoma coerulescens) -Wherever found	T	\$291,000	\$1,495	\$292,495	\$197,836	\$490,331
	113	Shearwater, Newell's Townsend's (Puffinus auricularis newelli) -Wherever found	T	\$834,806	\$6,100	\$840,906	\$0	\$840,906
	158	Sparrow, Cape Sable seaside (Ammodramus maritimus mirabilis) -Wherever found	E	\$237,500	\$300,000	\$537,500	\$0	\$537,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	207	Sparrow, Florida grasshopper (Ammodramus savannarum floridanus) - Wherever found	E	\$266,000	\$0	\$266,000	\$106,139	\$372,139
	1569	Sparrow, San Clemente sage (Amphispiza belli clementeae) -Wherever found	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	1639	Starling, Rothschild's (myna) (Leucopsar rothschildi) -Wherever found	E	\$0	\$1,000	\$1,000	\$0	\$1,000
	214	Stilt, Hawaiian (Himantopus mexicanus knudseni) -Wherever found	E	\$328,031	\$31,100	\$359,131	\$0	\$359,131
	209	Stork, wood (Mycteria americana) -U.S.A. (AL, FL, GA, MS, NC, SC)	T	\$327,500	\$43,005	\$370,505	\$300	\$370,805
	258	Storm-petrel, band-rumped (Oceanodroma castro) -USA (HI)	E	\$172,521	\$46,925	\$219,446	\$0	\$219,446
	309	Swiftlet, Mariana gray (Aerodramus vanikorensis bartschi) -Wherever found	E	\$144,700	\$0	\$144,700	\$0	\$144,700
	186	Tern, California least (Sterna antillarum browni) -Wherever found	E	\$415,400	\$5,300	\$420,700	\$1,000	\$421,700
	296	Tern, Roseate (Sterna dougallii dougallii) - U.S.A. (Atlantic Coast south to NC), Canada (Newf., N.S, Que.), Bermuda	E	\$126,000	\$10,000	\$136,000	\$19,900	\$155,900
	696	Tern, Roseate (Sterna dougallii dougallii) - Western Hemisphere and adjacent oceans, incl. U.S.A. (FL, PR, VI), where not listed as endangered	T	\$24,500	\$5,000	\$29,500	\$0	\$29,500
	987	Thrush, large Kauai (=kamao) (Myadestes myadestinus) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	987	Thrush, Molokai (Myadestes lanaiensis rutha) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	313	Thrush, small Kauai (=puaiohi) (Myadestes palmeri) -Wherever found	E	\$140,219	\$0	\$140,219	\$0	\$140,219
	350	Towhee, Inyo California (Pipilo crissalis eremophilus) -Wherever found	T	\$115,100	\$0	\$115,100	\$0	\$115,100
	182	Vireo, least Bell's (Vireo bellii pusillus) - Wherever found	E	\$366,200	\$71,148	\$437,348	\$0	\$437,348
	1541	Warbler (=wood), Bachman's (Vermivora bachmanii) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1541	Warbler, elfin-woods (<i>Setophaga angelae</i>) - Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	319	Warbler, nightingale reed (old world warbler) (<i>Acrocephalus luscini</i>) -Wherever found	E	\$133,000	\$0	\$133,000	\$0	\$133,000
	1402	White-eye, Bridled (<i>Zosterops conspicillatus</i>) -Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	809	White-eye, Rota bridled (<i>Zosterops rotensis</i>) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	553	Woodpecker, ivory-billed (<i>Campephilus principalis</i>) -Wherever found	E	\$45,500	\$0	\$45,500	\$0	\$45,500
	63	Woodpecker, red-cockaded (<i>Picoides borealis</i>) -Wherever found	E	\$1,727,500	\$128,335	\$1,855,835	\$393,194	\$2,249,029
Birds subtotal				\$32,520,999	\$20,548,403	\$53,069,402	\$1,762,826	\$54,832,228
Clams	907	Acornshell, southern (<i>Epioblasma othcaloogensis</i>) -Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	271	Bankclimber, purple (mussel) (<i>Elliptoideus sloatianus</i>) -Wherever found	T	\$194,000	\$0	\$194,000	\$3,493	\$197,493
	1285	Bean, Choctaw (<i>Obovaria choctawensis</i>) - Wherever found	E	\$5,500	\$0	\$5,500	\$3,493	\$8,993
	1608	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	328	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) -Wherever found; Except where listed as Experimental Populations	E	\$122,500	\$2,450	\$124,950	\$0	\$124,950
	821	Bean, purple (<i>Villosa perpurpurea</i>) - Wherever found	E	\$20,500	\$0	\$20,500	\$0	\$20,500
	681	Bean, rayed (<i>Villosa fabalis</i>) -Wherever found	E	\$28,500	\$1,635	\$30,135	\$0	\$30,135

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1541	Blossom, green (pearlymussel) (<i>Epioblasma torulosa</i> gubernaculum) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1608	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	1026	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) -Wherever found; Except where listed as Experimental Populations	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1338	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$8,000	\$0	\$8,000	\$0	\$8,000
	1111	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) -Wherever found; Except where listed as Experimental Populations	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1527	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$3,500	\$0	\$3,500	\$0	\$3,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1541	Blossom, yellow (pearlymussel) (Epioblasma florentina florentina) - Wherever found; Except where listed as Experimental Populations	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1361	Catspaw, white (pearlymussel) (Epioblasma perobliqua) -Wherever found	E	\$7,600	\$0	\$7,600	\$0	\$7,600
	1608	Clubshell (Pleurobema clava) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	434	Clubshell (Pleurobema clava) -Wherever found; Except where listed as Experimental Populations	E	\$75,100	\$1,787	\$76,887	\$300	\$77,187
	1624	Clubshell, black (Pleurobema curtum) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	1495	Clubshell, Canoe Creek (Pleurobema athearni) -Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	786	Clubshell, ovate (Pleurobema perovatum) - Wherever found	E	\$22,500	\$0	\$22,500	\$0	\$22,500
	525	Clubshell, southern (Pleurobema decisum) - Wherever found	E	\$51,000	\$0	\$51,000	\$0	\$51,000
	1152	Combshell, Cumberlandian (Epioblasma brevidens) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$10,000	\$0	\$10,000	\$0	\$10,000
	522	Combshell, Cumberlandian (Epioblasma brevidens) -Wherever found; Except where listed as Experimental Populations	E	\$49,000	\$3,085	\$52,085	\$0	\$52,085

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1416	Combshell, southern (<i>Epioblasma penita</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	919	Combshell, upland (<i>Epioblasma metastriata</i>) -Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	565	Ebonyshell, round (<i>Reginaia rotulata</i>) - Wherever found	E	\$41,000	\$0	\$41,000	\$3,493	\$44,493
	325	Elktoe, Appalachian (<i>Alasmodonta raveneliana</i>) -Wherever found	E	\$119,000	\$0	\$119,000	\$8,000	\$127,000
	611	Elktoe, Cumberland (<i>Alasmodonta atropurpurea</i>) -Wherever found	E	\$35,500	\$2,450	\$37,950	\$0	\$37,950
	1024	Fanshell (<i>Cyprogenia stegaria</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$12,700	\$0	\$12,700	\$0	\$12,700
	471	Fanshell (<i>Cyprogenia stegaria</i>) -Wherever found	E	\$60,000	\$1,787	\$61,787	\$0	\$61,787
	809	fatmucket, Arkansas (<i>Lampsilis powellii</i>) - Wherever found	T	\$9,000	\$0	\$9,000	\$12,000	\$21,000
	86	Heelsplitter, Carolina (<i>Lasmigona decorata</i>) - Wherever found	E	\$1,063,000	\$0	\$1,063,000	\$174,300	\$1,237,300
	537	heelsplitter, Inflated (<i>Potamilus inflatus</i>) - Wherever found	T	\$48,500	\$0	\$48,500	\$0	\$48,500
	267	Higgins eye (pearlymussel) (<i>Lampsilis higginsii</i>) -Wherever found	E	\$148,600	\$2,076	\$150,676	\$51,105	\$201,781
	696	Hornshell, Texas (<i>Popenaias popeii</i>) - Wherever found	E	\$20,000	\$9,500	\$29,500	\$0	\$29,500
	371	Kidneyshell, fluted (<i>Ptychobranhus subtentus</i>) -Wherever found	E	\$105,000	\$3,085	\$108,085	\$0	\$108,085
	1357	Kidneyshell, southern (<i>Ptychobranhus jonesi</i>) -Wherever found	E	\$4,500	\$0	\$4,500	\$3,493	\$7,993
	619	Kidneyshell, triangular (<i>Ptychobranhus greenii</i>) -Wherever found	E	\$37,000	\$0	\$37,000	\$0	\$37,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	533	Lampmussel, Alabama (<i>Lampsilis virescens</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$49,000	\$0	\$49,000	\$0	\$49,000
	1037	Lampmussel, Alabama (<i>Lampsilis virescens</i>) -Wherever found; Except where listed as Experimental Populations	E	\$11,500	\$635	\$12,135	\$0	\$12,135
	297	Lance, yellow (<i>Elliptio lanceolata</i>) - Wherever found	T	\$137,000	\$0	\$137,000	\$18,500	\$155,500
	613	Lilliput, pale (pearlymussel) (<i>Toxolasma cylindrellus</i>) -Wherever found	E	\$37,000	\$635	\$37,635	\$0	\$37,635
	1290	Mapleleaf, winged (<i>Quadrula fragosa</i>) - U.S.A. (AL-specified portions of the Tennessee River; see 17.85(a)(1))	EXPN	\$8,500	\$76	\$8,576	\$0	\$8,576
	597	Mapleleaf, winged (<i>Quadrula fragosa</i>) - Wherever found, except where listed as an experimental population	E	\$29,100	\$9,300	\$38,400	\$800	\$39,200
	667	Moccasinshell, Alabama (<i>Medionidus acutissimus</i>) -Wherever found	T	\$28,000	\$0	\$28,000	\$3,493	\$31,493
	654	Moccasinshell, Coosa (<i>Medionidus parvulus</i>) -Wherever found	E	\$33,000	\$0	\$33,000	\$0	\$33,000
	780	Moccasinshell, Gulf (<i>Medionidus penicillatus</i>) -Wherever found	E	\$19,500	\$0	\$19,500	\$3,493	\$22,993
	772	Moccasinshell, Ochlockonee (<i>Medionidus simpsonianus</i>) -Wherever found	E	\$20,500	\$0	\$20,500	\$3,493	\$23,993
	617	Moccasinshell, Suwannee (<i>Medionidus walkeri</i>) -Wherever found	T	\$34,000	\$0	\$34,000	\$3,493	\$37,493
	878	Monkeyface, Appalachian (pearlymussel) (<i>Theliderma sparsa</i>) -Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1608	Monkeyface, Cumberland (pearlymussel) (Theliderma intermedia) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	513	Monkeyface, Cumberland (pearlymussel) (Theliderma intermedia) -Wherever found; Except where listed as Experimental Populations	E	\$53,000	\$635	\$53,635	\$0	\$53,635
	482	Mucket, Neosho (Lampsilis rafinesqueana) - Wherever found	E	\$58,800	\$0	\$58,800	\$0	\$58,800
	1541	Mucket, oranenacre (Hamiota perovalis) - Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	256	Mucket, pink (pearlymussel) (Lampsilis abrupta) -Wherever found	E	\$220,500	\$4,237	\$224,737	\$0	\$224,737
	1608	Mussel, oyster (Epioblasma capsaeformis) - U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	382	Mussel, oyster (Epioblasma capsaeformis) - Wherever found; Except where listed as Experimental Populations	E	\$99,500	\$2,450	\$101,950	\$0	\$101,950
	414	Mussel, scaleshell (Leptodea leptodon) - Wherever found	E	\$84,000	\$635	\$84,635	\$0	\$84,635
	265	Mussel, sheepnose (Plethobasus cyphus) - Wherever found	E	\$188,200	\$1,863	\$190,063	\$13,000	\$203,063
	163	Mussel, snuffbox (Epioblasma triquetra) - Wherever found	E	\$480,900	\$18,339	\$499,239	\$4,300	\$503,539
	542	Pearlshell, Alabama (Margaritifera marrianae) -Wherever found	E	\$47,000	\$0	\$47,000	\$0	\$47,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	149	Pearlshell, Louisiana (<i>Margaritifera hembeli</i>) -Wherever found	T	\$572,000	\$0	\$572,000	\$0	\$572,000
	1608	Pearllymussel, birdwing (<i>Lemiox rimosus</i>) - U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	791	Pearllymussel, birdwing (<i>Lemiox rimosus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$21,500	\$635	\$22,135	\$0	\$22,135
	1608	Pearllymussel, cracking (<i>Hemistena lata</i>) - U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	1608	Pearllymussel, cracking (<i>Hemistena lata</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	774	Pearllymussel, cracking (<i>Hemistena lata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$23,000	\$635	\$23,635	\$0	\$23,635
	838	Pearllymussel, Curtis (<i>Epioblasma florentina curtisii</i>) -Wherever found	E	\$19,700	\$0	\$19,700	\$0	\$19,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1608	Pearl mussel, dromedary (<i>Dromus dromas</i>) - U.S.A. (AL; The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	532	Pearl mussel, dromedary (<i>Dromus dromas</i>) - Wherever found; Except where listed as Experimental Populations	E	\$46,000	\$3,085	\$49,085	\$0	\$49,085
	679	Pearl mussel, littlewing (<i>Pegias fabula</i>) - Wherever found	E	\$28,000	\$2,450	\$30,450	\$0	\$30,450
	438	Pearl mussel, slabside (<i>Pleuroaia dolabelloides</i>) - Wherever found	E	\$75,500	\$635	\$76,135	\$0	\$76,135
	465	Pigtoe, Atlantic (<i>Fusconaia masoni</i>) - Wherever found	T	\$41,500	\$3,200	\$44,700	\$18,500	\$63,200
	1002	Pigtoe, Cumberland (<i>Pleuroaia gibber</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1495	Pigtoe, dark (<i>Pleurobema furvum</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1608	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) - U.S.A. (AL; The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	1663	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$600	\$600	\$0	\$600
	506	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$54,500	\$635	\$55,135	\$0	\$55,135

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1569	Pigtoe, flat (<i>Pleurobema marshalli</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1285	Pigtoe, fuzzy (<i>Pleurobema strodeanum</i>) - Wherever found	T	\$5,500	\$0	\$5,500	\$3,493	\$8,993
	1041	Pigtoe, Georgia (<i>Pleurobema hanleyianum</i>) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	634	Pigtoe, heavy (<i>Pleurobema taitianum</i>) - Wherever found	E	\$35,500	\$0	\$35,500	\$0	\$35,500
	1126	Pigtoe, narrow (<i>Fusconaia escambia</i>) - Wherever found	T	\$7,500	\$0	\$7,500	\$3,493	\$10,993
	470	Pigtoe, oval (<i>Pleurobema pyriforme</i>) - Wherever found	E	\$58,500	\$0	\$58,500	\$3,493	\$61,993
	513	Pigtoe, rough (<i>Pleurobema plenum</i>) - Wherever found	E	\$52,000	\$1,635	\$53,635	\$0	\$53,635
	1588	Pigtoe, shiny (<i>Fusconaia cor</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$600	\$2,300	\$0	\$2,300
	559	Pigtoe, shiny (<i>Fusconaia cor</i>) -Wherever found; Except where listed as Experimental Populations	E	\$44,500	\$635	\$45,135	\$0	\$45,135
	645	Pigtoe, southern (<i>Pleurobema georgianum</i>) - Wherever found	E	\$34,500	\$0	\$34,500	\$0	\$34,500
	1148	Pigtoe, tapered (<i>Fusconaia burkei</i>) - Wherever found	T	\$7,000	\$0	\$7,000	\$3,493	\$10,493
	443	Pimpleback, orangefoot (pearlymussel) (<i>Plethobasus cooperianus</i>) -Wherever found	E	\$73,000	\$635	\$73,635	\$0	\$73,635
	546	Pocketbook, fat (<i>Potamilus capax</i>) - Wherever found	E	\$46,400	\$0	\$46,400	\$0	\$46,400
	727	Pocketbook, finelined (<i>Hamiota altilis</i>) - Wherever found	T	\$26,500	\$0	\$26,500	\$0	\$26,500
	749	Pocketbook, Ouachita rock (<i>Arcidens wheeleri</i>) -Wherever found	E	\$15,000	\$0	\$15,000	\$10,000	\$25,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	678	Pocketbook, shinyrayed (<i>Hamiota subangulata</i>) -Wherever found	E	\$27,000	\$0	\$27,000	\$3,493	\$30,493
	629	Pocketbook, speckled (<i>Lampsilis streckeri</i>) -Wherever found	E	\$30,500	\$0	\$30,500	\$5,500	\$36,000
	1608	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata</i>) - U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$0	\$1,700	\$0	\$1,700
	497	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$56,500	\$152	\$56,652	\$0	\$56,652
	229	Rabbitsfoot (<i>Quadrula cylindrica cylindrica</i>) -Wherever found	T	\$267,100	\$16,965	\$284,065	\$15,300	\$299,365
	749	Rabbitsfoot, rough (<i>Quadrula cylindrica strigillata</i>) -Wherever found	E	\$25,000	\$0	\$25,000	\$0	\$25,000
	588	Riffleshell, northern (<i>Epioblasma rangiana</i>) -Wherever found	E	\$38,500	\$2,152	\$40,652	\$300	\$40,952
	449	Riffleshell, tan (<i>Epioblasma florentina walkeri</i> (=E. walkeri)) -Wherever found	E	\$66,000	\$2,450	\$68,450	\$0	\$68,450
	598	Ring pink (mussel) (<i>Obovaria retusa</i>) -Wherever found	E	\$37,500	\$1,635	\$39,135	\$0	\$39,135
	1148	Sandshell, Southern (<i>Hamiota australis</i>) -Wherever found	T	\$7,000	\$0	\$7,000	\$3,493	\$10,493
	1060	Slabshell, Chipola (<i>Elliptio chipolaensis</i>) -Wherever found	T	\$8,500	\$0	\$8,500	\$3,493	\$11,993
	262	Spectaclecase (mussel) (<i>Cumberlandia monodonta</i>) -Wherever found	E	\$205,100	\$6,161	\$211,261	\$1,100	\$212,361
	952	Spinymussel, Altamaha (<i>Elliptio spinosa</i>) -Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	187	Spinymussel, James (<i>Parvaspina collina</i>) -Wherever found	E	\$419,500	\$0	\$419,500	\$1,600	\$421,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	426	Spinymussel, Tar River (Parvaspina steinstansana) -Wherever found	E	\$57,500	\$0	\$57,500	\$23,500	\$81,000
	1569	Stirrupshell (Quadrula stapes) -Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	507	Threeridge, fat (mussel) (Amblema neislerii) -Wherever found	E	\$51,500	\$0	\$51,500	\$3,493	\$54,993
	1662	Wartyback, white (pearlymussel) (Plethobasus cicatricosus) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$635	\$635	\$0	\$635
	868	Wartyback, white (pearlymussel) (Plethobasus cicatricosus) -Wherever found	E	\$18,000	\$0	\$18,000	\$0	\$18,000
	168	Wedgemussel, dwarf (Alasmidonta heterodon) -Wherever found	E	\$464,000	\$3,200	\$467,200	\$23,300	\$490,500
	1680	No Common Name (Fusconaia subrotunda) -	T	\$0	\$100	\$100	\$0	\$100
	1680	No Common Name (Obovaria subrotunda) -	T	\$0	\$100	\$100	\$0	\$100
Clams subtotal				\$7,170,700	\$115,590	\$7,286,290	\$437,293	\$7,723,583
Conifers and Cycads	1416	Cypress, Gowen (Cupressus goveniana ssp. goveniana) -Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	1021	Cypress, Santa Cruz (Cupressus abramsiana) -Wherever found	T	\$12,900	\$0	\$12,900	\$0	\$12,900
	362	Fadang (Cycas micronesica) -Wherever found	T	\$111,657	\$0	\$111,657	\$0	\$111,657
	1647	Torreya, Florida (Torreya taxifolia) -	E	\$900	\$0	\$900	\$0	\$900
	146	No Common Name (Pinus albicaulis) -	T	\$0	\$602,692	\$602,692	\$0	\$602,692
Conifers and Cycads subtotal				\$131,957	\$602,692	\$734,649	\$0	\$734,649
Corals	131	coral, Boulder star (Orbicella franksi) -Wherever found	T	\$0	\$731,200	\$731,200	\$0	\$731,200
	110	Coral, elkhorn (Acropora palmata) -Wherever found	T	\$0	\$858,000	\$858,000	\$0	\$858,000
	122	Coral, Lobed Star (Orbicella annularis) -Wherever found	T	\$0	\$780,700	\$780,700	\$0	\$780,700
	114	Coral, Mountainous Star (Orbicella faveolata) -Wherever found	T	\$0	\$838,700	\$838,700	\$0	\$838,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	127	coral, Pillar (<i>Dendrogyra cylindrus</i>) - Wherever found	T	\$0	\$738,700	\$738,700	\$0	\$738,700
	140	Coral, Rough Cactus (<i>Mycetophyllia ferox</i>) - Wherever found	T	\$0	\$656,200	\$656,200	\$0	\$656,200
	111	Coral, staghorn (<i>Acropora cervicornis</i>) - Wherever found	T	\$0	\$851,700	\$851,700	\$0	\$851,700
	200	No common name (<i>Acropora globiceps</i>) - Wherever found	T	\$0	\$378,500	\$378,500	\$0	\$378,500
	203	No common name (<i>Acropora jacquelinae</i>) - Wherever found	T	\$0	\$377,500	\$377,500	\$0	\$377,500
	201	No common name (<i>Acropora retusa</i>) - Wherever found	T	\$0	\$378,000	\$378,000	\$0	\$378,000
	203	No common name (<i>Acropora speciosa</i>) - Wherever found	T	\$0	\$377,500	\$377,500	\$0	\$377,500
	203	No common name (<i>Euphyllia paradivisa</i>) - Wherever found	T	\$0	\$377,500	\$377,500	\$0	\$377,500
	201	No common name (<i>Isopora crateriformis</i>) - Wherever found	T	\$0	\$378,000	\$378,000	\$0	\$378,000
	194	No common name (<i>Seriatopora aculeata</i>) - Wherever found	T	\$0	\$393,000	\$393,000	\$0	\$393,000
Corals subtotal				\$0	\$8,115,200	\$8,115,200	\$0	\$8,115,200
Crustaceans	614	Amphipod, Hay's Spring (<i>Stygobromus hayi</i>) -Wherever found	E	\$37,500	\$0	\$37,500	\$0	\$37,500
	1152	Amphipod, Illinois cave (<i>Gammarus acherondytes</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	760	Amphipod, Kauai cave (<i>Spelaeorchestia koloana</i>) -Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	1591	Amphipod, Noel's (<i>Gammarus desperatus</i>) - Wherever found	E	\$0	\$2,000	\$2,000	\$0	\$2,000
	809	crayfish, Benton County cave (<i>Cambarus aculabrum</i>) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	266	Crayfish, Big Sandy (<i>Cambarus callainus</i>) - Wherever found	T	\$202,000	\$0	\$202,000	\$0	\$202,000
	519	Crayfish, Guyandotte River (<i>Cambarus veteranus</i>) -Wherever found	E	\$51,000	\$0	\$51,000	\$2,187	\$53,187
	715	Crayfish, Hell Creek Cave (<i>Cambarus zophonastes</i>) -Wherever found	E	\$28,000	\$0	\$28,000	\$0	\$28,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	573	Crayfish, Nashville (Orconectes shoupi) - Wherever found	E	\$33,000	\$7,000	\$40,000	\$3,000	\$43,000
	316	Crayfish, Panama City (Procambarus econfinae) -Wherever found	T	\$72,500	\$0	\$72,500	\$64,627	\$137,127
	259	Crayfish, Shasta (Pacifastacus fortis) - Wherever found	E	\$210,000	\$5,601	\$215,601	\$0	\$215,601
	859	crayfish, Slenderclaw (Cambarus cracens) - Wherever found	E	\$18,500	\$0	\$18,500	\$0	\$18,500
	301	Fairy shrimp, Conservancy (Branchinecta conservatio) -Wherever found	E	\$143,100	\$8,436	\$151,536	\$0	\$151,536
	705	Fairy shrimp, longhorn (Branchinecta longiantenna) -Wherever found	E	\$22,400	\$6,546	\$28,946	\$0	\$28,946
	1493	Fairy shrimp, Riverside (Streptocephalus woottoni) -Wherever found	E	\$4,800	\$0	\$4,800	\$0	\$4,800
	1527	Fairy shrimp, San Diego (Branchinecta sandiegonensis) -Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	137	Fairy shrimp, vernal pool (Branchinecta lynchi) -Wherever found	T	\$648,087	\$38,001	\$686,088	\$0	\$686,088
	792	Isopod, Lee County cave (Lirceus usdagalun) -Wherever found	E	\$22,000	\$0	\$22,000	\$0	\$22,000
	935	Isopod, Madison Cave (Antrolana lira) - Wherever found	T	\$15,000	\$0	\$15,000	\$0	\$15,000
	1639	Isopod, Socorro (Thermosphaeroma thermophilus) -Wherever found	E	\$0	\$1,000	\$1,000	\$0	\$1,000
	1416	Shrimp, Alabama cave (Palaemonias alabamae) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	707	Shrimp, anchialine pool (Procaris hawaiana) -Wherever found	E	\$21,500	\$7,204	\$28,704	\$0	\$28,704
	708	Shrimp, anchialine pool (Vetericaris chaceorum) -Wherever found	E	\$28,500	\$0	\$28,500	\$0	\$28,500
	1494	Shrimp, California freshwater (Syncaris pacificica) -Wherever found	E	\$4,600	\$0	\$4,600	\$0	\$4,600
	1064	Shrimp, Kentucky cave (Palaemonias ganteri) -Wherever found	E	\$10,500	\$1,000	\$11,500	\$0	\$11,500
	1637	Shrimp, Squirrel Chimney Cave (Palaemonetes cummingi) -Wherever found	T	\$1,100	\$0	\$1,100	\$0	\$1,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	147	Tadpole shrimp, vernal pool (<i>Lepidurus packardii</i>) -Wherever found	E	\$547,300	\$36,401	\$583,701	\$0	\$583,701
Crustaceans subtotal				\$2,186,887	\$113,189	\$2,300,076	\$69,814	\$2,369,890
Ferns and Allies	1026	aumakua, Palapalai (<i>Dryopteris crinalis</i> var. <i>podosorus</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1132	diellia, Asplenium-leaved (<i>Asplenium dielirectum</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1680	Fern, Alabama streak-sorus (<i>Thelypteris pilosa</i> var. <i>alabamensis</i>) -Wherever found	T	\$100	\$0	\$100	\$0	\$100
	1292	Fern, Aleutian shield (<i>Polystichum aleuticum</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	768	Fern, American hart's-tongue (<i>Asplenium scolopendrium</i> var. <i>americanum</i>) -Wherever found	T	\$24,000	\$0	\$24,000	\$0	\$24,000
	1527	Fern, Elfin tree (<i>Cyathea dryopteroides</i>) -Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1152	Fern, pendant kihi (<i>Adenophorus periens</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	809	Hohiu (<i>Dryopteris glabra</i> var. <i>pusilla</i>) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	1132	Ihi`ihi (<i>Marsilea villosa</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	672	Kupukupu makalii (<i>Cyclosorus boydiae</i>) -Wherever found	E	\$20,500	\$10,241	\$30,741	\$0	\$30,741
	1292	No common name (<i>Asplenium dielfalcatum</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1041	No common name (<i>Asplenium diellaciniatum</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	952	No common name (<i>Asplenium dielmannii</i>) -Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	1064	No common name (<i>Asplenium dielpallidum</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	608	No common name (<i>Asplenium peruvianum</i> var. <i>insulare</i>) -Wherever found	E	\$27,930	\$10,241	\$38,171	\$0	\$38,171
	1416	No common name (<i>Asplenium unisorum</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	967	No common name (<i>Deparia kaalaana</i>) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	809	No common name (Diplazium molokaiense) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	1002	No common name (Doryopteris angelica) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1416	No common name (Doryopteris takeuchii) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1464	No common name (Elaphoglossum serpens) -Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	987	No common name (Huperzia stemmermanniae) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	695	No common name (Microlepia strigosa var. mauiensis) -Wherever found	E	\$19,500	\$10,241	\$29,741	\$0	\$29,741
	1451	No common name (Polystichum calderonense) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	799	No common name (Pteris lidgatei) -Wherever found	E	\$21,500	\$0	\$21,500	\$0	\$21,500
	1451	No common name (Tectaria estremerana) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1541	No common name (Thelypteris inabonensis) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1541	No common name (Thelypteris verecunda) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1541	No common name (Thelypteris yaucoensis) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1152	olua (Hypolepis hawaiiensis var. mauiensis) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	768	Pauoa (Ctenitis squamigera) -Wherever found	E	\$24,000	\$0	\$24,000	\$0	\$24,000
	1473	Quillwort, black spored (Isoetes melanospora) -Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	840	Quillwort, Louisiana (Isoetes louisianensis) -Wherever found	E	\$19,500	\$0	\$19,500	\$0	\$19,500
	1507	Quillwort, mat-forming (Isoetes tegetiformans) -Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	887	Wawae'iole (Huperzia mannii) -Wherever found	E	\$7,000	\$10,241	\$17,241	\$0	\$17,241

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1292	Wawae`iole (Huperzia nutans) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	689	fern, Florida bristle (Trichomanes punctatum ssp. floridanum) -	E	\$30,000	\$0	\$30,000	\$0	\$30,000
Ferns and Allies subtotal				\$445,530	\$40,964	\$486,494	\$0	\$486,494
Fishes	153	Bocaccio (Sebastes paucispinis) -Puget Sound-Georgia Basin DPS See 50 CFR 224.101	E	\$0	\$559,000	\$559,000	\$0	\$559,000
	288	Bonytail (Gila elegans) -Wherever found	E	\$22,800	\$142,564	\$165,364	\$0	\$165,364
	1064	Cavefish, Alabama (Speoplatyrhinus poulsoni) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	808	Cavefish, Ozark (Amblyopsis rosae) -Wherever found	T	\$15,500	\$0	\$15,500	\$5,600	\$21,100
	1639	Chub, Chihuahua (Gila nigrescens) -Wherever found	T	\$0	\$1,000	\$1,000	\$0	\$1,000
	874	Chub, Gila (Gila intermedia) -Wherever found	E	\$0	\$17,924	\$17,924	\$0	\$17,924
	132	Chub, humpback (Gila cypha) -Wherever found	T	\$23,600	\$700,691	\$724,291	\$0	\$724,291
	986	Chub, Hutton tui (Gila bicolor ssp.) -Wherever found	T	\$13,600	\$0	\$13,600	\$0	\$13,600
	1330	Chub, Owens Tui (Gila bicolor ssp. snyderi) -Wherever found	E	\$8,400	\$0	\$8,400	\$0	\$8,400
	490	Chub, Pahrnagat roundtail (Gila robusta jordani) -Wherever found	E	\$56,700	\$468	\$57,168	\$0	\$57,168
	864	chub, Peppered (Macrhybopsis tetranema) -Wherever found	E	\$13,300	\$0	\$13,300	\$5,000	\$18,300
	1653	Chub, slender (Erimystax cahni) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(s)(1)(i))	EXPN	\$700	\$0	\$700	\$0	\$700
	868	Chub, slender (Erimystax cahni) -Wherever found	T	\$18,000	\$0	\$18,000	\$0	\$18,000
	1569	Chub, spotfin (Erimonax monachus) -U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$2,500	\$0	\$2,500	\$0	\$2,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1653	Chub, spotfin (Erimonax monachus) -U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$700	\$0	\$700	\$0	\$700
	1653	Chub, spotfin (Erimonax monachus) -U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$700	\$0	\$700	\$0	\$700
	383	Chub, spotfin (Erimonax monachus) - Wherever found, except where listed as an experimental population	T	\$94,000	\$5,235	\$99,235	\$1,100	\$100,335
	260	Chub, Virgin River (Gila seminuda (=robusta)) -Wherever found	E	\$197,600	\$17,573	\$215,173	\$0	\$215,173
	116	Cui-ui (Chasmistes cujus) -Wherever found	E	\$809,700	\$0	\$809,700	\$0	\$809,700
	967	Dace, Ash Meadows speckled (Rhinichthys osculus nevadensis) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	436	Dace, blackside (Phoxinus cumberlandensis) -Wherever found	T	\$77,000	\$0	\$77,000	\$0	\$77,000
	375	Dace, Clover Valley speckled (Rhinichthys osculus oligoporus) -Wherever found	E	\$104,700	\$0	\$104,700	\$0	\$104,700
	283	Dace, desert (Eremichthys acros) -Wherever found	T	\$27,800	\$150,000	\$177,800	\$0	\$177,800
	1471	Dace, Independence Valley speckled (Rhinichthys osculus lethoporus) -Wherever found	E	\$5,200	\$0	\$5,200	\$0	\$5,200
	907	Dace, Kendall Warm Springs (Rhinichthys osculus thermalis) -Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	1451	dace, Laurel (Chrosomus saylori) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	379	Dace, Moapa (Moapa coriacea) -Wherever found	E	\$73,800	\$30,000	\$103,800	\$0	\$103,800
	828	Darter, amber (Percina antesella) -Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	702	Darter, bayou (Etheostoma rubrum) - Wherever found	T	\$4,000	\$0	\$4,000	\$25,000	\$29,000
	868	darter, bluemask (Etheostoma akatulo) - Wherever found	E	\$18,000	\$0	\$18,000	\$0	\$18,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1569	Darter, boulder (<i>Etheostoma wapiti</i>) -Shoal Creek (from Shoal Creek mile 41.7 (66.7 km)) at the mouth of Long Branch, Lawrence County, TN, downstream to the backwaters of Wilson Reservoir (Shoal Creek mile 14 (22 km)) at Goose Shoals, Lauderdale County, AL, including the lower 5 miles (8 km) of all tributaries that enter this reach.	EXPN	\$2,500	\$0	\$2,500	\$0	\$2,500
	159	Darter, boulder (<i>Etheostoma wapiti</i>) - Wherever found	E	\$532,000	\$0	\$532,000	\$0	\$532,000
	279	Darter, candy (<i>Etheostoma osburni</i>) - Wherever found	E	\$155,000	\$1,500	\$156,500	\$28,571	\$185,071
	809	Darter, Cherokee (<i>Etheostoma scotti</i>) - Wherever found	T	\$21,000	\$0	\$21,000	\$0	\$21,000
	1292	Darter, Cumberland (<i>Etheostoma susanae</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	596	Darter, diamond (<i>Crystallaria cincotta</i>) - Wherever found	E	\$38,500	\$200	\$38,700	\$651	\$39,351
	1653	Darter, duskytail (<i>Etheostoma percnurum</i>) - The Tellico River, between the backwaters of the Tellico Reservoir (approximately Tellico River mile 19 (30.4 kilometers) and Tellico River mile 33 (52.8 kilometers), near the Tellico Ranger Station, Monroe County, Tennessee.	EXPN	\$700	\$0	\$700	\$0	\$700
	1653	Darter, duskytail (<i>Etheostoma percnurum</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(q)(1)(ii))	EXPN	\$700	\$0	\$700	\$0	\$700
	435	Darter, duskytail (<i>Etheostoma percnurum</i>) - Wherever found	E	\$65,000	\$12,012	\$77,012	\$0	\$77,012
	476	Darter, Etowah (<i>Etheostoma etowahae</i>) - Wherever found	E	\$60,000	\$0	\$60,000	\$0	\$60,000
	715	Darter, goldline (<i>Percina aurolineata</i>) - Wherever found	T	\$28,000	\$0	\$28,000	\$0	\$28,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	427	Darter, Kentucky arrow (<i>Etheostoma spilotum</i>) - Wherever found	T	\$80,500	\$0	\$80,500	\$0	\$80,500
	782	Darter, leopard (<i>Percina pantherina</i>) - Wherever found	T	\$11,500	\$0	\$11,500	\$11,400	\$22,900
	341	Darter, Niangua (<i>Etheostoma nianguae</i>) - Wherever found	T	\$9,000	\$0	\$9,000	\$110,000	\$119,000
	689	Darter, Okaloosa (<i>Etheostoma okaloosae</i>) - Wherever found	T	\$30,000	\$0	\$30,000	\$0	\$30,000
	261	Darter, Pearl (<i>Percina aurora</i>) - Wherever found	T	\$213,500	\$0	\$213,500	\$0	\$213,500
	848	Darter, relict (<i>Etheostoma chienense</i>) - Wherever found	E	\$19,000	\$0	\$19,000	\$0	\$19,000
	987	Darter, rush (<i>Etheostoma phytophilum</i>) - Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	760	darter, Sickie (<i>Percina williamsi</i>) - Wherever found	T	\$24,500	\$0	\$24,500	\$0	\$24,500
	557	Darter, slackwater (<i>Etheostoma boschungii</i>) - Wherever found	T	\$45,000	\$360	\$45,360	\$0	\$45,360
	220	Darter, trispot (<i>Etheostoma trisella</i>) - Wherever found	T	\$333,000	\$0	\$333,000	\$0	\$333,000
	868	Darter, vermilion (<i>Etheostoma chermocki</i>) - Wherever found	E	\$18,000	\$0	\$18,000	\$0	\$18,000
	878	Darter, watercress (<i>Etheostoma nuchale</i>) - Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500
	1525	Darter, yellowcheek (<i>Etheostoma moorei</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$1,300	\$3,800
	153	Eulachon (<i>Thaleichthys pacificus</i>) - Southern DPS - See 50 CFR 223.102	T	\$0	\$559,000	\$559,000	\$0	\$559,000
	1591	Gambusia, Big Bend (<i>Gambusia gaigei</i>) - Wherever found	E	\$0	\$2,000	\$2,000	\$0	\$2,000
	1541	Gambusia, Pecos (<i>Gambusia nobilis</i>) - Wherever found	E	\$0	\$3,000	\$3,000	\$0	\$3,000
	178	Goby, tidewater (<i>Eucyclogobius newberryi</i>) - Wherever found	E	\$442,800	\$16,000	\$458,800	\$0	\$458,800
	935	Logperch, Conasauga (<i>Percina jenkinsi</i>) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	189	Logperch, Roanoke (<i>Percina rex</i>) -Wherever found	E	\$400,500	\$0	\$400,500	\$4,000	\$404,500
	580	Madtom, Carolina (<i>Noturus furiosus</i>) - Wherever found	E	\$18,500	\$3,200	\$21,700	\$20,000	\$41,700
	484	Madtom, chunky (<i>Noturus crypticus</i>) - Wherever found	E	\$58,500	\$0	\$58,500	\$0	\$58,500
	694	Madtom, Neosho (<i>Noturus placidus</i>) - Wherever found	T	\$29,900	\$0	\$29,900	\$0	\$29,900
	1653	Madtom, pygmy (<i>Noturus stanauli</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))	EXPN	\$700	\$0	\$700	\$0	\$700
	896	Madtom, pygmy (<i>Noturus stanauli</i>) - Wherever found	E	\$16,500	\$0	\$16,500	\$0	\$16,500
	1602	Madtom, smoky (<i>Noturus baileyi</i>) -The Tellico River, between the backwaters of the Tellico Reservoir (approximately Tellico River mile 19 (30.4 kilometers) and Tellico River mile 33 (52.8 kilometers), near the Tellico Ranger Station, Monroe County, Tennessee.	EXPN	\$1,900	\$0	\$1,900	\$0	\$1,900
	731	Madtom, smoky (<i>Noturus baileyi</i>) - Wherever found	E	\$16,000	\$10,012	\$26,012	\$0	\$26,012
	1653	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$700	\$0	\$700	\$0	\$700
	595	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(e)(1)(ii))	EXPN	\$39,500	\$0	\$39,500	\$0	\$39,500
	594	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - Wherever found, except where listed as an experimental population	T	\$29,700	\$10,012	\$39,712	\$0	\$39,712
	1002	Minnow, Devils River (<i>Dionda diaboli</i>) - Wherever found	T	\$0	\$0	\$0	\$13,000	\$13,000
	671	Minnow, loach (<i>Tiaroga cobitis</i>) -Wherever found	E	\$0	\$30,876	\$30,876	\$0	\$30,876

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1569	Minnow, Rio Grande Silvery (Hybognathus amarus) -Wherever found, except where listed as an experimental population	E	\$0	\$2,500	\$2,500	\$0	\$2,500
	951	pikeminnow, Colorado (Ptychocheilus lucius) -Salt and Verde R. drainages, AZ	EXPN	\$12,000	\$2,667	\$14,667	\$0	\$14,667
	304	pikeminnow, Colorado (Ptychocheilus lucius) -Wherever found, except where listed as an experimental population	E	\$19,100	\$132,039	\$151,139	\$0	\$151,139
	454	Poolfish, Pahrump (Empetrichthys latos) - Wherever found	E	\$58,500	\$9,254	\$67,754	\$0	\$67,754
	1413	Pupfish, Ash Meadows Amargosa (Cyprinodon nevadensis mionectes) - Wherever found	E	\$6,900	\$0	\$6,900	\$0	\$6,900
	275	Pupfish, desert (Cyprinodon macularius) - Wherever found	E	\$5,900	\$183,894	\$189,794	\$0	\$189,794
	106	Pupfish, Devils Hole (Cyprinodon diabolis) - Wherever found	E	\$408,100	\$500,000	\$908,100	\$0	\$908,100
	484	Pupfish, Owens (Cyprinodon radiosus) - Wherever found	E	\$31,500	\$27,000	\$58,500	\$0	\$58,500
	1446	Pupfish, Warm Springs (Cyprinodon nevadensis pectoralis) -Wherever found	E	\$6,300	\$0	\$6,300	\$0	\$6,300
	153	rockfish, yelloweye (Sebastes ruberrimus) - Puget Sound - Georgia Basin DPS See 50 CFR 223.102	T	\$0	\$559,000	\$559,000	\$0	\$559,000
	43	salmon Atlantic (Salmo salar) -Gulf of Maine DPS - See 50 CFR 224.101	E	\$3,316,000	\$400,000	\$3,716,000	\$0	\$3,716,000
	321	salmon Atlantic (Salmo salar) -Kennebec River DPS	E	\$131,500	\$0	\$131,500	\$0	\$131,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	25	salmon Atlantic (<i>Salmo salar</i>) -U.S.A. ME Gulf of Maine Atlantic Salmon Distinct Population Segment, which includes all naturally reproducing wild populations and those river-specific hatchery populations of Atlantic salmon having historical, river-specific characteristics found north of and including tributaries of the lower Kennebec River to, but not including, the mouth of the St. Croix River at the U.S.-Canda border. To date, the Services have determined that these populations are found in the Dennys, East Machias, Machias, Pleasant, Narraguagus, Sheepscot, and Ducktrap Rivers and in Cove Brook, Maine	E	\$0	\$6,800,000	\$6,800,000	\$0	\$6,800,000
	35	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -California Coastal ESU - See 50 CFR 223.102	T	\$0	\$4,667,000	\$4,667,000	\$0	\$4,667,000
	38	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Central Valley spring-run ESU See 50 CFR 223.102	T	\$0	\$4,567,500	\$4,567,500	\$0	\$4,567,500
	1507	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Central Valley spring-run ESU XN - See 50 CFR 223.102	EXPN	\$4,000	\$0	\$4,000	\$0	\$4,000
	9	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Lower Columbia River ESU See 50 CFR 223.102	T	\$1,030,500	\$17,302,272	\$18,332,772	\$0	\$18,332,772
	23	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Puget Sound ESU See 50 CFR 223.102	T	\$1,451,109	\$5,571,500	\$7,022,609	\$0	\$7,022,609
	33	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Sacramento River winter-run ESU - See 50 CFR 224.101	E	\$169,800	\$4,566,000	\$4,735,800	\$0	\$4,735,800
	8	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River fall-run ESU See 50 CFR 223.102	T	\$307,400	\$18,094,182	\$18,401,582	\$0	\$18,401,582

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	3	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River spring/summer-run ESU See 50 CFR 223.102	T	\$156,125	\$36,290,929	\$36,447,054	\$0	\$36,447,054
	5	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Columbia River spring-run ESU See 50 CFR 224.101	E	\$336,669	\$26,933,995	\$27,270,664	\$0	\$27,270,664
	14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Willamette River ESU See 50 CFR 223.102	T	\$0	\$11,310,060	\$11,310,060	\$0	\$11,310,060
	17	Salmon, chum (<i>Oncorhynchus keta</i>) -Columbia River ESU See 50 CFR 223.102	T	\$0	\$9,680,097	\$9,680,097	\$0	\$9,680,097
	29	Salmon, chum (<i>Oncorhynchus keta</i>) -Hood Canal summer-run ESU - See 50 CFR 223.102	T	\$693,609	\$5,319,000	\$6,012,609	\$0	\$6,012,609
	28	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Central California Coast ESU - See 50 CFR 224.101	E	\$1,153,500	\$5,030,590	\$6,184,090	\$0	\$6,184,090
	15	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Lower Columbia River ESU - See 50 CFR 223.102	T	\$145,500	\$11,080,345	\$11,225,845	\$0	\$11,225,845
	10	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Oregon Coast ESU - See 50 CFR 223.102	T	\$218,000	\$13,434,483	\$13,652,483	\$0	\$13,652,483
	18	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	T	\$98,500	\$9,376,564	\$9,475,064	\$0	\$9,475,064
	30	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) -Ozette Lake ESU - See 50 CFR 223.102	T	\$0	\$5,349,000	\$5,349,000	\$0	\$5,349,000
	11	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) -Snake River ESU - See 50 CFR 224.101	E	\$12,000	\$12,839,461	\$12,851,461	\$0	\$12,851,461
	88	Sawfish, smalltooth (<i>Pristis pectinata</i>) -US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$0	\$854,700	\$854,700	\$355,427	\$1,210,127

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	537	Sculpin, Grotto (<i>Cottus specus</i>) -Wherever found	E	\$48,000	\$0	\$48,000	\$500	\$48,500
	792	Sculpin, pygmy (<i>Cottus paulus</i> (=pygmaeus)) -Wherever found	T	\$22,000	\$0	\$22,000	\$0	\$22,000
	1608	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) -Central and Southwest Atlantic DPS	T	\$0	\$1,700	\$1,700	\$0	\$1,700
	846	Shiner, Arkansas River (<i>Notropis girardi</i>) - Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$12,000	\$0	\$12,000	\$7,240	\$19,240
	544	Shiner, blue (<i>Cyprinella caerulea</i>) - Wherever found	T	\$46,500	\$0	\$46,500	\$0	\$46,500
	1338	Shiner, Cahaba (<i>Notropis cahabae</i>) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	355	Shiner, Cape Fear (<i>Notropis mekistocholas</i>) - Wherever found	E	\$110,500	\$0	\$110,500	\$3,000	\$113,500
	952	Shiner, palezone (<i>Notropis albizonatus</i>) - Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	1473	Shiner, sharpnose (<i>Notropis oxyrhynchus</i>) - Wherever found	E	\$0	\$0	\$0	\$5,000	\$5,000
	1473	Shiner, smalleye (<i>Notropis buccula</i>) - Wherever found	E	\$0	\$0	\$0	\$5,000	\$5,000
	175	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$434,100	\$8,238	\$442,338	\$18,700	\$461,038
	1589	Silverside, Waccamaw (<i>Menidia extensa</i>) - Wherever found	T	\$1,700	\$0	\$1,700	\$500	\$2,200
	56	Smelt, delta (<i>Hypomesus transpacificus</i>) - Wherever found	T	\$2,655,800	\$12,219	\$2,668,019	\$0	\$2,668,019
	675	Spikedace (<i>Meda fulgida</i>) -Wherever found	E	\$0	\$30,552	\$30,552	\$0	\$30,552
	1063	Spinedace, Big Spring (<i>Lepidomeda mollispinis pratensis</i>) -Wherever found	T	\$10,400	\$1,206	\$11,606	\$0	\$11,606
	1601	Spinedace, Little Colorado (<i>Lepidomeda vittata</i>) -Wherever found	T	\$0	\$1,961	\$1,961	\$0	\$1,961
	626	Spinedace, White River (<i>Lepidomeda albivallis</i>) -Wherever found	E	\$36,300	\$0	\$36,300	\$0	\$36,300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	797	Springfish, Hiko White River (<i>Crenichthys baileyi grandis</i>) -Wherever found	E	\$13,400	\$8,591	\$21,991	\$0	\$21,991
	701	Springfish, Railroad Valley (<i>Crenichthys nevadae</i>) -Wherever found	T	\$28,500	\$592	\$29,092	\$0	\$29,092
	1019	Springfish, White River (<i>Crenichthys baileyi baileyi</i>) -Wherever found	E	\$12,100	\$804	\$12,904	\$0	\$12,904
	36	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -California Central Valley DPS - See 50 CFR 223.102	T	\$0	\$4,596,995	\$4,596,995	\$0	\$4,596,995
	34	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Central California Coast DPS - See 50 CFR 223.102	T	\$0	\$4,676,000	\$4,676,000	\$0	\$4,676,000
	13	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Lower Columbia River DPS - See 50 CFR 223.102	T	\$116,500	\$12,107,162	\$12,223,662	\$0	\$12,223,662
	1	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Middle Columbia River DPS - See 50 CFR 223.102	T	\$585,409	\$41,593,362	\$42,178,771	\$0	\$42,178,771
	31	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Northern California DPS - See 50 CFR 223.102	T	\$0	\$5,332,000	\$5,332,000	\$0	\$5,332,000
	26	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Puget Sound DPS - See 50 CFR 223.102	T	\$796,609	\$5,569,000	\$6,365,609	\$0	\$6,365,609
	2	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Snake River Basin DPS - See 50 CFR 223.102	T	\$148,325	\$38,473,139	\$38,621,464	\$21,000	\$38,642,464
	37	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -South-Central California Coast DPS - See 50 CFR 223.102	T	\$5,700	\$4,566,100	\$4,571,800	\$0	\$4,571,800
	39	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Southern California DPS - See 50 CFR 224.101	E	\$0	\$4,566,000	\$4,566,000	\$0	\$4,566,000
	7	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Upper Columbia River DPS - See 50 CFR 223.102	T	\$980,055	\$18,850,827	\$19,830,882	\$0	\$19,830,882

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	21	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Upper Willamette River DPS - See 50 CFR 223.102	T	\$0	\$8,271,923	\$8,271,923	\$0	\$8,271,923
	417	Stickleback, unarmored threespine (<i>Gasterosteus aculeatus williamsoni</i>) - Wherever found	E	\$83,900	\$0	\$83,900	\$0	\$83,900
	550	Sturgeon, Alabama (<i>Scaphirhynchus suttkusi</i>) -Wherever found	E	\$46,000	\$0	\$46,000	\$0	\$46,000
	80	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Carolina DPS - See 50 CFR 224.101	E	\$628,000	\$782,800	\$1,410,800	\$100	\$1,410,900
	109	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Chesapeake Bay DPS - See 50 CFR 224.101	E	\$14,000	\$845,500	\$859,500	\$0	\$859,500
	174	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Gulf of Maine DPS - See 50 CFR 223.102	T	\$0	\$470,500	\$470,500	\$0	\$470,500
	120	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -New York Bight DPS - See 50 CFR 224.101	E	\$0	\$792,564	\$792,564	\$0	\$792,564
	103	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -South Atlantic DPS - See 50 CFR 224.101	E	\$3,000	\$865,156	\$868,156	\$71,754	\$939,910
	142	sturgeon, Green (<i>Acipenser medirostris</i>) - Southern DPS - See 50 CFR 223.102	T	\$55,000	\$592,648	\$647,648	\$0	\$647,648
	89	Sturgeon, Gulf (<i>Acipenser oxyrinchus</i> (=oxyrhynchus) desotoi) -Wherever found	T	\$528,000	\$577,180	\$1,105,180	\$71,754	\$1,176,934
	156	Sturgeon, pallid (<i>Scaphirhynchus albus</i>) - Wherever found	E	\$279,200	\$196,965	\$476,165	\$79,881	\$556,046
	73	Sturgeon, shortnose (<i>Acipenser brevirostrum</i>) -Wherever found	E	\$18,500	\$1,135,940	\$1,154,440	\$492,755	\$1,647,195
	12	Sturgeon, white (<i>Acipenser transmontanus</i>) - U.S.A. (ID, MT), Canada (B.C.), (Kootenai R. system)	E	\$190,633	\$12,236,330	\$12,426,963	\$1,100	\$12,428,063
	599	Sucker, June (<i>Chasmistes liorus</i>) -Wherever found	T	\$39,000	\$0	\$39,000	\$0	\$39,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	27	Sucker, Lost River (<i>Deltistes luxatus</i>) - Wherever found	E	\$6,188,934	\$5,000	\$6,193,934	\$0	\$6,193,934
	264	Sucker, razorback (<i>Xyrauchen texanus</i>) - Wherever found	E	\$61,400	\$144,299	\$205,699	\$0	\$205,699
	245	Sucker, Santa Ana (<i>Catostomus santaanae</i>) - Los Angeles River basin, San Gabriel River basin, Santa Ana River basin	T	\$243,800	\$0	\$243,800	\$0	\$243,800
	20	Sucker, shortnose (<i>Chasmistes brevirostris</i>) - Wherever found	E	\$8,747,301	\$7,800	\$8,755,101	\$0	\$8,755,101
	107	Sucker, Warner (<i>Catostomus warnerensis</i>) - Wherever found	T	\$21,500	\$879,000	\$900,500	\$0	\$900,500
	1675	Sucker, Zuni bluehead (<i>Catostomus discobolus yarrowi</i>) -Wherever found	E	\$0	\$200	\$200	\$0	\$200
	1362	Sunfish, spring pygmy (<i>Elassoma alabamae</i>) -Wherever found	T	\$7,500	\$0	\$7,500	\$0	\$7,500
	479	Topminnow, barrens (<i>Fundulus julisia</i>) - Wherever found	E	\$59,500	\$0	\$59,500	\$0	\$59,500
	847	Topminnow, Gila (incl. Yaqui) (<i>Poeciliopsis occidentalis</i>) -U.S. Only	E	\$0	\$19,130	\$19,130	\$0	\$19,130
	827	Trout, Apache (<i>Oncorhynchus apache</i>) - Wherever found	T	\$20,000	\$200	\$20,200	\$0	\$20,200
	76	Trout, bull (<i>Salvelinus confluentus</i>) - Clackamas River subbasin and the mainstem Willamette River, from Willamette Falls to its points of confluence with the Columbia River, including Multnomah Channel	EXPN	\$1,582,209	\$0	\$1,582,209	\$0	\$1,582,209
	6	Trout, bull (<i>Salvelinus confluentus</i>) -U.S.A., conterminous, (lower 48 states)	T	\$7,470,623	\$15,293,109	\$22,763,732	\$447,000	\$23,210,732
	888	Trout, Gila (<i>Oncorhynchus gilae</i>) -Wherever found	T	\$17,000	\$200	\$17,200	\$0	\$17,200
	222	trout, Greenback Cutthroat (<i>Oncorhynchus clarkii stomias</i>) -Wherever found	T	\$31,200	\$53,000	\$84,200	\$244,358	\$328,558
	16	Trout, Lahontan cutthroat (<i>Oncorhynchus clarkii henshawi</i>) -Wherever found	T	\$7,443,400	\$2,905,634	\$10,349,034	\$0	\$10,349,034
	1541	Trout, Little Kern golden (<i>Oncorhynchus aguabonita whitei</i>) -Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1462	Trout, Paiute cutthroat (<i>Oncorhynchus clarkii seleniris</i>) -Wherever found	T	\$5,800	\$0	\$5,800	\$0	\$5,800
	403	Tui chub, Mohave (<i>Gila bicolor</i> ssp. <i>mohavensis</i>) -Wherever found	E	\$0	\$87,585	\$87,585	\$0	\$87,585
	285	Woundfin (<i>Plagopterus argentissimus</i>) -Gila R. drainage, AZ, NM	EXPN	\$170,000	\$0	\$170,000	\$0	\$170,000
	660	Woundfin (<i>Plagopterus argentissimus</i>) - Wherever found, except where listed as an experimental population	E	\$14,500	\$17,573	\$32,073	\$0	\$32,073
	118	grouper, Nassau (<i>Epinephelus striatus</i>) -	T	\$0	\$794,700	\$794,700	\$0	\$794,700
Fishes subtotal				\$54,287,210	\$400,528,043	\$454,815,253	\$2,050,691	\$456,865,944
Flowering Plants	361	`Ahinahina (<i>Argyroxiphium sandwicense</i> ssp. <i>macrocephalum</i>) -Wherever found	T	\$9,500	\$102,410	\$111,910	\$0	\$111,910
	1151	`Ahinahina (<i>Argyroxiphium sandwicense</i> ssp. <i>sandwicense</i>) -Wherever found	E	\$10,051	\$0	\$10,051	\$0	\$10,051
	1185	`Aiakeakua, popolo (<i>Solanum sandwicense</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	854	`Aiea (<i>Nothocestrum breviflorum</i>) -Wherever found	E	\$18,930	\$0	\$18,930	\$0	\$18,930
	713	`Aiea (<i>Nothocestrum latifolium</i>) -Wherever found	E	\$18,000	\$10,241	\$28,241	\$0	\$28,241
	1185	`Aiea (<i>Nothocestrum peltatum</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	`Akoko (<i>Euphorbia celastroides</i> var. <i>kaenana</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1292	`Akoko (<i>Euphorbia deppeana</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1064	`Akoko (<i>Euphorbia eleanoriae</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1362	`Akoko (<i>Euphorbia haeleeleana</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1185	`Akoko (<i>Euphorbia herbstii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1416	`Akoko (<i>Euphorbia kuwaleana</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1185	`Akoko (<i>Euphorbia remyi</i> var. <i>kauaiensis</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	878	`Akoko (Euphorbia remyi var. remyi) - Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500
	1416	`Akoko (Euphorbia rockii) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	380	`Akoko, Ewa Plains (Euphorbia skottsbergii var. skottsbergii) -Wherever found	E	\$102,500	\$0	\$102,500	\$0	\$102,500
	643	`aku (Cyanea tritomantha) -Wherever found	E	\$33,000	\$1,803	\$34,803	\$0	\$34,803
	1185	`aku`aku (Cyanea platyphylla) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	935	`Ala `ala wai nui (Peperomia subpetiolata) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	1185	`Anaunau (Lepidium arbuscula) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1108	`Anunu (Sicyos albus) -Wherever found	E	\$9,500	\$1,803	\$11,303	\$0	\$11,303
	757	`Anunu (Sicyos macrophyllus) -Wherever found	E	\$22,930	\$1,803	\$24,733	\$0	\$24,733
	1132	`Awikiwiki (Canavalia molokaiensis) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1185	`Awikiwiki (Canavalia napaliensis) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	727	`Awikiwiki (Canavalia pubescens) - Wherever found	E	\$26,500	\$0	\$26,500	\$0	\$26,500
	599	`Ena`ena (Pseudognaphalium sandwicensium var. molokaiense) - Wherever found	E	\$39,000	\$0	\$39,000	\$0	\$39,000
	1152	`Oha wai (Clermontia drepanomorpha) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	357	`Oha wai (Clermontia lindseyana) - Wherever found	E	\$111,177	\$1,803	\$112,980	\$0	\$112,980
	952	`Oha wai (Clermontia oblongifolia ssp. brevipes) -Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	1064	`Oha wai (Clermontia oblongifolia ssp. mauiensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	342	`Oha wai (Clermontia peleana) -Wherever found	E	\$117,177	\$1,803	\$118,980	\$0	\$118,980

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	360	`Oha wai (Clermontia pyralaria) -Wherever found	E	\$112,177	\$0	\$112,177	\$0	\$112,177
	474	`Oha wai (Clermontia samuelii) -Wherever found	E	\$9,500	\$51,208	\$60,708	\$0	\$60,708
	659	`Ohe (Joinvillea ascendens ascendens) -Wherever found	E	\$12,000	\$20,483	\$32,483	\$0	\$32,483
	1362	`Ohe`ohe (Polyscias gymnocarpa) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1292	A`e (Zanthoxylum dipetalum var. tomentosum) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1185	A`e (Zanthoxylum hawaiiense) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1292	A`e (Zanthoxylum oahuense) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1185	"Akoko (Euphorbia halemanui) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1152	Alani (Melicope adscendens) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1362	Alani (Melicope balloui) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1362	Alani (Melicope christophersenii) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	570	Alani (Melicope degeneri) -Wherever found	E	\$43,500	\$0	\$43,500	\$0	\$43,500
	1064	Alani (Melicope haupuensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1292	Alani (Melicope hiiakae) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1292	Alani (Melicope knudsenii) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1292	Alani (Melicope lydgatei) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1185	Alani (Melicope makahae) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1185	Alani (Melicope mucronulata) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1258	Alani (Melicope munroi) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	719	Alani (Melicope ovalis) -Wherever found	E	\$7,500	\$20,483	\$27,983	\$0	\$27,983
	1185	Alani (Melicope pallida) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	570	Alani (Melicope paniculata) -Wherever found	E	\$43,500	\$0	\$43,500	\$0	\$43,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	Alani (Melicope puberula) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1185	Alani (Melicope quadrangularis) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	Alani (Melicope reflexa) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1258	Alani (Melicope saint-johnii) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1258	Alani (Melicope zahlbruckneri) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1255	Allocarya, Calistoga (Plagiobothrys strictus) -Wherever found	E	\$9,200	\$0	\$9,200	\$0	\$9,200
	1504	Alopecurus, Sonoma (Alopecurus aequalis var. sonomensis) -Wherever found	E	\$4,000	\$250	\$4,250	\$0	\$4,250
	290	Amaranth, seabeach (Amaranthus pumilus) -Wherever found	T	\$71,500	\$90,102	\$161,602	\$900	\$162,502
	1653	Ambrosia, San Diego (Ambrosia pumila) -Wherever found	E	\$700	\$0	\$700	\$0	\$700
	809	Ambrosia, south Texas (Ambrosia cheiranthifolia) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	1130	amole, purple (Chlorogalum purpureum) -Wherever found	T	\$10,700	\$0	\$10,700	\$0	\$10,700
	1495	Amphianthus, little (Amphianthus pusillus) -Wherever found	T	\$4,500	\$0	\$4,500	\$0	\$4,500
	820	Aplokating-palaoan (Psychotria malaspinae) -Wherever found	E	\$20,551	\$0	\$20,551	\$0	\$20,551
	896	Arrowhead, bunched (Sagittaria fasciculata) -Wherever found	E	\$12,500	\$0	\$12,500	\$4,000	\$16,500
	786	Aster, decurrent false (Boltonia decurrens) -Wherever found	T	\$22,500	\$0	\$22,500	\$0	\$22,500
	1338	Aster, Ruth's golden (Pityopsis ruthii) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1362	Aupaka (Isodendron hosakae) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	967	Aupaka (Isodendron laurifolium) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	919	Aupaka (Isodendron longifolium) -Wherever found	T	\$15,500	\$0	\$15,500	\$0	\$15,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1022	Avens, spreading (<i>Geum radiatum</i>) - Wherever found	E	\$6,500	\$3,350	\$9,850	\$3,000	\$12,850
	1402	'Awiwi (<i>Kadua cookiana</i>) -Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1362	Awiwi (<i>Schenkia sebaeoides</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	809	Ayenia, Texas (<i>Ayenia limitaris</i>) -Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	799	Barbara's Buttons, Mohr's (<i>Marshallia mohrii</i>) -Wherever found	T	\$21,500	\$0	\$21,500	\$0	\$21,500
	1565	Barberry, island (<i>Berberis pinnata</i> ssp. <i>insularis</i>) -Wherever found	E	\$2,800	\$0	\$2,800	\$0	\$2,800
	1633	Barberry, Nevin's (<i>Berberis nevinii</i>) - Wherever found	E	\$1,300	\$0	\$1,300	\$0	\$1,300
	1258	Bariaco (<i>Trichilia triacantha</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	873	Beaked-rush, Knieskern's (<i>Rhynchospora knieskernii</i>) -Wherever found	T	\$14,500	\$3,456	\$17,956	\$0	\$17,956
	1258	bean, sea (<i>Mucuna sloanei</i> var. <i>persericea</i>) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	805	beardtongue, Parachute (<i>Penstemon debilis</i>) - Wherever found	T	\$13,000	\$0	\$13,000	\$8,300	\$21,300
	1258	Beardtongue, Penland (<i>Penstemon penlandii</i>) -Wherever found	E	\$4,000	\$5,000	\$9,000	\$0	\$9,000
	478	Bear-poppy, Dwarf (<i>Arctomecon humilis</i>) - Wherever found	E	\$28,000	\$31,648	\$59,648	\$0	\$59,648
	1416	Bedstraw, El Dorado (<i>Galium californicum ssp. sierrae</i>) -Wherever found	E	\$1,500	\$5,000	\$6,500	\$0	\$6,500
	888	Bedstraw, island (<i>Galium buxifolium</i>) - Wherever found	E	\$12,200	\$5,000	\$17,200	\$0	\$17,200
	619	Birch, Virginia round-leaf (<i>Betula uber</i>) - Wherever found	T	\$37,000	\$0	\$37,000	\$0	\$37,000
	929	Bird's beak, palmate-bracted (<i>Cordylanthus palmatus</i>) -Wherever found	E	\$8,900	\$6,549	\$15,449	\$0	\$15,449
	1624	Bird's-beak, Pennell's (<i>Cordylanthus tenuis ssp. capillaris</i>) -Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	726	Bird's-beak, salt marsh (<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>) -Wherever found	E	\$26,600	\$100	\$26,700	\$0	\$26,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	539	Bird's-beak, soft (<i>Cordylanthus mollis</i> ssp. <i>mollis</i>) -Wherever found	E	\$48,300	\$0	\$48,300	\$0	\$48,300
	1672	Bittercress, small-anthered (<i>Cardamine micranthera</i>) -Wherever found	E	\$300	\$0	\$300	\$0	\$300
	903	Bladderpod, Dudley Bluffs (<i>Lesquerella congesta</i>) -Wherever found	T	\$11,000	\$5,000	\$16,000	\$425	\$16,425
	967	Bladderpod, kodachrome (<i>Lesquerella tumulosa</i>) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1602	Bladderpod, lyrate (<i>Lesquerella lyrata</i>) -Wherever found	T	\$1,900	\$0	\$1,900	\$0	\$1,900
	432	bladderpod, Missouri (<i>Physaria filiformis</i>) -Wherever found	T	\$28,000	\$50,000	\$78,000	\$100	\$78,100
	520	Bladderpod, San Bernardino Mountains (<i>Lesquerella kingii</i> ssp. <i>bernardina</i>) -Wherever found	E	\$53,100	\$0	\$53,100	\$0	\$53,100
	650	Bladderpod, Short's (<i>Physaria globosa</i>) -Wherever found	E	\$33,500	\$0	\$33,500	\$300	\$33,800
	1026	Bladderpod, Spring Creek (<i>Lesquerella perforata</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1402	Bladderpod, white (<i>Physaria pallida</i>) -Wherever found	E	\$0	\$0	\$0	\$7,000	\$7,000
	251	Bladderpod, White Bluffs (<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>) -Wherever found	T	\$234,088	\$0	\$234,088	\$0	\$234,088
	803	Bladderpod, Zapata (<i>Physaria thamnophila</i>) -Wherever found	E	\$9,400	\$0	\$9,400	\$12,000	\$21,400
	682	Blazingstar, Ash Meadows (<i>Mentzelia leucophylla</i>) -Wherever found	T	\$100	\$30,000	\$30,100	\$0	\$30,100
	1291	Blazingstar, Heller's (<i>Liatris helleri</i>) -Wherever found	T	\$300	\$4,750	\$5,050	\$3,500	\$8,550
	1185	Bluegrass, Hawaiian (<i>Poa sandvicensis</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1185	Bluegrass, Mann's (<i>Poa mannii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1451	Bluegrass, Napa (<i>Poa napensis</i>) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1536	Blue-star, Kearney's (<i>Amsonia kearneyana</i>) - Wherever found	E	\$0	\$3,200	\$3,200	\$0	\$3,200
	1535	Bluet, Roan Mountain (<i>Hedyotis purpurea</i> var. <i>montana</i>) -Wherever found	E	\$800	\$0	\$800	\$2,500	\$3,300
	1402	Boxwood, Vahl's (<i>Buxus vahlii</i>) -Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1540	Brodiaea, Chinese Camp (<i>Brodiaea pallida</i>) - Wherever found	T	\$3,100	\$0	\$3,100	\$0	\$3,100
	1500	Buckwheat, cushenbury (<i>Eriogonum</i> ovalifolium var. <i>vineum</i>) -Wherever found	E	\$4,300	\$0	\$4,300	\$0	\$4,300
	1624	Buckwheat, Ione (incl. Irish Hill) (<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i>)) -Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	451	Buckwheat, steamboat (<i>Eriogonum</i> ovalifolium var. <i>williamsiae</i>) -Wherever found	E	\$13,300	\$54,833	\$68,133	\$0	\$68,133
	551	Bush-clover, prairie (<i>Lespedeza</i> <i>leptostachya</i>) -Wherever found	T	\$45,300	\$76	\$45,376	\$500	\$45,876
	1473	Bush-mallow, San Clemente Island (<i>Malacothamnus clementinus</i>) -Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1500	Bush-mallow, Santa Cruz Island (<i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i>) -Wherever found	E	\$1,300	\$3,000	\$4,300	\$0	\$4,300
	749	Buttercup, autumn (<i>Ranunculus aestivalis</i> (= <i>acriiformis</i>)) -Wherever found	E	\$25,000	\$0	\$25,000	\$0	\$25,000
	828	Butterweed, Layne's (<i>Senecio layneae</i>) - Wherever found	T	\$15,000	\$5,000	\$20,000	\$0	\$20,000
	1672	Button-celery, San Diego (<i>Eryngium</i> <i>aristulatum</i> var. <i>parishii</i>) -Wherever found	E	\$300	\$0	\$300	\$0	\$300
	1606	Cactus, Acuña (<i>Echinomastus</i> <i>erectocentrus</i> var. <i>acunensis</i>) -Wherever found	E	\$0	\$1,850	\$1,850	\$0	\$1,850
	1675	Cactus, Arizona hedgehog (<i>Echinocereus</i> <i>arizonicus</i> ssp. <i>arizonicus</i>) -Wherever found	E	\$0	\$200	\$200	\$0	\$200
	856	Cactus, Bakersfield (<i>Opuntia treleasei</i>) - Wherever found	E	\$18,800	\$0	\$18,800	\$0	\$18,800

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1111	Cactus, black lace (<i>Echinocereus reichenbachii</i> var. <i>albertii</i>) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1566	Cactus, Brady pincushion (<i>Pediocactus bradyi</i>) -Wherever found	E	\$0	\$2,700	\$2,700	\$0	\$2,700
	1541	Cactus, Chisos Mountain hedgehog (<i>Echinocereus chisoensis</i> var. <i>chisoensis</i>) -Wherever found	T	\$0	\$0	\$0	\$3,000	\$3,000
	282	Cactus, Colorado hookless (<i>Sclerocactus glaucus</i>) -Wherever found	T	\$64,500	\$112,701	\$177,201	\$1,225	\$178,426
	1414	Cactus, Fickeisen plains (<i>Pediocactus peeblesianus</i> ssp. <i>fickeiseniae</i>) -Wherever found	E	\$500	\$6,200	\$6,700	\$0	\$6,700
	1586	Cactus, Knowlton's (<i>Pediocactus knowltonii</i>) -Wherever found	E	\$0	\$1,000	\$1,000	\$1,475	\$2,475
	419	Cactus, Kuenzler hedgehog (<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>) -Wherever found	T	\$0	\$83,000	\$83,000	\$0	\$83,000
	1505	Cactus, Lloyd's Mariposa (<i>Sclerocactus mariposensis</i>) -Wherever found	T	\$0	\$1,200	\$1,200	\$3,000	\$4,200
	303	Cactus, Mesa Verde (<i>Sclerocactus mesae-verdae</i>) -Wherever found	T	\$3,000	\$146,847	\$149,847	\$1,475	\$151,322
	638	cactus, Pariette (<i>Sclerocactus brevispinus</i>) -Wherever found	T	\$0	\$35,122	\$35,122	\$0	\$35,122
	1131	Cactus, Pima pineapple (<i>Coryphantha scheeri</i> var. <i>robustispina</i>) -Wherever found	E	\$0	\$10,600	\$10,600	\$0	\$10,600
	561	Cactus, San Rafael (<i>Pediocactus despainii</i>) -Wherever found	E	\$20,000	\$25,063	\$45,063	\$0	\$45,063
	1147	Cactus, Siler pincushion (<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>) -Wherever found	T	\$1,000	\$9,496	\$10,496	\$0	\$10,496
	1002	Cactus, star (<i>Astrophytum asterias</i>) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1338	cactus, Tobusch fishhook (<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>) -Wherever found	T	\$0	\$0	\$0	\$8,000	\$8,000
	607	cactus, Uinta Basin hookless (<i>Sclerocactus wetlandicus</i>) -Wherever found	T	\$0	\$38,191	\$38,191	\$0	\$38,191

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	549	Cactus, Winkler (<i>Pediocactus winkleri</i>) - Wherever found	T	\$17,000	\$29,220	\$46,220	\$0	\$46,220
	518	Cactus, Wright fishhook (<i>Sclerocactus wrightiae</i>) -Wherever found	E	\$14,000	\$39,197	\$53,197	\$0	\$53,197
	1608	Campion, fringed (<i>Silene polypetala</i>) - Wherever found	E	\$1,700	\$0	\$1,700	\$0	\$1,700
	1541	Capa rosa (<i>Callicarpa ampla</i>) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	193	Catchfly, Spalding's (<i>Silene spaldingii</i>) - Wherever found	T	\$360,634	\$23,500	\$384,134	\$9,000	\$393,134
	935	Ceanothus, coyote (<i>Ceanothus ferrisiae</i>) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	1416	Ceanothus, Pine Hill (<i>Ceanothus roderickii</i>) - Wherever found	E	\$1,500	\$5,000	\$6,500	\$0	\$6,500
	919	Centaury, spring-loving (<i>Centaurium namophilum</i>) -Wherever found	T	\$12,500	\$3,000	\$15,500	\$0	\$15,500
	501	Chaff-flower, round-leaved (<i>Achyranthes splendens</i> var. <i>rotundata</i>) -Wherever found	E	\$56,000	\$0	\$56,000	\$0	\$56,000
	326	Chaffseed, American (<i>Schwalbea americana</i>) -Wherever found	E	\$124,000	\$0	\$124,000	\$2,500	\$126,500
	934	Checker-mallow, Keck's (<i>Sidalcea keckii</i>) - Wherever found	E	\$9,400	\$5,602	\$15,002	\$0	\$15,002
	1449	Checker-mallow, Kenwood Marsh (<i>Sidalcea oregana</i> ssp. <i>valida</i>) -Wherever found	E	\$6,200	\$0	\$6,200	\$0	\$6,200
	243	Checker-mallow, Nelson's (<i>Sidalcea nelsoniana</i>) -Wherever found	T	\$223,134	\$28,000	\$251,134	\$0	\$251,134
	302	Checkermallow, Wenatchee Mountains (<i>Sidalcea oregana</i> var. <i>calva</i>) -Wherever found	E	\$151,474	\$0	\$151,474	\$0	\$151,474
	1464	Chumbo, Higo (<i>Harrisia portoricensis</i>) - Wherever found	T	\$5,500	\$0	\$5,500	\$0	\$5,500
	1416	Chupacallos (<i>Pleodendron macranthum</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	783	Clarkia, Pismo (<i>Clarkia speciosa</i> ssp. <i>immaculata</i>) -Wherever found	E	\$22,600	\$100	\$22,700	\$0	\$22,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1258	Clarkia, Presidio (Clarkia franciscana) - Wherever found	E	\$8,200	\$800	\$9,000	\$0	\$9,000
	1026	Clarkia, Springville (Clarkia springvillensis) -Wherever found	T	\$12,500	\$0	\$12,500	\$0	\$12,500
	749	Clarkia, Vine Hill (Clarkia imbricata) - Wherever found	E	\$25,000	\$0	\$25,000	\$0	\$25,000
	1287	Cliffrose, Arizona (Purshia (=Cowania) subintegra) -Wherever found	E	\$0	\$8,900	\$8,900	\$0	\$8,900
	1563	Clover, Monterey (Trifolium trichocalyx) - Wherever found	E	\$2,900	\$0	\$2,900	\$0	\$2,900
	876	Clover, showy Indian (Trifolium amoenum) - Wherever found	E	\$17,500	\$200	\$17,700	\$0	\$17,700
	1416	Cobana negra (Stahlia monosperma) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	633	Coneflower, smooth (Echinacea laevigata) - Wherever found	T	\$30,000	\$400	\$30,400	\$5,500	\$35,900
	1451	Cory cactus, bunched (Coryphantha ramillosa) -Wherever found	T	\$0	\$3,000	\$3,000	\$3,000	\$6,000
	500	Cycladenia, Jones (Cycladenia humilis var. jonesii) -Wherever found	T	\$28,000	\$28,228	\$56,228	\$0	\$56,228
	876	daisy, Lakeside (Hymenoxys herbacea) - Wherever found	T	\$17,700	\$0	\$17,700	\$0	\$17,700
	560	Daisy, Parish's (Erigeron parishii) - Wherever found	T	\$43,100	\$2,000	\$45,100	\$0	\$45,100
	396	Daisy, Willamette (Erigeron decumbens) - Wherever found	E	\$70,500	\$23,000	\$93,500	\$0	\$93,500
	314	Dawn-flower, Texas prairie (Hymenoxys texana) -Wherever found	E	\$138,000	\$0	\$138,000	\$0	\$138,000
	293	Desert buckwheat, Umtanum (Eriogonum codium) -Wherever found	T	\$158,228	\$0	\$158,228	\$0	\$158,228
	891	Dogweed, ashy (Thymophylla tephroleuca) - Wherever found	E	\$17,000	\$0	\$17,000	\$0	\$17,000
	661	Dropwort, Canby's (Oxypolis canbyi) - Wherever found	E	\$29,500	\$0	\$29,500	\$2,500	\$32,000
	1038	Dudleya, Conejo (Dudleya abramsii ssp. parva) -Wherever found	T	\$12,100	\$0	\$12,100	\$0	\$12,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	984	Dudleya, marcescent (<i>Dudleya cymosa</i> ssp. <i>marcescens</i>) -Wherever found	T	\$13,400	\$300	\$13,700	\$0	\$13,700
	947	Dudleya, Santa Clara Valley (<i>Dudleya setchellii</i>) -Wherever found	E	\$14,900	\$0	\$14,900	\$0	\$14,900
	790	Dudleya, Santa Cruz Island (<i>Dudleya nesiotica</i>) -Wherever found	T	\$12,200	\$10,000	\$22,200	\$0	\$22,200
	1633	Dudleya, Santa Monica Mountains (<i>Dudleya cymosa</i> ssp. <i>ovatifolia</i>) -Wherever found	T	\$1,300	\$0	\$1,300	\$0	\$1,300
	375	Dudleya, Verity's (<i>Dudleya verityi</i>) -Wherever found	T	\$103,200	\$1,500	\$104,700	\$0	\$104,700
	384	Dwarf-flax, Marin (<i>Hesperolinon congestum</i>) -Wherever found	T	\$99,400	\$900	\$100,300	\$0	\$100,300
	1451	Erubia (<i>Solanum drymophilum</i>) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	430	Evening-primrose, Antioch Dunes (<i>Oenothera deltoides</i> ssp. <i>howellii</i>) -Wherever found	E	\$79,000	\$0	\$79,000	\$0	\$79,000
	967	Fescue, Guadalupe (<i>Festuca ligulata</i>) -Wherever found	E	\$0	\$1,000	\$1,000	\$13,000	\$14,000
	819	Fiddleneck, large-flowered (<i>Amsinckia grandiflora</i>) -Wherever found	E	\$14,400	\$6,547	\$20,947	\$0	\$20,947
	680	Flannelbush, Mexican (<i>Fremontodendron mexicanum</i>) -Wherever found	E	\$30,300	\$0	\$30,300	\$0	\$30,300
	437	Flannelbush, Pine Hill (<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>) -Wherever found	E	\$1,500	\$75,000	\$76,500	\$0	\$76,500
	1330	Fleabane, Zuni (<i>Erigeron rhizomatus</i>) -Wherever found	T	\$0	\$8,400	\$8,400	\$0	\$8,400
	268	Four-o'clock, MacFarlane's (<i>Mirabilis macfarlanei</i>) -Wherever found	T	\$93,302	\$108,000	\$201,302	\$0	\$201,302
	1469	Fringepod, Santa Cruz Island (<i>Thysanocarpus conchuliferus</i>) -Wherever found	E	\$4,900	\$400	\$5,300	\$0	\$5,300
	397	Fritillary, Gentner's (<i>Fritillaria gentneri</i>) -Wherever found	E	\$93,481	\$0	\$93,481	\$0	\$93,481

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1002	Gardenia (=Na`u), Hawaiian (Gardenia brighamii) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	708	Gerardia, sandplain (Agalinis acuta) -Wherever found	E	\$23,000	\$0	\$23,000	\$5,500	\$28,500
	1446	Gilia, Hoffmann's slender-flowered (Gilia tenuiflora ssp. hoffmannii) -Wherever found	E	\$1,300	\$5,000	\$6,300	\$0	\$6,300
	1360	Gilia, Monterey (Gilia tenuiflora ssp. arenaria) -Wherever found	E	\$7,800	\$0	\$7,800	\$0	\$7,800
	648	Glade cress, Kentucky (Leavenworthia exigua laciniata) -Wherever found	T	\$34,000	\$0	\$34,000	\$0	\$34,000
	967	Gladecress, Fleshy-fruit (Leavenworthia crassa) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	732	Gladecress, Texas golden (Leavenworthia texana) -Wherever found	E	\$19,000	\$0	\$19,000	\$7,000	\$26,000
	1416	Goetzea, beautiful (Goetzea elegans) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1663	Goldenrod, Blue Ridge (Solidago spithamaea) -Wherever found	T	\$300	\$0	\$300	\$300	\$600
	1041	Goldenrod, Houghton's (Solidago houghtonii) -Wherever found	T	\$12,000	\$0	\$12,000	\$0	\$12,000
	789	Goldenrod, Short's (Solidago shortii) -Wherever found	E	\$21,100	\$0	\$21,100	\$1,200	\$22,300
	1258	Goldfields, Burke's (Lasthenia burkei) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	318	Goldfields, Contra Costa (Lasthenia conjugens) -Wherever found	E	\$127,000	\$6,547	\$133,547	\$0	\$133,547
	963	Grass, Colusa (Neostapfia colusana) -Wherever found	T	\$8,700	\$5,602	\$14,302	\$0	\$14,302
	1445	Grass, Solano (Tuctoria mucronata) -Wherever found	E	\$700	\$5,602	\$6,302	\$0	\$6,302
	732	Grass, Tennessee yellow-eyed (Xyris tennesseensis) -Wherever found	E	\$26,000	\$0	\$26,000	\$0	\$26,000
	599	Ground-plum, Guthrie's (=Pyne's) (Astragalus bibullatus) -Wherever found	E	\$39,000	\$0	\$39,000	\$0	\$39,000
	682	Gumplant, Ash Meadows (Grindelia fraxinipratensis) -Wherever found	T	\$100	\$30,000	\$30,100	\$0	\$30,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1402	Ha`iwale (Cyrtandra crenata) -Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1362	Ha`iwale (Cyrtandra dentata) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1041	Ha`iwale (Cyrtandra filipes) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1288	Ha`iwale (Cyrtandra giffardii) -Wherever found	E	\$7,000	\$1,803	\$8,803	\$0	\$8,803
	1185	Ha`iwale (Cyrtandra hematos) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1111	Ha`iwale (Cyrtandra kaulantha) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1041	Ha`iwale (Cyrtandra limahuliensis) -Wherever found	T	\$12,000	\$0	\$12,000	\$0	\$12,000
	1111	Ha`iwale (Cyrtandra munroi) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1185	Ha`iwale (Cyrtandra oenobarba) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	Ha`iwale (Cyrtandra oxybapha) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1152	Ha`iwale (Cyrtandra polyantha) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1416	Ha`iwale (Cyrtandra sessilis) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1416	Ha`iwale (Cyrtandra subumbellata) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	987	Ha`iwale (Cyrtandra tintinnabula) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	1292	Ha`iwale (Cyrtandra viridiflora) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1111	haha (Cyanea acuminata) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1185	haha (Cyanea asarifolia) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	604	haha (Cyanea asplenifolia) -Wherever found	E	\$7,500	\$30,724	\$38,224	\$0	\$38,224
	1362	haha (Cyanea calycina) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1416	haha (Cyanea copelandii ssp. copelandii) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	526	haha (Cyanea copelandii ssp. haleakalaensis) -Wherever found	E	\$8,000	\$41,966	\$49,966	\$0	\$49,966
	1111	haha (Cyanea crispa) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1064	haha (Cyanea dolichopoda) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1258	haha (Cyanea dunbariae) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1258	haha (Cyanea duvalliorum) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1064	haha (Cyanea eleeleensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1064	haha (Cyanea gibsonii) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	865	haha (Cyanea glabra) -Wherever found	E	\$8,000	\$10,241	\$18,241	\$0	\$18,241
	1064	haha (Cyanea grimesiana ssp. grimesiana) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	haha (Cyanea grimesiana ssp. obatae) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1258	haha (Cyanea hamatiflora ssp. carlsonii) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	657	haha (Cyanea hamatiflora ssp. hamatiflora) -Wherever found	E	\$12,430	\$20,483	\$32,913	\$0	\$32,913
	1258	haha (Cyanea humboldtiana) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1064	haha (Cyanea kolekoleensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1292	haha (Cyanea koolauensis) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1064	haha (Cyanea kuhihewa) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	719	haha (Cyanea kunthiana) -Wherever found	E	\$7,500	\$20,483	\$27,983	\$0	\$27,983
	1152	haha (Cyanea lanceolata) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1064	haha (Cyanea lobata) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1362	haha (Cyanea longiflora) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1152	haha (Cyanea magnicalyx) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1362	haha (Cyanea mannii) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	825	haha (Cyanea maritae) -Wherever found	E	\$10,000	\$10,241	\$20,241	\$0	\$20,241
	631	haha (Cyanea marksii) -Wherever found	E	\$35,930	\$0	\$35,930	\$0	\$35,930
	1338	haha (Cyanea mauiensis) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1338	haha (Cyanea mceldowneyi) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1185	haha (Cyanea munroi) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1185	haha (Cyanea obtusa) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1258	haha (Cyanea pinnatifida) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1185	haha (Cyanea procera) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1185	haha (Cyanea profuga) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1292	haha (Cyanea purpurellifolia) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1041	haha (Cyanea recta) -Wherever found	T	\$12,000	\$0	\$12,000	\$0	\$12,000
	1064	haha (Cyanea remyi) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	696	haha (Cyanea rivularis) -Wherever found	E	\$29,500	\$0	\$29,500	\$0	\$29,500
	339	haha (Cyanea shipmanii) -Wherever found	E	\$118,177	\$1,803	\$119,980	\$0	\$119,980
	987	haha (Cyanea st.-johnii) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	737	haha (Cyanea stictophylla) -Wherever found	E	\$23,930	\$1,803	\$25,733	\$0	\$25,733
	1185	haha (Cyanea superba) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1152	haha (Cyanea truncata) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1064	haha (Cyanea undulata) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	590	Haiwale (Cyrtandra ferripilosa) -Wherever found	E	\$9,500	\$30,724	\$40,224	\$0	\$40,224
	1292	Haiwale (Cyrtandra gracilis) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	577	Haiwale (Cyrtandra nanawaleensis) -Wherever found	E	\$42,000	\$0	\$42,000	\$0	\$42,000
	1064	Haiwale (Cyrtandra paliku) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	669	Haiwale (Cyrtandra wagneri) -Wherever found	E	\$31,000	\$0	\$31,000	\$0	\$31,000
	1185	Haiwale (Cyrtandra waiolani) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	Hala pepe (Pleomele fernaldii) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1292	Hala pepe (Pleomele forbesii) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	914	Hala pepe (Pleomele hawaiiensis) -Wherever found	E	\$14,000	\$1,803	\$15,803	\$0	\$15,803

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	628	halomtano, Berenghenas (<i>Solanum guamense</i>) -Wherever found	E	\$36,051	\$0	\$36,051	\$0	\$36,051
	483	halumtano, Cebello (<i>Bulbophyllum guamense</i>) -Wherever found	T	\$58,551	\$0	\$58,551	\$0	\$58,551
	536	Harperella (<i>Ptilimnium nodosum</i>) -Wherever found	E	\$37,000	\$0	\$37,000	\$11,505	\$48,505
	1127	Hau kuahiwi (<i>Hibiscadelphus giffardianus</i>) -Wherever found	E	\$9,000	\$1,803	\$10,803	\$0	\$10,803
	1258	Hau kuahiwi (<i>Hibiscadelphus hualalaiensis</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1064	Hau kuahiwi (<i>Hibiscadelphus woodii</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	431	Heartleaf, dwarf-flowered (<i>Hexastylis naniflora</i>) -Wherever found	T	\$77,500	\$400	\$77,900	\$600	\$78,500
	821	Heather, mountain golden (<i>Hudsonia montana</i>) -Wherever found	T	\$20,500	\$0	\$20,500	\$0	\$20,500
	1064	Heau (<i>Exocarpos luteolus</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1064	Hibiscus, Clay's (<i>Hibiscus clayi</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1416	Higuero de sierra (<i>Crescentia portoricensis</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1132	Ho`awa (<i>Pittosporum napaliense</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1152	Hoawa (<i>Pittosporum halophilum</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	708	Hoawa (<i>Pittosporum hawaiiense</i>) -Wherever found	E	\$28,500	\$0	\$28,500	\$0	\$28,500
	1253	Holei (<i>Ochrosia haleakalae</i>) -Wherever found	E	\$7,500	\$1,803	\$9,303	\$0	\$9,303
	1362	Holei (<i>Ochrosia kilaueaensis</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1569	Holly, Cook's (<i>Ilex cookii</i>) -Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1150	Honohono (<i>Haplostachys haplostachya</i>) -Wherever found	E	\$8,500	\$1,803	\$10,303	\$0	\$10,303

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1362	Hulumoa (<i>Korthalsella degeneri</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	402	Iagu, Hayun (=Guam), Tronkon guafi (Rota)) (<i>Serianthes nelsonii</i>) -Wherever found	E	\$87,729	\$0	\$87,729	\$0	\$87,729
	479	Ihi (<i>Portulaca villosa</i>) -Wherever found	E	\$59,500	\$0	\$59,500	\$0	\$59,500
	1064	Iliau, dwarf (<i>Wilkesia hobbdi</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	622	Iris, dwarf lake (<i>Iris lacustris</i>) -Wherever found	T	\$36,600	\$0	\$36,600	\$0	\$36,600
	1602	Irisette, white (<i>Sisyrinchium dichotomum</i>) -Wherever found	E	\$300	\$0	\$300	\$1,600	\$1,900
	837	Ischaemum, Hilo (<i>Ischaemum byrone</i>) -Wherever found	E	\$9,500	\$10,241	\$19,741	\$0	\$19,741
	682	Ivesia, Ash Meadows (<i>Ivesia kingii</i> var. <i>eremica</i>) -Wherever found	T	\$100	\$30,000	\$30,100	\$0	\$30,100
	647	ivesia, Webber's (<i>Ivesia webberi</i>) -Wherever found	T	\$4,800	\$29,303	\$34,103	\$0	\$34,103
	455	Jewelflower, California (<i>Caulanthus californicus</i>) -Wherever found	E	\$13,000	\$54,583	\$67,583	\$0	\$67,583
	947	Jewelflower, Metcalf Canyon (<i>Streptanthus albidus</i> ssp. <i>albidus</i>) -Wherever found	E	\$14,900	\$0	\$14,900	\$0	\$14,900
	1624	Jewelflower, Tiburon (<i>Streptanthus niger</i>) -Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	543	Joint-vetch, Sensitive (<i>Aeschynomene virginica</i>) -Wherever found	T	\$34,000	\$12,096	\$46,096	\$800	\$46,896
	987	Kamakahala (<i>Labordia cyrtandrae</i>) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	1132	Kamakahala (<i>Labordia helleri</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1132	Kamakahala (<i>Labordia lydgatei</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1041	Kamakahala (<i>Labordia pumila</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1258	Kamakahala (<i>Labordia tinifolia</i> var. <i>lanaiensis</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1064	Kamakahala (<i>Labordia tinifolia</i> var. <i>wahiawaensis</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1292	Kamakahala (<i>Labordia triflora</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1258	Kamanomano (<i>Cenchrus agrimonoides</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	775	Kamapua`a (<i>Kadua fluviatilis</i>) -Wherever found	E	\$23,500	\$0	\$23,500	\$0	\$23,500
	1185	Kauai hau kuahiwi (<i>Hibiscadelphus distans</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	896	Kauila (<i>Colubrina oppositifolia</i>) -Wherever found	E	\$16,500	\$0	\$16,500	\$0	\$16,500
	878	Kaulu (<i>Pteralyxia kauaiensis</i>) -Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500
	1185	Kaulu (<i>Pteralyxia macrocarpa</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1292	Kio`ele (<i>Kadua coriacea</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	337	Kiponapona (<i>Phyllostegia racemosa</i>) -Wherever found	E	\$121,177	\$0	\$121,177	\$0	\$121,177
	1362	Ko`oko`olau (<i>Bidens amplexans</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	602	Ko`oko`olau (<i>Bidens campylotheca</i> ssp. <i>pentamera</i>) -Wherever found	E	\$8,000	\$30,724	\$38,724	\$0	\$38,724
	604	Ko`oko`olau (<i>Bidens campylotheca</i> ssp. <i>waihoiensis</i>) -Wherever found	E	\$7,500	\$30,724	\$38,224	\$0	\$38,224
	1338	Ko`oko`olau (<i>Bidens conjuncta</i>) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	961	Ko`oko`olau (<i>Bidens micrantha</i> ssp. <i>ctenophylla</i>) -Wherever found	E	\$14,000	\$409	\$14,409	\$0	\$14,409
	568	Ko`oko`olau (<i>Bidens micrantha</i> ssp. <i>kalealaha</i>) -Wherever found	E	\$13,500	\$30,724	\$44,224	\$0	\$44,224
	1362	Ko`oko`olau (<i>Bidens wiebkei</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	967	Ko`oloa`ula (<i>Abutilon menziesii</i>) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1152	Kohe malama malama o kanaloa (<i>Kanaloa kahoolawensis</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	Koholapehu (<i>Dubautia latifolia</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1258	Koki'o (<i>Kokia drynarioides</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1064	Koki'o (<i>Kokia kauaiensis</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1362	Koki'o ke'oke'o (<i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1026	Koki'o ke'oke'o (<i>Hibiscus waimeae</i> ssp. <i>hannerae</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1185	Koki'o, Cooke's (<i>Kokia cookei</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1041	Kolea (<i>Myrsine fosbergii</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1416	Kolea (<i>Myrsine juddii</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1185	Kolea (<i>Myrsine knudsenii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	848	Kolea (<i>Myrsine linearifolia</i>) -Wherever found	T	\$19,000	\$0	\$19,000	\$0	\$19,000
	1064	Kolea (<i>Myrsine mezii</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1152	Kolea (<i>Myrsine vaccinioides</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1185	kopa (<i>Kadua cordata remyi</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	967	Kopiko (<i>Psychotria grandiflora</i>) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1292	Kopiko (<i>Psychotria hexandra</i> ssp. <i>oahuensis</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1416	Kopiko (<i>Psychotria hobdyi</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1127	Kuahiwi laukahi (<i>Plantago hawaiiensis</i>) -Wherever found	E	\$9,000	\$1,803	\$10,803	\$0	\$10,803
	572	Kuahiwi laukahi (<i>Plantago princeps</i>) -Wherever found	E	\$12,500	\$30,724	\$43,224	\$0	\$43,224
	1041	Kuawawaenohu (<i>Schiedea lychnoides</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	868	Kula wahine noho (<i>Isodendrion pyrifolium</i>) -Wherever found	E	\$18,000	\$0	\$18,000	\$0	\$18,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1152	Kulu'i (Nototrichium humile) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	724	Ladies-tresses, Navasota (Spiranthes parksii) -Wherever found	E	\$27,000	\$0	\$27,000	\$0	\$27,000
	173	Ladies'-tresses, Ute (Spiranthes diluvialis) -Wherever found	T	\$254,733	\$215,570	\$470,303	\$2,800	\$473,103
	1255	Larkspur, Baker's (Delphinium bakeri) -Wherever found	E	\$9,200	\$0	\$9,200	\$0	\$9,200
	1473	Larkspur, San Clemente Island (Delphinium variegatum ssp. kinkiense) -Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1258	Larkspur, yellow (Delphinium luteum) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1185	Lau`ehu (Panicum niihauense) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1064	Laulihilihi (Schiedea stellarioides) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	503	Layia, beach (Layia carnosa) -Wherever found	T	\$53,500	\$2,000	\$55,500	\$0	\$55,500
	1258	Leather flower, Alabama (Clematis socialis) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1041	lehua makanoe (Lysimachia daphnoides) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	446	Lessingia, San Francisco (Lessingia germanorum (=L.g. var. germanorum)) -Wherever found	E	\$21,800	\$49,000	\$70,800	\$0	\$70,800
	1152	Liliwai (Acaena exigua) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	845	Lily, Minnesota dwarf trout (Erythronium propullans) -Wherever found	E	\$19,300	\$38	\$19,338	\$0	\$19,338
	702	Lily, Pitkin Marsh (Lilium pardalinum ssp. pitkinense) -Wherever found	E	\$29,000	\$0	\$29,000	\$0	\$29,000
	462	Lily, Western (Lilium occidentale) -Wherever found	E	\$35,600	\$29,000	\$64,600	\$0	\$64,600
	658	Liveforever, Santa Barbara Island (Dudleya traskiae) -Wherever found	E	\$2,500	\$30,000	\$32,500	\$0	\$32,500
	1329	Locoweed, Fassett's (Oxytropis campestris var. chartacea) -Wherever found	T	\$8,400	\$76	\$8,476	\$0	\$8,476

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	508	Lomatium, Cook's (Lomatium cookii) - Wherever found	E	\$54,707	\$0	\$54,707	\$0	\$54,707
	935	Loosestrife, rough-leaved (Lysimachia asperulaefolia) -Wherever found	E	\$12,000	\$0	\$12,000	\$3,000	\$15,000
	1473	lotus, San Clemente Island (Acmispon dendroideus var. traskiae) -Wherever found	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	452	Lousewort, Furbish (Pedicularis furbishiae) - Wherever found	E	\$68,000	\$0	\$68,000	\$0	\$68,000
	1292	Love grass, Fosberg's (Eragrostis fosbergii) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1536	Lupine, clover (Tidestrom"s) (Lupinus tidestromii) -Wherever found	E	\$2,200	\$1,000	\$3,200	\$0	\$3,200
	218	Lupine, Kincaid's (Lupinus sulphureus ssp. kincaidii) -Wherever found	T	\$148,511	\$192,710	\$341,221	\$0	\$341,221
	894	Lupine, Nipomo Mesa (Lupinus nipomensis) -Wherever found	E	\$16,800	\$0	\$16,800	\$0	\$16,800
	1002	Ma`o hau hele, (=native yellow hibiscus) (Hibiscus brackenridgei) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1041	Ma`oli`oli (Schiedea apokremnos) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	708	Ma`oli`oli (Schiedea hawaiiensis) - Wherever found	E	\$28,500	\$0	\$28,500	\$0	\$28,500
	1362	Ma`oli`oli (Schiedea kealiae) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1111	Ma`oli`oli (Schiedea pubescens) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1026	Mahoe (Alectryon macrococcus) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1023	Makou (Peucedanum sandwicense) - Wherever found	T	\$12,740	\$0	\$12,740	\$0	\$12,740
	352	Makou (Ranunculus hawaiiensis) -Wherever found	E	\$114,177	\$0	\$114,177	\$0	\$114,177
	1292	Makou (Ranunculus mauiensis) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1362	Malacothrix, island (Malacothrix squalida) - Wherever found	E	\$2,500	\$5,000	\$7,500	\$0	\$7,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1450	Malacothrix, Santa Cruz Island (Malacothrix indecora) -Wherever found	E	\$1,100	\$5,000	\$6,100	\$0	\$6,100
	1470	mallow, Gierisch (Sphaeralcea gierischii) -Wherever found	E	\$500	\$4,749	\$5,249	\$0	\$5,249
	1446	Mallow, Kern (Eremalche kernensis) -Wherever found	E	\$6,300	\$0	\$6,300	\$0	\$6,300
	1338	Manaca, palma de (Calyptronomia rivalis) -Wherever found	T	\$8,000	\$0	\$8,000	\$0	\$8,000
	907	Manioc, Walker's (Manihot walkerae) -Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	907	Manzanita, Franciscan (Arctostaphylos franciscana) -Wherever found	E	\$15,100	\$900	\$16,000	\$0	\$16,000
	1541	Manzanita, Ione (Arctostaphylos myrtifolia) -Wherever found	T	\$1,500	\$1,500	\$3,000	\$0	\$3,000
	867	Manzanita, Morro (Arctostaphylos morroensis) -Wherever found	T	\$18,000	\$100	\$18,100	\$0	\$18,100
	1663	Manzanita, pallid (Arctostaphylos pallida) -Wherever found	T	\$600	\$0	\$600	\$0	\$600
	1500	Manzanita, Presidio (Arctostaphylos hookeri var. ravenii) -Wherever found	E	\$3,600	\$700	\$4,300	\$0	\$4,300
	1399	Manzanita, Santa Rosa Island (Arctostaphylos confertiflora) -Wherever found	E	\$2,300	\$5,000	\$7,300	\$0	\$7,300
	1185	Mapele (Cyrtandra cyaneoides) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1624	Mariposa lily, Tiburon (Calochortus tiburonensis) -Wherever found	T	\$1,500	\$0	\$1,500	\$0	\$1,500
	906	Meadowfoam, Butte County (Limnanthes floccosa ssp. californica) -Wherever found	E	\$10,400	\$5,602	\$16,002	\$0	\$16,002
	473	meadowfoam, large-flowered woolly (Limnanthes pumila ssp. grandiflora) -Wherever found	E	\$61,500	\$0	\$61,500	\$0	\$61,500
	1183	Meadowfoam, Sebastopol (Limnanthes vinculans) -Wherever found	E	\$9,700	\$0	\$9,700	\$0	\$9,700
	1110	Meadowrue, Cooley's (Thalictrum cooleyi) -Wherever found	E	\$11,000	\$0	\$11,000	\$100	\$11,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1002	Mehamehame (Flueggea neowawraea) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	682	Mesa-mint, Otay (Pogogyne nudiuscula) - Wherever found	E	\$30,100	\$0	\$30,100	\$0	\$30,100
	1672	Mesa-mint, San Diego (Pogogyne abramsii) - Wherever found	E	\$300	\$0	\$300	\$0	\$300
	457	Milk-vetch, Applegate's (Astragalus applegatei) -Wherever found	E	\$67,200	\$0	\$67,200	\$0	\$67,200
	682	Milk-vetch, Ash meadows (Astragalus phoenix) -Wherever found	T	\$100	\$30,000	\$30,100	\$0	\$30,100
	527	Milk-vetch, Braunton's (Astragalus brauntonii) -Wherever found	E	\$48,500	\$1,000	\$49,500	\$0	\$49,500
	1624	Milk-vetch, Clara Hunt's (Astragalus clarianus) -Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	935	Milk-vetch, Coachella Valley (Astragalus lentiginosus var. coachellae) -Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	1500	Milk-vetch, Cushenbury (Astragalus albens) -Wherever found	E	\$4,300	\$0	\$4,300	\$0	\$4,300
	534	Milk-vetch, Fish Slough (Astragalus lentiginosus var. piscinensis) -Wherever found	T	\$2,900	\$46,000	\$48,900	\$0	\$48,900
	967	Milk-vetch, heliotrope (Astragalus montii) - Wherever found	T	\$14,000	\$0	\$14,000	\$0	\$14,000
	585	Milk-vetch, Holmgren (Astragalus holmgreniorum) -Wherever found	E	\$28,000	\$13,397	\$41,397	\$0	\$41,397
	859	Milk-vetch, Jesup's (Astragalus robbinsii var. jesupii) -Wherever found	E	\$18,500	\$0	\$18,500	\$0	\$18,500
	476	Milk-vetch, Lane Mountain (Astragalus jaegerianus) -Wherever found	E	\$60,000	\$0	\$60,000	\$0	\$60,000
	359	Milk-vetch, Mancos (Astragalus humillimus) -Wherever found	E	\$500	\$111,847	\$112,347	\$0	\$112,347
	424	milkvetch, Osterhout (Astragalus osterhoutii) -Wherever found	E	\$4,000	\$77,014	\$81,014	\$475	\$81,489
	592	Milk-vetch, Peirson's (Astragalus magdalenae var. peirsonii) -Wherever found	T	\$0	\$40,000	\$40,000	\$0	\$40,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1675	Milk-vetch, Sentry (Astragalus cremnophylax var. cremnophylax) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	558	Milk-vetch, Shivwits (Astragalus ampullarioides) -Wherever found	E	\$19,000	\$26,350	\$45,350	\$0	\$45,350
	405	Milk-vetch, triple-ribbed (Astragalus tricarinatus) -Wherever found	E	\$85,100	\$2,200	\$87,300	\$0	\$87,300
	552	Milk-vetch, Ventura Marsh (Astragalus pycnostachyus var. lanosissimus) -Wherever found	E	\$45,800	\$0	\$45,800	\$0	\$45,800
	418	Milkweed, Mead's (Asclepias meadii) - Wherever found	T	\$61,400	\$114	\$61,514	\$21,700	\$83,214
	489	Milkweed, Welsh's (Asclepias welshii) - Wherever found	T	\$12,000	\$45,200	\$57,200	\$0	\$57,200
	682	Monardella, willowy (Monardella viminea) - Wherever found	E	\$30,100	\$0	\$30,100	\$0	\$30,100
	745	monkey-flower, Michigan (Mimulus michiganensis) -Wherever found	E	\$24,000	\$1,200	\$25,200	\$0	\$25,200
	1036	Monkeyflower, Vandenberg (Diplacus vandenbergensis) -Wherever found	E	\$12,400	\$0	\$12,400	\$0	\$12,400
	576	Monkshood, northern wild (Aconitum noveboracense) -Wherever found	T	\$42,000	\$38	\$42,038	\$500	\$42,538
	1399	Morning-glory, Stebbins' (Calystegia stebbinsii) -Wherever found	E	\$1,500	\$5,800	\$7,300	\$0	\$7,300
	1563	mountainbalm, Indian Knob (Eriodictyon altissimum) -Wherever found	E	\$2,800	\$100	\$2,900	\$0	\$2,900
	746	Mustard, Penland alpine fen (Eutrema penlandii) -Wherever found	T	\$18,000	\$5,500	\$23,500	\$1,625	\$25,125
	1362	Na`ena`e (Dubautia herbstobatae) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1185	Na`ena`e (Dubautia imbricata ssp. imbricata) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1064	Na`ena`e (Dubautia pauciflora) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	Na`ena`e (Dubautia plantaginea ssp. humilis) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	Na`ena`e (Dubautia plantaginea ssp. magnifolia) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1064	Na`ena`e (Dubautia waialealae) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1064	Naenae (Dubautia kalalauensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1064	Naenae (Dubautia kenwoodii) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1292	Nani wai`ale`ale (Viola kauaiensis var. wahiawaensis) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1185	Nanu (Gardenia mannii) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	672	Nanu (Gardenia remyi) -Wherever found	E	\$20,500	\$10,241	\$30,741	\$0	\$30,741
	1026	Naupaka, dwarf (Scaevola coriacea) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1631	Navarretia, spreading (Navarretia fossalis) -Wherever found	T	\$1,300	\$100	\$1,400	\$0	\$1,400
	1185	nehe (Lipochaeta fauriei) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	nehe (Lipochaeta lobata var. leptophylla) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1064	nehe (Lipochaeta micrantha) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1362	nehe (Lipochaeta venosa) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1064	nehe (Lipochaeta waimeaensis) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	nehe (Melanthera kamolensis) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1002	nehe (Melanthera tenuifolia) -Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	952	Nioi (Eugenia koolauensis) -Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	540	Niterwort, Amargosa (Nitrophila mohavensis) -Wherever found	E	\$4,200	\$44,000	\$48,200	\$0	\$48,200
	1152	No common name (Abutilon eremitopetalum) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1362	No common name (Abutilon sandwicense) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	No common name (Achyranthes mutica) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1338	No common name (Agave eggersiana) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	715	No common name (Amaranthus brownii) - Wherever found	E	\$28,000	\$0	\$28,000	\$0	\$28,000
	1473	No common name (Aristida chaseae) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1451	No common name (Auerodendron pauciflorum) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	664	No common name (Bidens hillebrandiana ssp. hillebrandiana) -Wherever found	E	\$31,500	\$0	\$31,500	\$0	\$31,500
	828	No common name (Bonamia menziesii) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	1185	No common name (Calyptanthus thomasiana) -Wherever found	E	\$6,500	\$3,000	\$9,500	\$0	\$9,500
	1132	No common name (Catesbaea melanocarpa) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1292	No common name (Chamaecrista glandulosa var. mirabilis) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1507	No common name (Cordia bellonis) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1451	No common name (Cranichis ricartii) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1258	No common name (Cyanea kauaulaensis) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1338	No common name (Cyperus fauriei) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1152	No common name (Cyperus neokunthianus) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	689	No common name (Cyperus pennatiformis) - Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	1473	No common name (Daphnopsis helleriana) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	859	No common name (Delissea rhytidospema) -Wherever found	E	\$18,500	\$0	\$18,500	\$0	\$18,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	No common name (<i>Delissea undulata</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	345	No common name (<i>Dendrobium guamense</i>) - Wherever found	T	\$118,500	\$0	\$118,500	\$0	\$118,500
	809	No common name (<i>Eugenia bryanii</i>) - Wherever found	E	\$21,000	\$0	\$21,000	\$0	\$21,000
	1185	No common name (<i>Eugenia woodburyana</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1002	No common name (<i>Festuca hawaiiensis</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1338	No common name (<i>Festuca molokaiensis</i>) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	481	No common name (<i>Geocarpon minimum</i>) - Wherever found	T	\$57,000	\$0	\$57,000	\$2,000	\$59,000
	1541	No common name (<i>Gesneria pauciflora</i>) - Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	1416	No common name (<i>Gonocalyx concolor</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1362	No common name (<i>Gouania hillebrandii</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1185	No common name (<i>Gouania meyenii</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	967	No common name (<i>Gouania vitifolia</i>) - Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1152	No common name (<i>Hesperomannia arborescens</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1185	No common name (<i>Hesperomannia arbuscula</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1064	No common name (<i>Hesperomannia lydgatei</i>) -Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1569	No common name (<i>Ilex sintenisii</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1362	No common name (<i>Kadua degeneri</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1292	No common name (<i>Kadua haupuensis</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1362	No common name (<i>Kadua parvula</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1064	No common name (<i>Kadua st.-johnii</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1041	No common name (<i>Keysseria</i> (=Lagenifera) <i>erici</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1041	No common name (<i>Keysseria</i> (=Lagenifera) <i>helenae</i>) -Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1292	No common name (<i>Labordia lorenciana</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1111	No common name (<i>Lepidium orbiculare</i>) - Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1338	No common name (<i>Leptocereus grantianus</i>) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1416	No common name (<i>Lobelia koolauensis</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1152	No common name (<i>Lobelia monostachya</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1362	No common name (<i>Lobelia niihauensis</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	562	No common name (<i>Lobelia oahuensis</i>) - Wherever found	E	\$45,000	\$0	\$45,000	\$0	\$45,000
	1507	No common name (<i>Lyonia truncata</i> var. <i>proctorii</i>) -Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1416	No common name (<i>Lysimachia filifolia</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1064	No common name (<i>Lysimachia iniki</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	No common name (<i>Lysimachia lydgatei</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	967	No common name (<i>Lysimachia maxima</i>) - Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1064	No common name (<i>Lysimachia pendens</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	No common name (<i>Lysimachia scopulensis</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1185	No common name (<i>Lysimachia venosa</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	828	No common name (<i>Maesa walkeri</i>) - Wherever found	T	\$20,000	\$0	\$20,000	\$0	\$20,000
	1473	No common name (<i>Mitracarpus maxwelliae</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1111	No common name (<i>Mitracarpus polycladus</i>) - Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1402	No common name (<i>Myrcia paganii</i>) - Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1185	No common name (<i>Neraudia angulata</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	893	No common name (<i>Neraudia ovata</i>) - Wherever found	E	\$15,000	\$1,803	\$16,803	\$0	\$16,803
	1185	No common name (<i>Neraudia sericea</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	629	No common name (<i>Nervilia jacksoniae</i>) - Wherever found	T	\$36,000	\$0	\$36,000	\$0	\$36,000
	448	No common name (<i>Nesogenes rotensis</i>) - Wherever found	E	\$69,229	\$0	\$69,229	\$0	\$69,229
	447	No common name (<i>Osmoxylon mariannense</i>) - Wherever found	E	\$70,729	\$0	\$70,729	\$0	\$70,729
	530	No common name (<i>Phyllanthus saffordii</i>) - Wherever found	E	\$49,107	\$0	\$49,107	\$0	\$49,107
	494	No common name (<i>Phyllostegia bracteata</i>) - Wherever found	E	\$15,000	\$41,966	\$56,966	\$0	\$56,966
	307	No common name (<i>Phyllostegia brevidens</i>) - Wherever found	E	\$118,177	\$30,724	\$148,901	\$0	\$148,901
	583	No common name (<i>Phyllostegia floribunda</i>) - Wherever found	E	\$41,430	\$0	\$41,430	\$0	\$41,430
	1362	No common name (<i>Phyllostegia glabra</i> var. <i>lanaiensis</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	494	No common name (<i>Phyllostegia haliakalae</i>) - Wherever found	E	\$15,000	\$41,966	\$56,966	\$0	\$56,966
	1185	No common name (<i>Phyllostegia hirsuta</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	935	No common name (Phyllostegia hispida) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	1258	No common name (Phyllostegia kaalaensis) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1064	No common name (Phyllostegia knudsenii) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1152	No common name (Phyllostegia mannii) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1185	No common name (Phyllostegia mollis) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1041	No common name (Phyllostegia parviflora) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	935	No common name (Phyllostegia pilosa) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	1416	No common name (Phyllostegia renovans) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	935	No common name (Phyllostegia stachyoides) -Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	340	No common name (Phyllostegia velutina) - Wherever found	E	\$119,677	\$0	\$119,677	\$0	\$119,677
	1064	No common name (Phyllostegia waimeae) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1111	No common name (Phyllostegia warshaueri) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1064	No common name (Phyllostegia wawrana) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	967	No common name (Platanthera holochila) - Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	394	No common name (Platydesma cornuta var. cornuta) -Wherever found	E	\$95,000	\$0	\$95,000	\$0	\$95,000
	1362	No common name (Platydesma cornuta var. decurrens) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	654	No common name (Platydesma remyi) - Wherever found	E	\$33,000	\$0	\$33,000	\$0	\$33,000
	1185	No common name (Poa siphonoglossa) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	828	No common name (<i>Polyscias bisattenuata</i>) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	1064	No common name (<i>Polyscias flynnii</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1152	No common name (<i>Polyscias lydgatei</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1041	No common name (<i>Polyscias racemosa</i>) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1185	No common name (<i>Pritchardia aylmer-robinsonii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1064	No common name (<i>Pritchardia bakeri</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1185	No common name (<i>Pritchardia hardyi</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1152	No common name (<i>Pritchardia kaalae</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	828	No common name (<i>Pritchardia lanigera</i>) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	933	No common name (<i>Pritchardia maideniana</i>) - Wherever found	E	\$13,000	\$2,046	\$15,046	\$0	\$15,046
	1026	No common name (<i>Pritchardia munroi</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	935	No common name (<i>Pritchardia napaliensis</i>) - Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	501	No common name (<i>Pritchardia remota</i>) - Wherever found	E	\$56,000	\$0	\$56,000	\$0	\$56,000
	1002	No common name (<i>Pritchardia schattaueri</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	848	No common name (<i>Pritchardia viscosa</i>) - Wherever found	E	\$19,000	\$0	\$19,000	\$0	\$19,000
	1041	No common name (<i>Remya kauaiensis</i>) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1002	No common name (<i>Remya montgomeryi</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1292	No common name (<i>Sanicula mariversa</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1292	No common name (<i>Sanicula purpurea</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	604	No common name (<i>Sanicula sandwicensis</i>) - Wherever found	E	\$7,500	\$30,724	\$38,224	\$0	\$38,224
	878	No common name (<i>Santalum involutum</i>) - Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500
	1631	No common name (<i>Schiedea attenuata</i>) - Wherever found	E	\$1,400	\$0	\$1,400	\$0	\$1,400
	643	No common name (<i>Schiedea diffusa</i> ssp. <i>macraei</i>) -Wherever found	E	\$33,000	\$1,803	\$34,803	\$0	\$34,803
	547	No common name (<i>Schiedea diffusa</i> subsp. <i>diffusa</i>) -Wherever found	E	\$15,500	\$30,724	\$46,224	\$0	\$46,224
	547	No common name (<i>Schiedea haleakalensis</i>) - Wherever found	E	\$15,500	\$30,724	\$46,224	\$0	\$46,224
	1152	No common name (<i>Schiedea helleri</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1258	No common name (<i>Schiedea hookeri</i>) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	919	No common name (<i>Schiedea jacobii</i>) - Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	1064	No common name (<i>Schiedea kaalae</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	760	No common name (<i>Schiedea kauaiensis</i>) - Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	1111	No common name (<i>Schiedea laui</i>) - Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1362	No common name (<i>Schiedea lydgatei</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1041	No common name (<i>Schiedea membranacea</i>) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	878	No common name (<i>Schiedea nuttallii</i>) - Wherever found	E	\$17,500	\$0	\$17,500	\$0	\$17,500
	1132	No common name (<i>Schiedea obovata</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1258	No common name (<i>Schiedea salicaria</i>) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1292	No common name (<i>Schiedea sarmentosa</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1416	No common name (<i>Schiedea spergulina</i> var. <i>leiopoda</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1111	No common name (<i>Schiedea spergulina</i> var. <i>spergulina</i>) -Wherever found	T	\$11,000	\$0	\$11,000	\$0	\$11,000
	1362	No common name (<i>Schiedea trinervis</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	592	No common name (<i>Schiedea verticillata</i>) - Wherever found	E	\$40,000	\$0	\$40,000	\$0	\$40,000
	952	No common name (<i>Schiedea viscosa</i>) - Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	1292	No common name (<i>Schoepfia arenaria</i>) - Wherever found	T	\$8,500	\$0	\$8,500	\$0	\$8,500
	952	No common name (<i>Sicyos lanceoloideus</i>) - Wherever found	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	1132	No common name (<i>Silene alexandri</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	983	No common name (<i>Silene hawaiiensis</i>) - Wherever found	T	\$13,930	\$0	\$13,930	\$0	\$13,930
	1258	No common name (<i>Silene lanceolata</i>) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1132	No common name (<i>Silene perlmanii</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1258	No common name (<i>Spermolepis</i> <i>hawaiiensis</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	930	No common name (<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>) -Wherever found	E	\$13,500	\$1,803	\$15,303	\$0	\$15,303
	919	No common name (<i>Stenogyne bifida</i>) - Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	1292	No common name (<i>Stenogyne campanulata</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	664	No common name (<i>Stenogyne cranwelliae</i>) - Wherever found	E	\$31,500	\$0	\$31,500	\$0	\$31,500
	1185	No common name (<i>Stenogyne kaalae</i> ssp. <i>sherffii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1064	No common name (<i>Stenogyne kanehoana</i>) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	1152	No common name (<i>Stenogyne kauaulaensis</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1292	No common name (<i>Stenogyne kealiae</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1473	No common name (<i>Tabernaemontana</i> <i>rotensis</i>) -Wherever found	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	1569	No common name (<i>Ternstroemia</i> <i>subsessilis</i>) -Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1258	No common name (<i>Tetramolopium</i> <i>arenarium</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1362	No common name (<i>Tetramolopium</i> <i>filiforme</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1152	No common name (<i>Tetramolopium</i> <i>lepidotum</i> ssp. <i>lepidotum</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1132	No common name (<i>Tetramolopium remyi</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1132	No common name (<i>Tetramolopium rockii</i>) - Wherever found	T	\$10,500	\$0	\$10,500	\$0	\$10,500
	786	No common name (<i>Tinospora homosepala</i>) - Wherever found	E	\$22,500	\$0	\$22,500	\$0	\$22,500
	1362	No common name (<i>Trematolobelia</i> <i>singularis</i>) -Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	472	No common name (<i>Tuberolabium</i> <i>guamense</i>) -Wherever found	T	\$61,700	\$0	\$61,700	\$0	\$61,700
	521	No common name (<i>Varronia rupicola</i>) - Wherever found	T	\$52,500	\$0	\$52,500	\$0	\$52,500
	1507	No common name (<i>Vernonia proctorii</i>) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1185	No common name (<i>Vigna o-wahuensis</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1292	No common name (<i>Viola helenae</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1152	No common name (<i>Viola lanaiensis</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1292	No common name (<i>Viola oahuensis</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	806	No common name (<i>Wikstroemia villosa</i>) - Wherever found	E	\$11,000	\$10,241	\$21,241	\$0	\$21,241
	1002	No common name (<i>Xylosma crenatum</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	590	Nohoanu (<i>Geranium arboreum</i>) -Wherever found	E	\$9,500	\$30,724	\$40,224	\$0	\$40,224
	602	Nohoanu (<i>Geranium hanaense</i>) -Wherever found	E	\$8,000	\$30,724	\$38,724	\$0	\$38,724
	1185	Nohoanu (<i>Geranium hillebrandii</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	987	Nohoanu (<i>Geranium kauaiense</i>) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	875	Nohoanu (<i>Geranium multiflorum</i>) - Wherever found	E	\$7,500	\$10,241	\$17,741	\$0	\$17,741
	825	nui, haha (<i>Cyanea horrida</i>) -Wherever found	E	\$10,000	\$10,241	\$20,241	\$0	\$20,241
	1185	Oha (<i>Delissea subcordata</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	390	Ohai (<i>Sesbania tomentosa</i>) -Wherever found	E	\$94,500	\$1,803	\$96,303	\$0	\$96,303
	935	Olulu (<i>Brighamia insignis</i>) -Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	689	Onion, Munz's (<i>Allium munzii</i>) -Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	544	Opuhe (<i>Urera kaalae</i>) -Wherever found	E	\$46,500	\$0	\$46,500	\$0	\$46,500
	298	Orchid, eastern prairie fringed (<i>Platanthera leucophaea</i>) -Wherever found	T	\$151,500	\$38	\$151,538	\$1,100	\$152,638
	294	Orchid, western prairie fringed (<i>Platanthera praeclara</i>) -Wherever found	T	\$113,300	\$22,339	\$135,639	\$22,500	\$158,139
	1624	Orcutt grass, California (<i>Orcuttia californica</i>) -Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	915	Orcutt grass, hairy (<i>Orcuttia pilosa</i>) - Wherever found	E	\$10,200	\$5,602	\$15,802	\$0	\$15,802
	866	Orcutt grass, Sacramento (<i>Orcuttia viscida</i>) - Wherever found	E	\$11,600	\$6,547	\$18,147	\$0	\$18,147
	1337	Orcutt grass, San Joaquin Valley (<i>Orcuttia inaequalis</i>) -Wherever found	T	\$1,600	\$6,547	\$8,147	\$0	\$8,147

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	674	Orcutt grass, slender (<i>Orcuttia tenuis</i>) - Wherever found	T	\$25,100	\$5,602	\$30,702	\$0	\$30,702
	781	Owl's-clover, fleshy (<i>Castilleja campestris</i> ssp. <i>succulenta</i>) -Wherever found	T	\$17,300	\$5,602	\$22,902	\$0	\$22,902
	346	oxytheca, Cushenbury (<i>Oxytheca parishii</i> var. <i>goodmaniana</i>) -Wherever found	E	\$118,100	\$0	\$118,100	\$0	\$118,100
	1026	Pa`iniu (<i>Astelia waialealae</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1680	Paintbrush, ash-grey (<i>Castilleja cinerea</i>) - Wherever found	T	\$100	\$0	\$100	\$0	\$100
	180	paintbrush, golden (<i>Castilleja levisecta</i>) - Wherever found	T	\$443,217	\$4,000	\$447,217	\$0	\$447,217
	1473	Paintbrush, San Clemente Island (<i>Castilleja</i> <i>grisea</i>) -Wherever found	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	398	Paintbrush, soft-leaved (<i>Castilleja mollis</i>) - Wherever found	E	\$73,300	\$20,000	\$93,300	\$0	\$93,300
	1255	Paintbrush, Tiburon (<i>Castilleja affinis</i> ssp. <i>neglecta</i>) -Wherever found	E	\$8,700	\$500	\$9,200	\$0	\$9,200
	1541	Palo colorado (<i>Ternstroemia luquillensis</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1541	Palo de jazmin (<i>Styrax portoricensis</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1541	Palo de nigua (<i>Cornutia obovata</i>) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1451	Palo de ramon (<i>Banara vanderbiltii</i>) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	952	Palo de rosa (<i>Ottoschulzia rhodoxylon</i>) - Wherever found	T	\$14,500	\$0	\$14,500	\$0	\$14,500
	1132	Pamakani (<i>Tetramolopium capillare</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1185	Pamakani (<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1338	Panicgrass, Carter's (<i>Panicum fauriei</i> var. <i>carteri</i>) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1111	Papala (<i>Charpentiera densiflora</i>) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	535	Pauddo (Hedyotis megalantha) -Wherever found	E	\$48,607	\$0	\$48,607	\$0	\$48,607
	1473	Pelos del diablo (Aristida portoricensis) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1338	Penny-cress, Kneeland Prairie (Thlaspi californicum) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1129	Pennyroyal, Todsens (Hedeoma todsonii) - Wherever found	E	\$0	\$10,750	\$10,750	\$0	\$10,750
	315	Penstemon, blowout (Penstemon haydenii) - Wherever found	E	\$34,600	\$93,539	\$128,139	\$9,000	\$137,139
	1064	Pentachaeta, Lyon's (Pentachaeta lyonii) - Wherever found	E	\$4,000	\$7,500	\$11,500	\$0	\$11,500
	1527	Peperomia, Wheeler's (Peperomia wheeleri) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1000	Phacelia, clay (Phacelia argillacea) - Wherever found	E	\$11,000	\$2,060	\$13,060	\$0	\$13,060
	425	Phacelia, DeBeque (Phacelia submutica) - Wherever found	T	\$48,000	\$30,000	\$78,000	\$3,175	\$81,175
	1334	Phacelia, island (Phacelia insularis ssp. insularis) -Wherever found	E	\$1,300	\$7,000	\$8,300	\$0	\$8,300
	488	Phacelia, North Park (Phacelia formosula) - Wherever found	E	\$4,000	\$50,847	\$54,847	\$2,700	\$57,547
	609	Phlox, Texas trailing (Phlox nivalis ssp. texensis) -Wherever found	E	\$17,000	\$2,100	\$19,100	\$19,000	\$38,100
	654	phlox, Yreka (Phlox hirsuta) -Wherever found	E	\$33,000	\$0	\$33,000	\$0	\$33,000
	1111	pilo (Kadua laxiflora) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	1185	Pilo kea lau li'i (Platydesma rostrata) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	828	Piperia, Yadon's (Piperia yadonii) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	1607	Pitcher-plant, Alabama canebrake (Sarracenia rubra ssp. alabamensis) - Wherever found	E	\$1,800	\$0	\$1,800	\$0	\$1,800
	907	Pitcher-plant, green (Sarracenia oreophila) - Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1330	Pitcher-plant, mountain sweet (<i>Sarracenia rubra</i> ssp. <i>jonesii</i>) -Wherever found	E	\$5,500	\$400	\$5,900	\$2,500	\$8,400
	931	Po'e (<i>Portulaca sclerocarpa</i>) -Wherever found	E	\$13,430	\$1,803	\$15,233	\$0	\$15,233
	758	Polygonum, Scotts Valley (<i>Polygonum hickmanii</i>) -Wherever found	E	\$24,600	\$0	\$24,600	\$0	\$24,600
	492	Pondberry (<i>Lindera melissifolia</i>) -Wherever found	E	\$49,500	\$0	\$49,500	\$7,500	\$57,000
	372	popcornflower, rough (<i>Plagiobothrys hirtus</i>) -Wherever found	E	\$64,810	\$43,156	\$107,966	\$0	\$107,966
	1185	Popolo (<i>Cyanea solanacea</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	404	Popolo (<i>Solanum nelsonii</i>) -Wherever found	E	\$87,500	\$0	\$87,500	\$0	\$87,500
	1185	Popolo ku mai (<i>Solanum incompletum</i>) -Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	987	Poppy, Sacramento prickly (<i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i>) -Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	1451	Poppy-mallow, Texas (<i>Callirhoe scabriuscula</i>) -Wherever found	E	\$0	\$0	\$0	\$6,000	\$6,000
	1622	Potentilla, Hickman's (<i>Potentilla hickmanii</i>) -Wherever found	E	\$0	\$1,600	\$1,600	\$0	\$1,600
	987	Prickly-ash, St. Thomas (<i>Zanthoxylum thomasianum</i>) -Wherever found	E	\$10,500	\$3,000	\$13,500	\$0	\$13,500
	1041	Primrose, Maguire (<i>Primula maguirei</i>) -Wherever found	T	\$12,000	\$0	\$12,000	\$0	\$12,000
	1258	Pu'uka'a (<i>Cyperus trachysanthos</i>) -Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1152	Pua `ala (<i>Brighamia rockii</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1338	Pussypaws, Mariposa (<i>Calyptridium pulchellum</i>) -Wherever found	T	\$8,000	\$0	\$8,000	\$0	\$8,000
	1464	Rattleweed, hairy (<i>Baptisia arachnifera</i>) -Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	1338	Reedgrass, Hillebrand's (<i>Calamagrostis hillebrandii</i>) -Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	857	Reedgrass, Maui (<i>Calamagrostis expansa</i>) - Wherever found	E	\$8,500	\$10,241	\$18,741	\$0	\$18,741
	965	Reed-mustard, Barneby (<i>Schoenocrambe barnebyi</i>) -Wherever found	E	\$6,000	\$8,237	\$14,237	\$0	\$14,237
	985	Reed-mustard, clay (<i>Schoenocrambe argillacea</i>) -Wherever found	T	\$13,000	\$691	\$13,691	\$0	\$13,691
	886	Reed-mustard, shrubby (<i>Schoenocrambe suffrutescens</i>) -Wherever found	E	\$12,000	\$5,486	\$17,486	\$0	\$17,486
	1292	Remya, Maui (<i>Remya mauiensis</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	848	Ridge-cress, Barneby (<i>Lepidium barnebyanum</i>) -Wherever found	E	\$19,000	\$0	\$19,000	\$0	\$19,000
	586	rock cress, Shale barren (<i>Boechera serotina</i>) - Wherever found	E	\$35,000	\$0	\$35,000	\$6,301	\$41,301
	652	Rock-cress, Braun's (<i>Arabis perstellata</i>) - Wherever found	E	\$33,500	\$0	\$33,500	\$0	\$33,500
	1602	Rockcress, Georgia (<i>Arabis georgiana</i>) - Wherever found	T	\$1,900	\$0	\$1,900	\$0	\$1,900
	878	Rock-cress, Hoffmann's (<i>Arabis hoffmannii</i>) -Wherever found	E	\$2,500	\$15,000	\$17,500	\$0	\$17,500
	918	Rock-cress, McDonald's (<i>Arabis macdonaldiana</i>) -Wherever found	E	\$15,551	\$0	\$15,551	\$0	\$15,551
	343	Rockcress, Santa Cruz Island (<i>Sibara filifolia</i>) -Wherever found	E	\$118,700	\$0	\$118,700	\$0	\$118,700
	1525	Rose-mallow, Neches River (<i>Hibiscus dasycalyx</i>) -Wherever found	T	\$3,800	\$0	\$3,800	\$0	\$3,800
	783	Rosemary, Cumberland (<i>Conradina verticillata</i>) -Wherever found	T	\$13,500	\$9,200	\$22,700	\$0	\$22,700
	556	roseroot, Leedy's (<i>Rhodiola integrifolia</i> ssp. leedyi) -Wherever found	T	\$45,400	\$38	\$45,438	\$0	\$45,438
	967	Rush-pea, slender (<i>Hoffmannseggia tenella</i>) - Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	1569	Rush-rose, island (<i>Helianthemum greenei</i>) - Wherever found	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	1132	Sandalwood, Lanai (= `iliah) (<i>Santalum haleakalae</i> var. <i>lanaiense</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1111	Sand-verbena, large-fruited (<i>Abronia macrocarpa</i>) -Wherever found	E	\$11,000	\$0	\$11,000	\$0	\$11,000
	627	Sandwort, Marsh (<i>Arenaria paludicola</i>) -Wherever found	E	\$35,600	\$500	\$36,100	\$0	\$36,100
	1152	Schiedea, Diamond Head (<i>Schiedea adamantis</i>) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	664	Sedge, golden (<i>Carex lutea</i>) -Wherever found	E	\$31,500	\$0	\$31,500	\$0	\$31,500
	966	Sedge, Navajo (<i>Carex specuicola</i>) -Wherever found	T	\$1,000	\$13,133	\$14,133	\$0	\$14,133
	498	Silversword, Mauna Loa (=Ka'u) (<i>Argyroxiphium kauense</i>) -Wherever found	E	\$11,500	\$45,000	\$56,500	\$0	\$56,500
	444	Skullcap, large-flowered (<i>Scutellaria montana</i>) -Wherever found	T	\$71,500	\$0	\$71,500	\$0	\$71,500
	739	Skyrocket, Pagosa (<i>Ipomopsis polyantha</i>) -Wherever found	E	\$21,000	\$0	\$21,000	\$4,650	\$25,650
	623	Sneezeweed, Virginia (<i>Helenium virginicum</i>) -Wherever found	T	\$36,500	\$76	\$36,576	\$0	\$36,576
	964	Spineflower, Ben Lomond (<i>Chorizanthe pungens</i> var. <i>hartwegiana</i>) -Wherever found	E	\$14,300	\$0	\$14,300	\$0	\$14,300
	935	Spineflower, Howell's (<i>Chorizanthe howellii</i>) -Wherever found	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	625	Spineflower, Monterey (<i>Chorizanthe pungens</i> var. <i>pungens</i>) -Wherever found	T	\$36,400	\$0	\$36,400	\$0	\$36,400
	747	Spineflower, Orcutt's (<i>Chorizanthe orcuttiana</i>) -Wherever found	E	\$25,100	\$0	\$25,100	\$0	\$25,100
	917	spineflower, Robust (<i>Chorizanthe robusta</i> var. <i>robusta</i>) -Wherever found	E	\$15,600	\$0	\$15,600	\$0	\$15,600
	758	spineflower, Scotts Valley (<i>Chorizanthe robusta</i> var. <i>hartwegii</i>) -Wherever found	E	\$24,600	\$0	\$24,600	\$0	\$24,600
	668	Spineflower, slender-horned (<i>Dodecahema leptoceras</i>) -Wherever found	E	\$31,200	\$0	\$31,200	\$0	\$31,200
	1569	Spineflower, Sonoma (<i>Chorizanthe valida</i>) -Wherever found	E	\$1,500	\$1,000	\$2,500	\$0	\$2,500
	381	Spiraea, Virginia (<i>Spiraea virginiana</i>) -Wherever found	T	\$44,500	\$54,966	\$99,466	\$2,509	\$101,975

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1020	Spurge, Hoover's (<i>Chamaesyce hooveri</i>) - Wherever found	T	\$7,300	\$5,602	\$12,902	\$0	\$12,902
	524	Stickseed, showy (<i>Hackelia venusta</i>) - Wherever found	E	\$51,042	\$0	\$51,042	\$0	\$51,042
	1473	stonecrop, Bartram's (<i>Graptopetalum bartramii</i>) -Wherever found	T	\$0	\$5,000	\$5,000	\$0	\$5,000
	573	Sumac, Michaux's (<i>Rhus michauxii</i>) - Wherever found	E	\$41,000	\$400	\$41,400	\$1,600	\$43,000
	1415	Sunburst, Hartweg's golden (<i>Pseudobahia bahiifolia</i>) -Wherever found	E	\$0	\$6,547	\$6,547	\$0	\$6,547
	1062	Sunburst, San Joaquin adobe (<i>Pseudobahia peirsonii</i>) -Wherever found	T	\$11,700	\$0	\$11,700	\$0	\$11,700
	1254	Sunflower, Pecos (=puzzle, =paradox) (<i>Helianthus paradoxus</i>) -Wherever found	T	\$0	\$9,250	\$9,250	\$0	\$9,250
	1335	Sunflower, San Mateo woolly (<i>Eriophyllum latilobum</i>) -Wherever found	E	\$8,200	\$0	\$8,200	\$0	\$8,200
	433	Sunflower, Schweinitz's (<i>Helianthus schweinitzii</i>) -Wherever found	E	\$71,000	\$0	\$71,000	\$6,500	\$77,500
	682	Sunray, Ash Meadows (<i>Enceliopsis nudicaulis</i> var. <i>corrugata</i>) -Wherever found	T	\$100	\$30,000	\$30,100	\$0	\$30,100
	1183	Sunshine, Sonoma (<i>Blennosperma bakeri</i>) - Wherever found	E	\$9,700	\$0	\$9,700	\$0	\$9,700
	1680	Taraxacum, California (<i>Taraxacum californicum</i>) -Wherever found	E	\$100	\$0	\$100	\$0	\$100
	491	Tarplant, Gaviota (<i>Deinandra increscens</i> ssp. <i>villosa</i>) -Wherever found	E	\$57,100	\$0	\$57,100	\$0	\$57,100
	569	Tarplant, Santa Cruz (<i>Holocarpha macradenia</i>) -Wherever found	T	\$43,900	\$0	\$43,900	\$0	\$43,900
	553	Thelypody, Howell's spectacular (<i>Thelypodium howellii</i> ssp. <i>spectabilis</i>) - Wherever found	T	\$45,500	\$0	\$45,500	\$0	\$45,500
	1536	Thistle, Chorro Creek bog (<i>Cirsium fontinale</i> var. <i>obispoense</i>) -Wherever found	E	\$3,100	\$100	\$3,200	\$0	\$3,200
	1335	Thistle, fountain (<i>Cirsium fontinale</i> var. <i>fontinale</i>) -Wherever found	E	\$8,200	\$0	\$8,200	\$0	\$8,200
	577	Thistle, La Graciosa (<i>Cirsium loncholepis</i>) - Wherever found	E	\$42,000	\$0	\$42,000	\$0	\$42,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	420	Thistle, Pitcher's (<i>Cirsium pitcheri</i>) - Wherever found	T	\$61,300	\$20,438	\$81,738	\$300	\$82,038
	614	Thistle, Suisun (<i>Cirsium hydrophilum</i> var. <i>hydrophilum</i>) -Wherever found	E	\$37,500	\$0	\$37,500	\$0	\$37,500
	639	Thornmint, San Diego (<i>Acanthomintha</i> <i>ilicifolia</i>) -Wherever found	T	\$35,100	\$0	\$35,100	\$0	\$35,100
	916	Townsendia, Last Chance (<i>Townsendia</i> <i>aprica</i>) -Wherever found	T	\$10,000	\$5,639	\$15,639	\$0	\$15,639
	1507	Trillium, persistent (<i>Trillium persistens</i>) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	516	Trillium, relict (<i>Trillium reliquum</i>) - Wherever found	E	\$53,500	\$0	\$53,500	\$0	\$53,500
	1024	Tuctoria, Greene's (<i>Tuctoria greenei</i>) - Wherever found	E	\$12,700	\$0	\$12,700	\$0	\$12,700
	903	Twinpod, Dudley Bluffs (<i>Physaria</i> <i>obcordata</i>) -Wherever found	T	\$11,000	\$5,000	\$16,000	\$425	\$16,425
	515	Ufa-halomtano (<i>Heritiera longipetiolata</i>) - Wherever found	E	\$53,607	\$0	\$53,607	\$0	\$53,607
	840	Uhi uhi (<i>Mezoneuron kawaiense</i>) -Wherever found	E	\$19,500	\$0	\$19,500	\$0	\$19,500
	1451	Uvillo (<i>Eugenia haematocarpa</i>) -Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1534	Vervain, Red Hills (<i>Verbena californica</i>) - Wherever found	T	\$2,600	\$800	\$3,400	\$0	\$3,400
	891	Vetch, Hawaiian (<i>Vicia menziesii</i>) - Wherever found	E	\$17,000	\$0	\$17,000	\$0	\$17,000
	1358	Wallflower, Ben Lomond (<i>Erysimum</i> <i>teretifolium</i>) -Wherever found	E	\$7,900	\$0	\$7,900	\$0	\$7,900
	1289	Wallflower, Contra Costa (<i>Erysimum</i> <i>capitatum</i> var. <i>angustatum</i>) -Wherever found	E	\$8,700	\$0	\$8,700	\$0	\$8,700
	736	Wallflower, Menzies' (<i>Erysimum menziesii</i>) -Wherever found	E	\$25,800	\$0	\$25,800	\$0	\$25,800
	1507	Walnut (=Nogal), West Indian (<i>Juglans</i> <i>jamaicensis</i>) -Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	511	Watercress, Gambel's (<i>Rorippa gambellii</i>) - Wherever found	E	\$54,200	\$0	\$54,200	\$0	\$54,200

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1541	Water-plantain, Kral's (<i>Sagittaria secundifolia</i>) -Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	512	wild buckwheat, clay-loving (<i>Eriogonum pelinophilum</i>) -Wherever found	E	\$26,500	\$26,432	\$52,932	\$1,150	\$54,082
	394	Wild-buckwheat, gypsum (<i>Eriogonum gypsophilum</i>) -Wherever found	T	\$0	\$95,000	\$95,000	\$0	\$95,000
	509	Wild-buckwheat, southern mountain (<i>Eriogonum kennedyi</i> var. <i>austromontanum</i>) -Wherever found	T	\$54,500	\$0	\$54,500	\$0	\$54,500
	1589	Wire-lettuce, Malheur (<i>Stephanomeria malheurensis</i>) -Wherever found	E	\$2,200	\$0	\$2,200	\$0	\$2,200
	235	Wooly-threads, San Joaquin (<i>Monolopia</i> (=Lembertia) <i>congdonii</i>) -Wherever found	E	\$58,600	\$209,493	\$268,093	\$0	\$268,093
	821	Yellowhead, desert (<i>Yermo xanthocephalus</i>) -Wherever found	T	\$5,000	\$15,500	\$20,500	\$0	\$20,500
	905	Yerba santa, Lompoc (<i>Eriodictyon capitatum</i>) -Wherever found	E	\$16,100	\$0	\$16,100	\$0	\$16,100
	1495	Bacora, marron (<i>Solanum conocarpum</i>) -	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1507	Rosemary, short-leaved (<i>Conradina brevifolia</i>) -	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	741	Rosemary, Apalachicola (<i>Conradina glabra</i>) -	E	\$25,500	\$0	\$25,500	\$0	\$25,500
	1416	Blazingstar, scrub (<i>Liatris ohlingerae</i>) -	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	913	Birds-in-a-nest, white (<i>Macbridea alba</i>) -	T	\$15,500	\$330	\$15,830	\$0	\$15,830
	1637	Pinkroot, gentian (<i>Spigelia gentianoides</i>) -	E	\$1,100	\$0	\$1,100	\$0	\$1,100
	1416	Spurge, deltoid (<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>) -	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1416	Fringe-tree, pygmy (<i>Chionanthus pygmaeus</i>) -	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	702	Aster, Florida golden (<i>Chrysopsis floridana</i>) -	E	\$29,000	\$0	\$29,000	\$0	\$29,000
	1507	Pigeon wings (<i>Clitoria fragrans</i>) -	T	\$4,000	\$0	\$4,000	\$0	\$4,000
	840	Gourd, Okeechobee (<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>) -	E	\$19,500	\$0	\$19,500	\$0	\$19,500
	353	Prairie-clover, leafy (<i>Dalea foliosa</i>) -	E	\$114,000	\$0	\$114,000	\$0	\$114,000
	1473	Beargrass, Britton's (<i>Nolina brittoniana</i>) -	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1002	No common name (<i>Phyllostegia helleri</i>) -	E	\$13,000	\$0	\$13,000	\$0	\$13,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1635	Butterwort, Godfrey's (<i>Pinguicula ionantha</i>) -	T	\$900	\$330	\$1,230	\$0	\$1,230
	270	Orchid, white fringeless (<i>Platanthera integrilabia</i>) -	T	\$198,500	\$0	\$198,500	\$0	\$198,500
	1569	Lead-plant, Crenulate (<i>Amorpha crenulata</i>) -	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1647	Milkpea, Small's (<i>Galactia smallii</i>) -	E	\$900	\$0	\$900	\$0	\$900
	1569	Mint, Garrett's (<i>Dicerandra christmanii</i>) -	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	634	leather flower, Morefield's (<i>Clematis morefieldii</i>) -	E	\$35,500	\$0	\$35,500	\$0	\$35,500
	1495	Harebells, Avon Park (<i>Crotalaria avonensis</i>) -	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	406	potato-bean, Price's (<i>Apios priceana</i>) -	T	\$81,000	\$6,000	\$87,000	\$0	\$87,000
	1507	Mint, scrub (<i>Dicerandra frutescens</i>) -	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1507	Mint, Lakela's (<i>Dicerandra immaculata</i>) -	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	740	Crabgrass, Florida pineland (<i>Digitaria pauciflora</i>) -	T	\$15,000	\$10,562	\$25,562	\$0	\$25,562
	1416	Whitlow-wort, papery (<i>Paronychia chartacea</i>) -	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	932	Grass, Eureka Dune (<i>Swallenia alexandrae</i>) -	T	\$100	\$15,000	\$15,100	\$0	\$15,100
	1569	Pawpaw, beautiful (<i>Deeringothamnus pulchellus</i>) -	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1473	Pawpaw, Rugel's (<i>Deeringothamnus rugelii</i>) -	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1569	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) -	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	1402	Snakeroot (<i>Eryngium cuneifolium</i>) -	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	486	Spurge, telephus (<i>Euphorbia telephioides</i>) -	T	\$58,000	\$0	\$58,000	\$0	\$58,000
	1569	Polygala, tiny (<i>Polygala smallii</i>) -	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	144	Peppergrass, Slickspot (<i>Lepidium papilliferum</i>) -	T	\$164,099	\$449,750	\$613,849	\$9,000	\$622,849
	821	Sandmat, pineland (<i>Chamaesyce deltoidea</i> <i>pinetorum</i>) -	T	\$20,500	\$0	\$20,500	\$0	\$20,500
	1587	Thoroughwort, Cape Sable (<i>Chromolaena frustrata</i>) -	E	\$900	\$1,500	\$2,400	\$0	\$2,400

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1402	Prairie-clover, Florida (<i>Dalea carthagenensis floridana</i>) -	E	\$5,000	\$2,000	\$7,000	\$0	\$7,000
	950	Sunflower, Whorled (<i>Helianthus verticillatus</i>) -	E	\$13,500	\$1,200	\$14,700	\$0	\$14,700
	1473	Cactus, Florida semaphore (<i>Consolea corallicola</i>) -	E	\$3,000	\$2,000	\$5,000	\$0	\$5,000
	651	Bully, Everglades (<i>Sideroxylon reclinatum ssp. austrofloridense</i>) -	T	\$19,000	\$14,562	\$33,562	\$0	\$33,562
	962	Heau (<i>Exocarpos menziesii</i>) -	E	\$12,500	\$1,803	\$14,303	\$0	\$14,303
	1506	eryngo, Arizona (<i>Eryngium sparganophyllum</i>) -	E	\$0	\$4,120	\$4,120	\$0	\$4,120
	1473	Bonamia, Florida (<i>Bonamia grandiflora</i>) -	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	274	Beauty, Harper's (<i>Harperocallis flava</i>) -	E	\$191,500	\$0	\$191,500	\$0	\$191,500
	1152	Cactus, Key tree (<i>Pilosocereus robinii</i>) -	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1473	Polygala, Lewton's (<i>Polygala lewtonii</i>) -	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	573	Seablite, California (<i>Suaeda californica</i>) -	E	\$43,000	\$0	\$43,000	\$0	\$43,000
	1591	Rosemary, Etonia (<i>Conradina etonia</i>) -	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	1185	No common name (<i>Wikstroemia skottsbergiana</i>) -	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1647	Mint, longspurred (<i>Dicerandra cornutissima</i>) -	E	\$900	\$0	\$900	\$0	\$900
	799	Lupine, scrub (<i>Lupinus aridorum</i>) -	E	\$21,500	\$0	\$21,500	\$0	\$21,500
	1647	Bellflower, Brooksville (<i>Campanula robinsiae</i>) -	E	\$900	\$0	\$900	\$0	\$900
	1527	Prickly-apple, fragrant (<i>Cereus eriophorus</i> var. <i>fragrans</i>) -	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1591	Spurge, Garber's (<i>Chamaesyce garberi</i>) -	T	\$2,000	\$0	\$2,000	\$0	\$2,000
	320	Pink, swamp (<i>Helonias bullata</i>) -	T	\$105,500	\$25,486	\$130,986	\$2,000	\$132,986
	1647	Jacquemontia, beach (<i>Jacquemontia reclinata</i>) -	E	\$900	\$0	\$900	\$0	\$900
	1507	Hypericum, highlands scrub (<i>Hypericum cumulicola</i>) -	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	230	Pogonia, small whorled (<i>Isotria medeoloides</i>) -	T	\$259,000	\$38,000	\$297,000	\$300	\$297,300
	1402	Water-willow, Cooley's (<i>Justicia cooleyi</i>) -	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	741	silverbush, Blodgett's (<i>Argythamnia blodgettii</i>) -	T	\$25,500	\$0	\$25,500	\$0	\$25,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	459	Skullcap, Florida (Scutellaria floridana) -	T	\$65,500	\$330	\$65,830	\$0	\$65,830
	896	Warea, wide-leaf (Warea amplexifolia) -	E	\$16,500	\$0	\$16,500	\$0	\$16,500
	1402	Mustard, Carter's (Warea carteri) -	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	840	Flax, sand (Linum arenicola) -	E	\$19,500	\$0	\$19,500	\$0	\$19,500
	1185	Flax, Carter's small-flowered (Linum carteri carteri) -	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1402	Wireweed (Polygonella basiramia) -	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1185	Sandlace (Polygonella myriophylla) -	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1362	Plum, scrub (Prunus geniculata) -	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	562	Rhododendron, Chapman (Rhododendron chapmanii) -	E	\$45,000	\$0	\$45,000	\$0	\$45,000
	760	Gooseberry, Miccosukee (Ribes echinellum) -	T	\$24,500	\$0	\$24,500	\$0	\$24,500
	777	Bulrush, Northeastern (Scirpus ancistrochaetus) -	E	\$23,000	\$0	\$23,000	\$100	\$23,100
	1623	Pawpaw, four-petal (Asimina tetramera) -	E	\$1,100	\$495	\$1,595	\$0	\$1,595
	1647	Prickly-apple, aboriginal (Harrisia (=Cereus) aboriginum (=gracilis)) -	E	\$900	\$0	\$900	\$0	\$900
	587	Pea, Big Pine partridge (Chamaecrista lineata keyensis) -	E	\$41,000	\$0	\$41,000	\$0	\$41,000
	840	Spurge, wedge (Chamaesyce deltoidea serpyllum) -	E	\$19,500	\$0	\$19,500	\$0	\$19,500
	1041	Ziziphus, Florida (Ziziphus celata) -	E	\$12,000	\$0	\$12,000	\$0	\$12,000
Flowering Plants subtotal				\$17,574,319	\$4,720,755	\$22,295,074	\$265,215	\$22,560,289
Insects	197	Beetle, American burying (Nicrophorus americanus) -Wherever found, except where listed as an experimental population	T	\$192,500	\$330	\$192,830	\$191,500	\$384,330
	363	Beetle, Casey's June (Dinacoma caseyi) - Wherever found	E	\$110,300	\$0	\$110,300	\$0	\$110,300
	1541	Beetle, Comal Springs dryopid (Stygoparnus comalensis) -Wherever found	E	\$0	\$0	\$0	\$3,000	\$3,000
	1040	Beetle, delta green ground (Elaphrus viridis) -Wherever found	T	\$6,400	\$5,602	\$12,002	\$0	\$12,002
	582	Beetle, Hungerford's crawling water (Brychius hungerfordi) -Wherever found	E	\$41,500	\$0	\$41,500	\$0	\$41,500
	777	Beetle, Mount Hermon June (Polyphylla barbata) -Wherever found	E	\$23,100	\$0	\$23,100	\$0	\$23,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	143	Beetle, valley elderberry longhorn (Desmocerus californicus dimorphus) - Wherever found	T	\$608,800	\$36,591	\$645,391	\$0	\$645,391
	248	bumble bee, Franklin's (Bombus franklini) - Wherever found	E	\$238,174	\$1,388	\$239,562	\$0	\$239,562
	121	Bumble bee, Rusty patched (Bombus affinis) -Wherever found	E	\$691,400	\$8,176	\$699,576	\$91,519	\$791,095
	727	Butterfly, Bartram's hairstreak (Strymon acis bartrami) -Wherever found	E	\$21,000	\$5,500	\$26,500	\$0	\$26,500
	648	Butterfly, bay checkerspot (Euphydryas editha bayensis) -Wherever found	T	\$34,000	\$0	\$34,000	\$0	\$34,000
	422	Butterfly, Behren's silverspot (Speyeria zerene behrensi) -Wherever found	E	\$62,500	\$19,200	\$81,700	\$0	\$81,700
	411	Butterfly, callippe silverspot (Speyeria callippe callippe) -Wherever found	E	\$85,800	\$0	\$85,800	\$0	\$85,800
	1527	Butterfly, El Segundo blue (Euphilotes battoides allyni) -Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	166	Butterfly, Fender's blue (Icaricia icarioides fenderi) -Wherever found	E	\$426,461	\$66,175	\$492,636	\$0	\$492,636
	676	Butterfly, Florida leafwing (Anaea troglodyta floridalis) -Wherever found	E	\$23,500	\$7,000	\$30,500	\$0	\$30,500
	277	Butterfly, Hermes copper (Lycaena hermes) - Wherever found	T	\$185,300	\$0	\$185,300	\$0	\$185,300
	206	Butterfly, Island marble (Euchloe ausonides insulanus) -Wherever found	E	\$300,804	\$73,000	\$373,804	\$0	\$373,804
	217	Butterfly, Karner blue (Lycaeides melissa samuelis) -Wherever found	E	\$198,100	\$876	\$198,976	\$143,650	\$342,626
	334	Butterfly, Lange's metalmark (Apodemia mormo langei) -Wherever found	E	\$122,800	\$0	\$122,800	\$0	\$122,800
	388	Butterfly, Mariana eight-spot (Hypolimnas octocula marianensis) -Wherever found	E	\$96,551	\$0	\$96,551	\$0	\$96,551
	689	Butterfly, Mariana wandering (Vagrans egistina) -Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	749	Butterfly, Miami Blue (Cyclargus (=Hemiargus) thomasi bethunebakeri) - Wherever found	E	\$25,000	\$0	\$25,000	\$0	\$25,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	263	Butterfly, mission blue (<i>Icaricia icarioides missionensis</i>) -Wherever found	E	\$31,100	\$175,000	\$206,100	\$0	\$206,100
	252	Butterfly, Mitchell's satyr (<i>Neonympha mitchellii mitchellii</i>) -Wherever found	E	\$230,800	\$76	\$230,876	\$0	\$230,876
	510	Butterfly, Mount Charleston blue (<i>Icaricia (Plebejus) shasta charlestonensis</i>) -Wherever found	E	\$54,300	\$0	\$54,300	\$0	\$54,300
	1026	Butterfly, Myrtle's silverspot (<i>Speyeria zerene myrtleae</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	134	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) -Wherever found	T	\$682,537	\$12,455	\$694,992	\$0	\$694,992
	1591	Butterfly, Palos Verdes blue (<i>Glaucopsyche lygdamus palosverdesensis</i>) -Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	1292	butterfly, Puerto Rican harlequin (<i>Atlantea tulita</i>) -Wherever found	T	\$8,500	\$0	\$8,500	\$0	\$8,500
	714	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. wrighti)) -Wherever found	E	\$28,200	\$0	\$28,200	\$0	\$28,200
	1401	Butterfly, Saint Francis' satyr (<i>Neonympha mitchellii francisci</i>) -Wherever found	E	\$7,000	\$0	\$7,000	\$100	\$7,100
	730	Butterfly, San Bruno elfin (<i>Callophrys mossii bayensis</i>) -Wherever found	E	\$18,700	\$7,500	\$26,200	\$0	\$26,200
	421	Butterfly, Schaus swallowtail (<i>Heraclides aristodemus ponceanus</i>) -Wherever found	E	\$27,000	\$55,000	\$82,000	\$0	\$82,000
	333	Butterfly, Smith's blue (<i>Euphilotes enoptes smithi</i>) -Wherever found	E	\$123,000	\$0	\$123,000	\$0	\$123,000
	890	Butterfly, Uncompahgre fritillary (<i>Boloria acrocema</i>) -Wherever found	E	\$5,000	\$12,100	\$17,100	\$0	\$17,100
	136	Checkerspot, Taylor's (=whulge) (<i>Euphydryas editha taylori</i>) -Wherever found	E	\$688,050	\$2,999	\$691,049	\$0	\$691,049
	464	Damselfly, blackline Hawaiian (<i>Megalagrion nigrohamatum nigrolineatum</i>) -Wherever found	E	\$63,704	\$0	\$63,704	\$0	\$63,704
	919	Damselfly, crimson Hawaiian (<i>Megalagrion leptodemas</i>) -Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	967	Damselfly, flying earwig Hawaiian (Megalagrion nesiotes) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	907	Damselfly, oceanic Hawaiian (Megalagrion oceanicum) -Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	378	Damselfly, orangeblack Hawaiian (Megalagrion xanthomelas) -Wherever found	E	\$97,893	\$6,609	\$104,502	\$0	\$104,502
	496	Damselfly, Pacific Hawaiian (Megalagrion pacificum) -Wherever found	E	\$16,000	\$40,825	\$56,825	\$0	\$56,825
	676	damselfly, Rota blue (Ischnura luta) -Wherever found	E	\$30,500	\$0	\$30,500	\$0	\$30,500
	273	Dragonfly, Hine's emerald (Somatochlora hineana) -Wherever found	E	\$191,300	\$647	\$191,947	\$600	\$192,547
	1668	Fly, Delhi Sands flower-loving (Rhaphiomidas terminatus abdominalis) -Wherever found	E	\$500	\$0	\$500	\$0	\$500
	1463	Grasshopper, Zayante band-winged (Trimerotropis infantilis) -Wherever found	E	\$5,600	\$0	\$5,600	\$0	\$5,600
	503	Moth, Blackburn's sphinx (Manduca blackburni) -Wherever found	E	\$55,500	\$0	\$55,500	\$0	\$55,500
	1472	Moth, Kern primrose sphinx (Euproserpinus euterpe) -Wherever found	T	\$5,000	\$100	\$5,100	\$0	\$5,100
	723	Naucorid, Ash Meadows (Ambrysus amargosus) -Wherever found	T	\$27,100	\$0	\$27,100	\$0	\$27,100
	775	picture-wing fly, Hawaiian (Drosophila aglaia) -Wherever found	E	\$23,500	\$0	\$23,500	\$0	\$23,500
	748	picture-wing fly, Hawaiian (Drosophila differens) -Wherever found	E	\$25,038	\$0	\$25,038	\$0	\$25,038
	577	picture-wing fly, Hawaiian (Drosophila digressa) -Wherever found	E	\$42,000	\$0	\$42,000	\$0	\$42,000
	409	picture-wing fly, Hawaiian (Drosophila hemipeza) -Wherever found	E	\$86,140	\$0	\$86,140	\$0	\$86,140
	767	picture-wing fly, Hawaiian (Drosophila heteroneura) -Wherever found	E	\$24,430	\$0	\$24,430	\$0	\$24,430
	440	picture-wing fly, Hawaiian (Drosophila montgomeryi) -Wherever found	E	\$75,653	\$0	\$75,653	\$0	\$75,653

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	722	picture-wing fly, Hawaiian (<i>Drosophila mulli</i>) -Wherever found	T	\$27,430	\$0	\$27,430	\$0	\$27,430
	919	picture-wing fly, Hawaiian (<i>Drosophila musaphilia</i>) -Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	919	picture-wing fly, Hawaiian (<i>Drosophila neoclavisetae</i>) -Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	1292	picture-wing fly, Hawaiian (<i>Drosophila obatai</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	902	picture-wing fly, Hawaiian (<i>Drosophila ochrobasis</i>) -Wherever found	E	\$16,430	\$0	\$16,430	\$0	\$16,430
	919	picture-wing fly, Hawaiian (<i>Drosophila sharpi</i>) -Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	1292	picture-wing fly, Hawaiian (<i>Drosophila substenoptera</i>) -Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	768	picture-wing fly, Hawaiian (<i>Drosophila tarphytrichia</i>) -Wherever found	E	\$24,000	\$0	\$24,000	\$0	\$24,000
	632	Skipper, Carson wandering (<i>Pseudocopaeodes eunus obscurus</i>) -Wherever found	E	\$13,400	\$22,507	\$35,907	\$0	\$35,907
	255	Skipper, Dakota (<i>Hesperia dacotae</i>) -Wherever found	T	\$174,300	\$42,338	\$216,638	\$11,005	\$227,643
	292	Skipper, Laguna Mountains (<i>Pyrgus ruralis lagunae</i>) -Wherever found	E	\$160,200	\$0	\$160,200	\$0	\$160,200
	1566	Skipper, Pawnee montane (<i>Hesperia leonardus montana</i>) -Wherever found	T	\$2,600	\$100	\$2,700	\$0	\$2,700
	287	skipperling, Poweshiek (<i>Oarisma poweshiek</i>) -Wherever found	E	\$161,200	\$76	\$161,276	\$4,500	\$165,776
	1569	Stonefly, meltwater lednian (<i>Lednia tumana</i>) -Wherever found	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	1001	Stonefly, western glacier (<i>Zapada glacier</i>) -Wherever found	T	\$3,300	\$9,750	\$13,050	\$0	\$13,050
	387	Tiger beetle, Miami (<i>Cicindelidia floridana</i>) -Wherever found	E	\$97,500	\$0	\$97,500	\$0	\$97,500
	329	Tiger beetle, Northeastern beach (<i>Habroscelimorpha dorsalis dorsalis</i>) -Wherever found	T	\$122,500	\$1,528	\$124,028	\$300	\$124,328

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1061	Tiger beetle, Ohlone (Cicindela ohlone) - Wherever found	E	\$11,800	\$0	\$11,800	\$0	\$11,800
	441	Tiger beetle, Puritan (Ellipsoptera puritana) - Wherever found	T	\$75,000	\$0	\$75,000	\$0	\$75,000
	429	Tiger beetle, Salt Creek (Cicindela nevadica lincolniana) -Wherever found	E	\$63,000	\$0	\$63,000	\$16,500	\$79,500
	349	Yellow-faced bee, anthracin (Hylaeus anthracinus) -Wherever found	E	\$116,634	\$0	\$116,634	\$0	\$116,634
	785	Yellow-faced bee, assimulans (Hylaeus assimulans) -Wherever found	E	\$22,606	\$0	\$22,606	\$0	\$22,606
	919	Yellow-faced bee, easy (Hylaeus facilis) - Wherever found	E	\$15,500	\$0	\$15,500	\$0	\$15,500
	741	yellow-faced bee, Hawaiian (Hylaeus kuakea) -Wherever found	E	\$25,500	\$0	\$25,500	\$0	\$25,500
	768	yellow-faced bee, Hawaiian (Hylaeus longiceps) -Wherever found	E	\$24,000	\$0	\$24,000	\$0	\$24,000
	749	yellow-faced bee, Hawaiian (Hylaeus mana) -Wherever found	E	\$25,000	\$0	\$25,000	\$0	\$25,000
	1064	Yellow-faced bee, hiliaris (Hylaeus hiliaris) - Wherever found	E	\$11,500	\$0	\$11,500	\$0	\$11,500
	386	Butterfly, Oregon silverspot (Speyeria zerene hippolyta) -	EXPN	\$98,900	\$0	\$98,900	\$0	\$98,900
Insects subtotal				\$7,619,835	\$613,448	\$8,233,283	\$462,674	\$8,695,957
Lichens	584	Lichen, rock gnome (Gymnoderma lineare) - Wherever found	E	\$41,000	\$0	\$41,000	\$400	\$41,400
	1533	Cladonia, Florida perforate (Cladonia perforata) -	E	\$3,000	\$495	\$3,495	\$0	\$3,495
Lichens subtotal				\$44,000	\$495	\$44,495	\$400	\$44,895
Mammals	1492	Addax (Addax nasomaculatus) -Wherever found	E	\$0	\$4,900	\$4,900	\$0	\$4,900
	863	Ass, Asian wild (Equus hemionus) - Wherever found	E	\$0	\$18,400	\$18,400	\$0	\$18,400
	167	bat, Florida bonneted (Eumops floridanus) - Wherever found	E	\$308,000	\$46,495	\$354,495	\$136,956	\$491,451
	92	Bat, gray (Myotis grisescens) -Wherever found	E	\$605,200	\$382,357	\$987,557	\$149,956	\$1,137,513

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	157	Bat, Hawaiian hoary (<i>Lasiurus cinereus semotus</i>) -Wherever found	E	\$513,487	\$40,825	\$554,312	\$0	\$554,312
	59	Bat, Indiana (<i>Myotis sodalis</i>) -Wherever found	E	\$1,634,800	\$322,356	\$1,957,156	\$526,700	\$2,483,856
	760	Bat, little Mariana fruit (<i>Pteropus tokudae</i>) -Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	331	Bat, Mariana fruit (=Mariana flying fox) (<i>Pteropus mariannus mariannus</i>) -Wherever found	T	\$123,600	\$0	\$123,600	\$0	\$123,600
	624	Bat, Mexican long-nosed (<i>Leptonycteris nivalis</i>) -Wherever found	E	\$0	\$36,541	\$36,541	\$0	\$36,541
	49	Bat, Northern long-eared (<i>Myotis septentrionalis</i>) -Wherever found	E	\$1,693,300	\$1,292,968	\$2,986,268	\$139,202	\$3,125,470
	439	Bat, Ozark big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii ingens</i>) -Wherever found	E	\$47,500	\$5,000	\$52,500	\$23,350	\$75,850
	369	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata rotensis</i>) -Wherever found	E	\$109,000	\$0	\$109,000	\$0	\$109,000
	642	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata semicaudata</i>) -American Samoa	E	\$34,930	\$0	\$34,930	\$0	\$34,930
	289	Bat, Virginia big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii virginianus</i>) -Wherever found	E	\$150,500	\$6,520	\$157,020	\$7,961	\$164,981
	24	Bear, grizzly (<i>Ursus arctos horribilis</i>) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$1,254,154	\$2,528,731	\$3,782,885	\$3,040,549	\$6,823,434
	75	Bear, polar (<i>Ursus maritimus</i>) -Wherever found	T	\$1,289,000	\$330,000	\$1,619,000	\$4,402	\$1,623,402
	128	Bison, wood (<i>Bison bison athabasca</i>) - U.S.A. (Alaska)	EXPN	\$27,000	\$0	\$27,000	\$711,632	\$738,632
	828	Bison, wood (<i>Bison bison athabasca</i>) -Wherever found	T	\$0	\$20,000	\$20,000	\$0	\$20,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	312	Caribou DPS, Southern Mountain (Rangifer tarandus ssp. caribou) -U.S.A. (wherever found), Canada (southeastern British Columbia)	E	\$140,339	\$0	\$140,339	\$1,000	\$141,339
	1639	Cat, black-footed (Felis nigripes) -Wherever found	E	\$0	\$1,000	\$1,000	\$0	\$1,000
	428	Cheetah (Acinonyx jubatus) -Wherever found	E	\$62,190	\$17,600	\$79,790	\$0	\$79,790
	179	Chimpanzee (Pan troglodytes) -Wherever found	E	\$445,992	\$1,800	\$447,792	\$0	\$447,792
	1541	Chimpanzee, pygmy (Pan paniscus) - Wherever found	E	\$0	\$3,000	\$3,000	\$0	\$3,000
	1668	Deer, Barbary (Cervus elaphus barbarus) - Wherever found	E	\$0	\$500	\$500	\$0	\$500
	70	Deer, Columbian white-tailed (Odocoileus virginianus leucurus) -Columbia River (Clark, Cowliz, Pacific, Skamania, and Wahkiakum Counties, WA., and Clatsop, Columbia, and Multnomah Counties, OR.)	T	\$1,761,500	\$119,384	\$1,880,884	\$0	\$1,880,884
	1507	Deer, Eld's brow-antlered (Cervus eldi) - Wherever found	E	\$0	\$4,000	\$4,000	\$0	\$4,000
	347	Deer, key (Odocoileus virginianus clavium) - Wherever found	E	\$118,000	\$0	\$118,000	\$0	\$118,000
	1109	Dog, African wild (Lycaon pictus) - Wherever found	E	\$0	\$11,300	\$11,300	\$0	\$11,300
	54	Elephant, African (Loxodonta africana) - Wherever found	T	\$2,721,046	\$2,400	\$2,723,446	\$0	\$2,723,446
	71	Elephant, Asian (Elephas maximus) - Wherever found	E	\$1,644,960	\$68,100	\$1,713,060	\$0	\$1,713,060
	61	Ferret, black-footed (Mustela nigripes) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$1,784,500	\$264,626	\$2,049,126	\$338,583	\$2,387,709
	99	Ferret, black-footed (Mustela nigripes) - Wherever found, except where listed as an experimental population	E	\$380,500	\$436,600	\$817,100	\$217,754	\$1,034,854
	210	Fisher (Pekania pennanti) -Southern Sierra Nevada Distinct Population Segment	E	\$57,800	\$309,740	\$367,540	\$0	\$367,540

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	105	Fisher (<i>Pekania pennanti</i>) -West Coast Distinct Population Segment	E	\$909,100	\$5,240	\$914,340	\$0	\$914,340
	126	Fox, San Joaquin kit (<i>Vulpes macrotis mutica</i>) -wherever found	E	\$698,300	\$46,359	\$744,659	\$0	\$744,659
	242	Fox, Santa Catalina Island (<i>Urocyon littoralis catalinae</i>) -Wherever found	T	\$241,700	\$13,500	\$255,200	\$0	\$255,200
	152	Fox, Sierra Nevada red (<i>Vulpes vulpes necator</i>) -Sierra Nevada DPS	E	\$467,000	\$98,664	\$565,664	\$0	\$565,664
	1668	Gazelle, Mhorr (<i>Gazella dama mhorr</i>) - Wherever found	E	\$0	\$500	\$500	\$0	\$500
	213	Gibbons (<i>Hylobates</i> spp. (including <i>Nomascus</i>)) -Wherever found	E	\$358,367	\$2,000	\$360,367	\$0	\$360,367
	145	Gorilla (<i>Gorilla gorilla</i>) -Wherever found	E	\$616,342	\$500	\$616,842	\$0	\$616,842
	645	Horse, Przewalski's (<i>Equus przewalskii</i>) - Wherever found	E	\$0	\$34,500	\$34,500	\$0	\$34,500
	135	Jaguar (<i>Panthera onca</i>) -Wherever found	E	\$678,344	\$13,510	\$691,854	\$0	\$691,854
	344	Jaguarundi, Gulf Coast (<i>Puma yagouaroundi cacomitli</i>) -Wherever found	E	\$30,000	\$88,600	\$118,600	\$0	\$118,600
	541	Kangaroo rat, Fresno (<i>Dipodomys nitratoide exilis</i>) -Wherever found	E	\$18,900	\$29,231	\$48,131	\$0	\$48,131
	364	Kangaroo rat, giant (<i>Dipodomys ingens</i>) - Wherever found	E	\$77,100	\$33,035	\$110,135	\$0	\$110,135
	1358	Kangaroo rat, Morro Bay (<i>Dipodomys heermanni morroensis</i>) -Wherever found	E	\$7,900	\$0	\$7,900	\$0	\$7,900
	195	Kangaroo rat, San Bernardino Merriam's (<i>Dipodomys merriami parvus</i>) -Wherever found	E	\$392,100	\$0	\$392,100	\$0	\$392,100
	257	Kangaroo rat, Stephens' (<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)) -Wherever found	T	\$220,500	\$0	\$220,500	\$0	\$220,500
	237	Kangaroo rat, Tipton (<i>Dipodomys nitratoide nitratoide</i>) -Wherever found	E	\$261,900	\$2,560	\$264,460	\$0	\$264,460
	1507	Leopard (<i>Panthera pardus</i>) -In Africa, in the wild, south of, and including, the following countries: Gabon, Congo, Zaire, Uganda, Kenya	T	\$0	\$4,000	\$4,000	\$0	\$4,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	715	Leopard, clouded (<i>Neofelis nebulosa</i>) - Wherever found	E	\$0	\$28,000	\$28,000	\$0	\$28,000
	461	Leopard, snow (<i>Uncia</i> (=Panthera) <i>uncia</i>) - Wherever found	E	\$65,180	\$0	\$65,180	\$0	\$65,180
	286	Lion (<i>Panthera leo</i> ssp. <i>leo</i>) -Wherever found	E	\$168,533	\$0	\$168,533	\$0	\$168,533
	57	Lynx, Canada (<i>Lynx canadensis</i>) -Wherever Found in Contiguous U.S.	T	\$1,670,130	\$525,419	\$2,195,549	\$321,913	\$2,517,462
	1675	Macaque, Japanese (<i>Macaca fuscata</i>) - Wherever found	T	\$0	\$200	\$200	\$0	\$200
	1668	Macaque, lion-tailed (<i>Macaca silenus</i>) - Wherever found	E	\$0	\$500	\$500	\$0	\$500
	22	Manatee, West Indian (<i>Trichechus manatus</i>) -Wherever found	T	\$1,943,500	\$26,320	\$1,969,820	\$6,178,427	\$8,148,247
	139	Marten, Pacific (<i>Martes caurina</i>) -Wherever found	T	\$456,678	\$202,805	\$659,483	\$0	\$659,483
	422	Mountain beaver, Point Arena (<i>Aplodontia</i> <i>rufa nigra</i>) -Wherever found	E	\$80,100	\$1,600	\$81,700	\$0	\$81,700
	291	Mouse, Alabama beach (<i>Peromyscus</i> <i>polionotus ammobates</i>) -Wherever found	E	\$160,500	\$990	\$161,490	\$0	\$161,490
	413	Mouse, Anastasia Island beach (<i>Peromyscus</i> <i>polionotus phasma</i>) -Wherever found	E	\$60,000	\$1,000	\$61,000	\$24,397	\$85,397
	408	Mouse, Choctawhatchee beach (<i>Peromyscus</i> <i>polionotus allopheys</i>) -Wherever found	E	\$70,000	\$0	\$70,000	\$16,265	\$86,265
	564	Mouse, Key Largo cotton (<i>Peromyscus</i> <i>gossypinus allapaticola</i>) -Wherever found	E	\$44,500	\$0	\$44,500	\$0	\$44,500
	400	Mouse, New Mexico meadow jumping (<i>Zapus hudsonius luteus</i>) -Wherever found	E	\$34,600	\$57,300	\$91,900	\$0	\$91,900
	385	Mouse, Pacific pocket (<i>Perognathus</i> <i>longimembris pacificus</i>) -Wherever found	E	\$100,200	\$0	\$100,200	\$0	\$100,200
	450	Mouse, Perdido Key beach (<i>Peromyscus</i> <i>polionotus trissyllepsis</i>) -Wherever found	E	\$46,500	\$5,625	\$52,125	\$16,265	\$68,390
	244	Mouse, Preble's meadow jumping (<i>Zapus</i> <i>hudsonius preblei</i>) -wherever found	T	\$184,000	\$14,447	\$198,447	\$50,107	\$248,554

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	150	Mouse, salt marsh harvest (Reithrodontomys raviventris) -wherever found	E	\$571,300	\$0	\$571,300	\$0	\$571,300
	368	Mouse, southeastern beach (Peromyscus polionotus niveiventris) -wherever found	T	\$79,500	\$5,330	\$84,830	\$24,397	\$109,227
	317	Mouse, St. Andrew beach (Peromyscus polionotus peninsularis) -Wherever found	E	\$118,000	\$0	\$118,000	\$16,265	\$134,265
	228	Ocelot (Leopardus (=Felis) pardalis) - wherever found	E	\$212,000	\$88,910	\$300,910	\$0	\$300,910
	95	Orangutan (Pongo pygmaeus) -Wherever found	E	\$1,079,000	\$0	\$1,079,000	\$0	\$1,079,000
	639	Oryx, scimitar-horned (Oryx dammah) - Wherever found	E	\$0	\$35,100	\$35,100	\$0	\$35,100
	177	Otter, Northern Sea (Enhydra lutris kenyoni) -Southwest Alaska, from Attu Island to Western Cook Inlet, including Bristol Bay, the Kodiak Archipelago, and the Barren Islands	T	\$431,500	\$0	\$431,500	\$27,408	\$458,908
	148	Otter, southern sea (Enhydra lutris nereis) - Wherever found	T	\$580,060	\$100	\$580,160	\$0	\$580,160
	706	Panda, giant (Ailuropoda melanoleuca) - Wherever found	E	\$0	\$28,900	\$28,900	\$0	\$28,900
	81	Panther, Florida (Puma (=Felis) concolor coryi) -Wherever found	E	\$760,000	\$37,500	\$797,500	\$583,861	\$1,381,361
	239	Pocket gopher, Olympia (Thomomys mazama pugetensis) -Wherever found	T	\$261,733	\$0	\$261,733	\$0	\$261,733
	356	Pocket gopher, Roy Prairie (Thomomys mazama glacialis) -Wherever found	T	\$113,008	\$0	\$113,008	\$0	\$113,008
	348	Pocket gopher, Tenino (Thomomys mazama tumuli) -Wherever found	T	\$117,868	\$0	\$117,868	\$0	\$117,868
	332	Pocket gopher, Yelm (Thomomys mazama yelmensis) -Wherever found	T	\$123,538	\$0	\$123,538	\$0	\$123,538
	138	Prairie dog, Utah (Cynomys parvidens) - Wherever found	T	\$120,000	\$551,181	\$671,181	\$0	\$671,181

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	949	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) -In Arizona, an area north of Interstate 8 and south of Interstate 10, bounded by the Colorado River on the west and Interstate 10 on the east; and an area south of Interstate 8, bounded by Highway 85 on the west, Interstates 10 and 19 on the east, and the U.S.-Mexico border on the south	EXPN	\$0	\$14,850	\$14,850	\$0	\$14,850
	280	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) -Wherever found, except where listed as an experimental population	E	\$35,500	\$148,453	\$183,953	\$0	\$183,953
	614	Rabbit, Lower Keys marsh (<i>Sylvilagus palustris hefneri</i>) -Wherever found	E	\$37,500	\$0	\$37,500	\$0	\$37,500
	108	Rabbit, Pygmy (<i>Brachylagus idahoensis</i>) - U.S.A. (WA-Douglas, Grant, Lincoln, Adams, Benton Counties)	E	\$872,269	\$11,000	\$883,269	\$0	\$883,269
	232	Rabbit, riparian brush (<i>Sylvilagus bachmani riparius</i>) -Wherever found	E	\$287,200	\$500	\$287,700	\$0	\$287,700
	62	Rhinoceros, black (<i>Diceros bicornis</i>) -Sub-Saharan Africa	E	\$2,342,863	\$38,500	\$2,381,363	\$0	\$2,381,363
	171	Rhinoceros, great Indian (<i>Rhinoceros unicornis</i>) -Nepal, India	E	\$487,463	\$0	\$487,463	\$0	\$487,463
	276	Rhinoceros, Javan (<i>Rhinoceros sondaicus</i>) - Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$186,000	\$0	\$186,000	\$0	\$186,000
	589	Rhinoceros, northern white (<i>Ceratotherium simum cottoni</i>) -Zaire, Uganda, Sudan, Central African Republic	E	\$40,295	\$0	\$40,295	\$0	\$40,295
	185	Rhinoceros, Sumatran (<i>Dicerorhinus sumatrensis</i>) -Wherever found	E	\$427,709	\$0	\$427,709	\$0	\$427,709
	619	Rice rat, silver (<i>Oryzomys palustris natator</i>) - Lower FL Keys (west of Seven Mile Bridge)	E	\$37,000	\$0	\$37,000	\$0	\$37,000
	32	Sea lion, Steller (<i>Eumetopias jubatus</i>) - Western DPS - See 50 CFR 224.101	E	\$0	\$4,685,000	\$4,685,000	\$335,944	\$5,020,944

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	67	Seal, bearded (<i>Erignathus barbatus nauticus</i>) -Beringia DPS - See 50 CFR 223.102	T	\$0	\$1,891,000	\$1,891,000	\$0	\$1,891,000
	254	Seal, Guadalupe fur (<i>Arctocephalus townsendi</i>) -Wherever found	T	\$0	\$227,800	\$227,800	\$0	\$227,800
	40	Seal, Hawaiian monk (<i>Monachus schauinslandi</i>) -Wherever found	E	\$88,000	\$4,274,000	\$4,362,000	\$0	\$4,362,000
	67	Seal, Ringed (<i>Phoca</i> (=Pusa) <i>hispida botnica</i>) -Wherever found	T	\$0	\$1,891,000	\$1,891,000	\$0	\$1,891,000
	338	Seal, Ringed (<i>Phoca</i> (=Pusa) <i>hispida hispida</i>) -Wherever found	T	\$0	\$120,000	\$120,000	\$0	\$120,000
	358	Sheep, Sierra Nevada bighorn (<i>Ovis canadensis sierrae</i>) -U.S.A. (CA) Sierra Nevada	E	\$19,500	\$93,080	\$112,580	\$0	\$112,580
	412	Shrew, Buena Vista Lake ornate (<i>Sorex ornatus relictus</i>) -Wherever found	E	\$71,600	\$14,000	\$85,600	\$0	\$85,600
	1568	Sloth, Brazilian three-toed (<i>Bradypus torquatus</i>) -Wherever found	E	\$2,666	\$0	\$2,666	\$0	\$2,666
	529	Squirrel, Carolina northern flying (<i>Glaucomys sabrinus coloratus</i>) -Wherever found	E	\$42,500	\$5,055	\$47,555	\$1,700	\$49,255
	208	Squirrel, Northern Idaho Ground (<i>Urocitellus brunneus</i>) -Wherever found	T	\$340,957	\$0	\$340,957	\$30,000	\$370,957
	741	Tapir, mountain (<i>Tapirus pinchaque</i>) -Wherever found	E	\$0	\$25,500	\$25,500	\$0	\$25,500
	87	Tiger (<i>Panthera tigris</i>) -Wherever found	E	\$1,195,678	\$25,500	\$1,221,178	\$0	\$1,221,178
	129	Vole, Amargosa (<i>Microtus californicus scirpensis</i>) -Wherever found	E	\$356,500	\$382,000	\$738,500	\$0	\$738,500
	1416	Vole, Florida salt marsh (<i>Microtus pennsylvanicus dukecampbelli</i>) -Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	46	whale, beluga (<i>Delphinapterus leucas</i>) -Cook Inlet DPS See 50 CFR 224.101	E	\$0	\$3,037,000	\$3,037,000	\$396,188	\$3,433,188
	85	Whale, blue (<i>Balaenoptera musculus</i>) -Wherever found	E	\$0	\$1,255,800	\$1,255,800	\$0	\$1,255,800
	188	Whale, bowhead (<i>Balaena mysticetus</i>) -Wherever found	E	\$0	\$407,000	\$407,000	\$0	\$407,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	240	whale, false killer (<i>Pseudorca crassidens</i>) - Main Hawaiian Islands Insular DPS See 50 CFR 224.101	E	\$0	\$261,000	\$261,000	\$0	\$261,000
	84	Whale, finback (<i>Balaenoptera physalus</i>) - Wherever found	E	\$0	\$1,260,000	\$1,260,000	\$0	\$1,260,000
	1608	Whale, gray (<i>Eschrichtius robustus</i>) - Western North Pacific DPS See 50 CFR 224.101	E	\$0	\$1,700	\$1,700	\$0	\$1,700
	1675	Whale, humpback (<i>Megaptera novaeangliae</i>) -Cape Verde Island/Northwest Africa	E	\$0	\$200	\$200	\$0	\$200
	215	Whale, humpback (<i>Megaptera novaeangliae</i>) -Central America DPS	E	\$0	\$357,800	\$357,800	\$0	\$357,800
	212	Whale, humpback (<i>Megaptera novaeangliae</i>) -Mexico DPS	T	\$0	\$361,000	\$361,000	\$0	\$361,000
	199	Whale, humpback (<i>Megaptera novaeangliae</i>) -Western North Pacific	E	\$0	\$381,000	\$381,000	\$0	\$381,000
	60	Whale, killer (<i>Orcinus orca</i>) -Southern Resident DPS See 50 CFR 224.101	E	\$0	\$2,402,100	\$2,402,100	\$0	\$2,402,100
	4	Whale, North Atlantic Right (<i>Eubalaena glacialis</i>) -Wherever found	E	\$0	\$28,572,500	\$28,572,500	\$867,695	\$29,440,195
	198	Whale, North Pacific Right (<i>Eubalaena japonica</i>) -Wherever found	E	\$0	\$381,100	\$381,100	\$0	\$381,100
	96	Whale, Sei (<i>Balaenoptera borealis</i>) - Wherever found	E	\$0	\$1,071,300	\$1,071,300	\$0	\$1,071,300
	97	Whale, sperm (<i>Physeter catodon</i> (=macrocephalus)) -Wherever found	E	\$0	\$1,071,100	\$1,071,100	\$0	\$1,071,100
	191	Wolf, gray (<i>Canis lupus</i>) -U.S.A. (MN)	T	\$399,500	\$0	\$399,500	\$552	\$400,052

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	64	Wolf, gray (<i>Canis lupus</i>) -U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of Highway 97 and Highway 17 north of Mesa and that portion of WA west of the centerline of Highway 395 south of Mesa). Mexico.	E	\$623,572	\$388,127	\$1,011,699	\$1,153,868	\$2,165,567
	531	Wolf, maned (<i>Chrysocyon brachyurus</i>) - Wherever found	E	\$0	\$49,100	\$49,100	\$0	\$49,100
	305	Wolf, Mexican (<i>Canis lupus baileyi</i>) -U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$0	\$150,910	\$150,910	\$0	\$150,910
	44	Wolf, Mexican (<i>Canis lupus baileyi</i>) - Wherever found, except where included in an experimental population as set forth in 17.84(k)	E	\$3,649,996	\$0	\$3,649,996	\$0	\$3,649,996
	373	Wolf, red (<i>Canis rufus</i>) -U.S.A. (portions of NC and TN see 17.84(c)(9))	EXPN	\$107,000	\$0	\$107,000	\$0	\$107,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	82	Wolf, red (Canis rufus) -Wherever found, except where listed as an experimental population	E	\$1,242,000	\$600	\$1,242,600	\$85,500	\$1,328,100
	444	Woodrat, Key Largo (Neotoma floridana smalli) -Wherever found	E	\$71,500	\$0	\$71,500	\$0	\$71,500
	1591	Woodrat, riparian (=San Joaquin Valley) (Neotoma fuscipes riparia) -Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
Mammals subtotal				\$44,534,547	\$63,515,144	\$108,049,691	\$15,428,797	\$123,478,488
Reptiles	65	Alligator, American (Alligator mississippiensis) -Wherever found	SAT	\$0	\$106,990	\$106,990	\$1,949,558	\$2,056,548
	1541	Anole, Culebra Island giant (Anolis roosevelti) -Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1338	Boa, Mona (Epicrates monensis monensis) -Wherever found	T	\$8,000	\$0	\$8,000	\$0	\$8,000
	353	Boa, Puerto Rican (Chilabothrus inornatus) -Wherever found	E	\$114,000	\$0	\$114,000	\$0	\$114,000
	760	Boa, Virgin Islands tree (Chilabothrus granti) -Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	246	Crocodile, American (Crocodylus acutus) - U.S.A. (FL)	T	\$83,000	\$160,500	\$243,500	\$0	\$243,500
	416	Crocodile, American (Crocodylus acutus) -Wherever found, except in U.S.A. (FL)	E	\$0	\$0	\$0	\$84,464	\$84,464
	1636	gartersnake, narrow-headed (Thamnophis rufipunctatus) -Wherever found	T	\$0	\$1,184	\$1,184	\$0	\$1,184
	393	gartersnake, northern Mexican (Thamnophis eques megalops) -Wherever found	T	\$0	\$95,582	\$95,582	\$0	\$95,582
	987	Iguana, Mona ground (Cyclura stejnegeri) -Wherever found	T	\$13,500	\$0	\$13,500	\$0	\$13,500
	236	Lizard, blunt-nosed leopard (Gambelia silus) -Wherever found	E	\$230,600	\$37,412	\$268,012	\$0	\$268,012
	1680	Lizard, Coachella Valley fringe-toed (Uma inornata) -Wherever found	T	\$100	\$0	\$100	\$0	\$100
	634	Lizard, St. Croix ground (Ameiva polops) -Wherever found	E	\$10,500	\$25,000	\$35,500	\$0	\$35,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	190	Massasauga (=rattlesnake), eastern (Sistrurus catenatus) -Wherever found	T	\$369,900	\$1,014	\$370,914	\$32,300	\$403,214
	700	piresnake, black (Pituophis melanoleucus lodingi) -Wherever found	T	\$29,000	\$330	\$29,330	\$0	\$29,330
	253	Pinesnake, Louisiana (Pituophis ruthveni) -Wherever found	T	\$227,300	\$500	\$227,800	\$3,000	\$230,800
	1646	Rattlesnake, New Mexican ridge-nosed (Crotalus willardi obscurus) -Wherever found	T	\$0	\$903	\$903	\$0	\$903
	90	Sea turtle, green (Chelonia mydas) -Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	T	\$356,011	\$809,418	\$1,165,429	\$0	\$1,165,429
	141	Sea turtle, green (Chelonia mydas) -Green sea turtles originating from the Central South Pacific Ocean, bounded by the following coordinates: 9 degrees N., 175 degrees W. in the northwest; 9 degrees N., 125 degrees W. in the northeast; 40 degrees S., 96 degrees W. in the southeast; 40 degrees S., 176 degrees E. in the southwest; and 13 degrees S., 171 degrees E. in the west	E	\$123,500	\$527,750	\$651,250	\$0	\$651,250

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	91	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Central West Pacific Ocean, bounded by the following coordinates: 41 degrees N., 146 degrees E. in the northwest; 41 degrees N., 169 degrees E. in the northeast; 9 degrees N., 175 degrees W. in the east; 13 degrees S., 171 degrees E. in the southeast; along the northern coast of the island of New Guinea; and 4.5 degrees N., 129 degrees E. in the west	E	\$117,000	\$1,026,000	\$1,143,000	\$0	\$1,143,000
	102	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the East Pacific Ocean, bounded by the following lines and coordinates: 41 degrees N., 143 degrees W. in the northwest; 41 degrees N. Lat. in the north; along the western coasts of the Americas; 40 degrees S. Lat. in the south; and 40 degrees S., 96 degrees W. in the southwest	T	\$4,500	\$936,500	\$941,000	\$0	\$941,000
	499	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Eastern Indian and Western Pacific Oceans, bounded by the following lines and coordinates: 41 degrees N. Lat. in the north, 41 degrees N., 146 degrees E. in the northeast; 4.5 degrees N., 129 degrees E. in the southeast; along the southern coast of the island of New Guinea; along the western coast of Australia (west of 142 degrees E. Long.); 40 degrees S. Lat. in the south; and 84 degrees E. Long. in the east	T	\$56,287	\$0	\$56,287	\$0	\$56,287

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	74	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	T	\$166,500	\$1,475,045	\$1,641,545	\$0	\$1,641,545
	779	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the North Indian Ocean, bounded by: Africa and Asia in the west and north; 84 degrees E. Long. in the east; and the equator in the south	T	\$23,000	\$0	\$23,000	\$0	\$23,000
	172	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	T	\$164,000	\$15,000	\$179,000	\$308,443	\$487,443
	52	Sea turtle, hawksbill (<i>Eretmochelys imbricata</i>) -Wherever found	E	\$421,787	\$2,061,116	\$2,482,903	\$308,443	\$2,791,346
	51	Sea turtle, Kemp's ridley (<i>Lepidochelys kempii</i>) -Wherever found	E	\$162,516	\$2,329,281	\$2,491,797	\$308,443	\$2,800,240
	41	Sea turtle, leatherback (<i>Dermochelys coriacea</i>) -Wherever found	E	\$1,014,834	\$2,863,412	\$3,878,246	\$308,443	\$4,186,689

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	807	Sea turtle, loggerhead (Caretta caretta) - Loggerhead sea turtles originating from the South Atlantic Ocean south of the equator, north of 60 degrees S. Lat., west of 20 degrees E. Long., and east of 67 degrees W. Long	T	\$21,000	\$200	\$21,200	\$0	\$21,200
	475	Sea turtle, loggerhead (Caretta caretta) - Loggerhead sea turtles originating from the South Pacific south of the equator, north of 60 degrees S. Lat., west of 67 degrees W. Long., and east of 141 degrees E. Long.	E	\$0	\$60,480	\$60,480	\$0	\$60,480
	365	Sea turtle, loggerhead (Caretta caretta) - Mediterranean Sea DPS - Loggerhead sea turtles originating from the Mediterranean Sea east of 5 degrees 36 minues W. Long.	E	\$110,000	\$0	\$110,000	\$0	\$110,000
	527	Sea turtle, loggerhead (Caretta caretta) - North Indian Ocean DPS - Loggerhead sea turtles originating from the North Indian Ocean north of the equator and south of 30? N. Lat	E	\$49,500	\$0	\$49,500	\$0	\$49,500
	77	Sea turtle, loggerhead (Caretta caretta) - North Pacific Ocean DPS - Loggerhead sea turtles originating from the North Pacific north of the equator and south of 60 degrees N. Lat	E	\$100,734	\$1,429,800	\$1,530,534	\$0	\$1,530,534
	183	Sea turtle, loggerhead (Caretta caretta) - Northeast Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northeast Atlantic Ocean north of the equator, south of 60 degrees N. Lat., and east of 40 degrees W. Long., except in the vicinity of the Strait of Gibraltar where the eastern boundary is 5 degrees 36 minutes W. Long	E	\$436,500	\$200	\$436,700	\$0	\$436,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	47	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	T	\$622,500	\$2,249,833	\$2,872,333	\$551,843	\$3,424,176
	224	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) -Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$90,720	\$234,500	\$325,220	\$0	\$325,220
	406	skink, blue-tailed mole (<i>Eumeces egregius lividus</i>) -Wherever found	T	\$87,000	\$0	\$87,000	\$0	\$87,000
	327	Skink, sand (<i>Neoseps reynoldsi</i>) -Wherever found	T	\$125,000	\$0	\$125,000	\$0	\$125,000
	492	skink, Slevin's (<i>Emoia slevini</i>) -Wherever found	E	\$57,000	\$0	\$57,000	\$0	\$57,000
	967	Snake, Atlantic salt marsh (<i>Nerodia clarkii taeniata</i>) -Wherever found	T	\$14,000	\$0	\$14,000	\$0	\$14,000
	612	Snake, copperbelly water (<i>Nerodia erythrogaster neglecta</i>) -Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$37,700	\$76	\$37,776	\$0	\$37,776
	161	Snake, eastern indigo (<i>Drymarchon couperi</i>) -Wherever found	T	\$389,500	\$3,055	\$392,555	\$129,669	\$522,224
	211	Snake, giant garter (<i>Thamnophis gigas</i>) - Wherever found	T	\$329,000	\$34,508	\$363,508	\$0	\$363,508
	377	Snake, San Francisco garter (<i>Thamnophis sirtalis tetrataenia</i>) -Wherever found	E	\$94,600	\$10,000	\$104,600	\$0	\$104,600
	896	Tortoise, Bolson (<i>Gopherus flavomarginatus</i>) -Wherever found	E	\$16,500	\$0	\$16,500	\$0	\$16,500
	370	Tortoise, desert (<i>Gopherus agassizii</i>) -AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$59,300	\$49,033	\$108,333	\$0	\$108,333
	53	Tortoise, desert (<i>Gopherus agassizii</i>) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$1,358,400	\$1,390,767	\$2,749,167	\$0	\$2,749,167
	196	Tortoise, gopher (<i>Gopherus polyphemus</i>) - Western DPS	T	\$385,000	\$1,480	\$386,480	\$0	\$386,480

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1639	Tortoise, Madagascar radiated (<i>Geochelone radiata</i>) -Wherever found	E	\$0	\$1,000	\$1,000	\$0	\$1,000
	738	Turtle, Alabama red-bellied (<i>Pseudemys alabamensis</i>) -Wherever found	E	\$25,000	\$660	\$25,660	\$0	\$25,660
	281	turtle, bog (<i>Glyptemys muhlenbergii</i>) - U.S.A. (GA, NC, SC, TN, VA)	SAT	\$148,000	\$3,800	\$151,800	\$28,000	\$179,800
	225	turtle, bog (<i>Glyptemys muhlenbergii</i>) - Wherever found, except GA, NC, SC, TN, VA	T	\$286,000	\$36,284	\$322,284	\$0	\$322,284
	804	Turtle, flattened musk (<i>Sternotherus depressus</i>) -Black Warrior R. system upstream from Bankhead Dam	T	\$21,000	\$350	\$21,350	\$0	\$21,350
	553	Turtle, Plymouth Redbelly (<i>Pseudemys rubriventris bangsi</i>) -Wherever found	E	\$42,000	\$0	\$42,000	\$3,500	\$45,500
	641	Turtle, ringed map (<i>Graptemys oculifera</i>) - Wherever found	T	\$24,000	\$0	\$24,000	\$11,000	\$35,000
	308	Turtle, Sonoyta mud (<i>Kinosternon sonoriense longifemorale</i>) -Wherever found	E	\$0	\$146,896	\$146,896	\$0	\$146,896
	1333	Turtle, yellow-blotched map (<i>Graptemys flavimaculata</i>) -Wherever found	T	\$8,000	\$330	\$8,330	\$0	\$8,330
	165	Whipsnake (=striped racer), Alameda (<i>Masticophis lateralis euryxanthus</i>) - Wherever found	T	\$484,100	\$10,218	\$494,318	\$0	\$494,318
Reptiles subtotal				\$9,145,389	\$18,136,407	\$27,281,796	\$4,027,106	\$31,308,902
Snails	566	[no common name] Snail (<i>Eua zebrina</i>) - Wherever found	E	\$39,430	\$5,000	\$44,430	\$0	\$44,430
	566	[no common name] Snail (<i>Ostodes strigatus</i>) -Wherever found	E	\$39,430	\$5,000	\$44,430	\$0	\$44,430
	125	Abalone, Black (<i>Haliotis cracherodii</i>) - Wherever found	E	\$0	\$751,100	\$751,100	\$0	\$751,100
	72	Abalone, White (<i>Haliotis sorenseni</i>) -North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$0	\$1,673,100	\$1,673,100	\$0	\$1,673,100
	1152	Campeloma, slender (<i>Campeloma decampi</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	828	Cavesnail, Tumbling Creek (<i>Antrobia culveri</i>) -Wherever found	E	\$12,500	\$0	\$12,500	\$7,500	\$20,000
	792	Elimia, lacy (snail) (<i>Elimia crenatella</i>) -Wherever found	T	\$22,000	\$0	\$22,000	\$0	\$22,000
	859	Hornsnail, rough (<i>Pleurocera foremani</i>) -Wherever found	E	\$18,500	\$0	\$18,500	\$0	\$18,500
	335	Limpet, Banbury Springs (<i>Lanx</i> sp.) -Wherever found	E	\$116,626	\$5,700	\$122,326	\$0	\$122,326
	1473	Lioplax, cylindrical (snail) (<i>Lioplax cyclostomaformis</i>) -Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1639	Marstonia, royal (snail) (<i>Marstonia ogmorhappe</i>) -Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	1464	Pebblesnail, flat (<i>Lepyrium showalteri</i>) -Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	749	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$25,000	\$0	\$25,000	\$0	\$25,000
	858	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - Wherever found; Except where listed as Experimental Populations	E	\$18,000	\$635	\$18,635	\$0	\$18,635
	967	Rocksnail, interrupted (=Georgia) (<i>Leptoxis foremani</i>) -Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	732	Rocksnail, painted (<i>Leptoxis taeniata</i>) -Wherever found	T	\$26,000	\$0	\$26,000	\$0	\$26,000
	1507	Rocksnail, plicate (<i>Leptoxis plicata</i>) -Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1338	Rocksnail, round (<i>Leptoxis ampla</i>) -Wherever found	T	\$8,000	\$0	\$8,000	\$0	\$8,000
	1132	Snail, armored (<i>Marstonia pachyta</i>) -Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	458	Snail, Bliss Rapids (<i>Taylorconcha serpenticola</i>) -Wherever found	T	\$57,587	\$8,600	\$66,187	\$0	\$66,187

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	848	Snail, Chittenango ovate amber (Novisuccinea chittenangoensis) -Wherever found	T	\$19,000	\$0	\$19,000	\$0	\$19,000
	1185	Snail, flat-spined three-toothed (Triodopsis platysayoides) -Wherever found	T	\$9,500	\$0	\$9,500	\$0	\$9,500
	391	Snail, fragile tree (Samoana fragilis) - Wherever found	E	\$96,000	\$0	\$96,000	\$0	\$96,000
	336	Snail, Guam tree (Partula radiolata) - Wherever found	E	\$122,000	\$0	\$122,000	\$0	\$122,000
	272	Snail, Humped tree (Partula gibba) - Wherever found	E	\$193,000	\$0	\$193,000	\$0	\$193,000
	1152	Snail, Iowa Pleistocene (Discus macclintocki) -Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	618	Snail, Lanai tree (Partulina semicarinata) - Wherever found	E	\$37,130	\$0	\$37,130	\$0	\$37,130
	637	Snail, Lanai tree (Partulina variabilis) - Wherever found	E	\$35,130	\$0	\$35,130	\$0	\$35,130
	389	Snail, Langford's tree (Partula langfordi) - Wherever found	E	\$96,500	\$0	\$96,500	\$0	\$96,500
	452	Snail, Morro shoulderband (=Banded dune) (Helminthoglypta walkeriana) -Wherever found	T	\$68,000	\$0	\$68,000	\$0	\$68,000
	669	Snail, Newcomb's (Erinna newcombi) - Wherever found	T	\$31,000	\$0	\$31,000	\$0	\$31,000
	732	snail, noonday (Mesodon clarki nantahala) - Wherever found	T	\$26,000	\$0	\$26,000	\$0	\$26,000
	516	Snail, painted snake coiled forest (Anguispira picta) -Wherever found	T	\$32,500	\$0	\$32,500	\$21,000	\$53,500
	1591	Snail, Pecos assiminea (Assiminea pecos) - Wherever found	E	\$0	\$2,000	\$2,000	\$0	\$2,000
	330	Snail, Snake River physa (Physa natricina) - Wherever found	E	\$122,581	\$0	\$122,581	\$1,600	\$124,181
	1591	Snail, Stock Island tree (Orthalicus reses (not incl. nesodryas)) -Wherever found	T	\$2,000	\$0	\$2,000	\$0	\$2,000
	799	Snail, tulotoma (Tulotoma magnifica) - Wherever found	T	\$21,500	\$0	\$21,500	\$0	\$21,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	233	Snails, Oahu tree (Achatinella spp.) - Wherever found	E	\$287,298	\$0	\$287,298	\$0	\$287,298
	467	Springsnail, Bruneau Hot (Pyrgulopsis bruneauensis) -Wherever found	E	\$57,587	\$5,300	\$62,887	\$0	\$62,887
	1591	Springsnail, Koster's (Juturnia kosteri) - Wherever found	E	\$0	\$2,000	\$2,000	\$0	\$2,000
	1591	Springsnail, Roswell (Pyrgulopsis roswellensis) -Wherever found	E	\$0	\$2,000	\$2,000	\$0	\$2,000
	721	Tree snail, Newcomb's (Newcombia cumingi) -Wherever found	E	\$27,630	\$0	\$27,630	\$0	\$27,630
<i>Snails subtotal</i>				<i>\$1,727,429</i>	<i>\$2,460,435</i>	<i>\$4,187,864</i>	<i>\$30,100</i>	<i>\$4,217,964</i>
Multi-Species				\$12,264,636	\$1,795,300	\$14,059,936	\$5,709,444	\$19,769,380
<i>Multi-Species subtotal</i>				<i>\$12,264,636</i>	<i>\$1,795,300</i>	<i>\$14,059,936</i>	<i>\$5,709,444</i>	<i>\$19,769,380</i>
Total Species Expenditures				\$201,284,591	\$522,279,985	\$723,564,576	\$30,694,047	\$754,258,623

**TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2022
REPORTED EXPENDITURES**

Species are arranged in descending rank order of total identifiable expenditures (Table 1). Cumulative totals of the expenditures totals are provided. The cumulative total column is the sum of all higher ranked species totals (from Table 1) and the selected species entry. This table does not include costs for land acquisition or multi-species conservation activities.

The species list in the report follows the same breakdown as in 50 CFR 17.11 and 17.12. All populations are itemized separately, not just Evolutionarily Significant Units (ESUs). Thus, it appears as if there are more listed species. Separate populations of the same species may be ranked differently. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

**TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2022 REPORTED EXPENDITURES,
NOT INCLUDING LAND ACQUISITION COSTS**

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Middle Columbia River DPS - See 50 CFR 223.102	T	\$42,178,771	\$42,178,771
2	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Snake River Basin DPS - See 50 CFR 223.102	T	\$38,642,464	\$80,821,235
3	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River spring/summer-run ESU See 50 CFR 223.102	T	\$36,447,054	\$117,268,289
4	Whale, North Atlantic Right (<i>Eubalaena glacialis</i>) -Wherever found	E	\$29,440,195	\$146,708,484
5	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Columbia River spring-run ESU See 50 CFR 224.101	E	\$27,270,664	\$173,979,148
6	Trout, bull (<i>Salvelinus confluentus</i>) -U.S.A., conterminous, (lower 48 states)	T	\$23,210,732	\$197,189,880
7	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Upper Columbia River DPS - See 50 CFR 223.102	T	\$19,830,882	\$217,020,762
8	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River fall-run ESU See 50 CFR 223.102	T	\$18,401,582	\$235,422,344
9	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Lower Columbia River ESU See 50 CFR 223.102	T	\$18,332,772	\$253,755,116
10	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Oregon Coast ESU - See 50 CFR 223.102	T	\$13,652,483	\$267,407,599
11	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) -Snake River ESU - See 50 CFR 224.101	E	\$12,851,461	\$280,259,060
12	Sturgeon, white (<i>Acipenser transmontanus</i>) -U.S.A. (ID, MT), Canada (B.C.), (Kootenai R. system)	E	\$12,428,063	\$292,687,123
13	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Lower Columbia River DPS - See 50 CFR 223.102	T	\$12,223,662	\$304,910,785
14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Willamette River ESU See 50 CFR 223.102	T	\$11,310,060	\$316,220,845
15	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Lower Columbia River ESU - See 50 CFR 223.102	T	\$11,225,845	\$327,446,690
16	Trout, Lahontan cutthroat (<i>Oncorhynchus clarkii henshawi</i>) -Wherever found	T	\$10,349,034	\$337,795,724
17	Salmon, chum (<i>Oncorhynchus keta</i>) -Columbia River ESU See 50 CFR 223.102	T	\$9,680,097	\$347,475,821
18	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	T	\$9,475,064	\$356,950,885
19	Owl, northern spotted (<i>Strix occidentalis caurina</i>) -Wherever found	T	\$9,062,422	\$366,013,307
20	Sucker, shortnose (<i>Chasmistes brevirostris</i>) -Wherever found	E	\$8,755,101	\$374,768,408

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
21	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Upper Willamette River DPS - See 50 CFR 223.102	T	\$8,271,923	\$383,040,331
22	Manatee, West Indian (<i>Trichechus manatus</i>) -Wherever found	T	\$8,148,247	\$391,188,578
23	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Puget Sound ESU See 50 CFR 223.102	T	\$7,022,609	\$398,211,187
24	Bear, grizzly (<i>Ursus arctos horribilis</i>) -U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$6,823,434	\$405,034,621
25	salmon Atlantic (<i>Salmo salar</i>) -U.S.A. ME Gulf of Maine Atlantic Salmon Distinct Population Segment, which includes all naturally reproducing wild populations and those river-specific hatchery populations of Atlantic salmon having historical, river-specific characteristics found north of and including tributaries of the lower Kennebec River to, but not including, the mouth of the St. Croix River at the U.S.-Canda border. To date, the Services have determined that these populations are found in the Dennys, East Machias, Machias, Pleasant, Narraguagus, Sheepscot, and Ducktrap Rivers and in Cove Brook, Maine	E	\$6,800,000	\$411,834,621
26	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Puget Sound DPS - See 50 CFR 223.102	T	\$6,365,609	\$418,200,230
27	Sucker, Lost River (<i>Deltistes luxatus</i>) -Wherever found	E	\$6,193,934	\$424,394,164
28	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) -Central California Coast ESU - See 50 CFR 224.101	E	\$6,184,090	\$430,578,254
29	Salmon, chum (<i>Oncorhynchus keta</i>) -Hood Canal summer-run ESU - See 50 CFR 223.102	T	\$6,012,609	\$436,590,863
30	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) -Ozette Lake ESU - See 50 CFR 223.102	T	\$5,349,000	\$441,939,863
31	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Northern California DPS - See 50 CFR 223.102	T	\$5,332,000	\$447,271,863
32	Sea lion, Steller (<i>Eumetopias jubatus</i>) -Western DPS - See 50 CFR 224.101	E	\$5,020,944	\$452,292,807
33	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Sacramento River winter-run ESU - See 50 CFR 224.101	E	\$4,735,800	\$457,028,607
34	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Central California Coast DPS - See 50 CFR 223.102	T	\$4,676,000	\$461,704,607
35	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -California Coastal ESU - See 50 CFR 223.102	T	\$4,667,000	\$466,371,607
36	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -California Central Valley DPS - See 50 CFR 223.102	T	\$4,596,995	\$470,968,602
37	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -South-Central California Coast DPS - See 50 CFR 223.102	T	\$4,571,800	\$475,540,402

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
38	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Central Valley spring-run ESU See 50 CFR 223.102	T	\$4,567,500	\$480,107,902
39	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Southern California DPS - See 50 CFR 224.101	E	\$4,566,000	\$484,673,902
40	Seal, Hawaiian monk (<i>Monachus schauinslandi</i>) -Wherever found	E	\$4,362,000	\$489,035,902
41	Sea turtle, leatherback (<i>Dermochelys coriacea</i>) -Wherever found	E	\$4,186,689	\$493,222,591
42	Salamander, California tiger (<i>Ambystoma californiense</i>) -U.S.A. (CA - Sonoma County)	E	\$4,139,700	\$497,362,291
43	salmon Atlantic (<i>Salmo salar</i>) -Gulf of Maine DPS - See 50 CFR 224.101	E	\$3,716,000	\$501,078,291
44	Wolf, Mexican (<i>Canis lupus baileyi</i>) -Wherever found, except where included in an experimental population as set forth in 17.84(k)	E	\$3,649,996	\$504,728,287
45	Frog, Oregon spotted (<i>Rana pretiosa</i>) -Wherever found	T	\$3,631,687	\$508,359,974
46	whale, beluga (<i>Delphinapterus leucas</i>) -Cook Inlet DPS See 50 CFR 224.101	E	\$3,433,188	\$511,793,162
47	Sea turtle, loggerhead (<i>Caretta caretta</i>) -Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	T	\$3,424,176	\$515,217,338
48	Murrelet, marbled (<i>Brachyramphus marmoratus</i>) -U.S.A. (CA, OR, WA)	T	\$3,277,122	\$518,494,460
49	Bat, Northern long-eared (<i>Myotis septentrionalis</i>) -Wherever found	E	\$3,125,470	\$521,619,930
50	`I'iwi (<i>Drepanis coccinea</i>) -Wherever found	T	\$2,823,998	\$524,443,928
51	Sea turtle, Kemp's ridley (<i>Lepidochelys kempii</i>) -Wherever found	E	\$2,800,240	\$527,244,168
52	Sea turtle, hawksbill (<i>Eretmochelys imbricata</i>) -Wherever found	E	\$2,791,346	\$530,035,514
53	Tortoise, desert (<i>Gopherus agassizii</i>) -Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$2,749,167	\$532,784,681
54	Elephant, African (<i>Loxodonta africana</i>) -Wherever found	T	\$2,723,446	\$535,508,127
55	parrotbill, Maui (<i>Kiwikiu</i>) (<i>Pseudonestor xanthophrys</i>) -Wherever found	E	\$2,693,535	\$538,201,662
56	Smelt, delta (<i>Hypomesus transpacificus</i>) -Wherever found	T	\$2,668,019	\$540,869,681
57	Lynx, Canada (<i>Lynx canadensis</i>) -Wherever Found in Contiguous U.S.	T	\$2,517,462	\$543,387,143
58	Akohekohe (<i>Palmeria dolei</i>) -Wherever found	E	\$2,514,060	\$545,901,203
59	Bat, Indiana (<i>Myotis sodalis</i>) -Wherever found	E	\$2,483,856	\$548,385,059
60	Whale, killer (<i>Orcinus orca</i>) -Southern Resident DPS See 50 CFR 224.101	E	\$2,402,100	\$550,787,159
61	Ferret, black-footed (<i>Mustela nigripes</i>) -U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$2,387,709	\$553,174,868
62	Rhinoceros, black (<i>Diceros bicornis</i>) -Sub-Saharan Africa	E	\$2,381,363	\$555,556,231
63	Woodpecker, red-cockaded (<i>Picoides borealis</i>) -Wherever found	E	\$2,249,029	\$557,805,260

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
64	Wolf, gray (<i>Canis lupus</i>) -U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of Highway 97 and Highway 17 north of Mesa and that portion of WA west of the centerline of Highway 395 south of Mesa). Mexico.	E	\$2,165,567	\$559,970,827
65	Alligator, American (<i>Alligator mississippiensis</i>) -Wherever found	SAT	\$2,056,548	\$562,027,375
66	Plover, piping (<i>Charadrius melodus</i>) -[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$1,915,104	\$563,942,479
67	Seal, bearded (<i>Erignathus barbatus nauticus</i>) -Beringia DPS - See 50 CFR 223.102	T	\$1,891,000	\$565,833,479
67	Seal, Ringed (<i>Phoca</i> (=Pusa) <i>hispida botnica</i>) -Wherever found	T	\$1,891,000	\$567,724,479
69	Cuckoo, yellow-billed (<i>Coccyzus americanus</i>) -Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern)); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	T	\$1,889,697	\$569,614,176
70	Deer, Columbian white-tailed (<i>Odocoileus virginianus leucurus</i>) -Columbia River (Clark, Cowliz, Pacific, Skamania, and Wahkiakum Counties, WA., and Clatsop, Columbia, and Multnomah Counties, OR.)	T	\$1,880,884	\$571,495,060
71	Elephant, Asian (<i>Elephas maximus</i>) -Wherever found	E	\$1,713,060	\$573,208,120
72	Abalone, White (<i>Haliotis sorenseni</i>) -North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$1,673,100	\$574,881,220
73	Sturgeon, shortnose (<i>Acipenser brevirostrum</i>) -Wherever found	E	\$1,647,195	\$576,528,415
74	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	T	\$1,641,545	\$578,169,960
75	Bear, polar (<i>Ursus maritimus</i>) -Wherever found	T	\$1,623,402	\$579,793,362

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
76	Trout, bull (<i>Salvelinus confluentus</i>) -Clackamas River subbasin and the mainstem Willamette River, from Willamette Falls to its points of confluence with the Columbia River, including Multnomah Channel	EXPN	\$1,582,209	\$581,375,571
77	Sea turtle, loggerhead (<i>Caretta caretta</i>) -North Pacific Ocean DPS - Loggerhead sea turtles originating from the North Pacific north of the equator and south of 60 degrees N. Lat	E	\$1,530,534	\$582,906,105
78	Plover, western snowy (<i>Charadrius nivosus nivosus</i>) -Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$1,508,501	\$584,414,606
79	Parrot, Puerto Rican (<i>Amazona vittata</i>) -Wherever found	E	\$1,430,000	\$585,844,606
80	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Carolina DPS - See 50 CFR 224.101	E	\$1,410,900	\$587,255,506
81	Panther, Florida (<i>Puma</i> (=Felis) <i>concolor coryi</i>) -Wherever found	E	\$1,381,361	\$588,636,867
82	Wolf, red (<i>Canis rufus</i>) -Wherever found, except where listed as an experimental population	E	\$1,328,100	\$589,964,967
83	sage-grouse, Gunnison (<i>Centrocercus minimus</i>) -Wherever found	T	\$1,299,443	\$591,264,410
84	Whale, finback (<i>Balaenoptera physalus</i>) -Wherever found	E	\$1,260,000	\$592,524,410
85	Whale, blue (<i>Balaenoptera musculus</i>) -Wherever found	E	\$1,255,800	\$593,780,210
86	Heelsplitter, Carolina (<i>Lasmigona decorata</i>) -Wherever found	E	\$1,237,300	\$595,017,510
87	Tiger (<i>Panthera tigris</i>) -Wherever found	E	\$1,221,178	\$596,238,688
88	Sawfish, smalltooth (<i>Pristis pectinata</i>) -US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$1,210,127	\$597,448,815
89	Sturgeon, Gulf (<i>Acipenser oxyrinchus</i> (=oxyrhynchus) <i>desotoi</i>) -Wherever found	T	\$1,176,934	\$598,625,749
90	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	T	\$1,165,429	\$599,791,178
91	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Central West Pacific Ocean, bounded by the following coordinates: 41 degrees N., 146 degrees E. in the northwest; 41 degrees N., 169 degrees E. in the northeast; 9 degrees N., 175 degrees W. in the east; 13 degrees S., 171 degrees E. in the southeast; along the northern coast of the island of New Guinea; and 4.5 degrees N., 129 degrees E. in the west	E	\$1,143,000	\$600,934,178
92	Bat, gray (<i>Myotis grisescens</i>) -Wherever found	E	\$1,137,513	\$602,071,691
93	Plover, piping (<i>Charadrius melodus</i>) -[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$1,119,133	\$603,190,824
94	Frog, California red-legged (<i>Rana draytonii</i>) -Wherever found	T	\$1,086,156	\$604,276,980

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
95	Orangutan (<i>Pongo pygmaeus</i>) -Wherever found	E	\$1,079,000	\$605,355,980
96	Whale, Sei (<i>Balaenoptera borealis</i>) -Wherever found	E	\$1,071,300	\$606,427,280
97	Whale, sperm (<i>Physeter catodon</i> (=macrocephalus)) -Wherever found	E	\$1,071,100	\$607,498,380
98	petrel, Hawaiian (<i>Pterodroma sandwichensis</i>) -Wherever found	E	\$1,045,789	\$608,544,169
99	Ferret, black-footed (<i>Mustela nigripes</i>) -Wherever found, except where listed as an experimental population	E	\$1,034,854	\$609,579,023
100	Flycatcher, southwestern willow (<i>Empidonax traillii extimus</i>) -Wherever found	E	\$1,010,358	\$610,589,381
101	rail, Eastern Black (<i>Laterallus jamaicensis</i> ssp. <i>jamaicensis</i>) -Wherever found	T	\$998,962	\$611,588,343
102	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the East Pacific Ocean, bounded by the following lines and coordinates: 41 degrees N., 143 degrees W. in the northwest; 41 degrees N. Lat. in the north; along the western coasts of the Americas; 40 degrees S. Lat. in the south; and 40 degrees S., 96 degrees W. in the southwest	T	\$941,000	\$612,529,343
103	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -South Atlantic DPS - See 50 CFR 224.101	E	\$939,910	\$613,469,253
104	Goose, Hawaiian (<i>Branta</i> (=Nesochen) <i>sandvicensis</i>) -Wherever found	T	\$917,310	\$614,386,563
105	Fisher (<i>Pekania pennanti</i>) -West Coast Distinct Population Segment	E	\$914,340	\$615,300,903
106	Pupfish, Devils Hole (<i>Cyprinodon diabolis</i>) -Wherever found	E	\$908,100	\$616,209,003
107	Sucker, Warner (<i>Catostomus warnerensis</i>) -Wherever found	T	\$900,500	\$617,109,503
108	Rabbit, Pygmy (<i>Brachylagus idahoensis</i>) -U.S.A. (WA-Douglas, Grant, Lincoln, Adams, Benton Counties)	E	\$883,269	\$617,992,772
109	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Chesapeake Bay DPS - See 50 CFR 224.101	E	\$859,500	\$618,852,272
110	Coral, elkhorn (<i>Acropora palmata</i>) -Wherever found	T	\$858,000	\$619,710,272
111	Coral, staghorn (<i>Acropora cervicornis</i>) -Wherever found	T	\$851,700	\$620,561,972
112	Crow, Mariana (=aga) (<i>Corvus kubaryi</i>) -Wherever found	E	\$845,897	\$621,407,869
113	Shearwater, Newell's Townsend's (<i>Puffinus auricularis newelli</i>) -Wherever found	T	\$840,906	\$622,248,775
114	Coral, Mountainous Star (<i>Orbicella faveolata</i>) -Wherever found	T	\$838,700	\$623,087,475
115	Crow, Hawaiian (=‘alala) (<i>Corvus hawaiiensis</i>) -Wherever found	E	\$827,695	\$623,915,170
116	Cui-ui (<i>Chasmistes cujus</i>) -Wherever found	E	\$809,700	\$624,724,870
117	Caracara, Audubon's crested (<i>Polyborus plancus audubonii</i>) -U.S.A. (FL)	T	\$809,500	\$625,534,370
118	grouper, Nassau (<i>Epinephelus striatus</i>) -	T	\$794,700	\$626,329,070
119	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. only, except where listed as an experimental population	E	\$794,490	\$627,123,560
120	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -New York Bight DPS - See 50 CFR 224.101	E	\$792,564	\$627,916,124
121	Bumble bee, Rusty patched (<i>Bombus affinis</i>) -Wherever found	E	\$791,095	\$628,707,219

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
122	Coral, Lobed Star (<i>Orbicella annularis</i>) -Wherever found	T	\$780,700	\$629,487,919
123	Salamander, California tiger (<i>Ambystoma californiense</i>) -U.S.A. (CA - Santa Barbara County)	E	\$764,137	\$630,252,056
124	Eider, spectacled (<i>Somateria fischeri</i>) -Wherever found	T	\$760,483	\$631,012,539
125	Abalone, Black (<i>Haliotis cracherodii</i>) -Wherever found	E	\$751,100	\$631,763,639
126	Fox, San Joaquin kit (<i>Vulpes macrotis mutica</i>) -wherever found	E	\$744,659	\$632,508,298
127	coral, Pillar (<i>Dendrogyra cylindrus</i>) -Wherever found	T	\$738,700	\$633,246,998
128	Bison, wood (<i>Bison bison athabasca</i>) -U.S.A. (Alaska)	EXPN	\$738,632	\$633,985,630
129	Vole, Amargosa (<i>Microtus californicus scirpensis</i>) -Wherever found	E	\$738,500	\$634,724,130
130	Knot, red (<i>Calidris canutus rufa</i>) -Wherever found	T	\$738,440	\$635,462,570
131	coral, Boulder star (<i>Orbicella franksi</i>) -Wherever found	T	\$731,200	\$636,193,770
132	Chub, humpback (<i>Gila cypha</i>) -Wherever found	T	\$724,291	\$636,918,061
133	Crane, Mississippi sandhill (<i>Grus canadensis pulla</i>) -Wherever found	E	\$708,000	\$637,626,061
134	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) -Wherever found	T	\$694,992	\$638,321,053
135	Jaguar (<i>Panthera onca</i>) -Wherever found	E	\$691,854	\$639,012,907
136	Checkerspot, Taylor's (=whulge) (<i>Euphydryas editha taylori</i>) -Wherever found	E	\$691,049	\$639,703,956
137	Fairy shrimp, vernal pool (<i>Branchinecta lynchi</i>) -Wherever found	T	\$686,088	\$640,390,044
138	Prairie dog, Utah (<i>Cynomys parvidens</i>) -Wherever found	T	\$671,181	\$641,061,225
139	Marten, Pacific (<i>Martes caurina</i>) -Wherever found	T	\$659,483	\$641,720,708
140	Coral, Rough Cactus (<i>Mycetophyllia ferox</i>) -Wherever found	T	\$656,200	\$642,376,908
141	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Central South Pacific Ocean, bounded by the following coordinates: 9 degrees N., 175 degrees W. in the northwest; 9 degrees N., 125 degrees W. in the northeast; 40 degrees S., 96 degrees W. in the southeast; 40 degrees S., 176 degrees E. in the southwest; and 13 degrees S., 171 degrees E. in the west	E	\$651,250	\$643,028,158
142	sturgeon, Green (<i>Acipenser medirostris</i>) -Southern DPS - See 50 CFR 223.102	T	\$647,648	\$643,675,806
143	Beetle, valley elderberry longhorn (<i>Desmocerus californicus dimorphus</i>) -Wherever found	T	\$645,391	\$644,321,197
144	Peppergrass, Slickspot (<i>Lepidium papilliferum</i>) -	T	\$622,849	\$644,944,046
145	Gorilla (<i>Gorilla gorilla</i>) -Wherever found	E	\$616,842	\$645,560,888
146	No Common Name (<i>Pinus albicaulis</i>) -	T	\$602,692	\$646,163,580
147	Tadpole shrimp, vernal pool (<i>Lepidurus packardii</i>) -Wherever found	E	\$583,701	\$646,747,281
148	Otter, southern sea (<i>Enhydra lutris nereis</i>) -Wherever found	T	\$580,160	\$647,327,441
149	Pearlshell, Louisiana (<i>Margaritifera hembeli</i>) -Wherever found	T	\$572,000	\$647,899,441
150	Mouse, salt marsh harvest (<i>Reithrodontomys raviventris</i>) -wherever found	E	\$571,300	\$648,470,741
151	Eider, Steller's (<i>Polysticta stelleri</i>) -U.S. Alaska breeding population only	T	\$570,600	\$649,041,341
152	Fox, Sierra Nevada red (<i>Vulpes vulpes necator</i>) -Sierra Nevada DPS	E	\$565,664	\$649,607,005

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
153	Bocaccio (<i>Sebastes paucispinis</i>) -Puget Sound-Georgia Basin DPS See 50 CFR 224.101	E	\$559,000	\$650,166,005
153	Eulachon (<i>Thaleichthys pacificus</i>) -Southern DPS - See 50 CFR 223.102	T	\$559,000	\$650,725,005
153	rockfish, yelloweye (<i>Sebastes ruberrimus</i>) -Puget Sound - Georgia Basin DPS See 50 CFR 223.102	T	\$559,000	\$651,284,005
156	Sturgeon, pallid (<i>Scaphirhynchus albus</i>) -Wherever found	E	\$556,046	\$651,840,051
157	Bat, Hawaiian hoary (<i>Lasiurus cinereus semotus</i>) -Wherever found	E	\$554,312	\$652,394,363
158	Sparrow, Cape Sable seaside (<i>Ammodramus maritimus mirabilis</i>) -Wherever found	E	\$537,500	\$652,931,863
159	Darter, boulder (<i>Etheostoma wapiti</i>) -Wherever found	E	\$532,000	\$653,463,863
160	Crane, whooping (<i>Grus americana</i>) -Wherever found, except where listed as an experimental population	E	\$530,219	\$653,994,082
161	Snake, eastern indigo (<i>Drymarchon couperi</i>) -Wherever found	T	\$522,224	\$654,516,306
162	Horned lark, streaked (<i>Eremophila alpestris strigata</i>) -Wherever found	T	\$511,233	\$655,027,539
163	Mussel, snuffbox (<i>Epioblasma triquetra</i>) -Wherever found	E	\$503,539	\$655,531,078
164	Kite, Everglade snail (<i>Rostrhamus sociabilis plumbeus</i>) -Wherever found	E	\$497,743	\$656,028,821
165	Whipsnake (=striped racer), Alameda (<i>Masticophis lateralis euryxanthus</i>) -Wherever found	T	\$494,318	\$656,523,139
166	Butterfly, Fender's blue (<i>Icaricia icarioides fenderi</i>) -Wherever found	E	\$492,636	\$657,015,775
167	bat, Florida bonneted (<i>Eumops floridanus</i>) -Wherever found	E	\$491,451	\$657,507,226
168	Wedgemussel, dwarf (<i>Alasmodonta heterodon</i>) -Wherever found	E	\$490,500	\$657,997,726
169	scrub-jay, Florida (<i>Aphelocoma coerulescens</i>) -Wherever found	T	\$490,331	\$658,488,057
170	Rail, California clapper (<i>Rallus longirostris obsoletus</i>) -Wherever found	E	\$490,102	\$658,978,159
171	Rhinoceros, great Indian (<i>Rhinoceros unicornis</i>) -Nepal, India	E	\$487,463	\$659,465,622
172	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	T	\$487,443	\$659,953,065
173	Ladies'-tresses, Ute (<i>Spiranthes diluvialis</i>) -Wherever found	T	\$473,103	\$660,426,168
174	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) -Gulf of Maine DPS - See 50 CFR 223.102	T	\$470,500	\$660,896,668
175	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) -Wherever found, except where listed as an experimental population	E	\$461,038	\$661,357,706
176	Rail, Yuma Ridgway's (<i>Rallus obsoletus yumanensis</i>) -Wherever found	E	\$460,331	\$661,818,037

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
177	Otter, Northern Sea (<i>Enhydra lutris kenyoni</i>) -Southwest Alaska, from Attu Island to Western Cook Inlet, including Bristol Bay, the Kodiak Archipelago, and the Barren Islands	T	\$458,908	\$662,276,945
178	Goby, tidewater (<i>Eucyclogobius newberryi</i>) -Wherever found	E	\$458,800	\$662,735,745
179	Chimpanzee (<i>Pan troglodytes</i>) -Wherever found	E	\$447,792	\$663,183,537
180	paintbrush, golden (<i>Castilleja levisecta</i>) -Wherever found	T	\$447,217	\$663,630,754
181	Owl, Mexican spotted (<i>Strix occidentalis lucida</i>) -Wherever found	T	\$440,513	\$664,071,267
182	Vireo, least Bell's (<i>Vireo bellii pusillus</i>) -Wherever found	E	\$437,348	\$664,508,615
183	Sea turtle, loggerhead (<i>Caretta caretta</i>) -Northeast Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northeast Atlantic Ocean north of the equator, south of 60 degrees N. Lat., and east of 40 degrees W. Long., except in the vicinity of the Strait of Gibraltar where the eastern boundary is 5 degrees 36 minutes W. Long	E	\$436,700	\$664,945,315
184	Frog, Sierra Nevada Yellow-legged (<i>Rana sierrae</i>) -Wherever found	E	\$435,607	\$665,380,922
185	Rhinoceros, Sumatran (<i>Dicerorhinus sumatrensis</i>) -Wherever found	E	\$427,709	\$665,808,631
186	Tern, California least (<i>Sterna antillarum browni</i>) -Wherever found	E	\$421,700	\$666,230,331
187	Spiny mussel, James (<i>Parvaspina collina</i>) -Wherever found	E	\$421,100	\$666,651,431
188	Whale, bowhead (<i>Balaena mysticetus</i>) -Wherever found	E	\$407,000	\$667,058,431
189	Logperch, Roanoke (<i>Percina rex</i>) -Wherever found	E	\$404,500	\$667,462,931
190	Massasauga (=rattlesnake), eastern (<i>Sistrurus catenatus</i>) -Wherever found	T	\$403,214	\$667,866,145
191	Wolf, gray (<i>Canis lupus</i>) -U.S.A. (MN)	T	\$400,052	\$668,266,197
192	Akikiki (<i>Oreomystis bairdi</i>) -Wherever found	E	\$398,560	\$668,664,757
193	Catchfly, Spalding's (<i>Silene spaldingii</i>) -Wherever found	T	\$393,134	\$669,057,891
194	No common name (<i>Seriatopora aculeata</i>) -Wherever found	T	\$393,000	\$669,450,891
195	Kangaroo rat, San Bernardino Merriam's (<i>Dipodomys merriami parvus</i>) -Wherever found	E	\$392,100	\$669,842,991
196	Tortoise, gopher (<i>Gopherus polyphemus</i>) -Western DPS	T	\$386,480	\$670,229,471
197	Beetle, American burying (<i>Nicrophorus americanus</i>) -Wherever found, except where listed as an experimental population	T	\$384,330	\$670,613,801
198	Whale, North Pacific Right (<i>Eubalaena japonica</i>) -Wherever found	E	\$381,100	\$670,994,901
199	Whale, humpback (<i>Megaptera novaeangliae</i>) -Western North Pacific	E	\$381,000	\$671,375,901
200	No common name (<i>Acropora globiceps</i>) -Wherever found	T	\$378,500	\$671,754,401
201	No common name (<i>Acropora retusa</i>) -Wherever found	T	\$378,000	\$672,132,401
201	No common name (<i>Isopora crateriformis</i>) -Wherever found	T	\$378,000	\$672,510,401
203	No common name (<i>Acropora jacquelineae</i>) -Wherever found	T	\$377,500	\$672,887,901
203	No common name (<i>Acropora speciosa</i>) -Wherever found	T	\$377,500	\$673,265,401
203	No common name (<i>Euphyllia paradivisa</i>) -Wherever found	T	\$377,500	\$673,642,901
206	Butterfly, Island marble (<i>Euchloe ausonides insulanus</i>) -Wherever found	E	\$373,804	\$674,016,705

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
207	Sparrow, Florida grasshopper (<i>Ammodramus savannarum floridanus</i>) -Wherever found	E	\$372,139	\$674,388,844
208	Squirrel, Northern Idaho Ground (<i>Uroditellus brunneus</i>) -Wherever found	T	\$370,957	\$674,759,801
209	Stork, wood (<i>Mycteria americana</i>) -U.S.A. (AL, FL, GA, MS, NC, SC)	T	\$370,805	\$675,130,606
210	Fisher (<i>Pekania pennanti</i>) -Southern Sierra Nevada Distinct Population Segment	E	\$367,540	\$675,498,146
211	Snake, giant garter (<i>Thamnophis gigas</i>) -Wherever found	T	\$363,508	\$675,861,654
212	Whale, humpback (<i>Megaptera novaeangliae</i>) -Mexico DPS	T	\$361,000	\$676,222,654
213	Gibbons (<i>Hylobates</i> spp. (including <i>Nomascus</i>)) -Wherever found	E	\$360,367	\$676,583,021
214	Stilt, Hawaiian (<i>Himantopus mexicanus knudseni</i>) -Wherever found	E	\$359,131	\$676,942,152
215	Whale, humpback (<i>Megaptera novaeangliae</i>) -Central America DPS	E	\$357,800	\$677,299,952
216	Frog, dusky gopher (<i>Rana sevosa</i>) -Wherever found	E	\$348,118	\$677,648,070
217	Butterfly, Karner blue (<i>Lycaeides melissa samuelis</i>) -Wherever found	E	\$342,626	\$677,990,696
218	Lupine, Kincaid's (<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>) -Wherever found	T	\$341,221	\$678,331,917
219	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) -Wherever found	T	\$335,531	\$678,667,448
220	Darter, trispot (<i>Etheostoma trisella</i>) -Wherever found	T	\$333,000	\$679,000,448
221	Akekee (<i>Loxops caeruleirostris</i>) -Wherever found	E	\$329,895	\$679,330,343
222	trout, Greenback Cutthroat (<i>Oncorhynchus clarkii stomias</i>) -Wherever found	T	\$328,558	\$679,658,901
223	Albatross, short-tailed (<i>Phoebastria</i> (=Diomedea) <i>albatrus</i>) -Wherever found	E	\$328,491	\$679,987,392
224	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) -Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$325,220	\$680,312,612
225	turtle, bog (<i>Glyptemys muhlenbergii</i>) -Wherever found, except GA, NC, SC, TN, VA	T	\$322,284	\$680,634,896
226	Crane, whooping (<i>Grus americana</i>) -U.S.A (Southwestern Louisiana)	EXPN	\$308,000	\$680,942,896
227	Gnatcatcher, coastal California (<i>Poliophtila californica californica</i>) -Wherever found	T	\$304,500	\$681,247,396
228	Ocelot (<i>Leopardus</i> (=Felis) <i>pardalis</i>) -wherever found	E	\$300,910	\$681,548,306
229	Rabbitsfoot (<i>Quadrula cylindrica cylindrica</i>) -Wherever found	T	\$299,365	\$681,847,671
230	Pogonia, small whorled (<i>Isotria medeoloides</i>) -	T	\$297,300	\$682,144,971
231	Kingfisher, Guam Micronesian (<i>Halcyon cinnamomina cinnamomina</i>) -Wherever found	E	\$292,508	\$682,437,479
232	Rabbit, riparian brush (<i>Sylvilagus bachmani riparius</i>) -Wherever found	E	\$287,700	\$682,725,179
233	Snails, Oahu tree (<i>Achatinella</i> spp.) -Wherever found	E	\$287,298	\$683,012,477
234	Toad, Wyoming (<i>Anaxyrus baxteri</i>) -Wherever found	E	\$270,370	\$683,282,847
235	Woolly-threads, San Joaquin (<i>Monolopia</i> (=Lembertia) <i>congdonii</i>) -Wherever found	E	\$268,093	\$683,550,940
236	Lizard, blunt-nosed leopard (<i>Gambelia silus</i>) -Wherever found	E	\$268,012	\$683,818,952
237	Kangaroo rat, Tipton (<i>Dipodomys nitratoide nitratoide</i>) -Wherever found	E	\$264,460	\$684,083,412
238	Crane, whooping (<i>Grus americana</i>) -U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$261,900	\$684,345,312
239	Pocket gopher, Olympia (<i>Thomomys mazama pugetensis</i>) -Wherever found	T	\$261,733	\$684,607,045
240	whale, false killer (<i>Pseudorca crassidens</i>) -Main Hawaiian Islands Insular DPS See 50 CFR 224.101	E	\$261,000	\$684,868,045

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
241	coot, Hawaiian (<i>Fulica americana alai</i>) -Wherever found	E	\$260,061	\$685,128,106
242	Fox, Santa Catalina Island (<i>Urocyon littoralis catalinae</i>) -Wherever found	T	\$255,200	\$685,383,306
243	Checker-mallow, Nelson's (<i>Sidalcea nelsoniana</i>) -Wherever found	T	\$251,134	\$685,634,440
244	Mouse, Preble's meadow jumping (<i>Zapus hudsonius preblei</i>) -wherever found	T	\$248,554	\$685,882,994
245	Sucker, Santa Ana (<i>Catostomus santaanae</i>) -Los Angeles River basin, San Gabriel River basin, Santa Ana River basin	T	\$243,800	\$686,126,794
246	Crocodile, American (<i>Crocodylus acutus</i>) -U.S.A. (FL)	T	\$243,500	\$686,370,294
247	Rail, Guam (<i>Rallus owstoni</i>) -Wherever found, except where listed as an experimental population	E	\$239,679	\$686,609,973
248	bumble bee, Franklin's (<i>Bombus franklini</i>) -Wherever found	E	\$239,562	\$686,849,535
249	Yosemite Toad (<i>Anaxyrus canorus</i>) -Wherever found	T	\$237,491	\$687,087,026
250	gallinule, Hawaiian common (<i>Gallinula galeata sandvicensis</i>) -Wherever found	E	\$234,140	\$687,321,166
251	Bladderpod, White Bluffs (<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>) -Wherever found	T	\$234,088	\$687,555,254
252	Butterfly, Mitchell's satyr (<i>Neonympha mitchellii mitchellii</i>) -Wherever found	E	\$230,876	\$687,786,130
253	Pinesnake, Louisiana (<i>Pituophis ruthveni</i>) -Wherever found	T	\$230,800	\$688,016,930
254	Seal, Guadalupe fur (<i>Arctocephalus townsendi</i>) -Wherever found	T	\$227,800	\$688,244,730
255	Skipper, Dakota (<i>Hesperia dacotae</i>) -Wherever found	T	\$227,643	\$688,472,373
256	Mucket, pink (pearlymussel) (<i>Lampsilis abrupta</i>) -Wherever found	E	\$224,737	\$688,697,110
257	Kangaroo rat, Stephens' (<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)) -Wherever found	T	\$220,500	\$688,917,610
258	Storm-petrel, band-rumped (<i>Oceanodroma castro</i>) -USA (HI)	E	\$219,446	\$689,137,056
259	Crayfish, Shasta (<i>Pacifastacus fortis</i>) -Wherever found	E	\$215,601	\$689,352,657
260	Chub, Virgin River (<i>Gila seminuda</i> (=robusta)) -Wherever found	E	\$215,173	\$689,567,830
261	Darter, Pearl (<i>Percina aurora</i>) -Wherever found	T	\$213,500	\$689,781,330
262	Spectaclecase (mussel) (<i>Cumberlandia monodonta</i>) -Wherever found	E	\$212,361	\$689,993,691
263	Butterfly, mission blue (<i>Icaricia icarioides missionensis</i>) -Wherever found	E	\$206,100	\$690,199,791
264	Sucker, razorback (<i>Xyrauchen texanus</i>) -Wherever found	E	\$205,699	\$690,405,490
265	Mussel, sheepnose (<i>Plethobasus cyphus</i>) -Wherever found	E	\$203,063	\$690,608,553
266	Crayfish, Big Sandy (<i>Cambarus callainus</i>) -Wherever found	T	\$202,000	\$690,810,553
267	Higgins eye (pearlymussel) (<i>Lampsilis higginsii</i>) -Wherever found	E	\$201,781	\$691,012,334
268	Four-o'clock, MacFarlane's (<i>Mirabilis macfarlanei</i>) -Wherever found	T	\$201,302	\$691,213,636
269	Palila (honeycreeper) (<i>Loxioides bailleui</i>) -Wherever found	E	\$200,341	\$691,413,977
270	Orchid, white fringeless (<i>Platanthera integrilabia</i>) -	T	\$198,500	\$691,612,477
271	Bankclimber, purple (mussel) (<i>Elliptioideus sloatianus</i>) -Wherever found	T	\$197,493	\$691,809,970
272	Snail, Humped tree (<i>Partula gibba</i>) -Wherever found	E	\$193,000	\$692,002,970
273	Dragonfly, Hine's emerald (<i>Somatochlora hineana</i>) -Wherever found	E	\$192,547	\$692,195,517
274	Beauty, Harper's (<i>Harperocallis flava</i>) -	E	\$191,500	\$692,387,017
275	Pupfish, desert (<i>Cyprinodon macularius</i>) -Wherever found	E	\$189,794	\$692,576,811

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
276	Rhinoceros, Javan (<i>Rhinoceros sondaicus</i>) -Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$186,000	\$692,762,811
277	Butterfly, Hermes copper (<i>Lycaena hermes</i>) -Wherever found	T	\$185,300	\$692,948,111
278	Duck, Hawaiian (=koloa) (<i>Anas wyvilliana</i>) -Wherever found	E	\$185,241	\$693,133,352
279	Darter, candy (<i>Etheostoma osburni</i>) -Wherever found	E	\$185,071	\$693,318,423
280	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) -Wherever found, except where listed as an experimental population	E	\$183,953	\$693,502,376
281	turtle, bog (<i>Glyptemys muhlenbergii</i>) -U.S.A. (GA, NC, SC, TN, VA)	SAT	\$179,800	\$693,682,176
282	Cactus, Colorado hookless (<i>Sclerocactus glaucus</i>) -Wherever found	T	\$178,426	\$693,860,602
283	Dace, desert (<i>Eremichthys acros</i>) -Wherever found	T	\$177,800	\$694,038,402
284	Toad, Dixie Valley (<i>Anaxyrus williamsi</i>) -	E	\$177,000	\$694,215,402
285	Woundfin (<i>Plagopterus argentissimus</i>) -Gila R. drainage, AZ, NM	EXPN	\$170,000	\$694,385,402
286	Lion (<i>Panthera leo</i> ssp. <i>leo</i>) -Wherever found	E	\$168,533	\$694,553,935
287	skipperling, Poweshiek (<i>Oarisma poweshiek</i>) -Wherever found	E	\$165,776	\$694,719,711
288	Bonytail (<i>Gila elegans</i>) -Wherever found	E	\$165,364	\$694,885,075
289	Bat, Virginia big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii virginianus</i>) -Wherever found	E	\$164,981	\$695,050,056
290	Amaranth, seabeach (<i>Amaranthus pumilus</i>) -Wherever found	T	\$162,502	\$695,212,558
291	Mouse, Alabama beach (<i>Peromyscus polionotus ammobates</i>) -Wherever found	E	\$161,490	\$695,374,048
292	Skipper, Laguna Mountains (<i>Pyrgus ruralis lagunae</i>) -Wherever found	E	\$160,200	\$695,534,248
293	Desert buckwheat, Umtanum (<i>Eriogonum codium</i>) -Wherever found	T	\$158,228	\$695,692,476
294	Orchid, western prairie fringed (<i>Platanthera praeclara</i>) -Wherever found	T	\$158,139	\$695,850,615
295	Rail, light-footed clapper (<i>Rallus longirostris levipes</i>) -Wherever found	E	\$156,600	\$696,007,215
296	Tern, Roseate (<i>Sterna dougallii dougallii</i>) -U.S.A. (Atlantic Coast south to NC), Canada (Newf., N.S, Que.), Bermuda	E	\$155,900	\$696,163,115
297	Lance, yellow (<i>Elliptio lanceolata</i>) -Wherever found	T	\$155,500	\$696,318,615
298	Orchid, eastern prairie fringed (<i>Platanthera leucophaea</i>) -Wherever found	T	\$152,638	\$696,471,253
299	Frog, mountain yellow-legged (<i>Rana muscosa</i>) -U.S.A., southern California	E	\$152,600	\$696,623,853
300	Frog, mountain yellow-legged (<i>Rana muscosa</i>) -U.S.A., northern California	E	\$152,207	\$696,776,060
301	Fairy shrimp, Conservancy (<i>Branchinecta conservatio</i>) -Wherever found	E	\$151,536	\$696,927,596
302	Checkermallow, Wenatchee Mountains (<i>Sidalcea oregana</i> var. <i>calva</i>) -Wherever found	E	\$151,474	\$697,079,070
303	Cactus, Mesa Verde (<i>Sclerocactus mesae-verdae</i>) -Wherever found	T	\$151,322	\$697,230,392
304	pikeminnow, Colorado (<i>Ptychocheilus lucius</i>) -Wherever found, except where listed as an experimental population	E	\$151,139	\$697,381,531
305	Wolf, Mexican (<i>Canis lupus baileyi</i>) -U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$150,910	\$697,532,441
306	salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) -Wherever found	E	\$149,531	\$697,681,972
307	No common name (<i>Phyllostegia brevidens</i>) -Wherever found	E	\$148,901	\$697,830,873

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
308	Turtle, Sonoyta mud (<i>Kinosternon sonoriense longifemorale</i>) -Wherever found	E	\$146,896	\$697,977,769
309	Swiftlet, Mariana gray (<i>Aerodramus vanikorensis bartschi</i>) -Wherever found	E	\$144,700	\$698,122,469
310	Salamander, Cheat Mountain (<i>Plethodon nettingi</i>) -Wherever found	T	\$143,637	\$698,266,106
311	Salamander, California tiger (<i>Ambystoma californiense</i>) -U.S.A. (CA - Central California)	T	\$142,700	\$698,408,806
312	Caribou DPS, Southern Mountain (<i>Rangifer tarandus ssp. caribou</i>) -U.S.A. (wherever found), Canada (southeastern British Columbia)	E	\$141,339	\$698,550,145
313	Thrush, small Kauai (=puaiohi) (<i>Myadestes palmeri</i>) -Wherever found	E	\$140,219	\$698,690,364
314	Dawn-flower, Texas prairie (<i>Hymenoxys texana</i>) -Wherever found	E	\$138,000	\$698,828,364
315	Penstemon, blowout (<i>Penstemon haydenii</i>) -Wherever found	E	\$137,139	\$698,965,503
316	Crayfish, Panama City (<i>Procambarus econfinae</i>) -Wherever found	T	\$137,127	\$699,102,630
317	Mouse, St. Andrew beach (<i>Peromyscus polionotus peninsularis</i>) -Wherever found	E	\$134,265	\$699,236,895
318	Goldfields, Contra Costa (<i>Lasthenia conjugens</i>) -Wherever found	E	\$133,547	\$699,370,442
319	Warbler, nightingale reed (old world warbler) (<i>Acrocephalus luscini</i>) -Wherever found	E	\$133,000	\$699,503,442
320	Pink, swamp (<i>Helonias bullata</i>) -	T	\$132,986	\$699,636,428
321	salmon Atlantic (<i>Salmo salar</i>) -Kennebec River DPS	E	\$131,500	\$699,767,928
322	Creeper, Hawaii (<i>Oreomystis mana</i>) -Wherever found	E	\$131,177	\$699,899,105
323	akepa, Hawaii (<i>Loxops coccineus</i>) -Wherever found	E	\$130,177	\$700,029,282
324	akiapolaau (<i>Hemignathus wilsoni</i>) -Wherever found	E	\$129,177	\$700,158,459
325	Elktoe, Appalachian (<i>Alasmidonta raveneliana</i>) -Wherever found	E	\$127,000	\$700,285,459
326	Chaffseed, American (<i>Schwalbea americana</i>) -Wherever found	E	\$126,500	\$700,411,959
327	Skink, sand (<i>Neoseps reynoldsi</i>) -Wherever found	T	\$125,000	\$700,536,959
328	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) -Wherever found; Except where listed as Experimental Populations	E	\$124,950	\$700,661,909
329	Tiger beetle, Northeastern beach (<i>Habroscelimorpha dorsalis dorsalis</i>) -Wherever found	T	\$124,328	\$700,786,237
330	Snail, Snake River physa (<i>Physa natricina</i>) -Wherever found	E	\$124,181	\$700,910,418
331	Bat, Mariana fruit (=Mariana flying fox) (<i>Pteropus mariannus mariannus</i>) -Wherever found	T	\$123,600	\$701,034,018
332	Pocket gopher, Yelm (<i>Thomomys mazama yelmensis</i>) -Wherever found	T	\$123,538	\$701,157,556
333	Butterfly, Smith's blue (<i>Euphilotes enoptes smithi</i>) -Wherever found	E	\$123,000	\$701,280,556
334	Butterfly, Lange's metalmark (<i>Apodemia mormo langei</i>) -Wherever found	E	\$122,800	\$701,403,356
335	Limpet, Banbury Springs (<i>Lanx</i> sp.) -Wherever found	E	\$122,326	\$701,525,682
336	Snail, Guam tree (<i>Partula radiolata</i>) -Wherever found	E	\$122,000	\$701,647,682
337	Kiponapona (<i>Phyllostegia racemosa</i>) -Wherever found	E	\$121,177	\$701,768,859
338	Seal, Ringed (<i>Phoca</i> (=Pusa) <i>hispida hispida</i>) -Wherever found	T	\$120,000	\$701,888,859
339	haha (<i>Cyanea shipmanii</i>) -Wherever found	E	\$119,980	\$702,008,839

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
340	No common name (Phyllostegia velutina) -Wherever found	E	\$119,677	\$702,128,516
341	Darter, Niangua (Etheostoma nianguae) -Wherever found	T	\$119,000	\$702,247,516
342	`Oha wai (Clermontia peleana) -Wherever found	E	\$118,980	\$702,366,496
343	Rockcress, Santa Cruz Island (Sibara filifolia) -Wherever found	E	\$118,700	\$702,485,196
344	Jaguarundi, Gulf Coast (Puma yagouaroundi cacomitli) -Wherever found	E	\$118,600	\$702,603,796
345	No common name (Dendrobium guamense) -Wherever found	T	\$118,500	\$702,722,296
346	oxytheca, Cushenbury (Oxytheca parishii var. goodmaniana) -Wherever found	E	\$118,100	\$702,840,396
347	Deer, key (Odocoileus virginianus clavium) -Wherever found	E	\$118,000	\$702,958,396
348	Pocket gopher, Tenino (Thomomys mazama tumuli) -Wherever found	T	\$117,868	\$703,076,264
349	Yellow-faced bee, anthricinan (Hylaeus anthracinus) -Wherever found	E	\$116,634	\$703,192,898
350	Towhee, Inyo California (Pipilo crissalis eremophilus) -Wherever found	T	\$115,100	\$703,307,998
351	Duck, Laysan (Anas laysanensis) -Wherever found	E	\$114,251	\$703,422,249
352	Makou (Ranunculus hawaiiensis) -Wherever found	E	\$114,177	\$703,536,426
353	Boa, Puerto Rican (Chilabothrus inornatus) -Wherever found	E	\$114,000	\$703,650,426
353	Prairie-clover, leafy (Dalea foliosa) -	E	\$114,000	\$703,764,426
355	Shiner, Cape Fear (Notropis mekistocholas) -Wherever found	E	\$113,500	\$703,877,926
356	Pocket gopher, Roy Prairie (Thomomys mazama glacialis) -Wherever found	T	\$113,008	\$703,990,934
357	`Oha wai (Clermontia lindseyana) -Wherever found	E	\$112,980	\$704,103,914
358	Sheep, Sierra Nevada bighorn (Ovis canadensis sierrae) -U.S.A. (CA) Sierra Nevada	E	\$112,580	\$704,216,494
359	Milk-vetch, Mancos (Astragalus humillimus) -Wherever found	E	\$112,347	\$704,328,841
360	`Oha wai (Clermontia pyralaria) -Wherever found	E	\$112,177	\$704,441,018
361	`Ahinahina (Argyroxiphium sandwicense ssp. macrocephalum) -Wherever found	T	\$111,910	\$704,552,928
362	Fadang (Cycas micronesica) -Wherever found	T	\$111,657	\$704,664,585
363	Beetle, Casey's June (Dinacoma caseyi) -Wherever found	E	\$110,300	\$704,774,885
364	Kangaroo rat, giant (Dipodomys ingens) -Wherever found	E	\$110,135	\$704,885,020
365	Hawk, Puerto Rican sharp-shinned (Accipiter striatus venator) -Wherever found	E	\$110,000	\$704,995,020
365	Salamander, Santa Cruz long-toed (Ambystoma macrodactylum croceum) -Wherever found	E	\$110,000	\$705,105,020
365	Sea turtle, loggerhead (Caretta caretta) -Mediterranean Sea DPS - Loggerhead sea turtles originating from the Mediterranean Sea east of 5 degrees 36 minues W. Long.	E	\$110,000	\$705,215,020
368	Mouse, southeastern beach (Peromyscus polionotus niveiventris) -wherever found	T	\$109,227	\$705,324,247
369	Bat, Pacific sheath-tailed (Emballonura semicaudata rotensis) -Wherever found	E	\$109,000	\$705,433,247
370	Tortoise, desert (Gopherus agassizii) -AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$108,333	\$705,541,580
371	Kidneyshell, fluted (Ptychobranhus subtentus) -Wherever found	E	\$108,085	\$705,649,665
372	popcornflower, rough (Plagiobothrys hirtus) -Wherever found	E	\$107,966	\$705,757,631
373	Wolf, red (Canis rufus) -U.S.A. (portions of NC and TN see 17.84(c)(9))	EXPN	\$107,000	\$705,864,631

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
374	Hellbender, Ozark (<i>Cryptobranchus alleganiensis bishopi</i>) -Wherever found	E	\$106,552	\$705,971,183
375	Dace, Clover Valley speckled (<i>Rhinichthys osculus oligoporus</i>) -Wherever found	E	\$104,700	\$706,075,883
375	Dudleya, Verity's (<i>Dudleya verityi</i>) -Wherever found	T	\$104,700	\$706,180,583
377	Snake, San Francisco garter (<i>Thamnophis sirtalis tetrataenia</i>) -Wherever found	E	\$104,600	\$706,285,183
378	Damselfly, orangeblack Hawaiian (<i>Megalagrion xanthomelas</i>) -Wherever found	E	\$104,502	\$706,389,685
379	Dace, Moapa (<i>Moapa coriacea</i>) -Wherever found	E	\$103,800	\$706,493,485
380	`Akoko, Ewa Plains (<i>Euphorbia skottsbergii</i> var. <i>skottsbergii</i>) -Wherever found	E	\$102,500	\$706,595,985
381	Spiraea, Virginia (<i>Spiraea virginiana</i>) -Wherever found	T	\$101,975	\$706,697,960
382	Mussel, oyster (<i>Epioblasma capsaeformis</i>) -Wherever found; Except where listed as Experimental Populations	E	\$101,950	\$706,799,910
383	Chub, spotfin (<i>Erimonax monachus</i>) -Wherever found, except where listed as an experimental population	T	\$100,335	\$706,900,245
384	Dwarf-flax, Marin (<i>Hesperolinon congestum</i>) -Wherever found	T	\$100,300	\$707,000,545
385	Mouse, Pacific pocket (<i>Perognathus longimembris pacificus</i>) -Wherever found	E	\$100,200	\$707,100,745
386	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) -	EXPN	\$98,900	\$707,199,645
387	Tiger beetle, Miami (<i>Cicindelia floridana</i>) -Wherever found	E	\$97,500	\$707,297,145
388	Butterfly, Mariana eight-spot (<i>Hypolimnas octocula marianensis</i>) -Wherever found	E	\$96,551	\$707,393,696
389	Snail, Langford's tree (<i>Partula langfordi</i>) -Wherever found	E	\$96,500	\$707,490,196
390	Ohai (<i>Sesbania tomentosa</i>) -Wherever found	E	\$96,303	\$707,586,499
391	Salamander, Jemez Mountains (<i>Plethodon neomexicanus</i>) -Wherever found	E	\$96,000	\$707,682,499
391	Snail, fragile tree (<i>Samoana fragilis</i>) -Wherever found	E	\$96,000	\$707,778,499
393	gartersnake, northern Mexican (<i>Thamnophis eques megalops</i>) -Wherever found	T	\$95,582	\$707,874,081
394	No common name (<i>Platydesma cornuta</i> var. <i>cornuta</i>) -Wherever found	E	\$95,000	\$707,969,081
394	Wild-buckwheat, gypsum (<i>Eriogonum gypsophilum</i>) -Wherever found	T	\$95,000	\$708,064,081
396	Daisy, Willamette (<i>Erigeron decumbens</i>) -Wherever found	E	\$93,500	\$708,157,581
397	Fritillary, Gentner's (<i>Fritillaria gentneri</i>) -Wherever found	E	\$93,481	\$708,251,062
398	Paintbrush, soft-leaved (<i>Castilleja mollis</i>) -Wherever found	E	\$93,300	\$708,344,362
399	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. (specific portions of Arizona, Nevada, and Utah)-see 17.84(j)	EXPN	\$92,800	\$708,437,162
400	Mouse, New Mexico meadow jumping (<i>Zapus hudsonius luteus</i>) -Wherever found	E	\$91,900	\$708,529,062
401	Moorhen, Mariana common (<i>Gallinula chloropus guami</i>) -Wherever found	E	\$91,000	\$708,620,062
402	Iagu, Hayun (=Guam), Tronkon guafi (Rota)) (<i>Serianthes nelsonii</i>) -Wherever found	E	\$87,729	\$708,707,791
403	Tui chub, Mohave (<i>Gila bicolor</i> ssp. <i>mohavensis</i>) -Wherever found	E	\$87,585	\$708,795,376
404	Popolo (<i>Solanum nelsonii</i>) -Wherever found	E	\$87,500	\$708,882,876
405	Milk-vetch, triple-ribbed (<i>Astragalus tricarinatus</i>) -Wherever found	E	\$87,300	\$708,970,176
406	skink, blue-tailed mole (<i>Eumeces egregius lividus</i>) -Wherever found	T	\$87,000	\$709,057,176
406	potato-bean, Price's (<i>Apios priceana</i>) -	T	\$87,000	\$709,144,176

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
408	Mouse, Choctawhatchee beach (<i>Peromyscus polionotus allopshrys</i>) -Wherever found	E	\$86,265	\$709,230,441
409	picture-wing fly, Hawaiian (<i>Drosophila hemipeza</i>) -Wherever found	E	\$86,140	\$709,316,581
410	Toad, Houston (<i>Bufo houstonensis</i>) -Wherever found	E	\$86,000	\$709,402,581
411	Butterfly, callippe silverspot (<i>Speyeria callippe callippe</i>) -Wherever found	E	\$85,800	\$709,488,381
412	Shrew, Buena Vista Lake ornate (<i>Sorex ornatus relictus</i>) -Wherever found	E	\$85,600	\$709,573,981
413	Mouse, Anastasia Island beach (<i>Peromyscus polionotus phasma</i>) -Wherever found	E	\$85,397	\$709,659,378
414	Mussel, scaleshell (<i>Leptodea leptodon</i>) -Wherever found	E	\$84,635	\$709,744,013
415	Megapode, Micronesian (<i>Megapodius laperouse</i>) -Wherever found	E	\$84,500	\$709,828,513
416	Crocodile, American (<i>Crocodylus acutus</i>) -Wherever found, except in U.S.A. (FL)	E	\$84,464	\$709,912,977
417	Stickleback, unarmored threespine (<i>Gasterosteus aculeatus williamsoni</i>) -Wherever found	E	\$83,900	\$709,996,877
418	Milkweed, Mead's (<i>Asclepias meadii</i>) -Wherever found	T	\$83,214	\$710,080,091
419	Cactus, Kuenzler hedgehog (<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>) -Wherever found	T	\$83,000	\$710,163,091
420	Thistle, Pitcher's (<i>Cirsium pitcheri</i>) -Wherever found	T	\$82,038	\$710,245,129
421	Butterfly, Schaus swallowtail (<i>Heracles aristodemus ponceanus</i>) -Wherever found	E	\$82,000	\$710,327,129
422	Butterfly, Behren's silverspot (<i>Speyeria zerene behrensii</i>) -Wherever found	E	\$81,700	\$710,408,829
422	Mountain beaver, Point Arena (<i>Aplodontia rufa nigra</i>) -Wherever found	E	\$81,700	\$710,490,529
424	milkvetch, Osterhout (<i>Astragalus osterhoutii</i>) -Wherever found	E	\$81,489	\$710,572,018
425	Phacelia, DeBeque (<i>Phacelia submutica</i>) -Wherever found	T	\$81,175	\$710,653,193
426	Spinymussel, Tar River (<i>Parvaspina steinstansana</i>) -Wherever found	E	\$81,000	\$710,734,193
427	Darter, Kentucky arrow (<i>Etheostoma spilotum</i>) -Wherever found	T	\$80,500	\$710,814,693
428	Cheetah (<i>Acinonyx jubatus</i>) -Wherever found	E	\$79,790	\$710,894,483
429	Tiger beetle, Salt Creek (<i>Cicindela nevadica lincolniiana</i>) -Wherever found	E	\$79,500	\$710,973,983
430	Evening-primrose, Antioch Dunes (<i>Oenothera deltoides</i> ssp. <i>howellii</i>) -Wherever found	E	\$79,000	\$711,052,983
431	Heartleaf, dwarf-flowered (<i>Hexastylis naniflora</i>) -Wherever found	T	\$78,500	\$711,131,483
432	bladderpod, Missouri (<i>Physaria filiformis</i>) -Wherever found	T	\$78,100	\$711,209,583
433	Sunflower, Schweinitz's (<i>Helianthus schweinitzii</i>) -Wherever found	E	\$77,500	\$711,287,083
434	Clubshell (<i>Pleurobema clava</i>) -Wherever found; Except where listed as Experimental Populations	E	\$77,187	\$711,364,270
435	Darter, duskytail (<i>Etheostoma percnurum</i>) -Wherever found	E	\$77,012	\$711,441,282
436	Dace, blackside (<i>Phoxinus cumberlandensis</i>) -Wherever found	T	\$77,000	\$711,518,282
437	Flannelbush, Pine Hill (<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>) -Wherever found	E	\$76,500	\$711,594,782
438	Pearlymussel, slabside (<i>Pleuonaia dolabelloides</i>) -Wherever found	E	\$76,135	\$711,670,917
439	Bat, Ozark big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii ingens</i>) -Wherever found	E	\$75,850	\$711,746,767
440	picture-wing fly, Hawaiian (<i>Drosophila montgomeryi</i>) -Wherever found	E	\$75,653	\$711,822,420
441	Millerbird, Nihoa (old world warbler) (<i>Acrocephalus familiaris kingi</i>) -Wherever found	E	\$75,000	\$711,897,420

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
441	Tiger beetle, Puritan (<i>Ellipsoptera puritana</i>) -Wherever found	T	\$75,000	\$711,972,420
443	Pimpleback, orangefoot (pearlymussel) (<i>Plethobasus cooperianus</i>) -Wherever found	E	\$73,635	\$712,046,055
444	Skullcap, large-flowered (<i>Scutellaria montana</i>) -Wherever found	T	\$71,500	\$712,117,555
444	Woodrat, Key Largo (<i>Neotoma floridana smalli</i>) -Wherever found	E	\$71,500	\$712,189,055
446	Lessingia, San Francisco (<i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i>)) -Wherever found	E	\$70,800	\$712,259,855
447	No common name (<i>Osmoxylon mariannense</i>) -Wherever found	E	\$70,729	\$712,330,584
448	No common name (<i>Nesogenes rotensis</i>) -Wherever found	E	\$69,229	\$712,399,813
449	Riffleshell, tan (<i>Epioblasma florentina walkeri</i> (=E. <i>walkeri</i>)) -Wherever found	E	\$68,450	\$712,468,263
450	Mouse, Perdido Key beach (<i>Peromyscus polionotus trissyllepsis</i>) -Wherever found	E	\$68,390	\$712,536,653
451	Buckwheat, steamboat (<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>) -Wherever found	E	\$68,133	\$712,604,786
452	Lousewort, Furbish (<i>Pedicularis furbishiae</i>) -Wherever found	E	\$68,000	\$712,672,786
452	Snail, Morro shoulderband (=Banded dune) (<i>Helminthoglypta walkeriana</i>) -Wherever found	T	\$68,000	\$712,740,786
454	Poolfish, Pahump (<i>Empetrichthys latos</i>) -Wherever found	E	\$67,754	\$712,808,540
455	Jewelflower, California (<i>Caulanthus californicus</i>) -Wherever found	E	\$67,583	\$712,876,123
456	Frog, Chiricahua leopard (<i>Rana chiricahuensis</i>) -Wherever found	T	\$67,565	\$712,943,688
457	Milk-vetch, Applegate's (<i>Astragalus applegatei</i>) -Wherever found	E	\$67,200	\$713,010,888
458	Snail, Bliss Rapids (<i>Taylorconcha serpenticola</i>) -Wherever found	T	\$66,187	\$713,077,075
459	Skullcap, Florida (<i>Scutellaria floridana</i>) -	T	\$65,830	\$713,142,905
460	Finch, Nihoa (honeycreeper) (<i>Telespyza ultima</i>) -Wherever found	E	\$65,500	\$713,208,405
461	Leopard, snow (<i>Uncia</i> (=Panthera) <i>uncia</i>) -Wherever found	E	\$65,180	\$713,273,585
462	Lily, Western (<i>Lilium occidentale</i>) -Wherever found	E	\$64,600	\$713,338,185
463	Hellbender, Eastern (<i>Cryptobranchus alleganiensis alleganiensis</i>) -Missouri DPS	E	\$64,500	\$713,402,685
464	Damselfly, blackline Hawaiian (<i>Megalagrion nigrohamatum nigrolineatum</i>) -Wherever found	E	\$63,704	\$713,466,389
465	Pigtoe, Atlantic (<i>Fusconaia masoni</i>) -Wherever found	T	\$63,200	\$713,529,589
466	Salamander, Shenandoah (<i>Plethodon shenandoah</i>) -Wherever found	E	\$63,000	\$713,592,589
467	Springsnail, Bruneau Hot (<i>Pyrgulopsis bruneauensis</i>) -Wherever found	E	\$62,887	\$713,655,476
468	Toad, arroyo (=arroyo southwestern) (<i>Anaxyrus californicus</i>) -Wherever found	E	\$62,600	\$713,718,076
469	elepaio, Oahu (<i>Chasiempis ibidis</i>) -Wherever found	E	\$62,000	\$713,780,076
470	Pigtoe, oval (<i>Pleurobema pyriforme</i>) -Wherever found	E	\$61,993	\$713,842,069
471	Fanshell (<i>Cyprogenia stegaria</i>) -Wherever found	E	\$61,787	\$713,903,856
472	No common name (<i>Tuberolabium guamense</i>) -Wherever found	T	\$61,700	\$713,965,556
473	meadowfoam, large-flowered woolly (<i>Limnanthes pumila</i> ssp. <i>grandiflora</i>) -Wherever found	E	\$61,500	\$714,027,056
474	`Oha wai (<i>Clermontia samuelii</i>) -Wherever found	E	\$60,708	\$714,087,764

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
475	Sea turtle, loggerhead (<i>Caretta caretta</i>) -Loggerhead sea turtles originating from the South Pacific south of the equator, north of 60 degrees S. Lat., west of 67 degrees W. Long., and east of 141 degrees E. Long.	E	\$60,480	\$714,148,244
476	Darter, Etowah (<i>Etheostoma etowahae</i>) -Wherever found	E	\$60,000	\$714,208,244
476	Milk-vetch, Lane Mountain (<i>Astragalus jaegerianus</i>) -Wherever found	E	\$60,000	\$714,268,244
478	Bear-poppy, Dwarf (<i>Arctomecon humilis</i>) -Wherever found	E	\$59,648	\$714,327,892
479	Ihi (<i>Portulaca villosa</i>) -Wherever found	E	\$59,500	\$714,387,392
479	Topminnow, barrens (<i>Fundulus julisia</i>) -Wherever found	E	\$59,500	\$714,446,892
481	No common name (<i>Geocarpon minimum</i>) -Wherever found	T	\$59,000	\$714,505,892
482	Mucket, Neosho (<i>Lampsilis rafinesqueana</i>) -Wherever found	E	\$58,800	\$714,564,692
483	halumtano, Cebello (<i>Bulbophyllum guamense</i>) -Wherever found	T	\$58,551	\$714,623,243
484	Madtom, chucky (<i>Noturus crypticus</i>) -Wherever found	E	\$58,500	\$714,681,743
484	Pupfish, Owens (<i>Cyprinodon radiosus</i>) -Wherever found	E	\$58,500	\$714,740,243
486	Finch, Laysan (honeycreeper) (<i>Telespyza cantans</i>) -Wherever found	E	\$58,000	\$714,798,243
486	Spurge, telephus (<i>Euphorbia telephioides</i>) -	T	\$58,000	\$714,856,243
488	Phacelia, North Park (<i>Phacelia formosula</i>) -Wherever found	E	\$57,547	\$714,913,790
489	Milkweed, Welsh's (<i>Asclepias welshii</i>) -Wherever found	T	\$57,200	\$714,970,990
490	Chub, Pahrnagat roundtail (<i>Gila robusta jordani</i>) -Wherever found	E	\$57,168	\$715,028,158
491	Tarplant, Gaviota (<i>Deinandra increscens</i> ssp. <i>villosa</i>) -Wherever found	E	\$57,100	\$715,085,258
492	Pondberry (<i>Lindera melissifolia</i>) -Wherever found	E	\$57,000	\$715,142,258
492	skink, Slevin's (<i>Emoia slevini</i>) -Wherever found	E	\$57,000	\$715,199,258
494	No common name (<i>Phyllostegia bracteata</i>) -Wherever found	E	\$56,966	\$715,256,224
494	No common name (<i>Phyllostegia haliakalae</i>) -Wherever found	E	\$56,966	\$715,313,190
496	Damselfly, Pacific Hawaiian (<i>Megalagrion pacificum</i>) -Wherever found	E	\$56,825	\$715,370,015
497	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata</i>) -Wherever found; Except where listed as Experimental Populations	E	\$56,652	\$715,426,667
498	Silversword, Mauna Loa (=Ka'u) (<i>Argyroxiphium kauense</i>) -Wherever found	E	\$56,500	\$715,483,167
499	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Eastern Indian and Western Pacific Oceans, bounded by the following lines and coordinates: 41 degrees N. Lat. in the north, 41 degrees N., 146 degrees E. in the northeast; 4.5 degrees N., 129 degrees E. in the southeast; along the southern coast of the island of New Guinea; along the western coast of Australia (west of 142 degrees E. Long.); 40 degrees S. Lat. in the south; and 84 degrees E. Long. in the east	T	\$56,287	\$715,539,454
500	Cycladenia, Jones (<i>Cycladenia humilis</i> var. <i>jonesii</i>) -Wherever found	T	\$56,228	\$715,595,682
501	Chaff-flower, round-leaved (<i>Achyranthes splendens</i> var. <i>rotundata</i>) -Wherever found	E	\$56,000	\$715,651,682
501	No common name (<i>Pritchardia remota</i>) -Wherever found	E	\$56,000	\$715,707,682
503	Layia, beach (<i>Layia carnosa</i>) -Wherever found	T	\$55,500	\$715,763,182

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
503	Moth, Blackburn's sphinx (<i>Manduca blackburni</i>) -Wherever found	E	\$55,500	\$715,818,682
505	Condor, California (<i>Gymnogyps californianus</i>) -U.S.A. (Oregon, and specific portions of northern California and northwest Nevada)	EXPN	\$55,261	\$715,873,943
506	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) -Wherever found; Except where listed as Experimental Populations	E	\$55,135	\$715,929,078
507	Threeridge, fat (mussel) (<i>Amblema neislerii</i>) -Wherever found	E	\$54,993	\$715,984,071
508	Lomatium, Cook's (<i>Lomatium cookii</i>) -Wherever found	E	\$54,707	\$716,038,778
509	Wild-buckwheat, southern mountain (<i>Eriogonum kennedyi</i> var. <i>austromontanum</i>) -Wherever found	T	\$54,500	\$716,093,278
510	Butterfly, Mount Charleston blue (<i>Icaricia (Plebejus) shasta charlestonensis</i>) -Wherever found	E	\$54,300	\$716,147,578
511	Watercress, Gambel's (<i>Rorippa gambellii</i>) -Wherever found	E	\$54,200	\$716,201,778
512	wild buckwheat, clay-loving (<i>Eriogonum pelinophilum</i>) -Wherever found	E	\$54,082	\$716,255,860
513	Monkeyface, Cumberland (pearlymussel) (<i>Theliderma intermedia</i>) -Wherever found; Except where listed as Experimental Populations	E	\$53,635	\$716,309,495
513	Pigtoe, rough (<i>Pleurobema plenum</i>) -Wherever found	E	\$53,635	\$716,363,130
515	Ufa-halomtano (<i>Heritiera longipetiolata</i>) -Wherever found	E	\$53,607	\$716,416,737
516	Snail, painted snake coiled forest (<i>Anguispira picta</i>) -Wherever found	T	\$53,500	\$716,470,237
516	Trillium, relict (<i>Trillium reliquum</i>) -Wherever found	E	\$53,500	\$716,523,737
518	Cactus, Wright fishhook (<i>Sclerocactus wrightiae</i>) -Wherever found	E	\$53,197	\$716,576,934
519	Crayfish, Guyandotte River (<i>Cambarus veteranus</i>) -Wherever found	E	\$53,187	\$716,630,121
520	Bladderpod, San Bernardino Mountains (<i>Lesquerella kingii</i> ssp. <i>bernardina</i>) -Wherever found	E	\$53,100	\$716,683,221
521	No common name (<i>Varronia rupicola</i>) -Wherever found	T	\$52,500	\$716,735,721
522	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) -Wherever found; Except where listed as Experimental Populations	E	\$52,085	\$716,787,806
523	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) -Wherever found, except where listed as an experimental population	E	\$51,450	\$716,839,256
524	Stickseed, showy (<i>Hackelia venusta</i>) -Wherever found	E	\$51,042	\$716,890,298
525	Clubshell, southern (<i>Pleurobema decisum</i>) -Wherever found	E	\$51,000	\$716,941,298
526	haha (<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>) -Wherever found	E	\$49,966	\$716,991,264
527	Milk-vetch, Braunton's (<i>Astragalus brauntonii</i>) -Wherever found	E	\$49,500	\$717,040,764
527	Sea turtle, loggerhead (<i>Caretta caretta</i>) -North Indian Ocean DPS - Loggerhead sea turtles originating from the North Indian Ocean north of the equator and south of 30° N. Lat	E	\$49,500	\$717,090,264
529	Squirrel, Carolina northern flying (<i>Glaucomys sabrinus coloratus</i>) -Wherever found	E	\$49,255	\$717,139,519
530	No common name (<i>Phyllanthus saffordii</i>) -Wherever found	E	\$49,107	\$717,188,626

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
531	Wolf, maned (<i>Chrysocyon brachyurus</i>) -Wherever found	E	\$49,100	\$717,237,726
532	Pearlymussel, dromedary (<i>Dromus dromas</i>) -Wherever found; Except where listed as Experimental Populations	E	\$49,085	\$717,286,811
533	Lampmussel, Alabama (<i>Lampsilis virescens</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$49,000	\$717,335,811
534	Milk-vetch, Fish Slough (<i>Astragalus lentiginos</i> var. <i>piscinensis</i>) -Wherever found	T	\$48,900	\$717,384,711
535	Paudedo (<i>Hedyotis megalantha</i>) -Wherever found	E	\$48,607	\$717,433,318
536	Harperella (<i>Ptilimnium nodosum</i>) -Wherever found	E	\$48,505	\$717,481,823
537	heelsplitter, Inflated (<i>Potamilus inflatus</i>) -Wherever found	T	\$48,500	\$717,530,323
537	Sculpin, Grotto (<i>Cottus specus</i>) -Wherever found	E	\$48,500	\$717,578,823
539	Bird's-beak, soft (<i>Cordylanthus mollis</i> ssp. <i>mollis</i>) -Wherever found	E	\$48,300	\$717,627,123
540	Niterwort, Amargosa (<i>Nitrophila mohavensis</i>) -Wherever found	E	\$48,200	\$717,675,323
541	Kangaroo rat, Fresno (<i>Dipodomys nitratoides exilis</i>) -Wherever found	E	\$48,131	\$717,723,454
542	Pearlshell, Alabama (<i>Margaritifera marrianae</i>) -Wherever found	E	\$47,000	\$717,770,454
543	Joint-vetch, Sensitive (<i>Aeschynomene virginica</i>) -Wherever found	T	\$46,896	\$717,817,350
544	Opuhe (<i>Urera kaalae</i>) -Wherever found	E	\$46,500	\$717,863,850
544	Shiner, blue (<i>Cyprinella caerulea</i>) -Wherever found	T	\$46,500	\$717,910,350
546	Pocketbook, fat (<i>Potamilus capax</i>) -Wherever found	E	\$46,400	\$717,956,750
547	No common name (<i>Schiedea diffusa</i> subsp. <i>diffusa</i>) -Wherever found	E	\$46,224	\$718,002,974
547	No common name (<i>Schiedea haleakalensis</i>) -Wherever found	E	\$46,224	\$718,049,198
549	Cactus, Winkler (<i>Pediocactus winkleri</i>) -Wherever found	T	\$46,220	\$718,095,418
550	Sturgeon, Alabama (<i>Scaphirhynchus suttkusi</i>) -Wherever found	E	\$46,000	\$718,141,418
551	Bush-clover, prairie (<i>Lespedeza leptostachya</i>) -Wherever found	T	\$45,876	\$718,187,294
552	Milk-vetch, Ventura Marsh (<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i>) -Wherever found	E	\$45,800	\$718,233,094
553	Thelypody, Howell's spectacular (<i>Thelypodium howellii</i> ssp. <i>spectabilis</i>) -Wherever found	T	\$45,500	\$718,278,594
553	Turtle, Plymouth Redbelly (<i>Pseudemys rubriventris bangsi</i>) -Wherever found	E	\$45,500	\$718,324,094
553	Woodpecker, ivory-billed (<i>Campephilus principalis</i>) -Wherever found	E	\$45,500	\$718,369,594
556	roseroot, Leedy's (<i>Rhodiola integrifolia</i> ssp. <i>leedyi</i>) -Wherever found	T	\$45,438	\$718,415,032
557	Darter, slackwater (<i>Etheostoma boschungii</i>) -Wherever found	T	\$45,360	\$718,460,392
558	Milk-vetch, Shivwits (<i>Astragalus ampullarioides</i>) -Wherever found	E	\$45,350	\$718,505,742
559	Pigtoe, shiny (<i>Fusconaia cor</i>) -Wherever found; Except where listed as Experimental Populations	E	\$45,135	\$718,550,877
560	Daisy, Parish's (<i>Erigeron parishii</i>) -Wherever found	T	\$45,100	\$718,595,977

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
561	Cactus, San Rafael (<i>Pediocactus despainii</i>) -Wherever found	E	\$45,063	\$718,641,040
562	No common name (<i>Lobelia oahuensis</i>) -Wherever found	E	\$45,000	\$718,686,040
562	Rhododendron, Chapman (<i>Rhododendron chapmanii</i>) -	E	\$45,000	\$718,731,040
564	Mouse, Key Largo cotton (<i>Peromyscus gossypinus allapaticola</i>) -Wherever found	E	\$44,500	\$718,775,540
565	Ebonyshell, round (<i>Reginaia rotulata</i>) -Wherever found	E	\$44,493	\$718,820,033
566	[no common name] Snail (<i>Eua zebrina</i>) -Wherever found	E	\$44,430	\$718,864,463
566	[no common name] Snail (<i>Ostodes strigatus</i>) -Wherever found	E	\$44,430	\$718,908,893
568	Ko'oko'olau (<i>Bidens micrantha</i> ssp. <i>kalealaha</i>) -Wherever found	E	\$44,224	\$718,953,117
569	Tarplant, Santa Cruz (<i>Holocarpha macradenia</i>) -Wherever found	T	\$43,900	\$718,997,017
570	Alani (<i>Melicope degeneri</i>) -Wherever found	E	\$43,500	\$719,040,517
570	Alani (<i>Melicope paniculata</i>) -Wherever found	E	\$43,500	\$719,084,017
572	Kuahiwi laukahi (<i>Plantago princeps</i>) -Wherever found	E	\$43,224	\$719,127,241
573	Crayfish, Nashville (<i>Orconectes shoupi</i>) -Wherever found	E	\$43,000	\$719,170,241
573	Sumac, Michaux's (<i>Rhus michauxii</i>) -Wherever found	E	\$43,000	\$719,213,241
573	Seablite, California (<i>Suaeda californica</i>) -	E	\$43,000	\$719,256,241
576	Monkshood, northern wild (<i>Aconitum noveboracense</i>) -Wherever found	T	\$42,538	\$719,298,779
577	Haiwale (<i>Cyrtandra nanawaleensis</i>) -Wherever found	E	\$42,000	\$719,340,779
577	picture-wing fly, Hawaiian (<i>Drosophila digressa</i>) -Wherever found	E	\$42,000	\$719,382,779
577	Thistle, La Graciosa (<i>Cirsium loncholepis</i>) -Wherever found	E	\$42,000	\$719,424,779
580	Madtom, Carolina (<i>Noturus furiosus</i>) -Wherever found	E	\$41,700	\$719,466,479
581	Falcon, Eurasian peregrine (<i>Falco peregrinus peregrinus</i>) -Wherever found	E	\$41,559	\$719,508,038
582	Beetle, Hungerford's crawling water (<i>Brychius hungerfordi</i>) -Wherever found	E	\$41,500	\$719,549,538
583	No common name (<i>Phyllostegia floribunda</i>) -Wherever found	E	\$41,430	\$719,590,968
584	Lichen, rock gnome (<i>Gymnoderma lineare</i>) -Wherever found	E	\$41,400	\$719,632,368
585	Milk-vetch, Holmgren (<i>Astragalus holmgreniorum</i>) -Wherever found	E	\$41,397	\$719,673,765
586	rock cress, Shale barren (<i>Boechera serotina</i>) -Wherever found	E	\$41,301	\$719,715,066
587	Pea, Big Pine partridge (<i>Chamaecrista lineata keyensis</i>) -	E	\$41,000	\$719,756,066
588	Riffleshell, northern (<i>Epioblasma rangiana</i>) -Wherever found	E	\$40,952	\$719,797,018
589	Rhinoceros, northern white (<i>Ceratotherium simum cottoni</i>) -Zaire, Uganda, Sudan, Central African Republic	E	\$40,295	\$719,837,313
590	Haiwale (<i>Cyrtandra ferripilosa</i>) -Wherever found	E	\$40,224	\$719,877,537
590	Nohoanu (<i>Geranium arboreum</i>) -Wherever found	E	\$40,224	\$719,917,761
592	Milk-vetch, Peirson's (<i>Astragalus magdalenae</i> var. <i>peirsonii</i>) -Wherever found	T	\$40,000	\$719,957,761
592	No common name (<i>Schiedea verticillata</i>) -Wherever found	E	\$40,000	\$719,997,761
594	Madtom, yellowfin (<i>Noturus flavipinnis</i>) -Wherever found, except where listed as an experimental population	T	\$39,712	\$720,037,473

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
595	Madtom, yellowfin (<i>Noturus flavipinnis</i>) -U.S.A. (TN-specified portions of the Tellico River; see 17.84(e)(1)(ii))	EXPN	\$39,500	\$720,076,973
596	Darter, diamond (<i>Crystallaria cincotta</i>) -Wherever found	E	\$39,351	\$720,116,324
597	Mapleleaf, winged (<i>Quadrula fragosa</i>) -Wherever found, except where listed as an experimental population	E	\$39,200	\$720,155,524
598	Ring pink (mussel) (<i>Obovaria retusa</i>) -Wherever found	E	\$39,135	\$720,194,659
599	'Ena`ena (<i>Pseudognaphalium sandwicense</i> var. <i>molokaiense</i>) -Wherever found	E	\$39,000	\$720,233,659
599	Ground-plum, Guthrie's (=Pyne's) (<i>Astragalus bibullatus</i>) -Wherever found	E	\$39,000	\$720,272,659
599	Sucker, June (<i>Chasmistes liorus</i>) -Wherever found	T	\$39,000	\$720,311,659
602	Ko`oko`olau (<i>Bidens campylothea</i> ssp. <i>pentamera</i>) -Wherever found	E	\$38,724	\$720,350,383
602	Nohoanu (<i>Geranium hanaense</i>) -Wherever found	E	\$38,724	\$720,389,107
604	haha (<i>Cyanea asplenifolia</i>) -Wherever found	E	\$38,224	\$720,427,331
604	Ko`oko`olau (<i>Bidens campylothea</i> ssp. <i>waihoiensis</i>) -Wherever found	E	\$38,224	\$720,465,555
604	No common name (<i>Sanicula sandwicensis</i>) -Wherever found	E	\$38,224	\$720,503,779
607	cactus, Uinta Basin hookless (<i>Sclerocactus wetlandicus</i>) -Wherever found	T	\$38,191	\$720,541,970
608	No common name (<i>Asplenium peruvianum</i> var. <i>insulare</i>) -Wherever found	E	\$38,171	\$720,580,141
609	Phlox, Texas trailing (<i>Phlox nivalis</i> ssp. <i>texensis</i>) -Wherever found	E	\$38,100	\$720,618,241
610	Pigeon, Puerto Rican plain (<i>Columba inornata wetmorei</i>) -Wherever found	E	\$38,000	\$720,656,241
611	Elktoe, Cumberland (<i>Alasmodonta atropurpurea</i>) -Wherever found	E	\$37,950	\$720,694,191
612	Snake, copperbelly water (<i>Nerodia erythrogaster neglecta</i>) -Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$37,776	\$720,731,967
613	Lilliput, pale (pearlymussel) (<i>Toxolasma cylindrellus</i>) -Wherever found	E	\$37,635	\$720,769,602
614	Amphipod, Hay's Spring (<i>Stygobromus hayi</i>) -Wherever found	E	\$37,500	\$720,807,102
614	Rabbit, Lower Keys marsh (<i>Sylvilagus palustris hefneri</i>) -Wherever found	E	\$37,500	\$720,844,602
614	Thistle, Suisun (<i>Cirsium hydrophilum</i> var. <i>hydrophilum</i>) -Wherever found	E	\$37,500	\$720,882,102
617	Moccasinshell, Suwannee (<i>Medionidus walkeri</i>) -Wherever found	T	\$37,493	\$720,919,595
618	Snail, Lanai tree (<i>Partulina semicarinata</i>) -Wherever found	E	\$37,130	\$720,956,725
619	Birch, Virginia round-leaf (<i>Betula uber</i>) -Wherever found	T	\$37,000	\$720,993,725
619	Kidneyshell, triangular (<i>Ptychobranhus greenii</i>) -Wherever found	E	\$37,000	\$721,030,725
619	Rice rat, silver (<i>Oryzomys palustris natator</i>) -Lower FL Keys (west of Seven Mile Bridge)	E	\$37,000	\$721,067,725
622	Iris, dwarf lake (<i>Iris lacustris</i>) -Wherever found	T	\$36,600	\$721,104,325
623	Sneezeweed, Virginia (<i>Helenium virginicum</i>) -Wherever found	T	\$36,576	\$721,140,901
624	Bat, Mexican long-nosed (<i>Leptonycteris nivalis</i>) -Wherever found	E	\$36,541	\$721,177,442
625	Spineflower, Monterey (<i>Chorizanthe pungens</i> var. <i>pungens</i>) -Wherever found	T	\$36,400	\$721,213,842
626	Spinedace, White River (<i>Lepidomeda albivallis</i>) -Wherever found	E	\$36,300	\$721,250,142
627	Sandwort, Marsh (<i>Arenaria paludicola</i>) -Wherever found	E	\$36,100	\$721,286,242

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
628	halomtano, Berenghenas (<i>Solanum guamense</i>) -Wherever found	E	\$36,051	\$721,322,293
629	No common name (<i>Nervilia jacksoniae</i>) -Wherever found	T	\$36,000	\$721,358,293
629	Pocketbook, speckled (<i>Lampsilis streckeri</i>) -Wherever found	E	\$36,000	\$721,394,293
631	haha (<i>Cyanea marksii</i>) -Wherever found	E	\$35,930	\$721,430,223
632	Skipper, Carson wandering (<i>Pseudocopaeodes eunus obscurus</i>) -Wherever found	E	\$35,907	\$721,466,130
633	Coneflower, smooth (<i>Echinacea laevigata</i>) -Wherever found	T	\$35,900	\$721,502,030
634	Lizard, St. Croix ground (<i>Ameiva polops</i>) -Wherever found	E	\$35,500	\$721,537,530
634	Pigtoe, heavy (<i>Pleurobema taitianum</i>) -Wherever found	E	\$35,500	\$721,573,030
634	leather flower, Morefield's (<i>Clematis morefieldii</i>) -	E	\$35,500	\$721,608,530
637	Snail, Lanai tree (<i>Partulina variabilis</i>) -Wherever found	E	\$35,130	\$721,643,660
638	cactus, Pariette (<i>Sclerocactus brevispinus</i>) -Wherever found	T	\$35,122	\$721,678,782
639	Oryx, scimitar-horned (<i>Oryx dammah</i>) -Wherever found	E	\$35,100	\$721,713,882
639	Thornmint, San Diego (<i>Acanthomintha ilicifolia</i>) -Wherever found	T	\$35,100	\$721,748,982
641	Turtle, ringed map (<i>Graptemys oculifera</i>) -Wherever found	T	\$35,000	\$721,783,982
642	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata semicaudata</i>) -American Samoa	E	\$34,930	\$721,818,912
643	`aku (<i>Cyanea tritomantha</i>) -Wherever found	E	\$34,803	\$721,853,715
643	No common name (<i>Schiedea diffusa</i> ssp. <i>macraei</i>) -Wherever found	E	\$34,803	\$721,888,518
645	Horse, Przewalski's (<i>Equus przewalskii</i>) -Wherever found	E	\$34,500	\$721,923,018
645	Pigtoe, southern (<i>Pleurobema georgianum</i>) -Wherever found	E	\$34,500	\$721,957,518
647	ivesia, Webber's (<i>Ivesia webberi</i>) -Wherever found	T	\$34,103	\$721,991,621
648	Butterfly, bay checkerspot (<i>Euphydryas editha bayensis</i>) -Wherever found	T	\$34,000	\$722,025,621
648	Glade cress, Kentucky (<i>Leavenworthia exigua laciniata</i>) -Wherever found	T	\$34,000	\$722,059,621
650	Bladderpod, Short's (<i>Physaria globosa</i>) -Wherever found	E	\$33,800	\$722,093,421
651	Bully, Everglades (<i>Sideroxylon reclinatum</i> ssp. <i>austrofloridense</i>) -	T	\$33,562	\$722,126,983
652	Rock-cress, Braun's (<i>Arabis perstellata</i>) -Wherever found	E	\$33,500	\$722,160,483
652	Spider, spruce-fir moss (<i>Microhexura montivaga</i>) -Wherever found	E	\$33,500	\$722,193,983
654	Moccasinshell, Coosa (<i>Medionidus parvulus</i>) -Wherever found	E	\$33,000	\$722,226,983
654	No common name (<i>Platydesma remyi</i>) -Wherever found	E	\$33,000	\$722,259,983
654	phlox, Yreka (<i>Phlox hirsuta</i>) -Wherever found	E	\$33,000	\$722,292,983
657	haha (<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>) -Wherever found	E	\$32,913	\$722,325,896
658	Liveforever, Santa Barbara Island (<i>Dudleya traskiae</i>) -Wherever found	E	\$32,500	\$722,358,396
659	`Ohe (<i>Joinvillea ascendens ascendens</i>) -Wherever found	E	\$32,483	\$722,390,879
660	Woundfin (<i>Plagopterus argentissimus</i>) -Wherever found, except where listed as an experimental population	E	\$32,073	\$722,422,952
661	Dropwort, Canby's (<i>Oxypolis canbyi</i>) -Wherever found	E	\$32,000	\$722,454,952
661	Salamander, Red Hills (<i>Phaeognathus hubrichti</i>) -Wherever found	T	\$32,000	\$722,486,952
663	Waterdog, Neuse River (<i>Necturus lewisi</i>) -Wherever found	T	\$31,700	\$722,518,652

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
664	No common name (<i>Bidens hillebrandiana</i> ssp. <i>hillebrandiana</i>) -Wherever found	E	\$31,500	\$722,550,152
664	No common name (<i>Stenogyne cranwelliae</i>) -Wherever found	E	\$31,500	\$722,581,652
664	Sedge, golden (<i>Carex lutea</i>) -Wherever found	E	\$31,500	\$722,613,152
667	Moccasinshell, Alabama (<i>Medionidus acutissimus</i>) -Wherever found	T	\$31,493	\$722,644,645
668	Spineflower, slender-horned (<i>Dodecahema leptoceras</i>) -Wherever found	E	\$31,200	\$722,675,845
669	Haiwale (<i>Cyrtandra wagneri</i>) -Wherever found	E	\$31,000	\$722,706,845
669	Snail, Newcomb's (<i>Erinna newcombi</i>) -Wherever found	T	\$31,000	\$722,737,845
671	Minnow, loach (<i>Tiaroga cobitis</i>) -Wherever found	E	\$30,876	\$722,768,721
672	Kupukupu makalii (<i>Cyclosorus boydiae</i>) -Wherever found	E	\$30,741	\$722,799,462
672	Nanu (<i>Gardenia remyi</i>) -Wherever found	E	\$30,741	\$722,830,203
674	Orcutt grass, slender (<i>Orcuttia tenuis</i>) -Wherever found	T	\$30,702	\$722,860,905
675	Spikedace (<i>Meda fulgida</i>) -Wherever found	E	\$30,552	\$722,891,457
676	Butterfly, Florida leafwing (<i>Anaea troglodyta floridalis</i>) -Wherever found	E	\$30,500	\$722,921,957
676	damselfly, Rota blue (<i>Ischnura lutea</i>) -Wherever found	E	\$30,500	\$722,952,457
678	Pocketbook, shinyrayed (<i>Hamiota subangulata</i>) -Wherever found	E	\$30,493	\$722,982,950
679	Pearlymussel, littlewing (<i>Pegias fabula</i>) -Wherever found	E	\$30,450	\$723,013,400
680	Flannelbush, Mexican (<i>Fremontodendron mexicanum</i>) -Wherever found	E	\$30,300	\$723,043,700
681	Bean, rayed (<i>Villosa fabalis</i>) -Wherever found	E	\$30,135	\$723,073,835
682	Blazingstar, Ash Meadows (<i>Mentzelia leucophylla</i>) -Wherever found	T	\$30,100	\$723,103,935
682	Gumplant, Ash Meadows (<i>Grindelia fraxinipratensis</i>) -Wherever found	T	\$30,100	\$723,134,035
682	Ivesia, Ash Meadows (<i>Ivesia kingii</i> var. <i>eremica</i>) -Wherever found	T	\$30,100	\$723,164,135
682	Mesa-mint, Otay (<i>Pogogyne nudiuscula</i>) -Wherever found	E	\$30,100	\$723,194,235
682	Milk-vetch, Ash meadows (<i>Astragalus phoenix</i>) -Wherever found	T	\$30,100	\$723,224,335
682	Monardella, willow (Monardella viminea) -Wherever found	E	\$30,100	\$723,254,435
682	Sunray, Ash Meadows (<i>Enceliopsis nudicaulis</i> var. <i>corrugata</i>) -Wherever found	T	\$30,100	\$723,284,535
689	Butterfly, Mariana wandering (<i>Vagrans egistina</i>) -Wherever found	E	\$30,000	\$723,314,535
689	Darter, Okaloosa (<i>Etheostoma okaloosae</i>) -Wherever found	T	\$30,000	\$723,344,535
689	No common name (<i>Cyperus pennatifolius</i>) -Wherever found	E	\$30,000	\$723,374,535
689	Onion, Munz's (<i>Allium munzii</i>) -Wherever found	E	\$30,000	\$723,404,535
689	fern, Florida bristle (<i>Trichomanes punctatum</i> ssp. <i>floridanum</i>) -	E	\$30,000	\$723,434,535
694	Madtom, Neosho (<i>Noturus placidus</i>) -Wherever found	T	\$29,900	\$723,464,435
695	No common name (<i>Microlepia strigosa</i> var. <i>mauiensis</i>) -Wherever found	E	\$29,741	\$723,494,176
696	haha (<i>Cyanea rivularis</i>) -Wherever found	E	\$29,500	\$723,523,676
696	Hornshell, Texas (<i>Popenaias popeii</i>) -Wherever found	E	\$29,500	\$723,553,176
696	Tern, Roseate (<i>Sterna dougallii dougallii</i>) -Western Hemisphere and adjacent oceans, incl. U.S.A. (FL, PR, VI), where not listed as endangered	T	\$29,500	\$723,582,676
699	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) -U.S.A (AZ, NM)	EXPN	\$29,400	\$723,612,076

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
700	piresnake, black (<i>Pituophis melanoleucus lodingi</i>) -Wherever found	T	\$29,330	\$723,641,406
701	Springfish, Railroad Valley (<i>Crenichthys nevadae</i>) -Wherever found	T	\$29,092	\$723,670,498
702	Darter, bayou (<i>Etheostoma rubrum</i>) -Wherever found	T	\$29,000	\$723,699,498
702	Lily, Pitkin Marsh (<i>Lilium pardalinum</i> ssp. <i>pitkinense</i>) -Wherever found	E	\$29,000	\$723,728,498
702	Aster, Florida golden (<i>Chrysopsis floridana</i>) -	E	\$29,000	\$723,757,498
705	Fairy shrimp, longhorn (<i>Branchinecta longiantenna</i>) -Wherever found	E	\$28,946	\$723,786,444
706	Panda, giant (<i>Ailuropoda melanoleuca</i>) -Wherever found	E	\$28,900	\$723,815,344
707	Shrimp, anchialine pool (<i>Procaris hawaiana</i>) -Wherever found	E	\$28,704	\$723,844,048
708	Coqui, Llanero (<i>Eleutherodactylus juanariveroi</i>) -Wherever found	E	\$28,500	\$723,872,548
708	Gerardia, sandplain (<i>Agalinis acuta</i>) -Wherever found	E	\$28,500	\$723,901,048
708	Hoawa (<i>Pittosporum hawaiiense</i>) -Wherever found	E	\$28,500	\$723,929,548
708	Ma`oli`oli (<i>Schiedea hawaiiensis</i>) -Wherever found	E	\$28,500	\$723,958,048
708	Shrimp, anchialine pool (<i>Vetericaris chaceorum</i>) -Wherever found	E	\$28,500	\$723,986,548
713	`Aiea (<i>Nothocestrum latifolium</i>) -Wherever found	E	\$28,241	\$724,014,789
714	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. <i>wrighti</i>)) -Wherever found	E	\$28,200	\$724,042,989
715	Crayfish, Hell Creek Cave (<i>Cambarus zophonastes</i>) -Wherever found	E	\$28,000	\$724,070,989
715	Darter, goldline (<i>Percina aurolineata</i>) -Wherever found	T	\$28,000	\$724,098,989
715	Leopard, clouded (<i>Neofelis nebulosa</i>) -Wherever found	E	\$28,000	\$724,126,989
715	No common name (<i>Amaranthus brownii</i>) -Wherever found	E	\$28,000	\$724,154,989
719	Alani (<i>Melicope ovalis</i>) -Wherever found	E	\$27,983	\$724,182,972
719	haha (<i>Cyanea kunthiana</i>) -Wherever found	E	\$27,983	\$724,210,955
721	Tree snail, Newcomb's (<i>Newcombia cumingi</i>) -Wherever found	E	\$27,630	\$724,238,585
722	picture-wing fly, Hawaiian (<i>Drosophila mulli</i>) -Wherever found	T	\$27,430	\$724,266,015
723	Naucorid, Ash Meadows (<i>Ambrysus amargosus</i>) -Wherever found	T	\$27,100	\$724,293,115
724	Ladies-tresses, Navasota (<i>Spiranthes parksii</i>) -Wherever found	E	\$27,000	\$724,320,115
724	Rail, Guam (<i>Rallus owstoni</i>) -Rota	EXPN	\$27,000	\$724,347,115
726	Bird's-beak, salt marsh (<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>) -Wherever found	E	\$26,700	\$724,373,815
727	`Awikiwiki (<i>Canavalia pubescens</i>) -Wherever found	E	\$26,500	\$724,400,315
727	Butterfly, Bartram's hairstreak (<i>Strymon acis bartrami</i>) -Wherever found	E	\$26,500	\$724,426,815
727	Pocketbook, finelined (<i>Hamiota altilis</i>) -Wherever found	T	\$26,500	\$724,453,315
730	Butterfly, San Bruno elfin (<i>Callophrys mossii bayensis</i>) -Wherever found	E	\$26,200	\$724,479,515
731	Madtom, smoky (<i>Noturus baileyi</i>) -Wherever found	E	\$26,012	\$724,505,527
732	Gladeceess, Texas golden (<i>Leavenworthia texana</i>) -Wherever found	E	\$26,000	\$724,531,527
732	Grass, Tennessee yellow-eyed (<i>Xyris tennesseensis</i>) -Wherever found	E	\$26,000	\$724,557,527
732	Rocksnailed, painted (<i>Leptoxis taeniata</i>) -Wherever found	T	\$26,000	\$724,583,527
732	snail, noonday (<i>Mesodon clarki nantahala</i>) -Wherever found	T	\$26,000	\$724,609,527

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
736	Wallflower, Menzies' (<i>Erysimum menziesii</i>) -Wherever found	E	\$25,800	\$724,635,327
737	haha (<i>Cyanea stictophylla</i>) -Wherever found	E	\$25,733	\$724,661,060
738	Turtle, Alabama red-bellied (<i>Pseudemys alabamensis</i>) -Wherever found	E	\$25,660	\$724,686,720
739	Skyrocket, Pagosa (<i>Ipomopsis polyantha</i>) -Wherever found	E	\$25,650	\$724,712,370
740	Crabgrass, Florida pineland (<i>Digitaria pauciflora</i>) -	T	\$25,562	\$724,737,932
741	Tapir, mountain (<i>Tapirus pinchaque</i>) -Wherever found	E	\$25,500	\$724,763,432
741	yellow-faced bee, Hawaiian (<i>Hylaeus kuakea</i>) -Wherever found	E	\$25,500	\$724,788,932
741	Rosemary, Apalachicola (<i>Conradina glabra</i>) -	E	\$25,500	\$724,814,432
741	silverbush, Blodgett's (<i>Argythamnia blodgettii</i>) -	T	\$25,500	\$724,839,932
745	monkey-flower, Michigan (<i>Mimulus michiganensis</i>) -Wherever found	E	\$25,200	\$724,865,132
746	Mustard, Penland alpine fen (<i>Eutrema penlandii</i>) -Wherever found	T	\$25,125	\$724,890,257
747	Spineflower, Orcutt's (<i>Chorizanthe orcuttiana</i>) -Wherever found	E	\$25,100	\$724,915,357
748	picture-wing fly, Hawaiian (<i>Drosophila differens</i>) -Wherever found	E	\$25,038	\$724,940,395
749	Buttercup, autumn (<i>Ranunculus aestivalis</i> (=acriiformis)) -Wherever found	E	\$25,000	\$724,965,395
749	Butterfly, Miami Blue (<i>Cyclargus</i> (=Hemiargus) thomasi bethunebakeri) -Wherever found	E	\$25,000	\$724,990,395
749	Clarkia, Vine Hill (<i>Clarkia imbricata</i>) -Wherever found	E	\$25,000	\$725,015,395
749	Pocketbook, Ouachita rock (<i>Arcidens wheeleri</i>) -Wherever found	E	\$25,000	\$725,040,395
749	Rabbitsfoot, rough (<i>Quadrula cylindrica strigillata</i>) -Wherever found	E	\$25,000	\$725,065,395
749	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$25,000	\$725,090,395
749	yellow-faced bee, Hawaiian (<i>Hylaeus mana</i>) -Wherever found	E	\$25,000	\$725,115,395
756	Eagle, harpy (<i>Harpia harpyja</i>) -Wherever found	E	\$24,818	\$725,140,213
757	Anunu (<i>Sicyos macrophyllus</i>) -Wherever found	E	\$24,733	\$725,164,946
758	Polygonum, Scotts Valley (<i>Polygonum hickmanii</i>) -Wherever found	E	\$24,600	\$725,189,546
758	spineflower, Scotts Valley (<i>Chorizanthe robusta</i> var. hartwegii) -Wherever found	E	\$24,600	\$725,214,146
760	Amphipod, Kauai cave (<i>Spelaeorchestia koloana</i>) -Wherever found	E	\$24,500	\$725,238,646
760	Bat, little Mariana fruit (<i>Pteropus tokudae</i>) -Wherever found	E	\$24,500	\$725,263,146
760	Boa, Virgin Islands tree (<i>Chilabothrus granti</i>) -Wherever found	E	\$24,500	\$725,287,646
760	darter, Sickie (<i>Percina williamsi</i>) -Wherever found	T	\$24,500	\$725,312,146
760	No common name (<i>Schiedea kauaiensis</i>) -Wherever found	E	\$24,500	\$725,336,646
760	Spider, Kauai cave wolf or pe'e pe'e maka 'ole (<i>Adelocosa anops</i>) -Wherever found	E	\$24,500	\$725,361,146
760	Gooseberry, Miccosukee (<i>Ribes echinellum</i>) -	T	\$24,500	\$725,385,646
767	picture-wing fly, Hawaiian (<i>Drosophila heteroneura</i>) -Wherever found	E	\$24,430	\$725,410,076

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
768	Fern, American hart's-tongue (<i>Asplenium scolopendrium</i> var. <i>americanum</i>) -Wherever found	T	\$24,000	\$725,434,076
768	Pauoa (<i>Ctenitis squamigera</i>) -Wherever found	E	\$24,000	\$725,458,076
768	picture-wing fly, Hawaiian (<i>Drosophila tarphytrichia</i>) -Wherever found	E	\$24,000	\$725,482,076
768	yellow-faced bee, Hawaiian (<i>Hylaeus longiceps</i>) -Wherever found	E	\$24,000	\$725,506,076
772	Moccasinshell, Ochlockonee (<i>Medionidus simpsonianus</i>) -Wherever found	E	\$23,993	\$725,530,069
773	Waterdog, black warrior (=Sipsey Fork) (<i>Necturus alabamensis</i>) -Wherever found	E	\$23,850	\$725,553,919
774	Pearl mussel, cracking (<i>Hemistena lata</i>) -Wherever found; Except where listed as Experimental Populations	E	\$23,635	\$725,577,554
775	Kamapua`a (<i>Kadua fluvialis</i>) -Wherever found	E	\$23,500	\$725,601,054
775	picture-wing fly, Hawaiian (<i>Drosophila aglaia</i>) -Wherever found	E	\$23,500	\$725,624,554
777	Beetle, Mount Hermon June (<i>Polyphylla barbata</i>) -Wherever found	E	\$23,100	\$725,647,654
777	Bulrush, Northeastern (<i>Scirpus ancistrochaetus</i>) -	E	\$23,100	\$725,670,754
779	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the North Indian Ocean, bounded by: Africa and Asia in the west and north; 84 degrees E. Long. in the east; and the equator in the south	T	\$23,000	\$725,693,754
780	Moccasinshell, Gulf (<i>Medionidus penicillatus</i>) -Wherever found	E	\$22,993	\$725,716,747
781	Owl's-clover, fleshy (<i>Castilleja campestris</i> ssp. <i>succulenta</i>) -Wherever found	T	\$22,902	\$725,739,649
782	Darter, leopard (<i>Percina pantherina</i>) -Wherever found	T	\$22,900	\$725,762,549
783	Clarkia, Pismo (<i>Clarkia speciosa</i> ssp. <i>immaculata</i>) -Wherever found	E	\$22,700	\$725,785,249
783	Rosemary, Cumberland (<i>Conradina verticillata</i>) -Wherever found	T	\$22,700	\$725,807,949
785	Yellow-faced bee, assimulans (<i>Hylaeus assimulans</i>) -Wherever found	E	\$22,606	\$725,830,555
786	Aster, decurrent false (<i>Boltonia decurrens</i>) -Wherever found	T	\$22,500	\$725,853,055
786	Clubshell, ovate (<i>Pleurobema perovatum</i>) -Wherever found	E	\$22,500	\$725,875,555
786	No common name (<i>Tinospora homosepala</i>) -Wherever found	E	\$22,500	\$725,898,055
789	Goldenrod, Short's (<i>Solidago shortii</i>) -Wherever found	E	\$22,300	\$725,920,355
790	Dudleya, Santa Cruz Island (<i>Dudleya nesiotica</i>) -Wherever found	T	\$22,200	\$725,942,555
791	Pearl mussel, birdwing (<i>Lemiox rimosus</i>) -Wherever found; Except where listed as Experimental Populations	E	\$22,135	\$725,964,690
792	Elimia, lacy (snail) (<i>Elimia crenatella</i>) -Wherever found	T	\$22,000	\$725,986,690
792	Isopod, Lee County cave (<i>Lirceus usdagalun</i>) -Wherever found	E	\$22,000	\$726,008,690
792	nukupuu, Maui (<i>Hemignathus affinis</i>) -Wherever found	E	\$22,000	\$726,030,690
792	Po`ouli (honeycreeper) (<i>Melamprosops phaeosoma</i>) -Wherever found	E	\$22,000	\$726,052,690
792	Sculpin, pygmy (<i>Cottus paulus</i> (=pygmaeus)) -Wherever found	T	\$22,000	\$726,074,690
797	Springfish, Hiko White River (<i>Crenichthys baileyi grandis</i>) -Wherever found	E	\$21,991	\$726,096,681
798	Ground-Dove, Friendly (<i>Gallicolumba stairi</i>) -American Samoa DPS	E	\$21,930	\$726,118,611
799	Barbara's Buttons, Mohr's (<i>Marshallia mohrii</i>) -Wherever found	T	\$21,500	\$726,140,111

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
799	No common name (<i>Pteris lidgatei</i>) -Wherever found	E	\$21,500	\$726,161,611
799	Snail, tulotoma (<i>Tulotoma magnifica</i>) -Wherever found	T	\$21,500	\$726,183,111
799	Lupine, scrub (<i>Lupinus aridorum</i>) -	E	\$21,500	\$726,204,611
803	Bladderpod, Zapata (<i>Physaria thamnophila</i>) -Wherever found	E	\$21,400	\$726,226,011
804	Turtle, flattened musk (<i>Sternotherus depressus</i>) -Black Warrior R. system upstream from Bankhead Dam	T	\$21,350	\$726,247,361
805	beardtongue, Parachute (<i>Penstemon debilis</i>) -Wherever found	T	\$21,300	\$726,268,661
806	No common name (<i>Wikstroemia villosa</i>) -Wherever found	E	\$21,241	\$726,289,902
807	Sea turtle, loggerhead (<i>Caretta caretta</i>) -Loggerhead sea turtles originating from the South Atlantic Ocean south of the equator, north of 60 degrees S. Lat., west of 20 degrees E. Long., and east of 67 degrees W. Long	T	\$21,200	\$726,311,102
808	Cavefish, Ozark (<i>Amblyopsis rosae</i>) -Wherever found	T	\$21,100	\$726,332,202
809	Ambrosia, south Texas (<i>Ambrosia cheiranthifolia</i>) -Wherever found	E	\$21,000	\$726,353,202
809	Ayenia, Texas (<i>Ayenia limitaris</i>) -Wherever found	E	\$21,000	\$726,374,202
809	crayfish, Benton County cave (<i>Cambarus aculabrum</i>) -Wherever found	E	\$21,000	\$726,395,202
809	Darter, Cherokee (<i>Etheostoma scotti</i>) -Wherever found	T	\$21,000	\$726,416,202
809	fatmucket, Arkansas (<i>Lampsilis powellii</i>) -Wherever found	T	\$21,000	\$726,437,202
809	Hawk, Puerto Rican broad-winged (<i>Buteo platypterus brunnescens</i>) -Wherever found	E	\$21,000	\$726,458,202
809	Hohiu (<i>Dryopteris glabra</i> var. <i>pusilla</i>) -Wherever found	E	\$21,000	\$726,479,202
809	No common name (<i>Diplazium molokaiense</i>) -Wherever found	E	\$21,000	\$726,500,202
809	No common name (<i>Eugenia bryanii</i>) -Wherever found	E	\$21,000	\$726,521,202
809	White-eye, Rota bridled (<i>Zosterops rotensis</i>) -Wherever found	E	\$21,000	\$726,542,202
819	Fiddleneck, large-flowered (<i>Amsinckia grandiflora</i>) -Wherever found	E	\$20,947	\$726,563,149
820	Aplokating-palaoan (<i>Psychotria malaspinae</i>) -Wherever found	E	\$20,551	\$726,583,700
821	Bean, purple (<i>Villosa perpurpurea</i>) -Wherever found	E	\$20,500	\$726,604,200
821	Heather, mountain golden (<i>Hudsonia montana</i>) -Wherever found	T	\$20,500	\$726,624,700
821	Yellowhead, desert (<i>Yermo xanthocephalus</i>) -Wherever found	T	\$20,500	\$726,645,200
821	Sandmat, pineland (<i>Chamaesyce deltoidea pinetorum</i>) -	T	\$20,500	\$726,665,700
825	haha (<i>Cyanea maritae</i>) -Wherever found	E	\$20,241	\$726,685,941
825	nui, haha (<i>Cyanea horrida</i>) -Wherever found	E	\$20,241	\$726,706,182
827	Trout, Apache (<i>Oncorhynchus apache</i>) -Wherever found	T	\$20,200	\$726,726,382
828	Bison, wood (<i>Bison bison athabasca</i>) -Wherever found	T	\$20,000	\$726,746,382
828	Butterweed, Layne's (<i>Senecio layneae</i>) -Wherever found	T	\$20,000	\$726,766,382
828	Cavesnail, Tumbling Creek (<i>Antrobia culveri</i>) -Wherever found	E	\$20,000	\$726,786,382
828	Darter, amber (<i>Percina antesella</i>) -Wherever found	E	\$20,000	\$726,806,382
828	No common name (<i>Bonamia menziesii</i>) -Wherever found	E	\$20,000	\$726,826,382
828	No common name (<i>Maesa walkeri</i>) -Wherever found	T	\$20,000	\$726,846,382

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
828	No common name (<i>Polyscias bisattenuata</i>) -Wherever found	E	\$20,000	\$726,866,382
828	No common name (<i>Pritchardia lanigera</i>) -Wherever found	E	\$20,000	\$726,886,382
828	Piperia, Yadon's (<i>Piperia yadonii</i>) -Wherever found	E	\$20,000	\$726,906,382
837	Ischaemum, Hilo (<i>Ischaemum byrone</i>) -Wherever found	E	\$19,741	\$726,926,123
838	Pearlymussel, Curtis (<i>Epioblasma florentina curtisii</i>) -Wherever found	E	\$19,700	\$726,945,823
839	penguin, emperor (<i>Aptenodytes forsteri</i>) -Wherever found	T	\$19,663	\$726,965,486
840	Quillwort, Louisiana (<i>Isoetes louisianensis</i>) -Wherever found	E	\$19,500	\$726,984,986
840	Uhi uhi (<i>Mezoneuron kavaiense</i>) -Wherever found	E	\$19,500	\$727,004,486
840	Gourd, Okeechobee (<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>) -	E	\$19,500	\$727,023,986
840	Flax, sand (<i>Linum arenicola</i>) -	E	\$19,500	\$727,043,486
840	Spurge, wedge (<i>Chamaesyce deltoidea serpyllum</i>) -	E	\$19,500	\$727,062,986
845	Lily, Minnesota dwarf trout (<i>Erythronium propullans</i>) -Wherever found	E	\$19,338	\$727,082,324
846	Shiner, Arkansas River (<i>Notropis girardi</i>) -Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$19,240	\$727,101,564
847	Topminnow, Gila (incl. Yaqui) (<i>Poeciliopsis occidentalis</i>) -U.S. Only	E	\$19,130	\$727,120,694
848	Darter, relict (<i>Etheostoma chienense</i>) -Wherever found	E	\$19,000	\$727,139,694
848	Kolea (<i>Myrsine linearifolia</i>) -Wherever found	T	\$19,000	\$727,158,694
848	No common name (<i>Pritchardia viscosa</i>) -Wherever found	E	\$19,000	\$727,177,694
848	Prairie-chicken, Attwater's greater (<i>Tympanuchus cupido attwateri</i>) -Wherever found	E	\$19,000	\$727,196,694
848	Ridge-cress, Barneby (<i>Lepidium barnebyanum</i>) -Wherever found	E	\$19,000	\$727,215,694
848	Snail, Chittenango ovate amber (<i>Novisuccinea chittenangoensis</i>) -Wherever found	T	\$19,000	\$727,234,694
854	(honeyeater), Mao (= maomao) (<i>Gymnomyza samoensis</i>) -Wherever found	E	\$18,930	\$727,253,624
854	Aiea (<i>Nothocestrum breviflorum</i>) -Wherever found	E	\$18,930	\$727,272,554
856	Cactus, Bakersfield (<i>Opuntia treleasei</i>) -Wherever found	E	\$18,800	\$727,291,354
857	Reedgrass, Maui (<i>Calamagrostis expansa</i>) -Wherever found	E	\$18,741	\$727,310,095
858	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) -Wherever found; Except where listed as Experimental Populations	E	\$18,635	\$727,328,730
859	crayfish, Slenderclaw (<i>Cambarus cracens</i>) -Wherever found	E	\$18,500	\$727,347,230
859	Hornsnail, rough (<i>Pleurocera foremani</i>) -Wherever found	E	\$18,500	\$727,365,730
859	Milk-vetch, Jesup's (<i>Astragalus robbinsii</i> var. <i>jesupii</i>) -Wherever found	E	\$18,500	\$727,384,230
859	No common name (<i>Delissea rhytidosperma</i>) -Wherever found	E	\$18,500	\$727,402,730
863	Ass, Asian wild (<i>Equus hemionus</i>) -Wherever found	E	\$18,400	\$727,421,130
864	chub, Peppered (<i>Macrhybopsis tetranema</i>) -Wherever found	E	\$18,300	\$727,439,430
865	haha (<i>Cyanea glabra</i>) -Wherever found	E	\$18,241	\$727,457,671
866	Orcutt grass, Sacramento (<i>Orcuttia viscida</i>) -Wherever found	E	\$18,147	\$727,475,818
867	Manzanita, Morro (<i>Arctostaphylos morroensis</i>) -Wherever found	T	\$18,100	\$727,493,918
868	Chub, slender (<i>Erimystax cahni</i>) -Wherever found	T	\$18,000	\$727,511,918

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
868	darer, bluemark (Etheostoma akatulo) -Wherever found	E	\$18,000	\$727,529,918
868	Darter, vermilion (Etheostoma chermocki) -Wherever found	E	\$18,000	\$727,547,918
868	Kula wahine noho (Isodendron pyrifolium) -Wherever found	E	\$18,000	\$727,565,918
868	Wartyback, white (pearlymussel) (Plethobasus cicatricosus) -Wherever found	E	\$18,000	\$727,583,918
873	Beaked-rush, Knieskern's (Rhynchospora knieskernii) -Wherever found	T	\$17,956	\$727,601,874
874	Chub, Gila (Gila intermedia) -Wherever found	E	\$17,924	\$727,619,798
875	Nohoanu (Geranium multiflorum) -Wherever found	E	\$17,741	\$727,637,539
876	Clover, showy Indian (Trifolium amoenum) -Wherever found	E	\$17,700	\$727,655,239
876	daisy, Lakeside (Hymenoxys herbacea) -Wherever found	T	\$17,700	\$727,672,939
878	`Akoko (Euphorbia remyi var. remyi) -Wherever found	E	\$17,500	\$727,690,439
878	Darter, watercress (Etheostoma nuchale) -Wherever found	E	\$17,500	\$727,707,939
878	Guajon (Eleutherodactylus cooki) -Wherever found	T	\$17,500	\$727,725,439
878	Kaulu (Pteralyxia kauaiensis) -Wherever found	E	\$17,500	\$727,742,939
878	Monkeyface, Appalachian (pearlymussel) (Theliderma sparsa) -Wherever found	E	\$17,500	\$727,760,439
878	No common name (Santalum involutum) -Wherever found	E	\$17,500	\$727,777,939
878	No common name (Schiedea nuttallii) -Wherever found	E	\$17,500	\$727,795,439
878	Rock-cress, Hoffmann's (Arabis hoffmannii) -Wherever found	E	\$17,500	\$727,812,939
886	Reed-mustard, shrubby (Schoenocrambe suffrutescens) -Wherever found	E	\$17,486	\$727,830,425
887	Wawae`iole (Huperzia mannii) -Wherever found	E	\$17,241	\$727,847,666
888	Bedstraw, island (Galium buxifolium) -Wherever found	E	\$17,200	\$727,864,866
888	Trout, Gila (Oncorhynchus gilae) -Wherever found	T	\$17,200	\$727,882,066
890	Butterfly, Uncompahgre fritillary (Boloria acrocneuma) -Wherever found	E	\$17,100	\$727,899,166
891	Dogweed, ashy (Thymophylla tephroleuca) -Wherever found	E	\$17,000	\$727,916,166
891	Vetch, Hawaiian (Vicia menziesii) -Wherever found	E	\$17,000	\$727,933,166
893	No common name (Neraudia ovata) -Wherever found	E	\$16,803	\$727,949,969
894	Lupine, Nipomo Mesa (Lupinus nipomensis) -Wherever found	E	\$16,800	\$727,966,769
895	`O`u (honeycreeper) (Psittirostra psittacea) -Wherever found	E	\$16,791	\$727,983,560
896	Arrowhead, bunched (Sagittaria fasciculata) -Wherever found	E	\$16,500	\$728,000,060
896	Blackbird, yellow-shouldered (Agelaius xanthomus) -Wherever found	E	\$16,500	\$728,016,560
896	Kauila (Colubrina oppositifolia) -Wherever found	E	\$16,500	\$728,033,060
896	Madtom, pygmy (Noturus stanauli) -Wherever found	E	\$16,500	\$728,049,560
896	Tortoise, Bolson (Gopherus flavomarginatus) -Wherever found	E	\$16,500	\$728,066,060
896	Warea, wide-leaf (Warea amplexifolia) -	E	\$16,500	\$728,082,560
902	picture-wing fly, Hawaiian (Drosophila ochrobasis) -Wherever found	E	\$16,430	\$728,098,990
903	Bladderpod, Dudley Bluffs (Lesquerella congesta) -Wherever found	T	\$16,425	\$728,115,415
903	Twinpod, Dudley Bluffs (Physaria obcordata) -Wherever found	T	\$16,425	\$728,131,840
905	Yerba santa, Lompoc (Eriodictyon capitatum) -Wherever found	E	\$16,100	\$728,147,940

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
906	Meadowfoam, Butte County (<i>Limnanthes floccosa</i> ssp. <i>californica</i>) -Wherever found	E	\$16,002	\$728,163,942
907	Acornshell, southern (<i>Epioblasma othcaloogensis</i>) -Wherever found	E	\$16,000	\$728,179,942
907	Dace, Kendall Warm Springs (<i>Rhinichthys osculus thermalis</i>) -Wherever found	E	\$16,000	\$728,195,942
907	Damselfly, oceanic Hawaiian (<i>Megalagrion oceanicum</i>) -Wherever found	E	\$16,000	\$728,211,942
907	Manioc, Walker's (<i>Manihot walkerae</i>) -Wherever found	E	\$16,000	\$728,227,942
907	Manzanita, Franciscan (<i>Arctostaphylos franciscana</i>) -Wherever found	E	\$16,000	\$728,243,942
907	Pitcher-plant, green (<i>Sarracenia oreophila</i>) -Wherever found	E	\$16,000	\$728,259,942
913	Birds-in-a-nest, white (<i>Macbridea alba</i>) -	T	\$15,830	\$728,275,772
914	Hala pepe (<i>Pleomele hawaiiensis</i>) -Wherever found	E	\$15,803	\$728,291,575
915	Orcutt grass, hairy (<i>Orcuttia pilosa</i>) -Wherever found	E	\$15,802	\$728,307,377
916	Townsendia, Last Chance (<i>Townsendia aprica</i>) -Wherever found	T	\$15,639	\$728,323,016
917	spineflower, Robust (<i>Chorizanthe robusta</i> var. <i>robusta</i>) -Wherever found	E	\$15,600	\$728,338,616
918	Rock-cress, McDonald's (<i>Arabis macdonaldiana</i>) -Wherever found	E	\$15,551	\$728,354,167
919	Aupaka (<i>Isodendron longifolium</i>) -Wherever found	T	\$15,500	\$728,369,667
919	Centaury, spring-loving (<i>Centaurium namophilum</i>) -Wherever found	T	\$15,500	\$728,385,167
919	Combshell, upland (<i>Epioblasma metastrata</i>) -Wherever found	E	\$15,500	\$728,400,667
919	Damselfly, crimson Hawaiian (<i>Megalagrion leptodemas</i>) -Wherever found	E	\$15,500	\$728,416,167
919	No common name (<i>Schiedea jacobii</i>) -Wherever found	E	\$15,500	\$728,431,667
919	No common name (<i>Stenogyne bifida</i>) -Wherever found	E	\$15,500	\$728,447,167
919	picture-wing fly, Hawaiian (<i>Drosophila musaphilia</i>) -Wherever found	E	\$15,500	\$728,462,667
919	picture-wing fly, Hawaiian (<i>Drosophila neoclavisetae</i>) -Wherever found	E	\$15,500	\$728,478,167
919	picture-wing fly, Hawaiian (<i>Drosophila sharpi</i>) -Wherever found	E	\$15,500	\$728,493,667
919	Yellow-faced bee, easy (<i>Hylaeus facilis</i>) -Wherever found	E	\$15,500	\$728,509,167
929	Bird's beak, palmate-bracted (<i>Cordylanthus palmatus</i>) -Wherever found	E	\$15,449	\$728,524,616
930	No common name (<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>) -Wherever found	E	\$15,303	\$728,539,919
931	Po'e (<i>Portulaca sclerocarpa</i>) -Wherever found	E	\$15,233	\$728,555,152
932	Grass, Eureka Dune (<i>Swallenia alexandrae</i>) -	T	\$15,100	\$728,570,252
933	No common name (<i>Pritchardia maideniana</i>) -Wherever found	E	\$15,046	\$728,585,298
934	Checker-mallow, Keck's (<i>Sidalcea keckii</i>) -Wherever found	E	\$15,002	\$728,600,300
935	`Ala `ala wai nui (<i>Peperomia subpetiolata</i>) -Wherever found	E	\$15,000	\$728,615,300
935	Ceanothus, coyote (<i>Ceanothus ferrisiae</i>) -Wherever found	E	\$15,000	\$728,630,300
935	Isopod, Madison Cave (<i>Antrolana lira</i>) -Wherever found	T	\$15,000	\$728,645,300
935	Logperch, Conasauga (<i>Percina jenkinsi</i>) -Wherever found	E	\$15,000	\$728,660,300
935	Loosestrife, rough-leaved (<i>Lysimachia asperulaefolia</i>) -Wherever found	E	\$15,000	\$728,675,300
935	Milk-vetch, Coachella Valley (<i>Astragalus lentiginosus</i> var. <i>coachellae</i>) -Wherever found	E	\$15,000	\$728,690,300
935	No common name (<i>Phyllostegia hispida</i>) -Wherever found	E	\$15,000	\$728,705,300

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
935	No common name (<i>Phyllostegia pilosa</i>) -Wherever found	E	\$15,000	\$728,720,300
935	No common name (<i>Phyllostegia stachyoides</i>) -Wherever found	E	\$15,000	\$728,735,300
935	No common name (<i>Pritchardia napaliensis</i>) -Wherever found	E	\$15,000	\$728,750,300
935	Olulu (<i>Brighamia insignis</i>) -Wherever found	E	\$15,000	\$728,765,300
935	Spineflower, Howell's (<i>Chorizanthe howellii</i>) -Wherever found	E	\$15,000	\$728,780,300
947	Dudleya, Santa Clara Valley (<i>Dudleya setchellii</i>) -Wherever found	E	\$14,900	\$728,795,200
947	Jewelflower, Metcalf Canyon (<i>Streptanthus albidus</i> ssp. <i>albidus</i>) -Wherever found	E	\$14,900	\$728,810,100
949	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) -In Arizona, an area north of Interstate 8 and south of Interstate 10, bounded by the Colorado River on the west and Interstate 10 on the east; and an area south of Interstate 8, bounded by Highway 85 on the west, Interstates 10 and 19 on the east, and the U.S.-Mexico border on the south	EXPN	\$14,850	\$728,824,950
950	Sunflower, Whorled (<i>Helianthus verticillatus</i>) -	E	\$14,700	\$728,839,650
951	pikeminnow, Colorado (<i>Ptychocheilus lucius</i>) -Salt and Verde R. drainages, AZ	EXPN	\$14,667	\$728,854,317
952	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>brevipes</i>) -Wherever found	E	\$14,500	\$728,868,817
952	akialoa, Kauai (honeycreeper) (<i>Akialoa stejnegeri</i>) -Wherever found	E	\$14,500	\$728,883,317
952	Nioi (<i>Eugenia koolauensis</i>) -Wherever found	E	\$14,500	\$728,897,817
952	No common name (<i>Asplenium dielmannii</i>) -Wherever found	E	\$14,500	\$728,912,317
952	No common name (<i>Schiedea viscosa</i>) -Wherever found	E	\$14,500	\$728,926,817
952	No common name (<i>Sicyos lanceoloideus</i>) -Wherever found	E	\$14,500	\$728,941,317
952	Palo de rosa (<i>Ottoschulzia rhodoxylon</i>) -Wherever found	T	\$14,500	\$728,955,817
952	Shiner, palezone (<i>Notropis albizonatus</i>) -Wherever found	E	\$14,500	\$728,970,317
952	Spinymussel, Altamaha (<i>Elliptio spinosa</i>) -Wherever found	E	\$14,500	\$728,984,817
961	Ko'oko'olau (<i>Bidens micrantha</i> ssp. <i>ctenophylla</i>) -Wherever found	E	\$14,409	\$728,999,226
962	Heau (<i>Exocarpos menziesii</i>) -	E	\$14,303	\$729,013,529
963	Grass, Colusa (<i>Neostapfia colusana</i>) -Wherever found	T	\$14,302	\$729,027,831
964	Spineflower, Ben Lomond (<i>Chorizanthe pungens</i> var. <i>hartwegiana</i>) -Wherever found	E	\$14,300	\$729,042,131
965	Reed-mustard, Barneby (<i>Schoenocrambe barnebyi</i>) -Wherever found	E	\$14,237	\$729,056,368
966	Sedge, Navajo (<i>Carex specuicola</i>) -Wherever found	T	\$14,133	\$729,070,501
967	Aupaka (<i>Isodendron laurifolium</i>) -Wherever found	E	\$14,000	\$729,084,501
967	Bladderpod, kodachrome (<i>Lesquerella tumulosa</i>) -Wherever found	E	\$14,000	\$729,098,501
967	Dace, Ash Meadows speckled (<i>Rhinichthys osculus nevadensis</i>) -Wherever found	E	\$14,000	\$729,112,501
967	Damselfly, flying earwig Hawaiian (<i>Megalagrion nesiotes</i>) -Wherever found	E	\$14,000	\$729,126,501
967	Fescue, Guadalupe (<i>Festuca ligulata</i>) -Wherever found	E	\$14,000	\$729,140,501
967	Gladecress, Fleshy-fruit (<i>Leavenworthia crassa</i>) -Wherever found	E	\$14,000	\$729,154,501
967	Ko'oloa'ula (<i>Abutilon menziesii</i>) -Wherever found	E	\$14,000	\$729,168,501
967	Kopiko (<i>Psychotria grandiflora</i>) -Wherever found	E	\$14,000	\$729,182,501
967	Milk-vetch, heliotrope (<i>Astragalus montii</i>) -Wherever found	T	\$14,000	\$729,196,501

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
967	No common name (<i>Deparia kaalaana</i>) -Wherever found	E	\$14,000	\$729,210,501
967	No common name (<i>Gouania vitifolia</i>) -Wherever found	E	\$14,000	\$729,224,501
967	No common name (<i>Lysimachia maxima</i>) -Wherever found	E	\$14,000	\$729,238,501
967	No common name (<i>Platanthera holochila</i>) -Wherever found	E	\$14,000	\$729,252,501
967	Rocksnail, interrupted (=Georgia) (<i>Leptoxis foremani</i>) -Wherever found	E	\$14,000	\$729,266,501
967	Rush-pea, slender (<i>Hoffmannseggia tenella</i>) -Wherever found	E	\$14,000	\$729,280,501
967	Snake, Atlantic salt marsh (<i>Nerodia clarkii taeniata</i>) -Wherever found	T	\$14,000	\$729,294,501
983	No common name (<i>Silene hawaiiensis</i>) -Wherever found	T	\$13,930	\$729,308,431
984	Dudleya, marcescent (<i>Dudleya cymosa</i> ssp. <i>marcescens</i>) -Wherever found	T	\$13,700	\$729,322,131
985	Reed-mustard, clay (<i>Schoenocrambe argillacea</i>) -Wherever found	T	\$13,691	\$729,335,822
986	Chub, Hutton tui (<i>Gila bicolor</i> ssp.) -Wherever found	T	\$13,600	\$729,349,422
987	`O`o, Kauai (honeyeater) (<i>Moho braccatus</i>) -Wherever found	E	\$13,500	\$729,362,922
987	Darter, rush (<i>Etheostoma phytophilum</i>) -Wherever found	E	\$13,500	\$729,376,422
987	Ha`iwale (<i>Cyrtandra tintinnabula</i>) -Wherever found	E	\$13,500	\$729,389,922
987	haha (<i>Cyanea st.-johnii</i>) -Wherever found	E	\$13,500	\$729,403,422
987	Iguana, Mona ground (<i>Cyclura stejnegeri</i>) -Wherever found	T	\$13,500	\$729,416,922
987	Kamakahala (<i>Labordia cyrtandrae</i>) -Wherever found	E	\$13,500	\$729,430,422
987	No common name (<i>Huperzia stemmermanniae</i>) -Wherever found	E	\$13,500	\$729,443,922
987	Nohoanu (<i>Geranium kauaiense</i>) -Wherever found	E	\$13,500	\$729,457,422
987	Poppy, Sacramento prickly (<i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i>) -Wherever found	E	\$13,500	\$729,470,922
987	Prickly-ash, St. Thomas (<i>Zanthoxylum thomasianum</i>) -Wherever found	E	\$13,500	\$729,484,422
987	Thrush, large Kauai (=kamao) (<i>Myadestes myadestinus</i>) -Wherever found	E	\$13,500	\$729,497,922
987	Thrush, Molokai (<i>Myadestes lanaiensis</i> <i>rutha</i>) -Wherever found	E	\$13,500	\$729,511,422
987	Toad, Puerto Rican crested (<i>Peltophryne lemur</i>) -Wherever found	T	\$13,500	\$729,524,922
1000	Phacelia, clay (<i>Phacelia argillacea</i>) -Wherever found	E	\$13,060	\$729,537,982
1001	Stonefly, western glacier (<i>Zapada glacier</i>) -Wherever found	T	\$13,050	\$729,551,032
1002	akepa, Maui (<i>Loxops ochraceus</i>) -Wherever found	E	\$13,000	\$729,564,032
1002	alauahio, Oahu (<i>Paroreomyza maculata</i>) -Wherever found	E	\$13,000	\$729,577,032
1002	Cactus, star (<i>Astrophytum asterias</i>) -Wherever found	E	\$13,000	\$729,590,032
1002	Creeper, Molokai (<i>Paroreomyza flammea</i>) -Wherever found	E	\$13,000	\$729,603,032
1002	Gardenia (=Na`u), Hawaiian (<i>Gardenia brighamii</i>) -Wherever found	E	\$13,000	\$729,616,032
1002	Ma`o hau hele, (=native yellow hibiscus) (<i>Hibiscus brackenridgei</i>) -Wherever found	E	\$13,000	\$729,629,032
1002	Mehamehame (<i>Flueggea neowawraea</i>) -Wherever found	E	\$13,000	\$729,642,032
1002	Minnow, Devils River (<i>Dionda diaboli</i>) -Wherever found	T	\$13,000	\$729,655,032
1002	nehe (<i>Melanthera tenuifolia</i>) -Wherever found	E	\$13,000	\$729,668,032
1002	No common name (<i>Doryopteris angelica</i>) -Wherever found	E	\$13,000	\$729,681,032
1002	No common name (<i>Festuca hawaiiensis</i>) -Wherever found	E	\$13,000	\$729,694,032

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1002	No common name (<i>Pritchardia schattaueri</i>) -Wherever found	E	\$13,000	\$729,707,032
1002	No common name (<i>Remya montgomeryi</i>) -Wherever found	E	\$13,000	\$729,720,032
1002	No common name (<i>Xylosma crenatum</i>) -Wherever found	E	\$13,000	\$729,733,032
1002	nukupuu, Kauai (<i>Hemignathus hanapepe</i>) -Wherever found	E	\$13,000	\$729,746,032
1002	Pigtoe, Cumberland (<i>Pleuroia gibber</i>) -Wherever found	E	\$13,000	\$729,759,032
1002	No common name (<i>Phyllostegia helleri</i>) -	E	\$13,000	\$729,772,032
1019	Springfish, White River (<i>Crenichthys baileyi baileyi</i>) -Wherever found	E	\$12,904	\$729,784,936
1020	Spurge, Hoover's (<i>Chamaesyce hooveri</i>) -Wherever found	T	\$12,902	\$729,797,838
1021	Cypress, Santa Cruz (<i>Cupressus abramsiana</i>) -Wherever found	T	\$12,900	\$729,810,738
1022	Avens, spreading (<i>Geum radiatum</i>) -Wherever found	E	\$12,850	\$729,823,588
1023	Makou (<i>Peucedanum sandwicense</i>) -Wherever found	T	\$12,740	\$729,836,328
1024	Fanshell (<i>Cyprogenia stegaria</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$12,700	\$729,849,028
1024	Tuctoria, Greene's (<i>Tuctoria greenei</i>) -Wherever found	E	\$12,700	\$729,861,728
1026	aumakua, Palapalai (<i>Dryopteris crinalis</i> var. <i>podosorus</i>) -Wherever found	E	\$12,500	\$729,874,228
1026	Bladderpod, Spring Creek (<i>Lesquerella perforata</i>) -Wherever found	E	\$12,500	\$729,886,728
1026	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) -Wherever found; Except where listed as Experimental Populations	E	\$12,500	\$729,899,228
1026	Butterfly, Myrtle's silverspot (<i>Speyeria zerene myrtleae</i>) -Wherever found	E	\$12,500	\$729,911,728
1026	Clarkia, Springville (<i>Clarkia springvillensis</i>) -Wherever found	T	\$12,500	\$729,924,228
1026	Koki'o ke'oke'o (<i>Hibiscus waimeae</i> ssp. <i>hannerae</i>) -Wherever found	E	\$12,500	\$729,936,728
1026	Mahoe (<i>Alectryon macrococcus</i>) -Wherever found	E	\$12,500	\$729,949,228
1026	Naupaka, dwarf (<i>Scaevola coriacea</i>) -Wherever found	E	\$12,500	\$729,961,728
1026	No common name (<i>Pritchardia munroi</i>) -Wherever found	E	\$12,500	\$729,974,228
1026	Pa'iniu (<i>Astelia waialealae</i>) -Wherever found	E	\$12,500	\$729,986,728
1036	Monkeyflower, Vandenberg (<i>Diplacus vanderbergensis</i>) -Wherever found	E	\$12,400	\$729,999,128
1037	Lampmussel, Alabama (<i>Lampsilis virescens</i>) -Wherever found; Except where listed as Experimental Populations	E	\$12,135	\$730,011,263
1038	Dudleya, Conejo (<i>Dudleya abramsii</i> ssp. <i>parva</i>) -Wherever found	T	\$12,100	\$730,023,363
1039	Curassow, blue-billed (<i>Crax alberti</i>) -Wherever found	E	\$12,062	\$730,035,425
1040	Beetle, delta green ground (<i>Elaphrus viridis</i>) -Wherever found	T	\$12,002	\$730,047,427
1041	Goldenrod, Houghton's (<i>Solidago houghtonii</i>) -Wherever found	T	\$12,000	\$730,059,427
1041	Ha'iwale (<i>Cyrtandra filipes</i>) -Wherever found	E	\$12,000	\$730,071,427
1041	Ha'iwale (<i>Cyrtandra limahuliensis</i>) -Wherever found	T	\$12,000	\$730,083,427
1041	haha (<i>Cyanea recta</i>) -Wherever found	T	\$12,000	\$730,095,427
1041	Kamakahala (<i>Labordia pumila</i>) -Wherever found	E	\$12,000	\$730,107,427
1041	Kolea (<i>Myrsine fosbergii</i>) -Wherever found	E	\$12,000	\$730,119,427

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1041	Kuawawaenuhu (<i>Schiedea lychnoides</i>) -Wherever found	E	\$12,000	\$730,131,427
1041	lehua makanoe (<i>Lysimachia daphnoides</i>) -Wherever found	E	\$12,000	\$730,143,427
1041	Ma`oli`oli (<i>Schiedea apokremnos</i>) -Wherever found	E	\$12,000	\$730,155,427
1041	No common name (<i>Asplenium diellaciniatum</i>) -Wherever found	E	\$12,000	\$730,167,427
1041	No common name (<i>Keysseria</i> (=Lagenifera) <i>erici</i>) -Wherever found	E	\$12,000	\$730,179,427
1041	No common name (<i>Keysseria</i> (=Lagenifera) <i>helenae</i>) -Wherever found	E	\$12,000	\$730,191,427
1041	No common name (<i>Phyllostegia parviflora</i>) -Wherever found	E	\$12,000	\$730,203,427
1041	No common name (<i>Polyscias racemosa</i>) -Wherever found	E	\$12,000	\$730,215,427
1041	No common name (<i>Remya kauaiensis</i>) -Wherever found	E	\$12,000	\$730,227,427
1041	No common name (<i>Schiedea membranacea</i>) -Wherever found	E	\$12,000	\$730,239,427
1041	Pigtoe, Georgia (<i>Pleurobema hanleyianum</i>) -Wherever found	E	\$12,000	\$730,251,427
1041	Primrose, Maguire (<i>Primula maguirei</i>) -Wherever found	T	\$12,000	\$730,263,427
1041	Ziziphus, Florida (<i>Ziziphus celata</i>) -	E	\$12,000	\$730,275,427
1060	Slabshell, Chipola (<i>Elliptio chipolaensis</i>) -Wherever found	T	\$11,993	\$730,287,420
1061	Tiger beetle, Ohlone (<i>Cicindela ohlone</i>) -Wherever found	E	\$11,800	\$730,299,220
1062	Sunburst, San Joaquin adobe (<i>Pseudobahia peirsonii</i>) -Wherever found	T	\$11,700	\$730,310,920
1063	Spinedace, Big Spring (<i>Lepidomeda mollispinis pratensis</i>) -Wherever found	T	\$11,606	\$730,322,526
1064	`Akoko (<i>Euphorbia eleanoriae</i>) -Wherever found	E	\$11,500	\$730,334,026
1064	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i>) -Wherever found	E	\$11,500	\$730,345,526
1064	Alani (<i>Melicope haupuensis</i>) -Wherever found	E	\$11,500	\$730,357,026
1064	Cavefish, Alabama (<i>Speoplatyrhinus poulsoni</i>) -Wherever found	E	\$11,500	\$730,368,526
1064	Crane, whooping (<i>Grus americana</i>) -U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$11,500	\$730,380,026
1064	haha (<i>Cyanea dolichopoda</i>) -Wherever found	E	\$11,500	\$730,391,526
1064	haha (<i>Cyanea eleeleensis</i>) -Wherever found	E	\$11,500	\$730,403,026
1064	haha (<i>Cyanea gibsonii</i>) -Wherever found	E	\$11,500	\$730,414,526
1064	haha (<i>Cyanea grimesiana</i> ssp. <i>grimesiana</i>) -Wherever found	E	\$11,500	\$730,426,026
1064	haha (<i>Cyanea kolekoleensis</i>) -Wherever found	E	\$11,500	\$730,437,526
1064	haha (<i>Cyanea kuhihewa</i>) -Wherever found	E	\$11,500	\$730,449,026
1064	haha (<i>Cyanea lobata</i>) -Wherever found	E	\$11,500	\$730,460,526
1064	haha (<i>Cyanea remyi</i>) -Wherever found	E	\$11,500	\$730,472,026
1064	haha (<i>Cyanea undulata</i>) -Wherever found	E	\$11,500	\$730,483,526
1064	Haiwale (<i>Cyrtandra paliku</i>) -Wherever found	E	\$11,500	\$730,495,026
1064	Hau kuahiwi (<i>Hibiscadelphus woodii</i>) -Wherever found	E	\$11,500	\$730,506,526
1064	Heau (<i>Exocarpos luteolus</i>) -Wherever found	E	\$11,500	\$730,518,026
1064	Hibiscus, Clay's (<i>Hibiscus clayi</i>) -Wherever found	E	\$11,500	\$730,529,526
1064	Iliau, dwarf (<i>Wilkesia hobydi</i>) -Wherever found	E	\$11,500	\$730,541,026

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1064	Kamakahala (<i>Labordia tinifolia</i> var. <i>wahiawaensis</i>) -Wherever found	E	\$11,500	\$730,552,526
1064	Koki'o (<i>Kokia kauaiensis</i>) -Wherever found	E	\$11,500	\$730,564,026
1064	Kolea (<i>Myrsine mezii</i>) -Wherever found	E	\$11,500	\$730,575,526
1064	Laulihilihi (<i>Schiedea stellarioides</i>) -Wherever found	E	\$11,500	\$730,587,026
1064	Na'ena'e (<i>Dubautia pauciflorula</i>) -Wherever found	E	\$11,500	\$730,598,526
1064	Na'ena'e (<i>Dubautia waialealae</i>) -Wherever found	E	\$11,500	\$730,610,026
1064	Naenae (<i>Dubautia kalalauensis</i>) -Wherever found	E	\$11,500	\$730,621,526
1064	Naenae (<i>Dubautia kenwoodii</i>) -Wherever found	E	\$11,500	\$730,633,026
1064	nehe (<i>Lipochaeta micrantha</i>) -Wherever found	E	\$11,500	\$730,644,526
1064	nehe (<i>Lipochaeta waimeaensis</i>) -Wherever found	E	\$11,500	\$730,656,026
1064	No common name (<i>Asplenium dielpallidum</i>) -Wherever found	E	\$11,500	\$730,667,526
1064	No common name (<i>Hesperomannia lydgatei</i>) -Wherever found	E	\$11,500	\$730,679,026
1064	No common name (<i>Kadua st.-johnii</i>) -Wherever found	E	\$11,500	\$730,690,526
1064	No common name (<i>Lysimachia iniki</i>) -Wherever found	E	\$11,500	\$730,702,026
1064	No common name (<i>Lysimachia pendens</i>) -Wherever found	E	\$11,500	\$730,713,526
1064	No common name (<i>Phyllostegia knudsenii</i>) -Wherever found	E	\$11,500	\$730,725,026
1064	No common name (<i>Phyllostegia waimeae</i>) -Wherever found	E	\$11,500	\$730,736,526
1064	No common name (<i>Phyllostegia wawrana</i>) -Wherever found	E	\$11,500	\$730,748,026
1064	No common name (<i>Polyscias flynnii</i>) -Wherever found	E	\$11,500	\$730,759,526
1064	No common name (<i>Pritchardia bakeri</i>) -Wherever found	E	\$11,500	\$730,771,026
1064	No common name (<i>Schiedea kaalae</i>) -Wherever found	E	\$11,500	\$730,782,526
1064	No common name (<i>Stenogyne kanehoana</i>) -Wherever found	E	\$11,500	\$730,794,026
1064	Pentachaeta, Lyon's (<i>Pentachaeta lyonii</i>) -Wherever found	E	\$11,500	\$730,805,526
1064	Shrimp, Kentucky cave (<i>Palaemonias ganteri</i>) -Wherever found	E	\$11,500	\$730,817,026
1064	Yellow-faced bee, hilaris (<i>Hylaeus hilaris</i>) -Wherever found	E	\$11,500	\$730,828,526
1108	Anunu (<i>Sicyos albus</i>) -Wherever found	E	\$11,303	\$730,839,829
1109	Dog, African wild (<i>Lycaon pictus</i>) -Wherever found	E	\$11,300	\$730,851,129
1110	Meadowrue, Cooley's (<i>Thalictrum cooleyi</i>) -Wherever found	E	\$11,100	\$730,862,229
1111	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) -Wherever found; Except where listed as Experimental Populations	E	\$11,000	\$730,873,229
1111	Cactus, black lace (<i>Echinocereus reichenbachii</i> var. <i>albertii</i>) -Wherever found	E	\$11,000	\$730,884,229
1111	Ha'iwale (<i>Cyrtandra kaulantha</i>) -Wherever found	E	\$11,000	\$730,895,229
1111	Ha'iwale (<i>Cyrtandra munroi</i>) -Wherever found	E	\$11,000	\$730,906,229
1111	haha (<i>Cyanea acuminata</i>) -Wherever found	E	\$11,000	\$730,917,229
1111	haha (<i>Cyanea crispa</i>) -Wherever found	E	\$11,000	\$730,928,229
1111	Ma'oli'oli (<i>Schiedea pubescens</i>) -Wherever found	E	\$11,000	\$730,939,229
1111	No common name (<i>Lepidium orbiculare</i>) -Wherever found	E	\$11,000	\$730,950,229

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1111	No common name (<i>Mitracarpus polycladus</i>) -Wherever found	E	\$11,000	\$730,961,229
1111	No common name (<i>Phyllostegia warshaueri</i>) -Wherever found	E	\$11,000	\$730,972,229
1111	No common name (<i>Schiedea laui</i>) -Wherever found	E	\$11,000	\$730,983,229
1111	No common name (<i>Schiedea spergulina</i> var. <i>spergulina</i>) -Wherever found	T	\$11,000	\$730,994,229
1111	Papala (<i>Charpentiera densiflora</i>) -Wherever found	E	\$11,000	\$731,005,229
1111	pilo (<i>Kadua laxiflora</i>) -Wherever found	E	\$11,000	\$731,016,229
1111	Sand-verbena, large-fruited (<i>Abronia macrocarpa</i>) -Wherever found	E	\$11,000	\$731,027,229
1126	Pigtoe, narrow (<i>Fusconaia escambia</i>) -Wherever found	T	\$10,993	\$731,038,222
1127	Hau kuahiwi (<i>Hibiscadelphus giffardianus</i>) -Wherever found	E	\$10,803	\$731,049,025
1127	Kuahiwi laukahi (<i>Plantago hawaiiensis</i>) -Wherever found	E	\$10,803	\$731,059,828
1129	Pennyroyal, Todsen's (<i>Hedeoma todsenii</i>) -Wherever found	E	\$10,750	\$731,070,578
1130	amole, purple (<i>Chlorogalum purpureum</i>) -Wherever found	T	\$10,700	\$731,081,278
1131	Cactus, Pima pineapple (<i>Coryphantha scheeri</i> var. <i>robustispina</i>) -Wherever found	E	\$10,600	\$731,091,878
1132	`Awikiwiki (<i>Canavalia molokaiensis</i>) -Wherever found	E	\$10,500	\$731,102,378
1132	diellia, Asplenium-leaved (<i>Asplenium dielirectum</i>) -Wherever found	E	\$10,500	\$731,112,878
1132	Ho`awa (<i>Pittosporum napaliense</i>) -Wherever found	E	\$10,500	\$731,123,378
1132	Ihi`ihi (<i>Marsilea villosa</i>) -Wherever found	E	\$10,500	\$731,133,878
1132	Kamakahala (<i>Labordia helleri</i>) -Wherever found	E	\$10,500	\$731,144,378
1132	Kamakahala (<i>Labordia lydgatei</i>) -Wherever found	E	\$10,500	\$731,154,878
1132	No common name (<i>Catesbaea melanocarpa</i>) -Wherever found	E	\$10,500	\$731,165,378
1132	No common name (<i>Schiedea obovata</i>) -Wherever found	E	\$10,500	\$731,175,878
1132	No common name (<i>Silene alexandri</i>) -Wherever found	E	\$10,500	\$731,186,378
1132	No common name (<i>Silene perlmannii</i>) -Wherever found	E	\$10,500	\$731,196,878
1132	No common name (<i>Tetramolopium remyi</i>) -Wherever found	E	\$10,500	\$731,207,378
1132	No common name (<i>Tetramolopium rockii</i>) -Wherever found	T	\$10,500	\$731,217,878
1132	Pamakani (<i>Tetramolopium capillare</i>) -Wherever found	E	\$10,500	\$731,228,378
1132	Sandalwood, Lanai (=`iliahi) (<i>Santalum haleakalae</i> var. <i>lanaiense</i>) -Wherever found	E	\$10,500	\$731,238,878
1132	Snail, armored (<i>Marstonia pachyta</i>) -Wherever found	E	\$10,500	\$731,249,378
1147	Cactus, Siler pincushion (<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>) -Wherever found	T	\$10,496	\$731,259,874
1148	Pigtoe, tapered (<i>Fusconaia burkei</i>) -Wherever found	T	\$10,493	\$731,270,367
1148	Sandshell, Southern (<i>Hamiota australis</i>) -Wherever found	T	\$10,493	\$731,280,860
1150	Honohono (<i>Haplostachys haplostachya</i>) -Wherever found	E	\$10,303	\$731,291,163
1151	`Ahinahina (<i>Argyroxiphium sandwicense</i> ssp. <i>sandwicense</i>) -Wherever found	E	\$10,051	\$731,301,214
1152	`Oha wai (<i>Clermontia drepanomorpha</i>) -Wherever found	E	\$10,000	\$731,311,214
1152	Alani (<i>Melicope adscendens</i>) -Wherever found	E	\$10,000	\$731,321,214
1152	Amphipod, Illinois cave (<i>Gammarus acherondytes</i>) -Wherever found	E	\$10,000	\$731,331,214
1152	Campeloma, slender (<i>Campeloma decampi</i>) -Wherever found	E	\$10,000	\$731,341,214

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1152	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$10,000	\$731,351,214
1152	Fern, pendant kihi (<i>Adenophorus periens</i>) -Wherever found	E	\$10,000	\$731,361,214
1152	Ha'iwale (<i>Cyrtandra polyantha</i>) -Wherever found	E	\$10,000	\$731,371,214
1152	haha (<i>Cyanea lanceolata</i>) -Wherever found	E	\$10,000	\$731,381,214
1152	haha (<i>Cyanea magnicalyx</i>) -Wherever found	E	\$10,000	\$731,391,214
1152	haha (<i>Cyanea truncata</i>) -Wherever found	E	\$10,000	\$731,401,214
1152	Hoawa (<i>Pittosporum halophilum</i>) -Wherever found	E	\$10,000	\$731,411,214
1152	Kohe malama malama o kanaloa (<i>Kanaloa kahoolawensis</i>) -Wherever found	E	\$10,000	\$731,421,214
1152	Kolea (<i>Myrsine vaccinioides</i>) -Wherever found	E	\$10,000	\$731,431,214
1152	Kulu'i (<i>Nototrichium humile</i>) -Wherever found	E	\$10,000	\$731,441,214
1152	Liliwai (<i>Acaena exigua</i>) -Wherever found	E	\$10,000	\$731,451,214
1152	No common name (<i>Abutilon eremitopetalum</i>) -Wherever found	E	\$10,000	\$731,461,214
1152	No common name (<i>Cyperus neokunthianus</i>) -Wherever found	E	\$10,000	\$731,471,214
1152	No common name (<i>Hesperomannia arborescens</i>) -Wherever found	E	\$10,000	\$731,481,214
1152	No common name (<i>Lobelia monostachya</i>) -Wherever found	E	\$10,000	\$731,491,214
1152	No common name (<i>Phyllostegia mannii</i>) -Wherever found	E	\$10,000	\$731,501,214
1152	No common name (<i>Polyscias lydgatei</i>) -Wherever found	E	\$10,000	\$731,511,214
1152	No common name (<i>Pritchardia kaalae</i>) -Wherever found	E	\$10,000	\$731,521,214
1152	No common name (<i>Schiedea helleri</i>) -Wherever found	E	\$10,000	\$731,531,214
1152	No common name (<i>Stenogyne kauaulaensis</i>) -Wherever found	E	\$10,000	\$731,541,214
1152	No common name (<i>Tetramolopium lepidotum</i> ssp. <i>lepidotum</i>) -Wherever found	E	\$10,000	\$731,551,214
1152	No common name (<i>Viola lanaiensis</i>) -Wherever found	E	\$10,000	\$731,561,214
1152	olua (<i>Hypolepis hawaiiensis</i> var. <i>mauiensis</i>) -Wherever found	E	\$10,000	\$731,571,214
1152	Pua`ala (<i>Brighamia rockii</i>) -Wherever found	E	\$10,000	\$731,581,214
1152	Schiedea, Diamond Head (<i>Schiedea adamantis</i>) -Wherever found	E	\$10,000	\$731,591,214
1152	Snail, Iowa Pleistocene (<i>Discus macclintocki</i>) -Wherever found	E	\$10,000	\$731,601,214
1152	Cactus, Key tree (<i>Pilosocereus robinii</i>) -	E	\$10,000	\$731,611,214
1183	Meadowfoam, Sebastopol (<i>Limnanthes vinculans</i>) -Wherever found	E	\$9,700	\$731,620,914
1183	Sunshine, Sonoma (<i>Blennosperma bakeri</i>) -Wherever found	E	\$9,700	\$731,630,614
1185	`Aiakeakua, popolo (<i>Solanum sandwicense</i>) -Wherever found	E	\$9,500	\$731,640,114
1185	`Aiea (<i>Nothoestrum peltatum</i>) -Wherever found	E	\$9,500	\$731,649,614
1185	`Akoko (<i>Euphorbia herbstii</i>) -Wherever found	E	\$9,500	\$731,659,114
1185	`Akoko (<i>Euphorbia remyi</i> var. <i>kauaiensis</i>) -Wherever found	E	\$9,500	\$731,668,614
1185	`aku`aku (<i>Cyanea platyphylla</i>) -Wherever found	E	\$9,500	\$731,678,114

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1185	`Anaunau (<i>Lepidium arbuscula</i>) -Wherever found	E	\$9,500	\$731,687,614
1185	`Awikiwiki (<i>Canavalia napaliensis</i>) -Wherever found	E	\$9,500	\$731,697,114
1185	A`e (<i>Zanthoxylum hawaiiense</i>) -Wherever found	E	\$9,500	\$731,706,614
1185	"Akoko (<i>Euphorbia halemanui</i>) -Wherever found	E	\$9,500	\$731,716,114
1185	Alani (<i>Melicope makahae</i>) -Wherever found	E	\$9,500	\$731,725,614
1185	Alani (<i>Melicope mucronulata</i>) -Wherever found	E	\$9,500	\$731,735,114
1185	Alani (<i>Melicope pallida</i>) -Wherever found	E	\$9,500	\$731,744,614
1185	Alani (<i>Melicope puberula</i>) -Wherever found	E	\$9,500	\$731,754,114
1185	Alani (<i>Melicope quadrangularis</i>) -Wherever found	E	\$9,500	\$731,763,614
1185	Bluegrass, Hawaiian (<i>Poa sandvicensis</i>) -Wherever found	E	\$9,500	\$731,773,114
1185	Bluegrass, Mann's (<i>Poa mannii</i>) -Wherever found	E	\$9,500	\$731,782,614
1185	Ha`iwale (<i>Cyrtandra hematos</i>) -Wherever found	E	\$9,500	\$731,792,114
1185	Ha`iwale (<i>Cyrtandra oenobarba</i>) -Wherever found	E	\$9,500	\$731,801,614
1185	haha (<i>Cyanea asarifolia</i>) -Wherever found	E	\$9,500	\$731,811,114
1185	haha (<i>Cyanea grimesiana</i> ssp. <i>obatae</i>) -Wherever found	E	\$9,500	\$731,820,614
1185	haha (<i>Cyanea munroi</i>) -Wherever found	E	\$9,500	\$731,830,114
1185	haha (<i>Cyanea obtusa</i>) -Wherever found	E	\$9,500	\$731,839,614
1185	haha (<i>Cyanea procera</i>) -Wherever found	E	\$9,500	\$731,849,114
1185	haha (<i>Cyanea profuga</i>) -Wherever found	E	\$9,500	\$731,858,614
1185	haha (<i>Cyanea superba</i>) -Wherever found	E	\$9,500	\$731,868,114
1185	Haiwale (<i>Cyrtandra waiolani</i>) -Wherever found	E	\$9,500	\$731,877,614
1185	Kauai hau kuahiwi (<i>Hibiscadelphus distans</i>) -Wherever found	E	\$9,500	\$731,887,114
1185	Kaulu (<i>Pteralyxia macrocarpa</i>) -Wherever found	E	\$9,500	\$731,896,614
1185	Koholapehu (<i>Dubautia latifolia</i>) -Wherever found	E	\$9,500	\$731,906,114
1185	Koki`o, Cooke's (<i>Kokia cookei</i>) -Wherever found	E	\$9,500	\$731,915,614
1185	Kolea (<i>Myrsine knudsenii</i>) -Wherever found	E	\$9,500	\$731,925,114
1185	kopa (<i>Kadua cordata</i> remyi) -Wherever found	E	\$9,500	\$731,934,614
1185	Lau `ehu (<i>Panicum niihauense</i>) -Wherever found	E	\$9,500	\$731,944,114
1185	Mapele (<i>Cyrtandra cyaneoides</i>) -Wherever found	E	\$9,500	\$731,953,614
1185	Na`ena`e (<i>Dubautia imbricata</i> ssp. <i>imbricata</i>) -Wherever found	E	\$9,500	\$731,963,114
1185	Na`ena`e (<i>Dubautia plantaginea</i> ssp. <i>humilis</i>) -Wherever found	E	\$9,500	\$731,972,614
1185	Na`ena`e (<i>Dubautia plantaginea</i> ssp. <i>magnifolia</i>) -Wherever found	E	\$9,500	\$731,982,114
1185	Nanu (<i>Gardenia mannii</i>) -Wherever found	E	\$9,500	\$731,991,614
1185	nehe (<i>Lipochaeta fauriei</i>) -Wherever found	E	\$9,500	\$732,001,114
1185	nehe (<i>Melanthera kamolensis</i>) -Wherever found	E	\$9,500	\$732,010,614
1185	No common name (<i>Achyranthes mutica</i>) -Wherever found	E	\$9,500	\$732,020,114
1185	No common name (<i>Calyptanthus thomasiana</i>) -Wherever found	E	\$9,500	\$732,029,614

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1185	No common name (<i>Delissea undulata</i>) -Wherever found	E	\$9,500	\$732,039,114
1185	No common name (<i>Eugenia woodburyana</i>) -Wherever found	E	\$9,500	\$732,048,614
1185	No common name (<i>Gouania meyenii</i>) -Wherever found	E	\$9,500	\$732,058,114
1185	No common name (<i>Hesperomannia arbuscula</i>) -Wherever found	E	\$9,500	\$732,067,614
1185	No common name (<i>Lysimachia lydgatei</i>) -Wherever found	E	\$9,500	\$732,077,114
1185	No common name (<i>Lysimachia scopulensis</i>) -Wherever found	E	\$9,500	\$732,086,614
1185	No common name (<i>Lysimachia venosa</i>) -Wherever found	E	\$9,500	\$732,096,114
1185	No common name (<i>Neraudia angulata</i>) -Wherever found	E	\$9,500	\$732,105,614
1185	No common name (<i>Neraudia sericea</i>) -Wherever found	E	\$9,500	\$732,115,114
1185	No common name (<i>Phyllostegia hirsuta</i>) -Wherever found	E	\$9,500	\$732,124,614
1185	No common name (<i>Phyllostegia mollis</i>) -Wherever found	E	\$9,500	\$732,134,114
1185	No common name (<i>Poa siphonoglossa</i>) -Wherever found	E	\$9,500	\$732,143,614
1185	No common name (<i>Pritchardia aylmer-robinsonii</i>) -Wherever found	E	\$9,500	\$732,153,114
1185	No common name (<i>Pritchardia hardyi</i>) -Wherever found	E	\$9,500	\$732,162,614
1185	No common name (<i>Stenogyne kaalae</i> ssp. <i>sherffii</i>) -Wherever found	E	\$9,500	\$732,172,114
1185	No common name (<i>Vigna o-wahuensis</i>) -Wherever found	E	\$9,500	\$732,181,614
1185	Nohoanu (<i>Geranium hillebrandii</i>) -Wherever found	E	\$9,500	\$732,191,114
1185	Oha (<i>Delissea subcordata</i>) -Wherever found	E	\$9,500	\$732,200,614
1185	Pamakani (<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>) -Wherever found	E	\$9,500	\$732,210,114
1185	Pilo kea lau li'i (<i>Platydesma rostrata</i>) -Wherever found	E	\$9,500	\$732,219,614
1185	Popolo (<i>Cyanea solanacea</i>) -Wherever found	E	\$9,500	\$732,229,114
1185	Popolo ku mai (<i>Solanum incompletum</i>) -Wherever found	E	\$9,500	\$732,238,614
1185	Snail, flat-spined three-toothed (<i>Triodopsis platysayoides</i>) -Wherever found	T	\$9,500	\$732,248,114
1185	No common name (<i>Wikstroemia skottsbergiana</i>) -	E	\$9,500	\$732,257,614
1185	Flax, Carter's small-flowered (<i>Linum carteri carteri</i>) -	E	\$9,500	\$732,267,114
1185	Sandlace (<i>Polygonella myriophylla</i>) -	E	\$9,500	\$732,276,614
1253	Holei (<i>Ochrosia haleakalae</i>) -Wherever found	E	\$9,303	\$732,285,917
1254	Sunflower, Pecos (=puzzle, =paradox) (<i>Helianthus paradoxus</i>) -Wherever found	T	\$9,250	\$732,295,167
1255	Allocarya, Calistoga (<i>Plagiobothrys strictus</i>) -Wherever found	E	\$9,200	\$732,304,367
1255	Larkspur, Baker's (<i>Delphinium bakeri</i>) -Wherever found	E	\$9,200	\$732,313,567
1255	Paintbrush, Tiburon (<i>Castilleja affinis</i> ssp. <i>neglecta</i>) -Wherever found	E	\$9,200	\$732,322,767
1258	Alani (<i>Melicope munroi</i>) -Wherever found	E	\$9,000	\$732,331,767
1258	Alani (<i>Melicope saint-johnii</i>) -Wherever found	E	\$9,000	\$732,340,767
1258	Alani (<i>Melicope zahlbruckneri</i>) -Wherever found	E	\$9,000	\$732,349,767
1258	Bariaco (<i>Trichilia triacantha</i>) -Wherever found	E	\$9,000	\$732,358,767
1258	bean, sea (<i>Mucuna sloanei</i> var. <i>persericea</i>) -Wherever found	E	\$9,000	\$732,367,767
1258	Beardtongue, Penland (<i>Penstemon penlandii</i>) -Wherever found	E	\$9,000	\$732,376,767

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1258	Clarkia, Presidio (<i>Clarkia franciscana</i>) -Wherever found	E	\$9,000	\$732,385,767
1258	Goldfields, Burke's (<i>Lasthenia burkei</i>) -Wherever found	E	\$9,000	\$732,394,767
1258	haha (<i>Cyanea dunbariae</i>) -Wherever found	E	\$9,000	\$732,403,767
1258	haha (<i>Cyanea duvalliorum</i>) -Wherever found	E	\$9,000	\$732,412,767
1258	haha (<i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i>) -Wherever found	E	\$9,000	\$732,421,767
1258	haha (<i>Cyanea humboldtiana</i>) -Wherever found	E	\$9,000	\$732,430,767
1258	haha (<i>Cyanea pinnatifida</i>) -Wherever found	E	\$9,000	\$732,439,767
1258	Hau kuahiwi (<i>Hibiscadelphus hualalaiensis</i>) -Wherever found	E	\$9,000	\$732,448,767
1258	Kamakahala (<i>Labordia tinifolia</i> var. <i>lanaiensis</i>) -Wherever found	E	\$9,000	\$732,457,767
1258	Kamanomano (<i>Cenchrus agrimonioides</i>) -Wherever found	E	\$9,000	\$732,466,767
1258	Koki'o (<i>Kokia drynarioides</i>) -Wherever found	E	\$9,000	\$732,475,767
1258	Larkspur, yellow (<i>Delphinium luteum</i>) -Wherever found	E	\$9,000	\$732,484,767
1258	Leather flower, Alabama (<i>Clematis socialis</i>) -Wherever found	E	\$9,000	\$732,493,767
1258	No common name (<i>Cyanea kauaulaensis</i>) -Wherever found	E	\$9,000	\$732,502,767
1258	No common name (<i>Phyllostegia kaalaensis</i>) -Wherever found	E	\$9,000	\$732,511,767
1258	No common name (<i>Schiedea hookeri</i>) -Wherever found	E	\$9,000	\$732,520,767
1258	No common name (<i>Schiedea salicaria</i>) -Wherever found	E	\$9,000	\$732,529,767
1258	No common name (<i>Silene lanceolata</i>) -Wherever found	E	\$9,000	\$732,538,767
1258	No common name (<i>Spermolepis hawaiiensis</i>) -Wherever found	E	\$9,000	\$732,547,767
1258	No common name (<i>Tetramolopium arenarium</i>) -Wherever found	E	\$9,000	\$732,556,767
1258	Pu'uka'a (<i>Cyperus trachysanthos</i>) -Wherever found	E	\$9,000	\$732,565,767
1285	Bean, Choctaw (<i>Obovaria choctawensis</i>) -Wherever found	E	\$8,993	\$732,574,760
1285	Pigtoe, fuzzy (<i>Pleurobema strodeanum</i>) -Wherever found	T	\$8,993	\$732,583,753
1287	Cliffrose, Arizona (<i>Purshia</i> (=Cowania) <i>subintegra</i>) -Wherever found	E	\$8,900	\$732,592,653
1288	Ha'iwale (<i>Cyrtandra giffardii</i>) -Wherever found	E	\$8,803	\$732,601,456
1289	Wallflower, Contra Costa (<i>Erysimum capitatum</i> var. <i>angustatum</i>) -Wherever found	E	\$8,700	\$732,610,156
1290	Mapleleaf, winged (<i>Quadrula fragosa</i>) -U.S.A. (AL-specified portions of the Tennessee River; see 17.85(a)(1))	EXPN	\$8,576	\$732,618,732
1291	Blazingstar, Heller's (<i>Liatris helleri</i>) -Wherever found	T	\$8,550	\$732,627,282
1292	`Akoko (<i>Euphorbia deppeana</i>) -Wherever found	E	\$8,500	\$732,635,782
1292	A'e (<i>Zanthoxylum dipetalum</i> var. <i>tomentosum</i>) -Wherever found	E	\$8,500	\$732,644,282
1292	A'e (<i>Zanthoxylum oahuense</i>) -Wherever found	E	\$8,500	\$732,652,782
1292	Alani (<i>Melicope hiiakae</i>) -Wherever found	E	\$8,500	\$732,661,282
1292	Alani (<i>Melicope knudsenii</i>) -Wherever found	E	\$8,500	\$732,669,782
1292	Alani (<i>Melicope lydgatei</i>) -Wherever found	E	\$8,500	\$732,678,282
1292	butterfly, Puerto Rican harlequin (<i>Atlantea tulita</i>) -Wherever found	T	\$8,500	\$732,686,782
1292	Darter, Cumberland (<i>Etheostoma susanae</i>) -Wherever found	E	\$8,500	\$732,695,282

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1292	Fern, Aleutian shield (<i>Polystichum aleuticum</i>) -Wherever found	E	\$8,500	\$732,703,782
1292	Ha'iwale (<i>Cyrtandra viridiflora</i>) -Wherever found	E	\$8,500	\$732,712,282
1292	haha (<i>Cyanea koolauensis</i>) -Wherever found	E	\$8,500	\$732,720,782
1292	haha (<i>Cyanea purpurellifolia</i>) -Wherever found	E	\$8,500	\$732,729,282
1292	Haiwale (<i>Cyrtandra gracilis</i>) -Wherever found	E	\$8,500	\$732,737,782
1292	Hala pepe (<i>Pleomele forbesii</i>) -Wherever found	E	\$8,500	\$732,746,282
1292	Kamakahala (<i>Labordia triflora</i>) -Wherever found	E	\$8,500	\$732,754,782
1292	Kio'ele (<i>Kadua coriacea</i>) -Wherever found	E	\$8,500	\$732,763,282
1292	Kopiko (<i>Psychotria hexandra</i> ssp. <i>oahuensis</i>) -Wherever found	E	\$8,500	\$732,771,782
1292	Love grass, Fosberg's (<i>Eragrostis fosbergii</i>) -Wherever found	E	\$8,500	\$732,780,282
1292	Makou (<i>Ranunculus mauiensis</i>) -Wherever found	E	\$8,500	\$732,788,782
1292	Nani wai'ale'ale (<i>Viola kauaiensis</i> var. <i>wahiawaensis</i>) -Wherever found	E	\$8,500	\$732,797,282
1292	Nightjar, Puerto Rican (<i>Caprimulgus noctitherus</i>) -Wherever found	E	\$8,500	\$732,805,782
1292	No common name (<i>Asplenium dielfalcatum</i>) -Wherever found	E	\$8,500	\$732,814,282
1292	No common name (<i>Chamaecrista glandulosa</i> var. <i>mirabilis</i>) -Wherever found	E	\$8,500	\$732,822,782
1292	No common name (<i>Kadua hauptensis</i>) -Wherever found	E	\$8,500	\$732,831,282
1292	No common name (<i>Labordia lorenciana</i>) -Wherever found	E	\$8,500	\$732,839,782
1292	No common name (<i>Sanicula mariversa</i>) -Wherever found	E	\$8,500	\$732,848,282
1292	No common name (<i>Sanicula purpurea</i>) -Wherever found	E	\$8,500	\$732,856,782
1292	No common name (<i>Schiedea sarmentosa</i>) -Wherever found	E	\$8,500	\$732,865,282
1292	No common name (<i>Schoepfia arenaria</i>) -Wherever found	T	\$8,500	\$732,873,782
1292	No common name (<i>Stenogyne campanulata</i>) -Wherever found	E	\$8,500	\$732,882,282
1292	No common name (<i>Stenogyne kealiae</i>) -Wherever found	E	\$8,500	\$732,890,782
1292	No common name (<i>Viola helenae</i>) -Wherever found	E	\$8,500	\$732,899,282
1292	No common name (<i>Viola oahuensis</i>) -Wherever found	E	\$8,500	\$732,907,782
1292	picture-wing fly, Hawaiian (<i>Drosophila obatai</i>) -Wherever found	E	\$8,500	\$732,916,282
1292	picture-wing fly, Hawaiian (<i>Drosophila substenoptera</i>) -Wherever found	E	\$8,500	\$732,924,782
1292	Remya, Maui (<i>Remya mauiensis</i>) -Wherever found	E	\$8,500	\$732,933,282
1292	Wawae'iole (<i>Huperzia nutans</i>) -Wherever found	E	\$8,500	\$732,941,782
1329	Locoweed, Fassett's (<i>Oxytropis campestris</i> var. <i>chartacea</i>) -Wherever found	T	\$8,476	\$732,950,258
1330	Chub, Owens Tui (<i>Gila bicolor</i> ssp. <i>snyderi</i>) -Wherever found	E	\$8,400	\$732,958,658
1330	Fleabane, Zuni (<i>Erigeron rhizomatus</i>) -Wherever found	T	\$8,400	\$732,967,058
1330	Pitcher-plant, mountain sweet (<i>Sarracenia rubra</i> ssp. <i>jonesii</i>) -Wherever found	E	\$8,400	\$732,975,458
1333	Turtle, yellow-blotched map (<i>Graptemys flavimaculata</i>) -Wherever found	T	\$8,330	\$732,983,788
1334	Phacelia, island (<i>Phacelia insularis</i> ssp. <i>insularis</i>) -Wherever found	E	\$8,300	\$732,992,088
1335	Sunflower, San Mateo woolly (<i>Eriophyllum latilobum</i>) -Wherever found	E	\$8,200	\$733,000,288
1335	Thistle, fountain (<i>Cirsium fontinale</i> var. <i>fontinale</i>) -Wherever found	E	\$8,200	\$733,008,488

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1337	Orcutt grass, San Joaquin Valley (<i>Orcuttia inaequalis</i>) -Wherever found	T	\$8,147	\$733,016,635
1338	Aster, Ruth's golden (<i>Pityopsis ruthii</i>) -Wherever found	E	\$8,000	\$733,024,635
1338	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$8,000	\$733,032,635
1338	Boa, Mona (<i>Epicrates monensis monensis</i>) -Wherever found	T	\$8,000	\$733,040,635
1338	cactus, Tobusch fishhook (<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>) -Wherever found	T	\$8,000	\$733,048,635
1338	haha (<i>Cyanea mauiensis</i>) -Wherever found	E	\$8,000	\$733,056,635
1338	haha (<i>Cyanea mceldowneyi</i>) -Wherever found	E	\$8,000	\$733,064,635
1338	Ko'oko'olau (<i>Bidens conjuncta</i>) -Wherever found	E	\$8,000	\$733,072,635
1338	Manaca, palma de (<i>Calyptronoma rivalis</i>) -Wherever found	T	\$8,000	\$733,080,635
1338	No common name (<i>Agave eggersiana</i>) -Wherever found	E	\$8,000	\$733,088,635
1338	No common name (<i>Cyperus fauriei</i>) -Wherever found	E	\$8,000	\$733,096,635
1338	No common name (<i>Festuca molokaiensis</i>) -Wherever found	E	\$8,000	\$733,104,635
1338	No common name (<i>Leptocereus grantianus</i>) -Wherever found	E	\$8,000	\$733,112,635
1338	Panicgrass, Carter's (<i>Panicum fauriei</i> var. <i>carteri</i>) -Wherever found	E	\$8,000	\$733,120,635
1338	Penny-cress, Kneeland Prairie (<i>Thlaspi californicum</i>) -Wherever found	E	\$8,000	\$733,128,635
1338	Pussypaws, Mariposa (<i>Calyptridium pulchellum</i>) -Wherever found	T	\$8,000	\$733,136,635
1338	Reedgrass, Hillebrand's (<i>Calamagrostis hillebrandii</i>) -Wherever found	E	\$8,000	\$733,144,635
1338	Rocksnail, round (<i>Leptoxis ampla</i>) -Wherever found	T	\$8,000	\$733,152,635
1338	Salamander, Salado (<i>Eurycea chisholmensis</i>) -Wherever found	T	\$8,000	\$733,160,635
1338	Shiner, Cahaba (<i>Notropis cahabae</i>) -Wherever found	E	\$8,000	\$733,168,635
1357	Kidneyshell, southern (<i>Ptychobranhus jonesi</i>) -Wherever found	E	\$7,993	\$733,176,628
1358	Kangaroo rat, Morro Bay (<i>Dipodomys heermanni morroensis</i>) -Wherever found	E	\$7,900	\$733,184,528
1358	Wallflower, Ben Lomond (<i>Erysimum teretifolium</i>) -Wherever found	E	\$7,900	\$733,192,428
1360	Gilia, Monterey (<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>) -Wherever found	E	\$7,800	\$733,200,228
1361	Catspaw, white (pearlymussel) (<i>Epioblasma perobliqua</i>) -Wherever found	E	\$7,600	\$733,207,828
1362	`Akoko (<i>Euphorbia celastroides</i> var. <i>kaenana</i>) -Wherever found	E	\$7,500	\$733,215,328
1362	`Akoko (<i>Euphorbia haelealeana</i>) -Wherever found	E	\$7,500	\$733,222,828
1362	`Ohe`ohe (<i>Polyscias gymnocarpa</i>) -Wherever found	E	\$7,500	\$733,230,328
1362	Alani (<i>Melicope balloui</i>) -Wherever found	E	\$7,500	\$733,237,828
1362	Alani (<i>Melicope christophersenii</i>) -Wherever found	E	\$7,500	\$733,245,328
1362	Alani (<i>Melicope reflexa</i>) -Wherever found	E	\$7,500	\$733,252,828
1362	Aupaka (<i>Isodendron hosakae</i>) -Wherever found	E	\$7,500	\$733,260,328
1362	Awiwi (<i>Schenkia sebaeoides</i>) -Wherever found	E	\$7,500	\$733,267,828
1362	Ha`iwale (<i>Cyrtandra dentata</i>) -Wherever found	E	\$7,500	\$733,275,328

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1362	Ha'iwale (Cyrtandra oxybapha) -Wherever found	E	\$7,500	\$733,282,828
1362	haha (Cyanea calycina) -Wherever found	E	\$7,500	\$733,290,328
1362	haha (Cyanea longiflora) -Wherever found	E	\$7,500	\$733,297,828
1362	haha (Cyanea mannii) -Wherever found	E	\$7,500	\$733,305,328
1362	Hala pepe (Pleomele fernaldii) -Wherever found	E	\$7,500	\$733,312,828
1362	Holei (Ochrosia kilaueaensis) -Wherever found	E	\$7,500	\$733,320,328
1362	Hulumoa (Korthalsella degeneri) -Wherever found	E	\$7,500	\$733,327,828
1362	Ko'oko'olau (Bidens amplexans) -Wherever found	E	\$7,500	\$733,335,328
1362	Ko'oko'olau (Bidens wiebkei) -Wherever found	E	\$7,500	\$733,342,828
1362	Koki'o ke'oke'o (Hibiscus arnottianus ssp. immaculatus) -Wherever found	E	\$7,500	\$733,350,328
1362	Ma'oli'oli (Schiedea kealiae) -Wherever found	E	\$7,500	\$733,357,828
1362	Malacothrix, island (Malacothrix squalida) -Wherever found	E	\$7,500	\$733,365,328
1362	Na'ena'e (Dubautia herbastobatae) -Wherever found	E	\$7,500	\$733,372,828
1362	nehe (Lipochaeta lobata var. leptophylla) -Wherever found	E	\$7,500	\$733,380,328
1362	nehe (Lipochaeta venosa) -Wherever found	E	\$7,500	\$733,387,828
1362	No common name (Abutilon sandwicense) -Wherever found	E	\$7,500	\$733,395,328
1362	No common name (Gouania hillebrandii) -Wherever found	E	\$7,500	\$733,402,828
1362	No common name (Kadua degeneri) -Wherever found	E	\$7,500	\$733,410,328
1362	No common name (Kadua parvula) -Wherever found	E	\$7,500	\$733,417,828
1362	No common name (Lobelia niihauensis) -Wherever found	E	\$7,500	\$733,425,328
1362	No common name (Phyllostegia glabra var. lanaiensis) -Wherever found	E	\$7,500	\$733,432,828
1362	No common name (Platydesma cornuta var. decurrens) -Wherever found	E	\$7,500	\$733,440,328
1362	No common name (Schiedea lydgatei) -Wherever found	E	\$7,500	\$733,447,828
1362	No common name (Schiedea trinervis) -Wherever found	E	\$7,500	\$733,455,328
1362	No common name (Tetramolopium filiforme) -Wherever found	E	\$7,500	\$733,462,828
1362	No common name (Trematolobelia singularis) -Wherever found	E	\$7,500	\$733,470,328
1362	Sunfish, spring pygmy (Elassoma alabamae) -Wherever found	T	\$7,500	\$733,477,828
1362	Plum, scrub (Prunus geniculata) -	E	\$7,500	\$733,485,328
1399	Manzanita, Santa Rosa Island (Arctostaphylos confertiflora) -Wherever found	E	\$7,300	\$733,492,628
1399	Morning-glory, Stebbins' (Calystegia stebbinsii) -Wherever found	E	\$7,300	\$733,499,928
1401	Butterfly, Saint Francis' satyr (Neonympha mitchellii francisci) -Wherever found	E	\$7,100	\$733,507,028
1402	'Awiwi (Kadua cookiana) -Wherever found	E	\$7,000	\$733,514,028
1402	Bladderpod, white (Physaria pallida) -Wherever found	E	\$7,000	\$733,521,028
1402	Boxwood, Vahl's (Buxus vahliae) -Wherever found	E	\$7,000	\$733,528,028
1402	Ha'iwale (Cyrtandra crenata) -Wherever found	E	\$7,000	\$733,535,028
1402	No common name (Myrcia paganii) -Wherever found	E	\$7,000	\$733,542,028
1402	White-eye, Bridled (Zosterops conspicillatus conspicillatus) -Wherever found	E	\$7,000	\$733,549,028

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1402	Snakeroot (<i>Eryngium cuneifolium</i>) -	E	\$7,000	\$733,556,028
1402	Prairie-clover, Florida (<i>Dalea carthagenensis floridana</i>) -	E	\$7,000	\$733,563,028
1402	Water-willow, Cooley's (<i>Justicia cooleyi</i>) -	E	\$7,000	\$733,570,028
1402	Mustard, Carter's (<i>Warea carteri</i>) -	E	\$7,000	\$733,577,028
1402	Wireweed (<i>Polygonella basiramia</i>) -	E	\$7,000	\$733,584,028
1413	Pupfish, Ash Meadows Amargosa (<i>Cyprinodon nevadensis mionectes</i>) -Wherever found	E	\$6,900	\$733,590,928
1414	Cactus, Fickeisen plains (<i>Pediocactus peeblesianus</i> ssp. <i>fickeiseniae</i>) -Wherever found	E	\$6,700	\$733,597,628
1415	Sunburst, Hartweg's golden (<i>Pseudobahia bahiifolia</i>) -Wherever found	E	\$6,547	\$733,604,175
1416	`Akoko (<i>Euphorbia kuwaleana</i>) -Wherever found	E	\$6,500	\$733,610,675
1416	`Akoko (<i>Euphorbia rockii</i>) -Wherever found	E	\$6,500	\$733,617,175
1416	Bedstraw, El Dorado (<i>Galium californicum</i> ssp. <i>sierrae</i>) -Wherever found	E	\$6,500	\$733,623,675
1416	Ceanothus, Pine Hill (<i>Ceanothus roderickii</i>) -Wherever found	E	\$6,500	\$733,630,175
1416	Chupacallos (<i>Pleodendron macranthum</i>) -Wherever found	E	\$6,500	\$733,636,675
1416	Cobana negra (<i>Stahlia monosperma</i>) -Wherever found	T	\$6,500	\$733,643,175
1416	Combshell, southern (<i>Epioblasma penita</i>) -Wherever found	E	\$6,500	\$733,649,675
1416	Coqui, golden (<i>Eleutherodactylus jasper</i>) -Wherever found	T	\$6,500	\$733,656,175
1416	Cypress, Gowen (<i>Cupressus goveniana</i> ssp. <i>goveniana</i>) -Wherever found	T	\$6,500	\$733,662,675
1416	Goetzea, beautiful (<i>Goetzea elegans</i>) -Wherever found	E	\$6,500	\$733,669,175
1416	Ha`iwale (<i>Cyrtandra sessilis</i>) -Wherever found	E	\$6,500	\$733,675,675
1416	Ha`iwale (<i>Cyrtandra subumbellata</i>) -Wherever found	E	\$6,500	\$733,682,175
1416	haha (<i>Cyanea copelandii</i> ssp. <i>copelandii</i>) -Wherever found	E	\$6,500	\$733,688,675
1416	Higuero de sierra (<i>Crescentia portoricensis</i>) -Wherever found	E	\$6,500	\$733,695,175
1416	Kolea (<i>Myrsine juddii</i>) -Wherever found	E	\$6,500	\$733,701,675
1416	Kopiko (<i>Psychotria hobdyi</i>) -Wherever found	E	\$6,500	\$733,708,175
1416	No common name (<i>Asplenium unisorum</i>) -Wherever found	E	\$6,500	\$733,714,675
1416	No common name (<i>Doryopteris takeuchii</i>) -Wherever found	E	\$6,500	\$733,721,175
1416	No common name (<i>Gonocalyx concolor</i>) -Wherever found	E	\$6,500	\$733,727,675
1416	No common name (<i>Lobelia koolauensis</i>) -Wherever found	E	\$6,500	\$733,734,175
1416	No common name (<i>Lysimachia filifolia</i>) -Wherever found	E	\$6,500	\$733,740,675
1416	No common name (<i>Phyllostegia renovans</i>) -Wherever found	E	\$6,500	\$733,747,175
1416	No common name (<i>Schiedea spergulina</i> var. <i>leiopoda</i>) -Wherever found	E	\$6,500	\$733,753,675
1416	Shrimp, Alabama cave (<i>Palaemonias alabamiae</i>) -Wherever found	E	\$6,500	\$733,760,175
1416	Vole, Florida salt marsh (<i>Microtus pennsylvanicus dukecampbelli</i>) -Wherever found	E	\$6,500	\$733,766,675
1416	Blazingstar, scrub (<i>Liatris ohlingerae</i>) -	E	\$6,500	\$733,773,175
1416	Spurge, deltoid (<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>) -	E	\$6,500	\$733,779,675
1416	Fringe-tree, pygmy (<i>Chionanthus pygmaeus</i>) -	E	\$6,500	\$733,786,175

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1416	Whitlow-wort, papery (<i>Paronychia chartacea</i>) -	T	\$6,500	\$733,792,675
1445	Grass, Solano (<i>Tuctoria mucronata</i>) -Wherever found	E	\$6,302	\$733,798,977
1446	Gilia, Hoffmann's slender-flowered (<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>) -Wherever found	E	\$6,300	\$733,805,277
1446	Mallow, Kern (<i>Eremalche kernensis</i>) -Wherever found	E	\$6,300	\$733,811,577
1446	Pupfish, Warm Springs (<i>Cyprinodon nevadensis pectoralis</i>) -Wherever found	E	\$6,300	\$733,817,877
1449	Checker-mallow, Kenwood Marsh (<i>Sidalcea oregana</i> ssp. <i>valida</i>) -Wherever found	E	\$6,200	\$733,824,077
1450	Malacothrix, Santa Cruz Island (<i>Malacothrix indecora</i>) -Wherever found	E	\$6,100	\$733,830,177
1451	Bluegrass, Napa (<i>Poa napensis</i>) -Wherever found	E	\$6,000	\$733,836,177
1451	Cory cactus, bunched (<i>Coryphantha ramillosa</i>) -Wherever found	T	\$6,000	\$733,842,177
1451	dace, Laurel (<i>Chrosomus saylori</i>) -Wherever found	E	\$6,000	\$733,848,177
1451	Erubia (<i>Solanum drymophilum</i>) -Wherever found	E	\$6,000	\$733,854,177
1451	No common name (<i>Auerodendron pauciflorum</i>) -Wherever found	E	\$6,000	\$733,860,177
1451	No common name (<i>Cranichis ricartii</i>) -Wherever found	E	\$6,000	\$733,866,177
1451	No common name (<i>Polystichum calderonense</i>) -Wherever found	E	\$6,000	\$733,872,177
1451	No common name (<i>Tectaria estremerana</i>) -Wherever found	E	\$6,000	\$733,878,177
1451	Palo de ramon (<i>Banara vanderbiltii</i>) -Wherever found	E	\$6,000	\$733,884,177
1451	Poppy-mallow, Texas (<i>Callirhoe scabriuscula</i>) -Wherever found	E	\$6,000	\$733,890,177
1451	Uvillo (<i>Eugenia haematocarpa</i>) -Wherever found	E	\$6,000	\$733,896,177
1462	Trout, Paiute cutthroat (<i>Oncorhynchus clarkii seleniris</i>) -Wherever found	T	\$5,800	\$733,901,977
1463	Grasshopper, Zayante band-winged (<i>Trimerotropis infantilis</i>) -Wherever found	E	\$5,600	\$733,907,577
1464	Chumbo, Higo (<i>Harrisia portoricensis</i>) -Wherever found	T	\$5,500	\$733,913,077
1464	Crow, white-necked (<i>Corvus leucognaphalus</i>) -Wherever found	E	\$5,500	\$733,918,577
1464	No common name (<i>Elaphoglossum serpens</i>) -Wherever found	E	\$5,500	\$733,924,077
1464	Pebblesnail, flat (<i>Lepyrium showalteri</i>) -Wherever found	E	\$5,500	\$733,929,577
1464	Rattleweed, hairy (<i>Baptisia arachnifera</i>) -Wherever found	E	\$5,500	\$733,935,077
1469	Fringepod, Santa Cruz Island (<i>Thysanocarpus conchuliferus</i>) -Wherever found	E	\$5,300	\$733,940,377
1470	mallow, Gierisch (<i>Sphaeralcea gierischii</i>) -Wherever found	E	\$5,249	\$733,945,626
1471	Dace, Independence Valley speckled (<i>Rhinichthys osculus lethoporus</i>) -Wherever found	E	\$5,200	\$733,950,826
1472	Moth, Kern primrose sphinx (<i>Euproserpinus euterpe</i>) -Wherever found	T	\$5,100	\$733,955,926
1473	Bush-mallow, San Clemente Island (<i>Malacothamnus clementinus</i>) -Wherever found	E	\$5,000	\$733,960,926
1473	Larkspur, San Clemente Island (<i>Delphinium variegatum</i> ssp. <i>kinkiense</i>) -Wherever found	E	\$5,000	\$733,965,926
1473	Lioplax, cylindrical (snail) (<i>Lioplax cyclostomaformis</i>) -Wherever found	E	\$5,000	\$733,970,926
1473	lotus, San Clemente Island (<i>Acemispion dendroideus</i> var. <i>traskiae</i>) -Wherever found	T	\$5,000	\$733,975,926
1473	No common name (<i>Aristida chaseae</i>) -Wherever found	E	\$5,000	\$733,980,926
1473	No common name (<i>Daphnopsis helleriana</i>) -Wherever found	E	\$5,000	\$733,985,926

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1473	No common name (<i>Mitracarpus maxwelliae</i>) -Wherever found	E	\$5,000	\$733,990,926
1473	No common name (<i>Tabernaemontana rotensis</i>) -Wherever found	T	\$5,000	\$733,995,926
1473	Paintbrush, San Clemente Island (<i>Castilleja grisea</i>) -Wherever found	T	\$5,000	\$734,000,926
1473	Pelos del diablo (<i>Aristida portoricensis</i>) -Wherever found	E	\$5,000	\$734,005,926
1473	Quillwort, black spored (<i>Isoetes melanospora</i>) -Wherever found	E	\$5,000	\$734,010,926
1473	Shiner, sharpnose (<i>Notropis oxyrhynchus</i>) -Wherever found	E	\$5,000	\$734,015,926
1473	Shiner, smalleye (<i>Notropis buccula</i>) -Wherever found	E	\$5,000	\$734,020,926
1473	stonecrop, Bartram's (<i>Graptopetalum bartramii</i>) -Wherever found	T	\$5,000	\$734,025,926
1473	Beargrass, Britton's (<i>Nolina brittoniana</i>) -	E	\$5,000	\$734,030,926
1473	Pawpaw, Rugel's (<i>Deeringothamnus rugelii</i>) -	E	\$5,000	\$734,035,926
1473	Cactus, Florida semaphore (<i>Consolea corallicola</i>) -	E	\$5,000	\$734,040,926
1473	Bonamia, Florida (<i>Bonamia grandiflora</i>) -	T	\$5,000	\$734,045,926
1473	Polygala, Lewton's (<i>Polygala lewtonii</i>) -	E	\$5,000	\$734,050,926
1492	Addax (<i>Addax nasomaculatus</i>) -Wherever found	E	\$4,900	\$734,055,826
1493	Fairy shrimp, Riverside (<i>Streptocephalus woottoni</i>) -Wherever found	E	\$4,800	\$734,060,626
1494	Shrimp, California freshwater (<i>Syncaris pacifica</i>) -Wherever found	E	\$4,600	\$734,065,226
1495	Amphianthus, little (<i>Amphianthus pusillus</i>) -Wherever found	T	\$4,500	\$734,069,726
1495	Clubshell, Canoe Creek (<i>Pleurobema atearni</i>) -Wherever found	E	\$4,500	\$734,074,226
1495	Pigtoe, dark (<i>Pleurobema furvum</i>) -Wherever found	E	\$4,500	\$734,078,726
1495	Bacora, marron (<i>Solanum conocarpum</i>) -	E	\$4,500	\$734,083,226
1495	Harebells, Avon Park (<i>Crotalaria avonensis</i>) -	E	\$4,500	\$734,087,726
1500	Buckwheat, cushenbury (<i>Eriogonum ovalifolium</i> var. <i>vineum</i>) -Wherever found	E	\$4,300	\$734,092,026
1500	Bush-mallow, Santa Cruz Island (<i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i>) -Wherever found	E	\$4,300	\$734,096,326
1500	Manzanita, Presidio (<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>) -Wherever found	E	\$4,300	\$734,100,626
1500	Milk-vetch, Cushenbury (<i>Astragalus albens</i>) -Wherever found	E	\$4,300	\$734,104,926
1504	Alopecurus, Sonoma (<i>Alopecurus aequalis</i> var. <i>sonomensis</i>) -Wherever found	E	\$4,250	\$734,109,176
1505	Cactus, Lloyd's Mariposa (<i>Sclerocactus mariposensis</i>) -Wherever found	T	\$4,200	\$734,113,376
1506	eryngo, Arizona (<i>Eryngium sparganophyllum</i>) -	E	\$4,120	\$734,117,496
1507	Deer, Eld's brow-antlered (<i>Cervus eldi</i>) -Wherever found	E	\$4,000	\$734,121,496
1507	Leopard (<i>Panthera pardus</i>) -In Africa, in the wild, south of, and including, the following countries: Gabon, Congo, Zaire, Uganda, Kenya	T	\$4,000	\$734,125,496
1507	No common name (<i>Cordia bellonis</i>) -Wherever found	E	\$4,000	\$734,129,496
1507	No common name (<i>Lyonia truncata</i> var. <i>proctorii</i>) -Wherever found	E	\$4,000	\$734,133,496
1507	No common name (<i>Vernonia proctorii</i>) -Wherever found	E	\$4,000	\$734,137,496
1507	Quillwort, mat-forming (<i>Isoetes tegetiformans</i>) -Wherever found	E	\$4,000	\$734,141,496
1507	Rocksnailed, plicate (<i>Leptoxis plicata</i>) -Wherever found	E	\$4,000	\$734,145,496

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1507	Salamander, Georgetown (<i>Eurycea naufragia</i>) -Wherever found	T	\$4,000	\$734,149,496
1507	Salamander, Jollyville Plateau (<i>Eurycea tonkawae</i>) -Wherever found	T	\$4,000	\$734,153,496
1507	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>) -Central Valley spring-run ESU XN - See 50 CFR 223.102	EXPN	\$4,000	\$734,157,496
1507	Spider, Tooth Cave (<i>Tayshaneta myopica</i>) -Wherever found	E	\$4,000	\$734,161,496
1507	Trillium, persistent (<i>Trillium persistens</i>) -Wherever found	E	\$4,000	\$734,165,496
1507	Walnut (=Nogal), West Indian (<i>Juglans jamaicensis</i>) -Wherever found	E	\$4,000	\$734,169,496
1507	Rosemary, short-leaved (<i>Conradina brevifolia</i>) -	E	\$4,000	\$734,173,496
1507	Pigeon wings (<i>Clitoria fragrans</i>) -	T	\$4,000	\$734,177,496
1507	Mint, scrub (<i>Dicerandra frutescens</i>) -	E	\$4,000	\$734,181,496
1507	Mint, Lakela's (<i>Dicerandra immaculata</i>) -	E	\$4,000	\$734,185,496
1507	Hypericum, highlands scrub (<i>Hypericum cumulicola</i>) -	E	\$4,000	\$734,189,496
1525	Darter, yellowcheek (<i>Etheostoma moorei</i>) -Wherever found	E	\$3,800	\$734,193,296
1525	Rose-mallow, Neches River (<i>Hibiscus dasycalyx</i>) -Wherever found	T	\$3,800	\$734,197,096
1527	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$3,500	\$734,200,596
1527	Butterfly, El Segundo blue (<i>Euphilotes battoides allyni</i>) -Wherever found	E	\$3,500	\$734,204,096
1527	Fairy shrimp, San Diego (<i>Branchinecta sandiegonensis</i>) -Wherever found	E	\$3,500	\$734,207,596
1527	Fern, Elfin tree (<i>Cyathea dryopteroides</i>) -Wherever found	E	\$3,500	\$734,211,096
1527	Peperomia, Wheeler's (<i>Peperomia wheeleri</i>) -Wherever found	E	\$3,500	\$734,214,596
1527	Prickly-apple, fragrant (<i>Cereus eriophorus</i> var. <i>fragrans</i>) -	E	\$3,500	\$734,218,096
1533	Cladonia, Florida perforate (<i>Cladonia perforata</i>) -	E	\$3,495	\$734,221,591
1534	Vervain, Red Hills (<i>Verbena californica</i>) -Wherever found	T	\$3,400	\$734,224,991
1535	Bluet, Roan Mountain (<i>Hedyotis purpurea</i> var. <i>montana</i>) -Wherever found	E	\$3,300	\$734,228,291
1536	Blue-star, Kearney's (<i>Amsonia kearneyana</i>) -Wherever found	E	\$3,200	\$734,231,491
1536	Lupine, clover (Tidestrom's) (<i>Lupinus tidestromii</i>) -Wherever found	E	\$3,200	\$734,234,691
1536	Thistle, Chorro Creek bog (<i>Cirsium fontinale</i> var. <i>obispoense</i>) -Wherever found	E	\$3,200	\$734,237,891
1539	macaw, Scarlet (<i>Ara macao</i> ssp. <i>cyanopterus</i>) -Wherever found	E	\$3,132	\$734,241,023
1540	Brodiaea, Chinese Camp (<i>Brodiaea pallida</i>) -Wherever found	T	\$3,100	\$734,244,123
1541	Anole, Culebra Island giant (<i>Anolis roosevelti</i>) -Wherever found	E	\$3,000	\$734,247,123
1541	Beetle, Comal Springs dryopid (<i>Stygoparnus comalensis</i>) -Wherever found	E	\$3,000	\$734,250,123
1541	Blossom, green (pearlymussel) (<i>Epioblasma torulosa gubernaculum</i>) -Wherever found	E	\$3,000	\$734,253,123
1541	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) -Wherever found; Except where listed as Experimental Populations	E	\$3,000	\$734,256,123

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1541	Cactus, Chisos Mountain hedgehog (<i>Echinocereus chisoensis</i> var. <i>chisoensis</i>) -Wherever found	T	\$3,000	\$734,259,123
1541	Capa rosa (<i>Callicarpa ampla</i>) -Wherever found	E	\$3,000	\$734,262,123
1541	Chimpanzee, pygmy (<i>Pan paniscus</i>) -Wherever found	E	\$3,000	\$734,265,123
1541	Gambusia, Pecos (<i>Gambusia nobilis</i>) -Wherever found	E	\$3,000	\$734,268,123
1541	Manzanita, Ione (<i>Arctostaphylos myrtifolia</i>) -Wherever found	T	\$3,000	\$734,271,123
1541	Mucket, orangenacre (<i>Hamiota perovalis</i>) -Wherever found	T	\$3,000	\$734,274,123
1541	No common name (<i>Gesneria pauciflora</i>) -Wherever found	T	\$3,000	\$734,277,123
1541	No common name (<i>Thelypteris inabonensis</i>) -Wherever found	E	\$3,000	\$734,280,123
1541	No common name (<i>Thelypteris verecunda</i>) -Wherever found	E	\$3,000	\$734,283,123
1541	No common name (<i>Thelypteris yaucoensis</i>) -Wherever found	E	\$3,000	\$734,286,123
1541	Palo colorado (<i>Ternstroemia luquillensis</i>) -Wherever found	E	\$3,000	\$734,289,123
1541	Palo de jazmin (<i>Styrax portoricensis</i>) -Wherever found	E	\$3,000	\$734,292,123
1541	Palo de nigua (<i>Cornutia obovata</i>) -Wherever found	E	\$3,000	\$734,295,123
1541	petrel, Bermuda (<i>Pterodroma cahow</i>) -Wherever found	E	\$3,000	\$734,298,123
1541	Trout, Little Kern golden (<i>Oncorhynchus aguabonita whitei</i>) -Wherever found	T	\$3,000	\$734,301,123
1541	Warbler (=wood), Bachman's (<i>Vermivora bachmanii</i>) -Wherever found	E	\$3,000	\$734,304,123
1541	Warbler, elfin-woods (<i>Setophaga angelae</i>) -Wherever found	T	\$3,000	\$734,307,123
1541	Water-plantain, Kral's (<i>Sagittaria secundifolia</i>) -Wherever found	T	\$3,000	\$734,310,123
1563	Clover, Monterey (<i>Trifolium trichocalyx</i>) -Wherever found	E	\$2,900	\$734,313,023
1563	mountainbalm, Indian Knob (<i>Eriodictyon altissimum</i>) -Wherever found	E	\$2,900	\$734,315,923
1565	Barberry, island (<i>Berberis pinnata</i> ssp. <i>insularis</i>) -Wherever found	E	\$2,800	\$734,318,723
1566	Cactus, Brady pincushion (<i>Pediocactus bradyi</i>) -Wherever found	E	\$2,700	\$734,321,423
1566	Skipper, Pawnee montane (<i>Hesperia leonardus montana</i>) -Wherever found	T	\$2,700	\$734,324,123
1568	Sloth, Brazilian three-toed (<i>Bradypus torquatus</i>) -Wherever found	E	\$2,666	\$734,326,789
1569	Chub, spotfin (<i>Erimonax monachus</i>) -U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$2,500	\$734,329,289
1569	Darter, boulder (<i>Etheostoma wapiti</i>) -Shoal Creek (from Shoal Creek mile 41.7 (66.7 km)) at the mouth of Long Branch, Lawrence County, TN, downstream to the backwaters of Wilson Reservoir (Shoal Creek mile 14 (22 km)) at Goose Shoals, Lauderdale County, AL, including the lower 5 miles (8 km) of all tributaries that enter this reach.	EXPN	\$2,500	\$734,331,789
1569	Holly, Cook's (<i>Ilex cookii</i>) -Wherever found	E	\$2,500	\$734,334,289
1569	Minnow, Rio Grande Silvery (<i>Hybognathus amarus</i>) -Wherever found, except where listed as an experimental population	E	\$2,500	\$734,336,789
1569	No common name (<i>Ilex sintenisii</i>) -Wherever found	E	\$2,500	\$734,339,289
1569	No common name (<i>Ternstroemia subsessilis</i>) -Wherever found	E	\$2,500	\$734,341,789

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1569	Pigtoe, flat (<i>Pleurobema marshalli</i>) -Wherever found	E	\$2,500	\$734,344,289
1569	Rush-rose, island (<i>Helianthemum greenei</i>) -Wherever found	T	\$2,500	\$734,346,789
1569	Sparrow, San Clemente sage (<i>Amphispiza belli clementeae</i>) -Wherever found	T	\$2,500	\$734,349,289
1569	Spineflower, Sonoma (<i>Chorizanthe valida</i>) -Wherever found	E	\$2,500	\$734,351,789
1569	Stirrupshell (<i>Quadrula stapes</i>) -Wherever found	E	\$2,500	\$734,354,289
1569	Stonefly, meltwater lednian (<i>Lednia tumana</i>) -Wherever found	T	\$2,500	\$734,356,789
1569	Lead-plant, Crenulate (<i>Amorpha crenulata</i>) -	E	\$2,500	\$734,359,289
1569	Mint, Garrett's (<i>Dicerandra christmanii</i>) -	E	\$2,500	\$734,361,789
1569	Pawpaw, beautiful (<i>Deeringothamnus pulchellus</i>) -	E	\$2,500	\$734,364,289
1569	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) -	T	\$2,500	\$734,366,789
1569	Polygala, tiny (<i>Polygala smallii</i>) -	E	\$2,500	\$734,369,289
1586	Cactus, Knowlton's (<i>Pediocactus knowltonii</i>) -Wherever found	E	\$2,475	\$734,371,764
1587	Thoroughwort, Cape Sable (<i>Chromolaena frustrata</i>) -	E	\$2,400	\$734,374,164
1588	Pigtoe, shiny (<i>Fusconaia cor</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$2,300	\$734,376,464
1589	Silverside, Waccamaw (<i>Menidia extensa</i>) -Wherever found	T	\$2,200	\$734,378,664
1589	Wire-lettuce, Malheur (<i>Stephanomeria malheurensis</i>) -Wherever found	E	\$2,200	\$734,380,864
1591	Amphipod, Noel's (<i>Gammarus desperatus</i>) -Wherever found	E	\$2,000	\$734,382,864
1591	Butterfly, Palos Verdes blue (<i>Glaucopsyche lygdamus palosverdesensis</i>) -Wherever found	E	\$2,000	\$734,384,864
1591	Gambusia, Big Bend (<i>Gambusia gaigei</i>) -Wherever found	E	\$2,000	\$734,386,864
1591	Snail, Pecos assiminea (<i>Assiminea pecos</i>) -Wherever found	E	\$2,000	\$734,388,864
1591	Snail, Stock Island tree (<i>Orthalicus reses</i> (not incl. <i>nesodryas</i>)) -Wherever found	T	\$2,000	\$734,390,864
1591	Springsnail, Koster's (<i>Juturnia kosteri</i>) -Wherever found	E	\$2,000	\$734,392,864
1591	Springsnail, Roswell (<i>Pyrgulopsis roswellensis</i>) -Wherever found	E	\$2,000	\$734,394,864
1591	Woodrat, riparian (=San Joaquin Valley) (<i>Neotoma fuscipes riparia</i>) -Wherever found	E	\$2,000	\$734,396,864
1591	Rosemary, Etonia (<i>Conradina etonia</i>) -	E	\$2,000	\$734,398,864
1591	Spurge, Garber's (<i>Chamaesyce garberi</i>) -	T	\$2,000	\$734,400,864
1601	Spinedace, Little Colorado (<i>Lepidomeda vittata</i>) -Wherever found	T	\$1,961	\$734,402,825
1602	Bladderpod, lyrata (<i>Lesquerella lyrata</i>) -Wherever found	T	\$1,900	\$734,404,725
1602	Irisette, white (<i>Sisyrinchium dichotomum</i>) -Wherever found	E	\$1,900	\$734,406,625
1602	Madtom, smoky (<i>Noturus baileyi</i>) -The Tellico River, between the backwaters of the Tellico Reservoir (approximately Tellico River mile 19 (30.4 kilometers) and Tellico River mile 33 (52.8 kilometers), near the Tellico Ranger Station, Monroe County, Tennessee.	EXPN	\$1,900	\$734,408,525

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1602	Rockcress, Georgia (<i>Arabis georgiana</i>) -Wherever found	T	\$1,900	\$734,410,425
1606	Cactus, Acuña (<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>) -Wherever found	E	\$1,850	\$734,412,275
1607	Pitcher-plant, Alabama canebreak (<i>Sarracenia rubra</i> ssp. <i>alabamensis</i>) -Wherever found	E	\$1,800	\$734,414,075
1608	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,415,775
1608	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,417,475
1608	Campion, fringed (<i>Silene polypetala</i>) -Wherever found	E	\$1,700	\$734,419,175
1608	Clubshell (<i>Pleurobema clava</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,420,875
1608	Monkeyface, Cumberland (pearlymussel) (<i>Theliderma intermedia</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,422,575
1608	Mussel, oyster (<i>Epioblasma capsaeformis</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,424,275
1608	Pearlymussel, birdwing (<i>Lemiox rimosus</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,425,975
1608	Pearlymussel, cracking (<i>Hemistena lata</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,427,675
1608	Pearlymussel, cracking (<i>Hemistena lata</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$1,700	\$734,429,375

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1608	Pearlymussel, dromedary (<i>Dromus dromas</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,431,075
1608	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,432,775
1608	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata</i>) -U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	EXPN	\$1,700	\$734,434,475
1608	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) -Central and Southwest Atlantic DPS	T	\$1,700	\$734,436,175
1608	Whale, gray (<i>Eschrichtius robustus</i>) -Western North Pacific DPS See 50 CFR 224.101	E	\$1,700	\$734,437,875
1622	Potentilla, Hickman's (<i>Potentilla hickmanii</i>) -Wherever found	E	\$1,600	\$734,439,475
1623	Pawpaw, four-petal (<i>Asimina tetramera</i>) -	E	\$1,595	\$734,441,070
1624	Bird's-beak, Pennell's (<i>Cordylanthus tenuis</i> ssp. <i>capillaris</i>) -Wherever found	E	\$1,500	\$734,442,570
1624	Buckwheat, Ione (incl. Irish Hill) (<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i>)) -Wherever found	E	\$1,500	\$734,444,070
1624	Clubshell, black (<i>Pleurobema curtum</i>) -Wherever found	E	\$1,500	\$734,445,570
1624	Jewelflower, Tiburon (<i>Streptanthus niger</i>) -Wherever found	E	\$1,500	\$734,447,070
1624	Mariposa lily, Tiburon (<i>Calochortus tiburonensis</i>) -Wherever found	T	\$1,500	\$734,448,570
1624	Milk-vetch, Clara Hunt's (<i>Astragalus clarianus</i>) -Wherever found	E	\$1,500	\$734,450,070
1624	Orcutt grass, California (<i>Orcuttia californica</i>) -Wherever found	E	\$1,500	\$734,451,570
1631	Navarretia, spreading (<i>Navarretia fossalis</i>) -Wherever found	T	\$1,400	\$734,452,970
1631	No common name (<i>Schiedea attenuata</i>) -Wherever found	E	\$1,400	\$734,454,370
1633	Barberry, Nevin's (<i>Berberis nevinii</i>) -Wherever found	E	\$1,300	\$734,455,670
1633	Dudleya, Santa Monica Mountains (<i>Dudleya cymosa</i> ssp. <i>ovatifolia</i>) -Wherever found	T	\$1,300	\$734,456,970
1635	Butterwort, Godfrey's (<i>Pinguicula ionantha</i>) -	T	\$1,230	\$734,458,200
1636	gartersnake, narrow-headed (<i>Thamnophis rufipunctatus</i>) -Wherever found	T	\$1,184	\$734,459,384
1637	Shrimp, Squirrel Chimney Cave (<i>Palaemonetes cummingi</i>) -Wherever found	T	\$1,100	\$734,460,484
1637	Pinkroot, gentian (<i>Spigelia gentianoides</i>) -	E	\$1,100	\$734,461,584
1639	Cat, black-footed (<i>Felis nigripes</i>) -Wherever found	E	\$1,000	\$734,462,584
1639	Chub, Chihuahua (<i>Gila nigrescens</i>) -Wherever found	T	\$1,000	\$734,463,584
1639	Isopod, Socorro (<i>Thermosphaeroma thermophilus</i>) -Wherever found	E	\$1,000	\$734,464,584
1639	Macaw, military (<i>Ara militaris</i>) -Wherever found	E	\$1,000	\$734,465,584

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1639	Marstonia, royal (snail) (<i>Marstonia ogmorhappe</i>) -Wherever found	E	\$1,000	\$734,466,584
1639	Starling, Rothschild's (mya) (<i>Leucopsar rothschildi</i>) -Wherever found	E	\$1,000	\$734,467,584
1639	Tortoise, Madagascar radiated (<i>Geochelone radiata</i>) -Wherever found	E	\$1,000	\$734,468,584
1646	Rattlesnake, New Mexican ridge-nosed (<i>Crotalus willardi obscurus</i>) -Wherever found	T	\$903	\$734,469,487
1647	Milkpea, Small's (<i>Galactia smallii</i>) -	E	\$900	\$734,470,387
1647	Torreya, Florida (<i>Torreya taxifolia</i>) -	E	\$900	\$734,471,287
1647	Mint, longspurred (<i>Dicerandra cornutissima</i>) -	E	\$900	\$734,472,187
1647	Bellflower, Brooksville (<i>Campanula robinsiae</i>) -	E	\$900	\$734,473,087
1647	Jacquemontia, beach (<i>Jacquemontia reclinata</i>) -	E	\$900	\$734,473,987
1647	Prickly-apple, aboriginal (<i>Harrisia</i> (=Cereus) aboriginum (=gracilis)) -	E	\$900	\$734,474,887
1653	Ambrosia, San Diego (<i>Ambrosia pumila</i>) -Wherever found	E	\$700	\$734,475,587
1653	Chub, slender (<i>Erimystax cahni</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(s)(1)(i))	EXPN	\$700	\$734,476,287
1653	Chub, spotfin (<i>Erimonax monachus</i>) -U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$700	\$734,476,987
1653	Chub, spotfin (<i>Erimonax monachus</i>) -U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$700	\$734,477,687
1653	Darter, duskytail (<i>Etheostoma percnurum</i>) -The Tellico River, between the backwaters of the Tellico Reservoir (approximately Tellico River mile 19 (30.4 kilometers) and Tellico River mile 33 (52.8 kilometers), near the Tellico Ranger Station, Monroe County, Tennessee.	EXPN	\$700	\$734,478,387
1653	Darter, duskytail (<i>Etheostoma percnurum</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(q)(1)(ii))	EXPN	\$700	\$734,479,087
1653	Madtom, pygmy (<i>Noturus stanauli</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))	EXPN	\$700	\$734,479,787
1653	Madtom, yellowfin (<i>Noturus flavipinnis</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$700	\$734,480,487
1653	penguin, African (<i>Spheniscus demersus</i>) -Wherever found	E	\$700	\$734,481,187
1662	Wartyback, white (pearlymussel) (<i>Plethobasus cicatricosus</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$635	\$734,481,822
1663	Goldenrod, Blue Ridge (<i>Solidago spithamea</i>) -Wherever found	T	\$600	\$734,482,422
1663	Ibis, northern bald (<i>Geronticus eremita</i>) -Wherever found	E	\$600	\$734,483,022
1663	Manzanita, pallid (<i>Arctostaphylos pallida</i>) -Wherever found	T	\$600	\$734,483,622
1663	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) -U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$600	\$734,484,222
1667	Tarantula, Sri Lankan Ornamental (<i>Poecilotheria fasciata</i>) -Wherever found	E	\$521	\$734,484,743
1668	Deer, Barbary (<i>Cervus elaphus barbarus</i>) -Wherever found	E	\$500	\$734,485,243

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1668	Fly, Delhi Sands flower-loving (<i>Rhaphiomidas terminatus abdominalis</i>) -Wherever found	E	\$500	\$734,485,743
1668	Gazelle, Mhorh (<i>Gazella dama mhorh</i>) -Wherever found	E	\$500	\$734,486,243
1668	Macaque, lion-tailed (<i>Macaca silenus</i>) -Wherever found	E	\$500	\$734,486,743
1672	Bittercress, small-anthered (<i>Cardamine micranthera</i>) -Wherever found	E	\$300	\$734,487,043
1672	Button-celery, San Diego (<i>Eryngium aristulatum</i> var. <i>parishii</i>) -Wherever found	E	\$300	\$734,487,343
1672	Mesa-mint, San Diego (<i>Pogogyne abramsii</i>) -Wherever found	E	\$300	\$734,487,643
1675	Cactus, Arizona hedgehog (<i>Echinocereus arizonicus</i> ssp. <i>arizonicus</i>) -Wherever found	E	\$200	\$734,487,843
1675	Macaque, Japanese (<i>Macaca fuscata</i>) -Wherever found	T	\$200	\$734,488,043
1675	Milk-vetch, Sentry (<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>) -Wherever found	E	\$200	\$734,488,243
1675	Sucker, Zuni bluehead (<i>Catostomus discobolus yarrowi</i>) -Wherever found	E	\$200	\$734,488,443
1675	Whale, humpback (<i>Megaptera novaeangliae</i>) -Cape Verde Island/Northwest Africa	E	\$200	\$734,488,643
1680	Fern, Alabama streak-sorus (<i>Thelypteris pilosa</i> var. <i>alabamensis</i>) -Wherever found	T	\$100	\$734,488,743
1680	Lizard, Coachella Valley fringe-toed (<i>Uma inornata</i>) -Wherever found	T	\$100	\$734,488,843
1680	Paintbrush, ash-grey (<i>Castilleja cinerea</i>) -Wherever found	T	\$100	\$734,488,943
1680	Taraxacum, California (<i>Taraxacum californicum</i>) -Wherever found	E	\$100	\$734,489,043
1680	No Common Name (<i>Fusconaia subrotunda</i>) -	T	\$100	\$734,489,143
1680	No Common Name (<i>Obovaria subrotunda</i>) -	T	\$100	\$734,489,243
Total Ranked Species Expenditures				\$734,489,243

**TABLE 3. FY 2022 REPORTED LAND ACQUISITION EXPENDITURES FOR
ENDANGERED AND THREATENED SPECIES**

Species are arranged in the same sequence as Table 1. These costs are not included in the expenditures displayed in Tables 1 and 2. Only those species that had land acquisition costs reported for FY 2022 are included here. Land rank is included to show the relative order of the species for which land expenditures are reported. Land acquisition costs for multi-species conservation activities are tabulated at the end of Table 3. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

TABLE 3. REPORTED LAND ACQUISITION EXPENDITURES FOR FY 2022 FOR ENDANGERED AND THREATENED SPECIES

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
Amphibians	216	Frog, dusky gopher (<i>Rana sevosa</i>) -Wherever found	E	\$0	\$0	\$0	\$4,123	\$4,123
	123	Salamander, California tiger (<i>Ambystoma californiense</i>) -U.S.A. (CA - Santa Barbara County)	E	\$3,611,197	\$0	\$3,611,197	\$0	\$3,611,197
	310	Salamander, Cheat Mountain (<i>Plethodon nettingi</i>) -Wherever found	T	\$62,500	\$0	\$62,500	\$0	\$62,500
	219	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) -Wherever found	T	\$313,344	\$0	\$313,344	\$42,229	\$355,573
	306	salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) -Wherever found	E	\$0	\$0	\$0	\$42,229	\$42,229
	365	Salamander, Santa Cruz long-toed (<i>Ambystoma macrodactylum croceum</i>) -Wherever found	E	\$3,700,000	\$0	\$3,700,000	\$0	\$3,700,000
	468	Toad, arroyo (=arroyo southwestern) (<i>Anaxyrus californicus</i>) -Wherever found	E	\$4,333,333	\$0	\$4,333,333	\$0	\$4,333,333
Amphibians Total				\$12,020,374	\$0	\$12,020,374	\$88,581	\$12,108,955
Birds	117	Caracara, Audubon's crested (<i>Polyborus plancus audubonii</i>) -U.S.A. (FL)	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	151	Eider, Steller's (<i>Polysticta stelleri</i>) -U.S. Alaska breeding population only	T	\$60,000	\$0	\$60,000	\$0	\$60,000
	523	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) -Wherever found, except where listed as an experimental population	E	\$4,101,935	\$0	\$4,101,935	\$0	\$4,101,935
	227	Gnatcatcher, coastal California (<i>Polioptila californica californica</i>) -Wherever found	T	\$5,247,234	\$0	\$5,247,234	\$0	\$5,247,234
	162	Horned lark, streaked (<i>Eremophila alpestris strigata</i>) -Wherever found	T	\$275,333	\$0	\$275,333	\$0	\$275,333
	164	Kite, Everglade snail (<i>Rostrhamus sociabilis plumbeus</i>) -Wherever found	E	\$0	\$0	\$0	\$5,491	\$5,491
	48	Murrelet, marbled (<i>Brachyramphus marmoratus</i>) - U.S.A. (CA, OR, WA)	T	\$74,045	\$0	\$74,045	\$0	\$74,045
	66	Plover, piping (<i>Charadrius melodus</i>) -[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$5,014,179	\$0	\$5,014,179	\$0	\$5,014,179

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	169	scrub-jay, Florida (Aphelocoma coerulescens) - Wherever found	T	\$1,818	\$0	\$1,818	\$42,754	\$44,572
	207	Sparrow, Florida grasshopper (Ammodramus savannarum floridanus) -Wherever found	E	\$0	\$0	\$0	\$93,454	\$93,454
	209	Stork, wood (Mycteria americana) -U.S.A. (AL, FL, GA, MS, NC, SC)	T	\$871,538	\$0	\$871,538	\$58,000	\$929,538
	296	Tern, Roseate (Sterna dougallii dougallii) -U.S.A. (Atlantic Coast south to NC), Canada (Newf., N.S, Que.), Bermuda	E	\$221,500	\$0	\$221,500	\$0	\$221,500
	63	Woodpecker, red-cockaded (Picoides borealis) - Wherever found	E	\$13,276,839	\$0	\$13,276,839	\$459,252	\$13,736,091
Birds Total				\$29,146,239	\$0	\$29,146,239	\$658,951	\$29,805,190
Clams	271	Banklimber, purple (mussel) (Elliptoideus sloatianus) - Wherever found	T	\$313,344	\$0	\$313,344	\$0	\$313,344
	537	heelsplitter, Inflated (Potamilus inflatus) -Wherever found	T	\$51,667	\$0	\$51,667	\$0	\$51,667
	267	Higgins eye (pearlymussel) (Lampsilis higginsii) - Wherever found	E	\$611,333	\$0	\$611,333	\$0	\$611,333
	619	Kidneyshell, triangular (Ptychobranhus greenii) - Wherever found	E	\$400,125	\$0	\$400,125	\$0	\$400,125
	256	Mucket, pink (pearlymussel) (Lampsilis abrupta) - Wherever found	E	\$1,055,167	\$0	\$1,055,167	\$0	\$1,055,167
	265	Mussel, sheepnose (Plethobasus cyphus) -Wherever found	E	\$47,333	\$0	\$47,333	\$0	\$47,333
	546	Pocketbook, fat (Potamilus capax) -Wherever found	E	\$1,055,167	\$0	\$1,055,167	\$0	\$1,055,167
	262	Spectaclecase (mussel) (Cumberlandia monodonta) - Wherever found	E	\$47,333	\$0	\$47,333	\$0	\$47,333
	168	Wedgemussel, dwarf (Alasmidonta heterodon) - Wherever found	E	\$128,720	\$0	\$128,720	\$0	\$128,720
Clams Total				\$3,710,189	\$0	\$3,710,189	\$0	\$3,710,189
Crustaceans	316	Crayfish, Panama City (Procambarus econfinae) - Wherever found	T	\$0	\$0	\$0	\$3,131	\$3,131
	301	Fairy shrimp, Conservancy (Branchinecta conservatio) - Wherever found	E	\$56,250	\$0	\$56,250	\$0	\$56,250
	137	Fairy shrimp, vernal pool (Branchinecta lynchi) - Wherever found	T	\$56,641	\$0	\$56,641	\$0	\$56,641

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	147	Tadpole shrimp, vernal pool (<i>Lepidurus packardii</i>) - Wherever found	E	\$56,641	\$0	\$56,641	\$0	\$56,641
Crustaceans Total				\$169,532	\$0	\$169,532	\$3,131	\$172,663
Fishes	715	Darter, goldline (<i>Percina aurolineata</i>) -Wherever found	T	\$400,125	\$0	\$400,125	\$0	\$400,125
	9	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Lower Columbia River ESU See 50 CFR 223.102	T	\$0	\$531,081	\$531,081	\$0	\$531,081
	8	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River fall-run ESU See 50 CFR 223.102	T	\$74,827	\$539,338	\$614,165	\$0	\$614,165
	3	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Snake River spring/summer-run ESU See 50 CFR 223.102	T	\$0	\$39,560	\$39,560	\$0	\$39,560
	5	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Columbia River spring-run ESU See 50 CFR 224.101	E	\$0	\$39,560	\$39,560	\$0	\$39,560
	14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) -Upper Willamette River ESU See 50 CFR 223.102	T	\$0	\$39,560	\$39,560	\$0	\$39,560
	17	Salmon, chum (<i>Oncorhynchus keta</i>) -Columbia River ESU See 50 CFR 223.102	T	\$0	\$530,314	\$530,314	\$0	\$530,314
	15	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Lower Columbia River ESU - See 50 CFR 223.102	T	\$0	\$531,081	\$531,081	\$0	\$531,081
	11	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU - See 50 CFR 224.101	E	\$0	\$39,560	\$39,560	\$0	\$39,560
	88	Sawfish, smalltooth (<i>Pristis pectinata</i>) -US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$0	\$0	\$0	\$3,131	\$3,131
	1338	Shiner, Cahaba (<i>Notropis cahabae</i>) -Wherever found	E	\$400,125	\$0	\$400,125	\$0	\$400,125
	36	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -California Central Valley DPS - See 50 CFR 223.102	T	\$349,769	\$0	\$349,769	\$0	\$349,769
	13	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) -Lower Columbia River DPS - See 50 CFR 223.102	T	\$0	\$531,081	\$531,081	\$0	\$531,081

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1	Steelhead (Oncorhynchus (=Salmo) mykiss) -Middle Columbia River DPS - See 50 CFR 223.102	T	\$0	\$39,560	\$39,560	\$0	\$39,560
	2	Steelhead (Oncorhynchus (=Salmo) mykiss) -Snake River Basin DPS - See 50 CFR 223.102	T	\$0	\$48,584	\$48,584	\$0	\$48,584
	7	Steelhead (Oncorhynchus (=Salmo) mykiss) -Upper Columbia River DPS - See 50 CFR 223.102	T	\$0	\$48,584	\$48,584	\$0	\$48,584
	21	Steelhead (Oncorhynchus (=Salmo) mykiss) -Upper Willamette River DPS - See 50 CFR 223.102	T	\$0	\$39,560	\$39,560	\$0	\$39,560
	103	Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - South Atlantic DPS - See 50 CFR 224.101	E	\$0	\$0	\$0	\$3,131	\$3,131
	142	sturgeon, Green (Acipenser medirostris) -Southern DPS - See 50 CFR 223.102	T	\$391	\$0	\$391	\$0	\$391
	89	Sturgeon, Gulf (Acipenser oxyrinchus (=oxyrhynchus) desotoi) -Wherever found	T	\$0	\$0	\$0	\$3,131	\$3,131
	156	Sturgeon, pallid (Scaphirhynchus albus) -Wherever found	E	\$833	\$0	\$833	\$0	\$833
	73	Sturgeon, shortnose (Acipenser brevirostrum) - Wherever found	E	\$0	\$0	\$0	\$3,131	\$3,131
	6	Trout, bull (Salvelinus confluentus) -U.S.A., conterminous, (lower 48 states)	T	\$0	\$1,708,392	\$1,708,392	\$0	\$1,708,392
Fishes Total				\$1,226,070	\$4,705,815	\$5,931,885	\$12,524	\$5,944,409
Flowering Plants	290	Amaranth, seabeach (Amaranthus pumilus) -Wherever found	T	\$2,400	\$0	\$2,400	\$0	\$2,400
	809	Ayenia, Texas (Ayenia limitaris) -Wherever found	E	\$7,430	\$0	\$7,430	\$0	\$7,430
	106	Baccharis, Encinitas (Baccharis vanessae) -Wherever found	T	\$913,900	\$0	\$913,900	\$0	\$913,900
	929	Bird's beak, palmate-bracted (Cordylanthus palmatus) - Wherever found	E	\$56,641	\$0	\$56,641	\$0	\$56,641
	1002	Cactus, star (Astrophytum asterias) -Wherever found	E	\$7,430	\$0	\$7,430	\$229,000	\$236,430
	243	Checker-mallow, Nelson's (Sidalcea nelsoniana) - Wherever found	T	\$275,333	\$0	\$275,333	\$0	\$275,333
	891	Dogweed, ashy (Thymophylla tephroleuca) -Wherever found	E	\$7,430	\$0	\$7,430	\$0	\$7,430

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	1602	Irisette, white (<i>Sisyrinchium dichotomum</i>) -Wherever found	E	\$0	\$0	\$0	\$984,500	\$984,500
	907	Manioc, Walker's (<i>Manihot walkerae</i>) -Wherever found	E	\$7,430	\$0	\$7,430	\$0	\$7,430
	915	Orcutt grass, hairy (<i>Orcuttia pilosa</i>) -Wherever found	E	\$56,250	\$0	\$56,250	\$0	\$56,250
	425	Phacelia, DeBeque (<i>Phacelia submutica</i>) -Wherever found	T	\$0	\$6,000	\$6,000	\$0	\$6,000
	492	Pondberry (<i>Lindera melissifolia</i>) -Wherever found	E	\$1,055,167	\$0	\$1,055,167	\$0	\$1,055,167
	1020	Spurge, Hoover's (<i>Chamaesyce hooveri</i>) -Wherever found	T	\$56,250	\$0	\$56,250	\$0	\$56,250
	1024	Tuctoria, Greene's (<i>Tuctoria greenei</i>) -Wherever found	E	\$56,641	\$0	\$56,641	\$0	\$56,641
	1416	Blazingstar, scrub (<i>Liatris ohlingerae</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1416	Fringe-tree, pygmy (<i>Chionanthus pygmaeus</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1507	Pigeon wings (<i>Clitoria fragrans</i>) -	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	1473	Beargrass, Britton's (<i>Nolina brittoniana</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1569	Mint, Garrett's (<i>Dicerandra christmanii</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1416	Whitlow-wort, papery (<i>Paronychia chartacea</i>) -	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	1569	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) -	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	1473	Bonamia, Florida (<i>Bonamia grandiflora</i>) -	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	1473	Polygala, Lewton's (<i>Polygala lewtonii</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	799	Lupine, scrub (<i>Lupinus aridorum</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	320	Pink, swamp (<i>Helonias bullata</i>) -	T	\$2,400	\$0	\$2,400	\$0	\$2,400
	1507	Hypericum, highlands scrub (<i>Hypericum cumulicola</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1402	Mustard, Carter's (<i>Warea carteri</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1402	Wireweed (<i>Polygonella basiramia</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1185	Sandlace (<i>Polygonella myriophylla</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	1362	Plum, scrub (<i>Prunus geniculata</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
	777	Bulrush, Northeastern (<i>Scirpus ancistrochaetus</i>) -	E	\$313,720	\$0	\$313,720	\$0	\$313,720
	1041	Ziziphus, Florida (<i>Ziziphus celata</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
Flowering Plants Total				\$2,847,510	\$6,000	\$2,853,510	\$1,213,500	\$4,067,010
Insects	143	Beetle, valley elderberry longhorn (<i>Desmocerus californicus dimorphus</i>) -Wherever found	T	\$391	\$0	\$391	\$0	\$391

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	714	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. wrighti)) -Wherever found	E	\$4,333,333	\$0	\$4,333,333	\$0	\$4,333,333
Insects Total				\$4,333,724	\$0	\$4,333,724	\$0	\$4,333,724
Lichens	1533	Cladonia, Florida perforate (<i>Cladonia perforata</i>) -	E	\$1,818	\$0	\$1,818	\$0	\$1,818
Lichens Total				\$1,818	\$0	\$1,818	\$0	\$1,818
Mammals	167	bat, Florida bonneted (<i>Eumops floridanus</i>) -Wherever found	E	\$0	\$0	\$0	\$3,131	\$3,131
	92	Bat, gray (<i>Myotis grisescens</i>) -Wherever found	E	\$400,958	\$0	\$400,958	\$33,133	\$434,091
	59	Bat, Indiana (<i>Myotis sodalis</i>) -Wherever found	E	\$1,385,587	\$0	\$1,385,587	\$41,875	\$1,427,462
	49	Bat, Northern long-eared (<i>Myotis septentrionalis</i>) -Wherever found	E	\$0	\$0	\$0	\$204,875	\$204,875
	57	Lynx, Canada (<i>Lynx canadensis</i>) -Wherever Found in Contiguous U.S.	T	\$128,720	\$0	\$128,720	\$0	\$128,720
	22	Manatee, West Indian (<i>Trichechus manatus</i>) -Wherever found	T	\$652,415	\$0	\$652,415	\$135,710	\$788,125
	368	Mouse, southeastern beach (<i>Peromyscus polionotus niveiventris</i>) -wherever found	T	\$156,250	\$0	\$156,250	\$0	\$156,250
	228	Ocelot (<i>Leopardus</i> (=Felis) <i>pardalis</i>) -wherever found	E	\$24,002	\$0	\$24,002	\$0	\$24,002
	81	Panther, Florida (<i>Puma</i> (=Felis) <i>concolor coryi</i>) -Wherever found	E	\$0	\$0	\$0	\$3,131	\$3,131
	32	Sea lion, Steller (<i>Eumetopias jubatus</i>) -Western DPS - See 50 CFR 224.101	E	\$74,045	\$0	\$74,045	\$0	\$74,045
	1608	Whale, gray (<i>Eschrichtius robustus</i>) -Western North Pacific DPS See 50 CFR 224.101	E	\$74,045	\$0	\$74,045	\$0	\$74,045
	60	Whale, killer (<i>Orcinus orca</i>) -Southern Resident DPS See 50 CFR 224.101	E	\$74,045	\$0	\$74,045	\$0	\$74,045
	4	Whale, North Atlantic Right (<i>Eubalaena glacialis</i>) -Wherever found	E	\$0	\$0	\$0	\$3,131	\$3,131

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	64	Wolf, gray (Canis lupus) -U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of Highway 97 and Highway 17 north of Mesa and that portion of WA west of the centerline of Highway 395 south of Mesa). Mexico.	E	\$13,000	\$0	\$13,000	\$0	\$13,000
Mammals Total				\$2,983,067	\$0	\$2,983,067	\$424,986	\$3,408,053
Reptiles	65	Alligator, American (Alligator mississippiensis) - Wherever found	SAT	\$0	\$0	\$0	\$4,729	\$4,729
	246	Crocodile, American (Crocodylus acutus) -U.S.A. (FL)	T	\$156,250	\$0	\$156,250	\$0	\$156,250
	416	Crocodile, American (Crocodylus acutus) -Wherever found, except in U.S.A. (FL)	E	\$0	\$0	\$0	\$3,131	\$3,131
	1680	Lizard, Coachella Valley fringe-toed (Uma inornata) - Wherever found	T	\$1,300,000	\$0	\$1,300,000	\$0	\$1,300,000
	190	Massasauga (=rattlesnake), eastern (Sistrurus catenatus) -Wherever found	T	\$0	\$0	\$0	\$205,000	\$205,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	90	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	T	\$693,596	\$0	\$693,596	\$0	\$693,596
	172	Sea turtle, green (<i>Chelonia mydas</i>) -Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	T	\$0	\$0	\$0	\$4,853	\$4,853
	52	Sea turtle, hawksbill (<i>Eretmochelys imbricata</i>) - Wherever found	E	\$380,252	\$0	\$380,252	\$4,853	\$385,105
	51	Sea turtle, Kemp's ridley (<i>Lepidochelys kempii</i>) - Wherever found	E	\$4,615,279	\$0	\$4,615,279	\$4,853	\$4,620,132
	41	Sea turtle, leatherback (<i>Dermochelys coriacea</i>) - Wherever found	E	\$868,596	\$0	\$868,596	\$4,853	\$873,449
	47	Sea turtle, loggerhead (<i>Caretta caretta</i>) -Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	T	\$0	\$0	\$0	\$4,853	\$4,853
	406	skink, blue-tailed mole (<i>Eumeces egregius lividus</i>) - Wherever found	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	327	Skink, sand (<i>Neoseps reynoldsi</i>) -Wherever found	T	\$1,818	\$0	\$1,818	\$0	\$1,818
	967	Snake, Atlantic salt marsh (<i>Nerodia clarkii taeniata</i>) - Wherever found	T	\$156,250	\$0	\$156,250	\$0	\$156,250
	612	Snake, copperbelly water (<i>Nerodia erythrogaster neglecta</i>) -Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$87,800	\$0	\$87,800	\$0	\$87,800

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	State Total	Species Total
	161	Snake, eastern indigo (Drymarchon couperi) -Wherever found	T	\$0	\$0	\$0	\$3,131	\$3,131
	211	Snake, giant garter (Thamnophis gigas) -Wherever found	T	\$3,336,641	\$0	\$3,336,641	\$0	\$3,336,641
	53	Tortoise, desert (Gopherus agassizii) -Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$1,300,000	\$2,511,000	\$3,811,000	\$0	\$3,811,000
	196	Tortoise, gopher (Gopherus polyphemus) -Western DPS	T	\$51,667	\$0	\$51,667	\$0	\$51,667
	641	Turtle, ringed map (Graptemys oculifera) -Wherever found	T	\$51,667	\$0	\$51,667	\$0	\$51,667
Reptiles Total				\$13,001,634	\$2,511,000	\$15,512,634	\$240,256	\$15,752,890
Snails	1473	Lioplax, cylindrical (snail) (Lioplax cyclostomaformis) -Wherever found	E	\$400,125	\$0	\$400,125	\$0	\$400,125
	1464	Pebblesnail, flat (Lepyrium showalteri) -Wherever found	E	\$400,125	\$0	\$400,125	\$0	\$400,125
	1338	Rocksail, round (Leptoxis ampla) -Wherever found	T	\$400,125	\$0	\$400,125	\$0	\$400,125
Snails Total				\$1,200,375	\$0	\$1,200,375	\$0	\$1,200,375
Multi-Species Total				\$0	\$0	\$0	\$9,000	\$9,000
Total Land Expenditures				\$70,640,532	\$7,222,815	\$77,863,347	\$2,650,929	\$80,514,276

TABLE 4. FY 2022 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES AND THE STATES

This table displays combined species expenditures and land acquisition costs by each reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species are included, as well as costs for multi-species conservation activities.

TABLE 4. FY 2022 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES AND THE STATES

Agency Type	Expense Type	Agency	Species Total	Land Total	Cumulative Total
Federal	Federal Domestic Expenditures	DOC National Oceanic and Atmospheric Administration	\$245,157,000		\$245,157,000
		DOD Army Corps of Engineers	\$617,760		\$617,760
		DOE Bonneville Power Administration	\$218,254,241	\$4,705,815	\$222,960,056
		DOE Western Area Power Administration	\$1,615,539		\$1,615,539
		DOI Bureau of Land Management	\$22,003,602	\$2,517,000	\$24,520,602
		DOI Fish and Wildlife Service	\$177,135,515	\$70,566,487	\$247,702,002
		DOI National Park Service	\$27,596,601		\$27,596,601
		Federal Aviation Administration	\$260,500		\$260,500
		Nuclear Regulatory Commission	\$97,100		\$97,100
		Smithsonian Institution	\$236,300		\$236,300
		Tennessee Valley Authority	\$520,262		\$520,262
		USDA Animal and Plant Health Inspection Service	\$252,400		\$252,400
	Federal Domestic Expenditures subtotal		\$693,746,820	\$77,789,302	\$771,536,122
	Federal Foreign Expenditures	DOC National Oceanic and Atmospheric Administration	\$3,392,000		\$3,392,000
		DOD Army Corps of Engineers	\$60,480		\$60,480
		DOI Fish and Wildlife Service	\$12,668,767	\$74,045	\$12,742,812
		DOI National Park Service	\$5,000		\$5,000
		Nuclear Regulatory Commission	\$5,800		\$5,800
		Smithsonian Institution	\$341,900		\$341,900
		USDA Animal and Plant Health Inspection Service	\$68,200		\$68,200
	Federal Foreign Expenditures subtotal		\$16,542,147	\$74,045	\$16,616,192
	Federal Multi-species	DOI Fish and Wildlife Service	\$11,480,309		\$11,480,309
		Federal Aviation Administration	\$42,000		\$42,000
		Nuclear Regulatory Commission	\$1,027,200		\$1,027,200
		Tennessee Valley Authority	\$110,000		\$110,000
		USDA Animal and Plant Health Inspection Service	\$616,100		\$616,100
	Federal Multi-species subtotal		\$13,275,609	\$0	\$13,275,609
	Federal Domestic, Foreign and Multi-species Expenditures subtotal		\$723,564,576	\$77,863,347	\$801,427,923
State	State Domestic Expenditures	State Governments	\$24,858,580	\$2,638,798	\$27,497,378
	State Foreign Species	State Governments	\$126,023	\$3,131	\$129,154
	State Multi-species	State Governments	\$5,709,444	\$9,000	\$5,718,444
	State Domestic, Foreign and Multi-species Expenditures subtotal		\$30,694,047	\$2,650,929	\$33,344,976
Total Federal and State Domestic and Foreign Expenditures			\$754,258,623	\$80,514,276	\$834,772,899

Appendix A. Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species -- FY 2022

Overview

This reporting guidance provides information on how to report expenditures attributable to individual species to the extent this information is reasonably available. The Congress requested information for this report on a primarily species-by-species basis. However, to provide the most accurate account of expenditures for the conservation of species listed under the Endangered Species Act, a “multi-species” reporting feature was added. This option allows the reporting of lump sum costs for multi-species conservation activities **when it is not possible to account accurately for individual species.**

The ESA Expenditures Report to Congress is an annual report and the current data call is requesting expenditures from Federal Fiscal Year 2022. The standard procedure for data submission would involve direct entry of expenditures into the ECOS Expenditures module. However, the system is not available for FY22 data entry. This guidance outlines the same method used for FY 2021 data, requesting data be submitted on the customized spreadsheet provided.

The FY 2022 spreadsheet is the year’s full Threatened & Endangered Species list that has been formatted with fields for General and Land Expenditures, and follows the format for the bulk upload. The bulk upload of all spreadsheet will be facilitated by FWS upon the close of the reporting window.

Data Reporting Instruction

FY 2022 Reporting Instruction

Each agency must report FY 2022 data using the customized spreadsheet (<https://ecos.fws.gov/docs/expenditures/ExpendituresSpeciesList2022.xlsx>). The customized spreadsheet will be locked to prevent changes to species information. Agencies will have access to enter expenditures for each species in the designated cells. Information on the scope of the activities to report is provided in the “Data Gathering Instruction” section below.

Completed FY 2022 spreadsheets should be emailed to fwhq_es_bbs@fws.gov with a subject line of “FY 2022 Expenditures Report Data Submission.”

Data Gathering Instruction

Background

- Section 18 of the Endangered Species Act of 1973 (16 U.S. 1531 *et seq.*) reads, in part, “...the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain--

(1) an accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act; and

(2) an accounting on a species-by-species basis for all reasonably identifiable expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act by States receiving grants under section 6."

- The conference report for the 1988 amendments made some of the following points concerning section 18 of the Act:
 - Provision should not become unduly burdensome or result in diversion of funds from the operation of the program.
 - The Secretary of the Interior is to make a good faith effort to develop and obtain data but is not to undertake extensive or extraordinary measures.
 - Generalized dollar estimates will suffice.
 - Amendment does not apply to expenditures for delisted, proposed, and candidate species.
 - Amendment is not to require new species-specific time sheets for the Service or other reporting agency.
 - Listing, consultation, and law enforcement are to be included to the extent they are "reasonably identifiable."
 - Only salary costs that can be reasonably identified as applicable to listed species need be reported.
 - For other Federal and State agencies, a good faith effort to collect information is expected.
 - Submission of data from any agency is not to be a precondition to receiving any contracts or grants or establishing other arrangements with the Fish and Wildlife Service.
 - Only one data set per agency should be submitted for any given fiscal year.

Data Format

- Species should be identified by the same scientific name as found in the most current list of endangered and threatened species.
- Amounts should be reported in whole dollars. Amounts above \$2,000 may be rounded to the nearest \$500 or \$1,000; smaller amounts may be rounded to the nearest \$100. **Do not report amounts lower than \$100.**

Scope of Expenditures to Report

- Only species on the Federal list of Endangered and Threatened Wildlife and Plants (50 CFR Part 17) at the end of Federal fiscal year 2022 (October 1, 2021 to September 30,

2022) are to be reported. Expenditures made prior to the actual listing date of a species, but still within the same year, may be reported (e.g., costs of public meetings, notices, surveys, initial recovery efforts). Funds expended for delisted, separate populations of listed vertebrates cannot be included in the report (e.g., Atlantic coast least terns, Alaska gray wolves). Expenditures for State-listed species are not reportable unless they are also federally listed. Amounts for foreign species on the list (e.g., grants or contracts carried out in another country) are reportable in the same manner as domestic species, but again, to the extent they are reasonably identifiable.

- Only reasonably identifiable expenditures for listed species will be compiled for this report. Extraordinary accounting to track funds expended on listed species is not expected. **Funds reported should be those actually expended, not obligated.**
- Examples of reportable expenditures include status surveys, habitat management, habitat acquisition, research, propagation (including surrogate species), recovery plan development or implementation, and mitigation. **Projects must primarily benefit listed species** and not other conservation goals, although if a readily identifiable portion of the expenditure is for the benefit of listed species (either individually or collectively), that portion of the expenditure may be reported.
- All habitat (land or water) acquisition costs are to be reported separately from all other identifiable expenditures. Such acquisitions must be primarily for the purpose of conserving federally listed species. As with other projects, those portions of habitat acquisition costs that are specifically for a listed species may be prorated out of the total acquisition costs when the other habitat is being acquired for other purposes (e.g., 25 acres of a 250-acre purchase will be set aside).
- Expenditures in a single project devoted to a number of listed species should be prorated by species, if possible. **When it is not possible to determine individual listed species costs in a multi-species project by prorating, those costs should be entered, according to the following instruction:**
 - **FY 2022:** Enter the cost in total, in the multi-species tab of the spreadsheet. The comment box, which is part of this feature, should be used to identify the species covered by this expenditure amount.
- General surveys or projects that cover a large number of species, some of which may not be listed, are reportable only to extent that the agency can reasonably identify the portion of costs that benefit listed species.
- Expenditures associated with consultations pursuant to Section 7 of the Act should be reported as for the benefit of a specific species to the extent they are readily identifiable to that particular species. Thus, a formal consultation dealing with a single species, or up to several species where the relative costs are easily divisible, should be reported under that or those species. **Consultation costs not readily attributable to individual species can be reported in the multi-species category.**

- Monitoring and survey costs should be reported when: (1) the cost of the monitoring or survey was incurred during the period the species was under a proposed listing and the species was actually listed in the same fiscal year that the monitoring and surveys were conducted; (2) the monitoring or survey covered both candidates and listed species, and the costs can be prorated to include only the listed species; or (3) the cost was readily identifiable to individual species. **Costs meeting the criteria in (1) and (2), but not readily identifiable to individual species can be reported in the multi-species category.**
- Salary and benefits of an employee working full-time on a single species or whose time is devoted to particular species that can be readily identified should be reported for those species. Travel costs should be treated similarly. **Staff costs for work on multi-species activities that cannot be identified to particular species can be reported in the multi-species category.**
- Any State or Federal project that incurs increased costs related directly to mitigation or other conservation efforts on behalf of federally listed species should report that added cost, by species, if readily identifiable, or **may report such costs in the multi-species category.**
- All State agencies (e.g., fish and wildlife, parks, heritage programs, forests, highways) should report their expenditures, but only a single, collective report from each State should be submitted. **Expenditures should include conservation costs for fish, wildlife, and plants.** For ESA section 6 grants or other Federal grants, States should report **only the State cost-share portion** of each grant. States should not include other sources of non-Federal cost share, e.g., third-party contributions, unless those contributions are donated to the State as a general donation.
- For reimbursable funds, the paying or disbursing agency should report the expenditure.
- **Litigation costs should not be reported.**

Appendix B. Reports Provided by Other Federal Agencies for Fiscal Year 2022

Department of Agriculture:

- Animal and Plant Health Inspection Service
- Farm Service Agency*
- Forest Service*
- Natural Resources Conservation Service*

Department of Commerce:

- National Oceanic and Atmospheric Administration

Department of Defense:

- Office of the Under Secretary
 - Representing: Air Force*, Army (includes Defense Logistics Agency)*, Marine Corps*, Navy*, and Army Corps of Engineers

Department of Energy:

- Bonneville Power Administration
- Southeastern Power Administration*
- Southwestern Power Administration*
- Western Area Power Administration

Department of Homeland Security

- Coast Guard*
- Customs and Border Protection*
- Federal Emergency Management Agency*
- Office of Science and Technology*

Department of the Interior:

- Bureau of Indian Affairs*
- Bureau of Land Management
- Bureau of Ocean Energy Management*
- Bureau of Reclamation*
- Bureau of Safety and Environmental Enforcement*
- National Park Service
- Office of Surface Mining*
- U.S. Geological Survey*

Department of Transportation:

- Federal Aviation Administration
- Federal Highway Administration*

Independent Agencies:

Environmental Protection Agency*
Federal Communications Commission*
Federal Energy Regulatory Commission*
Nuclear Regulatory Commission
Smithsonian Institution
Tennessee Valley Authority

*Agencies with no report provided for FY 2022

DOC National Oceanic and Atmospheric Administration - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$636,000		\$636,000
		Coral, elkhorn	Acropora palmata	Wherever found	\$636,000		\$636,000
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$636,000		\$636,000
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$636,000		\$636,000
		coral, Pillar	Dendrogyra cylindrus	Wherever found	\$636,000		\$636,000
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$636,000		\$636,000
		Coral, staghorn	Acropora cervicornis	Wherever found	\$636,000		\$636,000
	<i>Corals subtotal</i>				<i>\$4,452,000</i>		<i>\$4,452,000</i>
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound-Georgia Basin DPS See 50 CFR 224.101	\$559,000		\$559,000
		Eulachon	Thaleichthys pacificus	Southern DPS - See 50 CFR 223.102	\$559,000		\$559,000
		grouper, Nassau	Epinephelus striatus	(blank)	\$746,000		\$746,000
		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Georgia Basin DPS See 50 CFR 223.102	\$559,000		\$559,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		salmon Atlantic	Salmo salar	U.S.A. ME Gulf of Maine Atlantic Salmon Distinct Population Segment, which includes all naturally reproducing wild populations and those river-specific hatchery populations of Atlantic salmon having historical, river-specific characteristics found north of and including tributaries of the lower Kennebec River to, but not including, the mouth of the St. Croix River at the U.S.-Canda border. To date, the Services have determined that these populations are found in the Dennys, East Machias, Machias, Pleasant, Narraguagus, Sheepscot, and Ducktrap Rivers and in Cove Brook, Maine	\$6,800,000		\$6,800,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU - See 50 CFR 223.102	\$4,616,000		\$4,616,000
				Central Valley spring-run ESU See 50 CFR 223.102	\$4,566,000		\$4,566,000
				Lower Columbia River ESU See 50 CFR 223.102	\$6,401,000		\$6,401,000
				Puget Sound ESU See 50 CFR 223.102	\$5,319,000		\$5,319,000
				Sacramento River winter-run ESU - See 50 CFR 224.101	\$4,566,000		\$4,566,000
				Snake River fall-run ESU See 50 CFR 223.102	\$6,352,000		\$6,352,000
				Snake River spring/summer-run ESU See 50 CFR 223.102	\$8,052,000		\$8,052,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Upper Columbia River spring-run ESU See 50 CFR 224.101	\$4,338,000		\$4,338,000
				Upper Willamette River ESU See 50 CFR 223.102	\$3,930,000		\$3,930,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU See 50 CFR 223.102	\$5,819,000		\$5,819,000
				Hood Canal summer-run ESU - See 50 CFR 223.102	\$5,319,000		\$5,319,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU - See 50 CFR 224.101	\$4,566,000		\$4,566,000
				Lower Columbia River ESU - See 50 CFR 223.102	\$6,401,000		\$6,401,000
				Oregon Coast ESU - See 50 CFR 223.102	\$11,128,000		\$11,128,000
				Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	\$8,980,000		\$8,980,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU - See 50 CFR 223.102	\$5,319,000		\$5,319,000
				Snake River ESU - See 50 CFR 224.101	\$4,043,000		\$4,043,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$847,000		\$847,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS - See 50 CFR 223.102	\$4,566,000		\$4,566,000
				Central California Coast DPS - See 50 CFR 223.102	\$4,566,000		\$4,566,000
				Lower Columbia River DPS - See 50 CFR 223.102	\$6,401,000		\$6,401,000
				Middle Columbia River DPS - See 50 CFR 223.102	\$5,319,000		\$5,319,000
				Northern California DPS - See 50 CFR 223.102	\$5,316,000		\$5,316,000
				Puget Sound DPS - See 50 CFR 223.102	\$5,319,000		\$5,319,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Snake River Basin DPS - See 50 CFR 223.102	\$8,052,000		\$8,052,000
				South-Central California Coast DPS - See 50 CFR 223.102	\$4,566,000		\$4,566,000
				Southern California DPS - See 50 CFR 224.101	\$4,566,000		\$4,566,000
				Upper Columbia River DPS - See 50 CFR 223.102	\$3,914,000		\$3,914,000
				Upper Willamette River DPS - See 50 CFR 223.102	\$3,930,000		\$3,930,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS - See 50 CFR 224.101	\$781,000		\$781,000
				Chesapeake Bay DPS - See 50 CFR 224.101	\$843,000		\$843,000
				Gulf of Maine DPS - See 50 CFR 223.102	\$468,000		\$468,000
				New York Bight DPS - See 50 CFR 224.101	\$788,000		\$788,000
				South Atlantic DPS - See 50 CFR 224.101	\$698,000		\$698,000
		sturgeon, Green	Acipenser medirostris	Southern DPS - See 50 CFR 223.102	\$586,000		\$586,000
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$576,000		\$576,000
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$1,060,000		\$1,060,000
	<i>Fishes subtotal</i>				<i>\$172,100,000</i>		<i>\$172,100,000</i>
	Mammals	Sea lion, Steller	Eumetopias jubatus	Western DPS - See 50 CFR 224.101	\$4,685,000		\$4,685,000
		Seal, bearded	Erignathus barbatus nauticus	Beringia DPS - See 50 CFR 223.102	\$1,891,000		\$1,891,000
		Seal, Guadalupe fur	Arctocephalus townsendi	Wherever found	\$226,000		\$226,000
		Seal, Hawaiian monk	Monachus schauinslandi	Wherever found	\$4,264,000		\$4,264,000
		Seal, Ringed	Phoca (=Pusa) hispida botnica	Wherever found	\$1,891,000		\$1,891,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		whale, beluga	Delphinapterus leucas	Cook Inlet DPS See 50 CFR 224.101	\$3,037,000		\$3,037,000
		Whale, blue	Balaenoptera musculus	Wherever found	\$1,254,000		\$1,254,000
		Whale, bowhead	Balaena mysticetus	Wherever found	\$407,000		\$407,000
		whale, false killer	Pseudorca crassidens	Main Hawaiian Islands Insular DPS See 50 CFR 224.101	\$261,000		\$261,000
		Whale, finback	Balaenoptera physalus	Wherever found	\$1,255,000		\$1,255,000
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$356,000		\$356,000
		Whale, killer	Orcinus orca	Southern Resident DPS See 50 CFR 224.101	\$2,402,000		\$2,402,000
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$28,568,000		\$28,568,000
		Whale, North Pacific Right	Eubalaena japonica	Wherever found	\$381,000		\$381,000
		Whale, Sei	Balaenoptera borealis	Wherever found	\$1,071,000		\$1,071,000
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$1,071,000		\$1,071,000
	<i>Mammals subtotal</i>				\$53,020,000		\$53,020,000
	Reptiles	Sea turtle, green	Chelonia mydas	Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	\$745,000		\$745,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Green sea turtles originating from the Central South Pacific Ocean, bounded by the following coordinates: 9 degrees N., 175 degrees W. in the northwest; 9 degrees N., 125 degrees W. in the northeast; 40 degrees S., 96 degrees W. in the southeast; 40 degrees S., 176 degrees E. in the southwest; and 13 degrees S., 171 degrees E. in the west	\$527,000		\$527,000
				Green sea turtles originating from the Central West Pacific Ocean, bounded by the following coordinates: 41 degrees N., 146 degrees E. in the northwest; 41 degrees N., 169 degrees E. in the northeast; 9 degrees N., 175 degrees W. in the east; 13 degrees S., 171 degrees E. in the southeast; along the northern coast of the island of New Guinea; and 4.5 degrees N., 129 degrees E. in the west	\$1,026,000		\$1,026,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Green sea turtles originating from the East Pacific Ocean, bounded by the following lines and coordinates: 41 degrees N., 143 degrees W. in the northwest; 41 degrees N. Lat. in the north; along the western coasts of the Americas; 40 degrees S. Lat. in the south; and 40 degrees S., 96 degrees W. in the southwest	\$934,000		\$934,000
				Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	\$972,000		\$972,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$1,865,000		\$1,865,000
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,250,000		\$1,250,000
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$2,787,000		\$2,787,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS - Loggerhead sea turtles originating from the North Pacific north of the equator and south of 60 degrees N. Lat	\$1,429,000		\$1,429,000
				Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	\$1,402,000		\$1,402,000
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$230,000		\$230,000
	<i>Reptiles subtotal</i>				<i>\$13,167,000</i>		<i>\$13,167,000</i>
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$748,000		\$748,000
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$1,670,000		\$1,670,000
	<i>Snails subtotal</i>				<i>\$2,418,000</i>		<i>\$2,418,000</i>
Domestic subtotal					\$245,157,000		\$245,157,000
Foreign	Corals	No common name	Acropora globiceps	Wherever found	\$377,000		\$377,000
			Acropora jacquelineae	Wherever found	\$377,000		\$377,000
			Acropora retusa	Wherever found	\$377,000		\$377,000
			Acropora speciosa	Wherever found	\$377,000		\$377,000
			Euphyllia paradivisa	Wherever found	\$377,000		\$377,000
			Isopora crateriformis	Wherever found	\$377,000		\$377,000
			Seriatopora aculeata	Wherever found	\$393,000		\$393,000
	<i>Corals subtotal</i>				<i>\$2,655,000</i>		<i>\$2,655,000</i>
	Mammals	Whale, humpback	Megaptera novaeangliae	Central America DPS	\$356,000		\$356,000
				Western North Pacific	\$381,000		\$381,000
	<i>Mammals subtotal</i>				<i>\$737,000</i>		<i>\$737,000</i>
Foreign subtotal					\$3,392,000		\$3,392,000
Grand Total					\$248,549,000		\$248,549,000

DOD Army Corps of Engineers - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Birds	Knot, red	Calidris canutus rufa	Wherever found	\$44,928		\$44,928
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$19,008		\$19,008
		rail, Eastern Black	Laterallus jamaicensis ssp. jamaicensis	Wherever found	\$15,552		\$15,552
	<i>Birds subtotal</i>				\$79,488		\$79,488
	Fishes	Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS - See 50 CFR 224.101	\$165,456		\$165,456
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$73,440		\$73,440
	<i>Fishes subtotal</i>				\$238,896		\$238,896
	Flowering Plants	Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$19,008		\$19,008
		Beaked-rush, Knieskern's	Rhynchospora knieskernii	Wherever found	\$3,456		\$3,456
		Joint-vetch, Sensitive	Aeschynomene virginica	Wherever found	\$12,096		\$12,096
		Pink, swamp	Helonias bullata	(blank)	\$20,736		\$20,736
	<i>Flowering Plants subtotal</i>				\$55,296		\$55,296
	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$5,184		\$5,184
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$28,080		\$28,080
	<i>Mammals subtotal</i>				\$33,264		\$33,264
	Reptiles	Sea turtle, green	Chelonia mydas	Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	\$59,184		\$59,184
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$9,936		\$9,936

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$60,480		\$60,480
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$58,752		\$58,752
		turtle, bog	Glyptemys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$22,464		\$22,464
	<i>Reptiles subtotal</i>				<i>\$210,816</i>		<i>\$210,816</i>
	Domestic subtotal				\$617,760		\$617,760
Foreign	Reptiles	Sea turtle, loggerhead	Caretta caretta	Loggerhead sea turtles originating from the South Pacific south of the equator, north of 60 degrees S. Lat., west of 67 degrees W. Long., and east of 141 degrees E. Long.	\$60,480		\$60,480
	<i>Reptiles subtotal</i>				<i>\$60,480</i>		<i>\$60,480</i>
	Foreign subtotal				\$60,480		\$60,480
	Grand Total				\$678,240		\$678,240

DOE Bonneville Power Administration - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Fishes	Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU See 50 CFR 223.102	\$10,890,472	\$531,081	\$11,421,553
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU See 50 CFR 223.102	\$11,705,903	\$539,338	\$12,245,241
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU See 50 CFR 223.102	\$28,097,090	\$39,560	\$28,136,650
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia River spring-run ESU See 50 CFR 224.101	\$22,595,795	\$39,560	\$22,635,355
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU See 50 CFR 223.102	\$7,199,560	\$39,560	\$7,239,120
		Salmon, chum	Oncorhynchus keta	Columbia River ESU See 50 CFR 223.102	\$3,851,097	\$530,314	\$4,381,411
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU - See 50 CFR 223.102	\$4,658,545	\$531,081	\$5,189,626
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	\$242,630		\$242,630
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU - See 50 CFR 224.101	\$8,765,952	\$39,560	\$8,805,512
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS - See 50 CFR 223.102	\$5,695,362	\$531,081	\$6,226,443
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS - See 50 CFR 223.102	\$36,271,362	\$39,560	\$36,310,922
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS - See 50 CFR 223.102	\$30,323,801	\$48,584	\$30,372,385
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS - See 50 CFR 223.102	\$14,921,627	\$48,584	\$14,970,211
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS - See 50 CFR 223.102	\$3,929,643	\$39,560	\$3,969,203
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), (Kootenai R. system)	\$12,226,330		\$12,226,330
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, (lower 48 states)	\$14,528,387	\$1,708,392	\$16,236,779

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$2,350,685		\$2,350,685
	<i>Fishes subtotal</i>				<i>\$218,254,241</i>	<i>\$4,705,815</i>	<i>\$222,960,056</i>
<i>Domestic subtotal</i>					<i>\$218,254,241</i>	<i>\$4,705,815</i>	<i>\$222,960,056</i>
Grand Total					\$218,254,241	\$4,705,815	\$222,960,056

DOE Western Area Power Administration - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$15,996		\$15,996
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$8,437		\$8,437
	<i>Amphibians subtotal</i>				\$24,433		\$24,433
	Birds	Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$112,280		\$112,280
		Cuckoo, yellow-billed	Coccyzus americanus	Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	\$32,784		\$32,784
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$26,347		\$26,347
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$23,847		\$23,847
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$55,611		\$55,611
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$5,602		\$5,602
		Rail, Yuma Ridgway's	Rallus obsoletus yumanensis	Wherever found	\$5,000		\$5,000
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$26,579		\$26,579
		Tern, California least	Sterna antillarum browni	Wherever found	\$5,000		\$5,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$6,548		\$6,548
	<i>Birds subtotal</i>				\$299,598		\$299,598
	Crustaceans	Crayfish, Shasta	Pacifastacus fortis	Wherever found	\$5,601		\$5,601
		Fairy shrimp, Conservancy	Branchinecta conservatio	Wherever found	\$8,436		\$8,436

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$6,546		\$6,546
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$34,901		\$34,901
		Tadpole shrimp, vernal pool	Lepidurus packardii	Wherever found	\$34,901		\$34,901
	<i>Crustaceans subtotal</i>				\$90,385		\$90,385
	Fishes	Bonytail	Gila elegans	Wherever found	\$77,500		\$77,500
		Chub, humpback	Gila cypha	Wherever found	\$77,500		\$77,500
		pikeminnow, Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$77,500		\$77,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU See 50 CFR 223.102	\$19,779		\$19,779
				Snake River spring/summer-run ESU See 50 CFR 223.102	\$19,778		\$19,778
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU - See 50 CFR 224.101	\$24,490		\$24,490
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$12,219		\$12,219
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS - See 50 CFR 223.102	\$24,495		\$24,495
				Upper Willamette River DPS - See 50 CFR 223.102	\$19,780		\$19,780
		sturgeon, Green	Acipenser medirostris	Southern DPS - See 50 CFR 223.102	\$6,548		\$6,548
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$188,000		\$188,000
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$77,500		\$77,500
	<i>Fishes subtotal</i>				\$625,089		\$625,089
	Flowering Plants	Bird's beak, palmate-bracted	Cordylanthus palmatus	Wherever found	\$6,549		\$6,549
		Cactus, Mesa Verde	Sclerocactus mesae-verdae	Wherever found	\$10,847		\$10,847
		Cactus, Pima pineapple	Coryphantha scheeri var. robustispina	Wherever found	\$5,000		\$5,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Checker-mallow, Keck's	Sidalcea keckii	Wherever found	\$5,602		\$5,602
		Fiddleneck, large-flowered	Amsinckia grandiflora	Wherever found	\$6,547		\$6,547
		Goldfields, Contra Costa	Lasthenia conjugens	Wherever found	\$6,547		\$6,547
		Grass, Colusa	Neostapfia colusana	Wherever found	\$5,602		\$5,602
		Grass, Solano	Tuctoria mucronata	Wherever found	\$5,602		\$5,602
		Jewelflower, California	Caulanthus californicus	Wherever found	\$7,492		\$7,492
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$15,847		\$15,847
		Meadowfoam, Butte County	Limnanthes floccosa ssp. californica	Wherever found	\$5,602		\$5,602
		Milk-vetch, Mancos	Astragalus humillimus	Wherever found	\$10,847		\$10,847
		milkvetch, Osterhout	Astragalus osterhoutii	Wherever found	\$37,014		\$37,014
		Milkweed, Welsh's	Asclepias welshii	Wherever found	\$1,500		\$1,500
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$9,039		\$9,039
		Orcutt grass, hairy	Orcuttia pilosa	Wherever found	\$5,602		\$5,602
		Orcutt grass, Sacramento	Orcuttia viscida	Wherever found	\$6,547		\$6,547
		Orcutt grass, San Joaquin Valley	Orcuttia inaequalis	Wherever found	\$6,547		\$6,547
		Orcutt grass, slender	Orcuttia tenuis	Wherever found	\$5,602		\$5,602
		Owl's-clover, fleshy	Castilleja campestris ssp. succulenta	Wherever found	\$5,602		\$5,602
		Penstemon, blowout	Penstemon haydenii	Wherever found	\$9,039		\$9,039
		Phacelia, North Park	Phacelia formosula	Wherever found	\$10,847		\$10,847
		Spurge, Hoover's	Chamaesyce hooveri	Wherever found	\$5,602		\$5,602
		Sunburst, Hartweg's golden	Pseudobahia bahiifolia	Wherever found	\$6,547		\$6,547
		wild buckwheat, clay-loving	Eriogonum pelinophilum	Wherever found	\$10,847		\$10,847
		Woolly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$7,493		\$7,493
	<i>Flowering Plants subtotal</i>				\$219,912		\$219,912

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Insects	Beetle, delta green ground	Elaphrus viridis	Wherever found	\$5,602		\$5,602
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$36,291		\$36,291
	<i>Insects subtotal</i>				<i>\$41,893</i>		<i>\$41,893</i>
	Mammals	Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$45,000		\$45,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$9,039		\$9,039
		Fox, San Joaquin kit	Vulpes macrotis mutica	Wherever found	\$39,559		\$39,559
		Kangaroo rat, Fresno	Dipodomys nitratoide exilis	Wherever found	\$29,231		\$29,231
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$32,935		\$32,935
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$15,847		\$15,847
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	Wherever found	\$10,847		\$10,847
	<i>Mammals subtotal</i>				<i>\$182,458</i>		<i>\$182,458</i>
	Reptiles	gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$8,000		\$8,000
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$36,312		\$36,312
		Snake, giant garter	Thamnophis gigas	Wherever found	\$34,208		\$34,208
		Tortoise, desert	Gopherus agassizii	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$49,033		\$49,033
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$4,218		\$4,218
	<i>Reptiles subtotal</i>				<i>\$131,771</i>		<i>\$131,771</i>
Domestic subtotal					<i>\$1,615,539</i>		<i>\$1,615,539</i>
Grand Total					\$1,615,539		\$1,615,539

DOI Bureau of Land Management - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$35,000		\$35,000
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$67,365		\$67,365
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$13,600		\$13,600
		Hellbender, Ozark	Cryptobranchus alleganiensis bishopi	Wherever found	\$152		\$152
	<i>Amphibians subtotal</i>				<i>\$116,117</i>		<i>\$116,117</i>
	Birds	Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)- see 17.84(j)	\$70,000		\$70,000
				U.S.A. only, except where listed as an experimental population	\$4,570		\$4,570
		Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$7,688		\$7,688
		Cuckoo, yellow-billed	Coccyzus americanus	Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	\$1,296,600		\$1,296,600
		Eider, spectacled	Somateria fischeri	Wherever found	\$77,000		\$77,000
		Eider, Steller's	Polysticta stelleri	U.S. Alaska breeding population only	\$65,000		\$65,000
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	U.S.A (AZ, NM)	\$29,400		\$29,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$816,011		\$816,011
		Horned lark, streaked	Eremophila alpestris strigata	Wherever found	\$1,363		\$1,363
		Knot, red	Calidris canutus rufa	Wherever found	\$3,066		\$3,066
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$1,280,890		\$1,280,890
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$216,970		\$216,970
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$3,684,087		\$3,684,087
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$28,986		\$28,986
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$77,500		\$77,500
		rail, Eastern Black	Laterallus jamaicensis ssp. jamaicensis	Wherever found	\$2,310		\$2,310
		Rail, Yuma Ridgway's	Rallus obsoletus yumanensis	Wherever found	\$280,631		\$280,631
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$663,362		\$663,362
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$495		\$495
		Stork, wood	Mycteria americana	U.S.A. (AL, FL, GA, MS, NC, SC)	\$1,155		\$1,155
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$64,000		\$64,000
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$1,485		\$1,485
	<i>Birds subtotal</i>				<i>\$8,672,569</i>		<i>\$8,672,569</i>
	Clams	Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$152		\$152
		Fanshell	Cyprogenia stegaria	Wherever found	\$152		\$152

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$76		\$76
		Hornshell, Texas	Popenaias popeii	Wherever found	\$9,500		\$9,500
		Mapleleaf, winged	Quadrula fragosa	U.S.A. (AL-specified portions of the Tennessee River; see 17.85(a)(1))	\$76		\$76
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$152		\$152
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$228		\$228
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$304		\$304
		Purple Cat's paw (=Purple Cat's paw pearlymussel)	Epioblasma obliquata	Wherever found; Except where listed as Experimental Populations	\$152		\$152
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$330		\$330
		Riffleshell, northern	Epioblasma rangiana	Wherever found	\$152		\$152
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$76		\$76
	<i>Clams subtotal</i>				<i>\$11,350</i>		<i>\$11,350</i>
	Crustaceans	Amphipod, Noel's	Gammarus desperatus	Wherever found	\$2,000		\$2,000
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$3,000		\$3,000
		Isopod, Socorro	Thermosphaeroma thermophilus	Wherever found	\$1,000		\$1,000
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$1,500		\$1,500
	<i>Crustaceans subtotal</i>				<i>\$7,500</i>		<i>\$7,500</i>
	Fishes	Bonytail	Gila elegans	Wherever found	\$65,064		\$65,064
		Chub, Chihuahua	Gila nigrescens	Wherever found	\$1,000		\$1,000
		Chub, Gila	Gila intermedia	Wherever found	\$17,724		\$17,724
		Chub, humpback	Gila cypha	Wherever found	\$45,514		\$45,514
		Chub, Pahrnagat roundtail	Gila robusta jordani	Wherever found	\$268		\$268
		Chub, Virgin River	Gila seminuda (=robusta)	Wherever found	\$17,373		\$17,373
		Dace, desert	Eremichthys acros	Wherever found	\$150,000		\$150,000
		Dace, Moapa	Moapa coriacea	Wherever found	\$30,000		\$30,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Gambusia, Pecos	Gambusia nobilis	Wherever found	\$3,000		\$3,000
		Minnow, loach	Tiaroga cobitis	Wherever found	\$30,362		\$30,362
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$2,500		\$2,500
		pikeminnow, Colorado	Ptychocheilus lucius	Salt and Verde R. drainages, AZ	\$2,667		\$2,667
				Wherever found, except where listed as an experimental population	\$54,339		\$54,339
		Poolfish, Pahrump	Empetrichthys latos	Wherever found	\$9,254		\$9,254
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$32,562		\$32,562
		Pupfish, Owens	Cyprinodon radiosus	Wherever found	\$27,000		\$27,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU - See 50 CFR 223.102	\$45,000		\$45,000
				Lower Columbia River ESU See 50 CFR 223.102	\$10,800		\$10,800
				Snake River fall-run ESU See 50 CFR 223.102	\$6,500		\$6,500
				Snake River spring/summer-run ESU See 50 CFR 223.102	\$112,061		\$112,061
				Upper Willamette River ESU See 50 CFR 223.102	\$180,500		\$180,500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU - See 50 CFR 223.102	\$10,800		\$10,800
				Oregon Coast ESU - See 50 CFR 223.102	\$2,306,483		\$2,306,483
				Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	\$144,434		\$144,434
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU - See 50 CFR 224.101	\$30,509		\$30,509
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$38		\$38
		Spikedace	Meda fulgida	Wherever found	\$30,238		\$30,238

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Spinedace, Big Spring	Lepidomeda mollispinis pratensis	Wherever found	\$1,206		\$1,206
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$1,761		\$1,761
		Springfish, Hiko White River	Crenichthys baileyi grandis	Wherever found	\$8,591		\$8,591
		Springfish, Railroad Valley	Crenichthys nevadae	Wherever found	\$592		\$592
		Springfish, White River	Crenichthys baileyi baileyi	Wherever found	\$804		\$804
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS - See 50 CFR 223.102	\$10,800		\$10,800
				Middle Columbia River DPS - See 50 CFR 223.102	\$3,000		\$3,000
				Northern California DPS - See 50 CFR 223.102	\$10,000		\$10,000
				Snake River Basin DPS - See 50 CFR 223.102	\$97,338		\$97,338
				Upper Columbia River DPS - See 50 CFR 223.102	\$15,000		\$15,000
				Upper Willamette River DPS - See 50 CFR 223.102	\$382,500		\$382,500
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$660		\$660
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$8,365		\$8,365
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), (Kootenai R. system)	\$10,000		\$10,000
		Sucker, Lost River	Deltistes luxatus	Wherever found	\$5,000		\$5,000
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$60,099		\$60,099
		Sucker, shortnose	Chasmistes brevirostris	Wherever found	\$7,800		\$7,800
		Sucker, Warner	Catostomus warnerensis	Wherever found	\$879,000		\$879,000
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	U.S. Only	\$18,630		\$18,630
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, (lower 48 states)	\$316,869		\$316,869

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$100		\$100
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$534,949		\$534,949
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$15,000		\$15,000
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$17,373		\$17,373
	<i>Fishes subtotal</i>				\$5,771,427		\$5,771,427
	Flowering Plants	Beardtongue, Penland	Penstemon penlandii	Wherever found	\$5,000		\$5,000
		Bear-poppy, Dwarf	Arctomecon humilis	Wherever found	\$31,648		\$31,648
		Bedstraw, El Dorado	Galium californicum ssp. sierrae	Wherever found	\$5,000		\$5,000
		Birds-in-a-nest, white	Macbridea alba	(blank)	\$330		\$330
		Bladderpod, Dudley Bluffs	Lesquerella congesta	Wherever found	\$5,000		\$5,000
		Blazingstar, Ash Meadows	Mentzelia leucophylla	Wherever found	\$30,000		\$30,000
		Blue-star, Kearney's	Amsonia kearneyana	Wherever found	\$3,200		\$3,200
		Buckwheat, steamboat	Eriogonum ovalifolium var. williamsiae	Wherever found	\$54,833		\$54,833
		Bush-clover, prairie	Lespedeza leptostachya	Wherever found	\$76		\$76
		Butterweed, Layne's	Senecio layneae	Wherever found	\$5,000		\$5,000
		Butterwort, Godfrey's	Pinguicula ionantha	(blank)	\$330		\$330
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$2,000		\$2,000
		Cactus, Colorado hookless	Sclerocactus glaucus	Wherever found	\$112,701		\$112,701
		Cactus, Fickeisen plains	Pediocactus peeblesianus ssp. fickeiseniae	Wherever found	\$6,000		\$6,000
		Cactus, Knowlton's	Pediocactus knowltonii	Wherever found	\$1,000		\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Kuenzler hedgehog	Echinocereus fendleri var. kuenzleri	Wherever found	\$83,000		\$83,000
		Cactus, Mesa Verde	Sclerocactus mesae-verdae	Wherever found	\$136,000		\$136,000
		cactus, Pariette	Sclerocactus brevispinus	Wherever found	\$35,122		\$35,122
		Cactus, Pima pineapple	Coryphantha scheeri var. robustispina	Wherever found	\$5,600		\$5,600
		Cactus, San Rafael	Pediocactus despainii	Wherever found	\$25,063		\$25,063
		Cactus, Siler pincushion	Pediocactus (=Echinocactus,=Utahia) sileri	Wherever found	\$9,296		\$9,296
		cactus, Uinta Basin hookless	Sclerocactus wetlandicus	Wherever found	\$38,191		\$38,191
		Cactus, Winkler	Pediocactus winkleri	Wherever found	\$4,220		\$4,220
		Cactus, Wright fishhook	Sclerocactus wrightiae	Wherever found	\$18,197		\$18,197
		Catchfly, Spalding's	Silene spaldingii	Wherever found	\$23,500		\$23,500
		Ceanothus, Pine Hill	Ceanothus roderickii	Wherever found	\$5,000		\$5,000
		Centaury, spring-loving	Centaureum namophilum	Wherever found	\$3,000		\$3,000
		Checker-mallow, Nelson's	Sidalcea nelsoniana	Wherever found	\$28,000		\$28,000
		Cliffrose, Arizona	Purshia (=Cowania) subintegra	Wherever found	\$8,700		\$8,700
		Cycladenia, Jones	Cycladenia humilis var. jonesii	Wherever found	\$28,028		\$28,028
		Daisy, Willamette	Erigeron decumbens	Wherever found	\$23,000		\$23,000
		eryngo, Arizona	Eryngium sparganophyllum	(blank)	\$4,120		\$4,120
		Flannelbush, Pine Hill	Fremontodendron californicum ssp. decumbens	Wherever found	\$75,000		\$75,000
		Fleabane, Zuni	Erigeron rhizomatus	Wherever found	\$8,400		\$8,400
		Four-o'clock, MacFarlane's	Mirabilis macfarlanei	Wherever found	\$108,000		\$108,000
		Gumplant, Ash Meadows	Grindelia fraxinipratensis	Wherever found	\$30,000		\$30,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ivesia, Ash Meadows	Ivesia kingii var. eremica	Wherever found	\$30,000		\$30,000
		ivesia, Webber's	Ivesia webberi	Wherever found	\$29,303		\$29,303
		Jewelflower, California	Caulanthus californicus	Wherever found	\$46,991		\$46,991
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$199,723		\$199,723
		Lily, Minnesota dwarf trout	Erythronium propullans	Wherever found	\$38		\$38
		Lily, Western	Lilium occidentale	Wherever found	\$29,000		\$29,000
		Locoweed, Fassett's	Oxytropis campestris var. chartacea	Wherever found	\$76		\$76
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$192,710		\$192,710
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$4,549		\$4,549
		Manzanita, Ione	Arctostaphylos myrtifolia	Wherever found	\$1,500		\$1,500
		Milk-vetch, Ash meadows	Astragalus phoenix	Wherever found	\$30,000		\$30,000
		Milk-vetch, Fish Slough	Astragalus lentiginosus var. piscinensis	Wherever found	\$46,000		\$46,000
		Milk-vetch, Holmgren	Astragalus holmgreniorum	Wherever found	\$13,197		\$13,197
		Milk-vetch, Mancos	Astragalus humillimus	Wherever found	\$101,000		\$101,000
		milkvetch, Osterhout	Astragalus osterhoutii	Wherever found	\$40,000		\$40,000
		Milk-vetch, Peirson's	Astragalus magdalenae var. peirsonii	Wherever found	\$40,000		\$40,000
		Milk-vetch, Shivwits	Astragalus ampullarioides	Wherever found	\$24,120		\$24,120
		Milkweed, Mead's	Asclepias meadii	Wherever found	\$114		\$114
		Milkweed, Welsh's	Asclepias welshii	Wherever found	\$43,500		\$43,500
		Monkshood, northern wild	Aconitum noveboracense	Wherever found	\$38		\$38
		Morning-glory, Stebbins'	Calystegia stebbinsii	Wherever found	\$5,800		\$5,800
		Mustard, Penland alpine fen	Eutrema penlandii	Wherever found	\$5,500		\$5,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Niterwort, Amargosa	Nitrophila mohavensis	Wherever found	\$44,000		\$44,000
		Orchid, eastern prairie fringed	Platanthera leucophaea	Wherever found	\$38		\$38
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$3,300		\$3,300
		Pawpaw, four-petal	Asimina tetramera	(blank)	\$495		\$495
		Pennyroyal, Todsens	Hedeoma todsenii	Wherever found	\$10,750		\$10,750
		Penstemon, blowout	Penstemon haydenii	Wherever found	\$84,500		\$84,500
		Peppergrass, Slickspot	Lepidium papilliferum	(blank)	\$449,750		\$449,750
		Phacelia, clay	Phacelia argillacea	Wherever found	\$2,060		\$2,060
		Phacelia, DeBeque	Phacelia submutica	Wherever found	\$30,000	\$6,000	\$36,000
		Phacelia, North Park	Phacelia formosula	Wherever found	\$40,000		\$40,000
		popcornflower, rough	Plagiobothrys hirtus	Wherever found	\$43,156		\$43,156
		Reed-mustard, Barneby	Schoenocrambe barnebyi	Wherever found	\$3,237		\$3,237
		Reed-mustard, clay	Schoenocrambe argillacea	Wherever found	\$691		\$691
		Reed-mustard, shrubby	Schoenocrambe suffrutescens	Wherever found	\$5,486		\$5,486
		roseroot, Leedy's	Rhodiola integrifolia ssp. leedyi	Wherever found	\$38		\$38
		Sedge, Navajo	Carex specuicola	Wherever found	\$8,352		\$8,352
		Skullcap, Florida	Scutellaria floridana	(blank)	\$330		\$330
		Sneezeweed, Virginia	Helenium virginicum	Wherever found	\$76		\$76
		Sunflower, Pecos (=puzzle, =paradox)	Helianthus paradoxus	Wherever found	\$9,250		\$9,250
		Sunray, Ash Meadows	Enceliopsis nudicaulis var. corrugata	Wherever found	\$30,000		\$30,000
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$38		\$38
		Townsendia, Last Chance	Townsendia aprica	Wherever found	\$2,639		\$2,639

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Twinpod, Dudley Bluffs	Physaria obcordata	Wherever found	\$5,000		\$5,000
		Vervain, Red Hills	Verbena californica	Wherever found	\$800		\$800
		wild buckwheat, clay-loving	Eriogonum pelinophilum	Wherever found	\$15,585		\$15,585
		Wild-buckwheat, gypsum	Eriogonum gypsophilum	Wherever found	\$95,000		\$95,000
		Wooly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$202,000		\$202,000
		Yellowhead, desert	Yermo xanthocephalus	Wherever found	\$15,500		\$15,500
	<i>Flowering Plants subtotal</i>				\$2,946,785	\$6,000	\$2,952,785
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$330		\$330
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$76		\$76
		Butterfly, Behren's silverspot	Speyeria zerene behrensii	Wherever found	\$19,200		\$19,200
		Butterfly, Fender's blue	Icaricia icarioides fenderi	Wherever found	\$66,175		\$66,175
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$76		\$76
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$76		\$76
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$12,455		\$12,455
		Butterfly, Uncompahgre fritillary	Boloria acrocneuma	Wherever found	\$12,100		\$12,100
		Checkerspot, Taylor's (=whulge)	Euphydryas editha taylori	Wherever found	\$2,999		\$2,999
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$647		\$647
		Skipper, Carson wandering	Pseudocopaeodes eunus obscurus	Wherever found	\$22,507		\$22,507
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$42,038		\$42,038

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Skipper, Pawnee montane	Hesperia leonardus montana	Wherever found	\$100		\$100
		skipperling, Poweshiek	Oarisma poweshiek	Wherever found	\$76		\$76
	<i>Insects subtotal</i>				<i>\$178,855</i>		<i>\$178,855</i>
	Lichens	Cladonia, Florida perforate	Cladonia perforata	(blank)	\$495		\$495
	<i>Lichens subtotal</i>				<i>\$495</i>		<i>\$495</i>
	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$495		\$495
		Bat, gray	Myotis grisescens	Wherever found	\$457		\$457
		Bat, Indiana	Myotis sodalis	Wherever found	\$2,589		\$2,589
		Bat, Mexican long-nosed	Leptonycteris nivalis	Wherever found	\$33,745		\$33,745
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$134,642		\$134,642
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$324,651		\$324,651
		Bear, polar	Ursus maritimus	Wherever found	\$330,000		\$330,000
		Bison, wood	Bison bison athabascæ	Wherever found	\$20,000		\$20,000
		Deer, Columbian white-tailed	Odocoileus virginianus leucurus	Columbia River (Clark, Cowlitz, Pacific, Skamania, and Wahkiakum Counties, WA., and Clatsop, Columbia, and Multnomah Counties, OR.)	\$119,384		\$119,384
		Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$214,100		\$214,100
		Fisher	Pekania pennanti	West Coast Distinct Population Segment	\$5,240		\$5,240
		Jaguar	Panthera onca	Wherever found	\$12,310		\$12,310
		Kangaroo rat, Tipton	Dipodomys nitratoideus nitratoideus	Wherever found	\$2,560		\$2,560

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$483,772		\$483,772
		Manatee, West Indian	Trichechus manatus	Wherever found	\$1,320		\$1,320
		Marten, Pacific	Martes caurina	Wherever found	\$120,455		\$120,455
		Mountain beaver, Point Arena	Aplodontia rufa nigra	Wherever found	\$1,600		\$1,600
		Mouse, Alabama beach	Peromyscus polionotus ammobates	Wherever found	\$990		\$990
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$2,100		\$2,100
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	Wherever found	\$3,600		\$3,600
		Mouse, southeastern beach	Peromyscus polionotus niveiventris	Wherever found	\$330		\$330
		Ocelot	Leopardus (=Felis) pardalis	Wherever found	\$110		\$110
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$425,395		\$425,395
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	In Arizona, an area north of Interstate 8 and south of Interstate 10, bounded by the Colorado River on the west and Interstate 10 on the east; and an area south of Interstate 8, bounded by Highway 85 on the west, Interstates 10 and 19 on the east, and the U.S.-Mexico border on the south	\$14,850		\$14,850
				Wherever found, except where listed as an experimental population	\$130,000		\$130,000
		Rabbit, Pygmy	Brachylagus idahoensis	U.S.A. (WA-Douglas, Grant, Lincoln, Adams, Benton Counties)	\$11,000		\$11,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Seal, Ringed	Phoca (=Pusa) hispida hispida	Wherever found	\$120,000		\$120,000
		Sheep, Sierra Nevada bighorn	Ovis canadensis sierrae	U.S.A. (CA) Sierra Nevada	\$10,000		\$10,000
		Vole, Amargosa	Microtus californicus scirpensis	Wherever found	\$382,000		\$382,000
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of	\$3,315		\$3,315

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$150,710		\$150,710
	<i>Mammals subtotal</i>				<i>\$3,061,720</i>		<i>\$3,061,720</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$990		\$990
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$670		\$670
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$74,782		\$74,782
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$1,000		\$1,000
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$114		\$114
		piresnake, black	Pituophis melanoleucus lodingi	Wherever found	\$330		\$330
		Rattlesnake, New Mexican ridge-nosed	Crotalus willardi obscurus	Wherever found	\$903		\$903

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, green	Chelonia mydas	Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	\$990		\$990
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$660		\$660
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,320		\$1,320
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$660		\$660
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	\$1,320		\$1,320
		Snake, copperbelly water	Nerodia erythrogaster neglecta	Indiana north of 40 degrees north latitude, Michigan, Ohio	\$76		\$76
		Snake, eastern indigo	Drymarchon couperi	Wherever found	\$1,155		\$1,155
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$1,124,399	\$2,511,000	\$3,635,399

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tortoise, gopher	Gopherus polyphemus	Western DPS	\$825		\$825
		Turtle, Alabama red-bellied	Pseudemys alabamensis	Wherever found	\$660		\$660
		Turtle, yellow-blotched map	Gratemys flavimaculata	Wherever found	\$330		\$330
	<i>Reptiles subtotal</i>				<i>\$1,211,184</i>	<i>\$2,511,000</i>	<i>\$3,722,184</i>
	Snails	Limpet, Banbury Springs	Lanx sp.	Wherever found	\$5,700		\$5,700
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$8,600		\$8,600
		Snail, Pecos assiminea	Assiminea pecos	Wherever found	\$2,000		\$2,000
		Springsnail, Bruneau Hot	Pyrgulopsis bruneauensis	Wherever found	\$5,300		\$5,300
		Springsnail, Koster's	Juturnia kosteri	Wherever found	\$2,000		\$2,000
		Springsnail, Roswell	Pyrgulopsis roswellensis	Wherever found	\$2,000		\$2,000
	<i>Snails subtotal</i>				<i>\$25,600</i>		<i>\$25,600</i>
Domestic subtotal					\$22,003,602	\$2,517,000	\$24,520,602
Grand Total					\$22,003,602	\$2,517,000	\$24,520,602

DOI National Park Service - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$105,100		\$105,100
		Frog, mountain yellow-legged	Rana muscosa	U.S.A., northern California	\$150,907		\$150,907
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$215		\$215
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$325,907		\$325,907
		Hellbender, Ozark	Cryptobranchus alleganiensis bishopi	Wherever found	\$15,400		\$15,400
		Salamander, Jemez Mountains	Plethodon neomexicanus	Wherever found	\$96,000		\$96,000
		Salamander, Shenandoah	Plethodon shenandoah	Wherever found	\$60,000		\$60,000
		Yosemite Toad	Anaxyrus canorus	Wherever found	\$75,591		\$75,591
	<i>Amphibians subtotal</i>				<i>\$829,120</i>		<i>\$829,120</i>
	Birds	(honeyeater), Mao (= maomao)	Gymnomyza samoensis	Wherever found	\$500		\$500
		'I'iwi	Drepanis coccinea	Wherever found	\$2,409,875		\$2,409,875
		Akohekohe	Palmeria dolei	Wherever found	\$2,474,560		\$2,474,560
		Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)- see 17.84(j)	\$22,100		\$22,100
				U.S.A. only, except where listed as an experimental population	\$508,320		\$508,320
		coot, Hawaiian	Fulica americana alai	Wherever found	\$25,000		\$25,000
		Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$20,200		\$20,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	\$19,514		\$19,514
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	Wherever found, except where listed as an experimental population	\$250		\$250
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$4,900		\$4,900
		Goose, Hawaiian	<i>Branta</i> (=Nesochen) <i>sandvicensis</i>	Wherever found	\$267,239		\$267,239
		Ground-Dove, Friendly	<i>Gallicolumba stairi</i>	American Samoa DPS	\$500		\$500
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$5,000		\$5,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$30,608		\$30,608
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$222,136		\$222,136
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$180,496		\$180,496
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$587,301		\$587,301
		parrotbill, Maui (Kiwikiu)	<i>Pseudonestor xanthophrys</i>	Wherever found	\$2,474,560		\$2,474,560
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$370,902		\$370,902
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$935,442		\$935,442

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$35,700		\$35,700
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$112,000		\$112,000
		rail, Eastern Black	Laterallus jamaicensis ssp. jamaicensis	Wherever found	\$31,000		\$31,000
		Rail, Yuma Ridgway's	Rallus obsoletus yumanensis	Wherever found	\$5,600		\$5,600
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$78,720		\$78,720
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$1,000		\$1,000
		Sparrow, Cape Sable seaside	Ammodramus maritimus mirabilis	Wherever found	\$300,000		\$300,000
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$25,000		\$25,000
		Stork, wood	Mycteria americana	U.S.A. (AL, FL, GA, MS, NC, SC)	\$40,550		\$40,550
		Storm-petrel, band-rumped	Oceanodroma castro	USA (HI)	\$40,825		\$40,825
		Tern, Roseate	Sterna dougallii dougallii	U.S.A. (Atlantic Coast south to NC), Canada (Newf., N.S, Que.), Bermuda	\$10,000		\$10,000
				Western Hemisphere and adjacent oceans, incl. U.S.A. (FL, PR, VI), where not listed as endangered	\$5,000		\$5,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$500		\$500
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$125,900		\$125,900
		<i>Birds subtotal</i>			<i>\$11,371,198</i>		<i>\$11,371,198</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Clams	Bean, Cumberland (pearlymussel)	Villosa trabalis	Wherever found; Except where listed as Experimental Populations	\$2,450		\$2,450
		Bean, rayed	Villosa fabalis	Wherever found	\$1,000		\$1,000
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$1,000		\$1,000
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$2,450		\$2,450
		Elktoe, Cumberland	Alasmodonta atropurpurea	Wherever found	\$2,450		\$2,450
		Fanshell	Cyprogenia stegaria	Wherever found	\$1,000		\$1,000
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$2,000		\$2,000
		Kidneyshell, fluted	Ptychobranhus subtentus	Wherever found	\$2,450		\$2,450
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$9,300		\$9,300
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$3,450		\$3,450
		Mussel, oyster	Epioblasma capsaeformis	Wherever found; Except where listed as Experimental Populations	\$2,450		\$2,450
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$1,000		\$1,000
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$17,400		\$17,400
		No common name	Fusconaia subrotunda	(blank)	\$100		\$100
			Obovaria subrotunda	(blank)	\$100		\$100
		Pearlymussel, dromedary	Dromus dromas	Wherever found; Except where listed as Experimental Populations	\$2,450		\$2,450
		Pearlymussel, littlewing	Pegias fabula	Wherever found	\$2,450		\$2,450
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$1,000		\$1,000
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$16,000		\$16,000
		Riffleshell, northern	Epioblasma rangiana	Wherever found	\$2,000		\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Riffleshell, tan	Epioblasma florentina walkeri (=E. walkeri)	Wherever found	\$2,450		\$2,450
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$1,000		\$1,000
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$5,450		\$5,450
	<i>Clams subtotal</i>				<i>\$81,400</i>		<i>\$81,400</i>
	Conifers and Cycads	No common name	Pinus albicaulis	(blank)	\$602,692		\$602,692
	<i>Conifers and Cycads subtotal</i>				<i>\$602,692</i>		<i>\$602,692</i>
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$95,000		\$95,000
		Coral, elkhorn	Acropora palmata	Wherever found	\$222,000		\$222,000
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$144,500		\$144,500
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$202,500		\$202,500
		coral, Pillar	Dendrogyra cylindrus	Wherever found	\$102,500		\$102,500
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$20,000		\$20,000
		Coral, staghorn	Acropora cervicornis	Wherever found	\$215,500		\$215,500
	<i>Corals subtotal</i>				<i>\$1,002,000</i>		<i>\$1,002,000</i>
	Crustaceans	Shrimp, anchialine pool	Procaris hawaiana	Wherever found	\$7,204		\$7,204
		Shrimp, Kentucky cave	Palaemonias ganteri	Wherever found	\$1,000		\$1,000
	<i>Crustaceans subtotal</i>				<i>\$8,204</i>		<i>\$8,204</i>
	Ferns and Allies	Kupukupu makalii	Cyclosorus boydiae	Wherever found	\$10,241		\$10,241
		No common name	Asplenium peruvianum var. insulare	Wherever found	\$10,241		\$10,241
			Microlepia strigosa var. mauensis	Wherever found	\$10,241		\$10,241
		Wawae`iole	Huperzia mannii	Wherever found	\$10,241		\$10,241
	<i>Ferns and Allies subtotal</i>				<i>\$40,964</i>		<i>\$40,964</i>
	Fishes	Chub, humpback	Gila cypha	Wherever found	\$577,477		\$577,477
		Darter, candy	Etheostoma osburni	Wherever found	\$1,500		\$1,500
		Darter, diamond	Crystallaria cincotta	Wherever found	\$200		\$200
		Darter, duskytail	Etheostoma percnurum	Wherever found	\$10,012		\$10,012
		Darter, slackwater	Etheostoma boschungii	Wherever found	\$360		\$360
		Gambusia, Big Bend	Gambusia gaigei	Wherever found	\$2,000		\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$16,000		\$16,000
		grouper, Nassau	Epinephelus striatus	(blank)	\$48,500		\$48,500
		Madtom, smoky	Noturus baileyi	Wherever found	\$10,012		\$10,012
		Madtom, yellowfin	Noturus flavipinnis	Wherever found, except where listed as an experimental population	\$10,012		\$10,012
		Minnow, loach	Tiaroga cobitis	Wherever found	\$314		\$314
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$151,132		\$151,132
		Pupfish, Devils Hole	Cyprinodon diabolis	Wherever found	\$500,000		\$500,000
		salmon Atlantic	Salmo salar	Gulf of Maine DPS - See 50 CFR 224.101	\$400,000		\$400,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU - See 50 CFR 223.102	\$6,000		\$6,000
				Central Valley spring-run ESU See 50 CFR 223.102	\$1,500		\$1,500
				Puget Sound ESU See 50 CFR 223.102	\$252,500		\$252,500
				Snake River fall-run ESU See 50 CFR 223.102	\$10,000		\$10,000
				Snake River spring/summer-run ESU See 50 CFR 223.102	\$10,000		\$10,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU See 50 CFR 223.102	\$10,000		\$10,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU - See 50 CFR 224.101	\$440,000		\$440,000
				Lower Columbia River ESU - See 50 CFR 223.102	\$10,000		\$10,000
				Southern Oregon - Northern California Coast ESU - See 50 CFR 223.102	\$9,500		\$9,500
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU - See 50 CFR 223.102	\$30,000		\$30,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$6,000		\$6,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$8,200		\$8,200
		Spikedace	Meda fulgida	Wherever found	\$114		\$114
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS - See 50 CFR 223.102	\$6,500		\$6,500
				Central California Coast DPS - See 50 CFR 223.102	\$110,000		\$110,000
				Northern California DPS - See 50 CFR 223.102	\$6,000		\$6,000
				Puget Sound DPS - See 50 CFR 223.102	\$250,000		\$250,000
				Upper Willamette River DPS - See 50 CFR 223.102	\$10,000		\$10,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS - See 50 CFR 224.101	\$2,064		\$2,064
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$520		\$520
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$6,500		\$6,500
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	U.S. Only	\$300		\$300
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, (lower 48 states)	\$447,853		\$447,853
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$52,900		\$52,900
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$20,000		\$20,000
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$72,585		\$72,585
	<i>Fishes subtotal</i>				\$3,506,555		\$3,506,555
	Flowering Plants	`Ahinahina	Argyroxiphium sandwicense ssp. macrocephalum	Wherever found	\$102,410		\$102,410
		`Aiea	Nothocestrum latifolium	Wherever found	\$10,241		\$10,241
		`aku	Cyanea tritomantha	Wherever found	\$1,803		\$1,803
		`Anunu	Sicyos albus	Wherever found	\$1,803		\$1,803

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
			Sicyos macrophyllus	Wherever found	\$1,803		\$1,803
		`Oha wai	Clermontia lindseyana	Wherever found	\$1,803		\$1,803
			Clermontia peleana	Wherever found	\$1,803		\$1,803
			Clermontia samuelii	Wherever found	\$51,208		\$51,208
		`Ohe	Joinvillea ascendens ascendens	Wherever found	\$20,483		\$20,483
		Alani	Melicope ovalis	Wherever found	\$20,483		\$20,483
		Alopecurus, Sonoma	Alopecurus aequalis var. sonomensis	Wherever found	\$250		\$250
		Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$71,094		\$71,094
		Avens, spreading	Geum radiatum	Wherever found	\$3,350		\$3,350
		Bedstraw, island	Galium buxifolium	Wherever found	\$5,000		\$5,000
		bladderpod, Missouri	Physaria filiformis	Wherever found	\$50,000		\$50,000
		Blazingstar, Heller's	Liatris helleri	Wherever found	\$4,750		\$4,750
		Bully, Everglades	Sideroxylon reclinatum ssp. austrofloridense	(blank)	\$14,562		\$14,562
		Bush-mallow, Santa Cruz Island	Malacothamnus fasciculatus var. nesioticus	Wherever found	\$3,000		\$3,000
		Cactus, Acuña	Echinomastus erectocentrus var. acunensis	Wherever found	\$1,850		\$1,850
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$500		\$500
		Cactus, Florida semaphore	Consolea corallicola	(blank)	\$2,000		\$2,000
		Cactus, Lloyd's Mariposa	Sclerocactus mariposensis	Wherever found	\$1,200		\$1,200
		Cactus, Winkler	Pediocactus winkleri	Wherever found	\$25,000		\$25,000
		Cactus, Wright fishhook	Sclerocactus wrightiae	Wherever found	\$21,000		\$21,000
		Clarkia, Presidio	Clarkia franciscana	Wherever found	\$800		\$800
		Clover, showy Indian	Trifolium amoenum	Wherever found	\$200		\$200
		Cory cactus, bunched	Coryphantha ramillosa	Wherever found	\$3,000		\$3,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crabgrass, Florida pineland	<i>Digitaria pauciflora</i>	(blank)	\$10,562		\$10,562
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$2,000		\$2,000
		Dudleya, marcescent	<i>Dudleya cymosa</i> ssp. <i>marcescens</i>	Wherever found	\$300		\$300
		Dudleya, Santa Cruz Island	<i>Dudleya nesiotica</i>	Wherever found	\$10,000		\$10,000
		Dudleya, Verity's	<i>Dudleya verityi</i>	Wherever found	\$1,500		\$1,500
		Dwarf-flax, Marin	<i>Hesperolinon congestum</i>	Wherever found	\$900		\$900
		Fescue, Guadalupe	<i>Festuca ligulata</i>	Wherever found	\$1,000		\$1,000
		Fringepod, Santa Cruz Island	<i>Thysanocarpus conchuliferus</i>	Wherever found	\$400		\$400
		Gilia, Hoffmann's slender-flowered	<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>	Wherever found	\$5,000		\$5,000
		Grass, Eureka Dune	<i>Swallenia alexandrae</i>	(blank)	\$15,000		\$15,000
		Ha'iwale	<i>Cyrtandra giffardii</i>	Wherever found	\$1,803		\$1,803
		haha	<i>Cyanea asplenifolia</i>	Wherever found	\$30,724		\$30,724
			<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>	Wherever found	\$41,966		\$41,966
			<i>Cyanea glabra</i>	Wherever found	\$10,241		\$10,241
			<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>	Wherever found	\$20,483		\$20,483
			<i>Cyanea kunthiana</i>	Wherever found	\$20,483		\$20,483
			<i>Cyanea maritae</i>	Wherever found	\$10,241		\$10,241
			<i>Cyanea shipmanii</i>	Wherever found	\$1,803		\$1,803
			<i>Cyanea stictophylla</i>	Wherever found	\$1,803		\$1,803
		Haiwale	<i>Cyrtandra ferripilosa</i>	Wherever found	\$30,724		\$30,724
		Hala pepe	<i>Pleomele hawaiiensis</i>	Wherever found	\$1,803		\$1,803
		Hau kuahiwi	<i>Hibiscadelphus giffardianus</i>	Wherever found	\$1,803		\$1,803
		Heau	<i>Exocarpos menziesii</i>	(blank)	\$1,803		\$1,803
		Holei	<i>Ochrosia haleakalae</i>	Wherever found	\$1,803		\$1,803
		Honohono	<i>Haplostachys haplostachya</i>	Wherever found	\$1,803		\$1,803
		Ischaemum, Hilo	<i>Ischaemum byrone</i>	Wherever found	\$10,241		\$10,241
		Ko'oko'olau	<i>Bidens campylotheca</i> ssp. <i>pentamera</i>	Wherever found	\$30,724		\$30,724

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
			<i>Bidens campylotheca</i> ssp. <i>waihoiensis</i>	Wherever found	\$30,724		\$30,724
			<i>Bidens micrantha</i> ssp. <i>ctenophylla</i>	Wherever found	\$409		\$409
			<i>Bidens micrantha</i> ssp. <i>kalealaha</i>	Wherever found	\$30,724		\$30,724
		Kuahiwi laukahi	<i>Plantago hawaiiensis</i>	Wherever found	\$1,803		\$1,803
			<i>Plantago princeps</i>	Wherever found	\$30,724		\$30,724
		Layia, beach	<i>Layia carnosa</i>	Wherever found	\$2,000		\$2,000
		Lessingia, San Francisco	<i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i>)	Wherever found	\$49,000		\$49,000
		Liveforever, Santa Barbara Island	<i>Dudleya traskiae</i>	Wherever found	\$30,000		\$30,000
		Lupine, clover (Tidestrom"s)	<i>Lupinus tidestromii</i>	Wherever found	\$1,000		\$1,000
		Malacothrix, island	<i>Malacothrix squalida</i>	Wherever found	\$5,000		\$5,000
		Malacothrix, Santa Cruz Island	<i>Malacothrix indecora</i>	Wherever found	\$5,000		\$5,000
		Manzanita, Franciscan	<i>Arctostaphylos franciscana</i>	Wherever found	\$900		\$900
		Manzanita, Presidio	<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>	Wherever found	\$700		\$700
		Manzanita, Santa Rosa Island	<i>Arctostaphylos confertiflora</i>	Wherever found	\$5,000		\$5,000
		Milk-vetch, Braunton's	<i>Astragalus brauntonii</i>	Wherever found	\$1,000		\$1,000
		Milk-vetch, Shivwits	<i>Astragalus ampullarioides</i>	Wherever found	\$2,230		\$2,230
		Milk-vetch, triple-ribbed	<i>Astragalus tricarinatus</i>	Wherever found	\$2,200		\$2,200
		monkey-flower, Michigan	<i>Mimulus michiganensis</i>	Wherever found	\$1,200		\$1,200
		Nanu	<i>Gardenia remyi</i>	Wherever found	\$10,241		\$10,241
		No common name	<i>Calyptranthes thomasiana</i>	Wherever found	\$3,000		\$3,000
			<i>Neraudia ovata</i>	Wherever found	\$1,803		\$1,803
			<i>Phyllostegia bracteata</i>	Wherever found	\$41,966		\$41,966

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
			<i>Phyllostegia brevidens</i>	Wherever found	\$30,724		\$30,724
			<i>Phyllostegia haliakalae</i>	Wherever found	\$41,966		\$41,966
			<i>Pritchardia maideniana</i>	Wherever found	\$2,046		\$2,046
			<i>Sanicula sandwicensis</i>	Wherever found	\$30,724		\$30,724
			<i>Schiedea diffusa</i> ssp. <i>macraei</i>	Wherever found	\$1,803		\$1,803
			<i>Schiedea diffusa</i> subsp. <i>diffusa</i>	Wherever found	\$30,724		\$30,724
			<i>Schiedea haleakalensis</i>	Wherever found	\$30,724		\$30,724
			<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>	Wherever found	\$1,803		\$1,803
			<i>Wikstroemia villosa</i>	Wherever found	\$10,241		\$10,241
		Nohoanu	<i>Geranium arboreum</i>	Wherever found	\$30,724		\$30,724
			<i>Geranium hanaense</i>	Wherever found	\$30,724		\$30,724
			<i>Geranium multiflorum</i>	Wherever found	\$10,241		\$10,241
		nui, haha	<i>Cyanea horrida</i>	Wherever found	\$10,241		\$10,241
		Ohai	<i>Sesbania tomentosa</i>	Wherever found	\$1,803		\$1,803
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$10,000		\$10,000
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$4,000		\$4,000
		Paintbrush, soft-leaved	<i>Castilleja mollis</i>	Wherever found	\$20,000		\$20,000
		Paintbrush, Tiburon	<i>Castilleja affinis</i> ssp. <i>neglecta</i>	Wherever found	\$500		\$500
		Pentachaeta, Lyon's	<i>Pentachaeta lyonii</i>	Wherever found	\$7,500		\$7,500
		Phacelia, island	<i>Phacelia insularis</i> ssp. <i>insularis</i>	Wherever found	\$7,000		\$7,000
		Phlox, Texas trailing	<i>Phlox nivalis</i> ssp. <i>texensis</i>	Wherever found	\$2,100		\$2,100
		Pink, swamp	<i>Helonias bullata</i>	(blank)	\$4,750		\$4,750
		Po'e	<i>Portulaca sclerocarpa</i>	Wherever found	\$1,803		\$1,803
		Pogonia, small whorled	<i>Isotria medeoloides</i>	(blank)	\$38,000		\$38,000
		potato-bean, Price's	<i>Apios priceana</i>	(blank)	\$6,000		\$6,000
		Potentilla, Hickman's	<i>Potentilla hickmanii</i>	Wherever found	\$1,600		\$1,600

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Prairie-clover, Florida	Dalea carthagenensis floridana	(blank)	\$2,000		\$2,000
		Prickly-ash, St. Thomas	Zanthoxylum thomasianum	Wherever found	\$3,000		\$3,000
		Reedgrass, Maui	Calamagrostis expansa	Wherever found	\$10,241		\$10,241
		Reed-mustard, Barneby	Schoenocrambe barnebyi	Wherever found	\$5,000		\$5,000
		Rock-cress, Hoffmann's	Arabis hoffmannii	Wherever found	\$15,000		\$15,000
		Rosemary, Cumberland	Conradina verticillata	Wherever found	\$9,200		\$9,200
		Sandwort, Marsh	Arenaria paludicola	Wherever found	\$400		\$400
		Sedge, Navajo	Carex specuicola	Wherever found	\$4,781		\$4,781
		Silversword, Mauna Loa (=Ka'u)	Argyroxiphium kauense	Wherever found	\$45,000		\$45,000
		Spineflower, Sonoma	Chorizanthe valida	Wherever found	\$1,000		\$1,000
		Spiraea, Virginia	Spiraea virginiana	Wherever found	\$54,366		\$54,366
		stonecrop, Bartram's	Graptopetalum bartramii	Wherever found	\$5,000		\$5,000
		Sunflower, Whorled	Helianthus verticillatus	(blank)	\$1,200		\$1,200
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$20,400		\$20,400
		Thoroughwort, Cape Sable	Chromolaena frustrata	(blank)	\$1,500		\$1,500
		Townsendia, Last Chance	Townsendia aprica	Wherever found	\$3,000		\$3,000
	<i>Flowering Plants subtotal</i>				<i>\$1,493,762</i>		<i>\$1,493,762</i>
	Insects	bumble bee, Franklin's	Bombus franklini	Wherever found	\$1,388		\$1,388
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$1,000		\$1,000
		Butterfly, Bartram's hairstreak	Strymon acis bartrami	Wherever found	\$5,500		\$5,500
		Butterfly, Florida leafwing	Anaea troglodyta floridalis	Wherever found	\$7,000		\$7,000
		Butterfly, Island marble	Euchloe ausonides insulanus	Wherever found	\$73,000		\$73,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly, Karner blue	<i>Lycaeides melissa samuelis</i>	Wherever found	\$800		\$800
		Butterfly, mission blue	<i>Icaricia icarioides missionensis</i>	Wherever found	\$175,000		\$175,000
		Butterfly, San Bruno elfin	<i>Callophrys mossii bayensis</i>	Wherever found	\$7,500		\$7,500
		Butterfly, Schaus swallowtail	<i>Heraclides aristodemus ponceanus</i>	Wherever found	\$55,000		\$55,000
		Damselfly, orangeblack Hawaiian	<i>Megalagrion xanthomelas</i>	Wherever found	\$6,609		\$6,609
		Damselfly, Pacific Hawaiian	<i>Megalagrion pacificum</i>	Wherever found	\$40,825		\$40,825
		Stonefly, western glacier	<i>Zapada glacier</i>	Wherever found	\$9,750		\$9,750
		Tiger beetle, Northeastern beach	<i>Habroscelimorpha dorsalis dorsalis</i>	Wherever found	\$1,528		\$1,528
	<i>Insects subtotal</i>				<i>\$384,900</i>		<i>\$384,900</i>
	Mammals	bat, Florida bonneted	<i>Eumops floridanus</i>	Wherever found	\$46,000		\$46,000
		Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$75,030		\$75,030
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$40,825		\$40,825
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$126,332		\$126,332
		Bat, Mexican long-nosed	<i>Leptonycteris nivalis</i>	Wherever found	\$2,796		\$2,796
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$924,785		\$924,785
		Bat, Ozark big-eared	<i>Corynorhinus (=Plecotus) townsendii ingens</i>	Wherever found	\$5,000		\$5,000
		Bat, Virginia big-eared	<i>Corynorhinus (=Plecotus) townsendii virginianus</i>	Wherever found	\$6,400		\$6,400
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$2,195,041		\$2,195,041

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ferret, black-footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$264,426		\$264,426
				Wherever found, except where listed as an experimental population	\$75,000		\$75,000
		Fisher	<i>Pekania pennanti</i>	Southern Sierra Nevada Distinct Population Segment	\$309,740		\$309,740
		Fox, Sierra Nevada red	<i>Vulpes vulpes necator</i>	Sierra Nevada DPS	\$98,664		\$98,664
		Lynx, Canada	<i>Lynx canadensis</i>	Wherever Found in Contiguous U.S.	\$24,300		\$24,300
		Manatee, West Indian	<i>Trichechus manatus</i>	Wherever found	\$25,000		\$25,000
		Marten, Pacific	<i>Martes caurina</i>	Wherever found	\$82,350		\$82,350
		Mouse, Anastasia Island beach	<i>Peromyscus polionotus phasma</i>	Wherever found	\$1,000		\$1,000
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$55,000		\$55,000
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$5,625		\$5,625
		Mouse, southeastern beach	<i>Peromyscus polionotus niveiventris</i>	Wherever found	\$5,000		\$5,000
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$37,500		\$37,500
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$125,786		\$125,786
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$18,453		\$18,453
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$10,000		\$10,000
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	U.S.A. (CA) Sierra Nevada	\$83,080		\$83,080
		Squirrel, Carolina northern flying	<i>Glaucomys sabrinus coloratus</i>	Wherever found	\$5,055		\$5,055

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$1,000		\$1,000
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of	\$384,812		\$384,812
	<i>Mammals subtotal</i>				\$5,034,000		\$5,034,000
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$103,000		\$103,000
		Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$150,000		\$150,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$314		\$314
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$12,600		\$12,600
		Lizard, St. Croix ground	Ameiva polops	Wherever found	\$25,000		\$25,000
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$900		\$900
		Sea turtle, green	Chelonia mydas	Green sea turtles originating from the Central North Pacific Ocean, bounded by the following coordinates: 41 degrees N., 169 degrees E. in the northwest; 41 degrees N., 143 degrees W. in the northeast; 9 degrees N., 125 degrees W. in the southeast; and 9 degrees N., 175 degrees W. in the southwest Pacific coast of Mexico	\$5,234		\$5,234
				Green sea turtles originating from the Central South Pacific Ocean, bounded by the following coordinates: 9 degrees N., 175 degrees W. in the northwest; 9 degrees N., 125 degrees W. in the northeast; 40 degrees S., 96 degrees W. in the southeast; 40 degrees S., 176 degrees E. in the southwest; and 13 degrees S., 171 degrees E. in the west	\$750		\$750

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	\$498,055		\$498,055

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	\$12,000		\$12,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$182,020		\$182,020
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,012,481		\$1,012,481
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$10,500		\$10,500
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	\$840,513		\$840,513
		Snake, eastern indigo	Drymarchon couperi	Wherever found	\$1,500		\$1,500
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$10,000		\$10,000
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$190,568		\$190,568
		Tortoise, gopher	Gopherus polyphemus	Western DPS	\$655		\$655

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		turtle, bog	Glyptemys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$3,000		\$3,000
				Wherever found, except GA, NC, SC, TN, VA	\$13,820		\$13,820
		Turtle, Sonoyta mud	Kinosternon sonoriense longifemorale	Wherever found	\$146,896		\$146,896
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$6,000		\$6,000
	<i>Reptiles subtotal</i>				\$3,225,806		\$3,225,806
	Snails	[no common name] Snail	Eua zebrina	Wherever found	\$5,000		\$5,000
			Ostodes strigatus	Wherever found	\$5,000		\$5,000
		Abalone, Black	Haliotis cracherodii	Wherever found	\$3,000		\$3,000
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$3,000		\$3,000
	<i>Snails subtotal</i>				\$16,000		\$16,000
Domestic subtotal					\$27,596,601		\$27,596,601
Foreign	Corals	No common name	Acropora globiceps	Wherever found	\$1,500		\$1,500
			Acropora jacquelineae	Wherever found	\$500		\$500
			Acropora retusa	Wherever found	\$1,000		\$1,000
			Acropora speciosa	Wherever found	\$500		\$500
			Euphyllia paradivisa	Wherever found	\$500		\$500
			Isopora crateriformis	Wherever found	\$1,000		\$1,000
	<i>Corals subtotal</i>				\$5,000		\$5,000
Foreign subtotal					\$5,000		\$5,000
Grand Total					\$27,601,601		\$27,601,601

Federal Aviation Administration - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$300		\$300
		Waterdog, Neuse River	Necturus lewisi	Wherever found	\$3,200		\$3,200
	<i>Amphibians subtotal</i>				\$3,500		\$3,500
	Birds	Horned lark, streaked	Eremophila alpestris strigata	Wherever found	\$500		\$500
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$600		\$600
	<i>Birds subtotal</i>				\$1,100		\$1,100
	Clams	Pigtoe, Atlantic	Fusconaia masoni	Wherever found	\$3,200		\$3,200
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$3,200		\$3,200
	<i>Clams subtotal</i>				\$6,400		\$6,400
	Fishes	Madtom, Carolina	Noturus furiosus	Wherever found	\$3,200		\$3,200
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS - See 50 CFR 224.101	\$100		\$100
	<i>Fishes subtotal</i>				\$3,300		\$3,300
	Flowering Plants	Sumac, Michaux's	Rhus michauxii	Wherever found	\$400		\$400
	<i>Flowering Plants subtotal</i>				\$400		\$400
	Insects	Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$300		\$300
	<i>Insects subtotal</i>				\$300		\$300
	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$65,500		\$65,500
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$103,000		\$103,000
		Rabbit, riparian brush	Sylvilagus bachmani riparius	Wherever found	\$500		\$500
		Wolf, red	Canis rufus	Wherever found, except where listed as an experimental population	\$600		\$600
	<i>Mammals subtotal</i>				\$169,600		\$169,600
	Reptiles	Snake, giant garter	Thamnophis gigas	Wherever found	\$300		\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$75,600		\$75,600
	<i>Reptiles subtotal</i>				\$75,900		\$75,900
<i>Domestic subtotal</i>					<i>\$260,500</i>		<i>\$260,500</i>
Multi-species					\$42,000		\$42,000
<i>Multi-Species subtotal</i>					<i>\$42,000</i>		<i>\$42,000</i>
Grand Total					\$302,500		\$302,500

Nuclear Regulatory Commission - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$100		\$100
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$100		\$100
	<i>Amphibians subtotal</i>				\$200		\$200
	Birds	Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$100		\$100
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$100		\$100
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$400		\$400
				[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$600		\$600
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$100		\$100
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$100		\$100
		Tern, California least	Sterna antillarum browni	Wherever found	\$100		\$100
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$100		\$100
	<i>Birds subtotal</i>				\$1,600		\$1,600
	Clams	Pigtoe, finereyed	Fusconaia cuneolus	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$600		\$600

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, shiny	Fusconaia cor	U.S.A. (AL;The free-flowing reach of the Tennessee R. from the base of Wilson Dam downstream to the backwaters of Pickwick Reservoir [about 12 RM (19 km)] and the lower 5 RM [8 km] of all tributaries to this reach in Colbert and Lauderdale Cos., see 17.85(a))	\$600		\$600
	<i>Clams subtotal</i>				<i>\$1,200</i>		<i>\$1,200</i>
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$200		\$200
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$200		\$200
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$200		\$200
		coral, Pillar	Dendrogyra cylindrus	Wherever found	\$200		\$200
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$200		\$200
		Coral, staghorn	Acropora cervicornis	Wherever found	\$200		\$200
	<i>Corals subtotal</i>				<i>\$1,200</i>		<i>\$1,200</i>
	Crustaceans	Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$100		\$100
	<i>Crustaceans subtotal</i>				<i>\$100</i>		<i>\$100</i>
	Fishes	Chub, spotfin	Erimonax monachus	Wherever found, except where listed as an experimental population	\$600		\$600
		grouper, Nassau	Epinephelus striatus	(blank)	\$200		\$200
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia River spring-run ESU See 50 CFR 224.101	\$200		\$200
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU - See 50 CFR 224.101	\$100		\$100
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$1,700		\$1,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS - See 50 CFR 223.102	\$100		\$100
				Upper Columbia River DPS - See 50 CFR 223.102	\$200		\$200
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS - See 50 CFR 224.101	\$1,700		\$1,700
				Chesapeake Bay DPS - See 50 CFR 224.101	\$2,500		\$2,500
				Gulf of Maine DPS - See 50 CFR 223.102	\$2,500		\$2,500
				New York Bight DPS - See 50 CFR 224.101	\$2,500		\$2,500
				South Atlantic DPS - See 50 CFR 224.101	\$1,700		\$1,700
		sturgeon, Green	Acipenser medirostris	Southern DPS - See 50 CFR 223.102	\$100		\$100
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$600		\$600
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$2,500		\$2,500
	<i>Fishes subtotal</i>				<i>\$17,200</i>		<i>\$17,200</i>
	Flowering Plants	Bird's-beak, salt marsh	Cordylanthus maritimus ssp. maritimus	Wherever found	\$100		\$100
		Clarkia, Pismo	Clarkia speciosa ssp. immaculata	Wherever found	\$100		\$100
		Coneflower, smooth	Echinacea laevigata	Wherever found	\$400		\$400
		Heartleaf, dwarf-flowered	Hexastylis naniflora	Wherever found	\$400		\$400
		Jewelflower, California	Caulanthus californicus	Wherever found	\$100		\$100
		Manzanita, Morro	Arctostaphylos morroensis	Wherever found	\$100		\$100
		mountainbalm, Indian Knob	Eriodictyon altissimum	Wherever found	\$100		\$100
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$100		\$100
		Pitcher-plant, mountain sweet	Sarracenia rubra ssp. jonesii	Wherever found	\$400		\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sandwort, Marsh	Arenaria paludicola	Wherever found	\$100		\$100
		Spiraea, Virginia	Spiraea virginiana	Wherever found	\$600		\$600
		Thistle, Chorro Creek bog	Cirsium fontinale var. obispoense	Wherever found	\$100		\$100
	<i>Flowering Plants subtotal</i>				\$2,600		\$2,600
	Insects	Moth, Kern primrose sphinx	Euproserpinus euterpe	Wherever found	\$100		\$100
	<i>Insects subtotal</i>				\$100		\$100
	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$600		\$600
		Bat, Indiana	Myotis sodalis	Wherever found	\$600		\$600
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$6,000		\$6,000
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$100		\$100
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$100		\$100
		Seal, Guadalupe fur	Arctocephalus townsendi	Wherever found	\$1,800		\$1,800
		Whale, blue	Balaenoptera musculus	Wherever found	\$1,800		\$1,800
		Whale, finback	Balaenoptera physalus	Wherever found	\$5,000		\$5,000
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$5,000		\$5,000
		Whale, killer	Orcinus orca	Southern Resident DPS See 50 CFR 224.101	\$100		\$100
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$3,500		\$3,500
		Whale, North Pacific Right	Eubalaena japonica	Wherever found	\$100		\$100
		Whale, Sei	Balaenoptera borealis	Wherever found	\$300		\$300
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$100		\$100
	<i>Mammals subtotal</i>				\$25,100		\$25,100
	Reptiles	Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$10,500		\$10,500
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$100		\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, green	Chelonia mydas	Green sea turtles originating from the East Pacific Ocean, bounded by the following lines and coordinates: 41 degrees N., 143 degrees W. in the northwest; 41 degrees N. Lat. in the north; along the western coasts of the Americas; 40 degrees S. Lat. in the south; and 40 degrees S., 96 degrees W. in the southwest	\$2,500		\$2,500
				Green sea turtles originating from the North Atlantic Ocean, bounded by the following lines and coordinates: 48 degrees N. Lat. in the north, along the western coasts of Europe and Africa (west of 5.5 degrees W. Long.); north of 19 degrees N. Lat. in the east; bounded by 19 degrees N., 65.1 degrees W. to 14 degrees N., 65.1 degrees W. then 14 degrees N., 77 degrees W. in the south and west; and along the eastern coasts of the Americas (north of 7.5 degrees N., 77 degrees W.)	\$4,000		\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	\$3,000		\$3,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$3,500		\$3,500
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$5,000		\$5,000
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$6,500		\$6,500
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS - Loggerhead sea turtles originating from the North Pacific north of the equator and south of 60 degrees N. Lat	\$800		\$800
				Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	\$6,000		\$6,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$4,500		\$4,500
		Snake, eastern indigo	Drymarchon couperi	Wherever found	\$400		\$400
		turtle, bog	Glyptemys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$800		\$800
	<i>Reptiles subtotal</i>				<i>\$47,600</i>		<i>\$47,600</i>
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$100		\$100
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$100		\$100
	<i>Snails subtotal</i>				<i>\$200</i>		<i>\$200</i>
Domestic subtotal					\$97,100		\$97,100
Foreign	Fishes	Shark, Scalloped Hammerhead	Sphyrna lewini	Central and Southwest Atlantic DPS	\$1,700		\$1,700
	<i>Fishes subtotal</i>				<i>\$1,700</i>		<i>\$1,700</i>
	Mammals	Whale, gray	Eschrichtius robustus	Western North Pacific DPS See 50 CFR 224.101	\$1,700		\$1,700
		Whale, humpback	Megaptera novaeangliae	Cape Verde Island/Northwest Africa	\$200		\$200
				Central America DPS	\$1,800		\$1,800
	<i>Mammals subtotal</i>				<i>\$3,700</i>		<i>\$3,700</i>
	Reptiles	Sea turtle, loggerhead	Caretta caretta	Loggerhead sea turtles originating from the South Atlantic Ocean south of the equator, north of 60 degrees S. Lat., west of 20 degrees E. Long., and east of 67 degrees W. Long	\$200		\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
				Northeast Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northeast Atlantic Ocean north of the equator, south of 60 degrees N. Lat., and east of 40 degrees W. Long., except in the vicinity of the Strait of Gibraltar where the eastern boundary is 5 degrees 36 minutes W. Long	\$200		\$200
	<i>Reptiles subtotal</i>				\$400		\$400
Foreign subtotal					\$5,800		\$5,800
Multi-species					\$1,027,200		\$1,027,200
Multi-species subtotal					\$1,027,200		\$1,027,200
Grand Total					\$1,130,100		\$1,130,100

Smithsonian Institution - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Birds	Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$1,800		\$1,800
		Kingfisher, Guam Micronesian	Halcyon cinnamomina cinnamomina	Wherever found	\$36,800		\$36,800
		Rail, Guam	Rallus owstoni	Wherever found, except where listed as an experimental	\$15,900		\$15,900
	<i>Birds subtotal</i>				<i>\$54,500</i>		<i>\$54,500</i>
	Mammals	Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$147,500		\$147,500
		Fox, San Joaquin kit	Vulpes macrotis mutica	Wherever found	\$6,800		\$6,800
		Fox, Santa Catalina Island	Urocyon littoralis catalinae	Wherever found	\$13,500		\$13,500
		Shrew, Buena Vista Lake ornate	Sorex ornatus relictus	Wherever found	\$14,000		\$14,000
	<i>Mammals subtotal</i>				<i>\$181,800</i>		<i>\$181,800</i>
Domestic subtotal					\$236,300		\$236,300
Foreign	Mammals	Addax	Addax nasomaculatus	Wherever found	\$4,900		\$4,900
		Ass, Asian wild	Equus hemionus	Wherever found	\$18,400		\$18,400
		Deer, Eld's brow-antlered	Cervus eldi	Wherever found	\$4,000		\$4,000
		Dog, African wild	Lycaon pictus	Wherever found	\$11,300		\$11,300
		Elephant, Asian	Elephas maximus	Wherever found	\$64,400		\$64,400
		Horse, Przewalski's	Equus przewalskii	Wherever found	\$34,500		\$34,500
		Leopard, clouded	Neofelis nebulosa	Wherever found	\$28,000		\$28,000
		Oryx, scimitar-horned	Oryx dammah	Wherever found	\$35,100		\$35,100
		Panda, giant	Ailuropoda melanoleuca	Wherever found	\$28,900		\$28,900
		Rhinoceros, black	Diceros bicornis	Sub-Saharan Africa	\$38,300		\$38,300
		Tapir, mountain	Tapirus pinchaque	Wherever found	\$25,500		\$25,500
		Wolf, maned	Chrysocyon brachyurus	Wherever found	\$48,600		\$48,600
	<i>Mammals subtotal</i>				<i>\$341,900</i>		<i>\$341,900</i>
Foreign subtotal					\$341,900		\$341,900
Grand Total					\$578,200		\$578,200

Tennessee Valley Authority - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Waterdog, black warrior (=Sipsey Fork)	Necturus alabamensis	Wherever found	\$350		\$350
	<i>Amphibians subtotal</i>				\$350		\$350
	Birds	Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$10,000		\$10,000
		Stork, wood	Mycteria americana	U.S.A. (AL, FL, GA, MS, NC, SC)	\$700		\$700
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$350		\$350
	<i>Birds subtotal</i>				\$11,050		\$11,050
	Clams	Bean, rayed	Villosa fabalis	Wherever found	\$635		\$635
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Fanshell	Cyprogenia stegaria	Wherever found	\$635		\$635
		Kidneyshell, fluted	Ptychobranthus subtentus	Wherever found	\$635		\$635
		Lampmussel, Alabama	Lampsilis virescens	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Lilliput, pale (pearlymussel)	Toxolasma cylindrellus	Wherever found	\$635		\$635
		Monkeyface, Cumberland (pearlymussel)	Theliderma intermedia	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$635		\$635
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$635		\$635
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$635		\$635
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$635		\$635
		Pearlymussel, birdwing	Lemiox rimosus	Wherever found; Except where listed as Experimental Populations	\$635		\$635

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pearlymussel, cracking	Hemistena lata	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Pearlymussel, dromedary	Dromus dromas	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Pearlymussel, slabside	Pleuonaia dolabelloides	Wherever found	\$635		\$635
		Pigtoe, finerayed	Fusconaia cuneolus	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$635		\$635
		Pigtoe, shiny	Fusconaia cor	Wherever found; Except where listed as Experimental Populations	\$635		\$635
		Pimpleback, orangefoot (pearlymussel)	Plethobasus cooperianus	Wherever found	\$635		\$635
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$635		\$635
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$635		\$635
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$635		\$635
		Wartyback, white (pearlymussel)	Plethobasus cicatricosus	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$635		\$635
	<i>Clams subtotal</i>				\$15,240		\$15,240
	Crustaceans	Crayfish, Nashville	Orconectes shoupi	Wherever found	\$7,000		\$7,000
	<i>Crustaceans subtotal</i>				\$7,000		\$7,000
	Fishes	Chub, spotfin	Erimonax monachus	Wherever found, except where listed as an experimental population	\$4,635		\$4,635
		Darter, duskytail	Etheostoma percnurum	Wherever found	\$2,000		\$2,000
	<i>Fishes subtotal</i>				\$6,635		\$6,635
	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$306,270		\$306,270
		Bat, Indiana	Myotis sodalis	Wherever found	\$121,551		\$121,551
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$51,061		\$51,061
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$120		\$120

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Mammals subtotal</i>				\$479,002		\$479,002
	Reptiles	Turtle, flattened musk	Sternotherus depressus	Black Warrior R. system upstream from Bankhead Dam	\$350		\$350
	<i>Reptiles subtotal</i>				\$350		\$350
	Snails	Riversnail, Anthony's	Athearnia anthonyi	Wherever found; Except where listed as Experimental Populations	\$635		\$635
	<i>Snails subtotal</i>				\$635		\$635
<i>Domestic subtotal</i>					\$520,262		\$520,262
Multi-species					\$110,000		\$110,000
<i>Multi-species subtotal</i>					\$110,000		\$110,000
Grand Total					\$630,262		\$630,262

USDA Animal and Plant Health Inspection Service - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$200		\$200
	<i>Amphibians subtotal</i>				\$200		\$200
	Birds	Condor, California	Gymnogyps californianus	U.S.A. (Oregon, and specific portions of northern California and northwest Nevada)	\$1,500		\$1,500
				U.S.A. (specific portions of Arizona, Nevada, and Utah)-see 17.84(j)	\$200		\$200
		coot, Hawaiian	Fulica americana alai	Wherever found	\$6,100		\$6,100
		Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$4,500		\$4,500
		Cuckoo, yellow-billed	Coccyzus americanus	Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	\$200		\$200
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$2,400		\$2,400
		Eider, spectacled	Somateria fischeri	Wherever found	\$500		\$500
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	Wherever found, except where listed as an experimental population	\$200		\$200
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$14,100		\$14,100
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$2,400		\$2,400
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$2,400		\$2,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$200		\$200
		Rail, Yuma Ridgway's	<i>Rallus obsoletus yumanensis</i>	Wherever found	\$200		\$200
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$6,100		\$6,100
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$6,100		\$6,100
		Stork, wood	<i>Mycteria americana</i>	U.S.A. (AL, FL, GA, MS, NC, SC)	\$600		\$600
		Storm-petrel, band-rumped	<i>Oceanodroma castro</i>	USA (HI)	\$6,100		\$6,100
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$200		\$200
	<i>Birds subtotal</i>				<i>\$54,000</i>		<i>\$54,000</i>
	Fishes	Chub, Gila	<i>Gila intermedia</i>	Wherever found	\$200		\$200
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$200		\$200
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$200		\$200
		Chub, Virgin River	<i>Gila seminuda</i> (=robusta)	Wherever found	\$200		\$200
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$200		\$200
		pikeminnow, Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$200		\$200
		Pupfish, desert	<i>Cyprinodon macularius</i>	Wherever found	\$200		\$200
		Spikedace	<i>Meda fulgida</i>	Wherever found	\$200		\$200
		Spinedace, Little Colorado	<i>Lepidomeda vittata</i>	Wherever found	\$200		\$200
		Sucker, razorback	<i>Xyrauchen texanus</i>	Wherever found	\$200		\$200
		Sucker, Zuni bluehead	<i>Catostomus discobolus yarrowi</i>	Wherever found	\$200		\$200
		Topminnow, Gila (incl. Yaqui)	<i>Poeciliopsis occidentalis</i>	U.S. Only	\$200		\$200
		Trout, Apache	<i>Oncorhynchus apache</i>	Wherever found	\$200		\$200
		Trout, Gila	<i>Oncorhynchus gilae</i>	Wherever found	\$200		\$200
		Woundfin	<i>Plagopterus argentissimus</i>	Wherever found, except where listed as an experimental population	\$200		\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Fishes subtotal</i>				\$3,000		\$3,000
	Flowering Plants	Cactus, Arizona hedgehog	Echinocereus arizonicus ssp. arizonicus	Wherever found	\$200		\$200
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$200		\$200
		Cactus, Fickeisen plains	Pediocactus peeblesianus ssp. fickeiseniae	Wherever found	\$200		\$200
		Cactus, Siler pincushion	Pediocactus (=Echinocactus,=Utahia) sileri	Wherever found	\$200		\$200
		Cliffrose, Arizona	Purshia (=Cowania) subintegra	Wherever found	\$200		\$200
		Cycladenia, Jones	Cycladenia humilis var. jonesii	Wherever found	\$200		\$200
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$200		\$200
		Milk-vetch, Holmgren	Astragalus	Wherever found	\$200		\$200
		Milk-vetch, Sentry	Astragalus cremnophylax var. cremnophylax	Wherever found	\$200		\$200
		Milkweed, Welsh's	Asclepias welshii	Wherever found	\$200		\$200
	<i>Flowering Plants subtotal</i>				\$2,000		\$2,000
	Insects	Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$7,100		\$7,100
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$300		\$300
	<i>Insects subtotal</i>				\$7,400		\$7,400
	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$600		\$600
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$400		\$400
		Ferret, black-footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$200		\$200
		Jaguar	Panthera onca	Wherever found	\$1,200		\$1,200
		Jaguarundi, Gulf Coast	Puma yagouaroundi cacomitli	Wherever found	\$88,600		\$88,600
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$1,500		\$1,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$200		\$200
		Ocelot	Leopardus (=Felis) pardalis	Wherever found	\$88,800		\$88,800
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$200		\$200
	<i>Mammals subtotal</i>				<i>\$181,700</i>		<i>\$181,700</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$3,000		\$3,000
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$200		\$200
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$200		\$200
		Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$500		\$500
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$200		\$200
	<i>Reptiles subtotal</i>				<i>\$4,100</i>		<i>\$4,100</i>
Domestic subtotal					\$252,400		\$252,400
Foreign	Birds	Ibis, northern bald	Geronticus eremita	Wherever found	\$600		\$600
		Macaw, military	Ara militaris	Wherever found	\$1,000		\$1,000
		penguin, African	Spheniscus demersus	Wherever found	\$700		\$700
		Starling, Rothschild's (myna)	Leucopsar rothschildi	Wherever found	\$1,000		\$1,000
	<i>Birds subtotal</i>				<i>\$3,300</i>		<i>\$3,300</i>
	Mammals	Cat, black-footed	Felis nigripes	Wherever found	\$1,000		\$1,000
		Cheetah	Acinonyx jubatus	Wherever found	\$17,600		\$17,600
		Chimpanzee	Pan troglodytes	Wherever found	\$1,800		\$1,800
		Chimpanzee, pygmy	Pan paniscus	Wherever found	\$3,000		\$3,000
		Deer, Barbary	Cervus elaphus barbarus	Wherever found	\$500		\$500
		Elephant, African	Loxodonta africana	Wherever found	\$2,400		\$2,400
		Elephant, Asian	Elephas maximus	Wherever found	\$3,700		\$3,700
		Gazelle, Mhorr	Gazella dama mhorr	Wherever found	\$500		\$500
		Gibbons	Hylobates spp. (including Nomascus)	Wherever found	\$2,000		\$2,000
		Gorilla	Gorilla gorilla	Wherever found	\$500		\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Leopard	Panthera pardus	In Africa, in the wild, south of, and including, the following countries: Gabon, Congo, Zaire, Uganda, Kenya	\$4,000		\$4,000
		Macaque, Japanese	Macaca fuscata	Wherever found	\$200		\$200
		Macaque, lion-tailed	Macaca silenus	Wherever found	\$500		\$500
		Rhinoceros, black	Diceros bicornis	Sub-Saharan Africa	\$200		\$200
		Tiger	Panthera tigris	Wherever found	\$25,500		\$25,500
		Wolf, maned	Chrysocyon brachyurus	Wherever found	\$500		\$500
	<i>Mammals subtotal</i>				\$63,900		\$63,900
	Reptiles	Tortoise, Madagascar radiated	Geochelone radiata	Wherever found	\$1,000		\$1,000
	<i>Reptiles subtotal</i>				\$1,000		\$1,000
<i>Foreign subtotal</i>					<i>\$68,200</i>		<i>\$68,200</i>
Multi-species					\$616,100		\$616,100
<i>Multi-species subtotal</i>					<i>\$616,100</i>		<i>\$616,100</i>
Grand Total					\$936,700		\$936,700

USDA Animal and Plant Health Inspection Service - FY 2022

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Amphibians	Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$200		\$200
	<i>Amphibians subtotal</i>				\$200		\$200
	Birds	Condor, California	Gymnogyps californianus	U.S.A. (Oregon, and specific portions of northern California and northwest Nevada)	\$1,500		\$1,500
				U.S.A. (specific portions of Arizona, Nevada, and Utah)-see 17.84(j)	\$200		\$200
		coot, Hawaiian	Fulica americana alai	Wherever found	\$6,100		\$6,100
		Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$4,500		\$4,500
		Cuckoo, yellow-billed	Coccyzus americanus	Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	\$200		\$200
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$2,400		\$2,400
		Eider, spectacled	Somateria fischeri	Wherever found	\$500		\$500
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	Wherever found, except where listed as an experimental population	\$200		\$200
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$14,100		\$14,100
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$2,400		\$2,400
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$2,400		\$2,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$200		\$200
		Rail, Yuma Ridgway's	<i>Rallus obsoletus yumanensis</i>	Wherever found	\$200		\$200
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$6,100		\$6,100
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$6,100		\$6,100
		Stork, wood	<i>Mycteria americana</i>	U.S.A. (AL, FL, GA, MS, NC, SC)	\$600		\$600
		Storm-petrel, band-rumped	<i>Oceanodroma castro</i>	USA (HI)	\$6,100		\$6,100
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$200		\$200
	<i>Birds subtotal</i>				<i>\$54,000</i>		<i>\$54,000</i>
	Fishes	Chub, Gila	<i>Gila intermedia</i>	Wherever found	\$200		\$200
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$200		\$200
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$200		\$200
		Chub, Virgin River	<i>Gila seminuda</i> (=robusta)	Wherever found	\$200		\$200
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$200		\$200
		pikeminnow, Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$200		\$200
		Pupfish, desert	<i>Cyprinodon macularius</i>	Wherever found	\$200		\$200
		Spikedace	<i>Meda fulgida</i>	Wherever found	\$200		\$200
		Spinedace, Little Colorado	<i>Lepidomeda vittata</i>	Wherever found	\$200		\$200
		Sucker, razorback	<i>Xyrauchen texanus</i>	Wherever found	\$200		\$200
		Sucker, Zuni bluehead	<i>Catostomus discobolus yarrowi</i>	Wherever found	\$200		\$200
		Topminnow, Gila (incl. Yaqui)	<i>Poeciliopsis occidentalis</i>	U.S. Only	\$200		\$200
		Trout, Apache	<i>Oncorhynchus apache</i>	Wherever found	\$200		\$200
		Trout, Gila	<i>Oncorhynchus gilae</i>	Wherever found	\$200		\$200
		Woundfin	<i>Plagopterus argentissimus</i>	Wherever found, except where listed as an experimental population	\$200		\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Fishes subtotal</i>				\$3,000		\$3,000
	Flowering Plants	Cactus, Arizona hedgehog	Echinocereus arizonicus ssp. arizonicus	Wherever found	\$200		\$200
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$200		\$200
		Cactus, Fickeisen plains	Pediocactus peeblesianus ssp. fickeiseniae	Wherever found	\$200		\$200
		Cactus, Siler pincushion	Pediocactus (=Echinocactus,=Utahia) sileri	Wherever found	\$200		\$200
		Cliffrose, Arizona	Purshia (=Cowania) subintegra	Wherever found	\$200		\$200
		Cycladenia, Jones	Cycladenia humilis var. jonesii	Wherever found	\$200		\$200
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$200		\$200
		Milk-vetch, Holmgren	Astragalus	Wherever found	\$200		\$200
		Milk-vetch, Sentry	Astragalus cremnophylax var. cremnophylax	Wherever found	\$200		\$200
		Milkweed, Welsh's	Asclepias welshii	Wherever found	\$200		\$200
	<i>Flowering Plants subtotal</i>				\$2,000		\$2,000
	Insects	Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$7,100		\$7,100
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$300		\$300
	<i>Insects subtotal</i>				\$7,400		\$7,400
	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$600		\$600
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$400		\$400
		Ferret, black-footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$200		\$200
		Jaguar	Panthera onca	Wherever found	\$1,200		\$1,200
		Jaguarundi, Gulf Coast	Puma yagouaroundi cacomitli	Wherever found	\$88,600		\$88,600
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$1,500		\$1,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$200		\$200
		Ocelot	Leopardus (=Felis) pardalis	Wherever found	\$88,800		\$88,800
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$200		\$200
	<i>Mammals subtotal</i>				<i>\$181,700</i>		<i>\$181,700</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$3,000		\$3,000
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$200		\$200
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$200		\$200
		Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$500		\$500
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$200		\$200
	<i>Reptiles subtotal</i>				<i>\$4,100</i>		<i>\$4,100</i>
Domestic subtotal					\$252,400		\$252,400
Foreign	Birds	Ibis, northern bald	Geronticus eremita	Wherever found	\$600		\$600
		Macaw, military	Ara militaris	Wherever found	\$1,000		\$1,000
		penguin, African	Spheniscus demersus	Wherever found	\$700		\$700
		Starling, Rothschild's (myna)	Leucopsar rothschildi	Wherever found	\$1,000		\$1,000
	<i>Birds subtotal</i>				<i>\$3,300</i>		<i>\$3,300</i>
	Mammals	Cat, black-footed	Felis nigripes	Wherever found	\$1,000		\$1,000
		Cheetah	Acinonyx jubatus	Wherever found	\$17,600		\$17,600
		Chimpanzee	Pan troglodytes	Wherever found	\$1,800		\$1,800
		Chimpanzee, pygmy	Pan paniscus	Wherever found	\$3,000		\$3,000
		Deer, Barbary	Cervus elaphus barbarus	Wherever found	\$500		\$500
		Elephant, African	Loxodonta africana	Wherever found	\$2,400		\$2,400
		Elephant, Asian	Elephas maximus	Wherever found	\$3,700		\$3,700
		Gazelle, Mhor	Gazella dama mhor	Wherever found	\$500		\$500
		Gibbons	Hylobates spp. (including Nomascus)	Wherever found	\$2,000		\$2,000
		Gorilla	Gorilla gorilla	Wherever found	\$500		\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Leopard	Panthera pardus	In Africa, in the wild, south of, and including, the following countries: Gabon, Congo, Zaire, Uganda, Kenya	\$4,000		\$4,000
		Macaque, Japanese	Macaca fuscata	Wherever found	\$200		\$200
		Macaque, lion-tailed	Macaca silenus	Wherever found	\$500		\$500
		Rhinoceros, black	Diceros bicornis	Sub-Saharan Africa	\$200		\$200
		Tiger	Panthera tigris	Wherever found	\$25,500		\$25,500
		Wolf, maned	Chrysocyon brachyurus	Wherever found	\$500		\$500
	<i>Mammals subtotal</i>				\$63,900		\$63,900
	Reptiles	Tortoise, Madagascar radiated	Geochelone radiata	Wherever found	\$1,000		\$1,000
	<i>Reptiles subtotal</i>				\$1,000		\$1,000
<i>Foreign subtotal</i>					\$68,200		\$68,200
Multi-species					\$616,100		\$616,100
<i>Multi-species subtotal</i>					\$616,100		\$616,100
Grand Total					\$936,700		\$936,700

Appendix C. Expenditures Reported by States for FY 2022

Through the Association of Fish and Wildlife Agencies, States voluntarily provided annual ESA expenditures via the Service provided species reporting spreadsheet. This report includes collective expenditure data from all responding State agencies.

States Provided Expenditure Data for FY 2022	States Did Not Provided Expenditure Data for FY 2022
Alaska Arkansas Colorado Florida Idaho Indiana Iowa Maine Massachusetts Michigan Minnesota Mississippi Missouri Montana Nebraska New Hampshire North Carolina North Dakota Ohio Oklahoma South Carolina South Dakota Tennessee Texas West Virginia Wyoming	Alabama American Samoa Arizona California Connecticut Delaware District of Columbia Georgia Guam Hawaii Illinois Kentucky Kansas Louisiana Maryland Nevada New Jersey New Mexico New York Oregon Puerto Rico Northern Mariana Islands Pennsylvania Rhode Island U.S. Virgin Islands Utah Vermont Virginia Washington Wisconsin

Appendix C. Expenditures Reported by the States for FY 2022

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
Amphibians	Frog, dusky gopher - Wherever found	Rana sevosa	\$106,618	\$4,123
	Hellbender, Eastern - Missouri DPS	Cryptobranchus alleganiensis alleganiensis	\$25,500	
	Hellbender, Ozark - Wherever found	Cryptobranchus alleganiensis bishopi	\$40,500	
	Salamander, Cheat Mountain - Wherever found	Plethodon nettingi	\$3,137	
	Salamander, frosted flatwoods - Wherever found	Ambystoma cingulatum	\$40,531	\$42,229
	Salamander, Georgetown - Wherever found	Eurycea naufragia	\$4,000	
	Salamander, Jollyville Plateau - Wherever found	Eurycea tonkawae	\$4,000	
	salamander, Reticulated flatwoods - Wherever found	Ambystoma bishopi	\$40,531	\$42,229
	Salamander, Salado - Wherever found	Eurycea chisholmensis	\$8,000	
	Toad, Houston - Wherever found	Bufo houstonensis	\$61,000	
	Toad, Wyoming - Wherever found	Anaxyrus baxteri	\$105,370	
	Waterdog, Neuse River - Wherever found	Necturus lewisi	\$6,500	
<i>Amphibians subtotal</i>			<i>\$445,687</i>	<i>\$88,581</i>
Arachnids	Spider, Tooth Cave - Wherever found	Tayshaneta myopica	\$4,000	
<i>Arachnids subtotal</i>			<i>\$4,000</i>	
Birds	Crane, whooping - Wherever found, except where listed as an experimental population	Grus americana	\$158,851	
	Cuckoo, yellow-billed - Western DPS: U.S.A. (AZ, CA, CO (western), ID, MT (western), NM (western), NV, OR, TX (western), UT, WA, WY (western)); Canada (British Columbia (southwestern); Mexico (Baja California, Baja California Sur, Chihuahua, Durango (western), Sinaloa, Sonora)	Coccyzus americanus	\$40,000	
	Eider, spectacled - Wherever found	Somateria fischeri	\$16,983	
	Falcon, Eurasian peregrine - Wherever found	Falco peregrinus peregrinus	\$41,559	
	Kite, Everglade snail - Wherever found	Rostrhamus sociabilis plumbeus	\$125,743	\$5,491
	Knot, red - Wherever found	Calidris canutus rufa	\$10,438	
	Plover, piping - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	Charadrius melodus	\$59,176	
	Plover, piping - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	Charadrius melodus	\$115,325	
	rail, Eastern Black - Wherever found	Laterallus jamaicensis ssp. jamaicensis	\$48,600	
	sage-grouse, Gunnison - Wherever found	Centrocercus minimus	\$427,782	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	scrub-jay, Florida - Wherever found	Aphelocoma coerulescens	\$197,836	\$42,754
	Sparrow, Florida grasshopper - Wherever found	Ammodramus savannarum floridanus	\$106,139	\$93,454
	Stork, wood - U.S.A. (AL, FL, GA, MS, NC, SC)	Mycteria americana	\$300	\$58,000
	Tern, California least - Wherever found	Sterna antillarum browni	\$1,000	
	Tern, Roseate - U.S.A. (Atlantic Coast south to NC), Canada (Newf., N.S, Que.), Bermuda	Sterna dougallii dougallii	\$19,900	
	Woodpecker, red-cockaded - Wherever found	Picoides borealis	\$393,194	\$459,252
Birds subtotal			\$1,762,826	\$658,951
Clams	Bankclimber, purple (mussel) - Wherever found	Elliptoideus sloatianus	\$3,493	
	Bean, Choctaw - Wherever found	Obovaria choctawensis	\$3,493	
	Clubshell - Wherever found; Except where listed as Experimental Populations	Pleurobema clava	\$300	
	Ebonyshell, round - Wherever found	Reginaia rotulata	\$3,493	
	Elktoe, Appalachian - Wherever found	Alasmidonta raveneliana	\$8,000	
	fatmucket, Arkansas - Wherever found	Lampsilis powellii	\$12,000	
	Heelsplitter, Carolina - Wherever found	Lasmigona decorata	\$174,300	
	Higgins eye (pearlymussel) - Wherever found	Lampsilis higginsii	\$51,105	
	Kidneyshell, southern - Wherever found	Ptychobranchus jonesi	\$3,493	
	Lance, yellow - Wherever found	Elliptio lanceolata	\$18,500	
	Mapleleaf, winged - Wherever found, except where listed as an experimental population	Quadrula fragosa	\$800	
	Moccasinshell, Alabama - Wherever found	Medionidus acutissimus	\$3,493	
	Moccasinshell, Gulf - Wherever found	Medionidus penicillatus	\$3,493	
	Moccasinshell, Ochlockonee - Wherever found	Medionidus simpsonianus	\$3,493	
	Moccasinshell, Suwannee - Wherever found	Medionidus walkeri	\$3,493	
	Mussel, sheepnose - Wherever found	Plethobasus cyphus	\$13,000	
	Mussel, snuffbox - Wherever found	Epioblasma triquetra	\$4,300	
	Pigtoe, Atlantic - Wherever found	Fusconaia masoni	\$18,500	
	Pigtoe, fuzzy - Wherever found	Pleurobema strodeanum	\$3,493	
	Pigtoe, narrow - Wherever found	Fusconaia escambia	\$3,493	
	Pigtoe, oval - Wherever found	Pleurobema pyriforme	\$3,493	
	Pigtoe, tapered - Wherever found	Fusconaia burkei	\$3,493	
	Pocketbook, Ouachita rock - Wherever found	Arcidens wheeleri	\$10,000	
	Pocketbook, shinyrayed - Wherever found	Hamiota subangulata	\$3,493	
	Pocketbook, speckled - Wherever found	Lampsilis streckeri	\$5,500	
	Rabbitsfoot - Wherever found	Quadrula cylindrica cylindrica	\$15,300	
	Riffleshell, northern - Wherever found	Epioblasma rangiana	\$300	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	Sandshell, Southern - Wherever found	Hamiota australis	\$3,493	
	Slabshell, Chipola - Wherever found	Elliptio chipolaensis	\$3,493	
	Spectaclecase (mussel) - Wherever found	Cumberlandia monodonta	\$1,100	
	Spinymussel, James - Wherever found	Parvaspina collina	\$1,600	
	Spinymussel, Tar River - Wherever found	Parvaspina steinstansana	\$23,500	
	Threeridge, fat (mussel) - Wherever found	Amblema neislerii	\$3,493	
	Wedgemussel, dwarf - Wherever found	Alasmidonta heterodon	\$23,300	
Clams subtotal			\$437,293	
Crustaceans	Crayfish, Guyandotte River - Wherever found	Cambarus veteranus	\$2,187	
	Crayfish, Nashville - Wherever found	Orconectes shoupi	\$3,000	
	Crayfish, Panama City - Wherever found	Procambarus econfinae	\$64,627	\$3,131
Crustaceans subtotal			\$69,814	\$3,131
Fishes	Cavefish, Ozark - Wherever found	Amblyopsis rosae	\$5,600	
	chub, Peppered - Wherever found	Macrhybopsis tetranema	\$5,000	
	Chub, spotfin - Wherever found, except where listed as an experimental population	Erimonax monachus	\$1,100	
	Darter, bayou - Wherever found	Etheostoma rubrum	\$25,000	
	Darter, candy - Wherever found	Etheostoma osburni	\$28,571	
	Darter, diamond - Wherever found	Crystallaria cincotta	\$651	
	Darter, leopard - Wherever found	Percina pantherina	\$11,400	
	Darter, Niangua - Wherever found	Etheostoma nianguae	\$110,000	
	Darter, yellowcheek - Wherever found	Etheostoma moorei	\$1,300	
	Logperch, Roanoke - Wherever found	Percina rex	\$4,000	
	Madtom, Carolina - Wherever found	Noturus furiosus	\$20,000	
	Minnow, Devils River - Wherever found	Dionda diaboli	\$13,000	
	Sawfish, smalltooth - US DPS - Smalltooth sawfish originating from U.S. waters.	Pristis pectinata	\$355,427	\$3,131
	Sculpin, Grotto - Wherever found	Cottus specus	\$500	
	Shiner, Arkansas River - Arkansas River Basin (AR, KS, NM, OK, TX)	Notropis girardi	\$7,240	
	Shiner, Cape Fear - Wherever found	Notropis mekistocholas	\$3,000	
	Shiner, sharpnose - Wherever found	Notropis oxyrhynchus	\$5,000	
	Shiner, smalleye - Wherever found	Notropis buccula	\$5,000	
	Shiner, Topeka - Wherever found, except where listed as an experimental population	Notropis topeka (=tristis)	\$18,700	
	Silverside, Waccamaw - Wherever found	Menidia extensa	\$500	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	Steelhead - Snake River Basin DPS - See 50 CFR 223.102	Oncorhynchus (=Salmo) mykiss	\$21,000	
	Sturgeon, Atlantic - Carolina DPS - See 50 CFR 224.101	Acipenser oxyrinchus oxyrinchus	\$100	
	Sturgeon, Atlantic - South Atlantic DPS - See 50 CFR 224.101	Acipenser oxyrinchus oxyrinchus	\$71,754	\$3,131
	Sturgeon, Gulf - Wherever found	Acipenser oxyrinchus (=oxyrhynchus) desotoi	\$71,754	\$3,131
	Sturgeon, pallid - Wherever found	Scaphirhynchus albus	\$79,881	
	Sturgeon, shortnose - Wherever found	Acipenser brevirostrum	\$492,755	\$3,131
	Sturgeon, white - U.S.A. (ID, MT), Canada (B.C.), (Kootenai R. system)	Acipenser transmontanus	\$1,100	
	Trout, bull - U.S.A., conterminous, (lower 48 states)	Salvelinus confluentus	\$447,000	
	trout, Greenback Cutthroat - Wherever found	Oncorhynchus clarkii stomias	\$244,358	
Fishes subtotal			\$2,050,691	\$12,524
Flowering Plants	Amaranth, seabeach - Wherever found	Amaranthus pumilus	\$900	
	Arrowhead, bunched - Wherever found	Sagittaria fasciculata	\$4,000	
	Avens, spreading - Wherever found	Geum radiatum	\$3,000	
	beardtongue, Parachute - Wherever found	Penstemon debilis	\$8,300	
	Bladderpod, Dudley Bluffs - Wherever found	Lesquerella congesta	\$425	
	bladderpod, Missouri - Wherever found	Physaria filiformis	\$100	
	Bladderpod, Short's - Wherever found	Physaria globosa	\$300	
	Bladderpod, white - Wherever found	Physaria pallida	\$7,000	
	Bladderpod, Zapata - Wherever found	Physaria thamnophila	\$12,000	
	Blazingstar, Heller's - Wherever found	Liatris helleri	\$3,500	
	Bluet, Roan Mountain - Wherever found	Hedyotis purpurea var. montana	\$2,500	
	Bulrush, Northeastern -	Scirpus ancistrochaetus	\$100	
	Bush-clover, prairie - Wherever found	Lespedeza leptostachya	\$500	
	Cactus, Chisos Mountain hedgehog - Wherever found	Echinocereus chisoensis var. chisoensis	\$3,000	
	Cactus, Colorado hookless - Wherever found	Sclerocactus glaucus	\$1,225	
	Cactus, Knowlton's - Wherever found	Pediocactus knowltonii	\$1,475	
	Cactus, Lloyd's Mariposa - Wherever found	Sclerocactus mariposensis	\$3,000	
	Cactus, Mesa Verde - Wherever found	Sclerocactus mesae-verdae	\$1,475	
	Cactus, star - Wherever found	Astrophytum asterias	\$0	\$229,000
	cactus, Tobusch fishhook - Wherever found	Sclerocactus brevihamatus ssp. tobuschii	\$8,000	
	Catchfly, Spalding's - Wherever found	Silene spaldingii	\$9,000	
	Chaffseed, American - Wherever found	Schwalbea americana	\$2,500	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	Coneflower, smooth - Wherever found	Echinacea laevigata	\$5,500	
	Cory cactus, bunched - Wherever found	Coryphantha ramillosa	\$3,000	
	Dropwort, Canby's - Wherever found	Oxypolis canbyi	\$2,500	
	Fescue, Guadalupe - Wherever found	Festuca ligulata	\$13,000	
	Gerardia, sandplain - Wherever found	Agalinis acuta	\$5,500	
	Gladecress, Texas golden - Wherever found	Leavenworthia texana	\$7,000	
	Goldenrod, Blue Ridge - Wherever found	Solidago spithamaea	\$300	
	Goldenrod, Short's - Wherever found	Solidago shortii	\$1,200	
	Harperella - Wherever found	Ptilimnium nodosum	\$11,505	
	Heartleaf, dwarf-flowered - Wherever found	Hexastylis naniflora	\$600	
	Irisette, white - Wherever found	Sisyrinchium dichotomum	\$1,600	\$984,500
	Joint-vetch, Sensitive - Wherever found	Aeschynomene virginica	\$800	
	Ladies'-tresses, Ute - Wherever found	Spiranthes diluvialis	\$2,800	
	Loosestrife, rough-leaved - Wherever found	Lysimachia asperulaefolia	\$3,000	
	Meadowrue, Cooley's - Wherever found	Thalictrum cooleyi	\$100	
	milkvetch, Osterhout - Wherever found	Astragalus osterhoutii	\$475	
	Milkweed, Mead's - Wherever found	Asclepias meadii	\$21,700	
	Monkshood, northern wild - Wherever found	Aconitum noveboracense	\$500	
	Mustard, Penland alpine fen - Wherever found	Eutrema penlandii	\$1,625	
	No common name - Wherever found	Geocarpum minimum	\$2,000	
	Orchid, eastern prairie fringed - Wherever found	Platanthera leucophaea	\$1,100	
	Orchid, western prairie fringed - Wherever found	Platanthera praeclara	\$22,500	
	Penstemon, blowout - Wherever found	Penstemon haydenii	\$9,000	
	Peppergrass, Slickspot -	Lepidium papilliferum	\$9,000	
	Phacelia, DeBeque - Wherever found	Phacelia submutica	\$3,175	
	Phacelia, North Park - Wherever found	Phacelia formosula	\$2,700	
	Phlox, Texas trailing - Wherever found	Phlox nivalis ssp. texensis	\$19,000	
	Pink, swamp -	Helonias bullata	\$2,000	
	Pitcher-plant, mountain sweet - Wherever found	Sarracenia rubra ssp. jonesii	\$2,500	
	Pogonia, small whorled -	Isotria medeoloides	\$300	
	Pondberry - Wherever found	Lindera melissifolia	\$7,500	
	Poppy-mallow, Texas - Wherever found	Callirhoe scabriuscula	\$6,000	
	rock cress, Shale barren - Wherever found	Boechera serotina	\$6,301	
	Skyrocket, Pagosa - Wherever found	Ipomopsis polyantha	\$4,650	
	Spiraea, Virginia - Wherever found	Spiraea virginiana	\$2,509	
	Sumac, Michaux's - Wherever found	Rhus michauxii	\$1,600	
	Sunflower, Schweinitz's - Wherever found	Helianthus schweinitzii	\$6,500	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	Thistle, Pitcher's - Wherever found	Cirsium pitcheri	\$300	
	Twinpod, Dudley Bluffs - Wherever found	Physaria obcordata	\$425	
	wild buckwheat, clay-loving - Wherever found	Eriogonum pelinophilum	\$1,150	
Flowering Plants subtotal			\$265,215	\$1,213,500
Insects	Beetle, American burying - Wherever found, except where listed as an experimental population	Nicrophorus americanus	\$191,500	
	Beetle, Comal Springs dryopid - Wherever found	Stygoparnus comalensis	\$3,000	
	Bumble bee, Rusty patched - Wherever found	Bombus affinis	\$91,519	
	Butterfly, Karner blue - Wherever found	Lycacides melissa samuelis	\$143,650	
	Butterfly, Saint Francis' satyr - Wherever found	Neonympha mitchellii francisci	\$100	
	Dragonfly, Hine's emerald - Wherever found	Somatochlora hineana	\$600	
	Skipper, Dakota - Wherever found	Hesperia dacotae	\$11,005	
	skipperling, Poweshiek - Wherever found	Oarisma poweshiek	\$4,500	
	Tiger beetle, Northeastern beach - Wherever found	Habroscelimorpha dorsalis dorsalis	\$300	
	Tiger beetle, Salt Creek - Wherever found	Cicindela nevadica lincolniana	\$16,500	
Insects subtotal			\$462,674	
Lichens	Lichen, rock gnome - Wherever found	Gymnoderma lineare	\$400	
Lichens subtotal			\$400	
Mammals	bat, Florida bonneted - Wherever found	Eumops floridanus	\$136,956	\$3,131
	Bat, gray - Wherever found	Myotis grisescens	\$149,956	\$33,133
	Bat, Indiana - Wherever found	Myotis sodalis	\$526,700	\$41,875
	Bat, Northern long-eared - Wherever found	Myotis septentrionalis	\$139,202	\$204,875
	Bat, Ozark big-eared - Wherever found	Corynorhinus (=Plecotus) townsendii ingens	\$23,350	
	Bat, Virginia big-eared - Wherever found	Corynorhinus (=Plecotus) townsendii virginianus	\$7,961	
	Bear, grizzly - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	Ursus arctos horribilis	\$3,040,549	
	Bear, polar - Wherever found	Ursus maritimus	\$4,402	
	Bison, wood - U.S.A. (Alaska)	Bison bison athabasca	\$711,632	
	Caribou DPS, Southern Mountain - U.S.A. (wherever found), Canada (southeastern British Columbia)	Rangifer tarandus ssp. caribou	\$1,000	
	Ferret, black-footed - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	Mustela nigripes	\$338,583	
	Ferret, black-footed - Wherever found, except where listed as an experimental population	Mustela nigripes	\$217,754	
	Lynx, Canada - Wherever Found in Contiguous U.S.	Lynx canadensis	\$321,913	
	Manatee, West Indian - Wherever found	Trichechus manatus	\$6,178,427	\$135,710

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
	Mouse, Anastasia Island beach - Wherever found	<i>Peromyscus polionotus phasma</i>	\$24,397	
	Mouse, Choctawhatchee beach - Wherever found	<i>Peromyscus polionotus allophrys</i>	\$16,265	
	Mouse, Perdido Key beach - Wherever found	<i>Peromyscus polionotus trissyllepsis</i>	\$16,265	
	Mouse, Preble's meadow jumping - wherever found	<i>Zapus hudsonius preblei</i>	\$50,107	
	Mouse, southeastern beach - wherever found	<i>Peromyscus polionotus niveiventris</i>	\$24,397	
	Mouse, St. Andrew beach - Wherever found	<i>Peromyscus polionotus peninsularis</i>	\$16,265	
	Otter, Northern Sea - Southwest Alaska, from Attu Island to Western Cook Inlet, including Bristol Bay, the Kodiak Archipelago, and the Barren Islands	<i>Enhydra lutris kenyoni</i>	\$27,408	
	Panther, Florida - Wherever found	<i>Puma (=Felis) concolor coryi</i>	\$583,861	\$3,131
	Sea lion, Steller - Western DPS - See 50 CFR 224.101	<i>Eumetopias jubatus</i>	\$335,944	
	Squirrel, Carolina northern flying - Wherever found	<i>Glaucomys sabrinus coloratus</i>	\$1,700	
	Squirrel, Northern Idaho Ground - Wherever found	<i>Urocitellus brunneus</i>	\$30,000	
	whale, beluga - Cook Inlet DPS See 50 CFR 224.101	<i>Delphinapterus leucas</i>	\$396,188	
	Whale, North Atlantic Right - Wherever found	<i>Eubalaena glacialis</i>	\$867,695	\$3,131
	Wolf, gray - U.S.A. (MN)	<i>Canis lupus</i>	\$552	
	Wolf, gray - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA as follows: (1) Northern AZ (that portion north of the centerline of Interstate Highway 40); (2) Northern NM (that portion north of the centerline of Interstate Highway 40); (3) Western OR (that portion of OR west of the centerline of Highway 395 and Highway 78 north of Burns Junction and that portion of OR west of the centerline of Highway 95 south of Burns Junction); (4) Most of Utah (that portion of UT south and west of the centerline of Highway 84 and that portion of UT south of Highway 80 from Echo to the UT/WY Stateline); and (5) Western WA (that portion of WA west of the centerline of Highway 97 and Highway 17 north of Mesa and that portion of WA west of the centerline of Highway 395 south of Mesa). Mexico.	<i>Canis lupus</i>	\$1,153,868	
	Wolf, red - Wherever found, except where listed as an experimental population	<i>Canis rufus</i>	\$85,500	

Group Name	Common Name	Scientific Name	General Expenses	Land Expenses
<i>Mammals subtotal</i>			<i>\$15,428,797</i>	<i>\$424,986</i>
Reptiles	Alligator, American - Wherever found	Alligator mississippiensis	\$1,949,558	\$4,729
	Crocodile, American - Wherever found, except in U.S.A. (FL)	Crocodylus acutus	\$84,464	\$3,131
	Massasauga (=rattlesnake), eastern - Wherever found	Sistrurus catenatus	\$32,300	\$205,000
	Pinesnake, Louisiana - Wherever found	Pituophis ruthveni	\$3,000	
	Sea turtle, green - Green sea turtles originating from the South Atlantic Ocean, bounded by the following lines and coordinates: along the northern and eastern coasts of South America (east of 7.5 degrees N., 77 degrees W.); 14 degrees N., 77 degrees W. to 14 degrees N., 65.1 degrees W. to 19 degrees N., 65.1 degrees W. in the north and west; 19 degrees N. Lat. in the northeast; 40 degrees S., 19 degrees E. in the southeast; and 40 degrees S. Lat. in the south	Chelonia mydas	\$308,443	\$4,853
	Sea turtle, hawksbill - Wherever found	Eretmochelys imbricata	\$308,443	\$4,853
	Sea turtle, Kemp's ridley - Wherever found	Lepidochelys kempii	\$308,443	\$4,853
	Sea turtle, leatherback - Wherever found	Dermochelys coriacea	\$308,443	\$4,853
	Sea turtle, loggerhead - Northwest Atlantic Ocean DPS - Loggerhead sea turtles originating from the Northwest Atlantic Ocean west of 40 degrees W. Long	Caretta caretta	\$551,843	\$4,853
	Snake, eastern indigo - Wherever found	Drymarchon couperi	\$129,669	\$3,131
	turtle, bog - U.S.A. (GA, NC, SC, TN, VA)	Glyptemys muhlenbergii	\$28,000	
	Turtle, Plymouth Redbelly - Wherever found	Pseudemys rubriventris bangsi	\$3,500	
	Turtle, ringed map - Wherever found	Graptemys oculifera	\$11,000	
Reptiles subtotal			\$4,027,106	\$240,256
Snails	Cavesnail, Tumbling Creek - Wherever found	Antrobia culveri	\$7,500	
	Snail, painted snake coiled forest - Wherever found	Anguispira picta	\$21,000	
	Snail, Snake River physa - Wherever found	Physa natricina	\$1,600	
<i>Snails subtotal</i>			<i>\$30,100</i>	
<i>Multi-Species subtotal</i>			<i>\$5,709,444</i>	<i>\$9,000</i>
Total State Expenditures			\$30,694,047	\$2,650,929