

Is habitat restoration targeting relevant ecological needs for endangered species? Using Pacific Salmon as a case study

KATIE A. BARNAS,^{1,†} STEPHEN L. KATZ,² DAVID E. HAMM,³ MONICA C. DIAZ,⁴ AND CHRIS E. JORDAN¹

¹*Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, Seattle, Washington 98112 USA*

²*School of the Environment, Washington State University, Pullman, Washington 99164 USA*

³*Hamm Consulting, Seattle, Washington 98117 USA*

⁴*Pacific States Marine Fisheries Commission, Seattle, Washington 98112 USA*

Citation: Barnas, K. A., S. L. Katz, D. E. Hamm, M. C. Diaz, and C. E. Jordan. 2015. Is habitat restoration targeting relevant ecological needs for endangered species? Using Pacific Salmon as a case study. *Ecosphere* 6(7):110. <http://dx.doi.org/10.1890/ES14-00466.1>

Abstract. Conservation and recovery plans for endangered species around the world, including the US Endangered Species Act (ESA), rely on habitat assessments for data, conclusions and planning of short and long-term management strategies. In the Pacific Northwest of the United States, hundreds of millions of dollars (\$US) per year are spent on thousands of restoration projects across the extent of ESA-listed Pacific salmon—often without clearly connecting restoration actions to ecosystem and population needs. Numerous decentralized administrative units select and fund projects based on agency/organization needs or availability of funds with little or no centralized planning nor post-project monitoring. The need therefore arises for metrics to identify whether ecosystem and species level restoration needs are being met by the assemblage of implemented projects. We reviewed habitat assessments and recovery plans to identify ecological needs and statistically compared these to the distribution of co-located restoration projects. We deployed two metrics at scales ranging from the sub-watershed to ESA listing units; one describes the unit scale match/mismatch between projects and ecological concerns, the other correlates ecological need with need treated by projects across units. Populations with more identified ecological concerns contained more restoration effort, but the frequency of ecological concerns in recovery plans did not correlate with their frequency as restoration targets. Instead, restoration projects were strongly biased towards less expensive types. Many ESA-listed salmon populations (78%) had a good match between need and action noted in their recovery plan, but fewer (31%) matched at the smaller sub-watershed scale. Further, a majority of sub-watersheds contained a suite of projects that matched ecological concerns no better, and often worse, than a random pick of all project types. These results suggest considerable room for gains in restoration funding and placement even in the absence of centralized planning. This analytical approach can be applied to any species for which habitat management is a principle tactic, and in particular can help improve efficiencies in matching identified needs with explicit management actions.

Key words: conservation plan; decentralized management; ecoinformatics; ecological concern; endangered species; habitat assessment; habitat restoration; limiting factor; Pacific salmon; recovery plan; text mining.

Received 24 November 2014; revised 12 February 2015; accepted 21 February 2015; final version received 31 March 2015; **published** 15 July 2015. Corresponding Editor: D. P. C. Peters.

Copyright: © 2015 Barnas et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. <http://creativecommons.org/licenses/by/3.0/>

† **E-mail:** Katie.Barnas@noaa.gov

INTRODUCTION

Around the world, restoration and protection of habitat form the foundation of long-term conservation strategies for threatened species recovery. In many countries, protected species laws require recovery and management plans to incorporate habitat restoration (Bottrill et al. 2011). These plans, in turn, provide the scientific framework to inform restoration project design and prioritization (Bernhardt et al. 2007, Miller and Hobbs 2007, Beechie et al. 2008). Moreover, aquatic ecosystems and species worldwide generally face more threats than terrestrial ecosystems (Richter et al. 1997, Magurran 2009), thus restoration actions more commonly target freshwater and estuarine habitats (Bernhardt et al. 2005, Roni et al. 2008). However, species or watershed plans do not always guide aquatic restoration. A survey of river restoration practitioners in the United States reported that although one third of habitat restoration projects could be considered part of a larger plan, only 16% of projects were initiated within the specific context of a watershed management plan (Bernhardt et al. 2007). Such fragmentation of restoration effort, in which individual projects proceed independent of a centralized plan, may not effectively recover species with ranges larger than that of a single project.

This observed fragmentation, or decentralized decision making with respect to restoration effort, is rationalized as having lower transaction costs, increased equity for local agents and better matching local knowledge with decision making (e.g., Berkes et al. 1989, Sewell 1996, Sharma 2003, Berkes 2007). However, fragmented restoration decision making has also been criticized (e.g., Prud'Homme 1995, Mody 2004) for reinforcing existing dis-equity and inefficiencies (Lebel et al. 2004, Lane and Corbett 2005). In a striking case, resource management in the U.S.'s Columbia River Basin has achieved a sufficient level of complexity that its water management policy has been seen to move to both greater and lesser centralization simultaneously (Wandschneider 1984). For example, the overwhelming majority of management funding for habitat restoration and protection in the Columbia River Basin comes from a small number of Federal sources, while over 65 local, regional and tribal

agents coordinate, direct and prioritize that funding (G.A.O. 2002). In this study we will look at the efficiency of habitat restoration decisions in efforts to recover endangered Pacific salmon in the Northwest United States given the complex management structure of this region.

Since 1991, 18 Evolutionarily Significant Units (ESU; Waples 1991) and Distinct Population Segments (DPS; *O. mykiss*) of anadromous Pacific salmonids (*Oncorhynchus* spp.) have been listed under the Endangered Species Act (ESA) in Washington, Oregon and Idaho covering a footprint of over 290,000 km², an area larger than the state of Oregon. As with most endangered species, the ESA listings (16 U.S.C. §§ 1531–1544) for Pacific salmon identified habitat degradation as a cause of decline. Thus habitat restoration has been widely applied to improve spawning and rearing habitat in hopes of increasing salmon numbers to meet recovery goals (National Research Council et al. 1996). The Pacific Northwest now contains one of the highest densities of freshwater restoration projects in the U.S. (Bernhardt et al. 2005), and is essentially the largest freshwater restoration effort ever undertaken on behalf of an endangered species with billions of dollars spent to date (G.A.O. 2002, NMF. 2013). Despite this extraordinary financial investment, projects to restore freshwater and estuarine habitat are assessed for effectiveness at the individual project level rather than in the context of the greater recovery efforts, if post-project monitoring is funded at all (Bash and Ryan 2002). Many projects are funded piece-meal by various administrative units based on the priorities of the funding agency, availability of funds, ease of implementation and buy-in by landowners and stakeholders rather than needs outlined in management plans (Miller and Hobbs 2007, Kondolf et al. 2008, Katz 2009). Indeed, in 2002 the U.S. Government Accountability office estimated that 66% of non-mainstem and non-hatchery federal expenditures (i.e., the fraction available for habitat restoration and protection) were distributed directly to local, non-Federal entities for salmon recovery management (US \$537.2M from 1997 to 2002; G.A.O. 2002).

In light of the lack of coordination across the restoration enterprise (Bernhardt et al. 2007, Kondolf et al. 2008), there is a critical need to

evaluate how well habitat restoration actions match the impaired habitat conditions, especially when targeting a threatened or endangered species like Pacific Salmon. Such evaluations establish accountability for the use of public and private funds and establish reasonable performance expectations for the considerable efforts and resources applied to restoration (G.A.O. 2002, Katz et al. 2007). In recent years databases have been developed to inventory aquatic restoration actions (Bernhardt et al. 2005, Katz et al. 2007); unfortunately, monitoring-derived data to characterize the impairment of aquatic habitat conditions is not similarly forthcoming (Rumps et al. 2007, O'Donnell and Galat 2008, Hamm 2012). Thus, assessing the match between actions needed across the landscape and actions completed must resort to more creative approaches.

Acknowledging that recovery planning could be improved by a better linkage between basic ecology and management actions (Clark et al. 2002, Palmer 2009, Dickens and Suding 2013), and that restoration success at species relevant spatial scales requires better coordinated approaches (Paulsen and Fisher 2005, Kondolf et al. 2008), here we develop methods to assess how restoration expenditures for Pacific Salmon align with ecological needs. We have adopted two approaches: relating expressed habitat concerns to completed restoration with a presence/absence metric on a unit scale, and correlations of need and project frequencies across units. We assay expressed ecological need by surveying management plans (i.e., data sets) encompassing various spatial scales relevant to species management, combine this information with data on the restoration actions in those same spatial units, and ask (1) does restoration address ecological concerns at the sub-watershed scale within a salmon population, and (2) does restoration address ecological concerns at the scale of a salmon population within an Endangered Species Act listing unit (ESU/ DPS)? This method provides an objective way to retrospectively assess types of restoration projects and their placement on landscapes, and evaluate the appropriateness of proposed projects for a given species or population based on documented ecological concerns.

METHODS

Identifying ecological concerns

We assembled ecological concern data by surveying documents from two hierarchical programs addressing watershed management and recovery planning for statements of ecological need. While these programs have large conceptual overlap, they use different semantics, cover different geographic footprints, and use different source data. As a consequence, we had to standardize the datasets before synthesizing; each data set is described in turn.

We reviewed subbasin plans for the Columbia Cascade Ecoprovince (<http://www.nwccouncil.org/fw/subbasinplanning>), located in North Central Washington State (Fig. 1). The 2005 subbasin plans prioritize habitat restoration actions and provide guidance for funding restoration as a part of the Northwest Power and Conservation Council's Columbia River Basin Fish and Wildlife Program, enacted to mitigate the impacts of hydropower and to fulfill legal obligations under the Endangered Species Act and Clean Water Act (Northwest Power and Conservation Council 2005). The Columbia Cascade Ecoprovince encompasses six subbasins: Chelan, Entiat, Methow, Okanogan, Upper Middle Columbia and Wenatchee and some smaller watersheds that drain directly to the Columbia River (Fig. 1). The subbasin plans divide the Columbia Cascade into 86 smaller spatial units, what we refer to as sub-watershed assessment units. This region historically produced healthy runs of anadromous salmonids (e.g., Mullan et al. 1992) but the Upper Columbia Chinook ESU and Upper Columbia Steelhead DPS now are listed as Endangered.

As the subbasin plans are only available for the Columbia River Basin, which excludes a number of ESA listed salmon ESU/DPSs, we also evaluated NOAA Fisheries final or public draft recovery plans for Pacific salmon in Washington, Oregon, and Idaho (<http://www.nmfs.noaa.gov/pr/recovery/plans.htm>). The recovery plans cover 219 populations nested within 16 ESU/DPSs and were developed from 2000 to the present (Fig. 1). The Oregon Coast Coho ESU and Puget Sound Steelhead DPS recovery plans are currently in development so could not be used in this analysis. Four additional populations were not evaluated in the recovery plan for a given ESU/




Fig. 1. ESA listed ESU/DPSs of Pacific Salmon cover large portions of the states of Washington, Oregon, and Idaho, outlined in black. Populations with completed recovery plans are shown gray. The Columbia Cascade Ecoprovince (patterned) is within the state of Washington and overlaps with the Upper Columbia ESU/DPSs.

DPS, and so were excluded from our analyses (Upper Willamette Steelhead–West Side Tributaries, Mid-Columbia Steelhead–Willow Creek, Upper Columbia Steelhead–Crab Creek, Puget Sound Chinook–Skykomish). We also excluded extirpated populations if they were not covered in their respective recovery plans (Snake Spring Summer Chinook–Clearwater populations and Snake River Steelhead–Clearwater and Hells Canyon populations) (Fig. 1).

For each spatial unit described in the plans we polled projects from the Pacific Northwest Salmon Habitat Project Database (Katz et al. 2007, NMFS 2014). All restoration projects in the PHSHP database are attributed with a project type and subtype as well as geo-referenced which enabled sampling at any scale using GIS. We defined a project as a unique location and project subtype combination. We confined our

queries to projects completed in the 20 years since the first Pacific Salmon ESA listing (1992–2011, 36,895 projects) and to actions involving physical changes to freshwater or estuarine habitat. We evaluated completed projects recognizing that in total, it can take up to 10 years from the application for funding and permits through project implementation, completion and final reporting before inclusion in a database (Katz et al. 2007). While the PNSHP restoration database contains the vast majority of known projects it is not considered a complete census (Barnas and Katz 2010).

Semantic synthesis

Among the challenges to the current work, the subbasin and recovery plans lack not only a prioritization of habitat impairment, but lack the common language, definitions and spatial units

to characterize such impairments (Hamm 2012). Prior work to address the lack of common semantics produced a standardized dictionary of ecological concerns, which classifies degraded salmon habitat in the Pacific Northwest (Hamm 2012) and has been incorporated into the most recent salmon recovery plans. An ecological concern was defined as: changes to the ecological conditions essential for maintaining the long-term viability of a given salmonid population, causing mortality, injury, reduced health or reduced reproduction (Hamm 2012) and consists of 10 types and 34 nested subtypes. Each of the source documents in this study were reviewed for all references to ecological concerns as defined by Hamm (2012) (e.g., habitat limiting factors, impaired habitat etc.) within each spatial unit.

Sub-watersheds generally followed the watershed boundaries of approximated fifth or sixth field United States Geological Survey (USGS) Hydrologic Cataloging Unit Codes (HUCs), or in some cases an ad hoc mixture of the two. In general the sub-watersheds did not map one-to-one with the population units described in Pacific salmon recovery plans. The spatial footprint of salmon populations within an ESU/DPS range in size from 40 to 12,000 km², or approximately a seventh field HUC to a third field HUC.

Crosswalk development

To relate restoration project types to the ecological concerns they treat, we developed crosswalks defined as “a table that maps the relationships and equivalencies between two or more metadata schemes” (Dublin Core Metadata Initiative: glossary; <http://dublincore.org/documents/usageguide/glossary.shtml>), using the ecological concerns (Hamm 2012) and PNSHP data dictionaries (Katz et al. 2007). Both the ecological concerns data dictionary and the PNSHP database are structured with general types and nested subtypes allowing us to create detailed crosswalks. Ideally each project subtype would map to a single ecological concern subtype; in reality, a restoration project type or subtype may address many ecological needs, e.g., a riparian planting project may treat bank stabilization, sedimentation, provide riparian cover, or all three. In response to this potential ambiguity, we created separate crosswalks to

obtain upper (Broad) and lower (Narrow) bounds to relate restoration projects to ecological need. The bounding crosswalks then allowed us to create a third “Intermediate” crosswalk.

In the “Broad” crosswalk (Appendix A: Tables A1–A4), we attributed projects to ecological concerns even if the likelihood of success was highly optimistic, contingent on large-scale implementation or on long-term maintenance for project effectiveness. For example, “riparian planting” addresses riparian condition as well as decreases stream temperatures due to shading, buffers sediment runoff, enhances nutrient subsidy, reduces bank erosion, augments the recruitment of large woody debris, and increases stream complexity due to mature trees falling into the river. For the “Narrow” crosswalk (Appendix B: Tables B1–B4), we attributed project types with addressing only the most probable goal. In this case, a “riparian planting” project is credited with addressing only “riparian conditions”.

The “Intermediate” crosswalk attributed projects with all concerns suggested by the literature unless cause and effect are separated by many steps, require coordinated implementation across large spatial scales or require time scales of decades for success (Appendix C: Tables C1–C4). For example in the “Intermediate” crosswalk, riparian planting addresses water temperature (through future shading), decreases sediment runoff (by stabilizing banks and slowing water runoff) but does not receive credit for altering channel form or increasing instream habitat complexity. These channel changes would result only if riparian plantings matured and fell into the river in sufficient quantities, a process that takes decades (Beechie et al. 2010). We used the “Intermediate” crosswalk to assign ecological concerns to restoration projects for our analyses.

Analysis

We initially surveyed subbasin and restoration plans for prioritization of ecological needs screening for terms such as ‘primary’ or ‘major’ within a sub-watershed or salmon population. This would have provided the data to correlate the highest priority needs with the most numerous or intensive restoration type(s). This level of relative prioritization was not available across all documents. Secondly, we looked for spatially explicit information on ecological need (e.g., lat/

long coordinates for needed barrier removals). This too was lacking. In response, we developed a metric of match/mismatch based simply on presence or absence. For each spatial assessment unit, using Geographic Information Systems (GIS) we compared restoration projects from PNSHP to ecological concerns gathered from subbasin plans (sub-watershed) or recovery plans (population). When an ecological concern was treated by one or more projects based on the intermediate crosswalk it was considered a match.

We calculated the percentage of projects matching the ecological concerns (Appendix D: Table D1), and the percentage of ecological concerns matched by at least one project for each unit resulting in the Salmon Habitat Assessment and Project Evaluator metric (SHAPE):

$$\text{SHAPE} = \frac{\text{No. concerns addressed}}{\text{No. concerns} - \frac{\text{No. restoration actions not matched}}{\text{No. restoration actions}}}.$$

The metric ranges from -1 , if projects failed to match any ecological concerns, to 1 , if all projects were appropriate and all ecological concerns were addressed by at least one project. A zero is reported if there are no projects, whether or not ecological concerns are present. If no ecological concerns are identified but there are projects, the SHAPE score is -1 . We considered a “good” SHAPE score anything above the average of all (sub-watershed and population combined) SHAPE scores (0.56) and a “poor” score anything below the lower quartile for all scores (0.40).

To evaluate SHAPE metric performance, we resampled with replacement the projects found in each sub-watershed from all possible project types, compared these project types to the ecological concerns using the crosswalk, and recalculated the SHAPE. We ran this resampling 5000 times to test if projects completed in each spatial unit better matched ecological concerns than project types selected at random. We then compared the mean permuted SHAPE score to the measured SHAPE score.

The metric values are continuous, but the ratios are bounded and based on presence/absence data. Therefore all statistical comparisons used non-parametric techniques such as Kendall's rank correlations. All statistics were

estimated using the R statistical computing environment (R Core Team 2013).

RESULTS

ESA listing units and populations within listing units

Ecological concern types were not equally likely to be documented in recovery plans across the salmon recovery domain; there were six relatively common and three relatively rare types (Fig. 2A). The six most common types made up almost 85% of the total, and were problems with: Peripheral and Transitional Habitats, Channel Structure and Form, Water Quantity, Riparian Condition, Water Quality and Sediment Conditions (see Hamm, 2012 for ecological concerns definitions). The aggregate distribution of restoration projects was distinct from ecological need in being dominated by a couple project types with the rest being relatively rare (Fig. 2A). Projects that targeted Water Quality and Sediment Conditions together represented over half of all projects (52%). Many restoration types address Water Quality and Sediment Condition (Appendix C: Tables C1–C4) so the allocation of projects to those categories may in part be a product of the crosswalk. The least common projects targeted Food Limitation and Injury and Mortality (5% of total), these categories are also the least commonly expressed ecological needs (Fig. 2A). Over the first two decades of salmon recovery we found no change in the project types utilized with the same types making up the majority of projects completed (1990s vs 2000s, Kendall's $\tau = 0.88$, $p = 0.00001$).

The number of ecological concern types and subtypes in a salmon population varied from 0 to 17 with the majority of populations having 9–12 concerns. We found a wide range in the number of projects within a given population, 0–2577, though the majority of populations had project numbers in the hundreds rather than thousands (median = 138). We see a positive relationship between the number of ecological concerns and the number of restoration projects across salmon populations ($r^2 = 0.20$, $p = 2.8 \cdot 10^{-12}$; Fig. 3). While this result supports the inference that more ecological concerns result in more restoration effort, it does not speak to the appropriateness of the restoration or other potential factors. We




Fig. 2. The frequency of expressed ecological concern (gray) and restoration to address that ecological concern (black) for (A) the 219 salmonid populations with ecological concern data from recovery plans and (B) the 86 sub-watersheds within the Columbia Cascade Ecoprovince subbasin plans.

found a number of stated ecological concern subtypes untreated by restoration in 10 or more populations: Predation (Injury and Mortality), Instream Structure and Bed and Channel Form (Channel Structure and Form), Side Channel, Floodplain, and Nearshore (Peripheral and Tran-




Fig. 3. Plot of the log of total number of restoration projects as a function of number of ecological concerns within each salmon population ($F_s = 54.92$, $df = 1,217$, $p = 2.8E-12$).

sitional Habitats), Decreased Water Quantity (Water Quantity), and Large Woody Debris Recruitment (Riparian). When we evaluate appropriateness at the individual population level using the SHAPE metric, scores covered the range from -1.0 to 1.0 . The distribution skewed toward higher values (mean = 0.75 , median = 0.87) with 79% of populations ($174/219$) scoring in the “good” range (>0.56), while 27 fell into the “poor” range (<0.40) (Fig. 4A).

The Columbia Cascade and sub-watershed units

In the Columbia Cascade Ecoprovince, ‘Channel Structure and Form’ was the most frequently documented ecological concern category (Fig. 2B) although the top six concern categories were similarly frequent (12.5–17.5% of concerns): Habitat Quantity, Water Quantity, Water Quality, Peripheral and Transitional Habitats, and Sediment Conditions (Fig. 2B). The Columbia Cascade Ecoprovince contains a few relatively frequent project types and a larger number of rarer types similar to the distribution of restoration projects across the salmon recovery domain (Fig. 2). Projects addressing Water Quality were the most numerous and constituted 24% of all projects in the Columbia Cascade. Water Quality along with Riparian Condition, Sediment Condition and Water Quantity, represented almost 75% of all projects (Fig. 2B).

Over both the entire salmon recovery domain and the Columbia Cascade Ecoprovince, we

found a poor rank-order correlation between ecological concerns and restoration projects (Kendall’s $\tau = 0.22$, $p = 0.46$ salmon populations; $\tau = 0.25$, $p = 0.40$ Columbia Cascade sub-watersheds). Interestingly, we also found a poor rank-order correlation between the expressed ecological concerns in the Columbia Cascade Ecoprovince compared to the larger salmon recovery domain ($\tau = 0.30$, $p = 0.29$). Despite this, the frequencies of restoration project types were highly correlated between the Columbia Cascade and the salmon recovery domain ($\tau = 0.88$, $p = 0.001$).

The 86 sub-watersheds within the Columbia Cascade Ecoprovince contained between 0 and 12 ecological concerns (median = 3) and 0 to 188 restoration projects (median = 6). Of the Columbia Cascade sub-watersheds, 7% had ecological concerns but no projects, and an additional 47% of sub-watersheds had one or more ecological concerns not matched by a project. Of sub-watershed units, 31% ($27/86$) scored in the “good range” (SHAPE > 0.56), compared to 79% of salmon populations (Fig. 4A). Over 86% of populations and 43% of sub-watersheds had a SHAPE score that was above “poor” (SHAPE < 0.40 ; Fig. 4A). Unlike the salmon population SHAPE scores however, the Columbia Cascade distribution has two distinct modes; 14 sub-watersheds scored 0.0, 13 of which had no restoration projects, and 24 sub-watersheds scored a -1.0 , with 67.0% ($56/86$) scoring “poorly” (mean = 0.06 , median = 0.08).

When ecological concerns and project numbers were held constant, but project type chosen at random, 10 sub-watersheds (14%) scored the same between the measured SHAPE score and the re-sampled SHAPE score, 53% of sub-watersheds scored higher and 33% lower. The mean difference was 0.19 higher for re-sampled SHAPE scores over measured SHAPE scores.

Effect of scale

The assessment units in this study varied three orders of magnitude, yet are all management units directly applicable to restoration planning for Pacific Salmon recovery. Smaller units averaged more ecological concerns and restoration projects per unit area ($<500 \text{ km}^2$ unit: 0.9 EC/km^2 , $2.1 \text{ projects/km}^2$; $>5000 \text{ km}^2$ unit: 0.002 EC/km^2 , $0.06 \text{ projects/km}^2$), but larger units on




Fig. 4. SHAPE score by (A) assessment unit type binned as the percentage of all assessment units: Columbia Cascade sub-watersheds (light grey; $n = 86$), salmon populations (dark grey; $n = 219$); ESU/DPSs (black; $n = 16$) and SHAPE score (B) as a function of assessment unit size (km^2) for sub-watersheds and populations. The inset includes scores for ESU/DPSs. We considered a “good” SHAPE score anything above the average, 0.56, and a “poor” score anything below the lower quartile for all scores, 0.40.

average had more total restoration. Thus, we evaluated how the match and mismatch of restoration effort and ecological concern interacted with size of unit.

Plotting the SHAPE scores for all assessment units reveals a “dust bunny” distribution (McCune et al. 2002), where data distribute along the axes (Fig. 4B). We found the greatest range of

SHAPE scores in the smallest units and the greatest range in assessment unit size at the highest SHAPE scores. The ESU/DPSs make up almost all of the values from 12,000 to 86,000 km^2 (Fig. 4B inset), so we removed the ESU/DPSs to evaluate the remaining data from smaller units (Fig. 4B). The Columbia Cascade Ecoprovince SHAPE score modes of -1.0 and 0.0 (Fig. 4B) are

almost entirely from very small sub-watersheds (<500 km²), although numerous small sub-watersheds also received higher SHAPE scores. However, no large assessment units (>5000 km², n = 22) scored poorly and all but one scored in the 'good' range.

We further evaluated the manner in which the SHAPE score responds to scale by looking at how the components of the SHAPE score are affected by the size of the assessment unit. When salmon populations are binned into five size classes, SHAPE scores start above 0 and increase when going from small to large units (Fig. 5A). Over this range of unit size, the percent of ecological concerns (arcsin transformed) addressed increases with unit size and shows a decreasing trend in the variance across unit sizes (Fig. 5B). While the percent of restoration actions not addressing a need (arcsin transformed) is about the same for all population sizes, the variance decreases markedly for larger populations (Fig. 5C). This supports the idea that the SHAPE score variance has a scale-dependence that may bias toward higher values in larger units.

Where salmon species and assessment units overlap, unit size and species specific ecological concerns can lead to differing SHAPE scores (Fig. 6A–E). For example, much of central Idaho has a "good" SHAPE score (>0.56) with respect to steelhead (*O. mykiss*), but "poor" scores (<0.40) for Chinook populations (*O. tshawytscha*). All species except sockeye salmon (*O. nerka*) had an average SHAPE score above 0.56. Despite this, 27/219 populations still fell between –1 and 0.04, indicating a poor relationship between actions completed and expressed habitat need (Fig. 6A–D).

Effect of cost

Water Quality projects were the most expensive, averaging \$2.3M while Sediment Reduction and Upland Management projects were cheapest, averaging \$55,000 and \$85,000 respectively. Projects that matched an ecological concern had similar costs to those of unmatched projects with one exception; Water Quality projects averaged almost \$200K more per project in assessment units with a water quality concern (Fig. 7). Water Quality and Estuary/Nearshore projects were both the most expensive and the least numerous,

while the inverse was true of Riparian and Sediment Reduction projects. Thus, we found a negative association between cost and abundance of project types.

Effect of crosswalk

The sensitivity of our results depends on the choice of crosswalk. Both the percentage of projects treating a concern ($\tau = 0.67$, $p < 0.001$) and the percent of concerns addressed ($\tau = 0.62$, $p < 0.001$) differed significantly between the two bounding crosswalks. Predictably, the Intermediate crosswalk results fell in the middle with an assessment unit median of 74% (39% Narrow, 80% Broad) of Columbia Cascade projects matching an expressed ecological concern, and 50% (35% Narrow, 78% Broad) of sub-watersheds with ecological concerns treated by restoration. The median SHAPE score for the Columbia Cascade sub-watersheds varied between 0.0 (Narrow), .07 (Intermediate) and .28 (Broad).

DISCUSSION

Do habitat restoration projects address the ecological needs identified within sub-watersheds or salmon populations for ESA-listed Pacific salmon? Our results show variability in restoration decision making across the extent of salmon recovery and suggest room for tangible gains in restoration efficiency. Given the increasing availability of restoration project spatial data, the lack of consistently reported and prioritized ecological needs is the principle limit in our ability to evaluate project placement. That these data were not forthcoming is perhaps surprising as the Endangered Species Act offers guidance in the development of recovery plans that includes prioritizing actions taken in recovery (Stanford Environmental Law Society 2001). Importantly however, the ESA does not require explicit prioritization of ecological concerns, nor an explicit connection between need and action. We were left with using two approaches to the question, operating at different scales, with different abilities to evaluate decision making, and supporting different inferences about the appropriateness of restoration actions. Being a retrospective look at the question, both approaches are ultimately constrained by the properties of the available data and represent




Fig. 5. Box and whisker plots as a function of five assessment unit size categories for (A) SHAPE Score, (B) the ratio of expressed ecological concerns addressed by restoration (arcsin transformed) to the total number of expressed ecological concerns within each assessment unit (i.e., the left side of Eq. 1) and (C) the ratio of restoration projects that do not match an ecological need (arcsin transformed) to the total number of restoration projects within each assessment unit (i.e., 1 minus the right side of Eq. 1). In all panels of the figure, the median values are plotted as dots.

an attempt to extract as much information as possible from a limited resource.

Aggregating the data from all assessment units, we found a weaker than expected, and

non-significant correlation between project type and frequency of ecological concern. This suggests a lack of connection between ecological need and the use of restoration across the spatial




Fig. 6. Maps of SHAPE score for each population of ESA-listed Pacific salmon (A) Chinook salmon populations (B) Chum and Sockeye salmon populations (C) Coho salmon populations (D) Steelhead trout populations and for the (E) Columbia Cascade Ecoprovince sub-watersheds. In each case, poor SHAPE scores (<0.40) are colored in shades of red, good SHAPE scores (>0.56) are green and blue, and intermediate scores are yellow or orange.

extent of either management process (Fig. 2). However, looking both at individual SHAPE scores and accumulating SHAPE scores across the region suggests that restoration types are

often appropriately placed despite fragmentation of restoration efforts. The majority of populations and close to half of the sub-watersheds had a SHAPE score that was above




Fig. 7. Average cost for Pacific Northwest Salmon Habitat Restoration Database projects by type found within ESU/DPSs. The cost for restoration projects that matched an ecological need within a population (black) is distinguished from the cost for those projects of the same type that did not address an expressed need (gray). Small squares show total project number for each type.

“poor” (Figs. 4A, 6).

Mismatches between stated need and action do exist. Almost half of the Columbia Cascade sub-watersheds had one or more ecological concerns not matched by a project. When we randomly resampled project types for Columbia Cascade sub-watersheds, SHAPE scores for over half of sub-watersheds were higher than when we calculated SHAPE using real project data. This suggests a poor connection between habitat assessment and restoration decision making. Further, across the extent of ESA listed salmon in the Pacific Northwest, we found over 7000 projects that did not match an expressed ecological concern for a salmon population in spite of generally positive SHAPE scores. Thus a good SHAPE score does not mean all projects have been appropriately placed, but a poor SHAPE score identifies the areas that could be prioritized for further investigation and restoration imple-

mentation improvements.

Spatial scale and patterns

Whether aggregating SHAPE scores from assessment units or looking at single unit comparisons, scale drives some of the observed patterns in this analysis. Fig. 5 suggests the SHAPE metric is best suited to the HUC 4 to HUC 6 scale (i.e., $< \sim 2,000 \text{ km}^2$), as this is where the dynamic range of the SHAPE score is the greatest and this scale aligns well with other conservation analyses. The HUC 5–6 is the “local” spatial scale used by other groups to determine limiting factors for Pacific Salmon in order to “review and prioritize restoration activities and guide future funding decisions” (Oregon Coast Coho Conservation Plan for the State of Oregon, March 16, 2007) and to calculate population-status metrics such as the Conservation Success Index (Williams et al. 2007). Other

studies have used the larger USGS HUC 3 and 4 units to assess watershed conservation value (Pinsky et al. 2009).

Populations with more ecological concerns also have more total restoration effort (Fig. 3). This suggests that net restoration effort is appropriately distributed based on ecological need, although the specific project types may not match the stated needs. This pattern may reflect the common view of restoration funders that the cumulative impact of multiple restoration projects will result in enhanced ecosystem function, and in turn, improve salmon population survival and abundance (NMFS 2000). Restoration effort alone may indeed have other benefits for salmon (Allen et al. 1997). For example, Paulsen and Fisher (2005) found that watersheds with more restoration effort correlated with higher juvenile Chinook parr to smolt survival. These findings do not discriminate between more effort being more effective in aggregate (efficiency neutral), and more effort generating more appropriate, and thus more effective project types by chance (efficiency decreasing).

Restoration projects often don't identify a target species, thus where species spatially overlap a project may address an ecological concern for one species, but not address the ecological needs of others. However, we found this had little impact on SHAPE scores. Where identical population boundaries cover more than one species or multiple run-timings of a single species (71 populations, 27 comparisons), SHAPE scores varied among species/run-timing just over half time, but by small amounts (<0.05) in most cases. The observed low impact in part motivated our consideration of multiple, overlapping populations. By conducting an analysis at the scales used to evaluate salmon recovery, the ESU/DPS and populations within, we have done a triage, which will allow more localized analyses targeted at the specific areas where they are most needed (e.g., areas with few or no projects matching one or more ecological concerns).

Other patterns in restoration usage

A suite of common project types appear to be implemented throughout the Western U.S. for Pacific Salmon regardless of habitat need, likely a result of decentralized decision making regarding restoration. We found no change in the

project types utilized when comparing the first decade of salmon recovery (1990s) to the more data rich 2000s. The most common project types in this study mirror those identified by Katz et al. (2007), who found that sediment reduction, riparian planting and instream structures are both the least expensive project types and the most common based on the PNSHP database (Figs. 2, 7). Sediment reduction (via road repair), riparian stabilization and instream structures were also the most common project types in an analysis of the Russian River basin, CA (Christian-Smith and Merenlender 2010). Further, the high correlation between frequency of project types in the Columbia Cascade sub-watersheds and the salmon populations on the one hand, and the poor correlations between frequency of project type and ecological need on the other, reinforces the idea that there is a default suite of restoration actions. The differences in size, diversity and character of the spatial units covered in these planning processes and the fact that the ecological need frequencies were poorly correlated between the two planning efforts suggest that while the assessment authors were likely acting independently, the ultimate restoration types implemented were still similar.

Whatever the underlying process, the resulting pattern of commonly used restoration actions leaves some ecological concerns less likely to be treated and others perhaps over treated. Sediment reduction, riparian planting and fish passage were both the most common project types and the project types least likely to match a known concern, with 73%, 75%, 72% of projects matching stated needs respectively (Fig. 7). Fish passage was the third most common project type, and although a majority of fish passage projects did match a known habitat quantity concern, that still left 1492 completed fish passage projects in 74 populations without a stated fish passage issue. In these cases, project sponsors may have been relying on guidance that in the absence of detailed information, restoration types with a high success rate and quick response time, like barrier removal, should be employed first (Beechie et al. 2008).

In previous work (Katz et al. 2007) and the present study, project cost most strongly predicts the use of restoration types, suggesting cost drives decision making by funders and project

sponsors. Ideally, funding agencies would direct restoration efforts at identified ecological concerns, given that most funders have stated goals of restoring ecosystem function, maintaining populations and adhering to relevant laws (Clean Water Act, ESA). That cost is a significant driver of decision making is not surprising however. Social constraints such as landownership, public acceptance, and funder priorities influence project type and placement (Halle 2007, Miller and Hobbs 2007, Kondolf et al. 2008, Christian-Smith and Merenlender 2010).

Some restoration project types have functional connections with land use that constrain their utility across a diverse landscape. For example, fish screens keep fish out of surface water diversions (e.g., hydroelectric facilities, irrigation, municipal and industrial water withdrawal projects). Fish screens are also relatively expensive and custom fabricated so are unlikely to be utilized where not essential. We found that 99% of fish screening projects matched an ecological concern and only two populations that needed fish screening projects did not have at least one. Thus, at the salmon population scale fish screening projects are, for the most part, efficiently funded and placed.

Implications for species recovery

Pacific salmon cover an extremely large area, matched only by other highly-migratory animals, or plants with large ranges. Pacific Salmon ranges overlap management jurisdictions that span municipal to international scales. The scale of restoration effort is equally as large, as the total number of completed projects in PNSHP doubled from around 7000 in the 1990s to over 14000 in the 2000s as a result of the Pacific Salmon ESA listings (NMFS 2014). This considerable restoration effort took place at the same time the subbasin and recovery plans were being drafted. Though the management documents lag behind the rush to implement restoration to conserve a species, our findings provide a baseline to inform adaptive management and suggest where restoration types are underutilized based on need (Runge 2011).

Widely distributed species, or those with large migration corridors present a number of unique recovery challenges including decentralized and overlapping multi-agency management, in-

creased diversity of restoration funders, and increasingly complex ecological threats—each of these factors demand increased cost, longer-term management, and consequently greater accountability (Boyd et al. 2014, Carroll et al. 2014). The tools developed here address some of these challenges by examining large amounts of project data from multiple sources, over large spatial extents in a fast, consistent and transparent fashion. To do so however, required the development of a new project appropriateness measure since there were no existing mechanisms in use across the scale of salmon recovery that linked restoration actions to local ecosystem needs. If restoration deployment in the future is to be more efficient in addressing ecological need, common metrics will need to be incorporated into the regional management decision making frameworks. The approach developed here makes restoration appropriateness transparent to all decision makers from the landowner proposing a restoration project to the federal, state, local, tribal, and private entities involved in restoration funding, habitat assessment, and conservation planning.

While the SHAPE score evaluates restoration activity appropriateness, we have not assessed if the restoration was ultimately successful or if enough restoration has been done to alleviate an ecological concern. Ultimately, even appropriately placed projects cannot be deemed successful without proper monitoring (O'Donnell and Galat 2008, Dickens and Suding 2013, Palmer et al. 2007). In addition, even when projects are identified and population response metrics are available, high variability limits statistical power to inferences about effectiveness at only the largest spatial scales if at all (Paulsen and Fisher 2003, 2005). To obtain reliable inferences of management action effectiveness on the scale of a salmon population would require either data that does not currently exist over that scale, including restoration project success criteria, habitat monitoring, and spatially explicit habitat assessments, or application to a species with smaller spatial scales. For threatened species with smaller spatial footprints however, it is likely that the complexity of fragmented and decentralized management is less of an issue, ultimately making explicit effectiveness assessments easier and metrics such as a SHAPE score, less useful.

The scale at which that complexity horizon exists is likely hard to predict, highly variable, and species specific.

Numerous recent papers have called for improved data standardization, reporting and monitoring of restoration, and approaches to restoration that facilitate adaptive management leading to improved understanding of species and ecosystem responses to restoration (Bernhardt et al. 2005, Palmer et al. 2005, Beechie et al. 2008, Runge 2011). Indeed, increased data standardization and sharing will continue to improve the scientific study of restoration (Palmer et al. 2007, Palmer 2009, Dickens and Suding 2013), and importantly empower higher resolution analysis than is possible with a presence/absence metric such as the SHAPE score. In the absence of standardization, different assessment methods can lead to biases in habitat evaluation (Al-Chokhachy and Roper 2010). Spatially referenced assessments with consistent methods would greatly aid in analyses and future restoration planning for ESA listed species; generating such a data system would entail significant up front cost, but those costs would be offset to some degree by increased management efficiencies. With the present low-availability and high-cost restoration effectiveness data, our metrics address the near-term need for an accountability mechanism in decentralized endangered species habitat management. While designed with salmon in mind, these methods are generally applicable to any imperiled species with a habitat assessment or recovery plan that identifies habitat concerns and can to improve information accessibility for project planning and placement across the diversity of stakeholders involved in habitat restoration and conservation planning.

ACKNOWLEDGMENTS

We would like to thank George Pess, Jeff Jorgensen, Rodney Sayler, Matthew Carroll and Stephanie Hampton for thoughtful insights on earlier drafts.

LITERATURE CITED

- Al-Chokhachy, R., and B. B. Roper. 2010. Quantifying the effects of sampling error in stream habitat data on the conservation and management of salmonids. *Fisheries* 35:476–488.
- Allen, E. B., W. W. Covington, and D. A. Falk. 1997. Developing the conceptual basis for restoration ecology. *Restoration Ecology* 5:275–276.
- Barnas, K., and S. L. Katz. 2010. The challenges of tracking habitat restoration at various spatial scales. *Fisheries* 35:232–241.
- Bash, J. S., and C. M. Ryan. 2002. Stream restoration and enhancement projects: Is anyone monitoring? *Environmental Management* 29:877–885.
- Beechie, T., G. Pess, P. Roni, and G. Giannico. 2008. Setting river restoration priorities: a review of approaches and a general protocol for identifying and prioritizing actions. *North American Journal of Fisheries Management* 28:891–905.
- Beechie, T. J., D. A. Sear, J. D. Olden, G. R. Pess, J. M. Buffington, H. Moir, P. Roni, and M. M. Pollock. 2010. Process-based principles for restoring river ecosystems. *BioScience* 60:209–222.
- Berkes, F. 2007. Community-based conservation in a globalized world. *Proceedings of the National Academy of Sciences* 104:15188–15193.
- Berkes, F., D. Feeny, B. J. McCay, and J. M. Acheson. 1989. The benefits of the commons. *Nature* 340:91–93.
- Bernhardt, E. S., E. B. Sudduth, M. A. Palmer, J. D. Allan, J. L. Meyer, G. Alexander, J. Follstad-Shah, B. Hassett, R. Jenkinson, and R. Lave. 2007. Restoring rivers one reach at a time: results from a survey of US river restoration practitioners. *Restoration Ecology* 15:482–493.
- Bernhardt, E. S., et al. 2005. Synthesizing US river restoration efforts. *Science* 308:636–637.
- Bottrill, M. C., J. C. Walsh, J. E. Watson, L. N. Joseph, A. Ortega-Argueta, and H. P. Possingham. 2011. Does recovery planning improve the status of threatened species? *Biological Conservation* 144:1595–1601.
- Boyd, C. S., D. D. Johnson, J. D. Kerby, T. J. Svejcar, and K. W. Davies. 2014. Of grouse and golden eggs: Can ecosystems be managed within a species-based regulatory framework? *Rangeland Ecology and Management* 67:358–368.
- Carroll, C., D. J. Rohlf, Y.-W. Li, B. Hartl, M. K. Phillips, and R. F. Noss. 2014. Connectivity conservation and endangered species recovery: a study in the challenges of defining conservation-reliant species. *Conservation Letters*. doi: 10.1111/conl.12102
- Christian-Smith, J., and A. M. Merenlender. 2010. The disconnect between restoration goals and practices: a case study of watershed restoration in the Russian River Basin, California. *Restoration Ecology* 18:95–102.
- Clark, J. A., J. M. Hoekstra, P. D. Boersma, and P. Kareiva. 2002. Improving US Endangered Species Act recovery plans: key findings and recommendations of the SCB recovery plan project. *Conservation Biology* 16:1510–1519.
- Dickens, S. J. M., and K. N. Suding. 2013. Spanning the

- science-practice divide: why restoration scientists need to be more involved with practice. *Ecological Restoration* 31:134–140.
- G.A.O. 2002. Columbia River Basin salmon and steelhead: federal agencies' recovery responsibilities, expenditures and actions. U.S. General Accounting Office, Washington, D.C., USA.
- Halle, S. 2007. Science, art, or application—the “karma” of restoration ecology. *Restoration Ecology* 15:358–361.
- Hamm, D. E. 2012. Development and evaluation of a data dictionary to standardize salmonid habitat assessments in the Pacific Northwest. *Fisheries* 37:6–18.
- Katz, E. 2009. The big lie: human restoration of nature. Page 443 in D. M. Kaplan, editor. *Readings in the philosophy of technology*. Rowman and Littlefield, Lanham, Maryland, USA.
- Katz, S. L., K. Barnas, R. Hicks, J. Cowen, and R. Jenkinson. 2007. Freshwater habitat restoration actions in the Pacific Northwest: a decade's investment in habitat improvement. *Restoration Ecology* 15:494–505.
- Kondolf, G. M., P. L. Angermeier, K. Cummins, T. Dunne, M. Healey, W. Kimmerer, P. B. Moyle, D. Murphy, D. Patten, and S. Railsback. 2008. Projecting cumulative benefits of multiple river restoration projects: an example from the Sacramento-San Joaquin river system in California. *Environmental Management* 42:933–945.
- Lane, M. B., and T. Corbett. 2005. The tyranny of localism: indigenous participation in community-based environmental management. *Journal of Environmental Policy and Planning* 7:141–159.
- Lebel, L., A. Contreras, S. Pasing, and P. Garden. 2004. Nobody knows best: alternative perspectives on forest management and governance in Southeast Asia. *International Environmental Agreements* 4:111–127.
- Magurran, A. E. 2009. Threats to freshwater fish. *Science* 325:1215–1216.
- McCune, B., J. B. Grace, and D. L. Urban. 2002. *Analysis of ecological communities*. MjM Software Design, Gleneden Beach, Oregon, USA.
- Miller, J. R., and R. J. Hobbs. 2007. Habitat restoration—Do we know what we're doing? *Restoration Ecology* 15:382–390.
- Mullan, J. W., K. R. Williams, G. Rhodus, T. W. Hillman, and J. D. McIntyre. 1992. Production and habitat of salmonids in mid Columbia River tributary streams. U.S. Fish and Wildlife Service Monograph I:1–489.
- Mody, J. 2004. Achieving accountability through decentralization: lessons for integrated river basin management. World Bank Policy Research Working Paper 3346. World Bank, Washington, D.C., USA.
- National Research Council, Board on Environmental Studies and Toxicology, Committee on Protection and Management of Pacific Northwest Anadromous Salmonids, and Commission on Life Sciences. 1996. *Upstream: salmon and society in the Pacific Northwest*. National Academies Press, Washington, D.C., USA.
- NMFS [National Marine Fisheries Service]. 2000. Reinitiation of Operation of the Federal Columbia River Power System (FCRPS), Including the Juvenile Fish Transportation System, and 19 Bureau of Reclamation Projects in the Columbia Basin (COE.), Section 7 Consultation. NWR-1999-1901. Northwest Regional Office, Portland, Oregon, USA.
- NMFS [National Marine Fisheries Service]. 2014. Pacific Northwest salmon habitat project database. National Marine Fisheries Service, Northwest Fisheries Science Center. <https://www.webapps.nwfsc.noaa.gov/pnshp/>
- Northwest Power and Conservation Council. 2005. Columbia Cascade Subbasin Plans: Chelan, Entiat, Methow, Okanogan, Wenatchee, Upper Middle Columbia. In *Columbia River Basin Fish and Wildlife Program*. Northwest Power and Conservation Council, Portland, Oregon, USA. <https://www.nwccouncil.org/fw/subbasinplanning/home/>
- O'Donnell, T. K., and D. L. Galat. 2008. Evaluating success criteria and project monitoring in river enhancement within an adaptive management framework. *Environmental Management* 41:90–105.
- Palmer, M. A. 2009. Reforming watershed restoration: science in need of application and applications in need of science. *Estuaries and Coasts* 32:1–17.
- Palmer, M., J. D. Allan, J. Meyer, and E. S. Bernhardt. 2007. River restoration in the twenty-first century: data and experiential knowledge to inform future efforts. *Restoration Ecology* 15:472–481.
- Palmer, M. A., E. S. Bernhardt, J. D. Allan, P. S. Lake, G. Alexander, S. Brooks, J. Carr, S. Clayton, C. N. Dahm, and J. Follstad Shah. 2005. Standards for ecologically successful river restoration. *Journal of Applied Ecology* 42:208–217.
- Paulsen, C. M., and T. R. Fisher. 2005. Do habitat actions affect juvenile survival? An information-theoretic approach applied to endangered Snake River Chinook salmon. *Transactions of the American Fisheries Society* 134:68–85.
- Paulsen, C. M., and T. R. Fisher. 2003. Detecting juvenile survival effects of habitat actions: power analysis applied to endangered Snake River spring summer chinook (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* 60:1122–1132.
- Pinsky, M. L., D. B. Springmeyer, M. N. Goslin, and X. Augerot. 2009. Range-wide selection of catchments for Pacific salmon conservation. *Conservation*

- Biology 23:680–691.
- Prud'Homme, R. 1995. The dangers of decentralization. *World Bank Research Observer* 10:201–220.
- R Core Team. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Richter, B. D., D. P. Braun, M. A. Mendelson, and L. L. Master. 1997. Threats to imperiled freshwater fauna. *Conservation Biology* 11:1081–1093.
- Roni, P., K. Hanson, and T. Beechie. 2008. Global review of the physical and biological effectiveness of stream habitat rehabilitation techniques. *North American Journal of Fisheries Management* 28:856–890.
- Rumps, J. M., S. L. Katz, K. Barnas, M. D. Morehead, R. Jenkinson, S. R. Clayton, and P. Goodwin. 2007. Stream restoration in the Pacific Northwest: analysis of interviews with project managers. *Restoration Ecology* 15:506–515.
- Runge, M. C. 2011. An introduction to adaptive management for threatened and endangered species. *Journal of Fish and Wildlife Management* 2:220–233.
- Sewell, D. O. 1996. “The Dangers of Decentralization” according to Prud'homme: some further aspects. *World Bank Research Observer* 11:143–150.
- Sharma, R. 2003. Kerala's decentralization: idea in practice. *Economic and Political Weekly* September 6:3832–3850.
- Stanford Environmental Law Society. 2001. *The Endangered Species Act*. Stanford University Press, Stanford, California, USA.
- Wandschneider, P. R. 1984. Managing river systems: centralization versus decentralization. *Natural Resources Journal* 24:1043.
- Waples, R. S. 1991. Pacific salmon, *Oncorhynchus* spp., and the definition of “species” under the Endangered Species Act. *Marine Fisheries Review* 53:11–22.
- Williams, J. E., A. L. Haak, N. G. Gillespie, and W. T. Colyer. 2007. The Conservation Success Index: synthesizing and communicating salmonid condition and management needs. *Fisheries* 32:477–493.

SUPPLEMENTAL MATERIAL

APPENDIX A

Table A1. Broad crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 1–3 (see Hamm 2012 for definitions). In the Broad crosswalk we attributed projects to ecological concerns even if the likelihood of success was highly optimistic, contingent on large-scale implementation or on long-term maintenance for project effectiveness. Ecological Concern types are: 1, Habitat quantity; 1.1, Anthropogenic barriers; 1.2, Natural barriers; 2, Direct mortality; 2.1, Predation; 2.2, Pathogens; 3, Toxic contaminants; 3.1, Water; 3.2, Biota.

Project type and subtype	Ecological Concern type								
	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Barrier removal									
Culvert improvements/upgrade		X							
Culvert installation		X							
Culvert removal		X							
Culvert replacement		X							
Dam removal		X							
Push-up dam/diversion dam removal		X							
Fish by-pass		X							
Fish ladder		X							
Fish ladder improved		X							
Fish ladder installed		X							

Table A1. Continued.

Project type and subtype	Ecological Concern type								
	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Log jam/debris removal			X						
Other		X							
Tidegate		X							
Weir		X							
Diversion screens									
Fish screen				X					
Fish screen replacement				X					
Sediment reduction									
Erosion control structures									
Other									
Road closing/abandonment								X	
Road drainage								X	
Road relocation								X	
Rocked ford									
Sediment traps									
Restore stream complexity-channel complexity									
Bank stabilization									
Beaver introduction/management									
Channel connectivity		X							
Channel reconfiguration									
Dike reconfiguration									
Dike removal		X							
Off-channel habitat		X							
Off-channel habitat:alcove		X							
Off-channel habitat:pond									
Off-channel habitat:side channel		X							
Other									
Wetland creation									
Wetland improvement/enhancement									
Wetland restoration									
Restore stream complexity-instream structure									
Boulders									
Deflector									
Gravel placement									
Rock weir									
Log weir									
Weir									
Other									
Large woody debris									
Rootwads									
Structure/log jam									
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									
Restore instream flow									
Instream water rights		X							
Reduce/regulate water withdrawal		X							
Water quantity		X							
Restore riparian function									
Fencing								X	
Forestry practices									
Livestock removal								X	
Livestock rotation								X	
Livestock stream crossing									
Off-channel watering								X	
Other									
Plant installation/revegetation									
Plant removal/control									
Water quality improvement									
Other									
Refuse removal							X		
Temperature controls		X							
Toxic clean up							X		

Table A1. Continued.

Project type and subtype	Ecological Concern type								
	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Upland management									
Agriculture management									
Erosion structures									
Fencing								X	
Invasive plant control									
Livestock management								X	
Planting									
Slope stabilization									
Vegetation management									
Water development									
Other									
Other									
Bridge		X							
Culvert		X							
Fencing		X							
Flood control		X							
Irrigation diversion		X							
Other		X							
Road		X							
Water control structures		X							
Water development		X							
Wetland management		X							

Table A2. Broad crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 4–5 (see Hamm 2012 for definitions). In the Broad crosswalk we attributed projects to ecological concerns even if the likelihood of success was highly optimistic, contingent on large-scale implementation or on long-term maintenance for project effectiveness. Ecological Concern types are: 4, Food; 4.1, Altered primary productivity; 4.2, Competition; 4.3 Altered prey species composition and diversity; 5, Riparian; 5.1, Riparian condition; 5.2, LWD recruitment.

Project type and subtype	Ecological Concern type						
	4	4.1	4.2	4.3	5	5.1	5.2
Barrier removal							
Culvert improvements/upgrade							
Culvert installation							
Culvert removal							
Culvert replacement							
Dam removal							
Push-up dam/diversion dam removal							
Fish by-pass							
Fish ladder							
Fish ladder improved							
Fish ladder installed							
Log jam/debris removal							
Other							
Tidegate							
Weir							
Diversion screens							
Fish screen							
Fish screen replacement							
Sediment reduction							
Erosion control structures							
Other							
Road closing/abandonment							
Road drainage							
Road relocation							

Table A2. Continued.

Project type and subtype	Ecological Concern type						
	4	4.1	4.2	4.3	5	5.1	5.2
Rocked ford							
Sediment traps							
Restore stream complexity-channel complexity							
Bank stabilization							
Beaver introduction/management		X			X		
Channel connectivity		X			X		
Channel reconfiguration		X				X	
Dike reconfiguration		X				X	
Dike removal		X				X	
Off-channel habitat		X			X		
Off-channel habitat:alcove		X			X		
Off-channel habitat:pond		X			X		
Off-channel habitat:side channel		X			X		
Other					X		
Wetland creation		X					
Wetland improvement/enhancement		X					
Wetland restoration		X					
Restore stream complexity-instream structure							
Boulders							
Deflector							
Gravel placement							
Rock weir							
Log weir							
Weir							
Other							
Large woody debris							
Rootwads							
Structure/log jam							
Nutrient enrichment							
Carcass placement	X				X		
Fertilizer	X				X		
Other	X				X		
Restore instream flow							
Instream water rights					X		
Reduce/Regulate water withdrawal					X		
Water quantity					X		
Restore riparian function							
Fencing					X		
Forestry practices					X		
Livestock removal		X			X		
Livestock rotation		X			X		
Livestock stream crossing		X			X		
Off-channel watering		X			X		
Other		X			X		
Plant installation/revegetation		X		X	X		
Plant removal/control		X		X	X		
Water quality improvement							
Other							
Refuse removal							
Temperature controls							
Toxic clean up							
Upland management							
Agriculture management							
Erosion structures							
Fencing					X		
Invasive plant control	X			X	X		
Livestock management					X		
Planting					X		
Slope stabilization					X		
Vegetation management					X		
Water development					X		
Other							
Other							
Bridge							
Culvert							
Fencing							

Table A2. Continued.

Project type and subtype	Ecological Concern type						
	4	4.1	4.2	4.3	5	5.1	5.2
Flood control							
Irrigation diversion							
Other							
Road							
Water control structures							
Water development							
Wetland management							

Table A3. Broad crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 6–8 (see Hamm 2012 for definitions). In the Broad crosswalk we attributed projects to ecological concerns even if the likelihood of success was highly optimistic, contingent on large-scale implementation or on long-term maintenance for project effectiveness. Ecological Concern types are: 6, Peripheral habitat; 6.1, Side channel and wetland conditions; 6.2, Floodplain condition; 7, Channel structure and form; 7.1, Bed and channel form; 7.2, Instream structural complexity; 8, Sediment conditions; 8.1, Decreased sediment quantity; 8.2, Increased sediment quantity.

Project type and subtype	Ecological Concern type								
	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Barrier removal									
Culvert improvements/upgrade									
Culvert installation									
Culvert removal									
Culvert replacement									
Dam removal							X		
Push-up dam/diversion dam removal							X		
Fish by-pass									
Fish ladder									
Fish ladder improved									
Fish ladder installed									
Log jam/debris removal									
Other									
Tidegate									
Weir									
Diversion screens									
Fish screen									
Fish screen replacement									
Sediment reduction									
Erosion control structures						X	X		X
Other						X	X		X
Road closing/abandonment	X					X	X		X
Road drainage				X		X	X		
Road relocation	X					X	X		X
Rocked ford						X	X		
Sediment traps				X		X	X		X
Restore stream complexity-channel complexity									
Bank stabilization					X	X	X		
Beaver introduction/management	X			X	X	X			
Channel connectivity	X			X	X	X	X		
Channel reconfiguration	X			X	X	X	X		
Dike reconfiguration	X				X				
Dike removal	X				X		X		
Off-channel habitat	X				X	X	X		
Off-channel habitat:alcove		X	X		X	X	X		
Off-channel habitat:pond		X	X		X	X	X		
Off-channel habitat:side channel		X	X		X	X	X		
Other	X			X	X	X	X		
Wetland creation		X			X				

Table A3. Continued.

Project type and subtype	Ecological Concern type								
	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Wetland improvement/enhancement		X			X				
Wetland restoration		X			X				
Restore stream complexity-instream structure									
Boulders					X	X	X		
Deflector					X	X	X		X
Gravel placement					X	X	X		
Rock weir					X	X	X		
Log weir					X	X	X		
Weir					X	X	X		
Other				X	X	X	X		
Large woody debris					X	X	X		
Rootwads					X	X	X		
Structure/log jam					X	X	X		
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									
Restore instream flow									
Instream water rights		X	X				X		X
Reduce/Regulate water withdrawal		X	X				X		X
Water quantity		X	X				X		X
Restore riparian function									
Fencing		X			X		X		
Forestry practices							X		
Livestock removal		X			X		X		
Livestock rotation		X			X		X		
Livestock stream crossing					X		X		
Off-channel watering		X			X		8		
Other		X					X		
Plant installation/revegetation		X			X	X	X		
Plant removal/control		X				X			
Water quality improvement									
Other									
Refuse removal									
Temperature controls									
Toxic clean up									
Upland management									
Agriculture management							X		X
Erosion structures							X		X
Fencing		X			X		X		X
Invasive plant control		X							
Livestock management		X			X		X		X
Planting		X					X		
Slope stabilization		X		X			X		
Vegetation management		X		X			X		
Water development		X			X		X		
Other							X		
Other									
Bridge									
Culvert									
Fencing									
Flood control									
Irrigation diversion							8		
Other							X		
Road									
Water control structures									
Water development									
Wetland management									

Table A4. Broad crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 9–10 (see Hamm 2012 for definitions). In the Broad crosswalk we attributed projects to ecological concerns even if the likelihood of success was highly optimistic, contingent on large-scale implementation or on long-term maintenance for project effectiveness. Ecological Concern types are: 9, Water quality; 9.1, Temperature; 9.2, Oxygen; 9.3, Turbidity; 9.4, pH; 9.5, Salinity; 10, Water quantity; 10.1, Increased water quantity; 10.2, Decreased water quantity; 10.3, Altered flow timing.

Project type and subtype	9	9.1	9.2	9.3	9.4	9.5	10	10.1	10.2	10.3
Barrier removal										
Culvert improvements/upgrade										
Culvert installation										
Culvert removal										
Culvert replacement										
Dam removal	9									
Push-up dam/diversion dam removal	9									
Fish By-pass										
Fish ladder										
Fish ladder improved										
Fish ladder installed										
Log jam/debris removal										
Other										
Tidegate	9									
Weir										
Diversion screens										
Fish screen										
Fish screen replacement										
Sediment reduction										
Erosion control structures				X						
Other				X						
Road closing/abandonment				X			X			
Road drainage				X			X			
Road relocation				X			X			
Rocked ford				X						
Sediment traps				X						
Restore stream complexity-channel complexity										
Bank stabilization		X		X						
Beaver introduction/management		X	X	X			X			
Channel connectivity		X	X	X			X			
Channel reconfiguration		X	X	X			X			
Dike reconfiguration		X	X	X			X			
Dike removal		X	X	X			X			
Off-channel habitat		X	X	X			X			
Off-channel habitat:alcove		X	X	X			X			
Off-channel habitat:pond		X	X	X			X			
Off-channel habitat:side channel		X	X	X			X			
Other		X	X	X			X			
Wetland creation		X	X	X			X			
Wetland improvement/enhancement		X	X	X			X			
Wetland restoration		X	X	X			X			
Restore stream complexity-instream structure										
Boulders										
Deflector										
Gravel placement										
Rock weir										
Log weir										
Weir										
Other										
Large woody debris		X								
Rootwads		X								
Structure/log jam		X								
Nutrient enrichment										
Carcass placement										
Fertilizer										
Other										

Table A4. Continued.

Project type and subtype	9	9.1	9.2	9.3	9.4	9.5	10	10.1	10.2	10.3
Restore instream flow										
Instream water rights	9								X	
Reduce/regulate water withdrawal	9								X	
Water quantity	9								X	
Restore riparian function										
Fencing		X	X	X						
Forestry practices		X	X	X			X			
Livestock removal		X	X	X						
Livestock rotation		X	X	X						
Livestock stream crossing				X						
Off-channel watering		X	X	X						
Other		X		X						
Plant installation/revegetation		X		X			X			
Plant removal/control		X								
Water quality improvement										
Other	9									
Refuse removal										
Temperature controls		X								
Toxic clean up										
Upland management										
Agriculture management		X	X	X			X			
Erosion structures		X	X	X						
Fencing		X	X	X						
Invasive plant control										
Livestock management		X	X	X						
Planting		X	X	X			X			
Slope stabilization		X	X	X			X			
Vegetation management		X	X	X			X			
Water development		X	X	X						
Other										
Other										
Bridge										
Culvert										
Fencing										
Flood control										
Irrigation diversion	9									
Other	9									
Road										
Water control structures										
Water development										
Wetland management										

APPENDIX B

Table B1. Narrow crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 1–3 (see Hamm 2012 for definitions). In the Narrow crosswalk we attributed project types with addressing only the most probable goal. Ecological Concern types are: 1, Habitat quantity; 1.1, Anthropogenic barriers; 1.2, Natural barriers; 2, Direct mortality; 2.1, Predation; 2.2, Pathogens; 3, Toxic contaminants; 3.1, Water; 3.2, Biota.

Project type and subtype	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Barrier removal									
Culvert improvements/upgrade		X							
Culvert installation		X							
Culvert removal		X							
Culvert replacement		X							
Dam removal		X							
Push-up dam/diversion dam removal		X							
Fish by-pass		X							
Fish ladder		X							
Fish ladder improved		X							
Fish ladder installed		X							
Log jam/debris removal			X						
Other									
Tidegate		X							
Weir		X							
Diversion screens									
Fish screen				X					
Fish screen replacement				X					
Sediment reduction									
Erosion control structures									
Other									
Road closing/abandonment									
Road drainage									
Road relocation									
Rocked ford									
Sediment traps									
Restore stream complexity-channel complexity									
Bank stabilization									
Beaver introduction/management									
Channel connectivity		X							
Channel reconfiguration									
Dike reconfiguration									
Dike removal		X							
Off-channel habitat		X							
Off-channel habitat:alcove		X							
Off-channel habitat:pond									
Off-channel habitat:side channel		X							
Other									
Wetland creation									
Wetland improvement/enhancement									
Wetland restoration									
Restore stream complexity-instream structure									
Boulders									
Deflector									
Gravel placement									
Rock weir									
Log weir									
Weir									
Other									
Large woody debris									
Rootwads									
Structure/Log jam									
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									

Table B1. Continued.

Project type and subtype	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Restore instream flow									
Instream water rights									
Reduce/regulate water withdrawal									
Water quantity									
Restore riparian function									
Fencing								X	
Forestry practices									
Livestock removal									
Livestock rotation									
Livestock stream crossing									
Off-channel watering									
Other									
Plant installation/revegetation									
Plant removal/control									
Water quality improvement									
Other									
Refuse removal									
Temperature controls									
Toxic clean up							X		
Upland management									
Agriculture management									
Erosion structures									
Fencing									
Invasive plant control									
Livestock management									
Planting									
Slope stabilization									
Vegetation management									
Water development									
Other									
Other									
Bridge									
Culvert									
Fencing									
Flood control									
Irrigation diversion									
Other									
Road									
Water control structures									
Water development									
Wetland management									

Table B2. Narrow crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 4–5 (see Hamm 2012 for definitions). In the Narrow crosswalk we attributed project types with addressing only the most probable goal. Ecological Concern types are: 4, Food; 4.1, Altered primary productivity; 4.2, Competition; 4.3 Altered prey species composition and diversity; 5, Riparian; 5.1, Riparian condition; 5.2, LWD recruitment.

Project type and subtype	4	4.1	4.2	4.3	5	5.1	5.2
Barrier removal							
Culvert improvements/upgrade							
Culvert installation							
Culvert removal							
Culvert replacement							
Dam removal							
Push-up dam/diversion dam removal							
Fish By-pass							
Fish ladder							
Fish ladder improved							
Fish ladder installed							
Log jam/debris removal							
Other							
Tidegate							
Weir							
Diversion screens							
Fish screen							
Fish screen replacement							
Sediment reduction							
Erosion control structures							
Other							
Road closing/abandonment							
Road drainage							
Road relocation							
Rocked ford							
Sediment traps							
Restore stream complexity-channel complexity							
Bank stabilization							
Beaver introduction/management					X		
Channel connectivity					X		
Channel reconfiguration							
Dike reconfiguration						X	
Dike removal						X	
Off-channel habitat							
Off-channel habitat:alcove							
Off-channel habitat:pond							
Off-channel habitat:side channel							
Other							
Wetland creation							
Wetland improvement/enhancement							
Wetland restoration							
Restore stream complexity-instream structure							
Boulders							
Deflector							
Gravel placement							
Rock weir							
Log weir							
Weir							
Other							
Large woody debris							
Rootwads							
Structure/log jam							
Nutrient enrichment							
Carcass placement		X					
Fertilizer		X					
Other		X					
Restore instream flow							
Instream water rights							
Reduce/regulate water withdrawal							
Water quantity							

Table B2. Continued.

Project type and subtype	4	4.1	4.2	4.3	5	5.1	5.2
Restore riparian function							
Fencing					X		
Forestry practices					X		
Livestock removal					X		
Livestock rotation					X		
Livestock stream crossing					X		
Off-channel watering					X		
Other							
Plant installation/revegetation					X		
Plant removal/control				X	X		
Water quality improvement							
Other							
Refuse removal							
Temperature controls							
Toxic clean up							
Upland management							
Agriculture management							
Erosion structures							
Fencing							
Invasive plant control				X			
Livestock management							
Planting							
Slope stabilization							
Vegetation management							
Water development							
Other							
Other							
Bridge							
Culvert							
Fencing							
Flood control							
Irrigation diversion							
Other							
Road							
Water control structures							
Water development							
Wetland management							

Table B3. Narrow crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 6–8 (see Hamm 2012 for definitions). In the Narrow crosswalk we attributed project types with addressing only the most probable goal. Ecological Concern types are: 6, Peripheral habitat; 6.1, Side channel and wetland conditions; 6.2, Floodplain condition; 7, Channel structure and form; 7.1, Bed and channel form; 7.2, Instream structural complexity; 8, Sediment conditions; 8.1, Decreased sediment quantity; 8.2, Increased sediment quantity.

Project type and subtype	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Barrier removal									
Culvert improvements/upgrade									
Culvert installation									
Culvert removal									
Culvert replacement									
Dam removal									
Push-up dam/diversion dam removal									
Fish by-pass									
Fish ladder									
Fish ladder improved									
Fish ladder installed									
Log jam/debris removal									
Other									
Tidegate									
Weir									
Diversion screens									
Fish screen									
Fish screen replacement									
Sediment reduction									
Erosion control structures							X		
Other									
Road closing/abandonment							X		
Road drainage							X		
Road relocation							X		
Rocked ford							X		
Sediment traps							X		
Restore stream complexity-channel complexity									
Bank stabilization					X	X	X		
Beaver introduction/management	X				X				
Channel connectivity	X				X				
Channel reconfiguration				X	X	X			
Dike reconfiguration		X	X		X				
Dike removal		X	X		X				
Off-channel habitat	X				X				
Off-channel habitat:alcove			X		X				
Off-channel habitat:pond			X		X				
Off-channel habitat:side channel			X		X				
Other					X				
Wetland creation		X							
Wetland improvement/enhancement		X							
Wetland restoration		X							
Restore stream complexity-instream structure									
Boulders					X	X			
Deflector					X	X			X
Gravel placement									
Rock weir					X	X			
Log weir					X	X			
Weir					X	X			
Other									
Large woody debris					X	X			
Rootwads					X	X			
Structure/log jam					X	X			
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									
Restore instream flow									
Instream water rights		X							
Reduce/regulate water withdrawal		X							

Table B3. Continued.

Project type and subtype	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Water quantity									
Restore riparian function									
Fencing		X			X		X		
Forestry practices		X					X		
Livestock removal		X			X		X		
Livestock rotation		X			X		X		
Livestock stream crossing					X		X		
Off-channel watering		X			X		X		
Other							X		
Plant installation/revegetation		X			X	X	X		
Plant removal/control		X				X			
Water quality improvement									
Other									
Refuse removal									
Temperature controls									
Toxic clean up									
Upland management									
Agriculture management							X		
Erosion structures							X		
Fencing		X			X		X		
Invasive plant control									
Livestock management		X			X		X		
Planting							X		
Slope stabilization							X		X
Vegetation management							X		X
Water development		X			X		X		
Other									
Other									
Bridge									
Culvert									
Fencing									
Flood control									
Irrigation diversion									
Other									
Road									
Water control structures									
Water development									
Wetland management									

Table B4. Narrow crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 9–10 (see Hamm 2012 for definitions). In the Narrow crosswalk we attributed project types with addressing only the most probable goal. Ecological Concern types are: 9, Water quality; 9.1, Temperature; 9.2, Oxygen; 9.3, Turbidity; 9.4, pH; 9.5, Salinity; 10, Water quantity; 10.1, Increased water quantity; 10.2, Decreased water quantity; 10.3, Altered flow timing.

Project type and subtype	9	9.1	9.2	9.3	9	9.5	10	10.1	10.2	10.3
Barrier removal										
Culvert improvements/upgrade										
Culvert installation										
Culvert removal										
Culvert replacement										
Dam removal										
Push-up dam/diversion dam removal										
Fish by-pass										
Fish ladder										
Fish ladder improved										
Fish ladder installed										
Log jam/debris removal										
Other										
Tidegate										
Weir										
Diversion screens										
Fish screen										
Fish screen replacement										
Sediment reduction										
Erosion control structures			X	X						
Other										
Road closing/abandonment				X				X		
Road drainage				X				X		
Road relocation				X				X		
Rocked ford				X						
Sediment traps				X						
Restore stream complexity-channel complexity										
Bank stabilization				X						
Beaver introduction/management							X			
Channel connectivity								X		
Channel reconfiguration								X		
Dike reconfiguration								X		
Dike removal								X		
Off-channel habitat								X		
Off-channel habitat:alcove								X		
Off-channel habitat:pond								X		
Off-channel habitat:side channel								X		
Other										
Wetland creation										
Wetland improvement/enhancement										
Wetland restoration										
Restore stream complexity-instream structure										
Boulders										
Deflector										
Gravel placement										
Rock weir										
Log weir										
Weir										
Other										
Large woody debris										
Rootwads										
Structure/log jam										
Nutrient enrichment										
Carcass placement										
Fertilizer										
Other										
Restore instream flow										
Instream water rights	X								X	
Reduce/regulate water withdrawal	X								X	
Water quantity	X								X	

Table B4. Continued.

Project type and subtype	9	9.1	9.2	9.3	9	9.5	10	10.1	10.2	10.3
Restore riparian function										
Fencing		X		X						
Forestry practices		X		X				X		
Livestock removal		X		X						
Livestock rotation		X		X						
Livestock stream crossing		X		X						
Off-channel watering		X		X						
Other										
Plant installation/revegetation		X		X				X		
Plant removal/control		X								
Water quality improvement										
Other										
Refuse removal										
Temperature controls		X								
Toxic clean up										
Upland management										
Agriculture management		X	X	X						X
Erosion structures		X	X	X						
Fencing		X	X	X						
Invasive plant control										
Livestock management		X	X	X						
Planting				X				X		
Slope stabilization		X		X						
Vegetation management		X		X						
Water development		X	X	X						
Other										
Other										
Bridge										
Culvert										
Fencing										
Flood control										
Irrigation diversion										
Other										
Road										
Water control structures										
Water development										
Wetland management										

APPENDIX C

Table C1. The Intermediate crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 1–3 (see Hamm 2012 for definitions). The Intermediate crosswalk attributed projects with all concerns suggested by the literature unless cause and effect are separated by many steps, require coordinated implementation across large spatial scales or require time scales of decades for success. Ecological Concern types are: 1, Habitat quantity; 1.1, Anthropogenic barriers; 1.2, Natural barriers; 2, Direct mortality; 2.1, Predation; 2.2, Pathogens; 3, Toxic contaminants; 3.1, Water; 3.2, Biota.

Project type and subtype	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Barrier removal									
Culvert improvements/upgrade		X							
Culvert installation		X							
Culvert removal		X							
Culvert replacement		X							
Dam removal		X							
Push-up dam/diversion dam removal		X							
Fish By-pass		X							
Fish ladder		X							
Fish ladder improved		X							
Fish ladder installed		X							
Log jam/debris removal			X						
Other	X								
Tidegate		X							
Weir		X							
Diversion screens									
Fish screen				X					
Fish screen replacement				X					
Sediment reduction									
Erosion control structures									
Other									
Road closing/abandonment									
Road drainage									
Road relocation									
Rocked ford									
Sediment traps									
Restore stream complexity-channel complexity									
Bank stabilization									
Beaver introduction/management					X				
Channel connectivity		X			X				
Channel reconfiguration									
Dike reconfiguration									
Dike removal									
Off-channel habitat					X				
Off-channel habitat:alcove					X				
Off-channel habitat:pond					X				
Off-channel habitat:side channel					X				
Other									
Wetland creation									
Wetland improvement/enhancement									
Wetland restoration									
Restore stream complexity-instream structure									
Boulders									
Deflector									
Gravel placement									
Rock weir									
Log weir									
Weir									
Other									
Large woody debris					X				
Rootwads					X				
Structure/log jam					X				

Table C1. Continued.

Project type and subtype	1	1.1	1.2	2	2.1	2.2	3	3.1	3.2
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									
Restore instream flow									
Instream water rights									
Reduce/regulate water withdrawal									
Water quantity									
Fencing									
Forestry practices									
Livestock removal									
Livestock rotation									
Livestock stream crossing									
Off-channel watering									
Other									
Plant installation/revegetation									
Plant removal/control									
Water quality improvement									
Other									
Refuse removal							X		
Temperature controls									
Toxic clean up							X		
Upland management									
Agriculture management									
Erosion structures									
Fencing									
Invasive plant control									
Livestock management									
Planting									
Slope stabilization									
Vegetation management									
Water development									
Other									
Other									
Bridge									
Culvert									
Fencing									
Flood control									
Irrigation diversion									
Other									
Road									
Water control structures									
Water development									
Wetland management									

Table C2. The Intermediate crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 4–5 (see Hamm 2012 for definitions). The Intermediate crosswalk attributed projects with all concerns suggested by the literature unless cause and effect are separated by many steps, require coordinated implementation across large spatial scales or require time scales of decades for success. Ecological Concern types are: 4, Food; 4.1, Altered primary productivity; 4.2, Competition; 4.3 Altered prey species composition and diversity; 5, Riparian; 5.1, Riparian condition; 5.2, LWD recruitment.

Project type and subtype	4	4.1	4.2	4.3	5	5.1	5.2
Barrier removal							
Culvert improvements/upgrade							
Culvert installation							
Culvert removal							
Culvert replacement							
Dam removal							
Push-up dam/diversion dam removal							
Fish by-pass							
Fish ladder							
Fish ladder improved							
Fish ladder installed							
Log jam/debris removal							
Other							
Tidegate							
Weir							
Diversion screens							
Fish screen							
Fish screen replacement							
Sediment reduction							
Erosion control structures							
Other							
Road closing/abandonment							
Road drainage							
Road relocation							
Rocked ford							
Sediment traps							
Restore stream complexity-channel complexity							
Bank stabilization							
Beaver introduction/management							
Channel connectivity	X						
Channel reconfiguration							
Dike reconfiguration						X	
Dike removal						X	
Off-channel habitat							
Off-channel habitat:alcove							
Off-channel habitat:pond							
Off-channel habitat:side channel							
Other							
Wetland creation	X						
Wetland improvement/enhancement	X						
Wetland restoration	X						
Restore stream complexity-instream structure							
Boulders							
Deflector							
Gravel placement							
Rock weir							
Log weir							
Weir							
Other							
Large woody debris							
Rootwads							
Structure/log jam							
Nutrient enrichment							
Carcass placement		X				X	
Fertilizer		X				X	
Other		X				X	

Table C2. Continued.

Project type and subtype	4	4.1	4.2	4.3	5	5.1	5.2
Restore instream flow							
Instream water rights							
Reduce/regulate water withdrawal							
Water quantity							
Restore riparian function							
Fencing						X	
Forestry practices						X	
Livestock removal						X	
Livestock rotation						X	
Livestock stream crossing						X	
Off-channel watering						X	
Other						X	
Plant installation/revegetation						X	
Plant removal/control						X	
Water quality improvement							
Other							
Refuse removal							
Temperature controls							
Toxic clean up							
Upland management							
Agriculture management							
Erosion structures							
Fencing						X	
Invasive plant control				X		X	
Livestock management						X	
Planting							
Slope stabilization							
Vegetation management							
Water development						X	
Other							
Other							
Bridge							
Culvert							
Fencing							
Flood control							
Irrigation diversion							
Other							
Road							
Water control structures							
Water development							
Wetland management							

Table C3. The Intermediate crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 6–8 (see Hamm 2012 for definitions). The Intermediate crosswalk attributed projects with all concerns suggested by the literature unless cause and effect are separated by many steps, require coordinated implementation across large spatial scales or require time scales of decades for success. Ecological Concern types are: 6, Peripheral habitat; 6.1, Side channel and wetland conditions; 6.2, Floodplain condition; 7, Channel structure and form; 7.1, Bed and channel form; 7.2, Instream structural complexity; 8, Sediment conditions; 8.1, Decreased sediment quantity; 8.2, Increased sediment quantity.

Project type and subtype	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Barrier removal									
Culvert improvements/upgrade									
Culvert installation									
Culvert removal									
Culvert replacement									
Dam removal								X	
Push-up dam/diversion dam removal								X	
Fish by-pass									
Fish ladder									
Fish ladder improved									
Fish ladder installed									
Log jam/debris removal									
Other									
Tidegate									
Weir									
Diversion screens									
Fish screen									
Fish screen replacement									
Sediment reduction									
Erosion control structures							X		
Other							X		
Road closing/abandonment							X		
Road drainage							X		
Road relocation							X		
Rocked ford							X		
Sediment traps							X		
Restore stream complexity-channel complexity									
Bank stabilization							X		
Beaver introduction/management	X			X			X		
Channel connectivity	X								
Channel reconfiguration	X				X		X		
Dike reconfiguration		X	X		X				
Dike removal		X	X		X				
Off-channel habitat	X								
Off-channel habitat:alcove		X			X				
Off-channel habitat:pond		X			X				
Off-channel habitat:side channel		X	X		X				
Other					X				
Wetland creation		X							
Wetland improvement/enhancement		X							
Wetland restoration		X							
Restore stream complexity-instream structure									
Boulders					X	X	X		
Deflector					X	X	X		
Gravel placement						X	X		
Rock weir					X	X	X		
Log weir					X	X	X		
Weir					X	X	X		
Other					X	X	X		
Large woody debris			X		X	X	X		
Rootwads			X		X	X	X		
Structure/log jam			X		X	X	X		
Nutrient enrichment									
Carcass placement									
Fertilizer									
Other									

Table C3. Continued.

Project type and subtype	6	6.1	6.2	7	7.1	7.2	8	8.1	8.2
Restore instream flow									
Instream water rights									
Reduce/regulate water withdrawal									
Water quantity									
Restore riparian function									
Fencing							X		
Forestry practices							X		
Livestock removal							X		
Livestock rotation							X		
Livestock stream crossing							X		
Off-channel watering							X		
Other									
Plant installation/revegetation									
Plant removal/control									
Water quality improvement									
Other									
Refuse removal									
Temperature controls									
Toxic clean up									
Upland management									
Agriculture management							X		
Erosion structures							X		
Fencing							X		
Invasive plant control									
Livestock management							X		
Planting							X		
Slope stabilization							X		
Vegetation management							X		
Water development							X		
Other									
Other									
Bridge									
Culvert									
Fencing									
Flood control									
Irrigation diversion									
Other									
Road									
Water control structures									
Water development									
Wetland management									

Table C4. The Intermediate crosswalk table relating the Pacific Northwest Salmon Habitat Project Database project types and subtypes (see <https://www.webapps.nwfsc.noaa.gov/pnshp/> for definitions) to Ecological Concern types 9–10 (see Hamm 2012 for definitions). The Intermediate crosswalk attributed projects with all concerns suggested by the literature unless cause and effect are separated by many steps, require coordinated implementation across large spatial scales or require time scales of decades for success. Ecological Concern types are: 9, Water quality; 9.1, Temperature; 9.2, Oxygen; 9.3, Turbidity; 9.4, pH; 9.5, Salinity; 10, Water quantity; 10.1, Increased water quantity; 10.2, Decreased water quantity; 10.3, Altered flow timing.

Project type and subtype	9	9.1	9.2	9.3	9.4	9.5	10	10.1	10.2	10.3
Barrier removal										
Culvert improvements/upgrade										
Culvert installation										
Culvert removal										
Culvert replacement										
Dam removal										
Push-up dam/diversion dam removal										
Fish by-pass										
Fish ladder										
Fish ladder improved										
Fish ladder installed										
Log jam/debris removal										
Other										
Tidegate										
Weir										
Diversion screens										
Fish screen										
Fish screen replacement										
Sediment reduction										
Erosion control structures				X						
Other										
Road closing/abandonment										
Road drainage				X						
Road relocation				X						
Rocked ford				X						
Sediment traps				X						
Restore stream complexity-channel complexity										
Bank stabilization				X						
Beaver introduction/management				X				X		X
Channel connectivity		X								
Channel reconfiguration		X								
Dike reconfiguration								X		X
Dike removal								X		X
Off-channel habitat		X						X		
Off-channel habitat:alcove		X						X		
Off-channel habitat:pond		X						X		
Off-channel habitat:side channel		X						X		
Other										
Wetland creation										
Wetland improvement/enhancement										
Wetland restoration										
Restore stream complexity-instream structure										
Boulders										
Deflector		X								
Gravel placement										
Rock weir		X								
Log weir		X								
Weir		X								
Other										
Large woody debris		X						X		
Restore stream complexity-instream structure										
Rootwads		X						X		
Structure/log jam		X						X		
Nutrient enrichment										
Carcass placement										
Fertilizer										
Other										

Table C4. Continued.

Project type and subtype	9	9.1	9.2	9.3	9.4	9.5	10	10.1	10.2	10.3
Restore instream flow										
Instream water rights	X								X	
Reduce/regulate water withdrawal	X								X	
Water quantity	X								X	X
Restore riparian function										
Fencing				X						
Forestry practices				X			X			
Livestock removal				X						
Livestock rotation				X						
Livestock stream crossing				X						
Off-channel watering				X						
Other										
Plant installation/revegetation										
Plant removal/control										
Water quality improvement										
Other	X									
Refuse removal										
Temperature controls		X								
Toxic clean up										
Upland management										
Agriculture management				X						
Erosion structures				X						
Fencing				X						
Invasive plant control										
Livestock management				X						
Planting				X						
Slope stabilization				X						
Vegetation management				X						
Water development	X								X	
Other				X						
Other										
Bridge										
Culvert										
Fencing										
Flood control										
Irrigation diversion										
Other										
Road										
Water control structures										
Water development										
Wetland management										

APPENDIX D

Table D1. Example table of how we applied the crosswalk comparison rules.

Assessment unit	Ecological concerns for assessment unit	Ecological concern(s) addressed by restoration projects	Project(s) matched	Project(s) not matched	Concern(s) matched	Concern(s) not matched
Example 1	1, 2.1, 2.2, 3, 3.1	Proj 1: 1	X		1, 2.1, 2.2	
		Proj 2: 2.1, 2.2	X			
		Proj 3: 4, 3.1	X			
Example 2	1, 1.1, 4, 5, 5.2	Proj 1: 1.1	X		1, 1.1	5, 5.2
		Proj 2: 3, 3.1		X		
		Proj 3: 3, 4.1	X			
Example 3	2.1, 4	Proj 1: 1, 4	X		4	2.1
		Proj 2: 4	X			
		Proj 3: 2.2		X		