

Banded Rainbowfish (*Melanotaenia trifasciata*)

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, Web Version – 1/29/2018



Photo: OpenCage. Licensed under Creative Commons BY 2.5.

1 Native Range and Status in the United States

Native Range

From Froese and Pauly (2016):

“Oceania: Northern Territory and Queensland in Australia.”

Status in the United States

No records of *Melanotaenia trifasciata* in the United States were found.

Means of Introductions in the United States

No records of *Melanotaenia trifasciata* in the United States were found.

Remarks

No additional remarks.

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2016):

“Taxonomic Status:

Current Standing: valid”

“Kingdom Animalia

Subkingdom Bilateria

Infrakingdom Deuterostomia

Phylum Chordata

Subphylum Vertebrata

Infraphylum Gnathostomata

Superclass Osteichthyes

Class Actinopterygii

Subclass Neopterygii

Infraclass Teleostei

Superorder Acanthopterygii

Order Atheriniformes

Suborder Atherinoidei

Family Melanotaeniidae

Genus *Melanotaenia*

Species *Melanotaenia trifasciata* (Rendahl, 1922)”

From Eschmeyer et al. (2017):

“***trifasciata*, *Rhombosoma*** Rendahl [H.] 1922:182 [Nyt Magazin for Naturvidenskaberne v. 60 [...]] Mary River, Northern Territory, Australia. Holotype (unique): ZMUO J995. Type catalog: Pethon 1969:6. •Valid as *Melanotaenia trifasciata* (Rendahl 1922) -- (Allen & Cross 1982:61 [...], Paxton et al. 1989:352 [...], Larson & Martin 1990:27 [...], Unmack 2001:1061 [...], Allen et al. 2002:159 [...], Allen et al. 2006:694 [...], Allen et al. 2008:90 [...], Larson et al. 2013:61 [...], Allen et al. 2015:103 [...], Allen et al. 2015:103 [...], Martin & Barclay 2015:54 [...], Allen et al. 2016:4 [...]). **Current status:** Valid as *Melanotaenia trifasciata* (Rendahl 1922). Melanotaeniidae.”

Size, Weight, and Age Range

From Froese and Pauly (2016):

“Max length : 13.0 cm SL male/unsexed; [Allen et al. 2002]; common length: 8.0 cm SL male/unsexed; [Allen et al. 2002]”

Environment

From Froese and Pauly (2016):

“Freshwater; benthopelagic; pH range: 7.0 - 8.0; dH range: 9 - 19. [...]; 25°C - 30°C [assumed to be recommended aquarium temperature range] [Baensch and Riehl 1985]”

“Occurs [in the wild] at [water] temperature and pH ranging from 22°-30° C and 5.2-7.0 respectively.”

Climate/Range

From Froese and Pauly (2016):

“Tropical; [...]”

Distribution Outside the United States

Native

From Froese and Pauly (2016):

“Oceania: Northern Territory and Queensland in Australia.”

Introduced

No records of *Melanotaenia trifasciata* introductions were found.

Means of Introduction Outside the United States

No records of *Melanotaenia trifasciata* introductions were found.

Short Description

A short description of *Melanotaenia trifasciata* was not available.

Biology

From Froese and Pauly (2016):

“Distinct pairing [Breder and Rosen 1966].”

“Lives mainly in small streams and waterholes in clear to moderately turbid water. Occurs over rocky or gravel bottoms or in well-vegetated areas, frequently around submerged logs and branches [...]. Inhabits small creeks and waterholes in semi-rainforest or open grassland areas.

Occasionally abundant in partly stagnant waterholes with leaf litter bottoms during the dry season.”

Human Uses

From Froese and Pauly (2016):

“Species is easy to maintain and thrives in captivity [Allen et al. 2002].”

Diseases

No records of OIE reportable diseases were found.

From Corlis (2004):

“[...] *I[liocirrus]. trifasciatae*, *H[elicirrus]. mcivori* and *L[ongidigitis]. hopevalensis* [collected] from *M. trifasciata* (Rendahl, 1922) [...]”

Threat to Humans

From Froese and Pauly (2016):

“Harmless”

3 Impacts of Introductions

No records of *Melanotaenia trifasciata* introductions were found.

4 Global Distribution



Figure 1. Known global distribution of *Melanotaenia trifasciata*. Map from GBIF Secretariat (2016).

5 Distribution Within the United States

No records of *Melanotaenia trifasciata* in the United States were found.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match for *Melanotaenia trifasciata* was low for most of the west, north, and an area along the Gulf Coast. Southern areas centered on Texas and Florida had a medium match with smaller areas of high match in Florida and Texas. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the contiguous U.S. was 0.062, medium. The following states had individually high climate scores: Arizona, Florida, Oklahoma, and Texas.




Figure 2. RAMP (Sanders et al. 2014) source map of Australia showing weather stations selected as source locations (red) and non-source locations (grey) for *Melanotaenia trifasciata* climate matching. Source locations from GBIF Secretariat (2016)




Figure 3. Map of RAMP (Sanders et al. 2014) climate matches for *Melanotaenia trifasciata* in the contiguous United States based on source locations reported by GBIF Secretariat (2016). 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X < 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

The certainty of assessment is low. There was minimal information available for *Melanotaenia trifasciata*. No records of introductions were found.

8 Risk Assessment

Summary of Risk to the Contiguous United States

The history of invasiveness for *Melanotaenia trifasciata* is uncertain. No records of introduction were found. The climate match is medium; areas of higher climate match include Texas and Florida. The certainty of assessment is low. The overall risk assessment category is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): Uncertain**
- **Climate Match (Sec. 6): Medium**
- **Certainty of Assessment (Sec. 7): Low**
- **Remarks/Important additional information** No additional remarks.
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

Corlis, D. 2004. Taxonomy of monogenean parasites and their coevolution with Australian atheriniform fishes. Doctoral dissertation. James Cook University, Townsville, Australia.

Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2017. Catalog of fishes: genera, species, references. Available:
<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmaintest.asp>. (March 2017).

Froese, R., and D. Pauly, editors. 2016. *Melanotaenia trifasciata* (Rendahl, 1922). FishBase. Available: <http://www.fishbase.org/summary/Melanotaenia-trifasciata.html>. (March 2016).

GBIF Secretariat. 2016. GBIF backbone taxonomy: *Melanotaenia trifasciata* (Rendahl, 1922). Global Biodiversity Information Facility, Copenhagen. Available:
<http://www.gbif.org/species/2411686>. (March 2016).

ITIS (Integrated Taxonomic Information System). 2016. *Melanotaenia trifasciata* (Rendahl, 1922). Integrated Taxonomic Information System, Reston, Virginia. Available:
http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=630798. (March 2016).

Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk assessment mapping program: RAMP. U.S. Fish and Wildlife Service.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Allen, G. R., and N. J. Cross. 1982. Rainbowfishes of Australia and Papua New Guinea. T. F. H. Publications, New Jersey.

Allen, G. R., N. J. Cross, D. J. Bray, and D. F. Hoese. 2006. Various families. Pages 637–646, 689–695, 776–785, 881–895, 911–916, 969–974 in D. F. Hoese, D. J. Bray, J. R. Paxton, and G. R. Allen. 2006. Zoological catalogue of Australia, volume 35. Fishes. Parts 1–3. CSIRO Publishing, Collingwood, Australia.

Allen, G. R., R. K. Hadiaty, P. J. Unmack, and M. V. Erdmann. 2015. Rainbowfishes (*Melanotaenia*: Melanotaeniidae) of the Aru Islands, Indonesia, with description of five new species and redescription of *M. patoti* Weber and *M. senkenbergianus* Weber. *aqua. International Journal of Ichthyology* 21(2):66–108.

Allen, G. R., S. H. Midgley, and M. Allen. 2002. Field guide to the freshwater fishes of Australia. Western Australian Museum, Perth.

Allen, G. R., A. W. Storey, and M. Yarrao. 2008. Freshwater fishes of the Fly River Papua New Guinea. [Source material did not give full citation for this reference.]

Allen, G. R., P. J. Unmack, and R. K. Hadiaty. 2016. The Goldiei group of rainbowfishes (Melanotaeniidae) from the Birds Neck Region of New Guinea (Papua and West Papua Provinces, Indonesia) with descriptions of five new species and recognition of *Melanotaenia dumasi* Weber. *aqua. International Journal of Ichthyology* 22(1):1–31.

Baensch, H. A., and R. Riehl. 1985. Aquarien atlas. Band 2. Mergus, Verlag für Natur-und Heimtierkunde GmbH, Melle, Germany.

Breder, C. M., and D. E. Rosen. 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey.

Larson, H. K., and K. C. Martin. 1990. The freshwater fishes of the Northern Territory. Northern Territory Museum of Arts and Science, Darwin, Handbook Series 1:1–102.

Larson, H. K., R. S. Williams, and M. P. Hammer. 2013. An annotated checklist of the fishes of the Northern Territory, Australia. *Zootaxa* 3696(1):1–293.

Martin, K. C., and S. Barclay. 2015. Distribution and status of banded rainbowfish *Melanotaenia trifasciata* (Melanotaeniidae) populations in north-eastern Queensland. *International Journal of Ichthyology* 21(2):54–65.

- Paxton, J. R., D. F. Hoese, G. R. Allen, and J. E. Hanley. 1989. Zoological catalogue of Australia, volume 7. Pisces. Petromyzontidae to Carangidae. Australian Government Publishing Service, Canberra.
- Pethon, P. 1969. List of type specimens of fishes, amphibians and reptiles in the Zoological Museum, University of Oslo. *Rhizocrinus*, Occasional Papers of the Zoological Museum, University of Oslo 1(1):1–17.
- Rendahl, H. 1922. Meddelelser fra det Zoologiske Museum, Kristiania. Nr. 5. A contribution to the ichthyology of north-west Australia. *Nyt Magazin for Naturvidenskaberne* 60:163–197.
- Unmack, P. J. 2001. Biogeography of Australian freshwater fishes. *Journal of Biogeography* 28:1053–1089.