

Carnatic Carp (*Barbodes carnaticus*)

Ecological Risk Screening Summary

U.S. Fish & Wildlife Service, March 2015

Revised, September 2017, October 2017

Web Version, 8/28/2018



Photo: Dr. N. Basavaraja. Licensed under Creative Commons BY-NC 3.0. Available: <http://www.fishbase.org/photos/UploadedBy.php?autoctr=12615&win=uploaded>. (March 30, 2015).

1 Native Range and Status in the United States

Native Range

From Ali and Raghavan (2013):

“*Barbodes carnaticus* is endemic to the Western Ghats (Dahanukar et al. 2004). Known from rivers in the states of Kerala, Tamil Nadu and Karnataka including Cauvery, Krishna (Jayaram 1999), Moyar (Rajan 1963, Arunachalam et al. 2000), Kabini, Bhavani, Bharathapuzha,

Chalakudy, Periyar, Pambar, Muvattupuzha, Manimala, Pamba, Achenkovil, Karamana, Neyyar (Shaji and Easa 2003, Chhapgar and Mankadan 2000, Kurup et al. 2004), Chaliyar (R. Raghavan and A. Ali pers. obs.). Ooty Lake (Jayaram 1999). Besides it has also been reported from the water bodies inside the Mudumalai Wildlife Sanctuary (Manimekalan 1998), from the drainages in the Dharmapuri district of Tamil Nadu (Rema Devi and Raghunathan 1999) and from Kolli Hills of Eastern Ghats (Arunachalam and Johnson 1998). The record from Tambraparini (Johnsingh and Vickram 1987) is erroneous (Johnson and Arunachalam pers. comm.). The report from southern Kerala is also doubtful (M. Arunachalam pers. comm.).”

Status in the United States

No records of *Barbodes carnaticus* in the wild or in trade in the United States were found.

Means of Introductions in the United States

No records of *Barbodes carnaticus* in the United States were found.

Remarks

No additional remarks.

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

According to Eschmeyer et al. (2017), *Barbodes carnaticus* (Jerdon 1849) is the valid name for this species. It was originally described as *Barbus carnaticus* and has been previously known as *Systomus carnaticus*, *Puntius carnaticus*, and *Hypselobarbus carnaticus*.

From ITIS (2015):

“Kingdom Animalia
Subkingdom Bilateria
Infrakingdom Deuterostomia
Phylum Chordata
Subphylum Vertebrata
Infraphylum Gnathostomata
Superclass Osteichthyes
Class Actinopterygii
Subclass Neopterygii
Infraclass Teleostei
Superorder Ostariophysi
Order Cypriniformes
Superfamily Cyprinoidea
Family Cyprinidae
Genus *Barbodes*
Species *Barbodes carnatiucs* (Jerdon, 1849)”

Size, Weight, and Age Range

From Froese and Pauly (2014):

“Max length: 60.0 cm TL male/unsexed; [Gopalakrishnan and Ponniah 2000]; max. published weight: 12.0 kg [Talwar and Jhingran 1991]; max. reported age: 5 years”

From Ali and Raghavan (2013):

“Longevity of the species which was estimated as 4-5 years (Manojkumar and Kurup 2010) is doubtful (M. Arunachalam, R. Kumar pers. comm.).”

Environment

From Froese and Pauly (2014):

“Freshwater; benthopelagic; potamodromous [Riede 2004].”

Climate/Range

From Froese and Pauly (2014):

“Tropical”

Distribution Outside the United States

Native

From Ali and Raghavan (2013):

“*Barbodes carnaticus* is endemic to the Western Ghats (Dahanukar et al. 2004). Known from rivers in the states of Kerala, Tamil Nadu and Karnataka including Cauvery, Krishna (Jayaram 1999), Moyar (Rajan 1963, Arunachalam et al. 2000), Kabini, Bhavani, Bharathapuzha, Chalakudy, Periyar, Pambar, Muvattupuzha, Manimala, Pamba, Achenkovil, Karamana, Neyyar (Shaji and Easa 2003, Chhapgar and Mankadan 2000, Kurup et al. 2004), Chaliyar (R. Raghavan and A. Ali pers. obs.). Ooty Lake (Jayaram 1999). Besides it has also been reported from the water bodies inside the Mudumalai Wildlife Sanctuary (Manimekalan 1998), from the drainages in the Dharmapuri district of Tamil Nadu (Rema Devi and Raghunathan 1999) and from Kolli Hills of Eastern Ghats (Arunachalam and Johnson 1998). The record from Tambraparini (Johnsingh and Vickram 1987) is erroneous (Johnson and Arunachalam pers. comm.). The report from southern Kerala is also doubtful (M. Arunachalam pers. comm.).”

Introduced

No records of *Barbodes carnaticus* introductions were found.

Means of Introduction Outside the United States

No records of *Barbodes carnaticus* introductions were found.

Short Description

From India Biodiversity Portal (2015):

“A species of *Barbodes* with 28–32 lateral line scales; 10–12 predorsal scales; two pairs of barbels; dorsal fin inserted slightly nearer to tip of snout than base of caudal fin, its last undivided ray osseous, strong and smooth; ; rostral barbel black; four or five fine and faint longitudinal stripes above lateral line. (Mishra et al. 2013).”

Biology

From Froese and Pauly (2014):

“Found in large pools and riffles of rapid rivers and streams. Adults are observed to hide under bedrocks, boulders and undercuttings. Feeds on allochthonous fallen leaves and seeds [Arunachalam et al. 2000]. Does not breed in ponds but spawns in flooded rivers during monsoon months [Talwar and Jhingran 1991]. Spawns in July-August and fry are available in September to December [Arunachalam et al. 2000].”

From Ali and Raghavan (2013):

“The young of *B. carnaticus* are seen in groups along the banks of rivers and reservoirs, while the mature carps are rarely seen along the banks (Biju 2005).”

“*B. carnaticus* is threatened by a wide range of factors including decline in habitat quality due to destructive fishing practices such as poisoning and dynamiting, altered river flow due to construction of dams, competition with exotic and transplanted carps and pollution from point sources.”

Human Uses

From Froese and Pauly (2014):

“Fisheries: commercial; aquaculture: commercial”

From Ali and Raghavan (2013):

“*B. carnaticus* is a much preferred food fish and are caught from wherever they occur. They form minor fisheries in several reservoirs located in the Cauvery drainage.”

Diseases

No records of OIE-reportable diseases were found for *Barbodes carnaticus*.

Poelen et al. (2014) lists *Caballeroia bhavani*, *Falcaustra nilgiriensis*, *F. kaverii*, and *F. brevispiculata* as parasites of *B. carnaticus*.

Threat to Humans

From Froese and Pauly (2014):

“Harmless”

3 Impacts of Introductions

No records of *Barbodes carnaticus* introductions were found, therefore there is no information on impacts of introductions.

4 Global Distribution



Figure 1. Known global distribution of *Barbodes carnaticus*. Locations are in India. Map from GBIF Secretariat (2016).

The northern point is outside any other description of the range of *Barbodes carnaticus* and was not used as a source point for the climate match.

5 Distribution Within the United States

No records of *Barbodes carnaticus* in the United States were found.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match for *Barbodes carnaticus* was low across the entire contiguous United States. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the contiguous United States was 0.000, low, and no states had an individually high climate match.




Figure 2. RAMP (Sanders et al. 2014) source map showing weather stations in India selected as source locations (red) and non-source locations (gray) for *Barbodes carnaticus* climate matching. Source locations from GBIF Secretariat (2016).




Figure 3. Map of RAMP (Sanders et al. 2014) climate matches for *Barbodes carnaticus* in the contiguous United States based on source locations reported by GBIF Secretariat (2016). 0 = Lowest match, 10 = Highest match. Counts of climate match scores are tabulated on the left.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

The certainty of assessment is medium. There was adequate, quality biological and ecological information available for *Barbodes carnaticus*. No records of introductions were found. The georeferenced distribution used for the climate match was very limited based on the text based descriptions of the species' range.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Carnatic Carp (*Barbodes carnaticus*) is a species of carp native to river drainages in India. It is used as a food fish and is a small commercial fishery in some areas. The history of invasiveness is uncertain. There were no records of introductions found. The climate match is low; the climate 6 score was 0.000. The certainty of assessment is medium. The information used in the climate match was incomplete. The overall risk assessment is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): Uncertain**
- **Climate Match (Sec. 6): Low**
- **Certainty of Assessment (Sec. 7): Medium**
- **Remarks/Important additional information** No additional remarks.
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

- Ali, A., and R. Raghavan. 2013. *Barbodes carnaticus*. The IUCN red list of threatened species. Version 2014.3. Available: <http://www.iucnredlist.org/details/full/172418/0>. (March 2015).
- Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2017. Catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (September 2017).
- Froese, R., and D. Pauly, editors. 2014. *Barbodes carnaticus* (Jerdon, 1849). FishBase. Available: <http://www.fishbase.org/summary/Barbodes-carnaticus.html>. (March 2015).
- GBIF Secretariat. 2016. *Barbus carnaticus*. GBIF backbone taxonomy: *Barbodes carnaticus* (Jerdon, 1849). Global Biodiversity Information Facility, Copenhagen. Available: <https://www.gbif.org/species/2365890>. (September 2016).
- India Biodiversity Portal. 2015. *Barbodes carnaticus* (Jerdon, 1849). India Biodiversity Portal, species page. Available: <http://indiabiodiversity.org/species/show/231769>. (March 2015).
- ITIS (Integrated Taxonomic Information System). 2015. *Barbodes carnaticus* (Jerdon, 1849). Integrated Taxonomic Information System, Reston, Virginia. Available: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=688436. (March 2015).

Poelen, J. H., J. D. Simons, and C. J. Mungall. 2014. Global Biotic Interactions: an open infrastructure to share and analyze species-interaction datasets. *Ecological Informatics* 24:148–159.

Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk assessment mapping program: RAMP. U.S. Fish and Wildlife Service.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Arunachalam and Johnson. 1998. [Source material did not give full citation for this reference].

Arunachalam, M., J. A. Johnson, A. Manimekalan, A. Sankaranarayanan, and R. Sornam. 2000. Cultivable and ornamental fishes of Western Ghat Rivers of South India. Pages 205–214 *in* A. Gopalakrishnan, and A. G. Ponniah, editors. Endemic fish diversity of Western Ghats. National Bureau of Fish Genetic Resources, Lucknow, India.

Biju, C. R. 2005. Habitat and distribution of hillstream fishes of Northern Kerala. Department of Zoology, Mahatma Gandhi University.

Chhapgar, B. F., and R. Manakadan. 2000. Ecology of hill streams of Western Ghats with special reference to fish community. U.S. Fish and Wildlife Service and Bombay Natural History Society, Bombay, India.

Dahanukar, N., R. Raut, and A. Bhat. 2004. Distribution, endemism and threat status of freshwater fishes in the Western Ghats of India. *Journal of Biogeography* 31:123–136.

Gopalakrishnan, A., and A. G. Ponniah. 2000. Cultivable, ornamental, sport and food fishes endemic to peninsular India with special reference to Western Ghats. Pages 13–32 *in* A. G. Ponniah, and A. Gopalakrishnan, editors. Endemic fish diversity of Western Ghats. NBFGR-NATP Publication. National Bureau of Fish Genetic Resources, Lucknow, India.

Jayaram, K. C. 1999. The freshwater fishes of the Indian region. Narendra Publishing House, Delhi, India.

Jerdon, T. C. 1849. On the fresh-water fishes of southern India. *Madras Journal of Literature and Science* 15(2):302–346.

Johnsingh, A. J. T., and D. Vickram. 1987. Fishes of Mundanthurai wildlife sanctuary, Tamil Nadu. *Journal of the Bombay Natural History Society* 84:526–533.

- Kurup, B. M., K. V. Radhakrishnan, and T. G. Manojkumar. 2004. Biodiversity status of fishes inhabiting rivers of Kerala (South India) with special reference to endemism, threats and conservation measures. *In* R. L. Welcomme, and T. Petr, editors. Proceedings of the second international symposium on the management of large rivers for fisheries 2. Cambodia.
- Manimekalan, A. 1998. The fishes of Mudumalai Wildlife Sanctuary, Tamil Nadu, South India. *Journal of the Bombay Natural History Society* 95(3):431–443.
- Manojkumar, T. G., and B. M. Kurup. 2010. Age and growth of the Carnatic carp, *Puntius carnaticus* (Jerdon, 1849) from Chalakudy River, Kerala. *Indian Journal of Fisheries* 57(1):81–85.
- Mishra, S. S., L. Kosygin, P. T. Rajan, and K. C. Gopi. 2013. Zoological survey of India. Pages 1–235 *in* K. Venkataraman, A. Chattopadhyay, and K. A. Subramanian, editors. Endemic animals of India (vertebrates). Zoological Survey of India, Kolkata.
- Rajan, S. 1963. Ecology of the fishes of river Pykara and Moyar (Nilgiris), Tamil Nadu. *Proceedings of the Indian Academy of Sciences* 58(5):291–323.
- Rema Devi, K., and M. B. Raghunathan. 1999. The ichthyofauna of Dharmapuri district, Tamil Nadu. *Records of the Zoological Survey of India* 97(1):145–162.
- Riede, K. 2004. Global register of migratory species - from global to regional scales. Final Report, R&D-Projekt 808 05 081. Federal Agency for Nature Conservation, Bonn.
- Shaji, C. P., and P. S. Easa, editors. 2003. Freshwater fishes of Kerala. Kerala Forest Research Institute (KFRI), Thrissur, India.
- Talwar, P. K., and A. G. Jhingran. 1991. Inland fishes of India and adjacent countries, volume 1. A. A. Balkema, Rotterdam, Netherlands.