


***Hypostomus piratatu* (a catfish, no common name)**

Ecological Risk Screening Summary

U.S. Fish and Wildlife Service, January 2013

Revised, October 2018

Web Version, 7/17/2019



No Photo Available

1 Native Range and Status in the United States

Native Range

From Fricke et al. (2018):

“Eastern Paraguay River tributaries and upper Paraná River basin: Argentina and Paraguay.”

Martins et al. (2012) report *H. piratatu* from the Paraguay River basin in Mato Grosso State, Brazil.

Status in the United States

This species has not been reported as introduced or established in the United States. However, unidentified members of the genus are established in the United States.

From Nico et al. (2018):

“Several morphologically distinct but unidentified *Hypostomus* species have been recorded as established in the United States: these included populations in Indian Springs in Nevada; Hillsborough County in Florida; and the San Antonio River and San Felipe Creek in Texas (Courtenay and Deacon 1982; Courtenay et al. 1984, 1986; Courtenay and Stauffer 1990; Page

and Burr 1991; López-Fernández and Winemiller 2005). A population of an unidentified *Hypostomus* species is firmly established in Hawaii (Devick 1991a, b). Reported from Arizona, Colorado, Connecticut, Louisiana, and Pennsylvania. Failed in Connecticut, Massachusetts, and Pennsylvania.”

This species was not found for sale from U.S.-based online aquarium retailers, and it does not appear to be in trade in the United States.

Means of Introduction into the United States

This species has not been reported as introduced or established in the United States. However, unidentified members of the genus are established in the United States.

From Nico et al. (2018):

“Members of this genus have been introduced through a combination of fish farm escapes or releases, and aquarium releases (Courtenay and Stauffer 1990; Courtenay and Williams 1992). In Texas, the initial introduction occurred when *Hypostomus* entered local streams after escaping from pool and canal systems of the San Antonio Zoological Gardens in or before 1962 (Barron 1964); the Comal County introduction was probably due to an aquarium release (Whiteside and Berkhouse 1992).”

Remarks

From Nico et al. (2018):

“The genus *Hypostomus* contains about 116 species (Burgess 1989). Highlighting the serious need for additional taxonomic and systematic work, Armbruster (1997) concluded that it is currently impossible to identify most species in the genus. Several apparently different *Hypostomus* species have been collected in the United States but not definitively identified to species level (Page and Burr 1991; Courtenay and Stauffer 1990). Distinguishing characteristics of the genus and a key to loricariid genera were provided by Burgess (1989) and Armbruster (1997). Photographs appeared in Burgess (1989) and Ferraris (1991). *Hypostomus* has officially replaced the generic name *Plecostomus*. The genus was included in the key to Texas fishes of Hubbs et al. (1991) and several identifying traits were also given by Page and Burr (1991).”

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2018):

“Kingdom Animalia
Subkingdom Bilateria
Infrakingdom Deuterostomia
Phylum Chordata
Subphylum Vertebrata
Infraphylum Gnathostomata

Superclass Actinopterygii
Class Teleostei
Superorder Ostariophysi
Order Siluriformes
Family Loricariidae
Subfamily Hypostominae
Genus *Hypostomus*
Species *Hypostomus piratatu* Weber, 1986”

“Current Standing: valid”

Size, Weight, and Age Range

From Froese and Pauly (2018):

“Max length: 27.5 cm SL male/unsexed; [Weber 2003]”

Environment

From Froese and Pauly (2018):

“Freshwater; demersal.”

Climate/Range

From Froese and Pauly (2018):

“Tropical”

Distribution Outside the United States

Native

From Fricke et al. (2018):

“Eastern Paraguay River tributaries and upper Paraná River basin: Argentina and Paraguay.”

Martins et al. (2012) report *H. piratatu* from the Paraguay River basin in Mato Grosso State, Brazil.

Introduced

No introductions of this species have been reported.

Means of Introduction Outside the United States

No introductions of this species have been reported.

Short Description

No information available.

Biology

No information available.

Human Uses

No information available.

Diseases

From Pérez-Ponce de León et al. (2008):

“*Crassicutis intermedius* [helminth parasite of fishes] occurs in [...] *Hypostomus piratatu* Weber [...]”

No OIE-reportable diseases have been documented in this species.

Threat to Humans

From Froese and Pauly (2018):

“Harmless”

3 Impacts of Introductions

No introductions of *H. piratatu* have been reported as introduced outside its native range so no impacts of introduction are known. However, unidentified members of the genus are established in the United States.

From Nico et al. (2018):

“The effects of these loricariid catfish is largely unknown. In Texas, Hubbs et al. (1978) reported possible local displacement of algae-feeding native fishes such as *Camptostoma anomalum* by *Hypostomus*, and López-Fernández and Winemiller (2005) suggest that reductions in *Dionda diaboli* abundance in portions of San Felipe Creek are due to population increases of *Hypostomus*. Because of their abundance in Hawaii, introduced *Hypostomus*, *Pterygoplichthys*, and *Ancistrus* may compete for food and space with native stream species (Devick 1989; Sabaj and Englund 1999).”

4 Global Distribution



Figure 1. Known global distribution of *Hypostomus piratatu*, reported from Brazil and Paraguay. Map from GBIF Secretariat (2017). No georeferenced occurrences were available for the southern part of the species range in Argentina and Paraguay.

5 Distribution within the United States

There is currently no known distribution of *Hypostomus piratatu* within the United States; however, unidentified species of *Hypostomus* are established in Nevada, Florida, Texas, and Hawaii.

6 Climate Matching

Summary of Climate Matching Analysis

The climate match (Sanders et al. 2018; 16 climate variables; Euclidean Distance) was highest in southern Florida, with matches as high as 8 on a scale from 0 (low) to 10 (high; Fig. 3). Medium matches extended through northern Florida and much of Georgia and along the Texas coastline. However, the majority of the contiguous United States showed a low climate match. Climate 6 score indicated that the contiguous United States has a medium climate match overall. The range of scores indicating medium match is between 0.005 and 0.103; Climate 6 score of *H. piratatu* is 0.013. A higher score would be possible if georeferenced occurrences were available from the southern portion of the species range.




Figure 2. RAMP (Sanders et al. 2018) source map showing weather stations selected as source locations (red; Brazil) and non-source locations (gray) for *H. piratatu* climate matching. Source locations from GBIF Secretariat (2017).




Figure 3. Map of RAMP (Sanders et al. 2018) climate matches for *H. piratatu* in the contiguous United States based on source locations reported by GBIF Secretariat (2017). 0= Lowest match, 10= Highest match.

The “High”, “Medium”, and “Low” climate match categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

Limited information was available on the biology and ecology of *Hypostomus piratatu*. It has not been reported as introduced outside its native range, so no impacts of introduction are known. However, unidentified species of *Hypostomus* have become established in the United States, and it is possible that one or more of those populations could be identified later as *H. piratatu*. There is considerable uncertainty about the taxonomy of this genus and about species-level identification. Certainty of this assessment is low.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Hypostomus piratatu is a catfish native to Eastern Paraguay River tributaries and the upper Paraná River basin in Brazil, Paraguay, and Argentina. This species has no documented history of introduction in the United States or elsewhere outside its native range, and it is not known to be in trade. However, unidentified species of *Hypostomus* are established in the United States. Climate match to the contiguous United States was medium overall, with high match in southern Florida. It is possible that the climate match could increase if more georeferenced occurrences from the southern portion of the species range became available. Because of the lack of documented introduction history and substantial taxonomic uncertainty, certainty of this assessment is low and overall risk is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): Uncertain**
- **Climate Match (Sec. 6): Medium**
- **Certainty of Assessment (Sec. 7): Low**
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

- Fricke, R., W. N. Eschmeyer, and R. van der Laan, editors. 2018. Catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (October 2018).
- Froese, R., and D. Pauly, editors. 2018. *Hypostomus piratatu* Weber, 1986. FishBase. Available: <https://www.fishbase.de/summary/Hypostomus-piratatu.html>. (October 2018).
- GBIF Secretariat. 2017. GBIF backbone taxonomy: *Hypostomus piratatu* Weber, 1986. Global Biodiversity Information Facility, Copenhagen. Available: <https://www.gbif.org/species/5202175>. (October 2018).
- ITIS (Integrated Taxonomic Information System). 2018. *Hypostomus piratatu* Weber, 1986. Integrated Taxonomic Information System, Reston, Virginia. Available: https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=680221#null. (October 2018).
- Martins, F. O., M. M. F. Marinho, F. Langeani, and J. P. Serra. 2012. A new species of *Hypostomus* (Siluriformes: Loricariidae) from the Upper Rio Paraguay basin, Brazil. *Copeia* 2012(3):494-500.

Nico, L., P. Fuller, and M. Neilson. 2018. *Hypostomus* sp. Lacepède, 1803. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=762>. (October 2018).

Pérez-Ponce de León, G., U. Razo-Mendivil, R. Rosas-Valdez, B. Mendoza-Garfias, and H. Mejía-Madrid. 2008. Description of a new species of *Crassicutis* Manter, 1936, parasite of *Cichlasoma beani* Jordan (Osteichthyes: Cichlidae) in Mexico, based on morphology and sequences of the ITS1 and 28S ribosomal RNA genes. *Journal of Parasitology* 94(1):257-263.

Sanders, S., C. Castiglione, and M. Hoff. 2018. Risk Assessment Mapping Program: RAMP, version 3.1. U.S. Fish and Wildlife Service.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Armbruster, J. W. 1997. Phylogenetic relationships of the sucker-mouth armored catfishes (Loricariidae) with particular emphasis on the Ancistrinae, Hypostominae, and Neoplecostominae. Doctoral dissertation. University of Illinois, Champaign-Urbana, Illinois.

Barron, J. L. 1964. Reproduction and apparent over-winter survival of the sucker-mouth armoured catfish, *Plecostomus* sp., in the headwaters of the San Antonio River. *The Texas Journal of Science* 16:449.

Burgess, W. E. 1989. An atlas of freshwater and marine catfishes: a preliminary survey of the Siluriformes. Tropical Fish Hobbyist Publications, Inc., Neptune City, New Jersey.

Courtenay, W. R., Jr., and J. E. Deacon. 1982. Status of introduced fishes in certain spring systems in southern Nevada. *Great Basin Naturalist* 42(3):361-366.

Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1984. Distribution of exotic fishes in the continental United States. Pages 41-77 in W. R. Courtenay, Jr., and J. R. Stauffer, Jr, editors. *Distribution, biology, and management of exotic fishes*. John Hopkins University Press, Baltimore, Maryland.

Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1986. Distribution of exotic fishes in North America. Pages 675-698 in C. H. Hocutt, and E. O. Wiley, editors. *The zoogeography of North American freshwater fishes*. John Wiley and Sons, New York.

- Courtenay, W. R., Jr., and J. R. Stauffer. 1990. The introduced fish problem and the aquarium fish industry. *Journal of the World Aquaculture Society* 21(3):145-159.
- Courtenay, W. R., Jr., and J. D. Williams. 1992. Dispersal of exotic species from aquaculture sources, with emphasis on freshwater fishes. Pages 49-81 *in* A. Rosenfield, and R. Mann, editors. *Dispersal of living organisms into aquatic ecosystems*. Maryland Sea Grant, College Park, Maryland.
- Devick, W. S. 1989. Disturbances and fluctuations in the Wahiawa Reservoir ecosystem. Project no. F-14-R-13, Job 4, Study I. Division of Aquatic Resources, Hawaii Department of Land and Natural Resources.
- Devick, W. S. 1991a. Disturbances and fluctuations in the Wahiawa Reservoir ecosystem. Project no. F-14-R-15, Job 4 Study I. Division of Aquatic Resources, Hawaii Department of Land and Natural Resources.
- Devick, W.S. 1991b. Patterns of introductions of aquatic organisms to Hawaiian freshwater habitats. Pages 189-213 *in* *New directions in research, management and conservation of Hawaiian freshwater stream ecosystems*. Proceedings of the 1990 symposium on freshwater stream biology and fisheries management. Division of Aquatic Resources, Hawaii Department of Land and Natural Resources.
- Ferraris, C. J., Jr. 1991. *Catfish in the aquarium*. Tetra Press, Morris Plains, New Jersey.
- Hubbs, C., R. J. Edwards, and G. P. Garrett. 1991. An annotated checklist of freshwater fishes of Texas, with key to identification of species. *Texas Journal of Science*, Supplement 43(4):1-56.
- Hubbs, C., T. Luciere, G. P. Garrett, R. J. Edwards, S. M. Dean, and E. Marsh. 1978. Survival and abundance of introduced fishes near San Antonio, Texas. *The Texas Journal of Science* 30(4):369-376.
- López-Fernández, H., and K. O. Winemiller. 2005. Status of *Dionda diaboli* and report of established populations of exotic fish species in lower San Felipe Creek, Val Verde County, Texas. *Southwestern Naturalist* 50(2):246-251.
- Page, L. M., and B. M. Burr. 1991. *A field guide to freshwater fishes of North America north of Mexico*. The Peterson Field Guide Series, volume 42. Houghton Mifflin Company, Boston.
- Sabaj, M. H., and R. A. Englund. 1999. Preliminary identification and current distribution of two suckermouth armored catfishes (Loricariidae) introduced to Oahu streams. *Bishop Museum Occasional Papers* 59:50-55.

Weber, C. 2003. Loricariidae - Hypostominae (armored catfishes). Pages 351-372 *in* R. E. Reis, S. O. Kullander, and C. J. Ferraris, Jr., editors. Checklist of the freshwater fishes of South and Central America. EDIPUCRS, Porto Alegre, Brazil.

Whiteside, B. G., and C. Berkhouse. 1992. Some new collections locations for six fish species. The Texas Journal of Science 44(4):494.