

# Tanganyika Blackfin (*Altolamprologus calvus*)

## Ecological Risk Screening Summary

U.S. Fish & Wildlife Service, April 2013  
Revised, September 2017, October 2017  
Web Version, 8/21/2018



Photo: D Ross Robertson. Licensed under Creative Commons BY-NC 3.0. Available: [http://eol.org/data\\_objects/13236885](http://eol.org/data_objects/13236885). (April 11, 2013).

## 1 Native Range and Status in the United States

---

### Native Range

From Froese and Pauly (2017):

“Africa: Endemic to Lake Tanganyika, found in the southwestern [Democratic Republic of the Congo, Tanzania, Zambia] part of the lake [Konings and Dieckhoff 1992; Konings 1998].”

From Bigirimana (2006):

“Endemic to the southern part of Lake Tanganyika [Democratic Republic of the Congo, Tanzania, Zambia].”

### **Status in the United States**

No records of *Altolamprologus calvus* in the United States were found. *A. calvus* is in trade in the United States.

From Cichlids and Herps (2018):

“*Altolamprologus calvus* “Inkfin” – juvenile \$18.00 [...] *Altolamprologus calvus* Black Congo juvenile 1.5”-2” \$16.00”

### **Means of Introductions in the United States**

No records of *Altolamprologus calvus* in the United States were found.

### **Remarks**

No additional remarks.

## **2 Biology and Ecology**

---

### **Taxonomic Hierarchy and Taxonomic Standing**

According to Eschmeyer et al. (2017), *Altolamprologus calvus* (Poll 1978) is the valid name for this species. It was originally described as *Lamprologus calvus*.

From ITIS (2013):

“Kingdom Animalia  
Subkingdom Bilateria  
Infrakingdom Deuterostomia  
Phylum Chordata  
Subphylum Vertebrata  
Infraphylum Gnathostomata  
Superclass Osteichthyes  
Class Actinopterygii  
Subclass Neopterygii  
Infraclass Teleostei  
Superorder Acanthopterygii  
Order Perciformes  
Suborder Labroidei  
Family Cichlidae  
Genus *Altolamprologus* Poll, 1978  
Species *Altolamprologus calvus* (Poll, 1978)”

## **Size, Weight, and Age Range**

From Froese and Pauly (2017):

“Max length: 13.5 cm TL male/unsexed; [Maréchal and Poll 1991]”

From WildScreen (2013):

“Male length: c.15 cm [Smith 1998]  
Female length: c.10 cm [Smith 1998]”

## **Environment**

From Froese and Pauly (2017):

“Freshwater; benthopelagic; pH range: 8.0 - 9.0; dH range: 9 - 19. [...]; 23°C - 25°C [assumed to be recommended aquarium temperature] [Baensch and Riehl 1985]; [...]”

## **Climate/Range**

From Froese and Pauly (2017):

“Tropical; [...]; 3°S - 9°S”

## **Distribution Outside the United States**

### **Native**

From Froese and Pauly (2017):

“Africa: Endemic to Lake Tanganyika, found in the southwestern part of the lake [Democratic Republic of the Congo, Tanzania, Zambia] [Konings and Dieckhoff 1992; Konings 1998].”

From Bigirimana (2006):

“Endemic to the southern part of Lake Tanganyika [Democratic Republic of the Congo, Tanzania, Zambia].”

### **Introduced**

No records of *Altolamprologus calvus* introductions could be found.

## **Means of Introduction Outside the United States**

No records of *Altolamprologus calvus* introductions could be found.

## Short Description

From WildScreen (2013):

“*Altolamprologus calvus* belongs to the Cichlidae, a family of freshwater fish that have adapted to a wide range of narrow ecological niches, resulting in the evolution of a huge diversity of species that live in close association [Clabaut et al. 2007]. This species has a large mouth and a compressed body, with a large dorsal fin running along the entire length of the back [Hanke and Wilson 2006]. Like many cichlids, *Altolamprologus calvus* occurs in a number of different colour variations, including black, light grey and yellow [Smith 1998, Clabaut et al. 2007].”

## Biology

From WildScreen (2013):

“In the case of *Altolamprologus calvus*, it has become adapted to living amongst rocky environments, where it feeds on tiny crustaceans found on rock surfaces (Sturmbauer et al. 1994) and the young of other fishes (P. V. Loiselle, personal communication). Its compressed body allows it to fit into narrow crevices and caves; an ability that has led to the development of an innovative breeding strategy. The female locates a crevice or cave that is too small for the male to enter, and spawns up to 300 eggs. The male then fertilises the eggs by lying over the crevice entrance and releasing sperm into the water. Both sexes guard the developing embryos; the male patrols the outside of the crevice, occasionally leaving to feed, while the female remains in the crevice at all times until the eggs hatch (Smith 1998). The young continue to be protected as long as they remain within their parents’ territory (P. V. Loiselle, personal communication).”

“*Altolamprologus calvus* occupies rocky regions along the shoreline between depths of 3 and 41 metres [Smith 1998].”

From Froese and Pauly (2017):

“Deep and laterally very compressed body with [sic] permits it to enter narrow cracks and shallow caves; feeds mainly on shrimps and other crustaceans; stalker which cruises through rocky habitat maintaining a distance of between 30 and 100 cm between themselves and substrate [Konings 1998].”

## Human Uses

From Froese and Pauly (2017):

“Fisheries: ; aquarium: commercial”

## Diseases

**No records of OIE reportable diseases were found.**

From Froese and Pauly (2017):

“Fish tuberculosis (FishMB), Bacterial diseases”

## Threat to Humans

From Froese and Pauly (2017):

“Harmless”


## 3 Impacts of Introductions

---

No records of *Altolamprologus calvus* introductions could be found.

## 4 Global Distribution

---



**Figure 1.** Known global distribution of *Altolamprologus calvus*. Locations are in the southern end of Lake Tanganyika in Democratic Republic of the Congo and Zambia. Map from GBIF Secretariat (2017).

## 5 Distribution Within the United States


---

No records of *Altolamprologus calvus* in the United States were found.


## 6 Climate Matching

### Summary of Climate Matching Analysis

The climate match for *Altolamprologus calvus* was low for most of the United States. Southwest Florida and the southern tip of Texas had a medium match; it was low everywhere else. The Climate 6 score (Sanders et al. 2014; 16 climate variables; Euclidean distance) for the contiguous United States was 0.000, low, and no States had an individually medium or high climate match.



**Figure 2.** RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red; Democratic Republic of the Congo, Tanzania, Zambia) and non-source locations (grey) for *Altolamprologus calvus* climate matching. Source locations from GBIF Secretariat (2017).



**Figure 3.** Map of RAMP (Sanders et al. 2014) climate matches for *Altolamprologus calvus* in the contiguous United States based on source locations reported by GBIF Secretariat (2017). 0 = Lowest match, 10 = Highest match. Counts of climate match scores are tabulated on the left.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
$\geq 0.103$	High

## 7 Certainty of Assessment

The certainty of this assessment is low. General species information is easily available for *Altolamprologus calvus*. There was no information in regard to introductions outside of its native range or any impacts that it would have.

## 8 Risk Assessment

---

### Summary of Risk to the Contiguous United States

Tanganyika Blackfin (*Altalamprologus calvus*) is a cichlid species native to the southern end of Lake Tanganyika. It is used in the aquarium industry. The history of invasiveness is uncertain. No records of introduction were found. *A. calvus* is in trade in the United States. No information on the volume or duration of this species in trade was found. The climate match is 0.000, low. *A. calvus* is endemic to Lake Tanganyika and has a very narrow climate range. The certainty of assessment is low. The overall risk assessment category is uncertain.

### Assessment Elements

- **History of Invasiveness (Sec. 3): Uncertain**
- **Climate Match (Sec. 6): Low**
- **Certainty of Assessment (Sec. 7): Low**
- **Remarks/Important additional information** No additional remarks.
- **Overall Risk Assessment Category: Uncertain**

## 9 References

---

**Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.**

Bigirimana, C. 2006. *Altalamprologus calvus*. The IUCN Red List of Threatened Species 2006: e.T60455A12368559. Available: <http://www.iucnredlist.org/details/full/60455/0>. (April 2013).

Cichlid and Herps Store. 2018. Shop – *Altalamprologus* – Lake Tanganyika. Available: <https://cichlidsandherps.net/t/lake-tanganyika---altalamprologus>. (August 2018).

Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2017. Catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (September 2017).

Froese, R., and D. Pauly, editors. 2017. *Altalamprologus calvus* (Poll, 1978). FishBase. Available: <http://www.fishbase.org/summary/Altalamprologus-calvus.html>. (September 2017).

GBIF Secretariat. 2017. GBIF Backbone Taxonomy: *Altalamprologus calvus* (Poll, 1978). Global Biodiversity Information Facility, Copenhagen. Available: <https://www.gbif.org/species/2372761>. (September 2017).



ITIS (Integrated Taxonomic Information System). 2013. *Altolamprologus calvus* (Poll, 1978). Integrated Taxonomic Information System, Reston, Virginia. Available: [http://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=648250](http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=648250). (April 2013).

Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk assessment mapping program: RAMP. U.S. Fish and Wildlife Service.

WildScreen. 2003. Arkive. Available: <http://www.arkive.org/tanganyika-blackfin/altolamprologus-calvus/>. (April 2013).

## 10 References Quoted But Not Accessed

---

**Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.**

Baensch, H. A., and R. Riehl. 1985. Aquarien atlas. Band 2. Mergus, Verlag für Natur-und Heimtierkunde GmbH, Melle, Germany.

Clabaut, C., P. M. E. Bunje, W. Salzburger, and A. Meyer. 2007. Geometric morphometric analyses provide evidence for the adaptive character of the Tanganyikan cichlid fish radiations. *Evolution* 61:560–578.

Hanke, G. F., and M. V. H. Wilson. 2006. Anatomy of the early Devonian acanthodian *Brochoadmones milesi* based on nearly complete body fossils, with comments on the evolution and development of paired fins. *Journal of Vertebrate Paleontology* 26:526–537.

Konings, A. 1998. Tanganyika cichlids in their natural habitat. Cichlid Press.

Konings, A., and H. W. Dieckhoff. 1992. Tanganyika secrets. Cichlid Press.

Maréchal, C., and M. Poll. 1991. *Altolamprologus*. Pages 4–5 in J. Daget, J.-P. Gosse, G. G. Teugels, and D. F. E. Thys van den Audenaerde, editors. Checklist of freshwater fishes of Africa (CLOFFA). ISNB, Brussels; MRAC, Tervuren, Belgium; and ORSTOM, Paris.

Smith, M. 1998. Lake Tanganyikan cichlids: everything about purchase, care, nutrition, behavior, and aquarium maintenance. Barron's Educational Series, New York.

Sturmbauer, C., E. Verheyen, and A. Meyer. 1994. Mitochondrial phylogeny of the Lamprologini, the major substrate spawning lineage of cichlid fishes from Lake Tanganyika in eastern Africa. *Molecular Biology and Evolution* 11:691–703.