

**FINAL DRAFT**

Avian Conservation Implementation Plan  
Timucuan Ecological and Historic Preserve and  
Fort Caroline National Memorial

National Park Service  
Southeast Region



Evan McClung, The Nature Conservancy

Compiled by J. Keith Watson  
U.S. Fish and Wildlife Service  
In cooperation with

TIMU Resource Management Staff, National Park Service  
And Bird Conservation Partners  
January 2005

## TABLE OF CONTENTS

Introduction.....	3
Background .....	3
The North American Bird Conservation Initiative .....	4
The Southeastern Bird Conservation Initiative: National Park Service .....	5
Role of NPS in Avian Conservation.....	5
Park Description .....	7
Avian Resources of the South Atlantic Coastal Plain.....	8
Avian Conservation in TIMU.....	13
Park Identified Needs for Avian Conservation.....	14
Coordination with Regional Conservation Initiatives .....	15
North American Bird Conservation Initiative .....	15
North American Waterfowl Management Plan.....	15
Partners In Flight .....	17
United States Shorebird Conservation Plan .....	17
Waterbird Conservation for the Americas.....	17
Integration of NABCI Goals and Objectives into Park Planning and Operations: NABCI Implementation	
Recommendations .....	17
Inventory .....	18
Monitoring .....	19
Habitat Restoration .....	20
Threat Management.....	22
Research .....	23
Compliance .....	24
Outreach .....	25
Partners and Partnerships .....	25
Funding Opportunities .....	26
Contacts.....	28
Literature Cited.....	30
Appendixes	
High Priority Species in the South Atlantic Coastal Plain	
South Atlantic Coastal Plain Bird and Habitat Assemblages	
Florida’s Endangered Species, Threatened Species, and Species of Special Concern	
USFWS Species of Conservation Concern (2002) in the Southeastern Coastal Plain (BCR 27)	

## Introduction

This Avian Conservation Implementation Plan (ACIP) is provided to the staff at Timucuan Ecological and Historic Preserve (TIMU) and Fort Caroline National Memorial (FOCA) to help identify and prioritize bird conservation opportunities, and to provide information and guidance for the successful implementation of needed conservation activities. This plan may identify goals, strategies, partnerships, and perhaps specific projects allowing the park to participate in existing bird conservation planning and implementation efforts associated with the North American Bird Conservation Initiative (NABCI). Under the auspice of NABCI, appropriate bird and habitat conservation goals may be recommended as identified in the appropriate existing national or regional bird conservation efforts aligned with this initiative: Partners In Flight (PIF), North American Waterfowl Management Plan (NAWMP), US Shorebird Conservation Plan (USSCP), and Waterbird Conservation for the Americas (WCA). For example, parks in the Appalachians and the Cumberland Plateau, will have few if any high priority waterbird conservation issues at a regional landscape or greater scale. As such, little information regarding waterbird conservation will be presented in the ACIP, unless there is an identified park need for this species group, or other mandates, such as federal laws. Similarly, because TIMU (hereafter also includes FOCA) is primarily a coastal estuary/marsh system, most recommendations for bird conservation will be based on existing colonial waterbird, shorebird, and waterfowl plans. However, because TIMU has upland habitats as well, the PIF plan will be used to make recommendations as well.

Information and data presented in the ACIP have been obtained from several sources: 1) interviews with TIMU staff 2) TIMU bird conservation partners 3) the PIF South Atlantic Coastal Plain Bird Conservation Plan, Version 1.0 (Hunter et al. 2001), 4) NPS databases, 5) peer reviewed bird conservation and management literature, and 6) personal communications with bird conservation specialists throughout North America, especially in the southeastern United States. This plan has been reviewed by TIMU resource management staff and managers, Southeast Coast Inventory and Monitoring Network (SEC I&M) staff, and bird conservation partners and approved by TIMU management. Optimally, this plan will be incorporated into the park's Resource Management Plan (RMP) and updated annually to reflect completed projects, newly identified needs, and shifts in bird conservation priorities in the region.

**TIMU is not obligated to undertake any of the proposed actions in this plan. The plan is provided to offer guidance to TIMU to voluntarily support important park, regional, and perhaps national and international bird conservation projects for which TIMU is a primary participant in the proposed actions.**

## Background

During the past thirty years, monitoring programs across North America have documented declines of certain bird species populations and their habitats, often severe (Sauer et al. 2000). The decline has caused great concern among scientists, biologists,

biodiversity proponents, ecologists, land managers, etc., and the bird conservation community in general. Birds are recognized as critical components of local and global genetic, species, and population diversity, providing important and often critical ecological, social, economic, and cultural values. Their overall decline has stimulated a worldwide focus on conservation efforts, and North American interest in bird conservation is rapidly becoming a focus of government, non-government, industry, and private interests and expenditures.

Many state, federal, and non-governmental wildlife agencies and non-governmental organizations (NGO's) have recognized this alarming bird decline trend and have joined forces in several extensive partnerships to address the conservation needs of various bird groups and their habitats. The primary initiatives are:

- North American Waterfowl Management Plan
- Partners in Flight
- U.S. Shorebird Conservation Plan
- Waterbird Conservation for the Americas

**The North American Bird Conservation Initiative:** While efforts associated with these plans have generated some successes, it has been increasingly recognized that the overlapping conservation interests of these initiatives can be better served through more integrated planning and delivery of bird conservation. The *North American Bird Conservation Initiative (NABCI; <http://www.nabci-us.org/main2.html>)* arose out of this realization. The vision of NABCI is simply to see ***“populations and habitats of North America’s birds protected, restored and enhanced through coordinated efforts at international, national, regional, state and local levels, guided by sound science and effective management.”*** NABCI seeks to accomplish this vision through (1) broadening bird conservation partnerships, (2) working to increase the financial resources available for bird conservation in the U.S., and (3) enhancing the effectiveness of those resources and partnerships by facilitating integrated bird conservation (U.S. NABCI Committee 2000). The four bird conservation initiatives mentioned above, as well as several other local and regional partnerships, work collectively to pursue this vision.

NABCI is guided by a set of principles that establish an operational framework within which the Initiative and its partners may conduct integrated bird conservation in the U.S. These will articulate a common understanding of the relationship among NABCI, the individual bird conservation initiatives, and all partner entities to ensure recognition of existing federal legislative and international treaty obligations, state authorities, and respect for the identity and autonomy of each initiative. The fundamental components of the conservation approach to be used by NABCI are expressed within its goal:

***To deliver the full spectrum of bird conservation through regionally-based, biologically-driven, landscape-oriented partnerships.***

**The Southeastern Bird Conservation Initiative: National Park Service:** In 1999, the Southeast Region of the National Park Service (NPS) recognized the importance of coordinating existing bird conservation goals into planning and operations of national park units in the southeast, that is, integration of NABCI. In support of this recognition, the Southeast Regional Office NPS approved and allocated eighty-eight thousand dollars, cost sharing 1:1 with the US Fish and Wildlife Service (FWS) Region 4 (Southeast) to hire a biologist to conduct this two-year project (Interagency Agreement FS028 01 0368). This project is unique in the NPS, and perhaps the nation, and represents a potential model for better coordinating regional bird conservation programs and activities within and outside the NPS. It further represents a progressive action toward institutionalizing bird conservation as a programmatic priority in the Southeast Region of NPS and potentially the nation.

As envisioned, the integration of NABCI into the Southeastern NPS involves:

- 1) Development and delivery of Avian Conservation Implementation Plans,
- 2) Coordination with NPS Inventory and Monitoring Program,
- 3) Development of a web-based project site,
- 4) Establishment or enhancement of bird conservation partnerships,
- 5) Identification and exploration of potential funding opportunities, and
- 6) Technical guidance and assistance as needed or requested.

This ACIP fulfills one aspect of the plan outlined above and serves as a basis for future bird conservation actions in TIMU and with adjacent partners or landowners.

Concurrently, the development of a Memorandum of Understanding (MOU) between the FWS and the NPS to implement Presidential Executive Order (EO) 13186 (US Government 2000), Responsibilities of Federal Agencies to Protect Migratory Birds, calls for integration of programs and recommendations of existing bird conservation efforts into park planning and operations. Complementing each other, the MOU and the Southeastern Bird Conservation Initiative will advance bird conservation in the Southeast Region of the NPS beyond current regional NPS efforts.

## **Role of NPS in Avian Conservation**

The interagency agreement that facilitates this partnership supports both FWS and NPS management policies. Specifically for the NPS, the agreement supports and advances the Strategy for Collaboration, a visionary document developed and signed by the Southeast Natural Resource Leaders Advisory Group (SENRLAG 2000), a consortium of 13 land and resource management agencies in the Southeastern United States whose vision is to encourage and support cooperation in planning and managing the region's natural resources. Furthermore, the agreement is aligned with and implements a variety of NPS Management Policies (2001) including, but not limited to, External Threats and Opportunities, Environmental Leadership, Cooperative Planning, Land Protection, and especially Natural Resource Management that details policy and management guidelines which apply to bird conservation. Important policies in the

Natural Resource Management chapter include:

- Planning for Natural Resource Management
- Partnerships
- Restoration of Natural Systems
- Studies and Collection
- General Principles for Managing Biological Resources
- Plant and Animal Population Management Principles
- Management of Native Plants and Animals
- Management of Endangered Plants and Animals
- Management of Natural Landscapes
- Management of Exotic Species
- Pest Management
- Fire Management and
- Water Resource Management

The NPS is the fourth largest landowner in the United States, consisting of over 380 national park units covering 83 million acres of land and water with associated biotic resources ([www.nps.gov](http://www.nps.gov)). The 64 units in the Southeast Region of the NPS represent 16% of the total number of park units in the national park system and cover approximately 5% of the total land base in the entire system. Park units in the Southeast Region include national seashores (Canaveral National Seashore, Cumberland Island National Seashore), national parks (Great Smoky Mountains National Park, Everglades National Park), national recreation areas (Big South Fork National River and Recreation Area), national preserves (Big Cypress National Preserve), national battlefields (Cowpens National Battlefield, Fort Donelson National Battlefield), national monuments (Ocmulgee National Monument), and others such as the Blue Ridge Parkway, Obed Wild and Scenic River, and Timucuan Ecological and Historic Preserve.

Southeast NPS units provide habitat for over 400 species of migrating, breeding, and wintering birds and include a wide range of Federal and State listed threatened and endangered species. Likewise, these units also provide nest, migration, and winter habitat for most of the eastern species identified in the national bird conservation plans in need of conservation attention.

Additionally, the NPS attracts over 280 million visitors to the parks each year, 120 million of these in the Southeast Region, affording excellent recreational bird watching and opportunities to strengthen bird conservation interpretation, outreach, and education programs. These opportunities, the NPS mission, policies, and organization all lead to the conclusion that the NPS is an extremely valuable partner and contributor to bird conservation in the region.

Nationally, the status of birds in national parks is largely unknown, although many parks have adequate knowledge regarding bird occurrence in the parks (<http://www.npwrc.usgs.gov/resource/othrdata/chekbird/chekbird.htm>). Parks often

play a role in ongoing regional bird conservation efforts. Indeed many of these parks are often important to regional, national, or international bird conservation, and many have been designated as Important Bird Areas (IBA's) by the National Audubon Society. To date, there are approximately 64 NPS units that are designated IBA's, 35 of which are considered of global importance (<http://abcbirds.org/iba/aboutiba.htm>). In the Southeast Region, the NPS has 13 global IBA's.

The **NPS Inventory and Monitoring (I&M) Program** has been developed to provide management driven scientific information to national park managers so that resources can be adequately protected within national parks. One of the first phases of this program is to inventory vertebrates, including birds, within the 260 national park units in the program. Once completed, data from the inventories will provide an account of the occurrence and abundance of birds in all the national parks in the program. These records will be stored in the NPS I&M NPSpecies database (<http://www.nature.nps.gov/im/apps/npspp/>). Coordination with I&M network staff is important to developing long-term bird monitoring programs that fulfill both park and NABCI objectives.

**Park Flight** is a NPS international partnership initiative that directs funding toward a variety of NPS programs that involve conservation of Neotropical migratory birds whose life history range covers a US national park and a Latin American protected area. A relatively new program, Park Flight offers parks the opportunity to partner with a Latin American national park or protected area to cooperate on developing bird conservation and education projects (USDI NPS 2002).

Recent increases in NPS base funded programs such as inventory and monitoring, exotic species management, habitat restoration, and fire management all indicate that national park managers recognize that park lands are increasingly subject to a variety of threats and conditions that must be improved to provide the quality of national park experience articulated in the NPS Organic Act (1916). Programmatic funding in these areas will increase the ability of national parks to provide quality habitat and conditions for increased wildlife conservation, including birds. Furthermore, private interests and non-profit conservation organizations have initiated programs, including grant programs, to provide much needed funding to national parks to meet backlogs of identified yet unfunded needs.

## **Park Description**

Situated entirely within Duval County and the city limits of Jacksonville, FL, Timucuan Ecological and Historical Preserve covers approximately 18,600 ha (46,000 acres) between the St. Johns and Nassau rivers. The southern third of the Preserve lies at the mouth of the extensive St. Johns River watershed, which encompasses parts of Duval and several other counties for approximately 300 miles to the south. The St. Johns River is heavily impacted by agricultural, industrial and urban pollution; however, marine tidal waters near its mouth serve to ameliorate pollution through dilution and flushing. Water quality is considered relatively good in the Preserve due to this flushing action.

The northern two thirds of the Preserve lies within the Nassau River drainage basin, a small watershed that covers parts of Duval and Nassau counties. The Nassau River watershed has not yet experienced the concentrated urban and industrial growth found along the St. Johns River; still, portions of the watershed exhibit poor water quality. The area surrounding the Preserve to the west and north is predominantly marsh and low uplands utilized for timbering, residential and agricultural uses (USDI NPS 2000).

TIMU and FOCA are administered as one park. Fort Caroline NM includes approximately 138 acres located along the St. Johns River within the city of Jacksonville and Duval County, Florida. Located primarily on a bluff overlooking the river that rises to a height of nearly 90 feet, the park consists of mixed species forest with fresh water wetlands, preserving an enclave of representative species native to the North Florida-South Georgia community (USDI NPS 2000).

Duval is one of the fastest growing counties in Florida. The Preserve is located in an area that has historically experienced limited development and growth due to lack of easy and quick access. Development and recreational use pressures have increased, however, with the opening of a six-lane bridge in 1989 and ongoing construction of a major highway linking the bridge to the interstate highway system (USDI NPS 2000).

Throughout both watersheds, many residential homes operate private well and septic systems, the failure of which is a presently unquantified source of water pollution. An unknown amount of pesticide, herbicide, and fertilizers is transported by stormwater runoff to the marshes of the Preserve. Contaminated sediments are known to occur in some areas of the Preserve, but the extent of contamination and the effects of sediments resuspension are not known. This is of particular concern as several major dredging projects are proposed in the near future (USDI NPS 2000).

Exotic plants and animals are known to occur within the Preserve, but information on species, locations and potential threats is lacking. The Preserve presently has little information on vegetative and aquatic habitats, ecological processes, and current ecological conditions. Related to the issue of exotic species is the recent development of a prescriptive fire program, which is expected to be instrumental in returning native species to the numerous pine plantations within the Preserve (USDI NPS 2000).

Public lands within TIMU are co-managed by the NPS, State of Florida, City of Jacksonville, US Army Corps of Engineers, US Navy, etc. Many privately owned lands remain within the boundary of TIMU. This mix of land ownership presents some challenging management issues, but also establishes a partnership among many private and government interests that allow opportunities for creative implementation of TIMU's goals (USDI NPS 2000).


## **Avian Resources of South Atlantic Coastal Plain**

The South Atlantic Coastal Plain, consisting of about 10,121,457 million ha, includes parts of Virginia, North Carolina, South Carolina, Georgia, Alabama and Florida (see

# Partners in Flight (PIF) Regions

Southeast Region (SER)


National Park Service  
U.S. Department of the Interior



# Partners in Flight (PIF) Regions and NPS Locations

Southeast Region (SER)

National Park Service  
U.S. Department of the Interior



Produced by Southeast Region GIS, Atlanta, GA

November 2003

PIF and NPS location maps below). This physiographic area is one of four coastal plain divisions recognized by Partners in Flight. Although these coastal plain areas share many conservation issues, differences in key species and habitats exist. For instance, the South Atlantic Coastal Plain includes (1) the largest forested floodplains outside of the Mississippi Alluvial Plain, (2) unique non-alluvial wetlands (Dismal Swamp, pocosins, Carolina bays, Okefenokee Swamp), (3) the largest remnants of the former longleaf pine dominated ecosystems (especially flatwoods and sandhills, and to a lesser extent savannas), (4) the best remaining examples of "natural" barrier and sea islands and maritime forests in the Southeast, and (5) biologically rich Apalachicola Bluff forests. Also present within this physiographic area are extensive tidal wetlands and commercial forests. Physical characteristics include a predominantly flat, weakly dissected alluvial plain with active fluvial deposition and shore zone processes along coastlines. Elevation ranges from 0 feet increasing towards the fall line to 600 feet. Major blackwater rivers (with headwaters in the coastal plain) include Chowan, Waccamaw, Satilla, St. Mary's, Suwanee, and St. John's (originating in Peninsular Florida). Major brownwater rivers (with headwaters originating in the Southern Piedmont or Southern Blue Ridge) include Roanoke, Tar, Neuse, Cape Fear, Pee Dee, Santee-Cooper, Ashepoo-Combahee-Edisto (ACE), Savannah, Ogeechee, Altamaha, and Apalachicola (Chattahoochee and Flint). Average annual precipitation is 40-60 inches except on the Florida Gulf Coast where it is 52-64 inches.

Land conversion for both agricultural and urban expansion has resulted in a 40 percent loss of natural vegetation (closer to 65 percent along some coastlines) in this physiographic area. Potential natural vegetation (i.e., absent frequent disturbances) is referred to as "southern mixed" forests and oak/hickory/pine, with intervening southern floodplain forest and pocosins, as well as live oak/sea oats along coastlines. However, disturbances are frequent and therefore, upland forests historically were characterized by open pine (predominantly longleaf) forests. Today, predominant vegetation remains slash (Florida) and longleaf pines, with loblolly pine becoming common nearer to the Southern Piedmont and the northern portions of this physiographic area. Oak/gum/cypress forest cover type is common along floodplains and prevalent species include laurel oak, water tupelo, swamp tupelo, swamp chestnut oak, cherrybark oak, and baldcypress. Pond pine and Atlantic white cedar become important within the Lower Coastal Plain, especially in pocosin and other non-alluvial wetland types. Live oak becomes important along coastal areas and frequently is included with other coastal pines and hardwoods in various types of "hammocks."

Within the South Atlantic Coastal Plain, fire is the single most important driving disturbance force. Natural burns occur over medium to large size areas between natural barriers (e.g., floodplains, other wetlands) with moderate frequency and low intensity. Fires most often occurred during the growing season, in many cases started by lightning, and were essential for supporting numerous plant communities and dependent animals, including many bird species. In addition to fire, hurricanes, tornadoes, and floods are frequent as disturbance agents. Ice storms, though rare, are devastating where they occur. Finally, southern pine beetles are important disturbance agents.

Conservation issues within the South Atlantic Coastal Plain include:

- (1) management and conservation of forested floodplains and related wintering waterfowl and migratory landbird needs;
- (2) monitoring and protection of colonially nesting terns and skimmers, as well as vulnerable shorebirds, especially in areas with increased human disturbance and habitat loss;
- (3) research and protection of Wood Storks and White Ibises;
- (4) conservation of nongame waterbird habitats (under the purview of other bird conservation groups such as the Western Hemispheric Shorebird Reserve Network, Waterbird Society, North American Waterfowl Management Plan, and the International Association of Fish and Wildlife Agencies' Migratory Shore and Upland Gamebird Subcommittees);
- (5) best management practices for forested wetlands, maritime communities, southern pine forests, and upland hardwood (including riparian) forests; and
- (6) conservation and protection of vulnerable Nearctic Neotropical migratory landbirds.

Over 300 bird species occur annually in the South Atlantic Coastal plain as nesting, post nesting dispersers, transients, and /or wintering residents. Among these species, the South Atlantic Coastal Plain supports critically important populations for a number of extremely high priority bird species. Species in need of the greatest conservation attention include Henslow's Sparrow, Wood Stork, Bachman's Sparrow, Swallow-tailed Kite, Swainson's Warbler, Eastern Painted Bunting, Black-capped and Bermuda Petrels, Red-cockaded Woodpecker, Southeastern American Kestrel, Wayne's Black-throated Green Warbler, Saltmarsh Sharp-tailed Sparrow, Red Knot, Piping Plover, and Snowy Plover (Gulf Coast). Other priority species also of conservation interest include Florida Sandhill Crane, White Ibis, Loggerhead Shrike, Cerulean Warbler, Prothonotary Warbler, Seaside Sparrow, Brown-headed Nuthatch, American Woodcock, Northern Bobwhite, Common Ground-Dove, Yellow-throated Warbler, Rusty Blackbird, Black Skimmer, Least Tern, Black Rail, Peregrine Falcon, Bald Eagle, American Oystercatcher, Red-throated Loon, and most migrating and wintering shorebirds and rails, Brant, American Black Duck, Lesser and Greater Scaup, Tundra Swan, and Wood Duck.

Conservation objectives for the South Atlantic Coastal Plain revolve mostly around (1) stabilizing or increasing populations of high priority breeding bird species, (2) wintering species, (3) and increasing the quality and availability of stopover habitat for transient species. Although waterbirds are treated here, these species groups are

mostly the subjects of other planning efforts. For landbird species, the primary habitat objectives proposed in this plan include the following:

1. Retain and restore 526,000 ha (1.3 million acres) of native warm season grass habitats, with as much associated with longleaf pine as feasible.
2. Provide at least 121,457 ha (300,000 acres) of 5 -year idle lands, 121,457 (300,000 acres) acres of annuals, and 243,000 ha (600,000) acres of 10-20 year idle lands.
3. Maintain and improve the habitat quality of 8 forested wetland sites for Swallow-tailed Kite, maintain and stabilize at least 1 forested wetland site for Cerulean Warbler, at least 10 sites for Wayne's Black-throated Green Warbler, and 30 sites for Swainson's Warblers, which requires 10 patches over 40,485 ha (100,000 acres), 15 patches over 8,100 ha (20,000 acres), 7 patches over 4,050 ha (10,000 acres), and 30 patches over 2,400 ha (6,000 acres).
4. Protect 100% of remaining maritime communities and increase acreage wherever restoration is possible.
5. Increase longleaf pine forest acreage from 607,300 ha (1.5 million acres) to over 890,700 ha (2.2 million acres) and improve conditions favoring warm-season grassy ground cover, on at least 263,157 ha (650,000 acres) by year 2025.

## **Avian Conservation in TIMU**

**Avian Biodiversity:** TIMU is well known for its rich avifauna. At least 339 species have been documented from TIMU or in nearby Duval County (USDI NPS 2003). TIMU has a complete avian inventory and a recently updated checklist of birds that is available for the public. The variety of habitats and the rich bird fauna of TIMU establish this area as an important avian conservation area. Indeed, portions of TIMU have been recently recognized and designated an Important Bird Area of Florida.

Verified records of birds in TIMU have been entered into the NPS I&M program's database, NPSpecies, and may be viewed via the internet at <http://www.nature.nps.gov/im/app/npspp> with a user identification and password combination authorized by the NPS for NPS personnel and NPS cooperators. Many other avian observational data need to be verified and entered into the database.

**Inventory:** The park has recently completed an avian inventory boasting 339 species observed in the park and vicinity. Park managers realized this is the first step necessary to understanding the avifauna of the park and developing appropriate management strategies for birds and other wildlife in the park. TIMU has identified additional inventory needs that will provide further baseline information on which park conservation programs can be implemented.

**Threatened and Endangered Species:** Several Federally listed species occur in TIMU: Wood Stork and Bald Eagle occur as breeders, Piping Plover occurs during migration and winter, and Brown Pelican occurs during all seasons (but not breeding in the park). Furthermore, TIMU is an area where Critical Habitat has been established for the wintering population of the Piping Plover. Extirpated from the eastern US in the mid-1950's, the American Peregrine Falcon (now de-listed) occurs within TIMU during migration and in winter.

Several **Florida** listed species (Appendix C) occur in TIMU and include Swallow-tailed Kite, Southeastern American Kestrel, and Least Tern.

In addition PIF lists other high priority species in need of conservation action such as Red Knot, Reddish Egret, American Black Duck, White Ibis, Roseate Spoonbill, Mottled Duck, Wilson's Plover, Gull-billed Tern, Black Skimmer, American Oystercatcher, American Woodcock, Gray Kingbird, Brown-headed Nuthatch, Painted Bunting, Northern Parula, Hooded Warbler, Saltmarsh Sharp-tailed Sparrow, Seaside Sparrow and many other migrating and wintering shorebirds, warblers and sparrows, all of which occur in TIMU.

**Monitoring:** Currently, several avian monitoring projects are being conducted at TIMU:

- A Christmas Bird Count circle is located in the area and covers a portion of the park
- Shorebird surveys in cooperation with the US NAVY using International Shorebird Survey protocols
- Spring and fall migrations counts are conducted in association with the International Migratory Bird Day (IMBD)

**Research:** Scientific research is permitted within the park. Only one project is currently undertaken:

- Annual survival of the Breeding Population of Painted Bunting in the Southeastern United States

**Outreach:** TIMU does have educational/outreach/interpretive activities related to birds.

- Several bird walks are led each year, highlights of which are the Annual Woodcock Field Trip, Painted Bunting walk, and spring and fall walks
- Specially organized trips for birding groups

### **Park Identified Needs for Avian Conservation**

TIMU has identified several projects that will enhance protection of avian communities at the seashore:

*Inventory:*

- The park desires to have additional inventories, and status and trend data for resident birds in all habitats, migratory passerines and shorebirds, and wintering birds in all habitats

*Monitoring:*

- The park desires to increase its capability to monitor several key bird communities.

*Outreach:*

- The park desires to strengthen it's outreach and visitor education programs since many threats to birds are associated with visitors and their uses of the preserve

## **Coordination with Regional Conservation Initiatives**

**North American Bird Conservation Initiative:** NABCI bird conservation planning units, referred to as Bird Conservation Regions (BCR), are often larger than other planning units associated with other plans, such as Partners In Flight. For example, TIMU is within the NABCI Southeastern Coastal Plain BCR which extends from Virginia south to northern Florida and west to Louisiana north to western Kentucky, following the Atlantic and Gulf coastal plains (see NABCI BCR below) and encompasses several PIF physiographic areas (the planning unit for PIF)(compare to PIF map).


Several NABCI BCR's have coordinators whose primary responsibility is to coordinate all bird conservation planning in the BCR, across all agencies and organizations. Currently, the Southeastern Coastal Plain does not have a designated coordinator; however, a large portion of the BCR lies within the Atlantic Coast Joint Venture area (Maine to Florida and includes Puerto Rico) and the ACJV has several professional bird conservationists base throughout the region to assist partners in bird conservation efforts (see contacts below). This staff can provide valuable assistance to TIMU with implementation of aspects of this ACIP.

**North American Waterfowl Management Plan (NAWMP):** The NAWMP (<http://northamerican.fws.gov/NAWMP/nawmphp.htm>) is completed and has been revised several times, incorporating updated goals and strategies based on new information. This plan is one of the most successful bird conservation delivery programs in the United States, being monetarily supported by the North American Wetlands Conservation Act (NAWCA) and focused primarily on wetland and waterfowl protection, but increasingly these funds have also been utilized for upland non-game species protection. TIMU has several potential projects that could be funded by NAWCA (see Habitat Restoration and Threat Management recommendations and consult ACJV coordinator for qualifying projects).

# Bird Conservation Regions

Southeast Region (SER)

National Park Service  
U.S. Department of the Interior



Produced by Southeast Regional Office GIS, Atlanta, GA

March 2004

**Partners In Flight:** Goals and strategies for the South Atlantic Coastal Plain (SACP) can be found in the draft bird conservation plan, previously submitted to the park. The current plan identifies priority bird and habitat conservation goals that must be implemented in order to achieve bird conservation success in this region. TIMU being a coastal park with all major bird groups represented will utilize all the bird conservation initiative plans. Since shorebird and colonial waterbird plans have not been developed on a regional basis, and the SACP plan covers these species, many of the recommendations in this plan will be derived from the SACP priorities.

Similar to NABCI BCR's, PIF physiographic areas often do not have designated coordinators. However, state level non-game agencies with investment in PIF will establish key personnel to develop partnerships among cooperators in the physiographic area. The State of Florida does have a landbird conservation coordinator and will be instrumental in assisting TIMU to implement recommendations identified in this ACIP and projects important to bird conservation relative to Florida's role in implementation of the SACP plan.

**United States Shorebird Conservation Plan (USSCP):** The USSCP has been completed and is available on the World Wide Web (<http://shorebirdplan.fws.gov/>). A regional step down plan is in preparation by FWS personnel and should be available in 2003. The developing regional shorebird plan will be important for TIMU since many of TIMU's avian resources are related to its shoreline and shorebird use, primarily during migration and winter.

**Waterbird Conservation for the Americas (WCA):** The WCA plan has been completed and is available on the World Wide Web or can be ordered from the US Fish and Wildlife Service National Conservation Training Center (<http://www.waterbirdconservation.org/>). A regional step down plan is in preparation by FWS personnel and should be available in 2003.

## **Integration of NABCI Goals and Objectives into Park Planning and Operations**

### **NABCI Implementation Recommendations**

To successfully achieve park established goals and actively participate in NABCI, the park could implement a variety of projects in different NPS programs. Most of these projects would require some level of participation by many existing park programs and could either be achieved through NPS funding, or more likely, through establishing or improving partnerships with agencies and organizations that already have the necessary expertise to provide guidance, funding, and execution of these programs. Programmatic areas where bird conservation actions are likely to be focused are:

- Inventory
- Monitoring

- Habitat Restoration/Management
- Threat Management (includes exotic species, air quality, water quality, etc.)
- Research
- Compliance
- Outreach
- Partnerships

To the extent appropriate, each of these program areas will be discussed separately and within each, specific opportunities identified that, when implemented, will enable the park to meet its mandates (current and expected), as well as integrate NABCI into its planning and operations. With emphasis added; the park is not expected to implement any of these recommendations or be obligated to pursue any opportunity other than those the park is required to do by law or NPS program or policy. In other words, participation in this effort is currently voluntary. However, implementation of EO 13186 (US Government 2000), Responsibilities of Federal Agencies to Protect Migratory Birds, will require NPS to incorporate a wide range of bird conservation programs into planning and operations. The development of the MOU between the FWS and the NPS will establish a formal agreement to promote bird conservation within the agency by incorporating goals and strategies of existing bird conservation initiatives, plans, and goals into park planning and operations.

Should the park decide to implement any of these projects, further consultation with bird conservation contacts is encouraged to obtain updated information on the relevance of these opportunities in regional bird conservation.

High priority projects are identified in **bold** print. Priorities that the park is encouraged to seek NPS funding for are marked with an asterisk (\*). These projects are those that are critical to the stabilization or improvement of a bird population in the planning region.

**Inventory:** The park has inventoried its bird fauna exceptionally well. Although the park has documented its avifauna well, additional inventory is needed to fully understand the status of birds in the park so that conservation actions for birds can be implemented. Information regarding the status of high priority species (as identified in the Florida's Endangered Species, Threatened Species, and Species of Special Concern and the South Atlantic Coastal Plain bird conservation plan) is needed to effectively structure park management for the continued preservation and enhancement of the park's avifauna and habitats.

#### **Additional surveys are needed**

- **in salt marshes where several high priority species occur in both in summer and winter but data is lacking, especially for secretive marshbirds ("Worthington's" Marsh Wren, Nelson's and Saltmarsh Sharp-tailed Sparrows, Seaside Sparrow, Clapper Rail)\***

Additionally, TIMU is encouraged to:

- **partner with Florida Wildlife and Conservation Commission (FWCC), Cumberland Island National Seashore (CUIS), St. Johns River Water Management District (SJRWMD), The Nature Conservancy (TNC) and Duval Audubon Society (<http://www.duvalaudubon.org/>) to coordinate area inventory effort**
- **continue to partner with the US Navy at Mayport to conduct International Shorebird Surveys**
- **partner with Department of Environmental Protection to survey Pumpkin Hill Preserve State Park**
- **standardize inventory and monitoring methodology to conform to NPS and/or FWS recommended standards (Fancy and Sauer 2000; Hunter 2000)**

**Monitoring:** Efforts should be made to continue existing monitoring programs, striving to conform to established NPS or FWS survey protocols. The park is encouraged to consider establishing permanent monitoring stations in main habitat types to systematically collect data on the distribution and relative abundances of priority species. This information will be useful for documented potential changes in park avifauna resulting from habitat change or management activities. Links to literature detailing inventory and monitoring methodologies for various avian groups (e.g. songbirds, shorebirds, raptors, etc.) can be found at: <http://biology.dbs.umt.edu/landbird/mbcg/groups.htm>. Coordination with the Atlantic Coast Joint Venture Coordinator and FWCC is needed to further identify and implement high priority projects on park lands and to ensure that park efforts contribute to park or regional bird conservation rather than undertake an action or actions that are not needed or are better conducted in other areas. Specific recommendations are to:

- **establish a monitoring program to document nesting and productivity of federally listed Wood Stork and Bald Eagle, and state listed Swallow-tailed Kite, and Least Tern\***
- **establish a migration and wintering monitoring program for Piping Plover (<http://midwest.fws.gov/endangered/pipingplover/recplan-fnl.html>)\***
- **establish a Painted Bunting monitoring program\***
- **strengthen migration monitoring efforts by using standardized protocols for bird groups being monitored\***
- **establish forest point count monitoring to monitor resident landbirds and breeding Neotropical migrants\***

- **establish a marshbird monitoring program following protocols established by the US Geological Survey, Biological Resources Division (USGS-BRD)**
- **establish International Shorebird Survey (ISS) to document migration and winter usage of shorebird habitats**
- **partner with Florida Wildlife and Conservation Commission (FWCC), Cumberland Island National Seashore (CUIS), St. Johns River Water Management District (SJRWMD), TNC, and Duval Audubon Society to coordinate area monitoring efforts**
- **submit monitoring information to appropriate databases (NPSpecies, eBird, South Atlantic Migratory Bird Initiative (SAMBI; <http://samigbird.fws.gov/>) for waterfowl and shorebirds [see Partnerships below])**
- **hire additional staff to support needed monitoring programs\***
- **standardize inventory and monitoring methodology to conform to NPS and/or FWS recommended standards (Fancy and Sauer 2000, Hunter 2000)**

**Habitat Restoration:** Landscape conditions in the Southeastern US have changed dramatically since early European explorers began documenting the area, its habitats, and its inhabitants. Historic landscapes were influenced by Native American burning, wildfire, bison, beaver, and elk, as well as by insect outbreaks and weather events (Hunter et al. 2001, Williams 2002), thus resulting in a landscape mosaic that supported a rich and diverse bird fauna in the Southeast (Barden 1997; Brawn et al. 2001). The arrival of Europeans and the subsequent change in landscape has dramatically effected bird habitat and bird populations. Bird conservationists have long recognized that habitat restoration is critical to restoration of bird populations, stabilizing or reversing bird declines, and removing birds from both State and Federal Threatened and Endangered Species lists. This is no exception for TIMU.

Recently, habitat restoration efforts have increased on NPS lands due to the increased restoration emphasis of the Management Policies (USDI NPS 2001). Parks may use a wide range of management tools to restore wetland, grassland, woodland, and other habitats. Restoration tools include, but are not limited to, forest management practices (e.g. silviculture), prescribed fire, exotic species management, and public use and recreation management. In addition, parks can coordinate infrastructure development (e.g. roads and buildings) with restoration activities to mitigate potential adverse impacts.

Due to the protected nature of TIMU lands, and generally those in the national park system, the condition of habitats for bird use may be of higher quality than other natural, developed, agricultural, or forest lands under other management regimes. However, national park lands are subject to a wide variety of threats, both inside and outside of the park, and habitats can be greatly improved for wildlife, and particularly bird use, by

restoring processes important for habitat formation, succession, and structural development. Largely, these processes have not been managed historically in the national park system, but current policy allows for active management of species, populations, and lands to provide for long-term conservation of park resources for the enjoyment of future generations. Protection, restoration, and enhancement of habitats in TIMU can greatly contribute to established habitat goals identified in the South Atlantic Coastal Plain bird conservation plan and regional shorebird and colonial waterbird conservation plans. TIMU provides excellent area and vegetative cover for a wide variety of birds for nesting, foraging, migrating, and wintering, but some habitats could be improved through management of recreational uses and habitat restoration, and use of prescribed fire in salt marshes and pine communities to mimic historic disturbances. Specific recommendations are to:

- **work with partners to eliminate Off Road Vehicles (ORV's) and associated disturbances to beach nesting birds or wherever ORV's are used**
- **manage recreational uses, including boat use, personal watercraft (PWC) to eliminate disturbance to birds nesting, foraging, and resting on the beach or tidal creeks (potentially prohibit use of PWC and boats if disturbance to birds can be documented)**
- **reintroduce historic disturbances such as fire to the landscape to improve habitat structure and productivity, especially in salt marshes, maritime forests (oak scrub), pine flatwoods, and longleaf pine forests**
- **restore longleaf pine/wiregrass communities**
- **acquire as much land management capability as possible either through land purchase or conservation easement (see US Fish and Wildlife Service private lands contact)**
- **preserve all remaining maritime forest and shrub-scrub areas for resident landbirds, Neotropical migratory birds for breeding and migration stopover**
- **eliminate or minimize conversion of shrub scrub and pine flatwoods to pine plantations either through purchase options or conservation easements**
- **work with partners maintain or restore natural character and function of the beach front and dune systems by allowing natural processes to shape landscape features**
- **work with partners to establish vehicle and pedestrian free areas for shorebird migration and winter resting and foraging areas**
- **restore hydrological processes where appropriate**

- **protect existing snag trees, where not identified as a safety hazard, as important to cavity nesting birds**
- **enhance or maintain water quality, especially in extensive marshes to support aquatic biota necessary to support existing aquatic invertebrates and fish as food sources for birds that forage in the marshes**
- **document all major habitat management activities, including the location (e.g. UTM coordinates) and a description of methods and of pre- and post-management habitat conditions. This information, when coupled with bird distribution and abundance data, is useful for assessing and replicating conservation actions**
- **assess historic landscape cover and determine feasibility of restoring landscape within the context of the park's enabling legislation**

**Threat Management:** Because of the wide variety of land ownerships within TIMU and the multi-agency co-managed nature of TIMU, unified natural resource management will be a challenge. However, within the scope of NPS's authority, TIMU is encouraged to:

- **work with partners, especially the Florida State Park system, to phase out ORV use plan that addresses impacts to birds that nest, rest, and forage on the beach or in other areas where ORV's travel\***
- **manage recreational uses of the seashore, including personal watercraft, kayaking, canoeing, kite boarding, etc. to avoid or minimize disturbance to nesting, foraging, migrating, and wintering birds\***
- **work with partners to identify vehicle and pedestrian free areas on the beach to protect nesting, foraging, and resting birds\***
- **protect Critical Habitat for wintering Piping Plover\***

Feral cats do occur in TIMU, but their impact on birds is unknown; however, feral cats in natural areas always result in some degree of impact to bird life. Additionally, unleashed dogs have been observed to disturb nesting, feeding and resting shorebirds. Staff has identified the need to manage feral cats and unleashed dogs. The park is encouraged to:

- **develop Feral Cat Management Plan similar to that developed at Cape Hatteras National Seashore (Altman2002, Harrison 2002)\***
- **develop and enforce a policy of no unleashed and unattended dogs in the preserve (Huguenot City Park has a potential problem with unleashed dogs and negative impacts to birds in the park)\***

Exotic plants species are not known to be negatively impacting habitat at TIMU. However, it is important to establish and continue inventory and monitoring for exotic plant species, especially the occurrence of exotic species and take appropriate management action before species do alter habitats. TIMU is encouraged to:

- **develop an exotic plant detection and monitoring program**

Additionally, the park is encouraged to:

- **prevent future installation of towers of any kind**
- **determine avian mortality from existing towers**
- **acquire as much land management capability either through land purchase or conservation easement (see US Fish and Wildlife Service private lands contact)**
- **place monofilament line disposal containers in fishing or beach access areas (contact Canaveral Seashore)**
- **hire additional park law enforcement and outreach/education staff to manage visitors and their activities near bird nesting areas**

#### **Research:**

TIMU has many research needs. Basic research on ecological processes and habitats are largely unknown and effects of increasing development and recreational pressures, exotic species, etc. all need investigation. Primary research projects and topics are to:

- **continue support of Painted Bunting research**
- **determine use of maritime and woodland habitats for Neotropical migratory bird breeding and migration\***
- **determine potential to manage habitat to encourage colonization by Southeastern American Kestrel**
- **encourage partners to assess impact of ORV and recreational use on beach nesting birds and foraging and resting shorebirds\***
- **quantify non-point pollution sources and effects from stormwater runoff that carries pesticides, herbicides, and fertilizers, etc. into marsh/estuary system**

- **determine the extent and nature of beached and dead birds along the beach**
- **list park needs and projects on Research Permit and Reporting System web site (RPRS)**
- **develop contact with Cooperative Ecosystem Studies Unit (CESU) at the University of Georgia**

**Compliance:** Park compliance with the Migratory Bird Treaty Act and the Executive Order 13186 (US Government 2000), Responsibilities of Federal Agencies to Protect Migratory Birds, is necessary to assure that park activities incorporate bird conservation into park planning and operations. Further, to ensure that migratory birds are considered in all phases of park planning processes, especially during the National Environmental Policy Act (NEPA) and the Director's Order #12 Compliance processes, the park should consider adding specific language in project evaluations that requires consideration and implications of park projects on migratory birds. The MOU being developed between the NPS and the FWS will likely contain specific language requiring a park to consider implications of park projects on migratory birds, especially those listed in the USFWS Species of Conservation Concern for the South Atlantic Coastal Plain (Appendix D). Compliance considerations for the park are for:

- **park staff to begin specific consideration of migratory birds during park planning processes and partnership planning processes\***
- park staff to attend USFWS training on implementation of EO 13186 at the National Conservation Training Center (NCTC) (when available) or other training on migratory bird conservation in North America. NCTC has several courses and training related to conservation of migratory birds (<http://training.fws.gov/courses.html>).

The USFWS NCTC offers and reserves two tuition free slots for National Park Service employees wishing to attend NCTC courses on a first come, first served basis. Additionally, discount lodging is also available while attending a NCTC course.

## **Outreach**

- **participate in International Migratory Bird Day (IMBD) events with a local partner (<http://birds.fws.gov/imbd.html>) such as Duval Audubon Society\* or with Little Talbot or Big Talbot Island State Park\***
- **develop outreach and educational programs to enhance visibility of bird conservation issues, which may include organized bird walks, owl prowls, and raptor surveys with the public \***
- **continue with construction of a bird observation tower for visitors**

- **support bird conservation by serving shade-grown coffees at meetings, events, and the office buildings in the park**  
(<http://www.americanbirding.org/programs/consbcof3.htm>)
- **develop guided bird walks and programs to be conducted a few times a month**
- subscribe to Florida Birds, an internet based forum for the exchange of information related to Florida birds
- publish articles in local newspapers related to bird protection and conservation activities
- link bird conservation and management literature from park to park's web site home page
- encourage accurate documentation and reporting from random outings by visitors (see Cornell University's eBird monitoring program 2002  
(<http://www.ebird.org/about/index.jsp>))
- encourage park interpretation/education staff to attend USFWS training on Migratory Bird Education at NCTC
- explore cultural affiliation of landscape to inhabitants, both historical and contemporary. Cultures are strongly tied to the landscape they inhabit and birds often play a role in a cultural tie to the landscape. When these connections are discovered and preserved, a greater appreciation for the landscape and its value to the culture can be achieved.

**Partners and Partnerships:** Partnerships for land conservation and protection will perhaps have the greatest positive influence on bird conservation above all other landscape scale planning. Specific recommendations are to:

- **develop and strengthen relationship with Duval Audubon Society for potential cooperation and implementation of segments of this plan\***
- **keep abreast of Duval County initiatives or programs that could impact park resources\***
- **meet with TIMU management partners to discuss implementation of various aspects of this plan\***
- **participate in the active conservation of birds and habitats with the South Atlantic Migratory Bird Initiative (SAMBI), an Atlantic Coast Joint Venture initiative\***

- **develop partnerships with FWCC to develop cooperative projects for bird conservation\***
- **establish partnership with the Longleaf Alliance to assist in plans to restore longleaf pine/wiregrass communities**  
[\(http://www.longleafalliance.org/\)](http://www.longleafalliance.org/)
- **contact the nearest Joint Venture office (see Funding section for explanation of Joint Ventures) or BCR coordinator to develop partnerships and funding proposals tiered to priorities established by the park, this ACIP, and the South Atlantic Coastal Plain bird conservation plan**
- evaluate local or regional land use data and plan potential for habitat protection across organizational boundaries and work with local communities to develop appropriate protection measures

**Funding Opportunities:** Internal NPS funding is often an effective source to obtain funding; however, the project will have to be a fairly high priority among the park's natural resource program to successfully compete for the limited funding available in the NPS. Therefore, partnerships and outside funding programs are often more productive for securing bird conservation funding. TIMU is encouraged to enter all high priority projects into the NPS Performance Management Information System (PMIS) database. Funding for conservation projects for Neotropical migrants is also available through the Park Flight program. Needed at TIMU is:

- **increased base funding to implement basic protection and management needs for birds and their habitats (habitat based management not only benefits the birds but other wildlife as well)**

With the exception of the North American Waterfowl Management Plan (NAWMP and its associated funding legislation, the North American Wetland Conservation Act), funding opportunities for bird conservation programs, plans, and initiatives have been lacking. Only within the last decade have other appropriate and specific sources for bird conservation funding been created and used. The NAWMP has been supported for approximately 14 years by the North American Wetlands Conservation Act (NAWCA 1989). This program has provided \$487 million in appropriated funds matched with \$1.7 billion for wetland and bird conservation projects since its inception. In 2002 alone, over \$70 million US dollars were awarded to US and Canadian agencies and organizations to enhance waterfowl populations by improving, restoring, or protecting wetland habitats. To adequately evaluate projects and distribute these funds, partnerships called Joint Ventures were established. Nationally, 14 (11 US, 3 Canada) joint ventures have been established to facilitate implementation of these programs. The Atlantic Coast Joint Venture is very active in bird conservation in the South Atlantic Coastal Plain and are a primary contact for potential funding (<http://www.acjv.org>) Additional information regarding Joint Ventures can be found at:

(<http://southwest.fws.gov/gulfcoastjv/ojvcontact.html>) and  
(<http://northamerican.fws.gov/NAWMP/jv.htm>).

Funding through NAWCA is highly underutilized by the NPS and any park unit that has wetland, water, or bird conservation needs associated with wetland are encouraged to investigate using this funding source. Naturally, there are certain requirements to be eligible for all grants and park managers are encouraged to consult with the nearest Joint Venture, BCR, or PIF Coordinator to learn how this program might be applicable to implementation of this plan, and other park wetland issues.

Internal FWS funding programs may be used to support projects, but no effective method of project proposal delivery to these sources is currently in place for the NPS. Current funding in these programs may result from FWS familiarity with NPS needs, or NPS participation in one of the area FWS Ecosystem Teams, where a project has been identified and proposed to be funded through the Ecosystem Team.

One largely unexplored yet potentially fruitful funding source for national parks is the myriad of grants through the FWS State Programs, where grants are awarded to private individuals engaged in habitat conservation projects. No funding is directly available to national parks, but identified projects with important or critical adjacent landowners can sometimes be funded through these sources. Similar programs are available if the adjacent landowner is a federally recognized American Indian tribe.

Specific congressional appropriations to protect migratory birds has recently been authorized under the Neotropical Migratory Bird Conservation Act (2000) (<http://www.nfwf.org/programs/nmbcapp.htm>). Appropriations through this Act are authorized up to \$5 million per year. However, in 2000, appropriation was approximately \$3.75 million and a majority of this funding was directed toward projects in Central and South America.

Many of the identified projects are eligible for funding under various grant programs of the National Fish and Wildlife Foundation <http://www.nfwf.org/programs/programs.htm>. Other prominent funding sources available to NPS managers for bird conservation are listed on this projects web site at: <http://southeast.fws.gov/birds/NPSHighlits.htm>.

Funding opportunities for migratory bird conservation are available yet most natural resource agencies are not fully aware of and/or understanding of how to use these sources. Perhaps a consolidated migratory bird funding source catalog will become available to managers in the future; this is needed.

## Contacts

Primary contacts within the region can be obtained by viewing the web site for the Southeastern Bird Conservation Initiative, National Park Service at <http://southeast.fws.gov/birds/npsbirds.htm>. This web site will provide contact information of the appropriate bird conservation coordinator in the region for park personnel. Primary contacts for TIMU are:

### US Fish and Wildlife Service

Keith Watson  
Asheville, NC 28801  
828-350-8228  
[Keith.Watson@fws.gov](mailto:Keith.Watson@fws.gov)

Dean Demarest  
Atlanta, GA  
404-679-7371  
[dean\\_demarest@fws.gov](mailto:dean_demarest@fws.gov)

Jennifer Wheeler  
Waterbird Conservation Plan  
Coordinator  
703-358-1714  
[Jennifer.A.Wheeler@fws.gov](mailto:Jennifer.A.Wheeler@fws.gov)

Craig Watson  
Atlantic Coast Joint Venture  
Charleston, SC  
843-727-4707 ext. 16  
[Craig.Watson@fws.gov](mailto:Craig.Watson@fws.gov)

Chuck Hunter  
Regional Refuge Biologist  
Atlanta, GA  
404-679-7130  
[Chuck.Hunter@fws.gov](mailto:Chuck.Hunter@fws.gov)

Jay Herrington  
Private Lands Biologist  
Jacksonville, FL  
904 232-2580  
[Jay.Herrington@fws.gov](mailto:Jay.Herrington@fws.gov)

### National Park Service

Richard Bryant  
Timucuan Ecological and Historic  
Preserve  
904 221-7567 x15  
[Richard.Bryant@nps.gov](mailto:Richard.Bryant@nps.gov)

Roger Clark  
Timucuan Ecological and Historic  
Preserve  
904 251-3626 x14  
[Roger.Clark@nps.gov](mailto:Roger.Clark@nps.gov)

Chris Furqueron  
Exotic Plant Management Coordinator  
National Park Service  
404-562-3113 ext 540  
[Chris.Furqueron@nps.gov](mailto:Chris.Furqueron@nps.gov)

John Yancy  
Piedmont – South Atlantic Coast  
Cooperative Ecosystem Studies Unit  
University of Georgia  
404-562-3279  
[John.Yancy@nps.gov](mailto:John.Yancy@nps.gov)

### State of Florida

Karl Miller  
Florida Fish and Wildlife Conservation  
Commission  
352-955-2230  
[karl.miller@fwc.state.fl.us](mailto:karl.miller@fwc.state.fl.us)

Jim Rodgers  
Florida Wildlife and Conservation  
Commission  
352-955-2230  
[james.rodgers@fwc.state.fl.us](mailto:james.rodgers@fwc.state.fl.us)

Jeff Gore  
Florida Fish and Wildlife Conservation  
Commission  
[Jeff.Gore@fwc.state.fl.us](mailto:Jeff.Gore@fwc.state.fl.us)

## Others

USDA APHIS Wildlife Services  
Bernice Constantin  
Florida Wildlife Services State Director  
352- 377-5556  
[bernice.u.constantin@aphis.usda.gov](mailto:bernice.u.constantin@aphis.usda.gov)

Joe Meyers  
University of Georgia  
Athens, GA  
706-542-1882  
[joe\\_meyers@usgs.gov](mailto:joe_meyers@usgs.gov)

## LITERATURE CITED

- Altman, J. 2002. Feral Cat Management Final Report for Cape Hatteras National Seashore, Wright Brothers National Memorial, and Fort Raleigh National Historic Site. April 15, 2002. 20p.
- Barden, L.S. 1997. Historic Prairie in the Piedmont of North and South Carolina, USA. *Natural Areas Journal* 17:149-152.
- Brawn, J.D., S.K. Robinson and F. R. Thomson III. 2001. The Role of Disturbance in the Ecology and Conservation of Birds. *Annu. Rev. Ecol. Syst.* 32:251-76.
- Conway, C.J. 2002. Standardized North American Marsh Bird Monitoring Protocols. U.S. Geological Survey, Biological Resources Division. Arizona Cooperative Fish and Wildlife Unit. December 2002. 23p.
- Cornell University and National Audubon Society. 2002. eBird Monitoring Program. Cornell University, Laboratory of Ornithology, Ithaca, NY.
- Fancy, S. and J. Sauer. 2000. Recommended Methods for Inventorying and Monitoring Landbirds in National Parks. National Park Service, Ft. Collins. 13p.
- Harrison, M. 2002. Outer Banks Feral Cat/Cats Indoors Outreach and Education Program. Draft Final Report to Cape Hatteras National Seashore in cooperation with The Nature Conservancy. June 17, 2002. 13p.
- Hunter, C. 2000. Bird Population Survey, Inventory, and Monitoring Standards for National Wildlife Refuges and Partners in the Southeastern U.S. USFWS. January 2002. 49p.
- Hunter, C., D. Buehler, R. Canterbury, J. Confer, and P. Hamel. 2001. Conservation of disturbance-dependent birds in eastern North America. *Wildlife Society Bulletin* 29:440-455.
- Hunter, C., L. Peoples, and J. Collazo. 2001. Partners in Flight Bird Conservation Plan for the South Atlantic Coastal Plain (Physiographic Area 03). American Bird Conservancy, May 2001. 166p.
- Neotropical Migratory Bird Conservation Act. 2000. 114 Stat. 593. 16 U.S.C. 6102 Public Law 106247, enacted July 20, 2000.
- North American Wetlands Conservation Act. 1989. 103 Stat. 1968; 16 U.S.C. 4401-4412. Public Law 101-233, enacted December 13, 1989.
- Sauer, J.R., J.E. Hines, I. Thomas, J. Fallon, and G. Gough. 2000. The North American Breeding Bird Survey, Results and Analysis 1966-1999. Version 98.1, U.S.G.S. Patuxent Wildlife Research Center, Laurel, MD.

- Southeast Natural Resource Leaders Advisory Group (SENRLAG). 2000. A Strategy for Collaboration. Unpublished. 4p.
- USDI, Geological Survey. 2001. National Point Count Database. Biological Research Division, Patuxent Wildlife Research Station, Patuxent, MD.
- USDI, National Park Service. 1916. Organic Act. 39 Stat. 535 16 U.S.C. 1-4, enacted August 25, 1916.
- USDI, National Park Service. 2000. Southeast Coastal Inventory and Monitoring Network Draft Study Plan. December 2000. 148p.
- USDI, National Park Service. 2001. National Park Service Management Policies. 2001. December 2000. 137p.
- USDI, National Park Service. 1997. Resource Management Plan for Timucuan Ecological and Historic Preserve and Fort Caroline National Memorial. 135pp.
- USDI, National Park Service. 2002. Park Flight and Migratory Bird Conservation. National Park Service, Natural Resource Program Center. 2p.
- USDI, National Park Service. 2003. Birds of Timucuan Preserve: Field Checklist. Timucuan National Preserve, National Park Service. 16p.
- US Government. 2000. Presidential Executive Order 13186, Responsibilities of Federal Agencies to Protect Migratory Birds, January 2000.
- US NABCI Committee. 2000. The North American Bird Conservation Initiative in the United States: A Vision of American Bird Conservation. U.S. NABCI Committee, September 2000. 29p.
- Williams, G.W. 2002. Aboriginal Use of Fire: Are There Any "Natural" Plant Communities? USDA Forest Service, Washington D.C., June 2002. 18p.

APPENDIX A

HIGH PRIORITY SPECIES IN THE SOUTH ATLANTIC COASTAL PLAIN  
BIRD CONSERVATION REGION

(From Hunter et al. 2001, Table 1. Priority bird species for South Atlantic Coastal Plain:  
Entry criteria and selection rationale.)

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
1a.	Bewick ' s Wren Appalachian	35	5	5		C	Nearly extinct
	Kirtland ' s Warbler <sup>5</sup>	35	5	5		A	Mostly SC, GA
	Black-capped Petrel	32	5	5		P	Concentrations off NC
	Bermuda Petrel <sup>5</sup>	32	2	5		P	Increasingly regular off NC
	Red Knot South Atlantic	32	5	5		C	Mostly GA, FL
	Red-cockaded Woodpecker <sup>5</sup>	32	5	4	80.4*	R	
	Snowy Plover Southeast	31	3	5		D	St. Joseph Peninsula to Dog Island, FL Gulf
	Painted Bunting Eastern	31	5	5		B	GA, SC, n. FL, se NC
	Roseate Tern <sup>5</sup> North American	30	3	4		A	Highly Pelagic
	Black-throated Green Warbler Wayne ' s (Coastal)	30	5	4	100.0*	B	VA, NC, SC
	Bachman ' s Sparrow	30	5	5	36.6*	R	Primarily breeding
	Saltmarsh Sharp-tailed Sparrow	30	5	3		C	
	Wood Stork <sup>5</sup> Southeast	29	4	4	44.3?	D	FL, GA, se SC
	Henslow ' s Sparrow	29	5	4		D	Winters FL, GA, SC(?), breeding ne NC, se VA
	Swallow-tailed Kite North American	28	4	3	10.8	B	SC, GA, FL
	American Kestrel Southeastern	28	5	4		D	

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
	Piping Plover <sup>5</sup>	28	4	4		D	Mostly winter, breeding NC, possibly SC
	American Oystercatcher North American	28	5	3		D	
Ib.	Short-tailed Hawk Florida	27	2	3		B	St. Marks to Lower Suwannee, FL
	Black Rail	27	4	4		D	
	Sandhill Crane Florida	27	3	3		R	FL, GA
	Brown-headed Nuthatch	27	5	5	38.7*	R	
	Nelson ' s Sharp-tailed Sparrow	27	3	3		C	
	Audubon ' s Shearwater Caribbean	26	5	3		P	
	Yellow Rail	26	4	3		C	
	Wilson ' s Plover	26	4	3		D	Mostly breeds, irregular in winter in GA, FL
	Bicknell ' s Thrush	26	5	3		A	
	Swainson ' s Warbler	26	4	1	15.9	B	
	Seaside Sparrow	26	5	3		D	Atl. and Gulf pops. may represent full species
	Whimbrel	25	5	5		A	
	Buff-breasted Sandpiper	25	3	4		A	
	Black-throated Blue Warbler	25	5	3		A	
	Cerulean Warbler	25	2	3		B	Roanoke River, NC; elsewhere?
	Brown Pelican Southeast	24	5	1		R	
	Marbled Godwit	24	3	4		C	
	Bobolink	24	5	5		A	
	Buff-breasted Sandpiper	24	3	3		A	
	Brant	23	3	5		C	Mostly NC
	King Rail	23	5	4		D	

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
	Sandhill Crane Greater	23	5	3		C	FL, GA
Ib (cont.).	White Ibis	23	5	4 <sup>4</sup>		D	
	Stilt Sandpiper	23	4	5		A	
	Solitary Sandpiper	23	5	3		A	
	American Woodcock	23	5	4		D	Mostly winter, some breeding
	Wood Thrush	23	3	5	8.5*	B	
	Northern Parula	23	5	5	23.7*	B	
	Cape May Warbler	23	5	3		A	
	Worm-eating Warbler	23	3	2	14.7	B	
	Connecticut Warbler	23	5	3		A	
	Hooded Warbler	23	4	4	15.0*	B	
	Cory ' s Shearwater	22	5	3		P	
	White Ibis	22	4	4	15.7?	D	
	American Black Duck	22	3	5		D	Breeds VA, NC; formerly wintered to GA
	Clapper Rail	22	5	3		D	
	Semipalmated Sandpiper	22	5	5		A	
	Purple Sandpiper	22	4	2		C	
	Short-billed Dowitcher	22	5	5		A	Many winter
	Short-eared Owl	22	3	5		C	
	Black Tern	22	5	5		A	
	Sedge Wren	22	4	2		C	
	Veery	22	5	5		A	
	Yellow-throated Warbler	22	4	3	25.5*	D	Mostly breeding, some winter coastal GA, ne FL
	Prairie Warbler	22	3	4	17.9*	B	
	Bay-breasted Warbler	22	3	3		A	
	Louisiana Waterthrush	22	4	2	8.1	B	
	Field Sparrow	22	5	5		D	Primarily winter
	Le Conte ' s Sparrow	22	3	2		C	Mostly GA, SC

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
Ila.	American Bittern	21	4	5		D	Most wintering, local breeding
	Canvasback	21	4	4		C	
	Northern Bobwhite	21	4	5		R	
	Black-bellied Plover	21	4	5		A	Many winter
	Willet	21	5	3		D	
	Ruddy Turnstone	21	5	5		A	Many winter
	Sanderling	21	5	5		A	Many winter
	Western Sandpiper	21	5	3		A	Many winter
	Gull-billed Tern	21	5	4	11.5?	D	
	Least Tern	21	5	5		B	
	Black Skimmer	21	4	5		D	
	Yellow-billed Cuckoo	21	4	5		B	
	Black-throated Green Warbler (all, including Wayne's)	21	5	3		A	
	Grasshopper Sparrow	21	5	5		D	Primarily migration, some breeding and wintering
	Least Bittern	20	5	3		B	
	Lesser Scaup	20	5	5		C	
	Black Scoter	20	4	5		C	
	Northern Harrier	20	4	4		C	
	American Avocet	20	3	3		C	
	Least Sandpiper	20	5	5		A	
	Dunlin	20	4	5		C	
	Sandwich Tern	20	5	3		B	
	Common Ground-Dove	20	3	5	17.6?	R	FL to se SC
	Palm Warbler	20	3	5		C	
	Eastern Towhee	20	5	5	24.5*	D	

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
IIb (cont.).	Red-throated Loon	19	5	4		C	Major concentrations from Back Bay, VA, to Cape Fear, NC, uncommon to rare elsewhere
	Common Loon	19	5	3		C	
	Greater Scaup	19	3	5		C	
	Greater Yellowlegs	19	5	3		A	Some winter
	Pectoral Sandpiper	19	5	3		A	
	Royal Tern	19	5	3	30.6?	D	
	Barn Owl	19	5	3		D	
	Least Flycatcher	19	3	5		A	
	Carolina Chickadee	19	4	4	11.4	R	
	Rusty Blackbird	19	3	5		C	
IIb.	Chuck-will ' s-widow	21	5	2	21.7*	B	
	Prothonotary Warbler	21	4	1	34.4*	B	
	Acadian Flycatcher	20	4	1	13.7	B	
	White-eyed Vireo	20	5	2	17.8	D	Primarily breeding
	Yellow-throated Vireo	19	4	1	10.8*	B	
	Pine Warbler	19	5	2	22.2*	D	
	Summer Tanager	19	5	2	18.6*	B	
	Orchard Oriole	19	5	2	12.9*	B	
IIIa.	Kentucky Warbler	19	2	1	2.5	B	
IIIb.	Bald Eagle <sup>5</sup>	17	3	2		D	

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes	
			Area Importance	Population Trend				
Regional Interest	Great Blue Heron	13	4	1		D		
	Great Egret	14	4	2		D		
	Snowy Egret	14	4	2		D		
	Little Blue Heron	15	4	2		D		
	Tricolored Heron	18	4	3		D		
	Black-crowned Night-Heron	17	4	5		D		
	Yellow-crowned Night-Heron	18	5	2		D		
	Glossy Ibis	17	4	3		D		
	Canada Goose Atlantic pops.	No Score				C	Mostly NC, SC	
	Tundra Swan	20	4	1		C	Mostly ne NC	
	Wood Duck	17	3	2		D		
	Mallard	15	5	3		D	Mostly winter	
	Blue-winged Teal	17	5	3		A	Some winter	
	Northern Pintail	16	3	5		C		
	Redhead	21	3	4		C		
	Ring-necked Duck	19	4	2		C		
	Surf Scoter	20	3	4		C	Mostly NC	
	White-winged Scoter	17	3	4		C	Mostly NC	
	Mississippi Kite	19	3	1		B	Most common FL to SC; Rare and local NC	
	Limpkin	16	2	2		R	Iso. pop. Apalachicola, FL	
	Semipalmated Plover	17	5	3		A	Many winter	
	Spotted Sandpiper	18	5	3		A	Many winter	
	Lesser Yellowlegs	18	5	3		A	Many winter	
	Common Tern	16	3	4		D	Of special concern VA, NC	
	Forster's Tern	19	2	3		D		
	Whip-poor-will	18	3	1		B		
	Red-headed Woodpecker	19	4	2		4.8	D	Primarily breeding

Priority Entry Criteria <sup>1</sup>	Species	Total PIF Priority Species Score	Score		Percent of BBS Population	Local Migratory Status <sup>2</sup>	Geographical or Historical Notes
			Area Importance	Population Trend			
Regional Interest (cont.).	Eastern Wood-Pewee	18	4	2		B	
	Eastern Kingbird	18	4	4		B	
	Loggerhead Shrike	19	3	4		D	Rare now in NC, VA Primarily breeding, rare winter coastal GA, FL
	Black-and-white Warbler	14	2	1		D	
	Yellow-breasted Chat	16	4	1		B	
	Eastern Meadowlark	16	2	5		D	

<sup>1</sup>Entry criteria:

- Ia. Overall Highest Priority Species. Species with total score 28-35. Ordered by total score. Consider deleting species with AI  $\leq$  2 confirmed to be of peripheral occurrence and not of local conservation interest, but retain species potentially undersampled by BBS or known to have greatly declined during this century.
- Ib. Overall High Priority Species. Species with total score 22-27. Ordered by total score. Consider deleting species with AI  $\leq$  2 confirmed to be of peripheral occurrence and not of local conservation interest, but retain species potentially undersampled by BBS or known to have greatly declined during this century.
- IIa. Area Priority Species. Species with slightly lower score total 19-21 with PT+AI=8+. Ordered by total score. These are overall moderate priority species.
- IIb. Species with High Percent of BBS Population. Species with score total 19-21 with percent of BBS population above a threshold established (based on relative size of physiographic area), not already listed above, ordered by total score (\*signifies highest percentage among physiographic area). These are overall moderate priority species.
- IIIa. Additional Species of Global Priority. Add WatchList species (Partners in Flight-National Audubon Society priority species at national level), not already listed in either I or II, with AI=2+. Order by total score. Consider deleting species with AI=2 if confirmed to be of peripheral occurrence and not of local conservation interest, but retain if a local population is viable and/or manageable. These are also overall moderate priority species.
- IIIb. Additional Federally Listed Species. Federal listed species if not already included above. Overall low priority, but appropriate legal obligations (Alegal priority species@) to protect through appropriate management and monitoring still apply. Only Bald Eagle meets this criterion in some Southeast physiographic areas.

Other Local or Regional Interest Species. Includes game or nongame species identified by State Working Groups. Also, may include species often meeting criteria for I or II within other physiographic areas and therefore of regional interest for monitoring throughout the Southeast. These are overall low priority species within physiographic area, but may be more important within one or more States (especially where multiple states have designated some special protective status on the species).

<sup>2</sup> Local Migratory Status, codes adapted from Texas Partners in Flight as follows:

- A = Breeds in temperate or tropical areas outside of region, and winters in temperate or tropics outside of region (*i.e.*, passage migrant).
- B = Breeds in temperate or tropical areas including the region, and winters exclusively in temperate or tropics outside the region (*i.e.*, includes both breeding and transient populations).
- C = Breeds in temperate or tropical areas outside of region, and winters in both the region and in temperate or tropical areas beyond area (*i.e.*, includes both transient and wintering populations).
- D = Breeds and winters in the region, with perhaps different populations involved, including populations moving through to winter beyond the region in temperate or tropical areas (*i.e.*, populations may be present throughout year, but may include a large number of passage migrants).
- E = Species reaching distributional limits within the region, either as short-distance or long-distance breeding migrants, but at population levels above peripheral status.
- F = Same as E except for wintering (non-breeding) migrants.
- R = Resident, generally non-migratory species (though there may be local movements).
- RP= Resident, non-migratory species, reaching distributional limits within the region, but at population levels above peripheral status.
- P = Pelagic, breeding grounds outside of region, but can occur during breeding season.
- PB = Post-breeding dispersal or non-breeding resident; species present during breeding season, but not known to be breeding in the region proper.

<sup>3</sup>Highest percent of breeding population recorded in temperate North America indicated by A\*@; ? indicates species widespread outside of temperate North America and/or waterbirds poorly sampled by Breeding Bird Survey within physio. area.

<sup>4</sup>AI or PT score revised from what was derived by BBS data, or lack thereof, based on better local information.

<sup>5</sup>Species listed as either Federal Endangered or Threatened.

## APPENDIX B

### SOUTH ATLANTIC COASTAL PLAIN BIRD ASSEMBLAGES AND HABITAT CONSERVATION PRIORITIES (from Hunter et al. 2001, Table 4. South Atlantic Coastal Plain Bird-Habitat Associations TB=threats breeding score, TN=threats non-breeding score)

	Total Score	TB	TN	Notes
<b>PRAIRIES, SAVANNAS, AND GRASSLANDS, OPEN COUNTRY</b>				
<u>Extremely High Priority</u>				
Bachman ' s Sparrow	30	4	4	Primarily breeding
Henslow ' s Sparrow	29		4	FL, GA, SC(?)
<u>High Priority</u>				
Sandhill Crane (Florida)	27	4	3	FL, GA
Henslow ' s Sparrow	26	4		NC, VA
Yellow Rail	26		4	
Bobolink	24		4	
Buff-breasted Sandpiper	24		3	Turf farms, airports, pastures
Sandhill Crane (Greater)	23		3	FL, GA
American Woodcock	23	3	3	Primarily winter
Northern Bobwhite	22	3	3	
Short-eared Owl	22		4	
Sedge Wren	22		3	
LeConte ' s Sparrow	22		4	Most in GA and SC
<u>Moderate Priority</u>				
Grasshopper Sparrow	21	3	3	Primarily migration
Loggerhead Shrike	20	4	3	Rare now in NC, VA
Palm Warbler	20		2	
Northern Harrier	20		3	
Barn Owl	19	3	3	
<u>Local or Regional Interest</u>				
Eastern Kingbird	18	3	2	
Eastern Meadowlark	17	3	3	
Bald Eagle	17	3	3	
<b>EARLY SUCCESSIONAL SHRUB-SCRUB</b>				
<u>Extremely High Priority</u>				
Bewick ' s Wren (Appalachian)	35		5	Nearing extinction
Painted Bunting (Eastern)	31	4		GA, SC, n. FL, se NC
Bachman ' s Sparrow	30	4	4	Primarily breeding
Henslow ' s Sparrow	29		4	FL, GA, SC (?)

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>EARLY SUCCESSIONAL (CONT.)</b>				
<u>High Priority</u>				
Henslow's Sparrow	26	4		NC, VA
American Woodcock	23	3	3	Primarily winter
Prairie Warbler	23	3		
Northern Bobwhite	22	3	3	
Field Sparrow	22	3	3	Primarily winter
<u>Moderate Priority</u>				
Common Ground-Dove	20	4	3	FL to se SC
Eastern Towhee	20	3	2	
Palm Warbler	20		2	
White-eyed Vireo	19	3	2	Primarily breeding
Orchard Oriole	19	3		
<u>Local or Regional Interest</u>				
Whip-poor-will	18	3		Ground nesting
Yellow-breasted Chat	16	3	2	
<b>SOUTHERN PINE (SAVANNAS, FLATWOODS, SANDHILLS)</b>				
<u>Extremely High Priority</u>				
Red-cockaded Woodpecker	32	5	5	Cavity nesting
Bachman's Sparrow	30	4	4	Primarily breeding, ground nesting
Henslow's Sparrow	29		4	Flatwoods, savannas, ground
American Kestrel (Southeast)	28	4	3	Primarily sandhills, cavity nesting
<u>High Priority</u>				
Brown-headed Nuthatch	27	3	3	Cavity nesting
Prairie Warbler	23	3		Understory
Northern Bobwhite	22	3	3	Ground
<u>Moderate Priority</u>				
Red-headed Woodpecker	21	3	3	Primarily breeding, cavity nesting
Chuck-will's-widow	21	3		Ground, open understory
Pine Warbler	19	2	2	

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>CONIFER-HARDWOOD GENERALISTS (INCLUDING SPECIES USING BOTH PINE DOMINATED AND HARDWOOD DOMINATED STANDS)</b>				
<u>Extremely High Priority</u>				
Black-throated Green Warbler	30	4		VA, NC, ne SC; canopy, often non-alluvial wetlands
<u>High Priority</u>				
Wood Thrush	24	3		Midstory nesting, ground foraging
Northern Parula	23	3		Canopy
Hooded Warbler	23	3		Understory
Worm-eating Warbler	23	3		Ground nesting
Yellow-throated Warbler	22	3		Mostly breeding, canopy
<u>Moderate Priority</u>				
Yellow-billed Cuckoo	21	3		Upper midstory
Carolina Chickadee	20	2	1	Cavity nesting
<u>ΔWatchlist@ Species</u>				
Kentucky Warbler	20	3		Ground nesting
<u>Local or Regional Interest</u>				
Acadian Flycatcher	20	3		Midstory
Summer Tanager	19	3		Canopy
Yellow-throated Vireo	19	3		Canopy
Eastern Wood-Pewee	18	3		Midstory
Black-and-white Warbler	14	2	2	Primarily breeding, ground nesting
<b>FORESTED WETLANDS (ALLUVIAL AND NON-ALLUVIAL, EXCEPT POND PINE [TALL] POCOSIN)</b>				
<u>Extremely High Priority</u>				
Swallow-tailed Kite (Southeast)	28	4		Nests in Δsuper-emergent@ trees
Swainson ' s Warbler	28	4		Understory, forages ground
<u>High Priority</u>				
Short-tailed Hawk (Florida)	27	4		St. Marks to Lower Suwannee, FL
Cerulean Warbler	25	4		Roanoke River, NC
American Woodcock	23	3	3	Understory, forages ground
American Black Duck	22	4	3	Breeds VA, NC; formerly wintered to GA

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>FORESTED WETLANDS (CONT.)</b>				
<u>Moderate Priority</u>				
Prothonotary Warbler	21	3		Cavity nesting
Louisiana Waterthrush	21	3		Streamside
Rusty Blackbird	19		3	Roosts in trees, forages ground
<u>Local or Regional Interest</u>				
Wood Duck	19	3	3	Cavity nesting
Mississippi Kite	19	3		Edge nesting
Bald Eagle <sup>1</sup>	17	3	3	
Limpkin (Florida)	17	3	3	Apalachicola, Suwannee
<b>POND PINE (TALL) POCOSIN</b>				
<u>Extremely High Priority</u>				
Red-cockaded Woodpecker	32	5	5	Cavity nesting
Swainson ' s Warbler	28	4		Understory, forages ground
<u>High Priority</u>				
Brown-headed Nuthatch	27	3	3	Cavity nesting
American Woodcock	23	3	3	Understory, forages ground
Prairie Warbler	23	3		Understory
Northern Bobwhite	22	3	3	Ground
Prothonotary Warbler	22	3		Cavity nesting
<u>Moderate Priority</u>				
Red-headed Woodpecker	21	3	3	Primarily breeding, cavity nesting
Rusty Blackbird	19		3	Roosts in trees, forages ground
Chuck-will ' s-widow	21	3		Ground, open understory
Louisiana Waterthrush	21	3		Streamside
Pine Warbler	19	2	2	
<u>Local or Regional Interest</u>				
Wood Duck	19	3	3	Cavity nesting
<b>MARITIME WOODLANDS</b> <b>(many of the same species under</b> <b>pine-hardwood, but also transient</b> <b>landbirds and 2 breeding species)</b>				
<u>Extremely High Priority</u>				
Kirtland ' s Warbler	35		5	
Painted Bunting (Eastern)	31	4		GA, SC, ne FL, se NC; edges

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>MARITIME WOODLANDS (CONT.)</b>				
<u>High Priority</u>				
Bicknell ' s Thrush	26		4	
Black-throated Blue Warbler	25		4	
Cape May Warbler	23		3	
Connecticut Warbler	23		2	
Veery	22		3	
Bay-breasted Warbler	22		3	
<u>Moderate Priority</u>				
Black-throated Green Warbler (All, including Wayne ' s)	21		3	
Common Ground-Dove	20	4	3	Ground nesting
Least Flycatcher	19		2	
<b>COLONIAL TREE AND/OR BRUSH NESTING WATERBIRDS (most species feed in emergent wetlands, open water, or mudflats)</b>				
<u>Extremely High Priority</u>				
Wood Stork (Southeast)	29	4	3	FL, GA, se SC
<u>High Priority</u>				
Brown Pelican (Southeast)	24	4	3	Coastal
White Ibis	22	4	2	
<u>Local or Regional Interest</u>				
Tricolored Heron	18	2	2	
Yellow-crowned Night-Heron	18	3	2	
Black-crowned Night-Heron	17	2	2	
Little Blue Heron	15	3	2	
Great Egret	14	2	2	
Snowy Egret	14	2	2	
Great Blue Heron	13	2	2	
<b>COLONIAL GROUND NESTING WATERBIRDS (most species feeding in open water or emergent wetlands)</b>				
<u>Moderate Priority</u>				
Black Skimmer	21	3	2	Beaches, dunes, rooftops
Gull-billed Tern	21	3		Marshes, protected islets
Least Tern	21	4		Beaches, dunes, rooftops
Sandwich Tern	20	3		Protected islets
Royal Tern	19	3	2	Protected islets

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>COLONIAL GROUND NESTING WATERBIRDS (CONT.)</b>				
<u>Local or Regional Interest</u>				
Forster ' s Tern	19	3	2	Marshes, NC
Glossy Ibis	17	3	2	Marshes
Common Tern	16	3	2	Protected islets, NC
<b>EMERGENT WETLANDS</b>				
<u>Extremely High Priority</u>				
Saltmarsh Sharp-tailed Sparrow	30		4	Coastal
<u>High Priority</u>				
Black Rail	27	4	4	
Nelson ' s Sharp-tailed Sparrow	27		4	Coastal
Yellow Rail	26		4	
Seaside Sparrow	26	3	3	Coastal
King Rail	23	3	3	
American Black Duck	22	4	3	Mostly NC, formerly to GA
Clapper Rail	22	3	3	Coastal
<u>Moderate Priority</u>				
American Bittern	21	3	3	Most wintering, local breeding
Least Bittern	20	3		
Northern Harrier	20		3	
<u>Local or Regional Interest</u>				
Peregrine Falcon	19		3	
Bald Eagle	17	3	3	
<b>BEACHFRONT</b>				
<u>Extremely High Priroity</u>				
Red Knot (South Atlantic)	32		4	Mostly GA, FL
Snowy Plover (Southeast Gulf)	31	5	4	St. Joseph Peninsula to Dog Island
Piping Plover	28	4	4	Mostly winter, local breeding NC (SC?)
American Oystercatcher (Eastern North America)	28	4	4	
<u>High Priority</u>				
Wilson ' s Plover	26	4	4	
Purple Sandpiper	22		3	Rocky coastal areas

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>BEACHFRONT (CONT.)</b>				
<u>Moderate Priority</u>				
Willet	21	3	2	
Black-bellied Plover	21		3	Many overwinter
Sanderling	21		4	Many overwinter
Ruddy Turnstone	21		4	Many overwinter
<u>Local or Regional Interest</u>				
Peregrine Falcon	19		3	Some overwinter
<b>ESTUARIES, MUDFLATS, AND IMPOUNDMENTS</b>				
<u>High Priority</u>				
Whimbrel	25		4	Some overwinter
Marbled Godwit	24		4	
Stilt Sandpiper	23		3	Mostly inland
Solitary Sandpiper	23		2	Mostly inland
Semipalmated Sandpiper	22		3	
Short-billed Dowitcher	22		3	Many winter
Buff-breasted Sandpiper	25		4	Mostly inland
Black Tern	22		3	
<u>Moderate Priority</u>				
Western Sandpiper	21		4	Many winter
American Avocet	20		4	
Dunlin	20		3	
Least Sandpiper	20		2	Many winter
Greater Yellowlegs	19		2	Some winter
Pectoral Sandpiper	19		2	Mostly inland
<u>High Percent of Continental Population</u>				
Semipalmated Plover	17		2	Many winter
Spotted Sandpiper	18		2	Many winter
Lesser Yellowlegs	18		2	Many winter
<b>OPEN WATER</b>				
<u>Extremely High Priority</u>				
Black-capped Petrel	32		3	Pelagic
Bermuda Petrel	32		5	Pelagic
Roseate Tern (North American)	30		3	Pelagic

Table 4 (cont.).

	Total Score	TB	TN	Notes
<b>OPEN WATER (CONT.)</b>				
<u>High Priority</u>				
Brant	23		3	Mostly NC
Audubon's Shearwater (Caribbean)	26		4	Pelagic
Cory's Shearwater	22		3	Pelagic
American Black Duck	22	4	3	Breeds VA, NC; formerly wintered to GA
<u>Moderate Priority</u>				
Canvasback	21		2	
Lesser Scaup	20		3	
Black Scoter	20		3	
Greater Scaup	19		3	
Common Loon	19		3	
Red-throated Loon	19		3	Major concentrations from Back Bay, VA, to Cape Fear, NC, uncommon to rare elsewhere
<u>Local or Regional Interest</u>				
Tundra Swan	20		3	NC (especially, Mattamuskeet NWR)
Wood Duck	19	3	3	
Mallard	15	2	2	Mostly winter
Blue-winged Teal	17		2	Some overwinter
Northern Pintail	16		2	
Redhead	21		3	
Rin-necked Duck	19		3	
Surf Scoter	20		3	
White-winged Scoter	17		3	
Canada Goose (Atlantic pop.)	???			

**APPENDIX C**  
 Wildlife Conservation Commission  
**FLORIDA'S ENDANGERED  
 SPECIES, THREATENED SPECIES  
 AND SPECIES OF SPECIAL CONCERN**  
*Official Lists*

Publication Date: 1 August 1997

This document consolidates the state and federal official lists of endangered species, threatened species, and other species categorized in some way by the respective jurisdictional agencies as meriting special protection or consideration. The state lists of animals are maintained by the Florida Game and Fresh Water Fish Commission and categorized as endangered, threatened and of special concern, and constitute Rules 39-27.003, 39-27.004 and 39-27.005, respectively, Florida Administrative Code (F.A.C.). The state lists of plants are categorized into endangered, threatened and commercially exploited, and are administered and maintained by the Florida Department of Agriculture and Consumer Services via Chapter 5B-40, F.A.C. The federal lists of animals and plants are administered by the U.S. Fish and Wildlife Service and categorized into endangered and threatened, and are published in 50 CFR 17 (animals) and 50 CFR 23 (plants). The abbreviations used in part one are:

- GFC = Florida Fish and Wildlife Conservation Commission
- FDA = Florida Department of Agriculture and Consumer Services
- FWS = United States Fish and Wildlife Service
- E = Endangered
- T = Threatened
- T(S/A) = Threatened/Similarity of Appearance
- T(E/P) = Threatened/Experimental Population
- SSC = Species of Special Concern
- C = Commercially Exploited

		Designated Status	
Scientific Name	Common Name(s)	GFC	FWS
<b><u>Birds</u></b>			
<i>Ajaia ajaja</i>	Roseate spoonbill	SSC	
<i>Ammodramus maritimus juncicolus</i>	Wakulla seaside sparrow	SSC	
<i>Ammodramus maritimus mirabilis</i>	Cape Sable seaside sparrow	E	E
<i>Ammodramus maritimus peninsulae</i>	Scott's seaside sparrow	SSC	
<i>Ammodramus savannarum floridanus</i>	Florida grasshopper sparrow	E	

		Designated Status	
Scientific Name	Common Name(s)	GFC	FWS
<i>Aphelocoma coerulescens</i>	Florida scrub-jay	T	T
<i>Aramus guarana</i>	Limpkin	SSC	
<i>Campephilus principalis</i>	Ivory-billed woodpecker	E	E
<i>Charadrius alexandrinus tenuirostris</i>	Southeastern snowy plover	T	
<i>Charadrius melodus</i>	Piping plover	T	T
<i>Cistothorus palustris griseus</i>	Worthington's marsh wren	SSC	
<i>Cistothorus palustris marianae</i>	Marian's marsh wren	SSC	
<i>Columba leucocephala</i>	White-crowned pigeon	T	
<i>Dendroica kirtlandii</i>	Kirtland's warbler	E	E
<i>Egretta caerulea</i>	Little blue heron	SSC	
<i>Egretta rufescens</i>	Reddish egret	SSC	
<i>Egretta thula</i>	Snowy egret	SSC	
<i>Egretta tricolor</i>	Tricolored (=Louisiana) heron	SSC	
<i>Eudocimus albus</i>	White ibis	SSC	
<i>Falco peregrinus tundrius</i>	Arctic peregrine falcon	E	
<i>Falco sparverius paulus</i>	Southeastern American kestrel	T	
<i>Grus americana</i>	Whooping crane	SSC	T(E/P)
<i>Grus canadensis pratensis</i>	Florida Sandhill crane	T	
<i>Haematopus palliatus</i>	American oystercatcher	SSC	
<i>Haliaeetus leucocephalus</i>	Bald eagle	T	T
<i>Mycteria americana</i>	Wood stork	E	E
<i>Pandion haliaetus</i>	Osprey	SSC*	

		Designated Status	
Scientific Name	Common Name(s)	GFC	FWS
<i>Pelecanus occidentalis</i>	Brown pelican	SSC	
<i>Picoides borealis</i>	Red-cockaded woodpecker	T	E
<i>Polyborus plancus audubonii</i>	Audubon's crested caracara	T	T
<i>Rostrhamus sociabilis</i>	Snail kite	E	E
<i>Rynchops niger</i>	Black skimmer	SSC	
<i>Speotyto cunicularia</i>	Burrowing owl	SSC	
<i>Sterna antillarum</i>	Least tern	T	
<i>Sterna dougallii</i>	Roseate tern	T	T
<i>Vermivora bachmanii</i>	Bachman's warbler	E	E
*Applicable in Monroe County only			

## APPENDIX D

### US FISH AND WILDLIFE SERVICE SPECIES OF CONSERVATION CONCERN (2002) in the SOUTHEASTERN COASTAL PLAIN (BCR 27)

Black-capped Petrel	Common Tern
Audubon's Shearwater	Least Tern (except where Endangered)
Little Blue Heron	Black Tern
Reddish Egret	Black Skimmer
Swallow-tailed Kite	Common Ground-Dove
Short-tailed Hawk	Burrowing Owl
American Kestrel (resident <i>paulus</i> ssp. only)	Chuck-will's-widow
<i>Peregrine Falcon</i>	Brown-headed Nuthatch
Yellow Rail	Bewick's Wren
Black Rail	Wood Thrush
Limpkin	Northern Parula
Snowy Plover	Black-throated Green Warbler
Wilson's Plover	Prairie Warbler
American Oystercatcher	Cerulean Warbler
Whimbrel	Swainson's Warbler
Marbled Godwit	Bachman's Sparrow
Red Knot	Henslow's Sparrow
Semipalmated Sandpiper	Le Conte's Sparrow
Stilt Sandpiper	Nelson's Sharp-tailed Sparrow
Buff-breasted Sandpiper	Saltmarsh Sharp-tailed Sparrow
Short-billed Dowitcher	Seaside Sparrow
Gull-billed Tern	Painted Bunting
	Orchard Oriole