


Interior Population of the Least Tern

Sterna Antillarum

Recovery Plan



RECOVERY PLAN FOR THE INTERIOR POPULATION OF THE

LEAST TERN (Sterna antillarum)

September 1990

Prepared by

John G. Sidle
U. S. Fish and Wildlife Service
203 West Second Street
Grand Island, Nebraska 68801

and

William F. Harrison
U. S. Fish and Wildlife Service
Federal Building, Fort Snelling
Twin Cities, Minnesota 55111

Approved: _____

Regional Director Date

EXECUTIVE SUMMARY OF THE RECOVERY PLAN FOR THE LEAST TERN

CURRENT STATUS: The interior population of the least tern (*Sterna antillarum*), a breeding migratory bird in mid-America, was listed as endangered on June 27, 1985 (50 Federal Register 21,784-21,792) and data currently indicate about 5,000 Interior least terns.

Habitat Requirements and Limiting Factors: Least terns breed in the Mississippi and Rio Grande River Basins from Montana to Texas and from eastern New Mexico and Colorado to Indiana and Louisiana. From late April to August they occur primarily on barren to sparsely vegetated riverine sandbars, dike field sandbar islands, sand and gravel pits, and lake shorelines. Threats to the survival of the species include the actual and functional loss of riverine sandbar habitat. Channelization and impoundment of rivers have directly eliminated nesting habitat. This recovery plan outlines recovery strategies to increase the interior population of the least tern to approximately 7,000 birds throughout its range.

Recovery Objective: **Delisting.**

Recovery Criteria: Assure the protection of essential habitat by removal of current threats and habitat enhancement, establish agreed upon management plans, and attain a population of 7,000 birds at the levels listed below.

1. Adult birds in the Missouri River system will increase to 2,100 and remain stable for 10 years.
2. Current number of adult birds (2,200-2,500) on the Lower Mississippi River will remain stable for 10 years.
3. Adult birds in the Arkansas River system will increase to 1,600 and remain stable for 10 years.
4. Adult birds in the Red River system will increase to 300 and remain stable for 10 years.
5. Current number of adult birds in the Rio Grande River system (500) will remain stable for 10 years.

Actions Needed:

1. Determine population trends and habitat requirements.
2. Protect, enhance and increase populations during breeding.
3. Manage reservoir and river water levels to the benefit of the species.
4. Develop public awareness and implement educational programs about the interior least tern.
5. Implement law enforcement actions at nesting areas in conflict with high public use.

Cost of Recovery: Estimated to be \$1,720,000 - \$2,000,000, to reach recovery criteria set out above, and complete subsequent monitoring for 10 years.

Date of Recovery: Delisting should be initiated in 2005, if recovery criteria have been met.

DISCLAIMER

Recovery **plans** delineate reasonable actions which are believed to be required to recover **and/or** protect listed species. **Plans** are published by the U.S. Fish and Wildlife Service, sometimes prepared **with** the assistance of **recovery teams**, contractors, State agencies, and others. Objectives will be attained and any necessary funds **made** available subject to budgetary constraints affecting the parties involved, as **well** as the need to address other priorities. Recovery plans do **not** necessarily represent the **views** nor the official positions or approval' of any individuals or agencies involved in the plan formulation, other than the U.S. Fish and Wildlife **Service**. They represent the **official** position of the **U. S. Fish and Wildlife Service** **only** after they have been signed by the Regional Director es ~~approved~~ recovery plans are subject to modification as dictated by new findings, changes in species **status**, and the completion of recovery **tasks**.

Literature Citation should read as follows:

U. S. Fish and Wildlife Service. 1990. Recovery plan for the interior population of the least tern (*Sterna antillarum*). U. S. Fish and Wildlife **Service**, Twin Cities, **Minnesota**. 91 pp.

Additional copies **may** be purchased from:

U. S. Fish and Wildlife Service Reference Service
5430 Grosvenor Lane, Suite 110
Bethesda, **Maryland** 20814 .
301/492-6403 or **1-800-582-3421**

The fee for the plan **varies** depending on the number of **pages** of the plan.

ACKNOWLEDGMENTS

The following individuals **are** gratefully **acknowledged for** reviewing drafts of the recovery plan and providing helpful assistance during the formulation of the plan: F. **Bagley**, D. Bowman, **J. Brabander**, **M. Dryer**, L. Hill, K. Higgins, E. **Kirsch**, N. **McPhillips**, **K. Nemec**, K. Smith, B. Williams, D. Christopherson, P. **Mayer**, D. Jordan, D. James, **J. Engel**, S. Hoffman, O. Bray, B. **Osmundson**, K. Russell, A. **Sappa**, **K. Keenlyne**, P. McKenzie, C. **Hintz**, and **R. Bovker**, We also thank P. Percy for her assistance with graphics and word processing. **An additional** list of individuals who also **reviewed** the plan and furnished **comments** is found in the appendices'.

TABLE OF CONTENTS

Page No.

Title page	i
Summary	ii
Disclaimer	iii
Acknowledgmentsiii
Table of Contents	iv
List of Figures and Tables	iv
I. INTRODUCTION	1
Description	2
Taxonomy	2
Distribution	3
Life History	12
Habitat Requirements	20
Reasons for Current Status	22
Conservation Efforts	23
II. RECOVERY	28
Recovery Objective	28
Step-down Outline	29
Narrative	32
References	51
III. IMPLEMENTATION	60
IV. APPENDICES	67
1. State Contact People	67
2. Agreements Necessary for Protection of Essential Habitat	69
3. Example of a Memorandum of Agreement to Protect Least Terns and their Habitat	70
4. Essential Breeding Habitat	73
5. List of Reviewers	89

LIST OF FIGURES

1. Breeding distribution of interior least terns in North America	11
--	----

LIST OF TABLES

1. Known breeding areas of interior least terns in the Missouri River system	6
2. Known breeding areas of interior least terns on the Mississippi and Ohio Rivers	8
3. Known breeding areas of interior least terns in the Arkansas River system	9
4. Known breeding areas of interior least terns in the Red River system	10
5. Known breeding areas of interior least terns in the Rio Grande River system	10
6. Productivity of interior least terns	13
7. Census data on interior least terns, 1985-1988	15
8. Recommended annual flow regime for Central Platte River, Nebraska	26

Description

Least terns (all currently recognized subspecies and populations) are the smallest members of the subfamily **Sterninae** and family **Laridae** of the order **Charadriiformes**, measuring about 21-24 cm long with a 51 cm wingspread. **Sexes** are alike, characterized by a black-capped **crow**, white forehead, grayish back and dorsal wing surfaces, **snowy white** undersurfaces, legs of various orange and yellow colors depending on the sex, and a black-tipped bill whose color also varies depending on sex (Watson 1966, Davis 1968, Boyd and Thompson 1985). Boyd and Thompson (1985) developed the following criteria to **distinguish** the sexes in the field based upon their work in Kansas:

- 1) Females usually have a **wing** chord less than 171 mm long while males usually have a **wing** chord greater than **174 mm**.
- 2) A **male's** feet are brighter than its mate's **'feet: the male's are** bright orange, while the female's feet are bright to pale yellow, or rarely **grey**.
- 3) A male's bill is larger than the female's; the female's bill depth at its widest point is 4.5 **mm** to 5.5 **mm**, while the male's is 6.0 **mm** or greater.
- 4) A male's bill is orange to bright yellow, whereas the female's bill is light or dull yellow, or-straw-colored.

Immature birds have darker plumage than adults, a dark bill, and dark **eye** stripes on their **white** foreheads. Jackson (1976) described the developmental stages of least tern chicks. Further details on plumage development and variation **were** presented by Massey and Atwood (1978) and Thompson and Slack (1983).

Taxonomy

The least tern (**Sterna antillarum**) in North America was described by Lesson in 1847 (Ridgway 1895, American Ornithologists' Union 1957, 1983). The least tern in interior North **America was** described later as a race (**Sterna albifrons athalassos**) of the Old World little tern (**Sterna albifrons**) (Burleigh and Lowery 1942). Two other **described** New World races **were** the eastern or coastal least tern (**Sterna albifrons antillarum**), and the California least tern (**Sterna albifrons browni**). The coastal least tern breeds along the Atlantic and Gulf coasts and the California least tern breeds along the California coast.

As a result of studies on vocalizations and behavior of this group of terns in the Old and New Worlds, the American Ornithologists' Union (1983) **now** treats the **New** World least terns as a distinct species, **Sterna antillarum**. Subspecies **of** New World least terns recognized by the American **Ornithologists'** Union. (1957, 1983) are the interior least tern (now **Sterna antillarum athalassos**), the **eastern** or coastal least tern (now **Sterna antillarum antillarum**), and the California least tern (now **Sterna antillarum browni**).

However, the validity of least tern subspecies has been questioned by several authors in recent years. Massey (1976) reported no consistent morphological, behavioral, or vocal differences between S. a. antillarum and S. a. browni. In Texas, where both S. a. antillarum and S. a. athalassos occur, electrophoretic analyses indicate little genetic differentiation between least terns produced on the Texas coast and Texas Panhandle rivers (McCament and Thompson 1987, McCament-Locknane 1988). Coastal least terns have populated interior breeding sites. Boyd and Thompson (1985) reported an incubating least tern at Quivira National Wildlife Refuge, Kansas, that originally had been banded as a chick on the Texas coast.

Originally, S. a. athalassos was proposed for endangered status. Because of the taxonomic uncertainty of least tern subspecies in North America, the U. S. Fish and Wildlife Service did not list the subspecies and instead designated as endangered those 'least terns occurring in interior North America. The California least tern has been listed as endangered since 1970 (U. S. Fish and Wildlife Service 1980).

Distribution

The interior least tern is migratory and historically bred along the Mississippi, Red and Rio Grande River systems and rivers of central Texas. The breeding range extended from Texas to Montana and from eastern Colorado and New Mexico to southern Indiana. It included the Red, Missouri, Arkansas, Mississippi, Ohio and Rio Grande River systems (American Ornithologists' Union 1957, Anderson 1971, Coues 1874, Burroughs 1961, Hardy 1957, Youngworth 1930, 1931, Ducey 1981). Incidental occurrences of least terns in Michigan, Minnesota, Wisconsin, Ohio and Arizona have been reported (Campbell 1935, Janssen 1986, Jung 1935, Hayfield 1943, Monson and Phillips 1981, Phillips et al. 1964).

Current Distribution

The interior least tern continues to breed in most of the aforementioned river systems, although its distribution generally is restricted to less altered river segments (Figure 1) (Tables 1-5).

Missouri River System: o r e r s , Lewis and Clark, observed the least terns along the Missouri River frequently and believed them to be "a native of this country and probably a constant resident" (Burroughs 1961). In the Dakotas, most interior least terns occur on those segments of the Missouri River and its tributaries that are not affected by impoundments or channelization. In South Dakota, the interior least tern nests primarily on flowing segments of the Missouri River and Cheyenne River (Nebraska Game and Parks Commission, Schwalbach 1988, Schwalbach et al. 1986, 1988). Breeding areas in North Dakota constitute about 192 km of the Missouri River from Garrison Dam to the mouth of the Cannonball River south of Bismarck (Dryer and Dryer 1985, Mayer and Dryer 1988), and about 29 km of the Yellowstone River in North Dakota from the Montana border to

the river's confluence with the **Missouri** River (Kreil and Dryer 1987). A few interior least terns nest on islands, shorelines and **sandbars** along the reservoir, Lake Oahe, an impoundment on-the Missouri River in North and South Dakota (Schwalbach 1988, **Mayer** and Dryer 1988). In Montana, breeding interior least terns recently have been recorded on the Yellowstone River, and on the Missouri River between **Fort Peck** Reservoir and North Dakota. A few interior least *terns* have been recorded on islands and shoreline within **the Fort Peck** Reservoir (Charles M. Russell National Wildlife Refuge). These locations are the western **most nesting** sites of the interior least tern. .

Interior least terns breed **along** the lower section of the Niobrara River, Nebraska, **from Keya Paha** and Rock Counties to the Xissouri River (Nebraska Game and Parks Commission **1985a**). Current distribution probably is similar to the historic distribution because the Niobrara River has been little changed by man (Ducey 1985). On the Platte River, Nebraska, interior least terns nest **on sandbars and** at **send** and gravel pits from the Missouri River to North Platte (Nebraska Game and Parks Commission 1987) and along the South Platte River as far west as Ogallala. On the **Loup** River, a tributary of the Platte River, interior least terns breed as far west as Arcadia but **are** most common between Saint Paul, Nebraska and the **Loup's** confluence with the Platte River at Columbus, Nebraska. A few interior least terns also occur along the **Elkhorn** River, another tributary of the Platte River.

The interior least tern no longer **nest**s in **the Missouri** reaches of the Missouri River (Smith **1985**, Sidle et al. 1988, Smith and **Renken** 1990). The hydrology of the River in **Missouri** has been drastically altered by channelization, and studies **show** that river levels are typically too high during the breeding season to expose **suitable nesting** habitat (Smith and Renken 1990).

Arkansas River System: Breeding interior least terns occur along the Arkansas River system in Colorado, Kansas, Oklahoma, **Arkansas** and Texas (Table 2). In Colorado, interior least terns nest at Adobe Creek reservoir (Blue Lake) and have been observed at Nee **Noshe** reservoir -(Carter 1989). Both reservoirs are located on small tributaries of the **Arkansas River**.

In **Kansas**, interior least terns nest on the **Cimarron** River in **Meade**, Comanche and Clark Counties, and **Quivira** National Wildlife Refuge, and in the **recent** past at Cheyenne Bottoms Wildlife **Management Area** (Boyd 1983, 1986, 1987; Schulenberg and **Ptacek** 1984).

'The interior least tern occurs on several **tributaries** of the Arkansas River in Oklahoma. It breeds along the Salt Fork of the Arkansas River at the Salt Plains **National** Wildlife Refuge (Hill 1985, Grover and Knopf 1982); Optima Reservoir at the fork of the Coldwater Creek and Beaver River in the Oklahoma Panhandle; and **on** the Cimarron River in Beaver, Harper, Woods, **Woodward, Major, Blaine**, Kingfisher, Logan, and Payne Counties (Boyd 1987, L. Hill personal communication).

Along the **Arkansas River** In Oklahoma, the interior least tern breeds in **Kay, Osage, Pawnee, Creek, Tulsa, Wagoner, Muskogee, and Sequoyah Counties** (Hoffman. 1986, L. Hill personal communication). In Arkansas, the breeding range on the Arkansas River is above Little Rock (Smith and Shepherd 1985, Smith et al. 1987, K. Smith 1986).

Along the Canadian River, interior least terns breed in Ellis, Roger Mills, **Dewey, Cleveland, McClain, Haskell,** and Sequoyah Counties, Oklahoma and in Hemphill, Roberts and Hutchinson Counties, Texas (**McCament** and Thompson 1985, 1987; U. S. Fish and Wildlife Service, unpublished data).

Mississippi and Ohio Rivers: The **Mississippi River**, interior least terns occur almost entirely in the lower valley south of Cairo, Illinois to Vicksburg, **Mississippi** (Sidle et al. 1988) (Table 3). Surveys by the U. S. Army Corps of Engineers (**Rumancik** 1985, 1986, 1987, and 1988; **M. Smith** 1986) and **Missouri** Department of Conservation (**J. Smith**.1985, 1986, 1987, and 1988, Smith and Renken 1990) indicate that about one-half of all interior least terns occur along 1100 km of the Lower Mississippi River.

On the Ohio River system, the interior least tern occurs just above the confluence of the Tennessee and Ohio Rivers and at one artificial site on the **Wabash River** in Indiana.

Red River System: Interior least terns are known to occur on the Prairie Dog Town Fork of the Red River in the eastern Texas Panhandle and along the Texas/Oklahoma boundary as far east as **Burkburnett, Texas** (**McCament** and Thompson 1985, 1987) (Table 4).

Rio Grande River System: Interior least terns occur at three reservoirs along the **Rio Grande River** and along the **Pecos River** at the **Bitter Lake National Wildlife Refuge, New Mexico** (**McCament** and Thompson 1985, 1987; Neck and Riskind 1981, Seibert 1951, **Marlatt** 1984, 1987) (Table 5).

Wintering Areas: The wintering area of interior least terns is unknown. However, least terns of unknown populations or subspecies are found during the winter along the Central American coast and the northern coast of South America from Venezuela to northeastern Brazil. Roger Boyd (personal communication 1986) reports that about 35 least terns have been recaptured in South America, mostly in **Guyana**. One interior least tern banded by Boyd, was captured in **El Salvador** two years later. Also, a banded California least tern was recaptured in Guatemala.

Table 1. Known breeding areas for interior least terns along the Missouri River system in 1985-1988.

State	County	Locations
Montana	Valley	Fort Peck Reservoir , Charles M. Russell National Wildlife Refuge
	Garfield	Fort Peck Reservoir , Charles M. Russell National Wildlife Refuge
	Prairie	Yellowstone River sandbars
	McCone Richland	Missouri River sandbars Missouri River sandbars
North Dakota	McLean	Hissouri River sandbars
	Burlrigh	Missouri River sandbars
	Oliver	Missouri River sandbars
	Horton	Missouri River sandbars
	Emmons	Lake Oahe
	Mercer	Missouri River sandbars
	sioux McKenzie	Missouri River sandbars Yellowstone River sandbars
South Dakota,	Charles Mi	Missouri River sandbars
	Bon Homme	Missouri River sandbars
	Yankton	Missouri River sandbars : .
	Clay	Missouri River sandbars
	Union	Hissouri River sandbars
	Sully	Lake Oahe
	Hughes	Lake Oahe
	Stanley	Lake Oahe
	Walworth	Lake Oahe
	Campbell	Lake Oahe.
	Corson	Lake Oahe
	Potter	Lake Oahe
	Dewey	Lake Oahe
	Ziebach Haakon	Cheyenne River sandbars Cheyenne River sandbars
Nebraska	Dixon	Missouri River sandbars
	Cedar	Missouri River sandbars
	Knox	Missouri River sandbars
	Howard	Loup River sandbars and sand/gravel pits
	Nance	Loup River sandbars and sand/gravel pits
	Sherman	Loup River sandbars and sand/gravel pits
	Platte	Loup River sandbars and sand/gravel pits
	Valley	Loup River sandbars and sand/gravel pits
	Douglas	Elkhorn River sandbars and sand/gravel pits
	Cumming	Elkhorn River sandbars and sand/gravel pits
	Stanton	Elkhorn River sandbars and sand/gravel pits
Boyd	Niobrara River sandbars	

Holt	Niobrara River sandbars
Keya Paha	Niobrara River sandbars
Brown	Niobrara River sandbars
Knox	Niobrara River ● 8ndb8re
Rock	Niobrara ☼✠♣□ ● 8ndb8re
Cars	Platte River eendbars 8nd sand/gravel pits
Sarpy	Platte River ● mdbere and sand/gravel pits
Saunders	Platte River eendbare 8nd sand/gravel pits
Douglas	Platte River eendbare 8nd sand/gravel pits
Dodge	Platte River eandbare end sand/gravel pits
Colfax	Platte River ● 8ndb8re ♣■Ω ● 8nd/gravel pits
Butler	☼●☼♣♣ ☼✠♣□ ● mdb8rs and sand/gravel pits
Platte	River sandbars 8nd send/gravel pits
Polk	Platte River sandbars 8nd sand/gravel pits
Hall	Platte River sandbars and sand/gravel pits
Buffalo	Platte River oandbars 8nd ● 8nd/gravel pits
Kearney	Platte River sandbars 8nd sand/gravel pits
Phe lps	Platte River sandbars 8nd sand/gravel pits
Dawson	Platte River ● andbare and ● and/gravel pits
Hamilton	Platte River sandbars end sand/gravel pits
Merrick	Platte River eandbare 8nd sand/gravel pits
Lincoln	Platte River sandbars ● ud sand/gravel pits
Lincoln	So. Platte River sandbars/sand/gravel pits
Keith	So. Platte River sandbars/sand/gravel pits

Iowa'

Woodbury Iowa Public **Service ash** ponds
Pottawattamie Iowa Power **and Light ash** ponds




Table 2. Known breeding areas for interior least terns along the Mississippi and Ohio Rivers, 1985-1988.

State	County or Parish	Location
Missouri	Pemiscott	Mississippi River sandbars 8nd dike fields
	New Madrid	Mississippi River ● ndb8rs and dike fields
	Mississippi	Mississippi River sandbars 8nd dike fields
	Scott	Mssissippi River sandbars and dike fields
Kentucky	Fulton	Mississippi River sandbars 8nd dike fields
	Hickman	Mssissippi River sandbars 8nd dike fields
	Carlisle	Mississippi River sandbars 8nd dike fields
Tennessee	Dyer	Mississippi River sandbars and dike fields
	Lake	Mississippi River sandbars 8nd dike fields
	Lauderdale	Mississippi River sandbars 8nd dike. fields
	Tipton	Mississippi River sandbars and dike fields
	Shelby	Mississippi River sandbars and dike fields
Arkansas	Mississippi	Mississippi River sandbars 8nd dike fields
	Crittenden	Mississippi River sandbars 8nd dike fields
	Lee	Mississippi River sandbars and dike fields
	Phillips	Mississippi River sandbars 8nd dike fields
	Desk8	Mississippi River sandbars and dike fields
	Chicot	Mississippi River sandbars 8nd dike fields
Mississippi	Deioto	Mississippi River sandbars 8nd dike fields
	Tunica	Mssissippi River sandbars and dike fields
	Coahoma	Mississippi River sandbars and dike fields
	Bolivar	Mississippi River sandbars 8nd dike fields
	Washington	Mississippi River sandbars 8nd dike fields
	Issaguena	Mississippi River sandbars 8nd dike fields
	Warren	Mississippi River sandbars and dike fields
Louisiana	East Carroll	Mississippi River sandbars and dike fields
	Madison	Mississippi River sandbars and dike fields
Illinois	Alexander	Mississippi River sandbars and dike fields
	Pulaski	Ohio River ● ndb8rs and dike fields
Indiana	Gibson	Public Power plant along Wabash River at East Mt. Carmel

Table 3. Known breeding areas for interior least terns along the Arkansas River system, 1985-1988.

State	county	Location
Arkansas	Pulaski	Arkansas River sandbars and dike fields
	Faulkner	Arkansas River sandbars and diks fields
	Conway	Arkansas River sandbars and dike fields
	Perry	Arkansas River sandbars and dike fields
	Pope	Arkansas River sandbars and dike fields
	Logan	Arkansas River sandbars and dike fields
	Johnson	Arkansas River sandbars and dike fields
	Sabastian	Arkansas River sandbars and dike fields
	Crawford	Arkansas River sandbars and dike fields
Oklahoma	O o a g e	Arkansas River sandbars
	Kay	Arkansas River sandbars
	Pawnee	Arkansas River sandbars
	Creek	Arkansas River sandbars
	Tulsa	Arkansas River sandbars
	Wagoner	Arkansas River sandbars
	Muskogee	Arkansas River sandbars
	Beaver	Cimarron River sandbars
	Harper	Cimarron River sandbars
	woods	Cimarron River sandbars
	Woodward	Cimarron River sandbars
	Major	Cimarron River sandbars
	Blaine	Cimarron River sandbars
	Kingfisher	Cimarron River sandbars
	Logan	Cimarron River sandbars
	Payne	Cimarron River sandbars
	Alfalpa	Salt Plains National Wildlife Refuge
	Texas	Optima Reservoir
	Ellis	Canadian River sandbars
	Roger Hills	Canadian River sandbars
Dewey	Canadian River sandbars	
Haskell	Sequoyah National Wildlife Refuge	
Sequoyah	Sequoyah National Wildlife Refuge	
Cleveland	Canadian River sandbars	
McClain	Canadian River sandbars	
Texas	Hemphill	Canadian River sandbars
	Roberts	Canadian River sandbars
	Hutchinson	Canadian River sandbars

Kansas	Comanche	Cimarron River sandbars
	Clark	Cimarron River sandbars
	Meade	Cimarron River sandbars
	Stafford	Quivira National Wildlife Refuge
Colorado	Kiowa	A&be Creek Reservoir
		Nee Noshe Reservoir
	Bent	Adobe Creek Reservoir

Table 4. **Known** breeding areas for interior least terns along the Red River system, 1985-1988.

State	County	Location
Texas	Childrest	Prairie Dog Town Fork sandbars
	Hall	Prairie Dog Town Fork sandbars
	Briscoe	Prairie Dog Town Fork sandbars

Table 5. **Known** breeding areas for interior least terns along the Rio Grande system, 1985-1988.

State	County	Location
Texas	Zapata	Falcon Reservoir
	Webb	the Casa Blanca
	Val Verde	Amistad Reservoir
New Mexico	Chaves	Bitter Lake National Wildlife Refuge




Fig. 1 Distribution of the interior least tern.

Life History

Breeding Behavior: Interior least terns spend about 4-5 months at their breeding sites. They arrive at breeding areas from late April to early June (Foanes 1983, Hardy 1957, U. S. Fish and Wildlife Service 1987a, Wilson 1984, Wycoff 1960, Youngworth 1930). Courtship behavior of least terns is similar throughout North America. Courtship occurs at the nesting site or at some distance from the nest site (Tomkins 1959). It includes the fish flight, an aerial display involving pursuit and maneuvers culminating in a fish transfer on the ground between two displaying birds. Other courtship behaviors include nest scraping, copulation and a variety of postures, and vocalizations (Ducey 1981, Hardy 1957, Wolk 1974).

The nest is a shallow and inconspicuous depression in an open, sandy area, gravelly patch, or exposed flat. Small stones, twigs, pieces of wood and debris usually lie near the nest. Least terns nest in colonies or ternerits, and nests can be as close as just a few meters apart or widely scattered up to hundreds of meters (Ducey 1988, Anderson 1983, Hardy 1957, Kirsch 1990, Smith and Renken 1990, Stiles 1939). The benefit of semi-colonial nesting in least terns may be related to anti-predator behavior and social facilitation (Burger 1988).

Interior least tern eggs are pale to olive buff and speckled or streaked with dark purplish-brown, chocolate, or blue-grey markings (Hardy 1957, Whitman 1988). Occasionally, eggs are pink instead of pale to olive buff (P. Mayer and M. Schwalbach, personal communication). The birds usually lay two or three eggs (Anderson 1983, Faanes 1983, Hardy 1957, Kirsch 1987-89, Sweet 1985, Smith 1985). The average clutch size for interior least terns nesting in the Mississippi River during 1986-1989 was 2.4 eggs (Smith and Renken 1990). Egg-laying begins by late May. Both sexes have short incubation which generally lasts 20-25 days but has ranged from 17 to 28 days (Faanes 1983, Hardy 1957, Moser 1940, Schwalbach 1988, G.R. Lingle, personal communication).

The precocial behavior of interior least tern chicks is similar to that of other least terns. They hatch within one day of each other, are brooded for about one week, and usually remain within the nesting territory but as they mature, wander further. Fledging occurs after three weeks, although parental attention continues until migration (Hardy 1957, Massey 1972, 1974; Tomkins 1959). Departure from colonies by both adults and fledglings varies but is usually complete by early September (Bent 1921, Hardy 1957, Stiles 1939). Thompson (1982) presented the following longevity data for coastal least terns revealed by band recoveries:

Percentage of Recoveries	
<u>Age (years)</u>	<u>Known and Assumed Dead (N)</u>
0-5	74 percent (58)
5-10	9 percent (7)
10-15	10 percent (8)
15-20	4 percent (3)
>20	3 percent

Population Biology: The interior least tern's annual reproductive success varies greatly along a given rivtr or shoreline (Table 6). Because tern's use tptmerol habitats, they art susceptible to frequent nest and chick loss. Consequently there are great local differences in productivity. In 1987, total number of interior least terns reached 4,800 (Table 7). This is considerably higher than the 1,200 interior least terns estimated by a partial survey in 1975 by Downing (1980). There are no comprehensive historic numbers to compare with these figures, although early qualitative descriptions indicate that the interior least tern was rather common (Burroughs 1961, Hardy 1957). Increased censusing efforts during the past few years probably account for the differences among recent census figures and earlier surveys.

Table 6. Some txamplts of the productivity of interior least terns.

Locations	Year	Nest Success	Flidgings per Pair	Frequency of Visits	% Population Monitored	Source
Missouri River	1988	0.62	0.42	7-10 days	100%	Mayer and Dryer 1989
	1989	0.56	0.21	"	"	
North Dakota						
Missouri Rivtr	1986		0.20	7-10 days	100%	Schwalbach 1988
	1987		0.64	"	"	
South Dakota						
Missouri River	1988	0.36	0.44	7-10 days	100%	Dirks 1990
	1989	0.51	0.55	"	"	
South Dakota						
Lower Platte River	1987	0.57	0.29	2-3 days	39%	Kirsch 1987-89
	1988	0.67	0.71	"	44%	
	1989	0.43	0.47	"	42%	
Nebraska						
Cimarron River	1982-83	0.18	1.09-0.56	--	--	Schulenberg and Ptactk 1984
Kansas						
Salt Plains NWR, Oklahoma	1987	0.44-0.33	0.44-0.15	1-3 days		Hill 1987

Dispersal Patterns: Breeding site fidelity of coastal and California least terns is very high (Atwood et al. 1984, Burger 1984). This may also be true for the interior least tern in its riverine environment. An interior least tern banded in 1988 as a breeding adult on the Missouri River in North Dakota returned in 1989 to breed on a Missouri River sandbar in North Dakota (Mayer and Dryer 1990). In the Mississippi River valley, a bird banded as a breeding adult in 1987 was observed nesting at the same site in 1989, and three others banded as breeding adults in 1988 returned to nest within the same stretch of the Mississippi River in 1989 (Smith and Rtnktn 1990). Two of those birds had returned to within 4.8 km of their former nesting site. Along the Platte River in Nebraska, interior least terns demonstrate a strong return pattern to previous nesting sites on the river and at sand and gravel pits regardless of reproductive success (E. Kirsch, C. Lingt, personal communication). One interior least tern captured in 1987 as a breeding adult at a Mississippi River ternery in Missouri had been banded as a chick in 1980 by Marsha Ualdron; this bird was nesting at a site 131 km upriver from its natal Ttnntsstt colony (Smith 1987, Smith and Renken 1990). Chick dispersal may be as far as that reported by Boyd and Thompson (1985) for a breeding Kansas bird that had been banded as a chick on the Texas coast.

Home Range and Territoriality: The interior least tern's home range during the breeding season usually is limited to a stretch of river near the sandbar nesting site. At Salt Plains National Wildlife Refuge, home ranges were highly variable, ranging from 11 to 1,015 ha (Talent and Hill 1985). Variation likely was due to food limitations and chick loss. The home range may change if re-nesting birds select a different breeding site. At sand and gravel pits along the central Platte River in Nebraska, nesting interior least terns utilize the pit or stretches of river as an adjacent stretch of river. Nesting territories are defended and birds defend any nest in the colony. In defending the territory, the incubating bird will fly up and give an obvious alarm call followed by repeated dives of the intruder (Hardy 1957). The strong defense of territories facilitates locating terneries during census surveys.



Table 7. Census data on the interior population of the least tern, 1985-1988'.

Location	Number of adult least terns				Approximate length of river stretch (km) where nesting least. terns intermittently occur	Source
	1985	1986	1987	1988		
Mississippi River Basin						
1. Ft. Peck Reservoir, Missouri River, Montana	-*	--**	4	2		(Alfonso, unpublished data, Montana Piping Plover Recovery Committee 1988)
2. Below Ft. Peck Reservoir, Missouri River, Montana		--	--	18	22	(D. Christopherson, unpublished data)'
3. Yellowstons River, Hontana	-	--	--	12	-	(Gorges, unpublished data)
4. Below Garrison Dam, Missouri River, North Dakota	114	169	175	142	192	(Dryer and Dryer 1985, Hayer and Dryer 1988)
5. Lake Sakakawea, Missouri River, North Dakota	-	-	-	7	-	(flayer and Dryer 1988)
6. Lake Oahe, Missouri River	-	-	-	7	-	(Hayer and Dryer 1988)
7. Yellowstone River, North Dakota		22	20	24	30	(Kreil and Dryer 1987, Hayer and Dryer 1988)
8. Cheyenne River, South Dakota	-	31	54	27	26	(Schwalbach et al. 1986, 1988; Schwalbach 1988)

Table 7 (continued)

location	Number of adult least terns				Approximate length of river stretch (km) where nesting least terns intermittently occur	Source
	1985	1986	1987	1988		
9. Lake Oahe. Missouri River, South Dakota	-	16	21	61		(Schwalbach et al. 1986, 1988; Schwalbach 1988)
10. Below Fort Randall and Gavins Point Dam, HO River, South Dakota to Ponca, NE	202	206	292	297	140	(Schwalbach et al. 1986, 1988; Schwalbach 1988)
11. Power plant ash lagoons near Council Bluffs, Iowa	18	28	22	22		(Dinsmore and Dinamore 1989, Wilson 1984)
12. Niobrara R., Nebraska	174	-	143	200	190	(Nebraska Game and Parks Commission 1985a)
13. Platte River, Nebraska	256	438	606	635	5 0 2	(Nebraska Game and Parks Commission 1988; G. R. Lingle , personal communication)
14. Loup River, Nebraska	-	-	100	155	70	(S. Gauthreaux and Nebraska Game and Parks Commission , unpublished data)
15. Elkhorn River, Nebraska	2	8	4	-	-	(J. Dinan , Nebraska Game and Parks Commission , personal communication)
16. Mississippi R., Cape Girardeau, Missouri to Vicksburg, Mississippi	1264	2244	2488	2356	1100	(Rumancik 1985, 1986; J.W. Smith 1985, 1986, 1987, 1988; M. Smith 1986; W. King personal communication; Smith and Renken 1990)
17. Power plant, Wabash River, E. Ht. Camel, IN	2	4	4	-	-	(Johnson 1987, Hills 1987)

Table 7 (continued)

Location	Number of adult least terns				Approximate length of river stretch (km) where nesting least terns intermittently occur	Source
	1985	1986	1987	1988		
18. Arkansas River, Arkansas (above Little Rock)	50	80	130	119	256	(Smith and Shepherd 1985, K. Smith 1986, Smith et al. 1987)
19. Arkansas River, Oklahoma	-	78	200	200	119	(Hoffman 1986, L. Hill personal communication)
20. Quivira National Wildlife Refuge, Kansas (Rattlesnake Creek of Arkansas River)	48	48	54	-	-	(Boyd 1986, 1987)
21. Adobe Creek Reservoir Colorado	-	-	6	10	-	(Barbara Campbell, personal communication)
22. Salt Plains National Wildlife Refuge, Oklahoma (Salt Fork of the Arkansas River)	-	140	210	-	-	(Boyd 1986, 1987)
23. Cinarron River, Kansas and Oklahoma	82	150	132	-	121	(Boyd 1986, 1987)
24. Optima Reservoir , Oklahoma (Beaver River)	46	52	60	38	-	(Boyd 1986, 1987; L. Hill)
25. Canadian River, western Oklahoma and Texas	127	182	20	16	2 5 3	(McCament and Thompson 1985, 1987; U. S. Fish and Wildlife Service, unpublished data)

Table 7 (continued)

Location	Number of adult least terns				Approximate length of river stretch (km) where nesting least terns intermittently occur	Source
	1985	1986	1987	1988		
26. Canadian River, Eufaula Dam to Arkansas River, including Sequoyah National Wildlife Refuge	-	-	105	34	43	(L. Hill personal communication)
27. Canadian River at Norman, Oklahoma	-	-	-	12	3	(L. Hill, personal communication)
28. Prairie Dog Town Fork of Red River, Texas	44	50	12	16	241	(McCament and Thompson 1985, 1987; B. Thompson, pers. commun.)
Rio Grande River Basin						
29. Falcon Reservoir, Rio Grands River	500	150	50	222	-	(McCament and Thompson 1985, 1987; B. Thompson, pers. commun.)
30. Lake Casa Blanca	5	-	14	50	-	(McCament and Thompson 1985, 1987; B. Thompson, pers. commun.)
31. Amistad Reservoir, Rio Grande River	20	9	-	14	-	(McCament and Thompson 1985, 1987; B. Thompson, pers. commun.)

Table 7 (continued)

Location	Number of adult least terns				Approximate length of river stretch (km) where nesting least terns intermittently occur	Source
	'1985	1986	1987	1988		
32. Bitter Lake National Wildlife Refuge, New Mexico (Pecos River)	-	8	6	6	-	(Shomo, 1988 and S. Williams, New Mexico Game and Fish Department, unpublished report)
Total	2952	4113	4932	4702	3308	

The census results should be viewed in light of the extent **and** frequency of census efforts. Increases or decreases from year to year may not be related to reproductive performance.

* no census conducted in that year.

** area surveyed but no birds found

Diet: The interior least tern is *is* plscivorous, **feeding** in shallow waters of rivers, streams and lakes. Other least terns also feed on crustaceans, insects, mollusks and **annelids** (Whitman 1988). The terns usually feed close to their nesting sites. Fish prey is small sized and important genera include **Fundulus, Notropis, Campostoma, Pimephales, Gambusia, Blonesox, Morone, Dorosoma, Lepomis** and **Carpiodes** (Grover 1979, Hardy 1957, Rumancik 1988, 1989; Schulenberg et al. 1980, Smith and Renken 1990, Wilson et al. 1989). Moseley (1976) believed least terns to be opportunistic feeders, exploiting any fish **within** a certain size range. Fishing occurs close to the riverine colony. Terns nesting at sand and gravel pits and other artificial habitats may fly up to 3.2 km to fish. Radio-tagged terns at Salt Plains National Wildlife Refuge often traveled **3.2-6.4** km to fish (Talent and Hill 1985). Fishing behavior involves hovering and diving over **standing or flowing water**.

Interspecific Interactions: Interior least terns are breeding associates of the piping plover (**Charadrius melodus**) in the Missouri River system (Dryer and Dryer 1985, Faanes 1983, Nebraska Game and Parks Commission 1987, Schvalbach 1988) and the snowy plover (**Charadrius alexandrius**) and American avocet (**Recurvirostra americana**) in the Arkansas River system (Grover and Knopf 1982, Hill 1985). Nesting piping plovers usually can be found within or near nesting interior least terns at sand and gravel pits and on riverine sandbars.

Habitat Requirements

Least terns throughout North America **nest** in areas with similar habitat attributes.

Coastal Areas: Coastal and California least terns usually nest on elevated portions of level, unvegetated substrates near foraging areas (Carreker 1985). **Beaches, sand pits, sandbars, islands** and peninsulas are the principal breeding habitats (Moseley 1976). Nesting **can be** close to water but is usually between the dune environment and the high tide line (Akers 1975, Blodget 1978). Unconsolidated substrate such **as** small stones, gravel, sand, debris and shells comprise the nesting substrate. A mixture of coarse sand, shells and other fragments may be preferred over fine-grained substrates **because** of better cryptic qualities, stability in wind, and water permeability (Burroughs 1966, Craig 1971, **Gochfeld 1983, Jernigan** et al. 1978, Soots and Parnell 1975, Svickard 1972, Thompson and Slack 1982).

Vegetation at California and coastal least tern nesting sites **is** sparse, scattered and short. Vegetation cover is usually less than 20% at the time of nesting (Craig 1971, Thompson and Slack 1982, Gochfeld 1983). Least tern colonies in denser vegetation may be a response to habitat loss or a function of strong site tenacity.

Rivers: The riverine nesting areas of interior least terns are sparsely vegetated sand and gravel bars within a **vi** & unobstructed river channel, or salt flats along lake shorelines. Nesting locations usually

are at the higher elevations and away *from* the water's edge because nesting starts when the river **flows are** high and small amounts *of* sand are exposed. The size of *nesting* areas depends on water levels and the extent of associated sandbars. **An** examination of the interior least **tern's** nesting ecology on the Missouri River (Schwalbach et al. 1988) illustrates the changes caused by varying river flows. Along one stretch of the Missouri River in South Dakota the average size of nesting sandbars was **12** and **31** ha in 1986 and 1987, respectively; nest elevation and nest to water distance differed by a factor of three in both years.

The Lower **Mississippi** River is very **wide** and carries a tremendous volume of water and sand. Sandbars form annually, are washed **away**, and shift **position**. **Many** sandbars are over **3.2 km** long and **1.2 km** wide. Nest sites are often several hundred meters *from* the water (**Rumancik** 1987, 1988). Thus, nesting areas **usually are** several hundred **hectares in** size. Mississippi River levels at the onset of nesting also influences the number *of* nests at a colony. **Smith and Renken** (1990) observed Mississippi River colonies that averaged 100 nests/colony when habitat was restricted by high water early in the nesting period, but which averaged only 19.3 nests/colony during a year *of* more moderate river **levels**.

Artificial Nesting Habitat: Least terns nest on artificial habitats such as sand and gravel pits and dredge islands (Dryer and Dryer 1985, **Haddon** and Knight 1983, Kirsch 1987-89, **Larkins** 1984, **Morris** 1980). In North America the coastal and California least terns commonly nest on a variety *of* artificial nesting habitats, even roof -tops (**Altman** and Cano 1984, Atwood' *et al.* 1979, Fisk 1975, 1978; **Jernigan** 1977, **Massey** and Atwood **1980, 1983**; Swickard 1974).

The interior least tern nests on dike **fields** along the **Mississippi** River (Smith and **Stucky** 1988; Smith and Renken 1990). at sand and gravel pits (Kirsch **1987-89**), ash disposal areas *of* power plants (**Dinsmore and Dinsmore** 1988, Johnson 1987, Wilson **1984**), along the shores of reservoirs (Boyd 1987, Chase and **Loeffler** 1978, Neck and **Riskind** 1981, Schwalbach 1988) and at other manmade sites (**Shomo** 1988). The percentage of interior least terns nesting on pits adjacent to the lower reach (Columbus -to **Plattsmouth**) *of* the Platte River varies depending on the flow and amount *of* exposed sandbar habitat (Kirsch 1987-89). Suitable nesting habitat in the upper Platte River channel has been severely reduced (Sidle et al. 1989) and in many stretches of the river, sand and gravel pits annually provide the only nesting habitat (Lingle 1989). It is unknown to what extent sand and gravel pits, dike fields, **reservoir** shorelines and other artificial habitats have replaced natural habitat. In the lower **Mississippi** River **alone**, 7,518 ha *of* bar and island habitat were lost in diked reaches between 1962 and 1976 (**Nunnally** and Beverly 1986, **Smith** and **Stucky** 1988).

Reasons For Current Status

Habitat alteration and destruction: Channelization, irrigation, and the construction of reservoirs and pools have contributed to the elimination of much of the tern's sandbar nesting habitat in the Missouri, Arkansas, and Red River systems (Punk and Robinsbn 1974, Hallbtr et al 1979, Sandhtinrich and Atchison 1986). **Ducey (1985)**, for example, describes the changes in the channel characteristics of the Missouri River since the early 1900s under the Missouri River **Bank Stabilization** and Navigation Project. The wide and braided character of the **Missouri River** was engineered into a single narrow navigation channel. Most sandbars virtually disappeared between Sioux City, Iowa and Saint Louis, Missouri (Sandhtinrich and Atchison 1986, Smith and **Stucky** 1988).

Where sandbars still occur along the Nebraska-South Dakota boundary (Missouri River), approximately 3,156 ha of sandbar habitat have been lost between 1956 and 1975 (Schmulbach et al. 1981). Sandbars along the **Nebraska-Iowa** Missouri River boundary have been virtually eliminated with the exception of 890 ha inventoried along the **80-km** Missouri National Recreation Area (Schmulbach et al. 1981).

Current regulation of Missouri River dam discharges post additional problems for interior least terns nesting in remaining habitats (Nebraska Game and Parks. Commission **1985c**, Schvalbach et al. 1988). Before regulation of river flows, summer flov patterns vtre more predictable. Peak flov occurred in **March** from local runoff and then again in **May** and **June** vhtn mountain snowmelt occurs. Flows then declined during the rest of the summer allowing interior least terns to nest as water levels dropped and sandbars became available (Stiles 1939, 'Hardy 1957). Currently, the main stem system is supposed to be regulated for **hydropower, navigation, water quality and supply, flood evacuation, irrigation, fish and wildlife conservation, and public recreation.** However, system releases are designed to provide equitable service to power and navigation demands, except vhtn they conflict with flood control functions of the system.

The demands are unpredictable and flows can fluctuate greatly. Flow regimes differ greatly from historic regimes. High flov periods may nov extend into the normal nesting period, thereby reducing the quality of existing nest sites and forcing fntrrior least terns to initiate nests in poor quality locations. Extreme fluctuations can flood existing nests, inundate potential nesting areas, or dtwater feeding areas. Interior least terns along the Arkansas River in Oklahoma and Arkansas contend with dam discharge problems similar to those on the Missouri River.

Along the **Lower Mississippi** River, and elaewhert, natural river discharge may exert considerable influence on reproductive success. A wet spring may delay river fall and habitat may not be available until later. Rises in the river during the spring and summer may inundate nests and vash away chicks (Rumancik 1986, 1989, Smith and Renktn 1990). **Renesting**, however, does occur and may be an adaptation to river fluctuations. Dike

construction has created many **sandbars** between the dikes and many nesting colonies are located on these sandbars (**Landin et al.** 1985, **Rumancik** 1986, 1987, 1988, 1989; **J. Smith** 1985, 1986, 1987). **The extent** to which these sandbars **are** attaching to the **riverbank** and reducing **tern** habitat is not known but according to Smith and **Stucky** (1988) the processes of dike field terrestrialization are well underway at several least tern colony sites in the lower **Mississippi** River.

Reservoir storage of flows responsible for scouring sandbars has resulted in the encroachment of **vegetation** along many rivers such as the Platte River, Nebraska and greatly reduced channel width (Currier et al. 1985, **O'Brien** and Currier 1987, **Eschner et al.** 1981, Lyons and Randle 1988, Sidle et al. 1989, **Stinnett et al.** 1987). In addition, river main stem reservoirs now trap much of the sediment load resulting in less aggradation and more degradation of the river bed and subsequently less formation of suitable sandbar nesting habitat. Riverine habitat along the central Platte River may require extensive vegetation clearing and other intensive management. In contrast, the lower Platte River (Columbus, Nebraska to the Missouri **River** confluence) has not undergone **as** extensive habitat changes as the central Platte. During 1987-1989, riverine sandbar habitat hosted 72% of the nests on the lower Platte and only 12% of the nests on the central Platte (**Kirsch** 1989, Linglt 1989).

Human Disturbance's have become the focus of recreational activities. Human presence reduces reproductive success (Mayer and Dryer 1988, Smith and Renken 1990). In **mid-America**, **sandbars** are **fast** becoming the recreational counterpart of coastal beaches. **Even sand** and gravel pits and other artificial nesting sites receive a high level of human disturbance.

Conservation Efforts

During the past few years there **has been a great** increase in the number of interior least tern surveys, research projects' and public relations endeavors to protect the birds **on the part of both public and** private conservation organizations. Proposed federal listing of the interior least tern prompted much of the interest **in** the northern Great Plains and elsewhere. **Today**, many state, federal **and** private organizations **are** collaborating to census the birds, curtail human disturbance and conduct research.

Under authority of Section 7 of the Endangered Species Act, the U. S. Fish and Wildlife Service is consulting with the U. S. Army Corps of Engineers on whether dam operations on the Missouri and Arkansas Rivers jeopardize the continued existence of the interior least tern (**U.S** Fish and Wildlife Service 1989, 1990). **The** outcome of these formal consultations is crucial to the recovery of the interior least tern. Areas of habitat along the Missouri River, for example, continue to degrade due to physical controls on the river **and** present water management schemes. Changes in **the** water release regime and physical manipulation of habitat will be necessary.

Aside from the Section 7 consultation on the **Missouri River**, the Corps Master **Manual** for river operations **is** under **review**. If upper Missouri River Basin states have their way *for* holding **water in** the reservoirs for recreation and fisheries, navigation in the **Missouri River** could be reduced and maintenance *of* the commercial navigation project above Omaha could become infeasible. The reach between Sioux City, **Iowa** and the mouth of the Platte River could'once more be available to interior least terns.

Montana **efforts** include surveys *to* determine the number and distribution of interior least terns **along the Missouri** and Yellowstone Rivers and along the shores of the Fort Peck **Reservoir**.

North Dakota: Censuring has been conducted along the **Missouri River** since 1982 and along the **Yellowstone River** since 1986. Habitat requirements are being estimated and recommendations are being made for the management *of* Missouri River habitat. Research continues on reproductive success and on methods *to* increase productivity. Resource agencies **are** involved with a variety *of* public relations efforts to curtail human disturbance on Missouri River sandbars and islands.

South Dakota: Detailed studies *of* interior least tern nesting ecology continue at **Missouri** and Cheyenne River sandbars and along the reservoir shoreline of Lake Oahe. **Resource** agencies *are* involved with public relations efforts to curtail human disturbance on the **Missouri River**. Management **activities** include the **posting** of nesting sites and **informational** signs at boat ramps and elsewhere. This has been complemented with enforcement actions being taken by state and federal officials. **Recent amendments** to South Dakota **law** prohibit the harassment of least **tern nesting** and **rearing sites** on the Missouri River.

Nebraska supports one *of* the largest breeding populations of interior least terns. Annual surveys have been carried **out since** 1979. Efforts **are underway** to quantify available nesting habitat on the Platte River at various river flows. Research on reproductive success, habitat **selection**, foraging ecology, predation and the value of sand and gravel pits continues along the Platte River (**Kirsch** 1987-89, **Lingle** 1989, **Wilson et al.** 1989).

Aflow management plan has been prepared for the **Missouri River** (Nebraska Game and Parks Commission 198%) **and certain instream flovs** have been determined on the Platte River for the interior least **tern**, its habitat and forage fish, and for other vildlife and resources (Table 8). In 1990 the Federal Energy Regulatory Commission (FERC) ordered the Nebraska Public Pover District to maintain the **instream** flovs In Table 8 for interior least terns (SO FERC Report (CM) 61,180) (Sidle et al. 1990). The District seeks a new license to operate diversion dams and other facilities associated with the **Lake McConaughy** reservoir on the North Platte River. **Lake McConaughy was** constructed in the late 1930s and licensed for 50 years. The dam, diversion structures, and other facilities have had a major impact on the downstream habitat of the

interior least tern. **When** granting a new power license the Federal Power Act requires **FERC** to give equal consideration to the protection, mitigation of damage to, and enhancement of, fish and wildlife.

Posting, extensive news media efforts, posters, brochures, information signs at river **entry** points, and law enforcement **patrols** are **some** of the additional activities being carried **out** in Nebraska. The Platte River Whooping Crane Habitat Trust is trying to rehabilitate sandbars in the central Platte River (Lexington to Grand Island) by removing vegetation over extensive **areas** of the river channel. **FERC** also ordered the Nebraska Public Power District to construct eight permanent five- to ten-acre sites for interior least tern nesting in the central Platte River where nesting habitat has been severely degraded, **in** part by the upstream Lake **McConaughy** and associated water diversion canals and offstream reservoirs.

Finally, Nebraska law requires state agencies to consult with the Nebraska Game and Parks Commission on any action authorized, funded, or carried out by the state agencies. **This** insures that such actions do not jeopardize the continued existence of endangered **or** threatened species or result in the destruction or adverse modification *of* habitat. The Commission reviews state sponsored or authorized projects that may impact endangered **or** threatened species and issues biological opinions to the state agencies.

Colorado: Interior least tern is known to breed **at** Adobe Creek reservoir and has been observed at Nee **Noshe** reservoir. Public relation efforts and other endeavors are underway to address fluctuating water levels, **human** disturbance, vegetation encroachment, and predation.

Iowa: Largely devoid *of* natural interior least tern habitat, Iowa's conservation efforts have focused on monitoring and protecting the few nest sites located on fly-ash disposal sites of two power generating stations along the Missouri *River* at Council Bluffs and Sioux City. Both sites are monitored to record the number of nesting pairs and reproductive success. The Council Bluffs nesting habitat also is protected by a management plan. The plan specifies 'that both people and heavy equipment will be kept out of the nesting area during the breeding season.


Interior least tern decoys have been set out at the **DeSoto** National Wildlife Refuge to attract terns which formerly nested *there* in the 1970s. Woody vegetation has been cleared and the **•** raas are **disked** to maintain open habitat.

Table 8. Recommended annual flow regime for Central Platte River, Nebraska¹

<u>Time Period</u>	<u>Flow(cfs²)</u>	<u>Species/Resources of Concern</u>	<u>Existing Median Flow(cfs) (1958-1985)</u>
Jan 1-Mar 22	1,100	Bald Eagle, yet meadow sandhill crane; waterfowl, least tern forage fish, sport fish	1,710
Mar 23-May 10	2,000	Whooping crane, sandhill crane, waterfowl, least tern forage fish, sport fish	1,823
May 11-May 14	800	Least tern forage fish, sport fish	1,433
May 15-Sep 15	800	Least tern, piping plover, tern forage fish, sport fish	781
Sep 16-Nov 15	2,000	Whooping crane, sandhill crane, waterfowl, least tern forage fish, sport fish	893
Nov 16-Dec 9	1,000	Waterfowl, least tern forage fish, sport fish	1,186
Dec 10- Dec 31	1,100	Bald eagle,, waterfowl, least tern forage fish, sport fish	1,253

¹As measured at the U. S. Geological Survey gage at Grand Island.

²Cubic feet per second



Missouri Department of Conservation maintains an active conservation, management and research program for interior least terns. The **Missouri** River has been thoroughly surveyed for potential habitat; **Mississippi** River colonies are closely monitored and under detailed study; and management plans have been developed. Regulations provide special protective status for least tern nesting areas on Department owned islands and sandbars. Public information programs about the interior least tern are widespread.

Kansas: The **Kansas** Department of Wildlife and Parks has funded research on distribution, reproductive success, banding and inter-colonial movements, foraging ecology, and predation since 1980. Annual surveys along the **Cimarron** River and at the **Quivira** National Wildlife Refuge have been conducted since 1980. Successful habitat alteration and management has been on-going since 1985. Studies also have focused on the issue of inadequate **instream** flows in both the **Cimarron** and **Arkansas** rivers in **Kansas**.

Oklahoma: The largest concentration of least terns in Oklahoma is at **Salt Plains** National Wildlife Refuge. This area has been studied **intermittently** since 1977. Research at river nesting sites has been on-going since 1982. The **Cimarron** and **Arkansas** rivers have received more survey and distribution effort than the **Red** and **Canadian** rivers. Various studies of reproductive success, **inter-colonial** movements and foraging ecology have been conducted at **Salt Plains**, **Optima** Reservoir and the western reaches of the **Cimarron** River. Posting, fencing and extensive news media efforts have been successful at **Optima** Reservoir and the western reaches of the **Cimarron** River. Nesting sites on the **Cimarron** River continue to be threatened by several river diversion and impoundment proposals. A memorandum of understanding has been developed between The Nature Conservancy, U. S. Army Corps of Engineers, Oklahoma Department of Wildlife Conservation, U. S. Fish and Wildlife Service, Tulsa Audubon Society, River Parks Authority and riverbed landowners for protection and management of essential habitat on the **Arkansas** River in **Tulsa** County.

Mississippi River States: The U. S. Army Corps of Engineers has undertaken extensive census work along the **Mississippi** River between **Illinois** and **Vicksburg**, **Mississippi**, and along the **Arkansas** River to the **Oklahoma** border. Their surveys have provided the only information on the tern on the **Mississippi** River below the State of **Missouri**. The locations of colonies are monitored and the information is used by regulatory personnel to evaluate permit applications and in planning operations and maintenance activities on the lower **Mississippi** River.

Texas and New Mexico: The **Texas** Parks and Wildlife Department has examined the numbers and distribution of interior least terns along the **Rio Grande** River and rivers in the **Texas** Panhandle, and investigated genetic characteristics of coastal and interior least terns. The **New Mexico** Department of Game and Fish has conducted several years of surveys and studies and developed management recommendations for interior least terns at and near the **Bitter Lake** National Wildlife Refuge along **Pecos** River (**Jungemann** 1988).

II. RECOVERY

Recovery objective

The purpose of this plan is to describe actions necessary to achieve recovery of **interior least terns**. The first **step** in this approach is to set a quantifiable goal (**i. e.**, recovery objective) that, **when** reached, will assure populations remain stable. **The** remainder of this plan outlines steps necessary to achieve **the** recovery objective. Recovery goals, objectives and **tasks** may **change** as **we** learn **more** about the interior least **terns**.

Recognizing that **the** interior least **tern** has a broad distribution, **the** recovery objective was **set** by taking into account: 1) current data on distribution and abundance of interior least **terns** in **each** river system; 2) knowledge of how thoroughly **each** river system has **been surveyed**; 3) historic population data, **when** available; 4) loss of viable habitat; 5) an assessment of **the** potential to increase breeding pairs at currently occupied sites; 6) assessment of **the** potential to establish breeding pairs at unoccupied sites. Technical experts and state and federal resource agencies **were** consulted to **determine the status** of current populations and habitats, as well as **the** potential for population **increase**.

Therefore, in order to **be** considered for removal from **the** endangered species list, interior least **tern** essential habitat will **be** properly protected and **managed** and populations will **have** increased to 7,000 birds:

I. Missouri River System

- A. Number of birds in the **Missouri** River system will increase to 2,100 adults.
- B. Essential breeding habitat (Appendix 4) will **be** protected, enhanced and restored.
- C. **The** breeding pairs will **be** maintained **in** the following distribution for 10 years (assuming at least four major -censuses will have **been conducted during this time**):
 - Montana** - 50 adults
 - North Dakota** - 250 **adults**
 - South Dakota** - 680 adults (includes 400 shared with **Nebraska** on **the Missouri** River).
 - Missouri River** below Gavin's Pt. Dun - 400 adults
 - Lake Oahe** - 100 adults
 - Missouri River** below Ft. Randall - 80 adults
 - Other Missouri River sites** - 20 adult8
 - Chytynt River** - 80 adults
 - Nebraska** - 1520 adults (includes 400 adults shared with South Dakota on **the Missouri River**).
 - Missouri River** - 400 adults
 - Niobrara River** - 200 adults
 - Loup River** - 170 **adults**
 - Platte River** - 750 adults
 - Missouri and Iowa** - Opportunities for habitat restoration 'and rttstablishmtnt of breeding pairs will be determined.

- II. **Mississippi** and Ohio Rivers
 - A. Current number of adult birds (**2,200-2,500**) on the Lower **Mississippi** River will remain stable **for** the next ten years.
 - B. Essential breeding habitat (Appendix 4) **will** be protected, enhanced, and restored.
- 111. Arkansas River System
 - A. Numbers of birds on the **Arkansas** River system will increase to 1,600 adults.
 - B. Essential breeding habitat (Appendix 4) **will** be protected, enhanced and restored.
 - C. The 1,600 breeding adults **will** be maintained in the following distribution for 10 years:
 - Arkansas River, Arkansas • **150** adults
 - Arkansas River, Oklahoma • 250 adults
 - Quivira** National Wildlife Refuge • 100 adults
 - Salt Plains National Wildlife Refuge • 300 adults
 - Cimarron River Basin • 400 adults
 - Canadian River • **300** adults
 - Beaver/ North Canadian River • 100 adults
- IV. Red River System
 - A. Number of birds in the Red River system will increase to **300** breeding adults.
 - B. Essential Breeding habitat (Appendix 4) will be protected, enhanced and restored.
 - C. The 300 adults **will** be distributed along the Prairie Dog Town Fork **where** interior least terns currently occur and **at** other essential habitat sites yet **to** be determined.
- v Rio Grande River System
 - A. Current number of adult birds (**500**) in the **Rio** Grande River system will remain stable for 10 years.
 - B. Essential breeding habitat will be protected, **enhanced** and restored.
 - C. The birds will be distributed along the Rio Grande and Pecos Rivers.

Step-Down Outline

The step-down outline lists tasks **necessary to meet the recovery** objective. Steps (or tasks) are not presented in order of importance. **Some** steps are **underway**, while others may take years before they are begun. An explanation **of there** steps is presented in the Narrative section of this plan. Following the Narrative, the Implementation Schedule lists and sets priorities to be taken in the next three **years**. The step-down outline is very similar to the step-down outline in the Great lakes/Northern Great Plains Piping Plover recovery plan (U. S. Fish and Wildlife Service 1988a) because both species breed in the same habitat areas in the **Missouri River system** and require similar recovery tasks.

- 1. Determine current distribution and population trends of the interior least tern.
 - 11. Assess status and distribution of breeding populations.
 - 111. Survey sandbars, **reservoir** shorelines, sand and gravel pits and other suitable habitats to determine breeding

- distribution. .
112. Develop A method for **standardization** of census techniques And timing of censuses.
 - 113. Census known And potential breeding sites.**
 111. Monitor reproductive success.
 115. Assess **dispersal patterns** And genetic diversity.
 116. **Assess mortality.**
 117. **Further** identify life history parameters And develop **population** models.
 12. **Assess status** And distribution **for the migration** period.
 13. **Assess status** and distribution during the Winter.
 131. Survey **beaches** And other **suitable habitat** to **determine** winter distribution.
 132. Census known wintering **areas.**
 133. Monitor movement of **birds between wintering** sites And Assess mixing of **populations.**
 - 134. Assess mortality on wintering areas.**
2. Determine current **habitat** requirements And **status.**
 21. Determine breeding **habitat** requirements And **status.**
 211. Assess **the characteristics**, including prey resources, of breeding **habitat.**
 212. **Quantify** And **evaluate available** breeding **habitat.**
 213. Examine historic **aerial photography and hydrographic surveys** of river systems to determine the previous extent of **potential habitat** And **vegetational changes.**
 22. Determine current migration habitat requirements And **status.**
 221. Assess **the characteristics**, including prey resources, of **migration habitat.**
 222. **Quantify** And **evaluate available** migration **habitat.**
 23. Determine **current habitat** requirements And **status** on wintering **areas.**
 231. Assess **the characteristics**, including prey resources, of winter **habitat.**
 232. **Quantify** And **evaluate** winter **habitat.**
 3. Protect, ● nhAnct, And increase interior **least tern** populations.
 31. Protect, enhance, And **increase populations** during **the** breeding **season.**
 311. **Increase** reproduction **And, survival at** occupied breeding sites.
 3111. **Evaluate predator impacts** on eggs and chicks and identify species responsible for **the** predation.
 3112. **Evaluate techniques** for predator management And implement where **appropriate.**
 3113. Restrict public use within **'nesting areas** And investigate enforcement options.
 3114. **Manage water** levels And river flows to reduce nest And chick loss.
 3115. **Modify or eliminate construction activities** that Adversely **impact** reproductive success.

- 3116. **Investigate** the effects of environmental contaminants at breeding **areas**.
- 32. Protect and enhance populations during migration and winter.
 - 321. **Manage** areas to maximize survival of birds during migration.
 - 322. Manage** winter areas to maximize survival of birds during winter.
 - 3221. Investigate the effects **of** human activities on winter **survival**.
 - 3222.. Investigate *the effects* of** environmental contaminants.
- 4. Preserve and enhance habitat.
 - 41. Provide protection and management of breeding habitat.
 - 411. Identify areas of essential breeding habitat.
 - 412. Continue to evaluate areas **for** consideration as essential breeding habitat.
 - 413. Establish liaison with agencies and organizations with land and water management **responsibilities**.
 - 414. **Revise**, establish, **or** utilize land and water laws and regulations to provide protection along rivers and lakes.
 - 415. Develop criteria and priorities **for** breeding habitat protection.
 - 416. Develop management plans for breeding habitat.
 - 4161. Determine direct, indirect and cumulative effects of manipulation of river hydraulics, **flow** regimes, and sediment discharge on breeding and foraging habitat.
 - 4162. Identify river flow regimes that will protect and enhance breeding and foraging habitat.
 - 4163.** Determine the relationship **of** existing **•** rtificfal breeding sites to river sites.
 - 4164.** Identify **need** and **techniques** of improving habitat by **management** of substrate and **by** vegetation control' through physical and/or non-toxic chemical **means**.
 - 4165. Study feasibility and determine **need** for creating new habitat and implement trials to determine **success** rates of creating new habitat.
 - 4166. Develop lake **and** reservoir control policies where existing and potential interior **least tern** habitat **is threatened**.
 - 4167. Identify **needs and techniques for** managing water levels.
 - 417. **Evaluate** success **of** protection **and** management techniques.
 - 42. **Provide protection** and management **of** migration habitat.
 - 43. Provide protection and management **of** winter habitat.
 - 431. Identify **areas** of essential winter habitat.
 - 432. Develop criteria **and** priorities **for** winter habitat protection.
 - 433.. Develop management techniques.
 - 434. **Modify** construction activities **that** may reduce or negatively alter winter habitat.
 - 435. Evaluate success **of** protection and management techniques.
- 5. Develop and **implement** an education program that publicizes information on the interior least tern, including its life history, **reasons for**

- current **status**, and options for **recovery**.
51. Inform and educate the public **on the** bird's plight and recovery efforts.
 511. Identify target audiences **among** the general public.
 512. Develop and distribute educational materials appropriate **to** various audiences.
 513. Develop materials for newspapers, radio, and television that highlight specific interior least tern projects.
 514. Provide controlled viewing opportunities if and when appropriate.
 52. Inform and educate public resource management agencies.
 521. Identify critical resource agency **constituents**.
 522. Develop educational materials appropriate to respective agencies and their management authority.
 523. Provide public resource agencies with periodic updates on the interior least tern's status and progress of **recovery** efforts.
 6. Coordinate **recovery** efforts.
 61. Designate a recovery plan coordinator.
 611. Coordinate research and **management activities** with **federal, state**, local and private organizations.
 612. Coordinate international **research and management activities**.
 613. **Coordinate** development of a public **information** program at the national and international level.

Narrative

The Narrative gives further details and justification for each task in the Step-Down Outline. The steps critical for recovery in the next three years are outlined and given priority in the Implementation Schedule.

1. Determine current distribution and population trends of the interior least tern.

The effectiveness of current conservation efforts **will not** be well-understood until comprehensive distribution and census data **have** been collected. Future plans **for recovery** also will be curtailed until a **more accurate** picture of the species status is defined.

1.1. Assess status and distribution of breeding DOD-

Most interior least tern **censusing** has been carried out during the breeding season. Results indicate interior least terns are widely distributed, as scattered pairs or in concentrations at breeding areas. The terns probably disperse great distances as suggested by Boyd and Thompson (1965). Continued search for new breeding areas and evaluation of known areas are necessary to complete our knowledge of the birds' status.

111. Survey sandbars, reservoir shorelines, sand and gravel pits and other suitable habitats to determine breeding distribution.

Currently, the distribution of the interior **least tern** on most of the **Missouri** River system is well-known and monitored, although reservoir shorelines in the Dakotas and **Montana** should be further surveyed for accurate **population estimates** especially during **low** water years when reservoir levels are low. Additional **survey work** is needed on the **Loup** River in Nebraska and elsewhere in the **Platte** River system. The **Arkansas** River system needs further **survey work** in **Arkansas, Kansas, Oklahoma** and Texas. The length of the **Red** River requires a thorough **survey** as does the **Rio Grande** River system and rivers in central Texas. Additional **survey work** is needed on the Lower **Mississippi** River to determine distribution when the river rises and floods nesting colonies. The **Missouri** Department of Conservation has a study in progress to address this need. The status of potential sites should be monitored and updated at least once every five years.

112. Develop a method for standardization of census techniques and timing.

The exposure of sandbars in the spring follows the reduction of **river flows**. The breeding cycle may commence at different times throughout the interior least tern's range. Differences in breeding chronology from south to north must be determined. Because of the length of **time** involved in surveying long stretches of rivers, **surveys** should be correlated with reported river levels and the exposure of sandbars. Surveys should account for **re-nesting** birds and later nesting by younger adults (Massey and, Atwood 1981, Smith and Renktn 1990).

113. Census known and potential breeding sites.

Once sites are identified as containing breeding pairs, annual censuses of breeding and non-breeding adults should be carried out at essential breeding habitat (Appendix 4) for several years. If the birds are established for several years, censusing should **continue** at least once **every year**.

114. Monitor reproductive success.

Census data provide an indication of an area's population size, but **estimates** of reproductive success are also necessary. More adults may be present in nesting areas than **actually** breed. **Frequent nest destruction** further lowers productivity of a site, rendering simple counts of **breeding** pairs **less** meaningful than censuses of adults and fledged chicks. Reproductive success or recruitment (measured in terms of number of chicks fledged per pair) should be monitored annually at essential sites and at **least** every **three years, on a rotating** basis, at other sites. **Causes** of reproductive failure should be identified whenever possible. Because of possible early fledgling departure from colonies, multiple counts of fledglings should be made for

determination of the **fledging rate** (Thompson 1982, Thompson and **Slack** 1983).

115. Assess dispersal patterns and genetic diversity.

Little is known about the interaction between coastal least terns and the interior least tern. **Boyd** and Thompson (1985) found a nesting least tern in Kansas which had been banded **as** a chick on the Texas coast. **It** would be useful to know if coastal least terns **serve** as a reservoir to replenish the interior least tern population; and **if** the status of the coastal least tern population **determines** the numbers and distribution of interior least terns. Monitoring **movements of** marked birds in major breeding areas **will fill** the gap in our understanding of dispersal. knowledge of **how new** nest sites are colonized, and where **new** birds originated **will** be useful in developing population management plans and **models**.

116. Assess mortality.

Factors such as human disturbance, predation, and water level regulation have reduced **success** of interior **least** tern eggs and chicks (**Mayer** and Dryer 1990). **Factors affecting** adult mortality, however, have never been fully addressed for any part of the annual cycle. Predation **is a problem** for some California and coastal least terns (**Burger** 1984, **Minsky** 1980, **Massey** 1981) and the closely allied little tern in Europe (**Had&n and Knight** 1983). During the breeding season, predation on interior least terns by coyote (**Canis latrans**), crow (**Corvus brachyrhynchos**), and raptors has been reported (**G. R. Lingle**, personal communication, **Hill** 1985, **Klrsch** 1990, **Mayer** and Dryer 1990) and predation on nesting adults by barred owls (**Strix varia**) has been recorded (**Smith** and **Renken** 1990). Predation **is** significant on the **Missouri National Recreational River** (U. S. Fish and Ulldlife **Service**, unpublished data). It is **important** to determine the extent and cause of adult and **juvenile** mortality during the breeding season.

117. Further identify life history parameters and develop population models.

Field studies of **interior** least terns should be carried out without reducing reproductive **success** or site tenacity. Future breeding studies only should be undertaken after researchers have identified specific critical factors that require resolution in order to rehabilitate the **species**. It would be **useful** to compile **all available** life history data and develop a model to estimate potential population trends.

12. Assess status and distribution for the migration period.

Less is known about the migratory ecology for the interior least tern than for any other phase of the annual cycle. Migratory routes have not been adequately described for spring or fall. It is not known **if** interior least terns follow **major** river systems **during** migration or **if they** migrate directly north and south. Further, **it is unknown if** interior least **terns** join coastal least **terns** prior to coastal least tern migration to Latin America or **if interior least** terns have their own migration route. Before

intensive individual field studies are undertaken, **it may be** beneficial to coordinate surveys of potential sites with natural resource employees or local birders to determine **if** interior least terns are stopping en route to wintering sites.

13. Assess status and distribution during the winter.

Interior least terns spend 6-7 months at wintering sites. Most field research, however, has been **carried** out on breeding birds. Factors limiting non-breeding birds may be as severe or worse than threats encountered during other times of the year.. Field studies should **begin to** ● t'least locate wintering sites.

131. Survey beaches and other suitable habitat to determine winter distribution.

Biologists familiar with the avifauna of Atlantic and Caribbean coastal Latin America should be contacted to assist in determining the winter distribution of least terns. **A** survey of the north **coast** of South America should be carried out to identify those habitat types used by least terns. However, the surveys may be difficult. Accessibility of coastal **areas** along **central** America and the northern coast of South America may be problematic for geographical and political **reasons**. Color-banded **individuals** would provide the **means** to distinguish interior least terns **from** other races or populations.

132. Census known wintering areas.

Once winter sites are known, censuses of important areas will provide an indication of **their** continuing importance and **status as post-breeding** sites.

133. Monitor movement of birds between wintering sites and assess mixing of populations.

It is not known if **post-breeding** interior least terns mix with coastal least terns at wintering sites. Once the habitat types **of** interior least terns **are** known, **habitat** protection can begin. Monitoring movements of birds between different sites will provide this information, as well as indicate the degree to which individuals from various breeding populations **mix** during the winter.

131. Assess mortality on wintering areas.

The extent and cause of mortality to post-breeding interior least terns has not been addressed. It is not clear if adults and juveniles suffer differential **mortality**, or if post-breeding **birds face greater threats** than do breeding birds. Any information leading **to** further delineation of threats to the species during this time will **be** important.

2. Determine current habitat requirements and status.

Habitat alteration has been identified as one of the principal causes of the current status of the interior least tern (U. S. Fish and Wildlife Service 1985, Whitman 1988). Recovery of the species will be affected substantially by the ability to identify and protect essential breeding habitat and to intensively manage that habitat to maximize productivity and survival. Setting priorities for protection of remaining sites and determining habitat management actions will require detailed knowledge of interior least tern habitat requirements and the availability and quality of existing sites.

21. Determine breeding habitat requirements and status.

Our knowledge of interior least tern breeding habitat requirements has increased greatly during the past five years. Data on seemingly adequate but unoccupied habitat is needed. Comparison of habitat conditions among used sites along with data on reproductive success will provide the information necessary to set priorities for protection, and determine site-specific management actions to enhance breeding habitat.

211. Assess the characteristics, including prey resources, of breeding habitat.

The characteristics of breeding habitat must be investigated across the entire range of the interior least tern. At riverine sites, habitat variables to be measured should include: nesting area and height above water level, vegetative cover and distribution, substrate type, and river level fluctuations. Other variables may be of particular interest at local breeding areas. Measurements taken and methods employed at various breeding sites should be standardized to allow comparisons among areas. Few data are available on food resources at interior least tern breeding areas. Information on prey species occurrence and abundance are needed, as are estimates of the likelihood of food being a limiting habitat factor. The goals of these investigations should be identification of the range of habitat conditions tolerated by interior least terns, determination of habitat factors that affect nest densities, and elucidation of habitat conditions that may be related to maximum reproductive success rates.

212. Quantify and evaluate available breeding habitat.

As habitat assessment is undertaken, efforts to quantify existing interior least tern habitat should be initiated. The first task should be quantification of known and potential breeding habitat. As habitat quality data become available, existing sites should be evaluated with respect to habitat adequacy and deficiencies. Based on this information, recommendations for site protection or management actions should be given priorities. Remote sensing techniques such as aerial videography (Sidle and Ziewitz 1990) can be useful to quantify and, if possible, rate interior least tern breeding habitat. Sandbars are easily visible on satellite imagery of the Mississippi and Missouri Rivers. A catalog or compendium of interior least

tern nesting areas should be developed.

213. Examine historic aerial photography and hydrographic surveys of river systems to determine the previous extent of potential habitat and vegetational changes.

For many rivers periodic aerial photographs and hydrographic surveys are available. It would be useful for predictive purposes to measure the change, if any, in the quantity and quality of sandbar habitat since photo and hydrographic coverage began (Hamel et al. in press, Rodekoher and Engelbrecht 1988, Sidle et al; 1989). Such an endeavor would allow an accurate forecast of habitat trends.

22. Determine current migration habitat requirements and status.

Because migration patterns of interior least terns are not understood, no information on habitat requirements or status is available. Once stop-over sites, if they exist, are determined, evaluation of habitat requirements should be undertaken.

221. Assess the characteristics, including prey resources, of migration habitat.

If stop-over sites are identified, the habitats used should be described and variables characterizing those habitats quantified. Quantification (time activity budgets) of how interior least terns use the available habitats and their length of stay at stop-over sites also should be determined.

222. Quantify and evaluate available migration habitat.

Once migratory habitats are identified and characterized, the availability of such habitats should be determined. Initially, habitat availability in the vicinity of known stop-over sites should be quantified and its quality assessed. If migratory habitat in the vicinity of current stop-over sites is limited, a large scale survey of available habitat along suspected migratory corridors should be made.

23. Determine current habitat requirements and status on wintering areas.

No data are available on interior least tern winter habitat requirements. This task should be undertaken followed by a determination of the extent to which wintering habitats are traditionally used. Information on the role of winter habitat abundance, distribution, and quality in interior least tern population dynamics is totally lacking. Data relating winter habitat conditions to population status are needed.

231. Assess the characteristics, including prey resources, of winter habitat.

As primary wintering areas are identified, characteristics of the habitats used by interior least terns must be quantified and variables affecting quality of those habitats elucidated. Winter habitats should be assessed with regard to interior least tern, prey abundance and distribution, roost site needs, and location of feeding and roosting habitat. Habitat characteristics near occupied sites, but not currently used by interior least terns, also should be assessed. Quantitative data on interior least tern use of winter habitats also are needed. Information on movements

among wintering areas, movements among habitats, time-activity budgets, the use of pre-migration staging tracts, etc., may provide important information on habitat quality. The goal of these studies should be identification of habitat features that affect winter survival of interior least terns, assure adequate pre-breeding condition, and favor mixing among individuals from local breeding populations.

232. Quantify and evaluate winter habitat.

After baseline information on habitat characteristics and quality is available, the amount and distribution of winter habitat should be determined. Additionally, the quality of existing habitat should be rated and deficiencies identified. This effort may involve development of remote sensing techniques to identify and monitor winter habitat. Based on data generated under steps 231 and 232 the likelihood of winter habitat quantity limiting the growth of the interior least tern population should be evaluated. If winter habitat is found to be limited, further recommendations should be developed on the need for habitat protection or management of specific sites.

233. Eliminate current or potential threats to winter habitat.

As winter habitat is identified, current and potential threats to each site should be determined. Priority should be given to sites currently used by interior least terns. It is important to not only identify threats that could destroy winter habitats, but also those that could result in lowering the quality of remaining sites. Habitat ownership will have to be taken into consideration when assessing threats to the species.

3. Protect, enhance, and increase interior least tern populations.

Legal protection is often not enough to ensure perpetuation of breeding populations. Active management actions, including predator management, restricted access, and water level management are critical components of a comprehensive protection plan.

31. Protect, enhance, and increase populations during the breeding season.

To date, breeding activity of interior least terns has been more thoroughly investigated than activities at other times of the year. Current surveys have now identified most of the nesting areas in the U. S. Extensive survey work and research investigations of various major breeding areas have helped delineate many factors contributing to the species' current status, thus enabling the development of specific recommendations that may enhance the species' survival during the reproductive season.

311. Increase reproduction and survival at occupied breeding sites.

Activities that reduce interior least tern reproductive success and survival on its breeding grounds are probably among the principal factors responsible for the species' current status. Actions directed at eliminating or

minimizing such impacts are essential to the interior least tern's recovery.

3111. Evaluate predator impacts on eggs and chicks and identify species responsible for the predation.

Predation can be high in California and coastal least tern colonies (Atwood et al. 1979, Burger 1984, Massey 1981). Surveys on the Lower Mississippi River revealed that nest predation, especially by coyotes, has substantially reduced reproductive success at certain colonies. The vulnerability of terneries to such predation increases when island habitat accretes to the shoreline during periods of low water (Smith and Renken 1990). Studies conducted in the Missouri River system have documented a high percentage of interior least tern egg and chick loss to predation (Nebraska Game and Parks Commission, unpublished data, Mayer and Dryer 1990). During 1987-1989, predation accounted for most of the nest losses on the Platte River except riverine nests on the central Platte where flooding caused the mortality (Kirsch 1990, Lingle 1989). Both avian and mammalian species are among the suspected predators. Further studies that document such losses should continue. Investigations that focus specifically on identifying predators, and the cues they use in locating nests and/or chicks, determining the time of predation, etc.; are necessary if egg and chick mortality are to be curtailed.

3112. Evaluate techniques for predator management and implement where appropriate

Lethal and non-lethal methods for managing mammalian predators have been extensively developed for other wildlife management purposes. They include: eliminating or relocating the animal, erecting electric fences, and developing taste aversions. Electric fences have been used to protect nesting California and coastal least terns (Massey and Atwood 1980, 1982; Minsky 1980). The applicability of these and other techniques (e. g. predator exclusion cages) to the interior least tern should be investigated. Few management efforts have focused on managing avian predators, such as common ravens (*Corvus corax*), American crows, great horned owls (*Bubo virginianus*), great blue herons (*Ardea herodias*), California gulls (*Larus californicus*), and ring-billed gulls (*L. delawarensis*). Appropriate management measures should be implemented at interior least tern sites that are now experiencing significant and repeated loss due to predation.

3113. Restrict public use within nesting areas and investigate enforcement options.

Disturbance of California and coastal **leas tern** colonies caused by foot traffic and recreational vehicles has **been** well-documented (Massey and Atwood 1979, Goodrich 1982, Burger 1984) and is also **true for** interior least **terns** (Schwalbach 1988, Kirsch 1987-90, Lingle 1989, Smith and Rtnktn 1990). Losses **incurred** by **these** activities can **be** direct, by destroying **eggs** and chicks, as **well** as indirect, by inhibiting **territory** establishment, feeding behavior, incubation and other reproductive behavior. A variety **of** techniques that restrict access to nesting **areas** **have been** successful in a **few** states and should **be implemented on a wider scale**. **These include** posting, restricted access, and fencing (Morris 1979, 1980; Larkins 1984, Massey and Atwood 1979). Because many interior least **tern** nesting **areas** are located in remote areas, strict enforcement of regulations is often impractical. Although the **site** may **restrict** substantial recreational **use**, budget restrictions rarely allow full-time monitoring by professional staff. It is essential, therefore, that **actions** to restrict recreational activities always **be** accompanied by an aggressive public relations effort that will effectively reach all potential visitors to an area and adequately explain **the** purport of **the** regulations. **Tern wardens** who patrol nesting areas to explain **the** restrictions, should **be** considered for particularly important breeding areas (McCulloch 1982). **The** U. S. Army Corps of Engineers, U. S. Fish and Wildlife **Service**, and **state** wildlife agencies could **become** involved in public relations efforts and patrols to protect interior least **tern** nesting **areas** on **the** river systems. Agents of **the** Missouri Department of Conservation maintain an active **enforcement** program at **Mississippi** River **territories**. Similar **state** and **federal** enforcement **endeavors** **have** begun on **the** Missouri River in North and South Dakota, and Nebraska, and on **the** Platte River in **Nebraska**. Field research on Interior **least terns** should **be** carefully examined **for** its effects on the reproductive success of **the** birds (Brubeck et al. 1981). **Research** proposals should **be** scrutinized **for** their benefit to interior least tern recovery.

3114. Manage water levels and river flows to reduce nest and chick loss.

A significant proportion of **the** interior least **tern** population resides along rivers **where** much habitat has **been** destroyed **by** reservoir construction, channelization, water depletion, vegetative

encroachment, and modification of flow regimes (Currier et al. 1985, Nebraska Game and Parks Commission 1985b, Schwalbach et al. 1986, 1988, Eschner et al. 1981, Smith and Stucky 1988, Sidle et al. 1989). This riverine habitat is subject to a number of additional threats, including untimely water releases from dams that flood sandbar nesting habitat (Dryer and Dryer 1985, Schwalbach et al. 1986, 1988; Schwalbach 1988, G. R. Lingle, personal communication). Managing water levels early in the spring along some rivers could help to resolve this problem. Nesting habitat, expected to be flooded late in the season, could be submerged when interior least terns begin establishing territories in early May, forcing them to seek higher grounds that would be safe throughout the nesting season. It is essential, however, that sufficient nesting habitat is available above the fluctuation zone. High waters in spring also helps keep sandbars devoid of vegetation by reducing sprouting of young herbaceous growth ● pd by increasing deposition of coarse sediments (Currier et al. 1985, O'Brien and Currier 1987).

Annual flow regimes need to be developed for many river segments where interior least terns occur. For example, along the central Platte River the Service has developed flow recommendations to support a variety of wildlife including least tern nesting habitat and the bird's forage fish (Table 8). These recommendations have been accepted by the Federal Energy Regulatory Commission as part of the annual relicensing of upstream water projects in Nebraska (Sidle et al. 1990). The water releases will occur on the North Platte River, far upstream of interior least tern nesting habitat. The Ohio River has a major effect on the availability of interior least tern habitat in the lower Mississippi River. Management of this river and other rivers throughout the bird's range need to be examined for their effect on the interior least tern and its habitat.

3115. Modify or eliminate construction activities that adversely impact reproductive success of interior least terns.

Recreational and residential development along river fronts should be discouraged in nesting areas. Proposals for maintenance or development activities that do not directly disturb breeding habitat but that occur in the vicinity of nest sites should be closely scrutinized for their potential impact.

3116. Investigate the effects of environmental contaminants during the breeding season.

Contaminant effects on interior least terns are unknown. It would be useful to at least collect addled eggs during surveys and field studies for later contaminant analysis.

32. Protect and enhance populations during migration and winter.

Each year, 30 percent or less of the interior least tern's time is spent on the breeding grounds. A comprehensive protection plan also should focus on the species survival during migration and winter. However, migration and winter are the most poorly understood stages of the bird's life cycle and little can be recommended until migratory patterns are determined. The delineation of key areas where interior least terns spend non-breeding months is a critical step to enable the protection measures necessary for the birds' survival year-round.

321. Manage areas to maximize survival during migration.

Nothing is currently known about either the extent or causes of mortality that interior least terns might encounter during migration. Work that focuses on delineating migration routes (Step 12) should be expanded to focus on causes of mortality as well. When appropriate, measures should then be taken to lessen the impact upon the species.

322. Manage winter areas to maximize survival during winter.

During winter, interior least terns probably use open habitats. Sand, gravel, and/or cobbled marine beaches may be selected, as well as intertidal beach bars and flats.

3221. Investigate effects of human activities on winter survival.

Recreational, residential, and industrial developments each pose a potential threat to interior least terns by increasing the level of human activity. Moreover, hunting of terns in the United States may be a factor. To date, research studies have focused primarily on describing the impacts of human activities on nesting grounds. Future efforts also should be directed at collecting similar data from wintering areas, once such areas are discovered.

3222. Investigate the effects of environmental contaminants in wintering areas.

During surveys for interior least tern wintering areas, attention should be paid to coastal pollution. Chemical use and its impacts on foreign wintering areas should be evaluated.

4. Preserve and enhance habitat.

Because of major habitat losses and increasing demands on available habitat, protecting and enhancing existing and potential interior least tern habitat is a major concern. Important breeding areas have been identified but enhancement and protection of essential habitat has been limited. Little is known about those areas along the migration route or on the wintering grounds.

41. Provide protection and management of breeding habitat.

Essential breeding habitat (Appendix 4) will need delineation,

protection, and enhancement to provide **for** recovery **of** the species. Efforts should include increased **management** activities to provide better use and protection of existing and potential areas. Compatibility of other uses (e.g., recreation) for **breeding** areas **should** be **defined**. All essential habitat **needs** permanent protection, **where** possible, through appropriate **fee** title acquisition, permanent easement, cooperative agreements, and **memorandums** of agreement or understanding among federal agencies and private organizations (Appendix 2).

411. Identify areas of essential breeding habitat

Essential Habitat is listed in Appendix 4 to highlight known areas to be protected.

412. Continue to evaluate areas for consideration as essential breeding habitat.

Recognizing the fragile nature of much of the interior least tern's breeding habitat, **cont**inued evaluation and designation of essential habitat in primary breeding areas will protect areas from detrimental development.

413. Establish liaison with agencies and organizations with land and water management responsibilities.

Due to increasing pressure for development and use of land and **water resources** to meet human needs, efforts should be made to **communicate** with agencies, organizations, and individuals whose **decisions** affect **the future of** interior least tern habitat. The purpose would be to resolve conflicts between known development **actions** and future **conflicts** through planning of land and water development.

414. Revise, establish, or utilize land and water laws and regulations to provide protection along rivers and lakes.

Increasing demands for agricultural land and urban development, **wetland** drainage, power generation, water for irrigation, recreational space, and operation of river **reservoirs** have threatened or destroyed interior least tern habitat. Enforcement of laws and regulations, particularly those involving **instream** flow protection, 404 permits, and endangered or threatened species habitat protection, is needed to restrict or modify **such** developments on the remaining essential interior least tern habitat. All land- and water-use legislation should be scrutinized for potential impact to interior least tern habitat. Undesirable legislation should be modified and laws enacted that **will** expand the consideration given wildlife during water and land development planning.

419. Develop criteria and priorities for breeding habitat protection.

To provide **adequate** protection, some habitat **will have to be** purchased in **fee** title, or placed under a protective **• ♦ M O M ■ ♦** or cooperative landowner agreement. Although permanent protection of **essential areas** usually **will** be preferred, in some instances, temporary protection of ephemeral nesting areas may be achieved **through agreements** with private parties and public authorities. Protection of

areas listed as essential habitat (Appendix 4) is based upon tradition of occupancy, number of birds present, **site** productivity, **proximity to other protected sites**, imminence of habitat destruction, and ephemeral nature of the site.

416. **Develop management plans for riverine breeding habitat.**

Techniques may vary front site to **site** depending on need and opportunity, but plans **should be** developed for management of essential riverine habitat (see Step 2).

4161. **Determine direct, indirect, and cumulative effects of manipulation of river hydraulics, flow regimes, and sediment discharge on breeding and foraging habitat.**

Manipulation of river flow regimes and river hydraulics through water diversion, storage of flows by dams, discharge from dams for power generation, navigation and irrigation demands, bank stabilization, and channelization has significantly altered the **natural** dynamic processes responsible for loss and **creation of sandbars used for nesting** (Nunnally and Beverly 1986, Sandheinrich and Atchison 1986, Smith and Stucky 1988). As a result, breeding habitat could be lost at a higher rate than what is being created. **Modifications of river flow regimes through operation of reservoirs and lock and dams** also has caused concern for long-term effects of riverbed degradation on interior least tern habitat. Although **many direct** effects of human manipulations have been identified, suspected indirect and cumulative impacts of **ongoing and future river developments need to be determined.** Under Section 7 of the Endangered Species Act the U. S. Fish and Wildlife Service and the U. S. Army Corps of Engineers have consulted on the effects of proposed dams in the Platte River system, and are consulting on the effects of main stem dam operations on **interior least terns** along the **Arkansas and Missouri Rivers** (U. S. Fish and Wildlife Service 1987b, 1987c, 1989, 1990). Section 7 consultation provides an opportunity to protect much of the interior least tern's breeding habitat.

4162. **Identify river flow regimes that will protect and enhance breeding and foraging habitat.**

Control of river flows is desirable to prevent inundation of nests and young (Nebraska Game and Parks Commission 1985c), discourage growth of woody vegetation, and to maintain a **river with a nutrient base** necessary for production of fish used as food by interior least terns. Proper **instream flow is a major** goal of ongoing Section 7 consultations regarding the interior least tern.

4163. **Determine the relationship of existing artificial breeding sites to river sites.**

California and coastal least terns readily use man-made habitats. Islands, spoil piles, and beaches formed by dredged sand and gravel, and located immediately adjacent to the Platte River in Nebraska and elsewhere are used by interior least terns. A large percentage of the Platte River breeding population of interior least terns nests at sand and gravel pits. Dike fields are commonly used along the Mississippi River (Hamelet al. in press, Landin et al. 1985, Rumancik 1987, Smith and Renken 1990). Terns may use barges filled with sand on river segments now devoid of sandbar habitat. The importance of artificial habitat to recovery of the species, and to what extent such habitat can replace lost natural sandbars, should be determined.

4164. Identify need and techniques of improving habitat by management of substrate and by vegetation control through physical and/or non-toxic chemical means.

Existing woody vegetation may have to be removed from sandbars to provide suitable nesting habitat through physical or chemical means. Annual control may be necessary. Dredging and spreading sand or gravel of particular particle size could improve substrates for nesting and increase the height of sandbars to: prevent continuous inundation. Currently, the U. S. Army Corps of Engineers and the Platte River Whooping Crane Habitat Maintenance Trust have been clearing islands on the Missouri and Platte Rivers, respectively.

4165. Study feasibility and determine need for creating new habitat and implement trials to determine success rates of creating new habitat.

A variety of techniques have been used to create artificial nesting sites for the California and coastal least terns and to attract terns to the sites (Massey 1981, Fancher 1984, Kotliar and Burger 1984). Creation of artificial habitat may be necessary in areas where manageable habitat is non-existent. This may be particularly important in areas where natural habitat has been lost to channelization and water diversion. For example, most of the lower Missouri River (Iowa, Kansas, Missouri, and Nebraska) is now a channel and artificially created sites (e.g., ash disposal sites at power stations in Iowa) (Wilson 1984, 1986; Dinsmore and Dinsmore 1989) are the only habitat available; As part of the annual relicensing effort for upstream water projects along the Platte River in Nebraska, restored least tern nesting habitat has been ordered by the Federal Energy Regulatory Commission for each bridge segment in the central Platte (Sidle et al. 1990). Additional restoration

will be needed elsewhere along the Platte River. Habitat on the **Cimarron** River appears to be progressively deteriorating from upstream to **downstream** as the channel narrows and woody vegetation encroaches. Vegetation control likely **will** be necessary to maintain essential habitat. Likewise, habitat restoration **will** be necessary if least terns are to recover **in** the Iowa and Missouri reaches **of the Missouri** River. In the **Mississippi** River, the **Missouri** Department **of Conservation** and the **U. s. Army Corps** of Engineers have developed a cooperative proposal to construct **two** artificial islands between St. **Louis** and Cape **Girardeau, Missouri**. **Smith and Stucky** (1988) discussed other recommendations, including modification of dike structures.

4166. Develop lake and reservoir control policies where existing and potential habitat is threatened.

Water levels affect interior least tern reproductive **success** by increasing or decreasing the amount of habitat available on the shoreline of reservoirs (**e. g.**, Lakes **Oahe** and **Sakakawea** in the Dakotas, and Salt Plains National Wildlife Refuge, Oklahoma) and in dike fields. Changes in these levels during critical periods may delay initiation of nesting, **flood** nest sites or feeding areas, or increase the distance from nest sites to the water's edge. **Lakes** and reservoirs with **interior least** tern habitat must **be identified** and any policies **controlling** water levels need to be scrutinized to determine the effect on interior least tern reproductive success.

4167. Identify needs and techniques for managing water levels.

Lakes and reservoirs currently supporting nesting interior least terns or that provide suitable nesting habitat should be evaluated to determine if water level management **is** feasible. **Where** feasible, techniques should **be** developed to manage water levels to improve reproductive success.

418. Evaluate success of protection and management techniques.

Monitoring must be sufficient to detect and measure the positive effects of protection and management and to avoid potentially detrimental impacts on interior least tern habitat. Daily and seasonal activity patterns of interior least terns, along with locations of specific nesting **areas**, will provide key measures of the birds' **response** to various **management** practices. **Monitoring** vegetation to determine where changing habitat conditions **exist and monitoring potential** predator **levels** in the area should be considered. **All techniques** used to **improve** interior least tern habitat should be evaluated **to** determine their cost-efficiency.

42. Provide protection and management of migration habitat.
 If migration sites are identified, their protection and enhancement will be essential. At that point, assessment of further needs of migrating interior least terns will be carried out. As stop-over habitats are identified, current and potential threats to those sites should be delineated. On publicly-owned sites, current land-use patterns or management actions that could conflict with interior least tern use of existing habitats should be identified. Feasibility of protecting major **privately-owned stop-over sites should be assessed.**
43. Provide protection and management of winter habitat.
 Survival and continued existence of "the species may depend on availability of suitable winter habitat. Furthermore, reproductive success of adults may partially be a function of their physical condition as they begin spring migration. Consequently, the quality and quantity of winter habitat may limit recovery of the species.
431. Identify areas of essential winter habitat.
 Essential winter habitat first needs to be identified by surveys in Latin America.
432. Develop criteria and priorities for winter habitat protection.
 Once further research is carried out in wintering areas, factors will be identified as being essential for winter habitat. At that point, a land protection strategy should be developed. Areas that support the greatest number of interior least terns, especially those supporting individuals from important sub-populations should be given priorities in a habitat management/protection plan.
433. Develop management techniques.
 Once actual and/or potential interior least tern wintering habitat is identified, methods of managing those habitats should be developed and improved so that wintering habitat is of sufficient quantity and quality to accommodate and promote expansion of interior least tern populations to more stable levels.
5. Develop and implement an education program that publicizes information about the interior least tern, including its life history, reasons for current status and options for recovery.
 Conservation of coastal least terns has benefitted greatly from public information endeavors (Jackson and Jackson 1985, Toups 1976). The interior least tern's successful recovery will depend on curtailing and/or redirecting human recreation and development activities. Therefore, resource managers and the general public should be provided with sufficient information to explain and justify changes in previous actions. Current efforts to develop a public information program have made an impressive start in this direction but must be intensified. These efforts also could benefit from better coordination at the national level to target specific audiences.
51. Inform and educate the public on the bird's plight and recovery efforts.
 The first priority in developing a public information program

should be to educate the general public about the significance and value of the interior-least tern. The public's support and cooperation ultimately will be essential **to** the species full recovery.

511. Identify target audiences among the general public.

Materials prepared to increase public awareness and appreciation of the interior least tern can be more effective if they are developed to **meet specific interests** and concerns of a particular audience. Time should be spent delineating which public groups are affected, either directly or indirectly, by interior least tern conservation **efforts** and how each audience can best be reached.

512. Develop and distribute educational materials appropriate for

v a r i o u s -

Current efforts should be expanded to make greater use of the various media, including newspapers, radio, and television. The primary **focus** of **this** task should be to provide background information describing the interior least tern's life history and habitat requirements and to describe how human activity/disturbance can threaten the survival of interior least terns. The public should also be made aware of the necessity to enact local regulations to protect the interior least tern. However, **information** materials should not increase the potential for observer disturbance to nesting birds. The **Service's** Tulsa office has produced an information brochure useful throughout the range of the interior least tern.

313. Develop materials for newspapers, radio, and television that highlight specific interior least tern projects.

In **several states**, cooperative **projects** between state and **federal** agencies, as well as private organizations and individuals are **underway** to protect interior least terns. Such **efforts** which generate public support should be applauded and widely publicized, particularly **at** the local level.

514. Provide controlled viewing opportunities if and when

Guided opportunities for **observing** interior least terns may be one of the best vehicles for generating public support and concern. **Led** by a qualified biologist under conditions that minimize or prevent disturbance to **the** birds, such trips can educate visitors first-hand about the need **for** strong protection and curtailment of **some** recreational activities.

52. Inform and educate public resource management agencies.

Some interior least terns occur on **lands that** are protected **and/or** managed by **state** and federal resource agencies. Recreation permitted on these **areas** (e.g., hiking, vehicle use, camping) can **reduce** the **bird's** reproductive success. In some **areas** an **agency's** own **activities** may also pose a **threat** (e.g., control of water levels in lakes and along rivers). Contact with

these agencies will facilitate better management of the areas for interior least terns.

521. Identify critical resource agency constituents.

Each resource agency-(including state, federal, and private organizations) whose activities can impact the interior least tern should be identified.

522. Develop educational materials appropriate to respective agencies and their management authority.

Resource managers need to be provided with basic life history information about the interior least tern as well as specific management information and recommendations directly pertinent to their area of responsibility.

523. Provide public resource agencies with periodic updates on the interior least tern's status and progress of recovery efforts.

It is important that each public agency responsible for ensuring the interior least tern's survival, either directly or indirectly, be kept abreast of the success of their efforts at both the local and national level. Periodic updates not only inform them of progress being made, but also remind them of their responsibilities to the conservation of interior least terns.

6. Coordinate recovery efforts.

Development of a recovery plan for interior least terns involves coordination of biologists, agencies, and governments so that the most comprehensive, up-to-date information is collected and disseminated in an efficient way. Proper coordination would also help ensure rapid implementation of actions necessary for full recovery.

61. Designate a recovery plan coordinator.

Designation of a coordinator is recommended. Duties of the coordinator would include: a) coordination of the implementation of the recovery plan; b) naming an individual in each state to coordinate and implement recovery tasks; c) monitoring execution of the plan's implementation schedule; d) maintaining collaboration with state, federal, and international agencies; disseminating critical annual data; and coordinating range-wide research activities for interior least terns. A least tern contact person should also be designated for each state.

611. Coordinate research and management activities with federal, state, local, and private organizations

Efficient achievement of recovery goal; will be enhanced through coordination of research and management with private and governmental agencies. For example, it would be useful to establish and coordinate an international banding scheme whereby birds can be easily identified throughout the annual cycle. The recovery plan outlines many facets of interior least tern conservation that require urgent investigation. Repetition of efforts due to lack of coordination will slow the recovery process and may cause undue disturbance to the birds.


612. Coordinate international research and management activities.

Development of population management plans on an

international scale may be necessary. Interior **least** terns probably winter **in** Latin America • d coordination with various nations and **international conservation organizations** **may** be necessary.

613. Coordinate development of a public information program at the national and international level.

Information **and** educational materials developed in one river system could be **of** equal benefit in other river systems. Some materials also **may** be helpful to **states that** support wintering populations. Coordination **at** the **federal** level will reduce duplication of effort • d encourage more efficient use of time and **money at** the state level. A coordinated approach to raising **an** • mreness **of** the interior least tern's plight **at** the **international** level would ensure protection throughout **its** range.



IV. REFERENCES

- Akers, J. W. 1975. The least tern in Virginia; breeding biology and population distribution. **M. A. thesis**, College of William and Mary, Williamsburg, Virginia. **75 pp.**
- Altman, R. L., and R. D. Gano, Jr. 1984. **Least** terns nest alongside Harrier jet pad. *Journal of Field Ornithology* **55**: 108-109.
- American Ornithologists' Union. 1957. Checklist of North American birds. Fifth edition. Baltimore, American Ornithologists' Union. 691 pp.
- American Ornithologists' Union. 1983. Checklist of North American birds. Sixth edition. Lawrence, Kansas, American Ornithologists' Union. 877 pp.
- Anderson, E. A. 1983. Nesting productivity of the interior least tern in Illinois. Unpublished report, **Cooperative** Wildlife Research Laboratory Southern Illinois University, Carbondale. 19 pp.
- Anderson, R. 1971. Nesting least terns. *Audubon Bulletin* **160:1718**.
- Atwood, J. L., R. A. Erickson, P. R. Kelly, and P. Unitt. 1979. California least tern census and nesting survey. California Department of Fish and Game. Job V-2.13.
- Atwood, J. L., and D. E. Minsky. 1983. **Least** tern foraging ecology at three major California breeding colonies. *Western Birds* **14:57-71**.
- Atwood, J. L., B. W. Massey, and C. T. Collins. 1984. Movement of the Huntington beach least tern colony: an assessment of possible impacts. U. S. Fish and Wildlife Service, Laguna Niguel, California. Unpublished report. **28 pp.**
- Atwood, J. L., and P. R. Kelly. 1984. Fish dropped on breeding colonies as indication of least tern food habits. *Wilson Bulletin* **96:34-47**.
- Bent, A. C. 1921. Life histories of North American gulls and terns. U. S. National Museum Bulletin 113. 345 pp.-
- Blodget, B. G. 1978. The effects of off-road vehicles on least terns and other shorebirds. **M. S. thesis**, University of Massachusetts, Amherst. **79 pp.**
- Boyd, R. L. 1983. Population ecology of snowy plover and least tern in Kansas. Unpublished report. 34 pp.
- Boyd, R. L. 1984. Population ecology of snowy plover and least tern in Kansas. Kansas Fish and Game Commission. Non-game wildlife project report. **17 pp.**
- Boyd, R. L. 1986. Habitat management and population ecology studies of the least tern in Kansas. Kansas Fish and Game Commission. Unpublished report.
- Boyd, R. L. 1987. Habitat management and population ecology studies of the least tern in Kansas. Kansas Fish and Game Commission. Unpublished report.
- Boyd, R. L., and B. C. Thompson. 1985. Evidence for reproductive mixing of least tern populations. *Journal of Field Ornithology* **56:405-406**.
- Brubeck, Ii. V., B. C. Thompson, and R. D. Slack. 1981. The effects of trapping, banding and patagial tagging on the parental behavior of least terns in Texas. *Colonial Waterbirds* **4:54-60**.
- Burger, J. 1984. Colony stability in least terns. *Con&r* **86:61-67**.

- Burger, J.** 1988. Social attraction in nesting least terns: effects of numbers, spacing and pairs. *Condor* **90:575-582.**
- Burleigh, T. D., and G. H. Lowery, Jr.** 1942. An inland race of *Sterna albifrons*. *Museum of Zoology, Occasional Papers*, number **10:173-177**, Louisiana State University.
- Burroughs, R. D., ed.** 1961. The natural history of the **Lewis** and Clark expedition. Michigan State University **Press**. 340 pp.
- Burroughs, R.** 1966. A study of the breeding biology of least terns on Nantucket Island. M. S. thesis, University of Massachusetts, **Amherst**. 87 pp.
- Campbell, L.** 1935. **Least** tern taken near **Toledo**, Ohio. *Auk* **52:87.**
- Carreker, R. G.** 1985. Habitat suitability in x models: least tern. U.S. Fish and Wildlife Service Biological report **82(10.103)**. 29 pp.
- Carter, M. F.** 1989. Breeding least tern inventory. Job progress report, project SE-12-1, Colorado.-
- Chase, C. and C. Loefflter.** 1978. **Arkansas** valley shorebird inventory. Colorado Division of Wildlife.
- Couts, E.** 1874. Birds of **the** northwest: a handbook of the ornithology of the region drained by **the Missouri** River and its tributaries. U. S. Geological Survey of **the** Territories, **Miscellaneous** publication number 3. 791 pp.
- Craig, A. M.** 1971. Survey of California least tern nesting sites. California Department of Fish and **Game**. Special Investigation, Project W-54-R-4, Job 11-5.1. 7pp.
- Currier, P. J., G. R. Lingle, and J. G. VanDerwalker.** 1985. Migratory bird habitat on **the** Platte and North Platte Rivers in Nebraska. The Platte River Whooping Crane Critical Habitat Maintenance Trust, Grand Island, Nebraska.
- Davis, M. E.** 1968. Nesting behavior of **the** least tern (*Sterna albifrons*). M.S. thesis, University of California, **Los Angeles**. 72 pp.
- Dinsmore, J. J., and S. J. Dinsmore.** 1989. Piping plover and least tern population and habitat in western Iowa. Unpublished report, 17 pp.
- Dirks, B. J.** 1990. Distribution and productivity of least terns and piping plovers along the Missouri and Cheyenne Rivers in South Dakota, M.S. Thesis, South Dakota State University, Brookings.
- Downing, R. L.** 1980. Survey of interior least tern nesting populations. *American Birds* **34:209-211.**
- Dryer, M. P., and P.J. Dryer.** 1985. Investigations into **the** population, breeding sites, habitat characteristics, threats, and productivity of the least tern in North Dakota. U. S. Fish and Wildlife Service. Resource **information** paper number 1. **Bismarck**, North Dakota. 17 pp.
- Ducey, J. E.** 1981. Interior least tern (*Sterna antillarum athalassos*). U.S. Fish and Wildlife Service, Pierre, South Dakota. Unpublished report. 56 pp.
- Ducey, J. E.** 1985. **The** historic breeding distribution of the least tern in Nebraska. *Nebraska Bird Review* **53(2):26-36.**
- Duccy, J.** 1988. Nest scrape characteristics of piping plover and least tern in Nebraska. *Nebraska Bird Review* **56:42-44.**
- Eschner, T., R. Hadley, and K. Crowley.** 1981. Hydrologic and morphologic changes in **the** Platte River Basin in Colorado, Wyoming and Nebraska: a historical perspective. U. S. Geological **Survey** open file report 81-1125. U. S. Geological **Survey**, Denver, Colorado.

- Faanes, C. A. 1983. **Aspects of the nesting ecology** of least terns and piping plovers in central Nebraska. *Prairie Naturalist* **15:145-154.**
- Fancher, J. M. 1984. A technique for making least tern decoys. *Journal of Field Ornithology* **55:241-243.**
- Fisk, E. J. 1975. **Least tern:** beleaguered, opportunistic and roof-nesting. *American Birds* **29:15-16.**
- Fisk, E. J. 1978. Roof-nesting **terns, skimmers, and plovers** in Florida. *Florida Field Naturalist* **6(1):1-22.**
- Funk, J. L., and J. W. Robinson. 1974. Changes in the channel of the lower **Missouri River** and effects on fish and **wildlife.** Aquatic Series No. 11. Missouri Department of **Conservation,** Jefferson City, **MO.**
- Gochfeld, M. 1983. Colony site selection by least terns: physical attributes of **sitar.** *Colonial Waterbirds* **6:205-213.**
- Goodrich, L. J. 1982. **The effects of disturbance on the reproductive success of the least tern.** M. S. **Thesis,** Rutgers State University, New **Brunswick, New Jersey.** 100 pp. **Grover, P. B. 1979.** Habitat requirements of charadriiform birds nesting on salt flats at Salt Plains National **Wildlife Refuge.** M. S. thesis, Oklahoma State University, Stillwater. **38 pp.**
- Grover, P. B., and F. L. **Knopf.** 1982. Habitat requirements and breeding **success of** charadriiform birds nesting at Salt Plains National Wildlife Refuge, Oklahoma. *Journal of Field Ornithology* **53:139-148.**
- Haddon, P. C., and R. C. **Knight.** 1983. A guide to little tern conservation. Royal Society for the Protection of Birds, Bedfordshire, Great Britain. 114 pp.
- Hallberg, G. R., J. M. Harbough, and P. M. Witniok.** 1979. Changes in the channel areas of the **Missouri River** in Iowa from 1879 to 1976. **Iowa Geological Survey Special Report, Series Number 1.**
- Hamel, P. B., M. L. Bierly, and M. G. Ualdron** (in press) **Habitat dynamics of least terns in Tennessee: a preliminary investigation.** *Colonial Waterbirds*
- Hardy, J. W. 1957. The least **tern in the Mississippi River.** Publication of the Museum, Michigan State University, *Biological Series* **1:1-60.**
- Hill, L. A. 1985. Breeding **ecology of** interior least terns, snowy plovers, and American avocets at Salt **Plains National Wildlife Refuge,** Oklahoma. M. S. thesis, Oklahoma State University, Stillwater.
- Hoffman, J. C. 1986. The distribution of the interior least tern in northeastern Oklahoma during the 1986 nesting season. Unpublished report.
- Jackson, J. A. 1976. **Some aspects of the nesting ecology** of least terns on the Mississippi Gulf coast. *Mississippi Kite* **6(2):25-35.**
- Jackson, J. A., and B. J. S. Jackson. 1985. **Status, dispersion and population changes of the least tern** in coastal **Mississippi.** *Colonial Waterbirds* **8(1):54-62.**
- Janssen, R. B.** 1986. **Least tern in Lyon County.** *Loon* **58:48-49.**
- Jrtnigan, L. S., Jr. 1977. Comparison of least **tern (Sterna albifrons antillarum)** population **size and nesting habitat** on barrier and dredge islands in the North **Carolina coastal zone.** M. S. thesis, North Carolina State University, Raleigh. 51 pp.

- Jernigan, L. S., J. F. Parnell, and T. Quay. 1978. Nesting habitats and breeding populations of the least tern (*Sterna albifrons antillarum*) in North Carolina. National Oceanic and Atmospheric Administration, Sea Grant Publication **UNC-SG-78-07**. 39 pp.
- Johnson, R. 1987. Least tern survey of the Wabash River, 1987 and evaluation of available habitat. Endangered species progress report E-1-1. Indiana Department of Natural Resources.
- Jung, C. 1935. Occurrence of the least tern (*Sterna antillarum*) in Wisconsin. **Auk 52:87**.
- Jungemann, P. K. 1988. Observations and management proposals for the interior least tern (*Sterna antillarum athalassos*) in southeastern New Mexico. Prepared by Professional Services Contractor for USFWS-Bitter Lake National Wildlife Refuge and the New Mexico Department of Game and Fish. 39 pp.
- Kirsch, E. M. 1987. Annual Report 1987: Least Tern and Piping Plover on the lower Platte River in Nebraska. Nebraska Game and Parks Commission. Unpublished report.
- Kirsch, E. M. 1988. Annual Report 1988: Least Tern and Piping Plover on the lower Platte River in Nebraska. Nebraska Game and Parks Commission. Unpublished report.
- Kirsch, E. M. 1989. Annual Report 1989: Least Tern and Piping Plover on the lower Platte River in Nebraska. Nebraska Game and Parks Commission. Unpublished report.
- Kirsch, E. M. 1990. Final report 1990: least Tern and Piping Plover on the lower Platte River in Nebraska. Nebraska Game and Parks Commission. Unpublished report.
- Kotliar, N. B., and J. Burger. 1984. The use of decoy to attract least terns (*Sterna antillarum*) to abandoned colony sites in New Jersey. 'Colonial Waterbirds 7:134-138.
- Krtil, R., and M. P. Dryer. 1987. Nesting of the interior least tern on the Yellowstone River in North Dakota. Prairie Naturalist 19:135-136.
- Landin, M. C., J. Rumancik, E. E. Parks, E. Scott Clark, and E. Buglevicz. 1985. Interior least terns in the lower Mississippi River and its tributaries: two years surveys. U. S. Army Corps of Engineers, Vicksburg, Mississippi. Unpublished report. 15 pp.
- Larkins, D. 1984. Little tern breeding colony on artificial site at Port Botany, New South Wales. **Cortilla 8(1):1-10**.
- Linglt, G. R. 1989. Least tern and piping plover nesting ecology along the central Platte River Valley, Nebraska. Progress report to U.S. Fish and Wildlife Service.
- Lyons, J., and T. Randlt. 1988. Platte River channel characteristics in the big bend reach. U. S. Bureau of Reclamation, Denver, Colorado. Unpublished report. 69 pp.
- Marlatt, S. L. 1984. History and management recommendations for the interior least tern in New Mexico. New Mexico Department of Game and Fish. Unpublished report. 42 pp.
- Marlatt, S. L. 1987. Observations and management recommendations for the interior least tern in Chaves County, New Mexico. New Mexico Department of Game and Fish. Unpublished report. 19 pp.
- Massey, B. W. 1972. The breeding biology of the California least tern. M. S. thesis, California State University, Long Beach. 101 pp.

- Massey, B. W. 1974. Breeding biology of California least tern. Proceedings of the Linnaean Society, Ntv York **72:124.**
- Massey, B. W. 1976. Vocal difftrencts between American least terns and the European little tern. Auk **93:760-773.**
- Massey, B. W. 1981. A least **ternmakes** a right turn. Natural History **90(11):62-71.**
- Massey, B. W., and J. L. Atwood. 1978. Plumages of **the least tern.** Bird-banding **49:360-370.**
- Massey, B. W., and J. L. Atwood. 1979. Application **of ecological** information to habitat management for **the California least tern.** Progress report number 1. U. S. Fish and Uildlift **Service, Laguna Niguel,** California.
- Massey, B. V., and J. L. Atwood. '1980. Application of **ecological** information to habitat **management** for **the California least tern.** Progress report number 2. U. S. Fish and Wildlife Service, **Laguna Niguel,** California.
- Massey, B. W., and J. L. Atwood. 1981. Second-vavt nesting of the California least **tern: age** composition and reproductive **success.** Auk **98:596-606.**
- Massey, B. W., and J. L. Atwood.' 1982. **Application of ecological** information to habitat management **for the California least tern.** progress report number 4. U. **S. Fish and Wildlife Service, Laguna Niguel, California.**
- Massey, B. W., and J. L. Atwood.** 1983. **Application of tcological** information to habitat mntngement **for the California least tern.** Progress report number 5. U. S. Fish and **Wildlife Service, Laguna Niguel,** California.
- Mayer, P. M., and M. P. Dryer.** 1988 Population biology of piping plovers and least terns on the Missouri **River in North Dakota and Montana: 1988 field season** report. U. S. Fish and **Wildlife Service, Bismarck,** North Dakota. Unpublished report.
- Mayer, P. **M., and M. P. Dryer.** 1990. Population biology **of** piping plovers and least terns on the Missouri River in North Dakota: 1989 field season report. U. S. Fish and **Wildlife Service, Bismarck,** North Dakota. Unpublished report.
- Mayfield, H.** 1943. Least **tern** in southeastern Michigan. Wilson Bulletin **55:245.**
- McCament, D., and B. C. Thompson.** 1985. Interior least **tern** distribution and taxonomy. Texas Parks and **Wildlife Department.** Annual performance report, Federal aid project number W-103-R-15, job number 54. 13 pp.
- McCament, D. and B. C. Thompson.** 1987. Interior least tern distribution and taxonomy. Texas Parks and Vildlift Department. Annual performance report Federal aid project **number** U-103-R-16.
- McCament-Locknane, D.** 1988. Interior least tern distribution and taxonomy. Texas Parks and Uildlift Department. **Final** report, Federal aid project number U-103-R-17.
- McCulloch, E. M.** 1982. Warden appointed for little **terns.** Bird Observer **611:89-91.**
- Mills, C. E. 1987. **Indiana's first least tern** nesting record; Indiana Audubon Quarterly **65:42-44.**
- Minsky, D. 1980. Preventing fox predation at a least **tern** colony with an electric fence. Journal of Field Ornithology **51:80-81.**

- Monson, G. and A. Phillips. 1981. Annotated checklist' of the birds of Arizona. University of Arizona Press, Tucson. 240 pp.
- Montana piping plover recovery committee. 1988. Results of surveys for piping plover (*Charadrius melodus*) and least tern (*Sterna antillarum*). Unpublished report. 24 pp.
- Morris, A. K. 1979. The declining status of the little tern in Nev South Wales. *Corell* 3(5):105-110.
- Morris, A. 1980. Little tern (*Sterna albifrons*). Parks and Wildlife, Endangered animals of Ntv South vales. August 1980 pages 33-37.
- Moseley, L. J. 1976. Behavior and communication in the least tern (*Sterna albifrons*). Ph. D. dissertation, University of North Carolina, Chapel Hill. 164 pp.
- Moser, R. 1940. The piping plover and least tern in Omaha. *Nebraska Bird Review* 8:92-94.
- Nebrasks Game and Parks Commission. 1985a. Niobrara River interior least tern and piping plover nesting survey. Unpublished report.
- Nebraska Game and Parks Commission. 1985b. Biological opinion: Little Blue-Catherland Project. Unpublished report. 96 pp.
- Nebraska Game and Parks Commission. 1985c. Missouri River least tern and piping plover habitat management proposal presented to the U. S. Army Corps of Engineers. Unpublished report. 4 pp.
- Nebraska Game and Parks Commission. 1987. Platte River interior least tern and piping plover nesting survey. Unpublished report. 33 pp.
- Nunnally, N. R., and L. B. Beverly. 1986. Morphologic affects of lover Mississippi River dike fields. *Miscellaneous Paper E-86-2*, U. S. Army Waterways Experiment Station, Vicksburg, Mississippi.
- Neck, R. W., and D. H. Riskind. 1981. Direct and indirect human impact on least tern nesting success at Falcon Reservoir, Zapata County, Texas. *Bulletin of the Texas Ornithological Society* 14:27-29.
- O'Brien, J. S., and P. J. Currier. 1987. Channal morphology, channel maintenance and riparian vegetation changes in the big bend reach of the Platte River in Nebraska. Unpublished report. 49 pp.
- Phillips, A., 3. Marshall, and G. Monson. 1964. The birds of Arizona. University Of Arizona Press, Tucson. 212 pp.
- Ridgway, R. 1895. The ornithology of Illinois, part I, descriptive catalogue. *Natural History Survey Illinois*. Springfield, Illinois. pages 247-248.
- Rodekohr, D. A., and K. W. Engtlbrtcht. 1988. Island and bank morphological changes detected in the Platte-River bounding the Papio Natural Resources District from 1949 through 1988. Ctntr for Advanced Land Management and Information Technologies. University of Nebraska-Lincoln. 28 pp.
- Rumancik, Jr. J. P. 1985. Survey of the interior ltut tern on the Mississippi River from Cape Girardeau, Missouri to Greenville, Mississippi. U. S. Army Corps of Engineers, Memphis District, Memphis, Tennessee. Unpublished report.
- Rumancik, Jr. J. P. 1986. Population survey of the interior least tern on the Mississippi River from Cape Girardeau, Missouri to Greenville, Mississippi, 1986. U. S. Army Corps of Engineers, Memphis District,

- Memphis, Tennessee. Unpublished report. 19 pp.
- Rumancik, Jr. J. P. 1987. Population survey of **the** interior least tern on **the** Mississippi River from Cape Girardaau, **Missouri** to Grtttvillt, -Mississippi, 1987. U. S. Army Corps of Engineers, Memphis District, Memphis, Tennessee. Unpublished report. 22 pp.
- Rumancik, Jr. J. P. 1988. Population survey of the interior least tern on **the** Mississippi River from Cape Girardeau, Missouri to **Greenville**, Mississippi, 198.8. U. S. Army **Corps** of Engineers, Memphis District, Memphis, Tennessee. Unpublished report. 25 pp. and appendices.
- Rumancik, Jr. **J. P.** 1989. Population survey of the interior least **tern** on the Mississippi River from Cape Girardeau, Missouri to Vicksburg, Mississippi, 1989. U. **S.** Army Corps of Engineers, **Memphis** District, Memphis, Tennessee. Unpublished report.
- Sandheinrich, **M. B.**, and **G. J.** Atchison. 1986. Environmental effects of dikes of dikes and revetments on large riverint systems. Technical Report **E86-5**, " U. S. Army Waterways Experiment Station, Vicksburg, Mississippi.
- Schmulbach**, J. C., J. J. **Schuckman**, and E. A. Nelson. 1981. Aquatic habitat inventory of the Missouri River from **Cavins** Point Dam to Ponca State Park, Nebraska. U. S. Army Corps of Engineers, **Omaha**. Unpublished report. **15 pp.**
- Schulenbrg, E., J. H. Schulenberg, and M. B. Schulenberg. 1980. Distribution and ecological study of **the** least **tern** in Kansas. Kansas Fish and Game Commission. Non-game **wildlife** project. 110 pp.
- Schultnberg, J. H., and **M. B.** Ptacek. 1984. Status of the interior least tern in **Kansas**. American Birds **38:975-981**.
- Schvalbach, **M.** 1988. Conservation of **least** terns and piping plovers along the Missouri River and its major vesttrn tributaries in South Dakota. M.S. thesis, South Dakota State University; **Brookings**.
- Schvalbach, M., G. Vandel, and K.Higgins. 1986. Status; distribution, and production of the interior least **tern** and piping plover along **the mainstem** Missouri River in South Dakota, 1986. Report number **86-10** to **the** U. S. Army Corps of Engineers, Missouri River Division, Omaha, Nebraska.
- Schvalbach, M., G. Vandel, and K. **Higgins**. 1988. Status, distribution, and production of the interior **least** tern and piping plover along the **mainstem** Missouri River in South Dakota, 1986-1987. Completion report to **the** U. S. Army Corps of Engineers, Missouri River Division, Omaha, Nebraska.
- Seibert, H. C. 1951. Least **terns** in southeastern Ntv **Mexico**. Condor **53:204**. Shomo, L. S. 1988. **Observations** on **the** interior least tern near **Roswell, New Mexico**, May-August 1987. Ntv Mexico Department of Game and Fish. Contract number 519-76-01. 22 pp.
- Sidele, J. G., J. J. **Dinan**, M. P. Dryer, J. P. Rumancik, Jr., and J. **W.** Smith. 1988. Distribution of **the** least **tern** in interior North America. American Birds **42:195-201**.
- Sidele, J. G., E. **D.** Miller, and P. J. Currier. **1989**. Changing habitats in the Platte River valley of Nebraska. Prairie **Naturalist** **21:91-104**. Sidele, J. G., and J. **W.** Ziewitz. 1990. Use of aerial videography in **wildlife** habitat studies. **Wildlife** Society Bulletin **18:56-62**
- .. Sidele, J. G., M. **LeValley**, and J. G. **VanDerwalker**. 1990. **FERC** attempts to protect Platte River. National Wetlands Newsletter **12(4):8-10**.

- Smith, J. V. 1985. Improving the status of endangered species in Missouri (interior least tern habitat and nest survey). Missouri Department of Conservation endangered species project **number SE-01-12**. 142 pp.
- Smith, J. V. 1986. 1986 survey of the interior least tern on **the** Mississippi River (Cape Girardeau, Missouri to Island **number 20, Tennessee**). Missouri Department of Conservation. Unpublished report.
- Smith, J. V. 1987. Improving the status of endangered species in Missouri: least tern investigations. Missouri Department of Conservation endangered species project **number SE-01-12**.
- Smith, J. V. 1988. Improving **the** status of **tndntg**red species in **Missouri**: least tern investigations. Missouri Department of **Conservation** endangered species project number SE-01-12.
- Smith, J. V., and N. P. **Stucky**. 1988. **Habitat** management for **interior least terns**: problems and opportunities in inland **waterways**. Pages 134-149 **in** **M. C. Landin**, cd. **Inland Waterways**: Proceedings national workshp On the **beneficial uses** of dredged material. TRD-88-8. U. S. **Army** Engineers Vaternvays Experiment Station, Vicksburg, **Mississippi**.
- Smith, J. V., and R. B. Renktn. 1990. Improving **the** status of endangered species in Missouri: least **tern** investigations. **Final** report, Jobs 1 and 2, Missouri Department of Conservation endangered species project SE-01-19.
- Smith, K. L., and V. M. Shepherd. 1985. A **survey** of **the** interior least tern on the **Arkansas** and **Whitt** Rivers in Arkansas. Arkansas Natural Heritage Commission. Unpublished report. 5 pp.
- Smith, K. L. 1986. Results of the 1986 survey of **the** Arkansas River for interior **least** terns. Arkansas **National** Heritage Commission. Unpublished report.
- Smith, K. L., S. Barklty, and C. Gates. 1987. A **survey** of interior least terns **on** the Arkansas River in Arkansas. Arkansas Natural Heritage Commission. Unpublished report.
- Smith, M. 1986; Field **survey** of **the** interior least **tern** (**Sterna antillarum athalassos**) on the Mississippi River and Red River, Louisiana. U. S. **Army** Corps of Engineers, Vicksburg District, Vicksburg, Mississippi. Unpublished report.
- Soots, R. F., Jr., and J. F. Parntll. 1975. Ecological succession of breeding birds in relation to plant succession on dredge islands in North Carolina estuaries. North Carolina Sea Grant Publication **UNC-SG-75-27, Raleigh**. 91 pp.
- Stiles, B. 1939. **The** least tern in **Iowa**. **Iowa** Bird Life **14:18-21**.
- Stinnett, D. P., R. V. Smith, and S. V. **Conrady**. 1987. Riparian areas of vesttrn Oklahoma: a special study **of** their status, trends and values. U.S. Fish and Vildlift Service, Tulsa, Oklahoma. Unpublished report. **80** P P.
- Sweet, M. J.** 1985. **Least** tern population **survey**, 1984. Illinois Department of **Conservation**. Unpublished report. Svickard, D. K. **1972**. Status of the least tern at Camp Pendleton, **California**. **California** Birds 3(3):49-58.
- Svickard, D. K. 1974. An evaluation of **two** artificial least **tern** nesting **sites**. **California** Fish and **Game** **60(2):88-90**.

- Talent, L. G. and L. A. Hill. 1985. Final report: breeding ecology of snowy plovers, American avocets, and interior least terns at Salt Plains National Wildlife Refuge, **Oklahoma**. Oklahoma State University, Stillwater. 186 pp.
- Thompson, B. C. 1982. Distribution, **colony characteristics**, and population status of least terns breeding on the Texas coast. Ph.D. Dissertation. Texas **A&M** University.
- Thompson, B. C., and R. D. Slack. 1982. **Physical** aspects of colony selection by least **terns** on the Texas coast.. Colonial Waterbirds **5:161-168**.
- Thompson, B. C., and R. D. Slack. 1983. Post-fledging departure from colonies by juvenile least **terns** in Texas: implication for estimating production. Wilson Bulletin **96:309-313**.
- Tomkins**, I. R. 1959. Lift history notes on **the** least **tern**.. Wilson Bulletin **71:313-322**.
- Toups**, J. 1976. A brief history of efforts to protect **the** least tern on the Mississippi coast. Mississippi **Kite** **6:22-24**.
- U. S. Fish and Wildlife Service. 1980. California least **tern** recovery plan. U. S. Fish and Wildlife Service. Portland, Oregon. 58 pp.
- U. S. Fish and Wildlife Service.. 1985. Interior population of the least **tern** determined to be endangered. Federal Register **50:21784-21792**.
- U. S. Fish and **Wildlife** Service. **1987a**. Least **tern** in: Endangered species information system (computer data base). U. S. Department of **the** Interior, Fish and Wildlife **Service**, Division of Endangered Species and Habitat Conservation, Washington, D. C.
- U. S. Fish and Wildlife Service. 1987b. Biological opinion on **the** Platte River off-site effects **of the** Wyoming Water Development Commission's proposed Deer **Creek** Dam and Reservoir project, **Letter** from U. S. Fish and Wildlife Service, **Denver**, Colorado to U. S. Army Corps of Engineers, **Omaha**, Nebraska **dated July 20, 1987**.
- U. S. Fish and **Wildlife** Service. **1987c**. **Biological** opinion on the Platte River off-site effects of **the** Denver Water Department's proposed Two Forks Dam project. **Letter** from U. S. Fish and Wildlife **Service**, Denver, Colorado to U. S. Army Corps of Engineers, Omaha, Nebraska dated October 14, 1987.
- U. S. Fish and Wildlife **Service**. 1988. Great lakes and Northern Great Plains Piping Plover recovery plan. U. S. Fish and Wildlife **Service**, Twin Cities, **Minnesota**. 160 pp.
- U. S. Fish and Wildlife Service. 1989. Draft biological opinion on the effects of Keystone and Kav dam operations in Oklahoma.
- U. S. Fish and Wildlife Service. 1990. *Draft* biological opinion on the effects of Missouri River dam operations.
- Watson, S. R. 1966. **Seabirds** of the tropical Atlantic Ocean. Smithsonian Press, Washington, D. C. 230 pp.
- Whitman, P. L. 1988. Biology and conservation of **the** endangered interior least tern: a literature review. U. S. Fish and Wildlife **Service** Biological report **88(3). 22 pp**.
- Williams**, S. O; 1988. Status of the least **tern** in Ntv Mexico. Paper presented to joint meeting of Colonial Waterbird Group and Pacific **Seabird** Group, 12-16 October 1988, Washington, D. C. (abstract).
- Wilson, B. L. 1984. 1984 search for piping plover and least **tern** in **Iowa**. Unpublished report. 10 pp.

- Wilson, B. L. 1986. Special birds of Council Bluffs-1986. Unpublished report.
- Wilson, E. C., S. H. Anderson, and V. A. Hubert. 1989. **Evaluation** of habitat suitability criteria proposed in **Armbruster** (1986) for the interior least **tern** on the Platte River, Nebraska, and adjacent sand pits during the 1989 nesting season. U.S. Fish and Wildlife **Service**, Wyoming Cooperative Fish and Wildlife Research Unit, **Laramie**, Wyoming. Unpublished report.
- Volk, R. G. 1974. Reproductive behavior of the least **tern**. Proceedings of the Linnatan **Society**, New York **72:44-62**.
- Wycoff, R. 1960. The least tern, Nebraska Bird **Review** **28:39-42**.
- Youngworth, V. 1930. Breeding of the least **tern** in Iowa. Wilson Bulletin **42:102-103**.
- Youngworth, V. 1931. The American egret and least tern in South Dakota. Wilson Bulletin **43:309-310**.

III. IMPLEMENTATION

The **Implementation** Schedule outlines and gives **priorities** to tasks deemed necessary to be undertaken in the next three years to maximize recovery of the interior least tern. This process **will** be reviewed every **three** years until the recovery objective is **met**. Therefore, priorities and tasks may change in the **future**.

KEY TO IMPLEMENTATION SCHEDULE

General Category (Column 1):

Information and Research (I,R)

1. Population status
2. Habitat status
3. Habitat requirements
4. Management techniques
5. Taxonomy
6. Demographic studies
7. Propagation
8. Migration
9. Vinttring
10. Predation
11. Compttition
12. Disease
13. Environmental contaminant
14. Reintroduction
15. Other information

Acquisition - A

1. **Lease**
2. **Easement**
3. **Management** agreement
4. Exchange
5. Withdralval
6. Fee title
- 7: Other

Management - M

1. Propagation
2. Reintroduction
3. Habitat maintenance and manipulation
4. predator and . . . competitor control
5. Depredation control
 6. Disease control
7. Pollution control
8. Public **information**
9. Other information

Priority (column 4):

1. Those actions absolutely necessary to prevent extinction of **the species** in the foreseeable **future**.
2. Those actions **necessary** to maintain **the** species' current population status.
3. All other actions necessary to provide for full recovery of the species.

Agency Responsibility (column 6):

USFWS Regional Office 2 - **Albuquerque**
3 - **Twin Cities**
4 - **Atlanta**
6 - **Denver**

USFWS Research - 8

USFWS Office of Migratory Bird Management - **OMBM**

USFWS Office of International Affairs - **IA**

SA - State **Wildlife** Agency

BR - Bureau of Reclamation

COE - U. S. Army Corps of Engineers

NPS - National Park Service

WCHT - Platte **River Whooping** Crane Habitat **Maintenance Trust**

CW - Colonial Waterbirds

MO - Missouri River System

MS - Mississippi River System

AR - Arkansas River System

RE - Red **River System**

RG - Rio Grande River System

IMPLEMENTATION SCHEDULE

Complete Implementation Schedule for First Three Years of Recovery Effort

General Category	Task	Task #	Priority #	Task Duration	Responsibility		Fiscal Year		
					Region (USFWS)	Other Agencies	1	7	3
I1	Survey, census and monitor breeding population	111-114	2 (MO)	Annual	Regions 3,6	SA, COE	\$15K	\$15K	\$15K
			2 (MS)	Annual	Regions 3,4	SA, COE	\$15K	\$15K	\$15K
			2 (AR)	Annual	Regions 2,4	SA, COE	\$20K	\$20K	\$20K
			2 (RE)	Annual	Regions 2,4	SA, COE	\$ 5K	\$ 5K	\$ 5K
			2 (RG)	Annual	Region 2	SA	\$10K	\$10K	\$10K
16, R6	Assets mortality and identify life history parameters (including population modeling)	116-117	3 (MO)	Annual	Regions 3,6	SA, WCHT	\$10K	\$10K	\$10K
			3 (MS)	Annual	Regions 3,4	SA, COE	\$10K	\$10K	\$10K
			3 (AR)	Annual	Regions 2,4,6	SA, COE	\$10K	\$10K	\$10K
			3 (RE)	Annual	Region 2,4	SA, COE	\$10K	\$10K	\$10K
			3 (RG)	Annual	Region 2	SA	\$10K	\$10K	\$10K
R9, R1 R6	Survey and census winter populations	131-132	2	Annual	8, OMB, IA	CW	\$35K	\$35K	\$15K
I2, R3	Quantify and evaluate breeding habitat and threats	211-213	2 (MO)	2 years	Regions 3,6	SA, BR, WCHT	\$15K	\$10K	\$10K
			2 (MS)	2 years	Regions 3,4	SA, COE	\$15K	\$15K	\$15K
			2 (AR)	2 years	Regions 2,4,6	SA, COE	\$15K	\$15K	\$15K
			2 (RE)	2 years	Regions 2,4	SA, COE	\$ 5K	\$ 5K	\$ 5K
			2 (RG)	2 years	Region 2	SA	\$10K	\$10K	\$10K
M4, R10	Evaluate predator impacts; evaluate predator management techniques and implement	3111-3112	2 (MO)	Annual	Regions 3,6	SA, WCHT	\$15K	\$15K	\$10K
			2 (MS)	Annual	Regions 3,4	SA, COE	\$10K	\$10K	\$10K
			2 (AR)	Annual	Regions 2,4,6	SA, COE	\$15K	\$15K	\$15K
			2 (RE)	Annual	Regions 2,4	SA	\$ 5K	\$ 5K	\$ 5K
			2 (RG)	Annual	Region 2	SA	\$ 5K	\$ 5K	\$ 5K

IMPLEMENTATION SCHEDULE


Complete Implementation Schedule for First Three Years of Recovery Effort

General Category	Task	Task #	Priority #	Task -Duration	Responsibility		Fiscal Year		
					Region (USFWS)	Other Agencies	c	o	s
M8, M9	Restrict human and vehicular access to nesting areas	3113	2 (MO)	Annual	Regions 3,6	SA, COE	\$15K	\$15K	\$15K
			2 (MS)	Annual	Regions 3,4	SA	\$10K	\$10K	\$10K
			2 (AR)	Annual	Regions 2,4,6	SA, COE	\$15K	\$15K	\$15K
			2 (RE)	Annual	Regions 2,4	SA	\$5K	\$5K	\$5K
			2 (RG)	Annual	Region 2	SA	\$5K	\$5K	\$5K
M3, M9	Manage water levels to reduce nest and chick loss	3114	1 (MO)	Annual	Regions 3,6	COE	\$20K	\$20K	\$20K
			1 (MS)	Annual	Regions 3,4	COE	\$15K	\$15K	\$15K
			1 (AR)	Annual	Regions 2,4,6	COE, BR	\$10K	\$10K	\$10K
			1 (RE)	Annual	Regions 2,4	COE	\$5K	\$5K	\$5K
			1 (RG)	Annual	Region 2	COE	\$5K	\$5K	\$5K
12	Identify essential breeding habitat	411-412	2 (MO)	Ongoing	Regions 3,6	SA			
			2 (MS)	Ongoing	Regions 3,4	SA			
			2 (AR)	Ongoing	Regions 2,4,6	SA			
			2 (RE)	Annual	Regions 2,4	SA			
			2 (RG)	Annual	Region 2	SA			
M3	Establish liaison to protect breeding habitat	413	3 (MO)	Annual	Regions 3,6	SA, COE, BR			
			3 (MS)	Annual	Regions 3,4	SA, COE			
			3 (AR)	Annual	Regions 2,4,6	SA, COE, BR			
			3 (RE)	Annual	Regions 2,4	SA, COE			
			3 (RG)	Annual	Region 2	SA			
M9	Revise or establish laws to protect breeding habitat	414	3 (MO)	Annual	Regions 3,6	SA			
			3 (MS)	Annual	Regions 3,4	SA			
			3 (AR)	Annual	Regions 2,4,6	SA			
			3 (RE)	Annual	Regions 2,4	SA			
			3 (RG)	Annual	Region 2	SA			

INPLENENTATION SCHEDULE
Complete Implementation Schedule for First Three Years of Recovery Effort

General Category	Task	Task #	Priority #	Task Duration	Responsibility		Fiscal Year costs		
					Region (USFWS)	Other Agencies	1	2	3
R2, R3	Develop criteria and priorities for habitat protection	415	3 (HO)	1 year	Regions 3,6	SA			
			3 (MS)	1 year	Regions 3,4	SA			
			3 (AR)	1 year	Regions 2,4,6	SA			
			3 (RE)	1 year	Regions 2,4	SA			
			3 (RG)	1 year	Region 2	SA			
R3, M3 \$15K	Develop river management plans	416	1 (MO)	Annual	Region 6	SA, COE, WCHT	\$15K	\$15K	
\$10K			1 (MS)	Annual	Region 4	SA, COB	\$10K	\$10K	
\$10K			1 (AR)	Annual-	Regions 2,4,6	SA, COE, BR	\$10K	\$10K	\$10K
5K			1 (RE)	Annual	Regions 2,4	SA, COB	\$5K	\$5K	\$5K
5K			1 (RG)	Annual	Region 2	SA, COE, BR	\$5K	\$5K	\$5K
R1, R2 \$25K	Determine effects of river hydraulics and sediment discharge on breeding habitat; identify flow regimes to protect habitat	4161-4162	1 (HO)	Annual	Region 6	SA, COE, BR	\$25K	\$25K	
\$20K			1 (US)	Annual	Region 4	WCHT SA, COE	\$20K	\$20K	
\$20K			1 (AR)	Annual	Region 2,6	SA, COE, BR	\$20K	\$20K	
\$10K			1 (RE)	Annual	Region 2	SA, COE	\$10K	\$10K	
\$10K			1 (RG)	Annual	Region 2	SA, COE	\$10K	\$10K	
R3 \$10K	Determine relationship of existing artificial breeding	4163	2 (MO)	2 years	Region 6	SA	\$10K	\$10K	
			2 (MS)	3 years	Region 4	SA, COE	\$10K	\$10K	

\$10K								
\$10K	sites to riverine sites		2 (AR)	2 years	Regions 2,6	SA	\$10K	\$10K
5K			2 (RE)	2 years	Region 2	SA	\$ 5K	\$ 5K \$
5K			2 (RG)	2 years	Region 2	SA	\$ 5K	\$ 5K \$
M3	Modify and/or eliminate	418	2 (MO)	Annual	Regions 3,6	SA, COE	\$ 5K	\$ 5K \$
5K	construction activities that		2 (MS)	Annual	Regions 3,4	SA, COE	\$ 5K	\$ 5K \$
SK	impact breeding habitat		2 (AR)	Annual	Regions 2,4,6	SA, COE	\$ 5K	\$ 5K \$
5K			2 (RE)	Annual	Regions 2,4	SA, COE	\$ 5K	\$ 5K \$
SK			2 (RG)	Annual	Region 2	SA, COB	\$ 5K	\$ 5K \$
5K								



IMPLEMENTATION SCHEDULE

Complete Implementation Schedule for First Three Years of Recovery Effort

General Category	Task	Task #	Priority #	Task Duration	Responsibility		Fiscal Year		
					Region (USFWS)	Other Agencies	1	2	3
M8	Inform and educate the public	511-513	2 (MO)	Annual	Regions 3,6	SA, COE	\$ 5K	\$ 5K	\$
5K			2 (MS)	Annual	Regions 3,4	SA, COE	\$ SK	\$ SK	\$
SK			2 (AR)	Annual	Regions 2,4,6	SA, COE, BR	\$ 5K	\$ 5K	\$
5K			2 (RE)	Annual	Regions 2,4	SA, COE	\$ 5K	\$ SK	\$
5K			2 (RG)	Annual	Region 2	SA, COE	\$ SK	\$ 5K	\$
SK	Inform and educate public resource management agencies	52	3 (MO)	Annual	Regions 3,6	SA, COE			
M8, M9			3 (MS)	Annual	Regions 3,4	SA, COE			
			3 (AR)	Annual	Regions 2,4,6	SA, COE			
			3 (RE)	Annual	Region 2	SA, COE			
			3 (RG)	Annual	Region 2	SA, COE			
M9	Coordinate recovery efforts	61	2	Annual	Regions 2,4,6	SA, COE			

APPENDIX1

contact People .

The following **individuals** have offered to provide **interested** parties with information pertaining to interior least terns in their **area**.

Roger Boyd
Biology Department
Baker University
Baldwin City, **Kansas**
913/594-6451

Dennis Christopherson
U. S. Fish and Wildlife **Service**
1501 14 St. West, Suite 230
Billings, MT 59102
406/657-6028

Mark Dryer or Paul Mayer
U. S. Fish and Wildlife Service
1500 Capitol Avenue
Bismarck, North Dakota 58501
701/255-4491

Paul B. **Hamel**
Tennessee Department of **Conservation**
701 Broadway
Nashville, Tennessee 37219-5237
615/742-6546

Laura A. Hill
U. S. Fish and **Wildlife** Service
222 South Houston, Suite A
Tulsa, Oklahoma 74127
918/581-7458

Gary R. **Lingle**
Platte River Whooping Crane Habitat Maintenance Trust
2550 N. **Diers** Ave.
Grand Island, Nebraska 68803
308/384-4663

Ross Lock
Nebraska **Game and** Parks Commission
P. O. Box 30370
Lincoln, Nebraska 68503
402/471-5438

Ren Lohofner
U. S. Fish and Wildlife Service
300 **Woodrow** Wilson, Suite 316
Jackson, US 39213
601-965-4900

Elizabeth N. **McPhillips**
U. S. Fish and Wildlife Service
Federal Building, Room 227
225 South Pierre
Pierre, South Dakota 57501
605/224-8693

Rochelle B. Renken
Fish and Wildlife Research Center
Missouri Department **of** Conservation
1110 S. College Avenue
Columbia, Missouri 65201
314/882-9880

John P. **Rumancik**, 'Jr.
U. S. Army Corps of Engineers
B-202 Clifford Davis Federal Building
Memphis, Tennessee 38103-1894
901/521-3857

Marvin Schvilling
Kansas Department of Wildlife and Parks
1407 College Drive
Emporia, Kansas 66801
316/342-1985

Kenneth Smith
Arkansas Natural Heritage Inventory
225 East **Markham**, Suite 200
Little Rock, **Arkansas** 72201
501/371/1706

Sartor O. Williams, III
Endangered **Species** Program
New Mexico Department of **Game** and Fish
State Capitol, Santa Fe, New **Mexico** 87503
505/827-9914

APPENDIX 2

Agreements Necessary For Protection Of **Essential Habitat**

1. Memorandum **of** Understanding should be developed between the U. S. Army Corps **of** Engineers, National Park **Service**, U. S. Fish and Wildlife **Service**, and the State wildlife agency, **for** permanent protection and management (vegetation clearing, law enforcement, public relations, etc.) of all essential habitat on the **Missouri** River in North Dakota, South Dakota, and Nebraska.
2. U. S. Fish and Wildlife **Service**, National Park **Service**, and U. S. Army Corps of Engineers should acquire easements and/or fee title of essential interior least tern habitat on the Missouri River in North Dakota, South Dakota, and Nebraska.
3. Memorandum of Understanding should be developed between the U. S. Army Corps of Engineers, U. S. Bureau of Reclamation, U. S. Fish and Wildlife Service, Platte River Whooping Crane Habitat Maintenance Trust, and the state wildlife agency, **for** the permanent protection and management of all essential habitat on the Platte River system **in** Nebraska.
4. The U. S. **Fish** and Wildlife Service should provide land protection of essential interior least **tern** habitat on the Platte River system.
5. Memorandum of Understanding should be developed **between** the U. S. Army Corps of Engineers, State natural resource **agency**, and the U. S. Fish and Wildlife Service **for the permanent** protection and management of essential habitat on the Mississippi and Ohio Rivers.
6. Memorandum of Understanding should be developed between the U. S. Fish and Wildlife Service, State wildlife agency, and the U. S. Army Corps of Engineers governing **the deposition** of dredge spoils on the **Mississippi** and Ohio Rivers for purposes **of** enhancing or creating interior least tern habitat.
7. Memorandum of Understanding should be developed between the U.S. Army Corps of Engineers, U.S. **Bureau of** Reclamation, U.S. Fish and Wildlife **Service**, U. S. Section **of** the International Boundary and Water Commission, State wildlife agencies, and appropriate agencies in Mexico for permanent protection and management of **all essential** habitat **in the Arkansas, Red, and Rio Grande Rivers basins in Kansas, Oklahoma, Arkansas, and Texas.**
8. U.S. Fish and Wildlife Service, U.S. **Army** Corps of Engineers, U.S. Bureau of Reclamation, and The Nature **Conservancy** should acquire easements and/or fee title **of** essential interior least tern habitat in the Arkansas, Red, and Rio Grande river **basins in Kansas, Oklahoma, Arkansas, and Texas.**

9. **Memorandum** of Understanding should be developed between the U.S. Fish and Wildlife **Service**, State wildlife agencies, and the U.S. Army Corps of Engineers governing removal and deposition of dredge spoil from the McClellan-Kerr Arkansas River Navigation System, in Oklahoma and Arkansas, for purposes **of** enhancing or creating least tern habitat.

Appendix 3. **Example** of a **memorandum of understanding**

MEMORANDUM OF UNDERSTANDING

The Nature Conservancy
U.S. **Army** Corps of Engineers
Oklahoma Department of Wildlife Conservation
U.S. Fish and Wildlife Service
Tulsa Audubon Society
River **Parks Authority**

WHEREAS _____ an Oklahoma corporation, ("**Owner**") has acquired certain lands and riverbeds on the Arkansas River floodplain **in** Tulsa County, Oklahoma, as more particularly shown on the plat attached hereto as Exhibit A (the "Property"); and

WHEREAS said Property has special value for wildlife including nesting populations of the endangered Interior **Least Tern**, ***Sterna antillarum*** **athalassos**; and

WHEREAS The Nature Conservancy ("**Conservancy**"), a private, nonprofit organization committed to the conservation and management **of** rare and endangered species, **communities**, and ecosystems, has **expressed an** interest to coordinate the efforts of local, state, and federal agencies in protecting the Least Tern; and

WHEREAS The United States **Army** Corps of Engineers ("Corps") has certain water management responsibilities on the Arkansas River that might affect the habitat of the **Least Tern**; and

WHEREAS the U.S. Fish and Wildlife Service, ("USFWS") *has federal* management **responsibilities over** federally-listed endangered species such as the Least Tern, and the Oklahoma **Department** of Wildlife Conservation ("**ODWC**") has state management responsibilities over state-listed endangered species such as the Least Tern; and

WHEREAS the Tulsa Audubon Society ("**TAS**"), a private, nonprofit organization, has expertise in the preservation of birds such as the Least Tern; and

WHEREAS the River Parks Authority ("**RPA**") is a **public** trust charged **with** the responsibility of protecting and enhancing **interalia**, natural **communities** and species along the Arkansas River and its environment in Tulsa County, Oklahoma.

WHEREAS the Owner, ODWC, **USFWS**, **Conservancy**, **TAS**, the Corps and RPA all have an interest in protecting nesting populations **of the** rare and endangered Interior Least Tern on the Arkansas River; and

WHEREAS The Owner is agreeable **to** manage jointly these lands to protect the Least Tern.

NOW THEREFORE, the Owner hereby grants to The River Parks Authority, an

exclusive license and permit, consisting of the following rights for the purposes described, in and to the lands described in Exhibit A attached hereto and made a part hereof, to-wit:

RIGHTS GRANTED TO THE RIVER PARKS AUTHORITY

1. The River Parks Authority shall have the right to enter upon • nd.uae said lands for the purpose of protecting all **Least** Tern nesting, fledging, feeding, resting and cover sites, located on said property. Said purposes shall include but not be limited to inspection, monitoring, research and, if deemed necessary, manipulation **of the** sites to enhance the Least Tern population. The River Parka Authority, upon consultation with the **USFWS**, may authorize personnel from the Corps, USPWS, ODWC, **TAS**, the Conservancy and others to enter said lands for the purposes described herein. Such consultation is necessary to alleviate potential for violations of the Endangered Species **Act**.
2. The River Parks Authority shall have the right to control and limit access to Least Tern nesting sites in breeding season, as necessary, and to erect and place any signs, posters, or other &vices to identify the land as a protected area.

SAID RIGHTS ARE SUBJECT TO THE **FOLLOWING LIMITATION**, HOWEVER:

1. No one will construct facilities on said premises nor modify the land surface or habitat thereon until a proposal thereof has been reviewed and approved by USFWS and Owner.
2. All existing **RPA** regulations (e.g., **no vehicle**, dogs on leash, **curfew** clauses) will apply.

OBLIGATIONS OF RIVER PARKS AUTHORITY

AS PARTIAL CONSIDERATION for the rights hereby granted by the Owner, **RPA** agrees to:

Solicit expert advice regarding the protection, management and enhancement of the Least Tern population on the lands **from the** agencies and organizations that are party to this agreement and from other sources available to it, and shall exercise its best efforts to implement said recommendations consistent with the terms of this **agreement**.

OBLIGATIONS OF THE OWNER

THE OWNER agrees that:

1. In its planning and use **of said lands**, it shall, whenever practicable, take into consideration protection **of** said preserve area for endangered bird species.
2. It shall exercise its best efforts to implement recommendations of the River Parks Authority.

GENERAL PROVISIONS

1. Neither Owner nor any other party to this agreement is required to

obligate or spend funds under this agreement, it being the intent of the parties that staff time and expertise be the primary contribution of each party to the effective implementation of this Agreement.

- 2. **This** permit may be terminated, in whole or in part, by the Owner or by the River Parks Authority upon 90 days written notice to the other party.
- 3. All notices required under this agreement shall be *effective* when mailed to the following persons:

To Owner:

To River Parks Authority:

Jackie Bubenik, Executive Director
 River Parks Authority
 707 South Houston, Suite **202**
 Tulsa, Oklahoma 74127

- 4. By their signatures hereto, the Corps, **USFWS**, ODWC, **TAS**, and the Conservancy agree to assist the **Owner and** The River Parks Authority by providing expertise and assistance toward the common goal of protecting, managing, and enhancing the Least Tern population on the lands described.

IN WITNESS WHEREOF, the parties hereto have subscribed their names as of the dates indicated:

By: _____
 Its: _____

Dated: _____

THE NATURE CONSERVANCY

Dated: _____

Attest:

By: _____
 Its Vice President

By: _____
 Its Assistant Secretary

U.S. **ARMY** CORPS OF *ENGINEERS*

Dated: _____

By: _____
 Its: _____

OKLAHOMA **DEPARTMENT OF WILDLIFE** CONSERVATION

By: _____
 Its: _____

Dated: _____

U.S. FISH AND WILDLIFE SERVICE

By: _____
 Its: _____

Dated: _____

TULSA AUDUBON SOCIETY

Dated: _____

A t t e s t :

By: _____
I t s : _____

RIVER PARKS AUTHORITY

By: _____
I t s : _____

Dated: _____

At t e s t : _____

By: _____
I t s : _____

Dated: _____

APPENDIX 4

Essential Breeding Habitat for Interior Least Terns

Riverine sandbars, river channel environment including open channel area, channel width, and appropriate **instream** flows, and lake shorelines and other habitats provide essential habitat for the interior least tern. The interior least tern is completely dependent on these habitats **for food** and nesting sites. Therefore, destruction or adverse modification of remaining habitats will cause continued reduction of the species range and eventually a reduction in population numbers. The areas described and mapped herein as essential habitat will provide the space necessary for continued existence and growth of interior least tern populations required to meet the recovery objective. The following maps depict essential habitat for the interior least tern. Hatch marks along river segments and certain national wildlife refuges indicate **the** areas where essential habitat intermittently occurs depending on water conditions. For example, sandbars and interior least terns do not **occur** along every kilometer of the indicated segments of rivers. Locations of nesting birds may change from year to **year** within the indicated segment.

I. Missouri River System

Montana • Missouri River between Fort Peck Dam and North Dakota

North Dakota • Yellowstone River and Missouri River between Garrison Dam and the Cannonball **River**.

South Dakota • Cheyenne River **from** the Belle Fourche River to Lake Oahe; **Missouri** River *from* Ft. Randall Dam to mouth of the Niobrara River and from Gavin's Pt. Dam to Ponca, Nebraska.

Nebraska • Missouri River from South **Dakota** to mouth of the Niobrara River **and** from Gavin's Pt. Dam to Ponca; Niobrara River **from** Highway 183 bridge to Missouri River; **Loup** River from St. Paul to Platte River; Platte River from Laxington to Chapman and from **Columbus** (Highway 81 bridge to, **Missouri** River.

II. Mississippi River • From Highway **146** bridge, **Missouri** and Illinois to Vicksburg, **Mississippi**

III. Arkansas River system

Kansas • Quivira National Wildlife Refuge and Cimarron River


Oklahoma • Salt Plains National Wildlife Refuge; from below **Kaw** Dam to Arkansas River and Arkansas River from Tulsa to Muskogee;


Cimarron River in Beaver, Harper, Woods, Woodward, Major, Kingfisher, Logan, and Payne counties; Canadian River in Ellis, Roger Mills, Dewey, Cleveland, **McClain**, Haskell, Pittsburgh, Hughes, Muskogee, and Sequoyah counties; Sequoyah National Wildlife Refuge;

Red River from Harmon county to Highway 277/281 bridge.


Texas • Canadian River *from* Sanford Dam *to* Oklahoma: Prairie Dog Town Fork/Red River from **Briscoe/Armstrong county** boundary to Burkburnett, Texas.

IV. **Pecos River** • Bitter Lake National Wildlife Refuge., **New Mexico.**






Essential Habitat in Montana:
Indicated Segments of the Missouri River





Essential Habitat in North Dakota:
Indicated Segments of the Missouri and Yellowstone Rivers



Essential Habitat in South Dakota:
Indicated Segments of the Cheyenne and Missouri Rivers


Essential Habitat in Nebraska: Indicated Segments of the Loup, Missouri, Niobrara, Missouri and Platte Rivers






MISSOURI
Indicated Segment of the Mississippi River

**Essential Habitat in Kentucky and Tennessee:
Indicated Segments of the Mississippi River**







Essential Habitat in Arkansas:
Indicated by thick line.


Essential Habitat in Mississippi

Indicated Segment of the Mississippi River






Essential Habitat in Louisiana:
Indicated Segment of the Mississippi River




Essential Habitats in Illinois:
Indicated Segment of the Mississippi River




Essential Habitat in Oklahoma:

Sequoyah & Salt Plains National Wildlife Refuge

Indicated Segments of the Arkansas, Canadian, Cimarron and Red Rivers



Essential Habitat in Texas:
Indicated Segments of the Canadian, Prairie Dog Town Fork and Red Rivers



Essential Habitat in Kansas:
Quivira National Wildlife Refuge and Cimarron River

Albuquerque .

PECOS
RIVER

Roswell . Bitter Lake
National Wild
Refuge

Carlsbad .

E s s e n t i a l · **H a b i t a t** · i n N e w M e x i c o :

Bitter Lake National Wildlife Refuge

Appendix 5

LIST OF **REVIEWERS**

Mr. Sam Barklty
Endangered Species Coordinator
Arkansas Game and Fish Commission
No. 2 Natural Resources Drive
Little Rock, Arkansas 72205

Dr. Dean Roosa
Iowa Department of Natural Resources
Wallace State Office Building
Des Moines, Iowa 50319

Ms. Susan **Lauzon**
Endangered Species Coordinator
Illinois DOC
Lincoln Tower Plaza
525 south Second Street
Springfield, Illinois 62706

Mr. Chris Iverson
Endangered Species Coordinator
Indiana DNR
608 State Office Building
Indianapolis, Indiana 46204

Mr. Marvin D. Schwilling
Kansas Fish and Game Commission
Box **54A**, Route 2
Pratt, Kansas 67124

Ms. Lynda **J. Andrews**
Kentucky Dept. of Fish & Wildlife
Resources
1 Game **Farm** Road
Frankfort, Kentucky 40601

Mr. Gary Lester
Louisiana Dept. of Wildlife
and Fisheries
P. O. Box 15570
Baton Rouge, Louisiana **70895**

Dr. John W. Smith
Missouri Department of Conservation
Fish and Wildlife Research Center
1100 college Avenue
Columbia, Missouri 65201

Mr. John P. **Rumancik** Jr. .
Department of Army
Corps of Engineers
B-202, Clifford David
Federal Building
Memphis, Tennessee 38103

Dr. Bruce C. Thompson
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, Texas 78744

Mr. Gary R. Linglt
Platte River Whooping Crane Trust
2550 North Diers **Avenue**, Suite **H**
Grand Island, Nebraska 68803

Mr. Ross Lock
Nebraska Game and Parks Commission
2200 North 33rd Street
P.O. Box 30370
Lincoln, Nebraska 68503

Mr. Clyde P. **Gates**
Department of the Army
Corps of Engineer8
Box 867
Little Rock, Arkansas **72203-0867**

Dr. Mary C. **Landin**
Waterways Experiment Station
Department of the Army
Corps of Engineers
Box 631
Vicksburg, Mississippi **39180-0631Mr.**

Hr. Paul **Hamel**
Tennessee Department of Conservation
701 Broadway
Nashville, Tennessee **37219**

Mr. **Ken** L. Smith
Arkansas Natural Heritage Inventory
225 **E.** Markham, Suite 200
Little Rock, Arkansas 72201

Mr. Gary Williams
Engineering and Research Center
Bureau of Reclamation
P.O. Box 25007
Building 67, Denver Federal Center
Denver, Colorado **80225-0007**

Dr. Stephen J. Chaplin
The Nature Conservancy
Midwest Regional Office
1313 Fifth Street S.E.
Minneapolis, Minnesota 55414

Mr. Robert D. Brown
Chief, Planning Division
Department **of** the Army
Corps of Engineers
P.O. Box 61
Tulsa, Oklahoma 74121-0061

Mr. Eugene Buglewicz
Environmental Analysis Branch
Department of the Army
Corps of **Engineers**
P.O. Box 80
Vicksburg, **Mississippi 39180-0080**

Mr. C. Gregory Schmitt
Wildlife Scientist
New Mexico Dept. of **Game** and Fish
State Capitol
Santa **Fe**, new Mexico 87503

James W. Flynn, Director
Montana Dept. of **Fish, Wildlife,**
Parks
Helena, Montana 59601

Dr. Brainard Palmer-Ball, Jr.
Kentucky Nature Preserves Commission
407 Broadway
Frankfort, Kentucky 40601

Dr. James H. Wilson
Mr. Michael Sweet
Missouri Department of Conservation
P.O. Box 180
Jefferson City, Missouri 65102

Mr. Robert M. Hatcher
Endangered Species Coordinator
Tennessee Wildlife Resources Agency
Ellington Agricultural Center
P.O. Box 40747
Nashville, Tennessee 37204

Mr. Dale L. Henegar, Commissioner.
North Dakota Game & Fish Dept.
100 N. Bismarck Expressway
Bismarck, North Dakota **58501-5095**

Mr. **William** Quisenberry
Mississippi Dept. of Uildlift
Conservation
P.O. Box **451**
Jackson, **Mississippi 39205-0451**

Mr. **Jim** Salyer
Wildlife Division Director
South Dakota Dept of
Game Fish & Parks
Sigurd Anderson Building
445 East Capitol
Pierre, South Dakota **57501-3185**

Mr. Charles D. Travis
Executive Director
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, Texas 78744

Mr. Steven Alan Lewis, Director .
Oklahoma Dept. of **Wildlife**
Conservation
1801 **North** Lincoln
Oklahoma City, Oklahoma 73105

Dr. Roger L. Boyd
Baker University
Baldwin City, Kansas **66006**

Mr. Gary **Willson**
Endangered Species Coordinator
Midwest Region-National Park **Service**
1709 Jackson Street
Omaha, Nebraska **68102-2571**

Mr. Conrad J. Keyes, Jr.
Principal Engineer, Planning
International Boundary and Water
Commission, United States and Mexico
The Commons, Building C, Suite 310
4171 North Mesa Street
El Paso, **Texas** 79902

Mr. Jot D. Kramer, Chief
Fisheries and Uildlift Division
Kansas Wildlife and Parks
RR 2, Box 54A
Pratt, **Kansas 67124**

Mr. John J. **Dinan**
Nebraska Game and Parks Commission
P.O. Box 30370
Lincoln, Nebraska 86503

Mr. Raymond E. Pettijohn
P.O. Box 46
Cedar Creek, Nebraska 68016

Mr. Gerald E. Jasmer
State Wildlife Biologist
Soil Conservation Service
Federal Building, Room 345
100 Centennial Mall North
Lincoln, Nebraska **68508-3866**

Mr. William **M.** Shepherd
Arkansas Natural Heritage
Commission
The Heritage Center, Suite 200
225 East **markham**
Little Rock, Arkansas 72201

Ms. Eileen **Dowd**
South Dakota **Natural** Heritage
South Dakota Dept. of Game, Fish &
Parks
445 East Capitol **Avenue**
Pierre, South Dakota **57501-3185**

Mr. Lloyd A. Jones
Commissioner
North Dakota Game & Fish Dept.
100 North Bismarck Expressway
Bismarck,, North Dakota **58501-5095**

Mr. Noel Caldwell
Planning Division
Lower Mississippi Valley Division
Dept. of the Army
Corps of Engineers
P.O. Box 80
Vicksburg, Mississippi **39180-0080**

Mr. Robert L. Jenkins
National Aquarium in Baltimore
Pier 3, 501 E. Pratt St.
Baltimore, Maryland 21202

Mr. William R. Ross
City **Manager**
P.O. Box 176
Yankton, South Dakota 57078

Mr. Michael Bean
Environmental Defense Fund
1616 P Street NW
Washington, DC 20036