

Draft Report

Environmental Contaminants in Water, Sediment and Biological Samples from Playa Lakes in Southeastern New Mexico - 1992

Environmental Contaminants Program

U.S. Fish and Wildlife Service
New Mexico Ecological Services Field Office
2105 Osuna NE
Albuquerque, New Mexico 87113

October 1998

Report Prepared by: Sky Bristol

Region 2 Environmental Contaminants Program
500 Gold Ave. SW
Albuquerque, New Mexico 87102

Abbreviations and Conversions

Abbreviations

liter	l
milliliter	ml
kilogram	kg
gram	gparts
per million	ppm
parts per billion	ppb
parts per trillion	ppt
milligram per kilogram	mg/kg
micrograms per gram	mcg/g
micrograms per milliliter	mcg/ml
micrograms per liter	mcg/l
nanograms per liter	ng/l

Conversion Factors

micrograms per gram	ppm
micrograms per milliliter	ppm
milligrams per kilogram	ppm
micrograms per liter	ppb
nanograms per liter	ppt

Conversions

$$\text{Wet weight} = (\text{dry weight})(1 - (\text{percent moisture}/100))$$

$$\text{Dry weight} = (\text{wet weight})/(1 - (\text{percent moisture}/100))$$

Table of Contents

Abbreviations and Conversions	-ii-
Table of Contents	-iii-
List of Tables	-iv-
Acknowledgments	-v-
Abstract	-vi-
Introduction	-3-
Study Area	-5-
Laguna Gatuna	-6-
Laguna Quattro	-6-
Laguna Tres	-6-
Laguna Uno	-6-
Laguna Walden	-7-
Lane Salt Lake	-7-
Middle Lake and Middle Lake Salt Playa	-7-
William's Sink	-7-
Sample Collection Methods	-7-
Water	-8-
Sediment	-8-
Invertebrates	-8-
Bird Tissue	-8-
Results	-8-
Contaminants in Water	-8-
Contaminants in Sediment	-9-
Contaminants in Aquatic Biota	-15-
Conclusions and Recommendations	-15-
Literature Cited	-18-

List of Tables

Table 1.	Sampling Sites , Analyses Performed, Matrices Collected, Date of Collection, and Current and Historic Disposal Status of Ten Sampling Sites in Southeastern New Mexico	-5-
Table 2.	Total PAH (ppm) in water samples from 8 playa sampling sites	-9-
Table 3.	Analytical Results for BTEX in 7 Sediment Samples (ug/g wet weight)	-9-
Table 4.	Percent total organic carbon in samples from 8 playa lakes	-11-
Table 5.	Total PAHs in sediment samples from 9 playa lakes	-12-
Table 6.	Criteria used to evaluate the potential hazard of trace elements in sediments and potential food items to birds and other organisms, expressed as mg/kg wet or dry weight (ww or dw; NA = not available).	-13-

Acknowledgments

Studies contributing to this report were conducted by Rick Roy, Tom O'Brien and other employees of the U.S. Fish and Wildlife Service. Special thanks to Rob Lee and Nick Chavez for help in the follow up observations of the playas and review of report materials.

Abstract

Sediment, water, bird tissue, and invertebrates were collected from 10 playa lakes in Southeastern New Mexico in 1991 and 1992. These samples were analyzed for a variety of constituents, including metals (trace elements, metals, and metalloids), polycyclic aromatic hydrocarbons (PAHs), alkanes, total organic carbon, BTEX (benzene, toluene, ethyl-benzene, p & m-xylene, and o-xylene), and specific cations and anions. Analytical results were compared to data available in the literature. Concentrations of most contaminants were less than water and sediment quality criteria, typical background concentrations, and avian dietary thresholds of adverse effects. Aluminum, cadmium, magnesium, mercury, selenium, vanadium, and zinc were elevated above dietary threshold concentrations in invertebrate and sediment samples at some or all of the sites where these constituents were analyzed. However, these constituents were not unilaterally elevated in bird tissues from the same sites, and the elevated concentrations of metals may reflect geologic sources in the region. In addition, total PAHs, including flouranthene, benzo[a]pyrene, benzo[g,h,i]perylene, benzo[b]flouranthene, benzo[k]flouranthene, and indeno[1,2,3-cd]pyrene, in several water samples were elevated above drinking water criteria for human health.

Introduction

Playas in southeast New Mexico provide wintering, breeding, feeding, and loafing habitat for many species of shorebirds, wading birds, waterfowl, and birds of prey. Over 120 species of migratory birds have been documented at playas in the more than 1,700 playas in the New Mexico playa lakes region, and as many as 15 million birds may migrate through the area each year with a third stopping to winter or feed. Playa lakes usually have no external drainage and often develop clay-rich bottom sediments which limit groundwater seepage or upwelling. Most water loss is therefore due to evapotranspiration, which can concentrate natural salts and contaminants, and in some basins, groundwater recharge. Depending on the size of the playa, the size of its watershed, rainfall, and temperature, a playa may hold water year-round or only for a few weeks.

The playas are also used as convenient disposal basins for produced water from oil and gas extraction activities and from potash mines. At present time, these industrial discharges are not regulated through the National Pollutant Discharge Elimination System (NPDES) administered by the Environmental Protection Agency. However, the disposal of produced water is regulated by the New Mexico Oil Conservation Division, and potash mine disposal activities are regulated by the New Mexico Environment Department, Ground Water Section. Neither agency requires the permittees to screen for EPA Priority Pollutants. These waters are classified by the New Mexico Water Quality Control Commission for the use of livestock and wildlife watering. Protection of this use by the State of New Mexico has not yet been enforced.

A significant number of migratory waterfowl were found dead during the fall and winter of 1992-93 on the shoreline of Laguna Toston. This playa has received slurry waste directly from a potash refining facility for nearly 40 years. During surveys of Laguna Toston in 1997, several whole migratory birds

were found dead on the shoreline and evidence of many more deaths ("feather spots") were observed on or near the shoreline. Migratory bird deaths have also been recorded at Laguna Gatuna and Laguna Quattro. Laguna Gatuna receives no potash disposal waters but has historically received oil brine disposal, and Laguna Quattro receives potash disposal directly through a chain of playas. Playas that receive produced water and/or potash mine brine support few if any shorebirds and invertebrates such as brine shrimp (Artemia spp.). Necropsies conducted by the Madison Wildlife Health Research Center in 1992 determined the cause of mortality of birds collected from these playa lakes as "probable salt toxicosis." In addition, a study conducted by the U.S. Geological Survey Biological Resources Division in 1994-95 (Dein et. al., 1997) concluded that the cause of death of migratory birds found in the study area along Laguna Toston was salt/sodium toxicity - water deprivation/dehydration.

Several samples of sediment, water, invertebrates and bird tissue were collected by the U.S. Fish and Wildlife Service in 1992 and analyzed for a variety of potential contaminants. The data collected in 1992 for this study was never compiled into a written report or analyzed for significance. The objective of this report is to summarize the findings of the study conducted in 1991-92 and determine if contaminants such as metals or PAHs exist in the playa lakes at levels posing potential risk to migratory birds and other wildlife. The results of this study may then be used in conjunction with the 1997 report by Dien and others to assess the impacts of potash waste disposal on the playa lakes and determine appropriate remedial and mitigative response.

The broad sampling scheme was adopted with the study began in 1992 and intended to survey a large area for potential contaminant problems. Acknowledged limitations to this study include: 1) the low sample size prevented statistical inferences about local population level effects; 2) the single sampling date for most sites represents only a single period in time; 3) for water and sediments, a single sampling location

or composite sample would not detect a heterogeneous distribution of contaminants; and 4) water, sediment, birds, and potential bird food items were not all sampled at every site, so little can be inferred about contaminant sources, pathways, or bioaccumulation pathways. Nonetheless, this study provides a screening-level analysis to focus future research on specific contaminants at identified locations.

Study Area

The playa lakes region of New Mexico is found in the southeast portion of the state near the city of Carlsbad. Data from ten different sampling sites is included in this report. Table 1 lists the 10 sampling sites and provides information on the sampling matrices collected, analytical constituents and anthropogenic discharge status.

Table 1. Sampling Sites , Analyses Performed, Matrices Collected, Date of Collection, and Current and Historic Disposal Status of Ten Sampling Sites in Southeastern New Mexico

Location	Analyses	Matrices	Date	Discharges
Laguna Gatuna	BTEX/PAH/TOC; Cat/An	Sediment; Water; Salt Encrustations	4/2/92	historic oil brine
Laguna Quattro	BTEX/PAH/TOC; Cat/An; Metals	Sediment; Water; Bird Liver/Kidney	5/6/92	indirect potash, historic oil brine
Laguna Tres	BTEX/PAH/TOC; Cat/An	Sediment; Water	5/6/92	indirect potash
Laguna Uno	BTEX/PAH/TOC; Cat/An	Sediment; Water	5/6/92	direct potash
Laguna Waldon	BTEX/PAH/TOC; Cat/An; Metals	Sediment; Water; Bird Liver; Invertebrate	5/6/92	oil brine
Lane Salt Lake	BTEX/PAH/TOC; Cat/An	Sediment; Water	4/15/92	oil brine
Lane Salt Lake (Burro)	BTEX/PAH/TOC; Cat/An; Metals	Sediment; Water; Bird Liver	4/15/92	oil brine
Middle Lake	Cat/An; Metals	Sediment; Bird Liver; Invertebrate	5/15/92	oil brine
Middle Lake Salt Playa	BTEX/PAH/TOC	Sediment; Water	4/15/92	oil brine
William's Sink	BTEX/PAH; Cat/An; Metals	Sediment; Water; Invertebrate	4/2/92	historic potash

Laguna Gatuna

Laguna Gatuna is located about 35 miles east of Carlsbad, New Mexico and just north of U.S. Highway 162. The playa has received historic oil brine disposal from several adjacent oil pumping operations but has received no direct potash waste disposal. However, it is possible that Laguna Gatuna may receive some indirect potash disposal water through groundwater from Laguna Toston, located approximately 4 miles upgradient.

Laguna Quattro

Laguna Quattro is located approximately 40 miles southeast of Carlsbad, New Mexico and just south of N.M. Highway 128. The playa receives potash waste disposal from IMC Corporation's processing facility at high water levels through a series of playas beginning with Laguna Uno. Laguna Quattro historically received illegal oil brine discharge directly from an operation by B&E Corporation. That operation was shut down in 1993.

Laguna Tres

Laguna Tres is located approximately 40 miles southeast of Carlsbad, New Mexico and just north of N.M. Highway 128. The playa may receives potash waste disposal from IMC Corporation's processing facility indirectly at high water levels through a series of playas beginning with Laguna Uno.

Laguna Uno

Laguna Uno is located approximately 40 miles southeast of Carlsbad, New Mexico and approximately 2 miles north of N.M. Highway 128. The playa receives direct potash waste disposal from IMC Corporation's processing facility. This water runs over at high levels into adjacent Lindsay Lake and then through a series of playas including Lagunas Tres and Quattro.

Laguna Walden

Laguna Walden is located approximately 40 miles southeast of Carlsbad, New Mexico and approximately 2 miles south of N.M. Highway 128. It is located just east of the Great Salt Lake that supports a salt mining operation. It is unknown if this playa is influenced by the potash waste disposal through the chain of lakes beginning with Laguna Uno and it may have received some historic oil brine disposal.

Lane Salt Lake

Lane Salt Lake is located approximately 5 miles northwest of Tatum, New Mexico. This playa has received no potash waste disposal, but has received oil brine disposal. The Burro Pipeline Outfall is a subsite within Lane Salt Lake at the outfall of a oil brine disposal pipeline.

Middle Lake and Middle Lake Salt Playa

Middle Lake and the associated salt playa are located approximately 4 miles northwest of Tatum, New Mexico. These playas have received no potash waste disposal, but have received oil brine disposal.

William's Sink

William's Sink is located about 30 miles east of Carlsbad, New Mexico and approximately 3 miles north of U.S. Highway 162. This playa has received historic potash waste disposal from a processing plant located to the east. Disposal to this playa ceased in 1985.

Sample Collection Methods

Samples of birds, sediment, water, invertebrates and salt encrustations from dead birds were collected in 1992. Collection of birds and other species proved difficult, therefore the sample sizes were limited to a few birds and small invertebrate samples from only some of the playas. Although an attempt was made to sample all available organisms at several of the playas, birds and/or potential avian food items were frequently absent during the sampling visits. Samples were collected and preserved as follows:

Water

Samples for polycyclic aromatic hydrocarbon analysis collected directly into 500-mL glass containers, and stored and transported on ice.

Sediment

Sediment was collected as composite samples (individual samples collected from various points around the playa) of the top 2 inches of sediment sampled with a dredge or stainless steel spoon, placed in a 500-mL glass jar, and transported on ice.

Invertebrates

Invertebrate samples were collected using fine mesh seines and dip nets or were picked up by hand from the shoreline where possible and placed in chemically clean glass jars and stored on ice in the field until frozen.

Bird Tissue

Birds were collected by shotgun (steel shot) and livers were removed with stainless steel instruments, placed in chemically clean jars and stored on ice in the field until frozen.

Results

Contaminants in Water

Polycyclic Aromatic Hydrocarbons

Total PAHs were elevated above PAH criteria for human health protection proposed by Eisler (1987) at 7 of the 8 sites sampled. Total PAHs were calculated as a total of flouranthene, benzo[a]pyrene, benzo[g,h,i]perylene, benzo[b]flouranthene, benzo[k]flouranthene, and indeno[1,2,3-cd]pyrene. No water samples were collected from Middle Lake. Lane Salt Lake was the only sampling site where total PAH levels did not exceed the 0.0135 - 0.2 ppm proposed criteria. Table 2 presents the results of PAH analyses for 8 playas.

Table 2. Total PAH (ppm) in water samples from 8 playa sampling sites

Sample Location	Total PAH ppm
Laguna Gatuna	44
Laguna Quatro	14575
Laguna Tres	10266
Laguna Uno	13817
Laguna Waldon	152
Lane Salt Lake	1
Lane Salt Lake (Burro pipeline)	344
Middle Lake Salt Playa	0
Williams Sink	67

Contaminants in Sediment


BTEX

BTEX analyses were conducted on sediment samples from 7 of the 10 playa sample sites. Sediment from Laguna Quattro contained the highest levels of BTEX, and a relatively high spike in toluene levels was observed in the sample from William's Sink. Overall, levels observed in the Playa Lakes were comparable to those seen in samples collected from the San Juan River area in Northwestern New Mexico where oil and gas production is prevalent.

Table 3. Analytical Results for BTEX in 7 Sediment Samples (ug/g wet weight)

Sample Location	Benzene	Ethyl Benzene	Toluene	m & p - Xylene	o - Xylene
Laguna Gatuna	0.46	0.75	1.14	1.08	0.69
Williams Sink	0.26	0.18	1.96	0.22	0.17
Lane Salt Lake	0.128	0.281	0.027	0.205	0.136
Laguna Quatro	0.65	2.25	3.29	4.64	2.94
Laguna Tres	0.20	0.74	0.33	1.13	1.05
Laguna Uno	0.09	0.09	0.11	0.31	0.11
Laguna Waldon	0.13	0.12	0.22	0.47	0.17

Results (ug/g) BTEX Analyses (7 of 9 playa lakes)



Total Organic Carbon


Percent total organic carbon (TOC) was measured in sediment samples from 8 of the 10 playa sample sites. The highest level (3.42% TOC) was reported at Lane Salt Lake at the Burro Pipeline outfall.

Overall, levels observed in the Playa Lakes were comparable to those seen in samples collected from the San Juan River area in Northwestern New Mexico where oil and gas production is prevalent.

Table 4. Percent total organic carbon in samples from 8 playa lakes

LOCATION	TOC
Laguna Gatuna	0.1%
Lane Salt Lake (Burro pipeline outfall)	3.4%
Laguna Quatro	0.2%
Lane Salt Lake	0.1%
Laguna Tres	0.5%
Laguna Uno	0.1%
Laguna Waldon	0.8%
Middle Lake Salt Playa	0.5%

**Total Organic Carbon in Sediment
(8 playa lakes)**



Polycyclic Aromatic Hydrocarbons


PAHs were analyzed in sediment samples from 9 of 10 playas where samples were collected. The highest level (15216.21 $\mu\text{g/g}$ wwt) of total PAHs (included all PAH analytes, see Table 5) was observed at Lane Salt Lake at the Burro Pipeline outfall. Overall, levels observed in the Playa Lakes were comparable to

those seen in samples collected from the San Juan River area in Northwestern New Mexico where oil and gas production is prevalent.

Table 5. Total PAHs in sediment samples from 9 playa lakes

Sample Location	Total PAH
Laguna Gatuna	209
Lane Salt Lake	77
Lane Salt Lake (Burro pipeline outfall)	15216
Middle Lake Salt Playa	79
Laguna Uno	34
Laguna Tres	5713
Laguna Quatro	819
Williams Sink	41
Laguna Waldon	5

Total PAH (ug/g wwt) in Sediment from 9 Playas



Trace Elements

Trace element analyses for sediment samples were only conducted on one sample from Laguna Quattro. Levels of aluminum (Al), barium (Ba), cadmium (Cd), mercury (Hg), magnesium (Mg), selenium (Se), and vanadium (V) exceeded avian dietary threshold criteria for adverse effects. Waterfowl normally ingest some soil and sediment while feeding. Significant ingestion of contaminated sediment and other food items can result in bioaccumulation of contaminants within an organism. Adverse effects can range from death to more chronic impacts such as reproductive impairment.

Aluminum, barium and magnesium are normally elevated in New Mexico soils, but the levels observed at Laguna Quattro were still above geometric means of 3736 mg aluminum/kg dry weight and 123 mg barium/kg dry weight found in other studies (O'Brien 1990). Because aluminum, magnesium and barium are common constituents of New Mexico soils, the elevated concentrations were most likely a consequence of local geologic sources. Cadmium, selenium, mercury, and vanadium, however, could originate from natural and/or anthropogenic sources. Anthropogenic sources may include storm runoff, roads and other non-point sources or point source discharges from mine wastes and associated runoff. Further sampling is required to assess natural background concentrations and potential anthropogenic sources of these elements near the playas.

Table 6. Criteria used to evaluate the potential hazard of trace elements in sediments and potential food items to birds and other organisms, expressed as mg/kg wet or dry weight (ww or dw; NA = not available).

Analyte	DTC (ww) ^a	Ontario MME Lowest Observable Effect Level (dw) ^b	Other Sediment Quality Criteria (dw) ^c	Rio Grande Geometric Mean- Background (dw) ^d
Silver	NA	NA	0.5	NA
Aluminum	200.0	NA	NA	3735.0
Arsenic	30.0	6.0	---	1.90
Barium	20.0	NA	20.0	123.01
Beryllium	NA	NA	NA	0.30
Boron	30.0	NA	NA	2.63
Cadmium	0.1	0.6	---	0.24

Analyte	DTC (ww) ^a	Ontario MME Lowest Observable Effect Level (dw) ^b	Other Sediment Quality Criteria (dw) ^c	Rio Grande Geometric Mean- Background (dw) ^d
Chromium	5.1	26.0	---	4.83
Cobalt	NA	NA	50.0	NA
Copper	300.0	16.0	---	5.53
Iron	NA	21,200 (2.0%)	---	6164.9
Mercury	0.1	0.2	---	0.00
Lead	50.0	31.0	---	6.74
Magnesium	3,000.0	NA	NA	2168.0
Manganese	2,000.0	460.0	---	840.8
Molybdenum	100.0	NA	4.0	1.78
Nickel	100.0	16.0	---	4.69
Selenium	0.8	NA	5.0	0.30
Strontium	3,000.0	NA	NA	72.67
Antimony	NA	NA	NA	NA
Vanadium	10.0	NA	NA	NA
Zinc	44.5	120.0	---	24.46

a Dietary Threshold Concentration; derived from various sources (NRC 1980, Eisler (USFWS Contaminant Report Series) 1985-94, Lemly and Smith 1987, and International Joint Commission 1993).

b Ontario Ministry of the Environment and Energy, Ontario, Sediment Quality Guidelines (1993).

c Derived from various sources cited in MacDonald (1994).

d From USFWS, Contaminants Investigation of Bitter Lake NWR, Roswell, NM (O'Brien 1990).

Cadmium and mercury (Hg) MDLs are too high to assess if these metals exceed avian DTCs. Most other sediment metals were not elevated above typical sediment background concentrations and should pose little risk to waterfowl and aquatic organisms. Of those metals that were above typical background concentrations, most are not directly hazardous to waterfowl. Overall, birds may be at risk from some sediment metals (Table 6) if: (1) sediment metals are bioaccumulating in food items or bird tissues, (2) a significant percentage of the bird's diet includes ingested sediment (ingested while feeding on benthic invertebrates and plants).

Contaminants in Aquatic Biota

Contaminant residues in aquatic biota and vegetation are an indicator of the bioaccumulation potential for birds eating fish, invertebrates, and vegetation. Little data is available to interpret the toxic potential of contaminant residues in plants, invertebrates and amphibians. Lethal or sublethal effects of elevated trace element concentrations in aquatic biota may translate to an indirect risk to waterfowl by way of reduction in the available food base. Adverse effects on the food base available to the birds must, therefore, be assessed by comparison to water and sediment quality criteria, or, by actual assessment of the biota present in each playa.

Trace Elements

Trace element analyses were conducted for samples of invertebrates collected from 3 playas, Laguna Walden, Middle Lake, and William's Sink and for samples of bird liver from 4 playas, Laguna Quattro, Laguna Walden, Lane Salt Lake (at Burro Pipeline outfall), and Middle Lake. One bird kidney sample was collected from Laguna Quattro and was analyzed for trace elements. In invertebrates, concentrations of aluminum, magnesium, and selenium exceeded avian dietary threshold concentrations at Middle Lake and William's Sink. Concentrations of these three elements in invertebrates from Laguna Walden were all below DTCs.

Conclusions and Recommendations

The presence of large bodies of water in the normally arid environment of Southeastern New Mexico have been shown to pose a significant hazard to migratory bird species. Many natural playa lake depressions that are normally dry during a significant portion of the spring and fall migratory seasons are kept constantly wet from brine waste discharges from mining and drilling operations. Many man made playa

lakes also exist in the area, some of which are relatively deep and contain a significant amount of water year round.

Both current and historic discharges into these playas of salt water waste, often containing a slurry of many other minerals, have created anthropogenic, attractive hazards for waterfowl and other migratory wildlife. In addition to the proven lethal doses of salt water available to these species, other hazards may exist from other contaminants such as selenium and arsenic. These contaminants are made available to species in natural and man-made water impoundments either directly as waste water from mining and drilling operations or through overland runoff and soil leaching. Forensic necropsy reports for migratory bird carcasses found in or around some of the playas have all indicated salt toxicosis as the cause of death.

The problem of the salt playas in this State and elsewhere has been hotly debated for years between State and Federal government agencies and various industry representatives. In the New Mexico playa lakes region, the U.S. Bureau of Land Management (BLM), the New Mexico Environment Department (NMED) and the New Mexico Oil Conservation Division (NMOCD) share governmental oversight and regulatory responsibility for both potash mining and oil and gas drilling. The BLM is responsible for the land and mineral extraction lease agreements and mining plans under which mining and drilling companies establish the methods they will practice. These agreements allow the unmitigated release of millions of gallons of waste water that is not regulated by the National Pollutant Discharge and Elimination System (NPDES) administered by the U.S. Environmental Protection Agency.

The Southeast New Mexico Playa Lakes Coordinating Committee was formed from members of State and Federal government and representatives from both the oil and gas and potash industries. Several plans coordinated by this group to keep birds off the largest problem playa (Laguna Toston), failed to significantly reduce or eliminate migratory bird deaths. No mitigative measures or modifications to current discharge practices have yet been proposed by the Committee. However, some significant

measures are required to eliminate the hazard to migratory wildlife and possibly mitigate for the historic losses to both migratory birds, who would otherwise not use this area, and to the habitat for other native species that has been destroyed by the playas.

Currently, negotiations are being conducted between the U.S. Fish and Wildlife Service, the BLM, State Government and various industry representatives to establish a viable solution. Both mitigative and practice modification techniques have been discussed. It is recommended that this report along with the June 1997 report from USGS be used in these continuing discussions.

Literature Cited

- Dein, Joshua F., Lauri A. Baetan, Carol U. Meteyer, Melody K. Moore, Michael D. Samuel, Paul D. Miller, Christopher Murphy, Steven Sissler, Clinton W. Jeske, Joseph R. Jehl, Jr., J.S. Yaeger, B. Bauer and Shiela A. Mahoney. 1997. Investigation into Avian Mortality in the Playa Lakes Region of Southeastern New Mexico: Final Report - June 1997. U.S. Geological Survey Biological Resources Division, National Wildlife Health Center, National Wetlands Research Center.
- Eisler, R., 1994, A review of arsenic hazards to plants and animals with emphasis on fishery and wildlife resources, in Nriagu, J.O, (ed.), Arsenic in the Environment, Part II: Human Health and Ecosystem Effects, p. 185-259.
- Eisler, R., 1993. Zinc Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review. Biological Report 10, Patuxent Wildlife Research Center U.S. Fish and Wildlife Service, Laurel, Maryland. 106 pp.
- Eisler, R. 1990. Boron hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish Wildl. Serv. Biol. Rep 85(1.20). 32 pp.
- Eisler, R. 1988. "Arsenic Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review." Prepared by U.S. Fish and Wildlife Service, Patuxent Wildlife Research Center, Laurel, MD. Sponsored by U.S. Department of the Interior. Biological Report 85(1.12), Washington, D.C., 80pp.
- Eisler, R., 1988b. Lead Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review. Biological Report 85(1.14), Patuxent Wildlife Research Center U.S. Fish and Wildlife Service, Laurel, Maryland. 134 pp.
- Eisler, R. 1987. Polycyclic Aromatic Hydrocarbon Hazards to Fish, Wildlife and Invertebrates: a Synoptic Review. U.S. Fish and Wildl. Serv. Biol. Rep., 85(1.11)
- Eisler, R. 1987. Mercury hazards to fish and wildlife and invertebrates: A synoptic review. U.S. Fish and Wildlife Service Biological Report 85(1.10), Washington, D.C., 90pp.
- Eisler, R., 1986. Chromium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review. Biological Report 85(1.6 , Patuxent Wildlife Research Center Laurel, U.S. Fish and Wildlife Service, Laurel, Maryland). 60 pp.
- Eisler, R., 1985. Cadmium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review. Biological Report 85(1.2), Patuxent Wildlife Research Center Laurel, U.S. Fish and Wildlife Service, Maryland. 46 pp.

- Eisler, R. 1985. Selenium Hazards to Fish, Wildlife, and Invertebrates: a Synoptic Review. U.S. Fish Wildl. Serv. Biol. Rep. 85(1.5). Contaminant Hazard Reviews Report No.5. Patuxent Wildlife Research Center, U.S. Fish and Wildlife Service, Laurel, Maryland 72.
- Lemly, A.D., and G.J. Smith. 1987. Aquatic Cycling of Selenium: Implications for Fish and Wildlife. Fish and Wildlife Leaflet No. 12. U.S. Fish and Wildlife Service, Washington, DC, 10pp.
- International Joint Commission. 1993. A strategy for virtual elimination of persistent toxic substances: Volume 1. Report of the virtual elimination task force. Windsor, Ontario. 72pp.
- MacDonald, D. D. 1994. Approach to the Assessment of Sediment Quality in Florida Coastal Waters, Florida Department of Environmental Protection, Tallahassee, Florida.
- National Research Council, Subcommittee on Mineral Toxicity in Animals, 1980, Mineral Tolerance of Domestic Animals: Washington, D.C., National Academy Press, 577 p.
- O'Brien, T.F. 1990. Organochlorine and heavy metal contaminant investigation for Bitter Lake National Wildlife Refuge, Roswell, New Mexico, 1986. U.S. Fish and Wildlife Service, New Mexico Ecological Services Field Office Contaminant Report, April 1990.
- Ontario MME (Ontario Ministry of the Environment and Energy). 1993. Development of the Ontario Provincial sediment quality guidelines for arsenic, cadmium, chromium, copper, iron, lead, manganese, mercury, nickel, and zinc. August 1993. 24 pp.