

**R
E
C
O
V
E
R
Y

S
A
N

J
U
A
N

R
I
V
E
R

B
A
S
I
N

I
M
P
L
E
M
E
N
T
A
T
I
O
N

P
R
O
G
R
A
M**

FISCAL YEAR 2009 ANNUAL BUDGET AND WORK PLAN



Approved: August 14, 2008

San Juan River Recovery Implementation Program FY2009 Funding

09 SOW	Title	Agency	Estimated Funding		
			Base	Capital	Other
Element 1 - Development, Integration, and Evaluation of Information for Recovery					
	Integrate biological/habitat data	RFP	\$100,000		
	Program Workshops	BOR/FWS	\$30,000		
6	Peer Review	BOR/FWS	\$40,000		
	Subtotal		\$170,000	\$0	\$0
Element 2 - Management and Augmentation of Populations and Protection of Genetic Integrity					
8	Stocking of Fingerling Colorado Pikeminnow	FWS, ABQ	\$19,344		
9	Colorado Pikeminnow Fingerling Production	FWS, DNFHTC	\$103,008		
10	Rearing Razorback Suckers Dexter	FWS, DNFHTC	\$73,853		
11	Razorback Sucker Production Uvalde	FWS, UNFH	\$127,272		
12	RBS Augmentation/NAPI Pond Management	FWS/NN	\$102,689		
	Subtotal		\$426,166	\$0	\$0
Element 3 - Protection, Management, and Augmentation of Habitat					
13	Maintenance and Operation of Model	BR, Durango	\$143,910		
14	Improve Stream Gaging and Flow Measurements	USGS	\$6,700		
	PNM O&M	PNM	\$25,000		
15	Operation of PNM Fish Passage Structure	FWS/NN	\$79,598		
16	Feasibility/Pre-planning for Improving Flow Effectiveness using Mechanical Manipulation	FWS/TNC	\$20,000		
	Capital Projects Management	BR		\$68,300	
	Capital Hogback Canal	BR		\$400,000	
	Subtotal		\$275,208	\$468,300	\$0
Element 4 - Interactions between Native and Non-Native Fish Species					
17	Nonnative Species Control - Upper San Juan River	FWS, ABQ	\$322,567		
18	Nonnative Species Control - Lower San Juan River	UDWR	\$181,044		
	Subtotal		\$503,611	\$0	\$0
Element 5 - Monitoring of Fish and Habitat					
19	Sub-Adult/Adult Large-Bodied Fish Comm. Monitoring	FWS, GJ	\$122,034		
20	YOY/Small-Bodied Fish Monitoring	NMDGF	\$89,479		\$40,000
21	RBS/CPM Larval Surveys (Combined SOW)	NMDGF	\$224,561		
22	Specimen Curation/Identification	UNM	\$30,117		
23	Fish-Habitat Relationship Surveys (2nd year of pilot project)	RFP	\$150,000		
24	Temperature Monitoring	RFP	\$12,000		
25	Long Term Channel Monitoring, Habitat Mapping	RFP	\$5,000		
26	River Videography		\$16,000		
27	PIT Tags	BR	\$45,000		
	Subtotal		\$694,191	\$0	\$40,000

Element 6 - Information and Education				
	Education and Outreach	FWS, ABQ	\$15,000	
	Subtotal		\$15,000	\$0
				\$0
Element 7 - Program Coordination				
28	Program Management FWS	FWS, ABQ	\$215,924	\$165,000
29	Base Fund Management BR	BR, SLC	\$137,135	
	Subtotal		\$353,059	\$0
				\$165,000
	SJRRIP Total		\$2,437,235	\$468,300
				\$205,000
	Actual Annual 2009 Base Funds		\$2,444,000	
	Estimated available 2009 funds to proposed expenditures		\$6,765	
	Projects Not Funded			
	Program Database		\$10,000	
	LT Geomorph Surveys		\$40,000	

2009 WORK PLAN PROJECTS**DELIVERABLES****DATE DUE****ELEMENT 1 - Development, Integration, and Evaluation of Information for Recovery****09-6****Peer Review**

Mark McKinstry
 U.S. Bureau of Reclamation
 125 S. State St.
 Salt Lake City, UT 84138-1147
 (801) 524-3835
mmckinstry@uc.usbr.gov

- Letter or verbal reports from each reviewer.

Upon request

ELEMENT 2 - Management and Augmentation of Populations and Protection of Genetic Integrity**09-8****Stocking of Fingerling Colorado Pikeminnow**

Weston Furr, Ernest Teller, and Jason Davis
 U.S. Fish and Wildlife Service
 New Mexico Fish and Wildlife Conservation Office
 3800 Commons N.E.
 Albuquerque, NM 87109
 (505) 342-9900
Jason_E_Davis@fws.gov
Weston_Furr@fws.gov
eteller@wildblue.net

- Database Files
- Draft Report
- Final Report

03-31-10

03-31-10

06-01-10

09-9**Colorado Pikeminnow Fingerling Production**

William Knight and Manuel E. Ulibarri
 Dexter National Fish Hatchery and Technology Center
 U.S. Fish and Wildlife Service
 PO Box 219, 7116 Hatchery Road
 Dexter, NM 88230-0219
 (505) 734-5910 Work
 (505) 734-6130 FAX
William_knight@fws.gov
Manuel_ulibarri@fws.gov

- Transport and distribute 300,000 fish age-0 fingerlings and 3,000 age-1 fish from Dexter to the San Juan River
- Draft Report
- Final Report

Annually

03-31-10

06-01-10

09-10**Razorback Sucker Production - Dexter**

William Knight and Manuel E. Ulibarri
 Dexter National Fish Hatchery and Technology Center
 U.S. Fish and Wildlife Service
 PO Box 219, 7116 Hatchery Road
 Dexter, NM 88230-0219
 (505) 734-5910 Work
 (505) 734-6130 FAX
William_knight@fws.gov
Manuel_ulibarri@fws.gov

- Delivery of 20,000, 200mm fish to Uvalde NWR and 10,000 200mm fish to NAPI ponds
- Draft Report
- Final Report

Annually

03-31-10

06-01-10

09-11**Razorback Sucker Production – Uvalde**

Grant L. Webber

Uvalde National Fish Hatchery

754 County Road 203

Uvalde, Texas 78801

830-278-2419

830-278-6042 Fax

Grant_Webber@fws.gov

- Delivery of 12,000, 300mm fish to the San Juan River Annually
- Draft Report 03-31-10
- Final Report 06-01-10

09-12**Razorback Sucker Augmentation at NAPI Ponds**

Jeffrey Cole, Albert Lapahie, and Viola Willetto

Navajo Nation Department of Fish and Wildlife

Box 1480

Window Rock, AZ 86515

(928) 871-7068

jcole@navajofishandwildlife.org,alapahie@navajofishandwildlife.orgvwilleto@navajofishandwildlife.org

- Harvest \geq 300 mm fish from 3 NAPI grow-out ponds and transport to San Juan River Annually
- Draft Report 03-31-10
- Final Report 06-01-10

Weston Furr, Ernest Teller, and Jason Davis

U.S. Fish and Wildlife Service

New Mexico Fish and Wildlife Conservation Office

3800 Commons N.E.

Albuquerque, NM 87109

(505) 342-9900

Jason_E_Davis@fws.govWeston_Furr@fws.goveteller@wildblue.net**ELEMENT 3 - Protection, Management, and Augmentation of Habitat****09-13****Maintenance & Operation of the San Juan River Basin Hydrology Model**

Katrina Grantz and Ryan Christianson

U.S. Bureau of Reclamation

835 E. 2nd Avenue, Suite #300

Durango, CO 81301

(970) 385-6590

kgrantz@uc.usbr.govrchristianson@uc.usbr.gov

- Hydrology analysis of water development scenarios or other scenarios in accordance with the HC's document entitled, "SJRIIP Recommendations for Conducting and/or Requesting Model Runs" Upon request

09-14**Improve Stream Gaging and Flow Measurements**

Pat Page

U.S. Bureau of Reclamation

835 E. 2nd Avenue, Suite #300

Durango, CO 81301

(970) 385-6560

ppage@uc.usbr.gov

- Technical presentation End of year

09-15**Operation PNM Fish Passage Structure**

Jeffrey Cole, Albert Lapahie, and Viola Willeto
Navajo Nation Department of Fish and Wildlife
Box 1480
Window Rock, AZ 86515
(928) 871-7068
jcole@navajofishandwildlife.org,
alapahie@navajofishandwildlife.org
vwilleto@navajofishandwildlife.org

- Monthly Fish Passage Reports
- Draft Annual Report 03-31-10
- Final Annual Report 06-30-10

Capital Improvement Program

Brent Uilenberg
U.S. Bureau of Reclamation
2764 Compass Dr., Suite 106
Grand Junction, CO 81506
(970) 248-0641
builenberg@uc.usbr.gov

- Financial reports presented to Coordination Committee giving status of Federal appropriations and non-Federal cost sharing contributions. Every Meeting

ELEMENT 4 - Interactions between Native and Non-Native Fish Species

09-17**Nonnative Species Monitoring and Control in the Upper San Juan River**

Jason E. Davis and D. Weston Furr
U.S. Fish and Wildlife Service
New Mexico Fish and Wildlife Conservation Office
3800 Commons N.E.
Albuquerque, NM 87109
(505) 342-9900
Jason_E_Davis@fws.gov
Weston_Furr@fws.gov

- PIT Tag Information 12-31-09
- All Data Files 03-31-10
- Interim Progress Report 03-31-10
- Draft Final Report 06-01-10

09-18**Nonnative Species Control in the Lower San Juan River**

Darek S. Elverud
Utah Division of Wildlife Resources
Moab Field Station
1165 S. Hwy 191, Suite 4
Moab, UT 84532
(435) 259-3782
darekelverud@utah.gov

- PIT Tag Information 12-31-09
- All Data Files 03-31-10
- Interim Progress Report 03-31-10
- Draft Final Report 06-01-10

ELEMENT 5 - Monitoring of Fish and Habitat

09-19**Sub-Adult & Adult Large-Bodied Fish Monitoring**

Dale Ryden and Chuck McAda
U.S. Fish and Wildlife Service
Colorado River Fishery Project
764 Horizon Drive, Building B
Grand Junction, CO 81506-3946
(970) 245-9319
cmcada@fws.gov
dryden@fws.gov

- PIT Tag Information 12-31-09
 - All Data Files 03-31-10
 - Interim Progress Report 03-31-10
 - Draft Final Report 06-01-10
-

09-20**YOY/Small Bodied Fish Monitoring**

David L. Propst and Yvette Paroz
 Conservation Services Division
 New Mexico Department of Game and Fish
 One Wildlife Way, P.O. Box 25112
 Santa Fe, NM 87504
 (505) 476-8103
dpropst@state.nm.us
yparoz@state.nm.us

- PIT Tag Information 12-31-09
- Draft Report including electronic database 03-31-10

09-21**Larval Colorado Pikeminnow and Larval Razorback Sucker Survey**

W. Howard Brandenburg and Michael A. Farrington
 American Southwest Ichthyological Researchers, LLC
 800 Enchino Place NE
 Albuquerque, NM 87102-2606
 (505) 247-9337
 (505) 247-2522 FAX
howard_brandenburg@asirllc.com
Michael_farrington@asirllc.com

- Draft Report 03-31-10
- Final Report 06-01-10

David L. Propst
 New Mexico Department of Game and Fish
 Conservation Services Division
 One Wildlife Way,
 Santa Fe, NM 87505
 (505) 476-8103
dpropst@state.nm.us

09-22**Specimen Curation/Identification**

Alexandra M. Snyder and Thomas F. Turner
 Division of Fishes - Museum of Southwestern Biology
 University of New Mexico
 Albuquerque, NM 87131
 (505) 277-6005
amsnyder@unm.edu
turnert@unm.edu

- Draft Report 03-31-10
- Final Report 06-01-10

09-23,24,25,26**Geomorphology and Habitat**

Ron Bliesner
 Keller-Bliesner Engineering
 78 East Center, Logan, UT 84321
 (435) 753-5651
bliesner@kelbli.com

- Element 1 – Temperature Monitoring**
- Draft Report 03-31-10
- Final Report and updated database 06-30-10
- Element 2 – Habitat Mapping**
- Base photography acquired 07-09
- Base maps prepared 08-09
- Archived digital image files 09-09
- Element 3 – Geomorphology**
- Draft Report 03-31-10
- Final Report 06-30-10
- Element 4 – Habitat Use of YOY Fish**
- Draft Report 03-31-10
- Final Report 06-30-10

09-27**Pit Tag and Readers**

Mark McKinstry

U.S. Bureau of Reclamation

125 S. State St.

Salt Lake City, UT 84138-1147

(801) 524-3835

mmckinstry@uc.usbr.gov

- Purchase as necessary

Annually

ELEMENT 7 - Program Coordination

09-28**Program Management – USFWS**

Program Coordinator and Program Support Assistant

U.S. Fish and Wildlife Service

2105 Osuna Road NE

Albuquerque, NM 87113

(505) 761-4745 david_campbell@fws.gov(505) 761-4753 sharon_whitmore@fws.gov(505) 761-4739 scott_durst@fws.gov

- Misc. tasks completed as per SOW and Program document

Annually

09-29**Base Funds Management - Reclamation**

Mark McKinstry

U.S. Bureau of Reclamation

125 S. State St.

Salt Lake City, UT 84138-1147

(801) 524-3835 mmckinstry@uc.usbr.gov

- Funding requests processed
- Draft Annual Report
- Final Annual Report

Upon request

03-31-10

06-01-10

**Peer Review for 2009
Fiscal Year 2009 Project Proposal**

Mark McKinstry UC-735
Bureau of Reclamation
125 South State Street, Room 6107
Salt Lake City, UT 84138-1147
Phone 801-524-3835
FAX 801-524-5499
mmckinstry@uc.usbr.gov

Background:

A Peer Review Panel was established in 1997 to assist the SJRIP with reports and plans for future studies. The four members of the panel participate in meetings and comment on pre-draft, draft, and final Scopes of Work, Workplans, reports, Integration analyses and reports, and other Program Documents.

This proposal provides for funding for the Peer Review Panel activities during 2009. It is anticipated that the Panel will meet with the Biology Committee at three meetings during the year; the December 2008 Planning meeting, the February/March, 2009 Researcher's meeting (combined with the Coordination Committee), and a May, 2009 meeting to finalize 2010 SOWs. Additionally, the Peer Reviewers will likely be asked to attend an additional meeting in conjunction with the Annual Researcher's meeting whereby they are asked to comment as a group on all aspects of the Program and Workplan. At this meeting the Peer Reviewer's will be tasked to review all activities of the SJRIP and possibly assist in the development of a Sufficient Progress Report. In 2009 the SJRIP will begin another Data Integration Review and the Program expects that the Peer Reviewers may be asked to participate in that activity.

Goal:

The goal of peer review is to provide additional scientific oversight over San Juan River Recovery Implementation Program technical studies and reporting. The Peer Review Panel will work with the Biology Committee and Program Staff to produce scientific credible documents and will assist the Biology Committee in maintaining a scientific basis for the Program.

Methods:

The Peer Review Panel will meet with the Biology Committee in FY2009 three times to review monitoring and research progress and to discuss scopes of work for 2010. They will provide verbal input during the meetings and provide written reviews of the progress of the Program. Their reviews will be provided to the Biology Committee through Mark McKinstry and David Campbell in letter form, or on the Biology Committee list server, and through discussions at the Biology Committee meetings. Biology Committee researchers may call Peer Review Panel members to ask for advice, and Peer Review Panel members may call Biology Committee researchers if they have questions concerning Program activities. All correspondence between the Biology Committee and the Peer Review Panel will be coordinated through either Mark McKinstry or David Campbell.

Products:

Peer review participation at 3 Biology Committee meetings, letter or verbal reports from each peer reviewer on an as-requested basis.

Primary Contact: Dr. Mark McKinstry
 Bureau of Reclamation
 125 South State Street, UC-735
 Salt Lake City, UT 84106
 Phone: 801/524-3835 FAX: 801-524-5499
 Email: mmckinstry@uc.usbr.gov

Personnel:

Dr. John Pitlick
 Department of Geology
 University of Colorado
 Boulder, CO 80309-0260
 Phone: 303-492-5906
 Email: pitlick@spot.colorado.edu

Dr. Stephen Ross
 Curator Emeritus of Fishes, Department of
 Biology and Museum of Southwestern
 Biology MSC 03-2020 University of New Mexico
 Albuquerque, NM 87131-0001
 Phone: 505-277-3893 Hm: 970-264-0158
 Email: stross1@unm.edu

Dr. Mel Warren Jr.
 Team Leader and Research Biologist
 Center for Bottomland Hardwoods Research
 Southern Research Station, USDA Forest Service
 1000 Front Street
 Oxford, MS 38655
 Phone: 662-234-2744, ext. 246
 Fax: 662-234-8318
 Email: mwarren01@fs.fed.us

Dr. Ron Ryel
 Department of Forest, Range, and Wildlife
 Utah State University
 5230 Old Main Hill
 Logan, UT 84322-5230
 Phone: 435-797-8119
 FAX: 435-797-3796
 Email: ron.ryel@usu.edu

Budget FY-09:

Payment for serving on the Peer Review Panel includes expenses for travel to and from meetings, and an hourly rate for services. It is anticipated that Panel Members will spend approximately 25-40 days each in 2009.

Due to the unknown commitment required for Data Integration, scheduled to begin in 2009, the budget for Peer Reviewers is estimated. The total budget is distributed among the four peer reviewers through individual Services Contracts with Reclamation.

Salaries:	\$30,000
Travel:	<u>\$10,000</u>
Total	\$40,000

Future use of the Peer Review Panel is not known but they likely will be used each year to provide guidance to the Biology Committee.

Estimated Outyear Funding:

2010	\$40,000
2011	\$40,000

Augmented Stockings of Age 0 and Age 1+ Colorado pikeminnow into the San Juan River Fiscal Year 2009 Project Proposal

Principal Investigators: D. Weston Furr, Ernest Teller, Sr. and Jason E. Davis
United States Fish and Wildlife Service
New Mexico Fish and Wildlife Conservation Office
3800 Commons N.E.
Albuquerque, N.M. 87109
(505) 342-9900

Weston_Furr@fws.gov eteller@wildblue.net

Jason_E_Davis@fws.gov

Background

Colorado pikeminnow (*Ptychocheilus lucius*) is a federally-listed endangered fish found in the San Juan River. The San Juan River Recovery Implementation Program (SJRIP) was initiated in 1992 to protect and recover populations of both Colorado pikeminnow and razorback sucker (*Xyrauchen texanus*) in the San Juan River Basin while water development proceeds in compliance with all applicable federal, state, and tribal laws (USFWS 2008). Recovery of Colorado pikeminnow, as listed in the recovery goals, is dependent on the maintenance of a wild population of at least 2,600 adults in the Green River subbasin and at least 700 adults in the upper Colorado River subbasin, as well as a target of 1,000 subadults in the San Juan River subbasin (USFWS 2002). Delisting criteria include a self sustaining population that exceeds 800 adults maintained in the San Juan River subbasin.

Fish community monitoring during the SJRIP seven year research period, 1991-1997, identified few wild Colorado pikeminnow inhabiting the San Juan River and prompted investigation into the feasibility and implementation of augmenting the population with hatchery reared fish. As a result of these findings, an experimental stocking of Colorado pikeminnow was conducted by Utah Department of Wildlife Resources in 1996 with the purposes of evaluating dispersal and retention of stocked Colorado pikeminnow and determining the availability, use, and selection of habitats by early life stages of Colorado pikeminnow (Ryden 2008). Stockings of larval, sub-adult and adult fish after this initial stocking resulted in the subsequent recapture of stocked fish suggesting that Colorado pikeminnow could survive in the San Juan River. In 2003, *An Augmentation Plan for Colorado Pikeminnow In The San Juan River* was finalized (Ryden 2003). This plan and later amendments calls for the annual stocking of $\geq 300,000$ age-0 and 3,000 age 1+ fish in the San Juan River until 2009. It is anticipated that this augmentation period will be extended to include continued augmentation past 2009.

Augmentation of the Colorado pikeminnow population in the San Juan River is related to the 2008 SJRIP Draft Long Range Plan (LRP). These activities are specifically addressed in the following Elements, Actions and Tasks:

Element 2. Management and Augmentation of Populations and Protection of Genetic Integrity

Action 2.1.2 Produce, rear and stock sufficient numbers of Colorado pikeminnow to meet stocking goals of augmentation plan.

Task 2.1.2.2 Annually stock $\geq 300,000$ age-0 Colorado pikeminnow.

Task 2.1.2.3 Annually stock 3,000 age-1+ Colorado pikeminnow.

Task 2.1.2.4 Opportunistically stock available Colorado pikeminnow in excess to those described above.

Action 2.1.4 Evaluate factors limiting Colorado pikeminnow population recovery

- Task 2.1.4.1 Identify strategies for improving survival and retention of stocked Colorado pikeminnow, including but not limited to acclimation, size of fish stocked, time of stocking, location of stocking, conditioning, and predator avoidance.

In addition to SJRIP Program priorities, the stocking of fish reared at U.S. Fish and Wildlife Service (Service) hatcheries in the Southwest Region (Region 2; New Mexico, Arizona, Texas and Oklahoma) are subject to Regional Policy No. 03-06, "Stocking of fish and other aquatic species". This policy applies to production, transport, and stocking for Service hatchery production and incorporates guidance and requirements from FWS Fish Health Policy (713 FWM 1-5), Policy for Controlled Propagation of Species Listed under the Endangered Species Act (Federal Register 65:183), and goals and objectives of the FWS Strategic Plan for the Fisheries Program. The FWS Fish and Wildlife Conservation Offices are the primary conduit for satisfaction of Policy requirements and ensures compliance with needs relative to fish health, stocking requests and priorities, deviation from approved stocking requests, pre-stocking treatments (e.g. nonnative fish removal from stocking sites), and applicable environmental compliance. The New Mexico Fish and Wildlife Conservation Office is the pertinent field office for the processing of SJRIP stocking requests under this policy directing the change in lead coordination and stocking responsibilities from FWS Region 6 to Region 2.

Objectives

1. Coordinate with Dexter National Fish Hatchery and Technology Center (NFH&TC) to procure and stock fish according to guidelines set forth in *An Augmentation Plan for Colorado Pikeminnow* (2003) in the San Juan River.
 - a. Annually stock 300,000 age-0 Colorado pikeminnow.
 - b. Annually stock 3,000 age-1+ Colorado pikeminnow.
2. Provide summarization report on timing and location of individual stockings, numbers, and age classes while relating information to fulfillment of recommended stocking numbers as outlined in the augmentation plan.

Methods and Approach

Objective 1.a. Age-0 Colorado pikeminnow will be annually reared and harvested by Dexter NFH&TC and delivered via standard distribution unit to the San Juan River. Fish will be stocked in the fall of each year, post irrigation season, to eliminate the risk of fish entrainment in irrigation canals. In accordance to the augmentation plan, multiple stocking sites will be used ranging from Fruitland Diversion Dam downstream to Shiprock Bridge. Additional sites downstream of Shiprock Bridge will be identified, and used as needed, for suitable use as temporary holding sites for acclimation (e.g. settling of blood chemistry post-delivery, flow, etc.) purposes.

Objective 1.b. Age-1+ Colorado pikeminnow will be annually reared and harvested by Dexter NFH&TC and delivered via standard distribution unit to the San Juan River. All age 1+ fish will be implanted with a Passive Implant Transponder (PIT) tag by Dexter NFH&TC personnel prior to delivery. Depending on availability, fish will be stocked in both the spring, prior to runoff, and in the fall of each year. During each stocking a subset of fish will be acclimated for up to 24 hours. Non-acclimated fish will be released at the same location and time. Subsequent recaptures of stocked fish will provide guidance on relative success of acclimation.

Objective 2. New Mexico FWCO will collate all pertinent stocking information including but not limited to timing, location, size and numbers and enter into a standardized database that will be provided to the Program Coordinators office for deposition. These data and subsequent recapture data will be used to evaluate acclimation effectiveness.

Products/Schedule

An electronic data file will be provided for inclusion in the centralized database by 31 March 2010. A draft summary report detailing findings will be submitted to the San Juan River Implementation Program, Biology Committee, by 31 March 2010. Revisions will be completed and a final annual report will be submitted by 1 June 2010.

Literature Cited

- Ryden, D.W. 2003. An augmentation plan for Colorado pikeminnow in the San Juan River. U.S. Fish and Wildlife Service, San Juan River Recovery Implementation Program, Albuquerque, NM. 63 pp. + appendices.
- Ryden, D.W. 2008. Augmentation of Colorado pikeminnow in the San Juan River: 2007. Interim Progress Report (Final) submitted to U.S. Fish and Wildlife Service, San Juan River Recovery Implementation Program, Albuquerque, NM. 6 pp. + appendices.
- San Juan River Basin Recovery Implementation Program. 2008. Long-range plan (Draft). San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, NM.
- U.S. Fish and Wildlife Service. 2002. Colorado pikeminnow (*Ptychocheilus lucius*) Recovery Goals: amendment and supplement to the Colorado Squawfish Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, CO.

FY 2009 Proposed Budget:**Personnel/Labor Costs (Federal Salary + Benefits)**

Fish Biologist (GS-7-2) –16 days @ \$218/day	\$ 7,848.00
Age-0 stockings (Objective 1.a): (1 person x 4 days/trip x 2 trips)	
Age-1 stockings (Objective 1.b): (1 person x 4 days/trip x 2 trips)	
Reporting/Data Management (Objective 2) – 20 days	

Bio. Science Technician (GS-8) – 16 days @ \$319/day	\$ 5,104.00
Age-0 stockings (Objective 1.a): (1 person x 4 days/trip x 2 trips)	
Age-1 stockings (Objective 1.b): (1 person x 4 days/trip x 2 trips)	

Sub-total	\$ 12,952.00
-----------	--------------

Travel and Per Diem (Based on Published FY-2008 Federal Per Diem Rates)

Hotel Costs – 12 nights (12 nights @ \$70/night – single occupancy = \$1,260)	\$ 840.00
Per Diem (Hotel Rate) – 16 days @ \$39/day	\$ 624.00

Sub-total	\$ 1,464.00
-----------	-------------

Equipment

Vehicle Maintenance & Gasoline 2,400 miles @ \$0.60/mile (based on Anticipated GSA rates established on 01 September 2008 and includes costs associated with gasoline/diesel fuel vehicle maintenance)	\$ 1,440.00
--	-------------

Sub-total	\$ 1,440.00
-----------	-------------

USFWS-NMFWCO Total	\$ 15,856.00
---------------------------	---------------------

USFWS Region 2 Regional Office Administrative Overhead (22.00%)	\$ 3,488.00
--	--------------------

USFWS Region 2 Total	\$ 19,344.00
-----------------------------	---------------------

Outyear funding (based on 5% Cost of Living Increase and extension of Augmentation program beyond 2009)

FY 2009	\$ 19,344
FY 2010	\$ 20,311
FY 2011	\$ 21,327

COLORADO PIKEMINNOW FINGERLING PRODUCTION
San Juan River
FY-2009

Principal Investigators- William Knight and Manuel E. Ulibarri
Dexter National Fish Hatchery and Technology Center
U.S. Fish and Wildlife Service
P.O. Box 219, 7116 Hatchery Road
Dexter, NM 88230-0219
June 02, 2008

505-734-5910 Work
505-734-6130 Fax
william_knight@fws.gov
manuel_ulibarri@fws.gov

Background

Once very common throughout the Colorado River Basin, Colorado pikeminnow have declined from historic levels and are now found primarily in the Upper basin of the Colorado River. Various factors have contributed to the decline of the specie including alteration of natural stream flows and temperature regimes, loss of habitat and habitat fragmentation as a result of water development and the introduction of nonnative fish species.

Colorado Pikeminnow are native to the San Juan River. Its historic distribution included the entire mainstem San Juan River up to Rosa, New Mexico, located approximately 25 miles upstream from present day Navajo Dam. Currently the species is considered extremely rare and the small population is estimated at less than 20 adults. This small group of fish has persisted in the San Juan River since the closure of Navajo Dam in 1962. Recent studies being conducted by the San Juan Recovery Implementation Program (SJRIP) indicate that the Colorado pikeminnow is reproducing and recruiting in the river to at least a limited degree, however the low numbers collected do not satisfy recovery goal requirements for the specie. The Recovery criteria calls for a target of 1,000 subadult's fish established by the end of a five year down listing period, and 800 adults maintained during the 7 year delisting period. The Upper Colorado River Endangered Fish Recovery Program has recommended that the wild population be increased by augmenting with hatchery produced fish. The **Augmentation Plan For Colorado Pikeminnow In The San Juan River**, (Ryden 2003) calls for annual stocking of age-0 fish over an eight year augmentation program (2002-2009). Year 2009 completes the 8th year of the age-0 and 4th year of the age-1 augmentation program. As per the modified work plan, dated 6 April 2005 age-1 fish have been produced at Dexter and will be delivered annually from 2006-2009 to the San Juan River for stocking, (Ryden 2005, Addendum #1 to Augmentation Plan For Colorado Pikeminnow In The San Juan River).

Dexter NFH & TC has been the leader in propagating and culturing Colorado pikeminnow (*Ptychocheilus lucius*) since 1981. The facility maintains several captive stocks as genetic reserves and has successfully produced fish for the Upper and Lower Colorado river basin programs and the SJ RIP. The major emphasis has been on the reproductive biology, broodstock development and culturing age-0, 1 and adults. This work plan proposes to continue the production of 300,000 age-0 fingerlings (50 mm

TL) and 3,000 age-1 (150mm TL) fish annually for reintroduction in the San Juan River. In order to meet the target size and number of age-1 fish, Dexter will maintain approximately 6,000, young-of-year fingerlings annually for growout.

Funding requested also covers costs associated with proper care of broodstock necessary to successfully carry out this study for future years and aide in restoration of the species.

Stocking will require coordination with New Mexico FWCO, CRFP-Grand Junction, New Mexico Department of Game and Fish, Colorado Division of Wildlife and Utah Department of Wildlife Resources.

Objectives

- (1) Produce 300,000 age-0 fingerlings (50 mm) and 3,000 age-1 (150mm) fish for stocking in the San Juan River in 2009.
- (2) Continue data collection on induced spawning of Colorado pikeminnow under controlled conditions.
- (3) Provide staff to PIT tagging age-1 fish.
- (4) Transport and distribute 300,000 age-0 and 3,000 age -1 Colorado pikeminnow from Dexter to the San Juan River.
- (5) Maintain 400 Colorado pikeminnow broodstock for recovery efforts.

Methods

Broodstock will consist of 400 (F1) adults. These fish are and 1991 and 1999 year-class progeny from wild adults collected from the Yampa, Green and Colorado Rivers, respectively.

A maximum of 40 paired matings (1 female X 1 male) will be spawned during 2009. Given the past history of hormonal induced ovulation, 30 females (75%) should produce viable eggs during a given year. All members of the broodstock are PIT tagged and records of spawning pairs will be maintained at Dexter.

Ovulation will be induced with intraperitoneal injections of common carp pituitary (CCP) at the rate of 4 mg/kg of body weight. When eggs can be expelled using slight pressure, a female will be stripped and milt added from one male. Each individual egg lot will be enumerated and kept separate in Heath trays until hatching occurs, about 96 hours following fertilization.

When eggs begin hatching, larvae will be transferred to hatchery tanks and held until swim-up occurs, approximately five to seven days. Fry will be enumerated and stocked into four earthen ponds ranging from .33 to .35 ha. Fry will be cultured in earthen ponds for 120 days and age-1 fish for 240. The target size fish will be stocked in the San Juan River in October and November of 2009.

Budget -Detailed Spending Plan 2009

Labor Costs:

Fishery Biologist - GS-9	11.0 Pay Periods	\$32,063
Fish Health Biologist –GS- 9	6.0 Pay Periods	\$14,607

Duties/Activities Descriptions:

Drain broodstock pond and transferring adults to fish culture building
Inject males and females with hormones
Prepare egg hatching system
Spawn broodstock and return to holding pond
Place eggs in hatching system and care for eggs
Prepared holding tanks for fry
Transfer fry from incubators to holding tanks
Prepare pond to receive fry
Pond management
Transfer swim-up fry from holding tanks to ponds
Daily feeding (including weekends and holidays)
Weekly dissolved oxygen, temperature and pH recordings
Drain fingerling ponds and transfer to fish culture building
Prepare holding tanks for fingerlings
Inventory (weights and numbers) for each pond
Annual fish health inspection and treat fish for parasites if required and/or needed
Prepare broodstock holding pond prior to spawning
Drain broodstock pond and transfer adults to over-winter pond

Subtotal **\$ 46,670.00**

Equipment and Supplies:

Liquid oxygen and compressed oxygen 12 cylinders @ 72.23, Airgas	927.00
Fish health sampling prior to stocking Lab supplies for bacti, viral and parasite testing.	1,500.00
Culture equipment (nets, seines, screens, etc.) Eager, Memphis Net & Twine	1,250.00
Pond management supplies, Casoron, \$250/50# bag Van Diest	3,500.00
Fish feed, .85/lb, 4,000 lbs Nelson & Sons	3,400.00
Cyclical Maintenance costs for: Tractors, mowers, gators, sweepers used in pond maintenance	1,500.00

Subtotal **\$ 12,077.00**

Utilities:

Pumping costs Electrical 311,222 kwh @ .064	19,918.00
--	-----------

Heating water for hatching eggs to swim-up	
Natural gas 1,515 ccf @ .85	1,287.75

Subtotal	\$21,205.75
-----------------	--------------------

Reintroduction Costs:

Salaries

WG-8 Engineering Equip Operator 819.99	
36 hrs @ \$22.775	820.00

WG-7 Maintenance Worker	
36 hrs @ \$20.02	720.00

Overtime

WG-8 Engineering Equip Operator - 6 hours @ \$26.70	161.00
---	--------

WG-7 Maintenance Worker - 6 hours @ \$24.51	74.00
---	-------

Per Diem \$86/day (Dexter to Farmington, NM and return)

\$101.50/trip x 3 trips x 3 employees =	913.50
---	--------

Fuel costs and truck maintenance 1200 miles @ \$4.50	5,400.00
--	----------

Subtotal	8,088.50
-----------------	-----------------

Annual Totals (O & M Direct Costs)	88,041.25
---	------------------

17% Administrative Overhead	14,967.01
------------------------------------	------------------

*per cost recovery rates and policy (d-1 category)

TOTAL REQUESTED FOR 2009	\$103,008.26
---------------------------------	---------------------

Out year funding

Expected budget requirements for 2010 is: (@ 5% annual inflation rate)

Fiscal Year 2010 \$108,158.41

Literature Cited:

Ryden, D. W. 2003. An Augmentation Plan For Colorado Pikeminnow In The San Juan River. U. S. Fish and Wildlife Service, Grand Junction , Co. 63 pp. + appendices.

Ryden, D. W. 2005. *Draft* Addendum #1, Stocking Age-1 Fish To Supplement Ongoing Augmentation Efforts. An Augmentation Plan For Colorado Pikeminnow In The San Juan River. U. S. Fish and Wildlife Service, Grand Junction , Co. 3 pages.

**Rearing Razorback Sucker Sub-Adults at Dexter National
Fish Hatchery and Technology Center
FY 2009**

Prepared for:
U.S. Bureau of Reclamation RFP 04-SF-40-2250 and
The San Juan Recovery Implementation Program

Principal Investigators – William Knight, Manuel E. Ulibarri
Dexter National Fish Hatchery and Technology Center
P.O. Box 219, 7116 Hatchery Road
Dexter, New Mexico 88230-0219
505-734-5910
505-734-6130 Fax

William_Knight@fws.gov
Manuel_Ulibarri@fws.gov

Background

The following scope of work identifies the facilities and methodologies that will be used at Dexter to continue producing 11,000, 200+ mm razorback sucker for use by the San Juan River Recovery Implementation Program (SJRIP) to meet its augmentation objectives for the species in the San Juan River. The primary purpose being the distribution of these fish to existing grow-out ponds located on the Navajo Indian Irrigation Project. Dexter has developed production guides for the species based on historical growth rates and produces large numbers of razorback sub-adults (300+mm) for stocking into Lake Mohave, Arizona, Lower Colorado River.

The U.S. Fish and Wildlife Service (USFWS) has developed extensive infrastructure and expertise at DNFH&TC to successfully contribute to recovery programs. The DNFH&TC program has been totally devoted to the maintenance, propagation and culture of threatened and endangered fish species for thirty years. During that period it has successfully cultured razorback sucker, bonytail and Colorado pikeminnow of the Colorado River system. Currently DNFH&TC maintains a large genetically diverse razorback sucker broodstock and over the years has developed successful spawning, culture and distribution methodologies for the species that are still used today. The facility utilizes an abundant water supply to produce over 1 million fish annually.

Location

Dexter National Fish Hatchery and Technology Center is located in the Pecos River Valley of southeastern New Mexico, 200 miles southeast of Albuquerque, 20 miles south of Roswell, and one mile east of Dexter on State Road 190 (Figure 1.). The hatchery was authorized under the White Act of 1930 (46 Stat. 371 - 05/21/30), to meet the demands for warmwater game fish throughout the southwest. The land, originally the property of the New Mexico State Game and Fish Commission, was acquired by the Federal Government on August 31, 1931. The station lands comprise a total of 640 acres in Section 16, T13S, R26E, Chaves County. Construction of the facility started immediately after purchase with assistance of the Public Works Administration and Progress Administration. Operations began in 1932.

Facilities

Situated on the northern fringes of the Chihuahu Desert, the elevation at Dexter is 3,500 feet, average rainfall is 12 inches, and the growing season of 180-200 days. Station facilities include: Administration/Laboratory Building; Fish Culture Building; Visitors Center; Maintenance/Shop Building; Vehicle Storage Building; Equipment Storage Building; Feed Building; General Storage Building.; three government houses; one mobile home, two RVs and one RV space.

Fish culture facilities in operation during FY-2004 consisted of 71 earthen/lined ponds ranging in size from 0.1-1.0 acres, four (6' X 40') fiberglass raceways, four (8' X 40') concrete raceways, Twenty (2' X 12') rectangular fiberglass tanks, forty (4') fiberglass circular tanks, fifty (3') fiberglass circular tanks and 80 ten-gallon and 20 forty-gallon aquariums. The facility utilizes three water reuse systems in the fish culture building. Phase III Facility Improvement Project was completed on June 5, 2003.

Water

An abundant supply of fish culture water is supplied by five shallow aquifer wells (150 feet in depth) capable of pumping a combined 2,000+ gallons per minute. The well water is a constant 64^o F, pH of 7.5-8.5, total hardness of 2,100 ppm, and total dissolved solids of 3,500 ppm. Water rights, allocated through the New Mexico State Engineer's Office, total 2,185.5 acre-feet per annum or 10,927.5 acre-feet per five-year water period. Waste water from all fish culture operations collects in two sumps on the southeastern area of the facility and provides year round water to the wetlands.

Lake Mohave Razorback Broodfish

Staff at Dexter National Fish Hatchery and Technology Center successfully propagate and maintain 16 federally listed fish species; and produces over 1.0 million fish annually for recovery and restoration programs throughout the southwest (Appendix 1 and 2). Razorback sucker have been maintained and cultured at facility since 1981. Captive broodstock representing the Lake Mohave population exist at DNFH&TC. According to station records, the initial broodstock was founded with progeny from 136 wild adult fish collected from Lake Mohave in 1981.

An additional 147 wild individuals were collected from the lake in 1982, spawned that year, and contributed fry to the stocking efforts in the Gila, Salt and Verde rivers and Lake Mohave, but were not incorporated into the captive broodstock. In 1984, Dexter's RBS captive broodstock consisted of 360 three-year old fish derived from the wild adults spawned at Dexter in 1981. Wild caught adults collected in 1981 and 1982 had expired by the end of 1985. The first captive broodstock of RBS at

DNFH&TC is referred to as the '81 broodstock. The '81 broodstock currently contains 112 adult fish (Table 2). Initial spawning of this broodstock occurred in 1984 (Hamman 1985). It should be noted that no progeny of the '81 broodstock are currently held as broodstock at any facility. Since the broodstock's inception, all offspring have been stocked to meet production commitments. Over the past 19 years, offspring from this stock have been stocked into Lake Mohave and Lake Havasu; Gila, Salt, Verde and San Juan rivers; Niland-Imperial Valley Hatchery, California; Page Springs SFH, Arizona; Buenos Aires, Cibola, Imperial, Havasu National Wildlife Refuges; and the Colorado River Fisheries Project (CRFP) at Vernal, Utah. The second broodstock is referred to as the Paired Matings (PM) broodstock. This stock, comprised of approximately 90 unique family groups is the product of paired matings of wild caught adults spawned at Willow Beach NFH from 1994 to 2004. Those efforts resulted in 1,000 fish currently held as PM future broodstock at DNFHTC (Table 2). A third broodstock has been developed at DNFHTC, and consists of six year classes of juvenile wild-caught fish from Lake Mohave. These fish were captured as fry from eight locations throughout Lake Mohave and given the designation of Wild Caught (WC) future broodstock (Table 2).

Table 2. Dexter NFH & TC Razorback Sucker Captive Broodstock

<u>Year Class</u>	<u>Origin</u>	<u>Numbers on hand</u>	<u>Founders Represented</u>	<u>Lot Designation</u>
1981	F ₁ Mohave	112	adults / Mohave	'81
1994-2003	Mohave	1000	90 / Mohave	PM
1999-2004	Mohave	800	fry /Mohave	WC
2003	F ₂ Mohave	500	25/ '81 captive stock	F ₂

'81-1981 year class, Mohave-Lake Mohave, AZ, PM-Pair Matings, WC-Wild Caught, WB-Willow Beach, P- Production.

From 2001-2008 production of subadult razorbacks at DNFH&TC yielded excellent survival and growth. The overall survival for razorback sucker grown to 300mm was 88.5%, while 44.8% of the fish achieved the target growout size of 300 mm. DNFH&TC's spawning and growing season consists of fish being spawned in the early spring and fry stocked in to earthen or lined ponds and grown out-door from April to October. Total dissolved oxygen and temperature are monitored daily and fish feed on phyto and zooplankton produced in fertilized ponds for approximately 45 days at which time they are offered a prepared razorback sucker diet. Fingerlings are routinely held and cultured in the Fish Culture building during the months of January - March to prevent mortalities associated with outdoor over wintering. In the fall of the year when the fish reach target size they are harvested from the ponds and transferred to the Fish Culture building for sorting and tagging. Following a 7 to 10 day rest and recovery period they are loaded into distribution trucks and hauled to their stocking locations. Dexter has successfully hauled 300+mm razorbacks and Bonytail to Lake Mohave, Arizona, in the lower Colorado River. These distribution trips log 660 miles (12 hours) of hauling time in one direction.

Production Plan

Objectives:

The main objective of this proposed work is to spawn razorback sucker adults and rear 11,000, 200+mm fish annually and deliver them to existing grow-out ponds located on the Navajo Indian Irrigation Project. Additional objectives of the work include:

- (1) Improve, maintain and staff facilities at DNFH&TC to rear and distribute the target # of fish.
- (2) Annually provide 25,000 RBS larvae to the Uvalde NFH for growout.

(3) Continue data collection on stocking densities in Dexter ponds for optimal growth of razorbacks and evaluate and adjust as necessary to meet required numbers and size.

(4) Maintain razorback sucker captive broodstock for recovery efforts.

Methods

DNFH&TC will conduct captive propagation activities that include spawning of a minimum of 25 pairs of broodstock, incubation of fertilized eggs, enumeration and stocking of swimup fry into DNFH&TC ponds, harvest of target sized fish from ponds, enumeration and distribution to Navajo Indian Irrigation Project.

The project will utilize indoor and outdoor facilities. All spawning and incubation activities will be conducted indoor in the fish culture building. Razorback sucker will be reared in four earthen and lined ponds at surface acres of 0.72, 0.79, 0.82, and 0.86.

Spawning

Broodfish will be harvested from ponds in early March and held indoor for spawning. Razorback sucker spawning protocols developed at DNFH&TC that will be used are listed in Appendix Table 3.

Rearing Ponds

To meet the production goal of 11,000 (200mm) fish, rearing ponds will be stocked at the following densities:

Age 0 Growth: (April thru May - 60 day growing period)

Pond 1- .72 acre @ 15,000 fry
Pond 2- .79 acre @ 15,000 fry

Age I Growth : (June thru October - 150 day growing period)

Harvest Age I fish; enumerate and stock fingerlings into 4 ponds.

Pond 1- .72 acre @ 6,000 fingerlings
Pond 2- .79 acre @ 6,000 fingerlings
Pond 3- .82 acre @ 6,000 fingerlings
Pond 4- .86 acre @ 6,000 fingerlings

Earthen and lined ponds will be used for production. In earthen ponds the bottoms will be packed and graded prior to receiving fish. Non-level pond bottoms can hinder fish harvest and aquatic vegetation can entrap fish at harvest time. Fertilization and slow filling of ponds will start 10 to 14 days prior to stocking. Staff will ensure that water quality is monitored. Temperature, dissolved oxygen and pH readings will be taken twice daily at 7:00am and 3:00 pm at the deepest part of the pond.

If the dissolved oxygen drops to ≤ 3 mg/l, supplemental aeration will be started. All feeding, fertilization and chemical applications will be stopped till adequate oxygen levels are restored. Aerators will be run all night for several days till the oxygen is back up to acceptable levels, (5-7 mg/l). Staff will avoid handling fish for 7 -10 days following a stress related circumstance.

Pond Vegetation Control and Fertilization

Sonar, Diuron or Barrier will be used in earthen ponds to control rooted aquatic vegetation. Staff will use granular form when possible and broadcast the entire pond bottom at the recommended rates.

Sonar - 20 lbs per acre (dry broadcast)
Barrier- 100 lbs per acre(dry broadcast)

Copper sulfate (CUSo4) will be used to control floating filamentous algae blooms. Treatments will began approximately 45 days after fish are stocked into the ponds and repeated every 30 days. Application rates in DNFH&TC ponds are 3 to 5 lbs per acre. A secondary benefit derived from using CUSo4 is its effectiveness in controlling external parasites.

Zooplankton and invertebrate insect populations are cultured with the proper fertilization regime. Four types of fertilizer will be used:

- 1) Alfalfa meal
- 2) Alfalfa pellets
- 3) Cottonseed meal
- 4) Super phosphate

Initial fertilization rates for earthen ponds are 100 lbs of cottonseed meal, 100 lbs of alfalfa meal or pellets and 3 lbs of super phosphate. Follow up rates are administered on Monday and Thursday with 10 lbs cottonseed meal, and 10 lbs, alfalfa meal or pellets.

Water temperature, dissolved oxygen (DO) and pH readings will be taken in all rearing ponds daily. All readings will be recorded on record charts. If morning DO readings are below 3.0 or above 13.0 all fertilization will be stopped until DO's are brought back to accepted levels. If pH readings are greater than 9.5 fertilization will be terminated.

Feeding Schedule

Fish will be sampled at the end of every month. Size, weight and over all condition will be recorded. Feed amounts will be adjusted and projected for the upcoming month. Razorback grower 0301 feed will be used and purchased from Nelson and Sons, Silver Cup, Murray, Utah. Fish will be fed twice daily, once at 10:00am and at 2:00pm.

Feeding rates are based on water temperature and fish densities in the ponds and will be calculated as follows:

- water temp ≥ 70 °F feed 3 % BW per day, Mon thru Fri.
- water temp 60-70 °F feed 2 % BW per day, Mon thru Fri.
- water temp < 60 °F feed 1.5 % BW per day, Mon, Wed, Fri.

Staff will use the following guide to determine the proper particle size to offer the fish. Feed sizes will be mixed at ½ rations of each size when making the transition to the next larger size feed.

<u>Fish Size</u>	<u>Particle Size</u>
2-3"	1.0 mm
4-6"	2.0 mm
6-8"	3.0 mm

Projected Harvest Dates and Delivery Date

Year 2009 marks the fourth year of razorback production at Dexter for distribution to the NAPI ponds. Since 2006, Dexter staff have stocked a total of 14,000 razorback's averaging 170mm in length into East and West Avocet and Hidden ponds. An additional 11,000 are scheduled to be stocked in the spring of 2009. Over the past two years the facility also provided over 150,000 razorback larvae to the Uvalde NFH for growout for the SJRIP.

Based on historical growth rates for razorback at Dexter, the production target of 11,000, 200+mm fish is achieved in a fifteen month period. In 2007 a new single cohort fish rearing strategy was adopted by the SJRIP for the NAPI ponds. Fish delivery will be in the spring of each year based on the new rotational production plan (single cohort). Approximately 11,000 fish will be stocked each trip and Dexter staff will

coordinate the deliveries with the Navajo Nation Department of Fish and Wildlife, BIA and USFWS FWCO personnel. The estimated duration of the program is scheduled for a total of 15 years (2005- 2020).

Predator Control

Historically, DNFH&TC has not experienced excessive avian or mammal predation on fish stocks. Salamander, crayfish, frog and turtle infestation of ponds are nonexistent. On an annual basis specific ponds are covered with bird netting during the winter months to eliminate predation by migrating birds. An additional strategy employed by the staff is the harvest and hold stocks of fish indoor during the winter months of November to March. Razorback reared for this project will be maintained indoor in two 40,000 gallon recirculating systems during the winter months. The recirculating systems contain biofiltration, supplemental aeration, temperature control and alarm systems.

Handling and Transport Protocol

Transport of all fish will follow guidelines described in the USFWS Protocols for Biological Investigations developed by Dr. Gary Carmichael, retired U.S. Fish & Wildlife Service employee. The protocol is as follows:

1. When razorback fingerlings, subadults and broodfish are handled they will be placed in a .5% salt bath to help in osmoregulation and reduce the effects of handling stress.
2. Temperature should be 5 degrees Fahrenheit lower in the hauling truck than in the river.
3. Drivers must be informed of and follow a specified route.
4. Transport water will contain 0.5 percent NaCl (18.9 grams per gallon) and 0.26 ml/L Stress Coat7 (1 ml per gallon).
5. Oxygen levels will be greater than 6.0 mg/L as determined with an oxygen meter.
6. Nets must be functional. Aeration equipment must be in place and must be used. A fish holding container will be a minimum of 5 gallons in size and fish densities will not exceed 1 lb of fish per gallon of water. Small delta mesh (1/8") will be present to transfer the fish from one container to another., although it is preferred to have water to water transfer. Oxygenation/aeration equipment will be in place and working.
7. Prior to transfer and after the fish are concentrated, they should be quickly placed in the transport tank. When using nets to place fish in transfer buckets or tanks, nets should not be overloaded. The fish on the bottom will be crushed. Using a wet transfer with buckets is preferable. When emptying the nets and buckets, care will be taken to avoid adding algae and mud to the transport tank. Before loading, dissolved oxygen levels should be at saturation.
8. Immediately after loading, all equipment on the transport vehicle should be re-checked and the vehicle should depart. Oxygen concentrations and temperatures should be monitored at a minimum of every hour.
9. During unloading tempering water should be present and functional, and thermometers should be used to match water temperatures. Hauling water temperatures should be equal to receiving water temperature.

***Acclimatizing the fish to the receiving water temperature will be conducted in increments of 2 degrees towards equalizing per 15 minutes time. Due to the high alkalinity and TDS of DNFH&TC water, staff will temper and acclimate the transported fish to the receiving water quality for a minimum of 1 hour prior to release. This process will allow sufficient time for the fish to osmoregulate to the receiving water quality. Tempering can be accomplished in the shipping tank by adding receiving water to the tank at given intervals.**

Fish Health Monitoring Protocols

All fish should be handled with the best animal husbandry practices available. A feeding schedule will be developed and followed daily. All tanks will be cleaned of uneaten food and feces daily. A daily log recording times of feeding, water temperature and comments on fish health will be maintained. If fish are maintained in a re-circulating system, all filters and pumps will be routinely cleaned and monitored. If fish are held in ponds O2 levels will be closely monitored. At least once a year, a fish health inspection will be conducted to examine fish for bacterial, viral and parasitic infections. Normally 60 fish per lot are sacrificed for an adequate sample. However, in the case of endangered or rare fish of genetic importance, numbers sampled may be less, depending upon availability. Non-lethal methods, if available, will be employed to obtain samples. Condition factors will be calculated on an annual basis and data added to a RBS database. Wet mounts will be examined for parasites and bacteria. Routine condition exams will be conducted and an examination will be conducted on all lots one month prior to delivery to the Navajo Indian Irrigation Project, SJRIP. Brood and refuge stock will have health checks annually and only when needed to minimize handling stress.

The U.S. Fish and Wildlife Service, Regional Fish Health Unit @ Dexter will provide bacterial and viral testing for razorback propagation and rearing activities. Treatment of disease will be the responsibility of the Dexter staff. Fish health experts are available to advise on proper treatment, and to examine fish for infection.

Budget

RE: RFP #04-SF-40-2250, Rearing Razorback Sucker Sub-Adults at Dexter National Fish Hatchery and Technology Center, Costs associated with rearing 11,000 – 200mm fish for NAPI ponds and producing 25,000 larvae for Uvalde NFH annually. Detailed Budget Spending Plan, 2009.

O&M Labor Costs

The labor costs identified in the 2009 Scope of Work are broken down as follows, and include fringe benefits and payroll additives for each position identified:

Dexter National Fish Hatchery & Technology Center

(1) Bio Science Tech (FR) (1,280 hours- 16pp) - GS 404-5 @	\$18.75/hr =	\$24,000
* Fish health and water quality monitoring, feeding, harvest and distribution.		
(1) Fishery Biologist (480 hours -3pp) - GS 482-9 @	\$28.50/hr =	\$6,840
* Supervision, spawning and distribution coordination.		
(1) Administrative Officer (320 hours- 2pp) - GS 341-9 @	\$26.95/hr =	\$4,321
* Budget tracking, purchasing, data base management & reporting.		
	Subtotal =	\$35,161

Materials and Supplies

Cost based on Dexter NFH&TC historical purchases:

Fish Health

-Lab supplies (pipets, petri dishes, slides, probes, markers)	\$	250
-Theriputents- salt, furacin, formalin, MS-222, stress coat	\$	500
- Liquid and compressed oxygen for fish distribution	\$	200

Feed

Production diet RBS0301 (1.5tons) 3000 lbs \$.81 per lb	\$	2,430
--	----	-------

Spawning Supplies

Hormones (HCG 10 vials @ \$ 50 per 10ml/vial)	\$	500
---	----	-----

Fertilizer		
Alfalfa pellets (1,000 lbs) .25/lb		\$ 250
Inorganic - Super Phosphate (10 bags) 7.50/bag		\$ 75

Chemicals Aquatic Vegetation Control		
Barrier- (5 bags) \$250/bag		\$ 1,250

Subtotal = \$ 5,455

Services

Utilities & Equipment Maintenance		
* Electrical, fuel and phone		\$ 2,750
* Boiler system, heat exchanger maintenance		\$ 1,000
*#4 well and water tower and pumping station maintenance		\$ 4,500

Subtotal = \$ 8,250

Travel

- Fish stocking/distribution.		
Dexter to Farmington & return- (820miles @ 4.50 per mile X 2 trips)=		\$ 7,380
Fuel and routine vehicle maintenance.		
Perdiem- \$120 per day X 2 trips X 2 individuals. =		\$ 480
Dexter to Uvalde & return- (960miles @ 4.50 per mile X 1 trip)=		\$ 4,320
Fuel and routine vehicle maintenance.		
Perdiem- \$120 per day X 1 trip X 1 individual. =		\$ 120

Subtotal = \$12,300

Annual Totals

O&M DIRECT COSTS		\$61,166
FY 2009 CPI-U (West Region, All items, 1982-1984=100) 3.2%		\$ 1,957
		\$63,123
INDIRECT COSTS (Admin Overhead @ 17%)		\$10,730
*per cost recovery rates and policy (d-1 category)		

TOTAL REQUESTED FOR 2009 \$73,853

Out Year Funding

Expected budget requirements for: @ CPI-U (West Region, All items, 1982-1984=100) 3.2%)		
Fiscal Year 2010	\$76,216	
Fiscal Year 2011	\$78,655	
Fiscal Year 2012	\$81,172	

Projected Duration Of Project:

This project was initiated in January 2005 in support of the SJRIP razorback augmentation effort (2004-2011) identified in the **Five-Year Augmentation Plan For Razorback Sucker In The San Juan River** (Ryden 1997). The rearing of razorback sucker subadults at Dexter NFH&TC could potentially continue till 2020 as per BOR RFP 04-SF-40-2250.

Reporting

Quarterly progress reports detailing fish culture and distribution activities will be completed and synthesized into a final accomplishment report available to the SJRIP by January 31, 2010.

Schedule

Broodfish will be spawned in March 2009 and the fish reared in earthen ponds for their first growing season (April - October 2009); held indoor during winter (November 2009- March 2010) stocked into lined ponds in March 2010. Target sized fish are available for distribution in spring and fall of each year.

Personnel Qualifications

Manuel E. Ulibarri, Center Director

Education:

B.S. 1985, Biology, Western New Mexico State University
1986 to 1988 Graduate work in Fisheries Science, New Mexico State University

Professional Experience:

Dexter NFH & TC - 2001 to present (EOD at Dexter NFH & TC 11/04/01)
Willow Beach NFH - 1998 to 2001
Uvalde NFH - 1991 to 1998
Mescalero NFH - 1986 to 1991
Rock Lake State Fish Hatchery, Santa Rosa, NM 1981 to 1984

Roger L. Hamman, Assistant Center Director

Education:

B.S. - 1975, Fishery Biology, Southeastern Oklahoma State University
B.S. - 1975, Wildlife Biology, Southeastern Oklahoma State University
A.S. - 1973, Murray State College

Professional Experience:

Dexter NFH & TC - 1986 to present
Dexter NFH - 1982 to 1986
Willow Beach NFH - 1982 to 1982
Leetown Fish Hatchery Manager Long Course - 1981 to 1982
Willow Beach NFH - 1978 to 1981
Tishomingo NFH - 1975 to 1978,

William Knight - Fisheries Biologist

Education:

B.S. 2005 – Fisheries and Wildlife Science, New Mexico State University

Professional Experience:

Uvalde NFH April – July 2006
NMSU A-Mountain Native Fish Facility 2001 to 2005

Janell Desmond - Administrative Officer

Education:

B. S. 1975- Education
Computer programming - college course, New Mexico Military Institute

Professional Experience:

Dexter NFH & TC - 2006 to present
Bitter lake National wildlife Refuge 1999 - 2006
Bureau of Land Management 1998 – 1999
Roswell Independent School district 1997 – 1998
Ozark National Scenic Riverways 1994 - 1997

Appendix Table 3.**Dexter National Fish Hatchery and Technology Center
Razorback Sucker Spawning Protocols: 2005**

Roger Hamman

March 1, 2005 ▪screen, board and start filling broodstock summer pond.	▪check eggs in incubators ▪individual egg lots can be moved at this time ▪prepare a minimum of two 12' tanks to receive fry
March 12, 2005 ▪start draining broodstock pond	March 19, 2005 ▪check eggs in incubators
March 13, 2005 ▪continue draining broodstock pond	March 20, 2005 ▪check eggs in incubators begin filling 12' tanks with heated water
March 14, 2005 ▪Harvest pond and bring all broodstock in to Fish Culture Building ▪sort males/females and place in separate tanks ▪record pit tag numbers, lengths, weights and take genetic samples of each fish ▪inject 25 females with 0.1cc HCG/lb in preparation for spawning ▪inject 25 males if necessary with 0.3 cc HCG/lb in preparation for spawning ▪Move all broodstock not used in spawning activities to summer pond.	March 21, 2005 ▪check incubators (morning and afternoon) and transfer fry to 12' tanks
March 15, 2005 ▪inject 25 females with 0.1cc HCG/lb	March 22, 2005 ▪check incubators (morning and afternoon) and transfer fry to 12' tanks
March 16, 2005 ▪inject 25 females with 0.1cc HCG/lb ▪prepare incubation system to receive eggs ▪gather other equipment and supplies needed for spawning trials	March 23, 2005 ▪transfer remaining fry to 12' tanks ▪clean incubators
March 17, 2005 ▪spawn razorbacks using 1 female X 1 male spawning procedure ▪inventory each individual spawn ▪place eggs in incubators	March 24, 2005 ▪observe fry in 12' tanks
March 18, 2005 ▪move spawned broodstock to summer pond	March 25, 2005 ▪observe fry in 12' tanks
	March 26, 2005 ▪observe fry in 12' tanks
	March 27, 2005 ▪observe fry in 12' tanks ▪clean 12' tanks in preparation for stocking fry into rearing ponds
	March 28, 2005 ▪fry stocked into rearing ponds at 20,000 per acre.

Literature Cited:

- Hamman, R. 1985. Induced spawning of hatchery-reared razorback sucker. Prog. Fish-Cult.. 47(3): 187-189
- Ryden, D. W. 2003. An augmentation plan for razorback sucker in the San Juan River: An addendum to the five-year augmentation plan for razorback sucker in the San Juan River (Ryden 1997). U. S. Fish and Wildlife Service, Grand Junction, CO. 32 pp.

FY 2009

**Rear 12,000-300mm Razorback Sucker and Assess Potential for Rearing
Bonytail at the Uvalde National Fish Hatchery, Uvalde, Texas**

Prepared for:
Biology Committee
The San Juan Recovery Implementation Program

Principal Investigator – Grant L. Webber
Uvalde National Fish Hatchery
754 County Road 203
Uvalde, Texas 78801
830-278-2419
830-278-6042 Fax

Grant_Webber@fws.gov

Introduction

Uvalde National Fish Hatchery (UNFH) submits the following proposal to rear 12,000 300mm razorback sucker sub-adults annually for the San Juan River Recovery Implementation Program (SJRIP) and conduct research activities related to rearing 300mm bonytail for the Lower Colorado River Multi-Species Conservation Program (LCRMSCP). The project will use 15 one acre ponds at the UNFH, Uvalde, Texas. Dexter National Fish Hatchery and Technology Center (DNFH&TC) will provide fry and/or fingerlings of both species to UNFH, as well as technical assistance with fish health and culture methods.

The following scope of work identifies the facilities and methodologies that will be used at UNFH to produce the target number of razorback sucker and conduct research on bonytail growth and performance. An initial production guide was developed for the species based on historical growth rates observed at Dexter, Willow Beach, and Achii Hanyo. The data generated from the past three years of work completed at Uvalde has been incorporated into the current razorback and bonytail production program. Funding is being requested for operations at UNFH. The UNFH will provide the infra-structure for stability in the production program. Fish hauling will be conducted by the Regional Distribution Unit (RDU) stationed at Inks Dam NFH; Burnet, Texas. Based on the number and size of fish to be stocked, when necessary, Uvalde will be available to assist in the transportation of the fishes.

Background

UNFH is located 3 miles Southwest of Uvalde, Texas on FM 481 and approximately 80 miles West of San Antonio. This is a large warm water fish culture facility that utilizes earthen and lined ponds, and intensive culturing raceways to produce fish.

The hatchery is situated on 100 acres of Mesquite Grasslands, in the Rio Grande Plain of Southwest Texas. There are 47 usable ponds totaling 50 surface acres of water. Five ponds were lined with high density polyethylene in FY 1987 and six more in FY 1993 for water conservation purposes. Buildings on the facility were renovated from 2001 to present, following a 100 year flood in 1998. Hatchery facilities include an office/fish culture building, shop/garage, fish holding house/nursery, feed room with cooler, two labs, and four living quarters with two double garages, two pump houses, and four concrete/two fiberglass raceways. Water for fish culture purposes is pumped from two deep wells. Two water towers provide a back up water source for intensive culture purposes.

Station Operations

Historically UNFH has been one of the top producing warm water fish culture facilities in the nation. During the mid 90's as many as 6 species were cultured producing 2.6 million fish, weighing 60,000 pounds. Over the past 15 years threatened and endangered fish species like Yaqui catfish, paddlefish, Comanche Springs pupfish and fountain darters have all been propagated and maintained successfully at the facility.

The climate in Southwest Texas provides 300 days (10 months) of growing season. Two deep wells provide 72° water year round.

Razorback sucker and bonytail have been reared at UNFH, since April 2006. On November 11th, 1,150 PIT tagged 300mm Age-1 razorbacks were stocked in the San Juan River (Hogback diversion area). The fish were marked with 134 kHz tags provided by the SJRIP. In 2006, 16% of all razorbacks stocked into Uvalde ponds reached the 300mm target size in six months. Approximately 75% of the remaining fish were 250+ mm in length. These fish were kept on station for future grow

out and eventual stocking in 2007. In 2007, Uvalde stocked approximately 5,000 razorbacks and with additional fish sent to Dexter NFH & TC in March of 2007, exceeded the 6,000 fish commitment. No fish were stocked in 2008 due to the facility testing positive for Largemouth Bass Virus. In February 2008, all contaminated sources were destroyed or removed from the hatchery. With two successful negative viral test results (July 2008 & July 2009), stocking of 12,000 razorback suckers is anticipated to commence during the fall of 2009, which is within this SOW activity period. Bonytail growth and survival was excellent as well. On November 16th, 2,397 300mm bonytail were stocked into the Lower Colorado River, near Park Moabi. Approximately 80% of all bonytail stocked into Uvalde ponds reached the 300mm target size in six months. Survival over the 180 day growing period for all ponds averaged 92%.

Facilities

This project will utilize up to 20 ponds to fulfill the production and research commitments of the proposal. These ponds require maintenance earth work to the bottoms and banks and are fully functional with water supplies, catch basins and drains.

Water

An abundant amount of fish culture water is supplied by two wells on station. The first and primary water supply (Spurgeon Well) derives its water from the Austin Chalk formation, which is capable of a deep aquifer well capable of pumping approximately 1,500 gallons per minute. The well water is a constant 72°F, pH of 7.5-8.05, total hardness of 496 ppm, and alkalinity of 224 ppm. A second well, named the Wilson Well is now operational and can produce an additional 1,500 gallons per minute. The well has a year round temperature of 75 °F, pH of approximately 7.1, total hardness of 380 ppm, and alkalinity of 245 ppm. The water system was recently designed to produce up to 3,000 gallons per minute flow throughout the hatchery. Due to the potential implications to threatened and endangered species utilizing the Edwards water throughout the system, the Service has established a limit on groundwater withdraws from the Edward's Aquifer. That self-imposed water right for Uvalde is 843 acre-feet per annum. Since the Spurgeon Well derives its water from a source other than the Edward's Aquifer (Austin Chalk formation), there are no groundwater pumping limits set for withdraws associated with the Spurgeon Well.

Lake Mohave Razorback and Bonytail Broodfish

The DNFH&TC has successfully propagated and maintained razorback sucker and bonytail broodstock at the facility since 1981. Captive broodstock representing the Lake Mohave population exist at DNFH&TC and will be spawned and their progeny will be transported to Uvalde in the form of fry/fingerlings for growout and research purposes.

Uvalde's growing scenario includes receiving fry and fingerlings from DNFH&TC in April. The fish are stocked into earthen or lined ponds and grown outdoors from April to November. Total dissolved oxygen, temperatures, and pH are monitored daily. Fish are fed phyto- and zooplankton produced in fertilized ponds for approximately 45 days at which time they are offered a prepared razorback sucker diet. In the fall of the year when the fish reach target size they will be harvested from the ponds and transferred to the fish culture building for sorting and tagging. Following a 7 to 10 day rest and recovery period they will be loaded into distribution trucks and hauled to their stocking locations by the Service's RDU. The RDU successfully hauled 300+mm bonytail to the Lower Colorado River in 2006 and 2007. These distribution trips log 660 miles (12 hours) of hauling in one direction.

Objectives

There are two main objectives of this SOW: 1) propagate 12,000 – 300mm razorback sucker sub-adults annually and deliver them to the San Juan River; and 2) determine optimal captive propagation densities for bonytail at UNFH that will enable the consistent production of 300mm fish annually for augmentation programs in the Lower Colorado River (LCRMSCP).

Additional objectives include:

1. Improve, maintain, and staff Uvalde NFH necessary to captively rear and distribute the target # of fish and conduct the target research;
2. Test effects of long distance hauling, water quality differences and elevation on RBS and BTC cultured at the UNFH;
3. Determine survival rates over time of fish hauled from UNFH to DNFH&TC.

Methods

DNFH&TC will conduct captive propagation activities that include spawning of a minimum of 25 pairs of broodstock, incubation of fertilized eggs, enumeration and providing swim up fry/fingerlings fish to UNFH for the completion of the objectives. UNFH will conduct the propagation and harvest of target sized fish from ponds, enumerate, tag and coordinate the distribution of these fish to the San Juan and Lower Colorado Rivers.

The project will utilize both indoor and outdoor facilities. At Dexter, all spawning and incubation activities will be conducted indoor in the fish culture building. At Uvalde, razorback sucker and bonytail will be reared in extensive culturing ponds and intensive culturing raceways to achieve target numbers and sizes.

Spawning

Broodfish will be harvested from DNFH&TC ponds in mid March and early April and held indoor for spawning. Over the next week eggs will be incubated and swim up fry/fingerlings shipped to Uvalde for rearing. This scenario will continue till adequate numbers of fingerlings are held at UNFH to rear to the target size. The facility will maintain at least 50,000 age I+ razorbacks and 30,000 bonytail in a given year to meet commitments for future years.

Ponds

Razorback Rearing

Sufficient numbers of razorback fingerlings were cultured at UNFH during the 2006 and 2007 growing season to meet requirements of a five year production program. Approximately 100,000 fry will be shipped every other year from Dexter to UNFH in order to continue the production cycle for the future. To meet the production goal of 12,000 (300mm) fish annually, the rearing ponds will be stocked at approximately the following densities:

Age 0 Growth: (April thru October -- approximate 214 day growing period)

3 acres @ 33,000 fry (every other year)

Age I Growth: (March thru November – approximate 244 day growing period)

Harvest Age I–fish from the ponds, enumerate and stock into 11 ponds.

Pond 7-	1 acre @ 4,500	fingerlings
Pond 8-	1 acre @ 3,600	fingerlings
Pond 10-	1 acre @ 3,500	fingerlings
Pond 11-	1 acre @ 3,400	fingerlings
Pond 18-	1 acre @ 2,300	fingerlings
Pond 19-	1 acre @ 3,600	fingerlings
Pond 20-	1 acre @ 3,600	fingerlings
Pond 26-	1 acre @ 6,900	fingerlings
Pond 30-	1 acre @ 5,700	fingerlings
Pond 36-	1 acre @ 3,600	fingerlings
Pond 37-	1 acre @ 3,600	fingerlings

Bonytail Research

Stocking densities per pond will be adjusted annually based on the previous year's fish growth and survival. (April thru November - 210 day growing period)

Pond 25-	1 acre @ 1,500	fingerlings
Pond 27-	1 acre @ 1,000	fingerlings
Pond 28-	1 acre @ 1,000	fingerlings
Pond 29-	1 acre @ 1,500	fingerlings
Pond 35-	1 acre @ 1,000	fingerlings
Pond 38-	1 acre @ 1,500	fingerlings

Any earthen pond used will be packed and graded prior to receiving fish. Non-level pond bottoms can hinder fish harvest and aquatic vegetation can entrap fish at harvest time. Lined ponds will be cleaned out every other year to reduce the amount of organic material in the ponds which could cause water quality deterioration. Fertilization and slow filling of ponds will start 10 to 14 days prior to stocking. Staff will ensure that water quality is monitored. Temperature, dissolved oxygen and pH readings will be taken twice daily at 7:00am and 3:00 pm at the deepest part of the pond.

If the dissolved oxygen drops to ≤ 3 mg/l, supplemental aeration will be started. All feeding, fertilization and chemical applications will be stopped till adequate oxygen levels are restored. Aerators will be run all night for several days till the oxygen is back up to acceptable levels, (5-7 mg/l). Staff will avoid handling fish for 7 -10 days following a stress related circumstance.

Pond Vegetation Control, Water Quality and Fertilization

Sonar, Diuron, Reward, Hydrothol, and cutrine plus and will be used in earthen ponds to control rooted aquatic vegetation. Staff will use granular form when possible and broadcast the entire pond bottom at the recommended rates.

- Sonar - 20 lbs per acre (dry broadcast)
- Diuron- 25 lbs per acre (dry broadcast)
- Citrine plus- 60 lbs per acre (dry broadcast)
- Hydrothol 191- 80 lbs acre (dry broadcast)

Copper sulfate (CuSo₄) will be used to control floating filamentous algae blooms. Treatments will began approximately 45 days after fish are stocked into the ponds and repeated every 30 days. Application rates in Uvalde ponds are 2 to 3lbs per acre. A secondary benefit derived from using CuSo₄ is its effectiveness in controlling external parasites.

Zooplankton and invertebrate insect populations for razorback and bonytail Age-0 will be cultured with the proper fertilization regime. Age-I fish are fed a prepared diet.

Four types of fertilizer will be used:

- 1) Alfalfa meal
- 2) Alfalfa pellets
- 3) Cottonseed meal
- 4) Super phosphate

Initial fertilization rates for earthen ponds are 100 lbs of cottonseed meal, 100 lbs of alfalfa meal or pellets and 3 lbs of super phosphate. Follow up rates are administered on Monday and Thursday with 10 lbs cottonseed meal, and 10 lbs, alfalfa meal or pellets.

Water temperature, dissolved oxygen (DO) and pH readings will be taken in all rearing ponds daily. All readings will be recorded on record charts. If morning DO readings are below 3.0 or above 13.0 all fertilization will be stopped until DO's are brought back to accepted levels. If pH readings exceed 9.5, fertilization will be terminated.

Escapement

Staff will reduce the potential for escapement by installing drain screens in the ponds when they are initially prepped to receive fish. Screen mesh size will be 250 micron in Age-0 ponds and ¼" in Age-I ponds. All Age-I fish will be graded prior to being stocked in the rearing ponds. Staff will monitor the ponds daily and insure there are no leaks in the dam boards. Sawdust will be used to stop all leaks that develop in the catch basin. Water levels will adjusted and maintained a minimum of six inches below the over flow mark until the fry average 30mm in length.

Feeding Schedule

Fish will be sampled at the end of every month. Size, weight and over all condition will be recorded. Feed amounts will be adjusted and projected for the upcoming month. Fry and fingerlings will receive a starter grower diet purchased from Nelson and Sons, Silver Cup, Murray, Utah. Fry will be fed 4 times daily and fingerlings twice daily, once at 10:00am and at 2:00pm.

Feeding rates are based on water temperature and fish densities in the ponds and will be calculated as follows:

- Water temp > = 70 °F (21 °C) feed 3 % BW per day, Mon thru Fri.
- Water temp 60-70 °F (16-21 °C) feed 2 % BW per day, Mon thru Fri.
- Water temp < 60 °F (16 °C) feed 1.5 % BW per day, Mon, Wed, Fri.

Staff will use the following guide to determine the proper particle size to offer the fish. Feed sizes will be mixed at ½ rations of each size when making the transition to the next larger size feed.

<u>Fish Size</u>	<u>Particle Size</u>
(Fry-2")	starter and #1, 2, & 3 crumbles
2-3"	1.0 mm

4-6"	2.0 mm
7-9"	3.0 mm
9-14"	4.0 mm

Projected Harvest Dates and Delivery Date

Based on harvest data from 2006 and 2007 the production target of 12,000- 300mm fish can be achieved in a fourteen month period. Fish will be harvested from the ponds and stocked into the San Juan River in October/November of each calendar year. Some fish may also be stocked in the spring months, should some achieve the target size sooner than expected. In order to establish a consistent long term production cycle UNFH will maintain 30,000 to 36,000 Age-I fish on station in a production year.

All bonytail will be harvested from the study ponds in the fall of each year and transferred to the fish holding house for enumeration. Length/weight, survival and fish health data will be collected, analyzed and adjustments made to the stocking densities for the following year. All fish achieving the target size of 300mm will be available for use in the augmentation effort of the LCRMSCP program. The projected date of stocking is November.

Predator Control

During the summer grow-out all ponds are monitored daily by on-site staff and predators are taken by rifle and traps. Fish remaining outdoors during the winter season will be under the protection of 2" X 2" block nylon bird predation netting. A minimum of 12,000 razorbacks and 6,000 bonytail reared for this project will be maintained in intensive culture raceways during the winter months. These intensive culture facilities contain 72°F flow through water, supplemental aeration, power back-ups, and a security alarm system.

Fish Health Monitoring Protocols

All fish will be handled with the best animal husbandry practices available. A feeding schedule will be developed and followed daily. All fish holding facilities on station will have their O₂ levels monitored daily. At least once a year, a fish health inspection will be conducted to examine fish for bacterial, viral and parasitic infections. Normally 60 fish per lot are sacrificed for an adequate sample. However, in the case of endangered or rare fish of genetic importance, numbers sampled may be less, depending upon availability. Non-lethal methods, if available, will be employed to obtain samples. Condition factors will be calculated on an annual basis and data added to a RBS and BTC database. Wet mounts will be examined for parasites and bacteria. Routine condition exams will be conducted and an on-site comprehensive examination will be conducted on all lots one month prior to delivery to the San Juan River. Brood and refuge stock will have health checks annually and only when needed to minimize handling stress.

The Region 2 Fish Health Unit @ Dexter will provide bacterial and viral testing for razorback propagation and rearing activities. Treatment of disease will be the responsibility of the Uvalde staff. Fish health experts at Dexter are available to advise on proper treatment and to examine fish for infection.

2009 Production and Distribution Schedule

Broodfish will be spawned in April, 2009 at Dexter NFH & TC; 100,000 Age-0 fry will be hauled to UNFH and stocked into three ponds to continue the production cycle. Age-I fish currently at Uvalde will be stocked into clean ponds in March/April 2009 and available for distribution in November 2008 (late fall and or as requested by the SJRIP).

Budget 2008 Fiscal Year

Rearing Razorback Sucker and Bonytail Sub-Adults at Uvalde National Fish Hatchery, Detailed Budget Spending Plan, 2009.

O&M Labor Costs

The labor costs identified in this proposal are broken down as follows, and include fringe benefits and payroll additives for each position identified:

Uvalde National Fish Hatchery

(1) Fishery Biologist (26 pp) - GS 482-9 @ \$29.35/hr =	\$62,000
* On-site fish rearing, water quality monitoring, vegetation treatment, fish marking and distribution coordination.	
(1) Animal Caretaker (26 pp) – WG-5048-5 @ \$18.09/hr =	<u>\$38,000</u>
-General fish husbandry activities such as fish feeding, chemical applications, and water quality data collections	
Subtotal =	\$100,000

Equipment, Materials and Supplies

Cost based on UNFH historical purchases:

Administrative supplies

- Paper, pens, folders, & miscellaneous office supplies \$ 2,000

Fish Health

-Water quality monitoring supplies (probes, meters, test pillows) \$ 1,000

-Therapeutants- salt, formalin, MS-222, Stress Coat, Nitrofurazone \$ 3,000

Feed

-Production diet RBS # 350 -25,000 lbs @ \$.85 per lb \$21,250

Fertilizer

-Alfalfa pellets (3,000 lbs) .25/lb \$1,500

-Inorganic - Super Phosphate (10 bags) \$10.00/bag \$ 200

Chemicals- Aquatic Vegetation Control and other water quality

-Hydrothol 191, Copper Sulfate, Citric Acid, Cutrine Plus, Navigate \$10,000

Subtotal = \$38,950

Services

-Utilities & Equipment Maintenance

* Electrical, fuel and phone

\$25,000

* Well and associated equipment

\$1,500

Subtotal = \$26,500

Travel

- Fish stocking/distribution (USFWS Regional Distribution Unit)

Haul tagged fish from Uvalde to Farmington & return to Burnett, Texas.

Fuel, routine vehicle maintenance, and salary, overtime and

Per diem (2 individuals)

\$ 10,000

Subtotal = \$10,000

TOTALS:

O&M DIRECT COSTS

\$175,450

INDIRECT COSTS (Admin Overhead @ 17%)

*per cost recovery rates and policy (d-1 category)

\$29,827

TOTAL O&M REQUESTED FOR 2008

\$205,277

San Juan Recovery Program portion (62%)

\$127,272

LCRMSCP portion (38%)

\$78,005

**Razorback Sucker Augmentation at NAPI Grow-Out Ponds
Fiscal Year 2009 Project Proposal**

Principal Investigators: Jeff Cole, Viola Willeto and Albert Lapahie
Navajo Nation Department of Fish and Wildlife
P.O. Box 1480
Window Rock, AZ 86515
(928) 871-7068

jcole@navajofishandwildlife.org ywilleto@navajofishandwildlife.org
alaphie@navajofishandwildlife.org

and

D.Weston Furr, Ernest Teller, Sr. and Jason E. Davis
United States Fish and Wildlife Service
New Mexico Fish and Wildlife Conservation Office
3800 Commons N.E.
Albuquerque, NM 87109
(505) 342-9900

Weston_Furr@fws.gov

eteller@wildblue.net

Jason_E_Davis@fws.gov

Background

The Long Range Plan for recovery of endangered fishes in the San Juan River calls for propagation and augmentation of razorback sucker (RBS). Nine ponds have been built on Navajo Agricultural Products Industry (NAPI) lands to grow out RBS for stocking into the San Juan River. The Coordination Committee has decided to only utilize three of the nine existing ponds on NAPI during FY 2009.

Avocet Pond was originally a single pond built for watering cattle. On March 2, 1998 Avocet was divided into 2 ponds known as Avocet East and West. Avocet West is 3.4 acres and holds 18 acre-feet of water. Avocet West has a siphon for draining the pond. Avocet East is 3.52 acres and holds 19.6 acre-feet of water. Avocet East has no siphon, so draining is accomplished by renting a battery of water pumps. A siphon will be installed in Avocet East during FY 2008.

In October of 1999, Hidden Pond was built to rear razorback sucker. Hidden Pond is 2.83 acres. The dam was breached due to a storm event and the fish were lost. The dam was re-built in FY 2000 and a toe drain and spillway were built to protect the dam. Hidden Pond was lined with bentonite and contoured and a kettle was installed to facilitate fish harvest. A siphon was installed in July 2003. A salamander fence was installed around the Hidden Pond perimeter in August of 2003 to exclude predatory tiger salamanders.

Responsibility for Management of the NAPI ponds was originally shared between the U.S. Fish and Wildlife Service (Service), Bureau of Indian Affairs (BIA), Keller-Bliesner Construction and Ecosystems Research Institute. The Service was responsible for determining which ponds would receive RBS and when. In addition, the Service conducted sample counts and harvested the ponds with the assistance of the BIA. Keller-Bliesner was responsible for design and construction of the Six Pack ponds and re-construction of Hidden Pond. The BIA was responsible for monitoring water quality and Ecosystems Research was responsible for fertilization of the ponds and for developing a pond management plan.

Original pond management was for multiple cohorts to be raised in the ponds. Harvesting would be done passively with fyke nets so that the ponds would not be drained on an annual basis. In FY 2007, it was determined to change pond management direction. All of the ponds would be drained and harvested and

single cohort management would replace the multiple cohort approach. During the first harvesting and draining of a Six-pack Pond, high mortality resulted when the number of fish remaining in the pond could not be removed before they succumbed to the rapidly warming water. Adjustments were made to reduce the mortality in future harvesting and draining events. The adjustments consisted of increasing the trapping effort prior to de-watering to reduce the number of fish remaining in the pond. In addition, the final fish removal would be accomplished with a higher pool of water to slow the warming of the water during the time of final harvest. This resulted in less mortality.

The Navajo Nation Department of Fish and Wildlife (NNDFW) was contracted to assume responsibility for daily management of the NAPI ponds in 2007. The Service assists the NNDFW with pond harvest as needed.

The ponds have been fenced and electric lines have been installed at each of the ponds. Aerators have been installed at each of the ponds to improve water quality. Water quality issues have caused fish mortalities in some of the ponds in the past. Water quality issues appear to have been resolved since installation of the aerators.

Objectives

(NAPI Ponds Management)

Cooperatively manage East Avocet, West Avocet, and Hidden ponds to provide an additional source of RBS to supplement the RBS augmentation program. Harvest, Passive Implant Transponder (PIT) tag, and stock razorback sucker from the three grow-out ponds into the San Juan River, in order to assist in fulfilling the tasks and objectives outlined in the current version of *An Augmentation Plan for Razorback Sucker in the San Juan River* (Ryden 2003).

- 1) Manage three grow-out ponds using a single cohort strategy; including passive and active harvest techniques.
- 2) Annually stock 3,500 ($\geq 200\text{mm}$) razorback sucker per pond.
- 3) Harvest all ponds on an annual basis.
 - a. Implant all razorback sucker with a PIT tag prior to stocking.
 - b. Stock all fish regardless of size at harvest.
 - c. Stock ~ 4,200 to 6,300 fish based on 40-60% return.

Location

The RBS grow-out ponds are located in Block III of Region 2 on NAPI lands, south of Farmington, New Mexico. Avocet East and West are located NW of the intersection of N 4062 and N 4087, which is approximately 3 miles southwest of the Ojo Amarillo NHA Housing Subdivision. Hidden Pond is located SE of the intersection of N 4087 and N 4095 approximately 1 mile northwest of the NAPI Region II Complex.

Methods/Approach

The Service, Region 2, will provide overall coordination for management of the grow-out ponds on NAPI. The NNDFW will be responsible for daily management of the three grow out ponds on NAPI with assistance by the Service, Region 2. Harvesting, tagging, and stocking will be conducted by the two Service Regions and NNDFW. Associated data management and reporting for the project will be handled by staff from the Service, Region 2.

Pond management requires that staff monitor and record water quality and quantity, and feed the fish on a daily basis. In addition, staff manages water quantity to ensure that water quality is optimal. Maintenance includes operating and repairing valves and aerators, evaluating the pond perimeters for erosion problems, operating the propane cannons to scare away predators, repairing fences, monitoring aquatic vegetation and maintaining a log book and database for management of the ponds.

During FY 2009, East Avocet, West Avocet, and Hidden ponds will be managed for a single cohort of RBS. NNDFW and Service staff will cooperatively trap, tag, and stock RBS into the SJR for several days prior to dewatering the ponds. As the ponds are dewatered, NNDFW and Service staff will work together to do the final RBS removal, tagging, and stocking into the SJR.

Whenever the ponds are drained, they will be evaluated for structural stability. Areas away from ponds that may be impacted by dewatering will also be evaluated. Staff will identify and document any structural damage to the ponds and dewatering areas if necessary. Feasibility will determine whether improvements are made or not. This proposal does not include any maintenance or repair work that is major and requires mobilization of heavy equipment and is outside of the constraints of this budget.

Products/Schedule

In the spring of 2009, Dexter National Fish Hatchery will deliver 10,500 \geq 200 mm RBS to the three NAPI grow-out ponds. In the fall of 2009, the NAPI ponds will be de-watered and the RBS, which are targeted to be \geq 300 mm will be harvested and transported to the San Juan River for stocking. A database summarizing numbers of fish, stocking locations and PIT tag numbers will be submitted to the SJRIP Program Coordinators Office by 31 March 2010. A draft report will be submitted by 31 March 2010 and finalized by 1 June 2010.

Budget Fiscal Year 2009

BUDGET WORKSHEET – Program Base Funding		
Razorback Sucker Augmentation at NAPI Grow-Out Ponds		
Personnel (salary/benefits)	USFWS NMFWCO	NNDFW
Daily Pond Management .30 FTE (GS-9-8) USFWS R2 and active/passive harvesting assistance .5 FTE NNDFW (19,000 x 32.15%)	\$ 25,168	\$ 25,109
Wildlife Technician .5 FTE NNDFW (11,000 x 32.15%)		\$ 14,537
Personnel Subtotal	\$ 25,168	\$ 39,646
Travel		
Per Diem Lodging and Meals	\$ 4,251	\$ 500
Vehicle Mileage and Maintenance	\$ 3,450	\$ 13,000
Travel Subtotal	\$ 7,701	\$ 13,500
Office Supplies and Equipment		\$ 500
General Operating Supplies (includes fish transport costs, i.e. oxygen, salt, stress coat, etc.)		\$ 2,500
Electricity Costs (Aeration)		\$ 2,500
Feed Cost (\$1.55/lb – 5,000 lbs)		\$ 7,750
Uniforms		\$ 500
Printing/Binding/Photocopying		\$ 100
Fuel – Propane/Cannon Guns		\$ 200
Repairs and Maintenance – Paint, sealant, lubricants, plumbing supplies, water quality probes, etc.		\$ 500
Support Subtotal	\$ -0-	\$ 14,550
Budget Subtotal	\$ 32,869	\$ 67,696
FY 2007 Carry over funds	-0-	-0-
Total	\$ 32,869	\$ 67,696
Administrative charge (22%)	\$ 7,231	\$ 14,893
USFWS/NNDFW Totals	\$ 40,100	\$ 82,589
Grand Total		\$ 122,689

Note: Budget was reduced to \$120,689 to account for \$20,000 of carry-over from FY08

Itemized budget for USFWS - NMFWCO:

Daily pond management activities .30 FTE (GS-8; \$55,553/year)	\$ 16,666.00
Active/Passive Harvest	
Fish Biologist (GS-7-2) - 39 days @ \$218/day (passive harvest – 5 days x 3 ponds x 2 trips) (active harvest – 3 days x 3 ponds x 1 trip)	\$ 8,502.00
Personnel subtotal	\$ 25,168.00
Travel and Per Diem (Based on Published FY-2008 Federal Per Diem Rates)	
Hotel Costs – 60 nights (39 nights @ \$70/night – single occupancy = \$2,730)	\$ 2,730.00
Per Diem (Hotel Rate) – 39 days @ \$39/day	\$ 1,521.00
Travel subtotal	\$ 4,251.00
Equipment	
Vehicle Maintenance & Gasoline 5,750 miles @ \$0.60/mile (based on Anticipated GSA rates established on 01 September 2008 and includes costs associated with gasoline/diesel fuel vehicle maintenance)	\$ 3,450.00
Equipment subtotal	\$ 3,450.00
USFWS – NMFWCO Total	\$ 32,869.00
USFWS Region 2 Regional Office Administrative Overhead (22.00%)	\$ 7,231.00
USFWS – Region 2 Total	\$ 40,100.00

FY 2009 Project Proposal
San Juan River Basin Hydrology Model Operation and Maintenance
SJRIP - Hydrology Committee

Katrina Grantz UC-432
Bureau of Reclamation
125 South State Street
Salt Lake City, UT 84138-1147
Phone 801-524-3635
Fax 801-524-3847
kgrantz@uc.usbr.gov

Ryan Christianson
Bureau of Reclamation
835 E. 2nd Avenue, Suite #300
Durango, CO 81301
Phone 970-385-6590
rchristianson@uc.usbr.gov

Relationship to SJRIP: Supports Program goals and management by developing, operating and maintaining a hydrology model of the San Juan Basin. The model is key to hydrological analysis of water development scenarios or other scenarios in relation to the flow recommendations.

Background:

The Third Generation San Juan Basin Hydrology Model (SJBHM) is a daily timestep hydrologic model of the San Juan Riverbasin. Though major model development was essentially completed in FY2004, revisions and modifications have been made throughout the multi-year validation and verification phase of model implementation. As of May 2008, the Third Generation SJBHM has not yet been approved by the San Juan River Basin Recovery Implementation Program. This scope of work includes the annual operation and maintenance of the model and necessary data. The Bureau of Reclamation has the primary responsibility for model O&M.

Once approved, the model will be available to generate and analyze runs associated with Section 7 Consultations and/or special requests from the Hydrology, Biology, or Coordination Committees related to the flow recommendations or other hydrological aspects of the Program.

Objective:

The objective for this work is to ensure that the San Juan Basin Hydrology Model is available for run requests. This will be accomplished by adjusting model configurations or operating rules to correct for errors or other changes, and evolving the data set forward through time. In addition, the FY2009 request includes funds to continue to provide technical transfer from the model developers to the model users and maintainers.

Task Description and Schedule:

Task 1: Data Maintenance Maintain data to evolve the data set forward through time. This includes an annual update of USGS data, Reclamation data, New Mexico non-irrigation data, New Mexico irrigation data, Arizona and Utah depletions, Colorado depletions, and climate data. Data must be obtained from various sources and processed for compatibility with the multiple data loaders.

Task 2: Model Maintenance Load updated data into the model, run and test the new data. Adjust model configuration, methodologies, or assumptions, as needed. Update and expand documentation to reflect current state of model.

Task 3: Software Maintenance Update and maintain data management interfaces and other software associated with the data and models. Apply all RiverWare updates and patches as they become available.

Task 4: Model Runs and Analyses Generate and analyze model runs associated with Section 7 consultations or special requests from the Biology, Hydrology, and/or Coordination Committees. A consultation run usually requires model reconfiguration and the implementation of operating criteria. Each consultation request will require approximately twelve staff days; each special run will require approximately five staff days.

Task 5: Program management and coordination Attend Hydrology Committee meetings to provide model status updates and present results on HC run requests. Develop the FY2010 budget and track FY2009 expenditures.

Task 6: Technology Transfer Provide technology transference to Reclamation's Upper Colorado Regional Office and Western Colorado Area Office staff in the details of maintaining the data and models, and in operating the models. Technology transfer will continue to ensure that several people are trained in the operation and maintenance of the model.

Budget:

Task 1 Data Maintenance

A) Labor

Task	Position	Salary total/hr	Total Hours	Total Cost
Annual Update of Data	UCRO Engineer ¹	\$55.00	160	\$8,800.00
	WCAO Engineer ²	\$70.00	40	\$2,800.00
Total				\$11,600.00

B) Travel

Position	Destination	Purpose	Days	Lodging per Day/ Total	Per Diem per Day/ Total	Airfare Other* Total	Total
WCAO Engineer	Salt Lake City	data update	2	\$110	\$55	\$60 \$1,000	\$1,390
Total				\$220	\$110	\$60 \$1,000	\$1,390

* Taxi \$20; Parking \$10; Rental car \$50/trip

Task 2 Model Maintenance

A) Labor

Task	Position	Salary total/hr	Total Hours	Total Cost
Annual Update of Model	UCRO Engineer	\$55.00	200	\$11,000.00
	TSC Engineer ³	\$125.00	40	\$5,000.00
Documentation	UCRO Engineer	\$55.00	200	\$11,000.00
	TSC Engineer	\$125.00	40	\$5,000.00
Total				\$32,000.00

B) Travel

No travel

Task 3 Software Maintenance**A) Labor**

Task	Position	Salary total/hr	Total Hours	Total Cost
Software updates	UCRO Engineer	\$55.00	56	\$3,080.00
Total				\$3,080.00

B) Travel

No travel

C) Other Costs

RiverWare Technical Support \$5,000

Task 4 Model Runs**A) Labor**

Task	Position	Salary total/hr	Total Hours	Total Cost
Model Runs and Analyses	UCRO Engineer	\$55.00	360	\$19,800.00
	WCAO Engineer	\$70.00	40	\$2,800.00
	TSC Engineer	\$125.00	80	\$10,000.00
Total				\$32,600.00

B) Travel

No travel

Task 5 Program Management Coordination**A) Labor**

Task	Position	Salary total/hr	Total Hours	Total Cost
Meetings	UCRO Engineer	\$55.00	80	\$4,400.00
	WCAO Engineer	\$70.00	80	\$5,600.00
	TSC Engineer	\$125.00	24	\$3,000.00
Budget	UCRO Engineer	\$55.00	40	\$2,200.00
Total				\$15,200.00

B) Travel

Position	Destination	Purpose	Trips/ Days per Trip	Lodging per Day/ Total	Per Diem per Day/ Total	Other*	Airfare Total	Total
UCRO Engineer	Farmington	HC Meetings	2	\$100	\$40			
				\$200	\$160	\$120	\$1,600	\$2,080
TSC Engineer	Farmington	HC Meetings	1	\$100	\$40			
			2	\$100	\$80	\$120	\$800	\$1,100
UCRO Engineer	Durango	CC Meeting	1	\$100	\$40			
			3	\$200	\$120	\$60	\$800	\$1,180
Total								\$4,360

* Taxi \$20; Parking \$10; Rental car \$50/trip

Task 6 Technology Transfer and Support**A) Labor**

Task	Position	Salary total/hr	Total Hours	Total Cost
Technology Transfer	UCRO Engineer	\$55.00	160	\$8,800.00
	WCAO Engineer	\$70.00	96	\$6,720.00
	TSC Engineer	\$125.00	160	\$20,000.00
Total				\$35,520.00

B) Travel

Position	Destination	Purpose	Days	Lodging per Day/ Total	Per Diem per Day/ Total	Other*	Airfare Total	Total
WCAO Engineer	Salt Lake City	tech transfer	2	\$110	\$55			
				\$220	\$110	\$60	\$800	\$1,190
UCRO Engineer	Denver	tech transfer	2	\$140	\$50			
				\$280	\$140	\$60	\$300	\$780
TSC Level 3 Eng.	Salt Lake City	tech transfer	2	\$110	\$55			
				\$220	\$110	\$60	\$800	\$1,190
Total								\$3,160

* Taxi \$20; Parking \$10; Rental car \$50/trip

Budget Summary**FY 2009**

Labor	Task 1 - Data Maintenance	\$11,600
	Task 2 - Model Maintenance	\$32,000
	Task 3 - Software Maintenance	\$3,080
	Task 4 - Model Runs	\$32,600
	Task 5 - Program Management	\$15,200
	Task 6 - Tech Transfer	\$35,520
Total labor		\$130,000
Travel	Task 1 - Data Maintenance	\$1,390
	Task 5 - Program Management	\$4,360
	Task 6 - Tech Transfer	\$3,160
Total Travel		\$8,910
Total Other - RiverWare Tech Support		\$5,000

Grand Total **\$143,910** *

* This total budget represents a 74% increase over the FY2008 budget.

FY-2010	\$151,106	**
FY-2011	\$158,661	**
FY-2012	\$166,594	**

** Includes ~5% adjustment

** assumes that tech transfer is continuing in out-years

¹ Upper Colorado Regional Office (Salt Lake City)

² Western Colorado Area Office (Durango)

³ Technical Services Center (Denver)

**Improve Stream Gaging and Flow Measurements
San Juan River Basin Recovery Implementation Program - Hydrology Committee
Fiscal Year 2008 Project Proposal**

Principal Investigator: Pat Page
Bureau of Reclamation
835 E. 2nd Avenue, Suite #300
Durango, CO 81301
(970) 385-6560 ppage@uc.usbr.gov

Background

There are five USGS streamflow gaging stations on the main stem of the San Juan River that are very important to the operation of the river and play an important role in the implementation of the flow recommendations. Four gages are in New Mexico and one is in Utah. These include:

- SJR @ Archuleta (NM) - a few miles below Navajo Dam; used to measure releases from the dam
- SJR @ Farmington (NM) - just downstream from Animas confluence; furthest upstream in critical habitat
- SJR @ Shiprock (NM) - in critical habitat
- SJR @ Four Corners (NM) - in critical habitat
- SJR @ Bluff (UT) - located at Mexican Hat; furthest downstream gage in critical habitat

Stream gaging data on the San Juan River are needed to attempt to reliably develop and implement flow recommendations.

Study Area

San Juan River Basin in New Mexico

Objective

Provide funding to the USGS to take additional flow measurements as needed at the four San Juan River gages in New Mexico. (Note: Base cost for operation of the stations is paid for by non-Program funds.)

Products

1. Improved flow measurement and more accurate gage readings.
2. Technical presentation at the end of the year from USGS summarizing the activities completed and the value of obtaining additional readings.

Budget Fiscal Year 2008

Objective: Provide funding to USGS for additional flow measurements at the 4 San Juan River Gages in NM.	Staff days	Labor	Travel	Equipment and supplies
Personnel	7.5	5,500		
Travel			1,200	
Equipment and supplies				
Total				\$6,700

Estimated Outyear Funding (Based on 5% allowance for inflation)

Fiscal Year 2009	\$7,000
Fiscal Year 2010	\$7,400
Fiscal Year 2011	\$7,800
Fiscal Year 2012	\$8,200

**Operation of Public Service Company of New Mexico Fish Passage Structure
Fiscal Year 2009 Project Proposal**

Principal Investigators: Jeffrey Cole, Albert Lapahie, Viola Willeto, Navajo Nation Department of Fish
and Wildlife Box 1480 Window Rock, AZ 86515 (928) 871-7068

jcole@navajofishandwildlife.org, alapahie@navajofishandwildlife.org

vwilleto@navajofishandwildlife.org

Background:

The Power Company of New Mexico (PNM) Diversion Dam was constructed in 1971. The 3.25-foot high diversion dam (weir) is located on the San Juan River about 12 miles downstream of Farmington, New Mexico near the town of Fruitland at River Mile 166.6. Facilities at the diversion include a concrete weir, a series of screened intake structures, an intake channel, a settling channel, and a pump house.

Water flows over the dam into a stilling basin created by a concrete apron. The stilling basin is the width of the river. The presence of the dam and the basin creates a barrier to fish moving upstream. As flows increase, the difference in the upstream and downstream water levels is reduced. Although water levels are reduced, water velocities increase and the weir provides an impediment to upstream fish movement. Recovery studies conducted as part of the SJRRIP have shown that some fish are able to move upstream past the weir but their specific method of movement is not known and the number of fish discouraged from upstream movement by the presence of the weir is also unknown. One possible method of upstream movement could occur during high river flows. When the flow in the San Juan River is above 7,000 cfs, some of the flow goes around the dam making it possible for fish to go around the dam at these higher flows.

A need has been identified by the San Juan River Basin Recovery Implementation Program (SJRRIP) to restore endangered fish passage upstream past the PNM Diversion Dam. The purpose of establishing fish passage was to protect and recover native Colorado pikeminnow (*Ptychocheilus lucius*) and razorback sucker (*Xyrauchen texanus*) populations in the San Juan Basin while water development proceeds in compliance with all applicable Federal and State laws, including fulfillment of Federal trust responsibilities to the Southern Ute Indian Tribe, Ute Mountain Ute Tribe, Jicarilla Apache Nation and the Navajo Nation. In addition, other native fish species would benefit from restored passage. The facility has been operated and maintained by the Navajo Nation Department of Fish and Wildlife (NNDFW) since it was built in 2003. The U.S. Fish and Wildlife Service (Service), Bureau of Reclamation (BOR), Bureau of Indian Affairs (BIA), Navajo Indian Irrigation Project (NIIP), Navajo Agricultural Products Industry (NAPI), and PNM have provided the NNDFW with technical assistance, planning assistance, environmental clearance, maintenance and improvements to the facility and its access points.

The fish passage has facilitated movement of pikeminnow and razorback suckers upstream into a 50 mile stretch of river, which is historical habitat of these species.

Study Area:

Public Service Company of New Mexico Diversion Dam is located at RM 166.6.

Methods/Approach:

The Fish Passage facility will be operated from April 1 to October 31, 2009. The fish passage traps fish attempting to move upstream of the facility. All fish that are caught in the trap are transported to a sorting tray. All fish are identified and enumerated. Non-endangered native fish are released upstream of the

facility. Rare native fishes are scanned for a pit tag, weighed and measured, marked with a pit tag if they do not have one and then released upstream of the facility. All non-native fishes are removed from the river system permanently. When feasible, channel catfish are transported to area fishing lakes that already have channel catfish in their systems to support the sport-fishing program.

Daily operation and maintenance includes cleaning of surface and submerged trash, debris, silt, and river-born algae from the trash racks and bar screens in the forebay of the fish passageway, and aluminum conduit screens in the fish trap. The amount of algae, debris, trash, and sediment that accumulates daily at this site is seasonally variable, depending upon flow magnitude and water volume during the water year. Maintenance also includes painting as necessary to control corrosion, lubrication of moving equipment, and checking fluid levels in gearboxes and cooling radiators, as necessary. Representatives from the NNDFW, BOR, PNM and the Service will perform an inspection of the facility every 3 years. In the event of a significant flood event, representatives from the NNDFW will notify BOR, PNM and FWS and appropriate parties will inspect the facility for damage, as necessary.

The Fish Passage Program maintains a database of all fish processed through the facility. Staff that operate this facility also have initiated a public outreach and education program that will continue in FY' 2009. School groups visit the facility to learn about the purpose of the facility and the endangered fish program on the San Juan River.

Objectives of this project are as follows:

- 1.) Determine the use of the fish passage by juvenile and adult native and nonnative fishes.
- 2.) Identify any Colorado pikeminnow congregations that may be related to the spawning period in the San Juan River.
- 3.) Maintain the facility in a manner that assures long-term benefit.

This proposal does not include any maintenance or repair work that is major and requires mobilization of heavy equipment and is outside of the constraints of this budget.

Products/Schedule:

The Fish Passage facility will be operated from April 1 to October 31, 2009. Data will include definitive numbers of species, numbers per species, and seasonal use and distribution by species.

NNDFW staff will prepare and submit monthly reports and one draft and final annual report. Service staff will assist NNDFW with data analysis and draft and final report preparation.

Program staff will attend SJRRIP Biology Committee meetings and provide reports as needed throughout the year. Service staff will assist the NNDFW in preparing presentations for the Biology Committee meetings.

Budget Fiscal Year – 2009

BUDGET WORKSHEET		
Operation of San Juan/PNM Fish Passage		
Personnel (salary and benefits)	USFWS Funding	Program Base Funding
.5 FTE Fisheries Biologist (19,000 X 32.15%)	\$	\$25,108
.5 FTE Wildlife Technician (11,000 X 32.15%)		\$14,536
Fringe Benefits \$39,644 X 32.15%		\$12,745
Personnel Subtotal		\$52,389
Travel		
1 Tribal Vehicle		\$15,000
Per Diem Lodging and Meals		\$2,500
Travel Subtotal	\$	\$17,500
Office Supplies		\$500
Office Equipment – LCD Projector and screen		\$1,500
General Operating Supplies Plumbing supplies, Hardware Supplies, Neoprene Waders, rubber boots, wet suit, landscaping supplies		\$2,500
Uniforms		\$500
Printing/Binding/Photocopying		\$100
Fuel – Gasoline for water pump		\$300
Sewage Services – Fish Passage		\$700
Repairs and Maintenance – Paint, sealant, lubricants, water pump repairs		\$1,000
Support Subtotal	\$	\$7,100
Training and Conference Registration		\$1,000
Consultant/ Professional Sub-Total		\$1,000
	USFWS Funding	Base Funding
Budget Subtotal		\$ 77,989 65,244
FY 2006 Carry over funds		0
Total		\$ 77,989 65,244
Administrative charge (22%)		\$ 17,157 14,354
Grand Total	\$	\$95,146 79,598

Note: Budget was adjusted to \$79,598

**San Juan Recovery and Implementation Program
Conservation Action Planning:
Analysis of Potential San Juan Ecosystem Restoration Sites
& Hazardous Material Threats
2008 Project Proposal**

Principal Investigators:
Adrian Oglesby & Rex Wahl
The Nature Conservancy
212 E. Marcy Street, Suite 200
Santa Fe, New Mexico 87501
505-280-7958
aoglesby@tnc.org

Background

The goals of the San Juan Recovery River Implementation Program (“SJ RIP”) Long Range Plan include providing suitable habitat and suitable water quality to support recovery of the Colorado pikeminnow and razorback sucker, including identification of types and sources of water quality contamination (Tasks 3.2.2, 3.4.1 and 3.4.3, SJ RIP Draft Long-Range Plan).

Accomplishing these goals remains pertinent to the recovery of the Colorado pikeminnow and razorback sucker. The SJ RIP is increasingly recognizing that flow regimes alone may not be sufficient to create and sustain the habitat necessary for the endangered fish. In addition, the 1994 San Juan Water Quality Contaminants and Review discusses sampling results from the early 1990’s that indicate San Juan fish have been exposed to hydrocarbons. Since that time development of both conventional oil and gas basins and coal bed methane deposits have continued.

In an effort to further the SJ RIP’s progress in achieving this goal, The Nature Conservancy proposes to initiate conservation planning activities that will include the evaluation of the need physical habitat enhancements and conduct an analysis of hazardous materials threats and impacts to the San Juan River ecosystems.

Goals

The primary goal of this scope of work shall be to utilize existing data and analyses of essential habitats, flow-habitat relationships, and endangered fish distributions to identify locations for habitat improvement projects that will use mechanical means to assist flows in the creation or improvement of endangered fish habitats.

A secondary goal shall be to analyze existing and proposed hazardous material threats to the San Juan River ecosystems; identify best management practices; and, make recommendations to minimize the risks of hazardous material impacts on the San Juan River.

Deliverable

Within one year of project initiation, a report will be provided with maps and GIS layers that identify and rank locations for potential mechanical habitat improvement projects.

If time and funding allows, a second report will be provided that identifies current hazardous material threats and impacts; the current local, State and Federal authorities over hazardous materials; and, provides recommendations for securing, monitoring and researching hazardous materials.

Study Area

San Juan River Basin, focusing in New Mexico.

Study Methods/Approach**Habitat Restoration Planning**

- 1) Review existing research, literature and regulatory databases pertaining to endangered species habitat and flow preferences in the San Juan;
- 2) Develop target habitat types;
- 3) Develop site evaluation criteria¹;
- 4) Conduct preliminary identification of potential habitat restoration sites based on aerial photographs and existing field data;
- 5) Conduct field surveys for potential habitat restoration sites and inspections of potential sites identified during data and aerial photography analysis;
- 6) Identify locations of discharges, heavy equipment use, tanks, drilling operations, storage pits, product pipelines and related infrastructure within the 100 year floodplain;
- 7) Consolidate analysis and develop site recommendations and priorities for mechanical habitat restoration projects;
- 8) Provide information consistent with related SJRIP activities (e.g., NMED River Ecosystem Restoration Initiative project).

If time and funding allow after completion of the above tasks, the following activities will be incorporated into this project as possible.

Hazardous Materials Threats Assessment

- 1) Review existing research, literature and regulatory databases pertaining to contaminants in the San Juan;
- 2) Identify all hazardous waste regulators with jurisdiction;
- 3) Provide overview of relevant regulators' authorizations;
- 4) Review and describe historic and current non-compliance issues;
- 5) Review and analyze State and Federal hazardous materials spills response plans and implementation procedures;
- 6) Consolidate data and analysis; and,

¹ Examples of potential evaluation criteria include suitability of habitat; landowner cooperation; potential longevity of restoration; accessibility for equipment, stocking, monitoring, and maintenance; engineering and construction costs; etc.

- 7) Develop recommendations for better identifying, securing, monitoring, and researching hazardous materials.

Budget

The salary, benefits and travel expenses of one Principal Investigator will be contributed by The Nature Conservancy.

Principal Investigator and GIS Professional Salary and Benefits	\$4,600
Supplies, GIS Shape Files, Aerial Photographs, and Database Reports	\$12,000
Travel	\$1,000
Indirect Costs (Overhead)	<u>\$2,400</u>
TOTAL	\$20,000

**Non-native species Monitoring and Control in the Upper San Juan River
Fiscal Year 2009 Project Proposal**

Principal Investigators: Jason E. Davis, D. Weston Furr
U.S. Fish and Wildlife Service
New Mexico Fishery Resources Office
3800 Commons N.E.
Albuquerque, NM 87109
(505)342-9900

[Jason E Davis@fws.gov](mailto:Jason_E_Davis@fws.gov) [Weston Furr@fws.gov](mailto:Weston_Furr@fws.gov)

and

Dale Ryden
U.S. Fish and Wildlife Service, Colorado River Fishery Project
764 Horizon Drive, Build B
Grand Junction, CO 81506-3946
(970) 245-9319
dale_ryden@fws.gov

Background

The August 1, 2001 Colorado pikeminnow and razorback sucker Recovery Goals identified predation by and/or competition by nonnative fish species as a primary threat to the continued existence or the reestablishment of self-sustaining populations of these endangered fishes. In addition, reducing the impacts of nonnative fishes has been identified as a critical element in the San Juan River Basin Recovery Implementation Programs Long Range Plan (Draft April 2008). Actions and Tasks associated with this Element encompassed within this scope of work include:

- Action 4.1.1 Develop, implement, and evaluate the most effective strategies for reducing problematic nonnative fish.
 - Task 4.1.1.4 Evaluate and refine alternative nonnative fish reduction methods.
 - Task 4.1.1.7 Evaluate effect of nonnative fish control on distribution, abundance, and demographics (e.g., fish size, age, sexual maturity) of nonnative fish populations.
 - Task 4.1.1.8 Evaluate effect of nonnative fish control on distribution, abundance, and demographics (e.g., fish size, age, sexual maturity) of the endangered fish populations and native fish community.
- Action 4.1.4 Establish criteria for reduction of target nonnative fish populations.
 - Task 4.1.4.1 Establish target criteria for reduction of problematic nonnative fish species to estimate time, effort, and cost for controlling nonnative fishes.

Intensive removal of non-native fishes, primarily channel catfish and common carp, has occurred in the upper reaches of the San Juan River since 2001. Between 2001 and 2003, removal trips focused on a 7.6 mile reach of river from PNM Weir (RM 166.6)

to Hogback Diversion (RM 159.0). Declines in catch rates, seasonal movement by channel catfish, and high abundance of non-native fishes downstream of Hogback Diversion prompted removal efforts to expand in 2003 to include the adjacent downstream reach, Hogback Diversion to Shiprock Bridge (RM 147.9). These efforts are ongoing with a total of eight (three passes/trip) trips divided between both reaches annually.

Multi-pass removal efforts were successful, to a degree, in suppressing non-native numbers within intensive removal reaches (Davis 2006; Jackson 2006). However, long term trend data collected during annual fall monitoring trips indicate an apparent increase in channel catfish abundance riverwide beginning in 2004. Much of this can be attributed to increased abundance of channel catfish in reaches that are between (RM 147.9 – 52.9) those where intensive removal efforts occur (Ryden 2006). Prior to 2006, non-native fishes within this portion of the San Juan River were only opportunistically removed during spring razorback sucker and fall annual monitoring trips.

Beginning in 2006, U.S. Fish and Wildlife Service (FWS) – New Mexico Fishery Resources Office (NMFRO) shifted removal efforts to include two trips from Shiprock, New Mexico to Montezuma Creek, Utah (RM 93.6). Removal efforts upstream of Shiprock Bridge were scaled back to accommodate non-native removal downstream to Montezuma Creek. In addition, at the direction of the San Juan River Recovery Implementation Program's (SJRIP) Biology Committee, trips specific to non-native removal were initiated in 2006 to encompass the Montezuma Creek to Mexican Hat, Utah portion of the river (22 February 2006 Biology Committee Meeting). Two trips were conducted from Montezuma Creek to Mexican Hat, Utah in 2006 by NMFRO and Utah Division of Wildlife Resources – Moab (UDWR).

It was determined at the February 2007 Biology Committee Meeting to increase efforts of nonnative removal from Shiprock to Mexican Hat to reflect similar efforts to intensively sampled reaches upstream. Therefore, we propose to continue our number of sampling trips to include four trips from Shiprock to Mexican Hat in FY 2009. These additional trips will allow removal crews to expand removal into areas of increased importance while maintaining sufficient effort in upstream reaches to maintain current accomplishments.

Description of Study Area

Intensive nonnative fishes removal will occur in the San Juan River, New Mexico-Colorado-Utah, including three distinct reaches of the upper and middle portions of the San Juan River. These sections include PNM Weir (RM 166.6) to Hogback Diversion (RM 159.0); Hogback Diversion to Shiprock Bridge (RM 147.9); and Shiprock Bridge to Montezuma Creek, Utah (RM 93.6). In addition, one sampling trip will continue downstream of Montezuma Creek to Mexican Hat, Utah (RM 52.9).

Objectives

1. Evaluate distribution and abundance patterns of non-native species to determine effects of mechanical removal.
2. Characterize distribution and abundance of endangered fish in the upper and middle reaches of the San Juan River.
3. Relate distribution and abundance patterns of both common and uncommon native fishes to nonnative removal.
4. Establish measurable criteria for evaluating removal effectiveness of nonnative fishes in the system to achieve recovery goals.
5. Continue and expand transplantation of channel catfish to closed impoundments isolated from the San Juan River with the assistance of New Mexico Department of Game and Fish, Navajo Nation Fish and Wildlife Service and the SWTFC.

Methods/Approach

Data Collection:

Mechanical removal will continue during the fall main channel monitoring efforts. During these sampling efforts, all non-native fishes collected will be sacrificed and data recorded for species identification and enumeration, ontogenetic stage (young-of-year, sub-adult, adult) at non-designated miles, and standard and total lengths and weight at designated miles.

Removal efforts from PNM Weir to Hogback and Hogback to Shiprock will be conducted by two electrofishing rafts and one support raft. Captured channel catfish will be measured (nearest 1 mm) for standard and total lengths, weighed (nearest 5 g), and, if not sacrificed for study purposes, transported by hatchery truck to isolated recreational angling impoundments. All other nonnative species sampled during these efforts will be sacrificed and appropriate data recorded for location, length, and mass.

Removal efforts from Shiprock to Mexican Hat will be conducted four times a year. Three of these four trips will be stand alone efforts while the fourth removal trip will be in concert with the Sub-adult/Adult Fish Community Monitoring conducted by FWS-GJ. Sampling for nonnative fishes will be conducted by four raft mounted electrofishing units. Two rafts will begin sampling approximately 1-2 hours after the initial two rafts begin essentially accomplishing two sampling passes per trip. Captured channel catfish will be measured (nearest 1 mm) for standard and total lengths, weighed (nearest 5 g), and, if not sacrificed for study purposes, transported by hatchery truck to isolated recreational angling impoundments. All other nonnative species sampled during these efforts will be sacrificed and appropriate data recorded for location, length, and mass.

In addition to nonnative fishes, all rare fishes seen will be netted. Rare fishes will be measured (nearest 1 mm) for standard and total lengths, weighed (nearest 5 g) and checked for the presence of a Passive Implant Transponder (PIT) tag. If no tag is present

and fish are ≥ 150 mm total length a tag will be implanted. At the time of collection, GPS coordinates will be recorded using a hand held GPS unit.

Data Analysis:

All available capture data will be analyzed independently by section and project (i.e. PNM to Hogback; Hogback to Shiprock; fall monitoring). To determine trends in distribution and abundance, mean CPUE and standard error will be calculated. Species CPUE represents the total number of fish collected divided by the total effort of sampling (hours of electrofishing). Data will be summarized by type of trip, year, section and by individual trips. If CPUE data meet the assumptions of normality and variance, a One Way Analysis of Variance (ANOVA) will be conducted to determine if significant differences exist. Multiple pairwise comparisons using Bonferroni post hoc tests will be used to determine where specific differences exist. All CPUE data that does not meet the assumptions of an ANOVA and transformations are unsuccessful in normalizing the data will first be analyzed using a non-parametric Kruskal-Wallis rank test. If significant differences are observed, among year comparisons of ranked data will be conducted using a Nemenyi post-hoc test (Sokal and Rohlf 1995). Statistical applications not mentioned here may be utilized if deemed appropriate.

Intensive removal trips by Section (FY 2009):

PNM to Hogback-	4 trips
Hogback to Shiprock-	4 trips
Shiprock to Mexican Hat	<u>4 trips</u>
Total # of trips-	12 trips in FY 2009

Products/Schedule

An electronic data file will be provided for inclusion in the centralized database by 31 March 2010. A draft summary report detailing findings will be submitted to the San Juan River Implementation Program, Biology Committee, by 31 March 2010. Revisions will be completed and a final annual report will be submitted by 1 June 2010.

Literature Cited

- Davis, J.E. 2006. *Non-native species monitoring and control in the upper San Juan River, New Mexico: 2005*. Final Report prepared for the San Juan River Recovery Implementation Program. U.S. Fish and Wildlife Service, Albuquerque New Mexico.
- Jackson, J.A. 2006. *Nonnative control in the lower San Juan River: 2005*. Final Report prepared for the San Juan River Recovery Implementation Program. U.S. Fish Wildlife Service, Albuquerque, New Mexico.
- Ryden, D.W. 2006. *Long term monitoring of sub-adult and adult large-bodied fishes in The San Juan River: 2005*. Prepared for the San Juan River Recovery Implementation Program. U.S. Fish and Wildlife Service, Albuquerque, New Mexico.

Sokal, R.R. and F.J. Rohlf. 1995. Biometry: the principles and practice of statistics in biological research. 3rd edition. W.H. Freeman and Company, New York.

Budget FY-2009:

Costs for participation of the U.S. Fish and Wildlife Service, New Mexico Fish and Wildlife Conservation Office in the expanded nonnative removal efforts in FY-2009.

Personnel/Labor Costs (Federal Salary + Benefits)

PNM Weir to Hogback Diversion:

Fish Biologist (GS-7-2) - 20 days @ \$218/day (1 person X 5 days/trip X 4 trips)	\$ 4,360.00
---	-------------

Bio. Science Tech (GS-8) – 20 days @ \$319/day (1 person X 5 days/trip X 4 trips)	\$ 6,380.00
	<u>\$ 10,740.00</u>

Hogback Diversion to Shiprock Bridge:

Principal Biologist (GS-12-3) – 20 days @ \$399/day (1 person X 5 days/trip X 4 trips)	\$ 7,980.00
---	-------------

Fish Biologist (GS-7-2) - 20 days @ \$218/day (1 person X 5 days/trip X 4 trips)	\$ 4,360.00
---	-------------

Bio. Science Tech (GS-8) – 20 days @ \$319/day (1 person X 5 days/trip X 4 trips)	\$ 6,380.00
	<u>\$ 18,720.00</u>

Shiprock Bridge to Mexican Hat, Utah:

Principal Biologist (GS-12-3) – 24 days @ \$399/day (1 person x 12 days/trip x 2 trips)	\$ 9,576.00
--	-------------

Fish Biologist (GS-7-2) – 24 days @ \$218/day (1 person x 12 days/trip x 2 trips)	\$ 5,232.00
--	-------------

Bio. Science Tech (GS-8) – 48 days @ \$319/day (1 person x 12 days/trip x 4 trips)	\$ 15,312.00
--	--------------

Fish Biologist (GS-5-1) – 48 days @ \$170/day (2 people x 12 days/trip x 4 trips)	\$ 8,160.00
---	-------------

Bio. Science Tech (GS-4-1) – 24 days @ \$152/day (2 people x 12 days/trip x 1 trip)	\$ 3,648.00
	<u>\$ 41,928.00</u>

Administrative Support (Federal Salary + Benefits)

Administrative Officer (GS-9-6) – 15 days @ \$268/day	\$ 4,020.00
	<u>\$ 4,020.00</u>

Reporting/Data Management (Federal Salary + Benefits)	
Principal Biologist (GS-12-3) – 60 days @ \$399/day	\$ 23,940.00
Fish Biologist (GS-7-2) – 25 days @ \$218/day	<u>\$ 5,450.00</u>
	\$ 29,390.00
Sub-Total for Personnel /Labor Costs	\$104,798.00
Travel and Per Diem (Based on Published FY-2007 Federal Per Diem Rates)	
Hotel Costs – 65 nights	\$ 4,550.00
(65 nights @ \$70/night – single occupancy = \$4,550)	
Per Diem (Hotel Rate) – 76 days @ \$39/day	\$ 2,964.00
Per Diem (Camp Rate) – 120 days @ \$29/day	<u>\$ 3,480.00</u>
	\$ 10,994.00
Equipment	
PNM Weir to Hogback/Shiprock	\$ 2,880.00
4,800 miles @ \$0.60/mile (based anticipated GSA rates established on 01 September 2008 and includes costs associated with gasoline/diesel fuel and vehicle maintenance)	
(400 miles roundtrip x 8 trips) +(1,600 shuttle miles)	
Shiprock to Mexican Hat, Utah	
8,400miles @ \$0.60/mile (based anticipated GSA rates established on 01 September 2008 and includes costs associated with gasoline/diesel fuel and vehicle maintenance)	\$ 5,040.00
(700 miles roundtrip x 4 trips) x 3 vehicles	
Generator fuel	
20 gallons/trip x 8 trips @ \$3.00/gallon; upper SJR trips and 110 gallons/trip x 2 trips @ \$3.00/gallon; camping trips)	\$ 1,140.00
Equipment Maintenance, Repair, & Replacement (e.g., life jackets, hip boots, generator repair, rubber gloves, dip nets, aluminum welding, raft repair, etc.)	<u>\$ 2,000.00</u>
	\$ 11,060.00
USFWS-NMFRO (Albuquerque) Total	\$126,852.00
USFWS Region 2 Regional Office Administrative Overhead (22.00%)	<u>\$ 27,907.00</u>
USFWS Region 2 Total	\$154,759.00
Funding for participating agencies:	
U.S. Fish and Wildlife Service – Grand Junction, CO	\$ 82,130.00
Utah Department of Wildlife Resources – Moab, UT	\$ 32,741.00
New Mexico Department of Game and Fish- Santa Fe, NM	\$ 52,937.00
Grand Total of FY 2009	
Expanded Nonnative Removal	\$322,567

Outyear funding (based on anticipated 5% Cost of Living Increase)

FY 2009	\$322,567
FY 2010	\$338,695
FY 2011	\$373,412
FY 2012	\$392,083

Costs for participation of the U.S. Fish and Wildlife Service, Colorado River Fishery Project (CRFP) office, Grand Junction, CO in the expanded nonnative removal efforts in FY-2009.

Personnel/Labor Costs (Federal Salary + Benefits)

Principal Biologist (GS-12) – 38 days @ \$449/day (1 person X 5 days/trip X 1 hotel trip) (1 person X 11 days/trip X 3 camping trips)	\$ 17,060.00
Biological Technicians (GS-7) - 49 days @ \$236/day (1 person X 5 days/trip X 1 hotel trip) (1 person X 11 days/trip X 4 camping trips)	\$ 11,565.00
Biological Technicians (GS-5) – 66 days @ \$131/day (3 people x 11 days/trips x 2 trips)	<u>\$ 8,645.00</u>
	\$ 37,270.00

Administrative Support (Federal Salary + Benefits)

Administrative Officer (GS-9) – 12 days @ \$297/day	<u>\$ 3,565.00</u>
	\$ 3,565.00

Reporting/Data Management (Federal Salary + Benefits)

Principal Biologist (GS-12) – 40 days @ \$449/day	<u>\$ 17,960.00</u>
	\$ 17,960.00

Travel and Per Diem (Based on Published FY-2007 Federal Per Diem Rates)

Hotel Costs – 18 nights (18 nights @ \$70/night – single occupancy = \$1,260)	\$ 1,260.00
Per Diem (Hotel Rate) – 16 days @ \$39/day	\$ 625.00
Per Diem (Camp Rate) – 80 days @ \$26/day	<u>\$ 2,080.00</u>
	\$ 3,965.00

Equipment

Vehicle Maintenance & Gasoline (@ \$0.60/mile) (600 miles round trip from Grand Junction, CO to Farmington, NM + 200 miles of shuttling per trip X 5 trips)	\$ 2,400.00
Generator Gasoline (110 gallons/trip x 2 trips @ \$3.00/gallon)	\$ 660.00
Equipment Maintenance, Repair, & Replacement (e.g., spark plugs and oil for electrofishing generators, generator repair, life jackets, hip boots, rubber gloves, dip nets, aluminum welding, raft repair, etc.)	<u>\$ 1,500.00</u>
	\$ 4,560.00

USFWS-CRFP (Grand Junction) Total \$ 67,320.00

USFWS Region 6 Regional Office Administrative Overhead (22.00%) \$ 14,810.00

USFWS Region 6 Total \$ 82,130.00

Costs for participation of Utah Division of Wildlife Resources, Moab, UT in the Upper and Middle San Juan nonnative removal FY-2009.

Personnel/Labor Costs (Salary + Benefits)

Principal Biologist– 45 days @ \$265/day (1 person X 5 days/trip X 2 hotel trips (1 person X 10 days/trip X 3 camping trips and 5 office days for trip prep, gear cleaning, etc.)	\$ 11,925.00
Biological Technicians - 45 days @ \$185/day (1 person X 5 days/trip X 2 hotel trips (1 person X 10 days/trip X 2 camping trips and 5 office days for trip prep, gear cleaning, etc.)	\$ 8,325.00
Project Leader – 2 days @ \$290/day	<u>\$ 580.00</u>
Subtotal	\$ 20,830.00

Travel and Per Diem (Based on Published FY-2007 Federal Per Diem Rates)

Hotel Costs – 18 nights @ \$70/night	\$ 1,260.00
Per Diem (Hotel Rate) – 22 days @ \$43/day	\$ 946.00
Per Diem (Camp Rate) – 60 days @ \$20/day	\$ 1,200.00
Vehicle Rent (1 truck @\$250 month for 2 months)	\$ 500.00
Vehicle Mileage (2,088 miles @ \$0.49 per mile)	\$ 1,023.00
Subtotal	\$ 4,929.00

Equipment Maintenance, Repair and Replacement

Generator Gasoline (30 gallons/trip x 5 trips @ \$3.50/gallon)	\$ 525.00
Maintenance (spark plugs, air filters, oil, generator repair)	\$ 400.00
Data Collection Supplies (spring scales, pencils, measuring boards)	\$ 200.00
Camping Gear (tents, sleeping pads, dry bags)	\$ 200.00
Rafting Supplies (oars, raft repair, life jackets, straps, pumps)	\$ 200.00
Subtotal	\$ 1,525.00

UDWR – Moab Total \$ **27,284.00**

UDWR - Moab Administrative Overhead (20 %) \$ **5,457.00**

UDWR – Moab Grand Total \$ **32,741.00**

Costs for participation of the New Mexico Department of Game and Fish, Santa Fe, N.M. in the expanded nonnative removal FY-2009.

Personnel/Labor Costs (State Salary + Benefits)

Biologists - 20 @ \$350/day (1 person x 5 days/trips x 4 trips)	\$ 7,000
Biologists– 88 days @ \$350/day (2 people X 11 days/trip X 4 trips; camping trips)	\$ 30,800
	<u>\$ 37,800</u>
Travel and Per Diem (Based on Published FY-2007 State Per Diem Rates)	
Per Diem – 104 days @ \$85/day	\$ 8,840
	<u>\$ 8,840</u>
Equipment	
Vehicle Maintenance & Gasoline (@ \$0.32/mile) (660 miles round trip from Albuquerque, NM to Blanding, UT x 4 trips and 500 miles round trip from Albuquerque, NM to Shiprock, NM x 4 trips)	\$ 1,485
	<u>\$ 1,485</u>
NMDGF – Santa Fe	Total
	\$ 48,125
Administrative Overhead (10%)	\$ 4,812
NMDGF – Santa Fe – Total Budget	\$ 52,937

**Nonnative Species Control in the Lower San Juan River
Fiscal Year 2009 Project Proposal**

Principal Investigator: Darek S. Elverud
Utah Division of Wildlife Resources, Moab Field Station
1165 S. Hwy 191- Suite 4, Moab, Utah 84532
(435) 259-3782
darekelverud@utah.gov

Background:

The lower San Juan River is particularly important in the recovery of the Colorado pikeminnow (*Ptychocheilus lucius*) and razorback sucker (*Xyrauchen texanus*) since it contains typical nursery habitat similar to what is present on the Green and Colorado rivers. Within the past six years, collections of endangered fish have been increasing in this section of river. The largest collection of razorback sucker larvae in 2002 was from Reach 2 (RM 21.2; Brandenburg et al. 2003) and the largest single collection of razorback sucker larvae in 2003 came from a backwater in Reach 1 at RM 8.1 (Brandenburg et al. 2004). Additionally, adult razorback sucker were found congregating around Slickhorn Rapid (RM 17.7) in the spring of 2002, apparently using this area for spawning (Jackson 2003). In spring of 2006, another congregation of adult razorback suckers and possible spawning area was located at river mile 23.4. Collections of adult Colorado pikeminnow in the San Juan River have been extremely rare. No wild adults have been collected since 2000 (Ryden 2003). From 2002 to 2004, Colorado pikeminnow adults and subadults, presumably from the 1996-1997 stocking efforts, have been found using the lower canyon (Reaches 1 and 2) of the San Juan River in the spring and summer (Jackson 2005). In 2003 and 2004, young-of-year Colorado pikeminnow stocked in the fall of the previous year near Farmington, NM, were also found using the lower portions of the San Juan River (Golden et al. 2005). One of the most encouraging findings from 2004 was the collection of two wild spawned Colorado pikeminnow larvae at RM 46.3 and 18.1 (Brandenburg et al. 2005).

Nonnative fish species remain prevalent in the lower San Juan River. Channel catfish (*Ictalurus punctatus*) and common carp (*Cyprinus carpio*) are typically the most abundant fish species collected during fall monitoring in Reaches 1 and 2 (Ryden 2003). Native and endangered fish are threatened by predation from adult channel catfish (Marsh and Brooks 1989, Brooks et al. 2000), and may compete for food and space with juvenile channel catfish. Additionally, Colorado pikeminnow have been found with channel catfish lodged in their throats in the San Juan (Ryden and Smith 2002) and Green (McAda 1983, personal observation) rivers. Common carp tend to feed on larval fish and eggs (Cooper 1987). In the spring and summer of 2004, recently stocked razorback sucker and Colorado pikeminnow were found in the stomachs of two different channel catfish (Jackson, 2005).

Since 1995, many nonnative species including striped bass (*Morone saxatilis*) and walleye (*Stizostedion vitreum*) have been able to move into the San Juan River from Lake Powell. From 1988 to 1995, a waterfall at approximately RM 0 acted as a barrier between the San Juan River and Lake Powell, preventing species from moving upstream. During 1995, rising lake levels inundated the waterfall. When lake levels receded in the winter of 1996, the waterfall did not reappear. Striped bass, walleye and threadfin shad (*Dorosoma petenense*), not previously documented in the San Juan River before waterfall inundation, were collected during large bodied fish sampling (Ryden 2001). Since then, striped bass and walleye have been collected periodically until 2000 when large numbers were collected near Farmington, NM (approximately 166 river miles upstream of Lake Powell). Many native suckers were found inside the stomachs of these striped bass (unpublished data from San Juan River database). The San Juan River Recovery Implementation Program (SJ RIP) determined in 2001 that control of striped bass and other nonnative species in the lower river was warranted. Utah Division of Wildlife Resources began nonnative fish control with the goal of removing striped bass and other nonnative species in the lower San Juan River, while documenting river and lake

conditions that may correlate to striped bass movement out of Lake Powell. It was anticipated that these correlations would provide information for determining the most effective time to remove striped bass. During 2002, Lake Powell water temperature was positively correlated with the highest catch of striped bass in June, in the lower San Juan River (Jackson, 2003). A new waterfall at RM -0.5 has prevented striped bass and other fish from moving from Lake Powell since 2003. No striped bass or walleye were observed in the lower San Juan River from 2003 to 2007. In 2006, two adult gizzard shad were captured below the waterfall indicating another possible nonnative fish of concern. In 2007, seine sampling below the waterfall collected hundreds of young-of-the-year gizzard shad below the waterfall. Additionally in 2007, two adult gizzard shad, two juvenile striped bass and two adult walleye were collected below the waterfall. Colorado pikeminnow and razorback suckers have also been collected during sampling efforts below the water.

Over 56,000 channel catfish and approximately 2,900 common carp were mechanically removed from the lower San Juan River from 2002 to 2007. A decrease in mean total length (TL) of channel catfish was observed between 2002 and 2004, indicating that removal efforts are causing a shift in the population size structure to smaller individuals. Additionally, shifts in sized structure of channel catfish have been reported further upstream (Davis 2005) and on a river-wide scale (Ryden 2005). Catch rate of adult channel catfish also decreased from 4.9 adult catfish per electrofishing hour in 2002 to 2.0 adult catfish per electrofishing hour in 2006 in the lower San Juan. Furthermore, similar shifts in yield and population structure have been observed in sport and commercial fisheries as the rate of exploitation increased (Bennet 1971; McHugh 1984, Pitlo 1997). Continued removal of all size classes of channel catfish in the San Juan River may eventually lead to decreased fecundity and a reduction of the overall population, therefore lessening the impact that these fish have on the native and endangered fish community.

A significant decline in catch rates of common carp was observed from 2002 to 2007. Between 2002 and 2007, catch rate of common carp decreased from over 5 fish per electrofishing hour to < 0.2 fish per electrofishing hour. However, it is unclear if this decline was directly related to removal efforts, the presence of the waterfall, or the low water conditions that have been present over the period of this project. It is probable that a combination of these factors is causative to some extent. The continuation of removal efforts for channel catfish and common carp will aid in the illumination of contributory factors and the evaluation of the success of this project and similar nonnative control efforts.

Over the course of this project, important information has been obtained on the progress of the endangered fish community as well. We have observed the apparent spawning aggregation of razorback sucker in spring 2002 at Slickhorn Rapid and collected some of the first wild spawned juvenile razorback sucker in 2003 and 2004. Since 2002, we have documented the distribution and abundance of Colorado pikeminnow in the lower San Juan River stocked from 2002 to 2006. Preliminary population estimates for juvenile Colorado pikeminnow residing in the lower San Juan River were generated in 2004, 2005, 2006 and 2007 from recapture data. In 2004, we documented the first cases of channel catfish predation on stocked juvenile razorback sucker and Colorado pikeminnow in the San Juan River. Sampling also documented the presence of Colorado pikeminnow and razorback sucker below the waterfall.

This work plan proposes the continuation of nonnative control, sub-element 4.1 of the Long Range Plan, in the lower San Juan River from Mexican Hat to Clay Hills, and sampling just below the waterfall at Piute Farms. Since striped bass and other fish are unable to navigate the waterfall, sampling below the waterfall will determine their presence or absence. If they are there, we can continue to document the riverine and lacustrine conditions related to their movement. This study will serve to determine the most effective time for removal actions, so that more intensive and specific removal efforts may be employed in the future when Lake Powell is once again influencing the lower San Juan River. The presence of the waterfall at Piute Farms may provide a rare opportunity to concentrate on removal of other nonnative fish while influx from the lake is eliminated. Continuing removal in the lower river above the waterfall will aid in removal efforts being conducted further upstream, and suppress predation and competition impacts on the endangered and native fish community by nonnative fish in the lower San Juan River.

In addition, we propose to continue to document the progress of Colorado pikeminnow and razorback sucker in the lower San Juan River. Recapture data for juvenile Colorado pikeminnow collected during nonnative removal will serve in determining population size, growth and movement of these fish in the lower San Juan River. Furthermore, conducting work below the waterfall will provide information on endangered fish that may be present and unable to move upstream.

Description of Study Area:

The study area for this project includes the San Juan River from Mexican Hat (RM 53) to Clay Hills (RM 2.9), Utah. Additionally, sampling will be conducted just below the waterfall at RM-0.5 and from the waterfall downstream to Lake Powell (approximately 15 to 20 river miles and within critical habitat). The river from Mexican Hat to RM 16 is part of Geomorphic Reach 2 and is primarily bedrock confined and dominated by riffle-type habitat. River mile 16 down to Clay Hills contains Geomorphic Reach 1 where the river is canyon bound with an active alluvial bed. Habitats within this section are heavily influenced by the shifting thalweg, changing river flow, and reservoir elevations. This section of river has been identified as important nursery habitat for native and endangered fish species.

Objectives

- 1.) Continue mechanical removal of large-bodied nonnative species in the lower portion of the San Juan River from Mexican Hat to Clay Hills.
- 2.) Generate a population estimate of channel catfish by mark-recapture data from Mexican Hat to Clay Hills.
- 3.) Characterize distribution and abundance of endangered fish in the lower San Juan River.
- 4.) Generate a population estimate of juvenile Colorado pikeminnow (>150 mm) by mark-recapture data from Mexican Hat to Clay Hills.
- 5.) Characterize abundance of endangered fish in the San Juan River just below the waterfall.
- 6.) Characterize abundance of reservoir immigrants (striped bass and walleye) moving out of Lake Powell into the San Juan River upstream to the new waterfall.
- 7.) Relate striped bass movement from Lake Powell into the San Juan River to lake and river conditions (including temperature, flows and turbidity).

Methods/Approach:

Mechanical removal of nonnative species will be conducted from Mexican Hat to Clay Hills, Utah. Sampling effort will be conducted via two raft mounted electrofishing boats. The entire study area will be electrofished in a downstream fashion with one boat on each shoreline. Each boat will have one netter and one rower. A third boat will follow behind to pick up nonnative fish missed by the electrofishing boats. These fish will not be included in catch rate calculations, so that comparisons can be made between trips and years. Nine five-day trips with 6 people are anticipated, and timing of sampling will be dependent on catch rate from past data. Bimonthly trips will be conducted, which will likely translate into every other week sampling from March through August. Data from the adult fall monitoring conducted by U.S. Fish and Wildlife Service- Grand

Junction in October will be incorporated into data analysis. In an average water year, this schedule would allow for sampling a variety of habitat conditions, including variable flows, temperatures, and turbidity. In addition, a variety of sampling methods will be used below the waterfall, including hoop and trammel netting, seining, hook and line, and electrofishing, if possible. Five sampling events will take place directly below the waterfall, most likely between April and August.

All nonnative fish collected will be identified, enumerated, measured to the nearest mm for total and standard length, weighed to the nearest gram, and removed from the river. Gender and reproductive status of lacustrine species will be determined and approximate location of capture by river mile recorded. Stomach contents of lacustrine species will be examined. Contents needing microscopic identification will be preserved. Any threatened and/or endangered fish encountered will be collected, identified, enumerated, measured to the nearest mm for total and standard length, weighed to the nearest gram, and scanned for a PIT tag. If a PIT tag is not present, one will be inserted. General condition of the fish will be recorded in addition to any parasites or abnormalities. All threatened and endangered fish collected will be returned to the river at the location in which they were caught. River mile and GPS coordinates will be recorded at the location in which threatened and endangered fish are collected. Catch rates for all fish will be calculated as number of fish caught per hour. Other native fish will not be netted.

Channel catfish collected during the first trip of the year will receive a floy tag and be returned to the river. Channel catfish collected on subsequent trips will be removed from the river. A Lincoln-Peterson population estimate will be generated for channel catfish captured during the first pass and recaptured in the second pass. Captures of channel catfish during subsequent trips will allow us to monitor ratios of marked to unmarked fish and use these ratios to calculate a rough population estimate thereafter. Ratios of marked fish to unmarked fish will help determine if assumptions of a closed population are being met.

Population estimates will be generated for juvenile Colorado pikeminnow (>150 mm) in the lower San Juan River using closed population models within program CAPTURE. Program CAPTURE will be used to determine confidence intervals around the estimate, the coefficient of variation, and the probability of capture. Population estimates between two passes will be calculated using the Lincoln-Peterson model. Conducting several trips in the lower San Juan River will allow for choosing the "mark" pass and the number of "recapture" passes. Use of different mark and recapture passes will allow for testing of the reality of the results generated. Furthermore, using several combinations of trips will allow for lessening the likelihood of violating assumptions of the models used.

General water quality parameters will be recorded including temperature, conductivity, salinity, and dissolved oxygen. Daily river discharge, temperature and turbidity will be compared to catch rates for striped bass to determine the relationship between river conditions and movement of these fish upstream.

Costs for other cooperating agencies that may provide personnel and equipment as needed are included in this budget.

Products/Schedule:

A draft report for the Nonnative Species Control in the Lower San Juan River activities will be prepared and distributed to the San Juan River Biology Committee for review by 31 March 2010. Historical information on nonnative fish species use of the lower San Juan River will be included; to the extent it is available. Upon receipt of written comments, that report will be finalized and forwarded to members of the San Juan River Biology Committee 1 June 2010. Electronic copies of the field and collection data will be transferred to the San Juan River database manager following the successful protocol previously employed.

Literature Cited:

- Bennet, G.W., 1971. Management of lakes and ponds, 2nd edition. Van Nostrand Rienhold, New York.
- Brandenburg, H.W., M.A. Farrington, S.J. Gottlieb. 2003. Razorback sucker larval fish survey in the San Juan River during 2002. Draft report to the San Juan River Basin Recovery Implementation Program. U.S. Fish and Wildlife Service, Albuquerque NM.
- Brandenburg, H.W., M.A. Farrington, S.J. Gottlieb. 2004. Razorback sucker larval fish survey in the San Juan River during 2003. Draft report to the San Juan River Basin Recovery Implementation Program. U.S. Fish and Wildlife Service, Albuquerque NM.
- Brandenburg, W.H., M.A. Farrington, S.J. Gottlieb. 2005. San Juan River 2004 Colorado pikeminnow and razorback sucker larval surveys. Draft Report. San Juan River Basin Recovery Implementation Program, USFWS, Albuquerque, NM
- Brooks, J.E., M.J. Buntjer, J.R. Smith. 2000. Non-native species interactions: Management implications to aid in the recovery of the Colorado pikeminnow (*Ptychocheilus lucius*) and razorback sucker (*Xyruachen texanus*) in the San Juan River, CO-NM-UT. U. S. Fish and Wildlife Service, Albuquerque, NM..
- Cooper, E.L. ed. 1987. Carp in North America. American Fisheries Society. Maryland, Bethesda
- Davis, J.E. 2005. Non-native species monitoring and control in the Upper San Juan River, New Mexico 2004 (Draft Report). Progress Report for the San Juan River Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, NM.
- Golden, M.E., P.B. Holden, S.K. Dahle. 2005. Retention, Growth, and Habitat Use of Stocked Colorado Pikeminnow in the San Juan River: 2002-2003 Draft Annual Report for the San Juan River Basin Recovery Implementation Program. U. S. Fish and Wildlife Service, Albuquerque NM.
- Jackson, J.A. 2003. Nonnative control in the lower San Juan River, 2002. Interim Progress Report for the San Juan River Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, NM.
- Jackson, J.A. 2005. Nonnative control in the lower San Juan River, 2004. Draft Progress Report for the San Juan River Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, NM.
- Marsh, P.C. and J.E. Brooks. 1989. Predation by ictalurid catfishes as a deterrent to re-establishment of hatchery-reared razorback suckers. Southwestern Naturalist 34(2):188-195.
- McAda, C.W. 1983. Colorado squawfish, *Ptychocheilus lucius* (Cyprinidae), with a channel catfish, *Ictalurus punctatus* (Ictaluridae), lodged in its throat. Southwestern Naturalist 28(1):119-120.
- McHugh, J.L. 1984. Industrial fisheries, pages 68-80 in R.T. Barber, C.N.K. Mooers, M.J. Bowman, and B. Zeitschel, editors. Lecture notes on coastal and estuarine studies. Springer-Verlag, New York.
- Pitlo, J.Jr. 1997. Response of upper Mississippi River channel catfish populations to changes in commercial harvest regulations. North American Journal of Fisheries Management 17: 848-859.
- Ryden, D. W. 2001. Long term monitoring of sub-adult and adult large-bodied fishes in the San Juan River, 2000. Interim Progress Report. U.S. Fish and Wildlife Service. Grand Junction, CO. 61 pp.

- Ryden, D. W. 2003. Long term monitoring of sub-adult and adult large-bodied fishes in the San Juan River: 1999-2001 Integration Report. U.S. Fish and Wildlife Service. Grand Junction, CO. 68 pp.
- Ryden, D. W. 2005. Long term monitoring of sub-adult and adult large-bodied fishes in the San Juan River, 2004. Interim Progress Report. U.S. Fish and Wildlife Service. Grand Junction, CO.
- Ryden, D. W. and J.R. Smith. 2002. Colorado pikeminnow with a channel catfish lodged in its throat in the San Juan River, Utah. *Southwestern Naturalist* 47:92-94.

Budget FY-2009:**Personnel / Labor Costs (State Salary + Benefits)**

Lead Biologist @ \$265/day for 152 man days total 82 field days (planning and organization, logistics, electrofishing) 70 other days (coordination, data entry, data analysis, administrative support, meeting attendance	\$40,280
Technicians @ \$185/day for 280 man days total preparation for field trips, equipment and gear maintenance, electrofishing	\$51,800
Project Leader @ \$290/day for 20 man days total office and administrative support, review of reports, logistical support, meeting attendance, electrofishing	\$ 5,800
Personnel / Labor Costs Subtotal	\$97,880

Travel and Per Diem

Mileage: Mexican Hat to Clay Hills trips-340 mi @ \$.49 per mi for 9 trips	\$ 3,165
shuttle of three vehicles @ \$350 per trip (9 trips)	\$ 3,150
Piute Farms- 336 mi @ \$.49 per mi for 5 trips	\$ 823
Vehicle rent (1 x 6 x \$250/month)	\$ 1,500
Per Diem:	
Camping rate-	
Mexican Hat to Clay Hills- 6 people @ \$20 per day for 45 days	\$ 5,400
Piute Farms- 2 people @ \$20 per day for 20 days	\$ 400
Hotel rate-	
Out-of-state per diem @ \$43 x 10 days	\$ 430
Hotel Costs	
4 meetings per year (\$70.00/night for 8 nights)	\$ 560
Travel and Per Diem Subtotal	\$15,428

Equipment Maintenance, Repair and Replacement

	Unit Price	\$ Total
Fuel for generators (30 gal/trip x 9 trips = 270 gallons)	\$3.50/gallon	\$ 945
Wiring replacement for electrofishing systems		\$ 100
Repair of electrofishing frame (aluminum welding)		\$ 300
Replacement of electrofishing equipment		
Dip nets	\$200	\$ 200
Foot switch	\$200	\$ 200

Life jackets	\$100	\$ 300
First aid supplies	\$ 80	\$ 80
Waders	\$100	\$ 200
Data collection supplies		
Paper, pencils, binders, staples, etc.		\$ 150
Measuring boards		\$ 100
Spring scales	\$ 40	\$ 200
Plungers, needles, alcohol for PIT tags		\$ 50
Floy tags for marking catfish		\$ 200
Tools		\$ 100
Repair of GPS units		\$ 50
Satellite phone charges (\$30/month for 6 months)		\$ 180
Repair of GPP		\$ 500
Repair of generators		\$ 500
Repair of trailers (bearings, axle, tires)		\$ 500
Repair and replacement of rafting supplies		
Oarlocks	\$ 30	\$ 60
Oars	\$200	\$ 400
River straps	\$ 5	\$ 125
Pumps	\$150	\$ 150
Raft repair (valves, d-rings, glue, patches)		\$ 100
Carabiners	\$ 10	\$ 100
Throw bags	\$ 50	\$ 100
Maintenance of generators (oil, sparks plugs, batteries)		\$ 100
Camping Equipment		
Tables	\$ 40	\$ 40
Tents	\$200	\$ 200
Drybags	\$ 50	\$ 200
Cookware		\$ 50
Chairs	\$ 20	\$ 20
Batteries		\$ 100
Toilet supplies		\$ 100
Charcoal	\$ 10	\$ 50
Cleaners		\$ 50
Food storage boxes		\$ 40
Propane		\$ 100
Groover disposal		\$ 50
Equipment Maintenance, Repair, and Replacement Subtotal		\$ 6,990
Subtotal of labor, travel, equipment, etc		\$ 120,298
Administrative Overhead (20%)		
20% of personnel cost for Salt Lake Office administration indirect cost, building operation costs for Moab Field Station (electricity, phone and computer lines, rent, etc.)		\$ 24,060
UDWR TOTAL		\$ 144,358

Funding for Participating Agencies

**Nonnative Species Control in the Lower San Juan River
Fiscal Year 2009 Project Proposal**

Principal Investigators: Darek Elverud
Utah Division of Wildlife Resources, Moab Field Station
1165 S. Hwy 191 – Suite 4, Moab, Utah 84532
(435) 259-3782
darekelverud@utah.gov

Under the heading “Funding for participation of other agencies.” Costs for participation of the New Mexico Game and Fish in FY-2009.

Personnel/Labor Costs (Salary, Benefits, Admin)

Fishery Biologist – 12 days @ \$350/day (1 person x 6 days per trip x 2 trips)	\$ 4,210
---	----------

Travel and Per Diem

(\$85 per day per person – 1 person - 6 days per trip x 2 trips)	\$ 1,020
--	----------

Equipment

Vehicle & Gasoline (\$0.35/mile) (700 miles round trip x 2 trips)	\$ 490
--	--------

Total	\$ 5,710
--------------	-----------------

Outyear funding including 5% increase:

Fiscal Year 2010	\$ 5,996
Fiscal Year 2011	\$ 6,295
Fiscal Year 2012	\$ 6,610

**Non-native Species Control in the *Lower San Juan River*
Fiscal Year 2009 Project Proposal**

Principal Investigator: Darek Elverud
Utah Division of Wildlife Resources, Moab Field Station
1165 S. Hwy 191- Suite 4, Moab, Utah 84532
(435) 259-3782
darekelverud@utah.gov

Budget FY-2009:

Under the heading “Funding for participating agencies.” Costs for participation of the U.S. Fish and Wildlife Service, Colorado River Fishery Project (CRFP) office, Grand Junction, CO in FY-2009.

Personnel/Labor Costs (Federal Salary + Benefits)	
Principal Biologist (GS-12) – 10 days @ \$449/day (1 person X 5 days/trip X 2 trips)	\$ 4,490.00
Biological Technicians (GS-7) - 20 days @ \$236/day (2 people X 5 days/trip X 2 trip s)	\$ 4,720.00
	\$ 9,210.00
Administrative Support (Federal Salary + Benefits)	
Administrative Officer (GS-9) – 4 days @ \$297/day	\$ 1,190.00
Travel and Per Diem (Based on Published FY-2007 Federal Per Diem Rates)	
Hotel Costs – 6 nights (6 nights @ \$70/night – single occupancy = \$420)	\$ 420.00
Per Diem (Hotel Rate) - 6 days @ \$39/day	\$ 235.00
Per Diem (Camping Rate) 30 days @ \$26/day	<u>\$ 780.00</u>
	\$ 1,435.00
Equipment	
Vehicle Maintenance & Gasoline (@ \$0.60/mile) (700 miles round trip from Grand Junction, CO to Clay Hills, UT X 2 trips)	\$ 850.00
Equipment Maintenance, Repair, & Replacement (e.g., spark plugs and oil for electrofishing generator, generator repair, life jackets, hip boots, rubber gloves, dip nets, aluminum welding, raft repair, etc.)	\$ 1,500.00
	\$ 2,350.00
USFWS-CRFP (Grand Junction) Total	\$ 14,185.00
USFWS Region 6 Regional Office Administrative Overhead (22.00%)	<u>\$ 3,120.00</u>
USFWS Region 6 Total	\$ 17,305.00
<u>Estimated Outyear Funding</u>	
Fiscal Year	Anticipated Outyear Cost
FY-2010	\$ 18,170
FY-2011	\$ 19,075
FY-2012	\$ 20,030

**Nonnative Species Control in the Lower San Juan River
Fiscal Year 2009 Project Proposal**

Principal Investigators: Darek Elverud
 Utah Division of Wildlife Resources, Moab Field Station
 1165 S. Hwy 191 – Suite 4, Moab, Utah 84532
 (435) 259-3782
darekelverud@utah.gov

Under the heading “Funding for participation of other agencies.” Costs for participation of the U.S. Fish and Wildlife Service, New Mexico Fish and Wildlife Conservation Office Albuquerque, NM in FY-2009.

Personnel/Labor Costs (Federal Salary + Benefits)	
Fish Biologist (GS-7-2) – 14 days @ \$218/day (1 person x 7 days x 2 trips)	\$ 3,052
Biological Technician (GS-8) – 14 days @ \$ 319/day (1 person x 7 days x 2 trips)	\$ 4,466
Administrative Officer (GS-9-6) – 2 days @ \$268/day	<u>\$ 536</u>
	<u>\$ 8,054</u>
Travel and Per Diem (Based on Published FY-2008 Federal Per Diem Rates)	
Hotel Costs – 4 nights (1 night x 2 rooms x 2 trips @ \$70/night; Bluff, UT)	\$ 280
Camping Costs – 20 nights @ \$29/night _____ (5 nights x 2 people x 2 trips)	<u>\$ 580</u>
	\$ 860
Equipment	
Vehicle Maintenance & Gasoline (@ \$0.60/mile) (660 miles round trip from Albuquerque, NM to Blanding, UT x 2 trips)	\$ 792
Equipment Maintenance, Repair, & Replacement (e.g., life jackets, hip boots, generator repair, rubber gloves, dip nets, aluminum welding, raft repair, etc.)	<u>\$ 1,500</u>
	<u>\$ 2,292</u>
USFWS-NMFRO (Albuquerque) Total	\$ 11,206
USFWS Region 2 Regional Office Administrative Overhead (22.00%)	<u>\$ 2,465</u>
USFWS Region 2 Total	\$ 13,671
Outyear funding including 5% COLA:	
Fiscal Year 2010	\$ 14,355
Fiscal Year 2011	\$ 15,073
Fiscal Year 2012	\$ 15,826

Sub-Adult & Adult Large-Bodied Fish Community Monitoring
(a.k.a. Adult/Juvenile Fish Community Monitoring)
Fiscal Year 2009 Project Proposal
19 May 2008

Principal Investigators: Dale Ryden and Chuck McAda
U. S. Fish and Wildlife Service, Colorado River Fishery Project
764 Horizon Drive, Building B
Grand Junction, Colorado 81506-3946
(970) 245-9319
dale_ryden@fws.gov and chuck_mcada@fws.gov

Background:

Studies performed before 1991 documented a native San Juan River fish fauna of eight species, including Colorado pikeminnow (previously known as Colorado squawfish), razorback sucker, and roundtail chub and provided baseline information on distribution and abundance of native and introduced fish species in the San Juan River. These studies indicated that at least one of the two endangered fish species (i.e., Colorado pikeminnow) was still a viable member of the San Juan River fish community.

Between 1991 and 1998, the Main Channel Fish Community Monitoring study (called “Adult Monitoring” for short), greatly refined our understanding of the San Juan River fish community. The main sampling technique employed during the 1991-1997 Adult Monitoring study was raft-borne electrofishing, although radio telemetry was also heavily employed. Data collected during the 1991-1997 Adult Monitoring study provided information on specific habitat usage by rare fish species. In addition, data gathered during the 1991-1997 Adult Monitoring study aided in the selection of specific sites for detailed hydrologic measurements and larval drift sampling. Integration of 1991-1997 Adult Monitoring data along with data from Colorado pikeminnow macrohabitat studies, razorback sucker experimental stocking studies, tributary and secondary channel studies, fish health studies, contaminants studies, habitat mapping studies, and non-native species interaction studies, helped provide a logical framework upon which to make flow recommendations for the reoperation of Navajo Reservoir that would benefit the San Juan River’s endangered fishes (as well as other members of the native fish community).

The Sub-Adult & Adult Large-Bodied Fish Community Monitoring study (also referred to as Adult Monitoring), which began in 1999, is a direct offshoot of the 1991-1997 Adult Monitoring study. This study is one of a suite of long-term monitoring efforts detailed in the San Juan River Monitoring Plan and Protocols (Propst et al. 2000) that are designed to help evaluate progress under the San Juan River Recovery Implementation Program (SJRIP) and the SJRIP’s Long Range Plan. The current Adult Monitoring study incorporates essentially the same monitoring protocols as did its 1991-1997 precursor study (e.g., sampling via raft-borne electrofishing). This allows for data collected during the current Adult Monitoring study to be validly combined with and compared to the older 1991-1997 Adult Monitoring data. The combination of these two data sets provides statistically-powerful, long-term trend data through which the SJRIP’s Biology Committee can view changes in the San Juan River’s large-bodied fish community over time. This long-term trend data allows the SJRIP Biology Committee to evaluate whether various management actions being implemented are having the desired effects on the San Juan River fish community. In addition, Adult Monitoring has proven to be a highly effective tool for monitoring populations of both stocked razorback sucker and Colorado pikeminnow.

Relationship to the Recovery Program:

At present, both the Long Range Plan and the long-term Monitoring Plan and Protocols are undergoing revision. However, the Adult Monitoring study provides data for or makes possible (at least in part) the following actions under element numbers 1-5 of the latest draft of the Long Range Plan (dated 18 April 2008): 1.1.1.1, 1.1.1.3, 1.1.4.3, 1.2.1.1, 1.2.3.1, 2.2.5.1, 2.2.5.2, 3.2.1.2, 4.1.1.7, 4.1.1.8, 5.1.1.2, 5.1.2.3, 5.1.2.4, 5.1.2.5, and 5.2.3.1. The monitoring protocols discussed in the Methods section of this workplan reflect those that are currently included in the latest draft of the revised long-term Monitoring Plan and Protocols.

Description of Study Area:

The study area for Adult Monitoring extends from river mile (RM) 180.0 (just downstream of the Animas River confluence) in Farmington, New Mexico, downstream to RM 2.9 (Clay Hills Landing) just upstream of Lake Powell in Utah.

Objectives:

- 1.) Monitor the San Juan River's fish community, specifically the large-bodied fish species, to identify shifts in fish community structure, species relative abundance and distribution, and length/weight frequencies that are occurring over time. Determine whether these shifts in fish community parameters correspond to management actions that are being implemented by the SJRIP. These include (but may not be limited to) the following:
 - a) Reoperation of water releases from Navajo Reservoir
 - b) Mechanical removal of nonnative fishes
 - c) Modification or removal of instream water diversion structures
 - d) Augmentation efforts for both federally-listed endangered fish species -- Colorado pikeminnow and razorback sucker
- 2.) Monitor population trends (e.g., distribution and abundance) of the rare San Juan River fish species -- Colorado pikeminnow, razorback sucker, and roundtail chub (both wild and stocked fish).
- 3.) Remove nonnative fish species which prey upon and may potentially compete with native fish species in the San Juan River.

Methods:

Objectives 1-3: One Adult Monitoring trip will take place in the fall of 2009. This trip will sample the entire study area, from near the Animas River confluence in New Mexico (RM 180.0) to Clay Hills Landing in Utah (RM 2.9). Raft-borne electrofishing will be the primary sampling technique. Sampling will begin in the second to third week of September and will be concluded by the second to third week of October.

Two oar-powered rafts, with one netter each, will electrofish in a continuous downstream fashion, with one raft on each shoreline. Netters will net all stunned fish that can possibly be collected, regardless of species or body size. Trailing or "chase" rafts will not be used to collect fish. No outboard motors will be used. Sampling crews will consist of approximately 8-10 people (4 for electrofishing, 2-3 for baggage rafts, and 2-3 for other research elements that are being done simultaneously with our sampling). Electrofishing will sample two out of every three miles (approximately 120 total sampled miles). All fish collected will be enumerated by species and life stage at the end of every sampled mile. Every fourth sampled mile (known as a "designated mile" or DM), all fish collected will be weighed and measured. All native fish collected will be returned alive to the river. All nonnative fish collected will be removed from the river. All nonnative predatory fishes (e.g. - walleye, striped bass, largemouth bass, smallmouth bass) collected will be weighed and measured, and may have stomach samples taken, before being removed from the river. Tag numbers, total length, and weight will be recorded on all recaptured, FLOY-tagged fish (both native and nonnative), as well as any rare fish collected. Colorado pikeminnow, razorback sucker, and roundtail chub greater than 200 mm TL will be implanted with 134 kHz PIT (Passive Integrated Transponder) tags. Notes will be kept on any parasites and/or abnormalities observed on collected fishes.

Electrofishing will follow the methods set forth above and in the SJRIP's long-term monitoring plan (as detailed in Propst et al. 2000). Alternate sampling techniques (e.g., seining, trammel netting, backpack electrofishing, etc.) may be employed where suitable habitat is available or if low-water conditions preclude the use of raft-mounted electrofishing, at the principal investigator's discretion.

The U.S. Fish and Wildlife Service will assume the lead responsibility for Adult Monitoring trips and other

cooperating agencies will provide personnel and equipment as needed. Costs for cooperating agencies are included in this budget.

Products:

An interim progress report for Adult Monitoring data collected during 2009 is scheduled to be available by 31 March 2010. The “draft final” of this interim progress report which incorporates comments received, is scheduled to be completed by 1 June 2010. Data files containing PIT tag information on the federally-listed endangered fish species (Colorado pikeminnow and razorback sucker) collected during this Adult Monitoring trip will be submitted for inclusion in the SJRIP’s integrated database by 31 December 2009. Data files containing the remainder of the information (e.g., data about common fish species) collected during this Adult Monitoring trip will be submitted for inclusion in the SJRIP’s integrated database by 31 March 2010.

Qualifications of Personnel Included in the Budget:

Principal Biologist (GS-12) -- Dale Ryden

Holds a BS degree. Has 18 years experience performing fisheries research/management in the Colorado River Basin. Dale has been performing fisheries research/management in the San Juan River Basin for the last 17 years.

Biological Technicians (GS-7) – Bio. Techs from USFWS-CRFP

All hold at least a BS degree in biology. Depending upon the individual, they have from 1-8 years experience performing fisheries research/management in the Colorado River Basin. Most have 1-3 years of experience performing fisheries research/management on the San Juan River.

Project Leader (GS-14) -- Chuck McAda

Holds an MS degree. Has 30 years experience performing fisheries research/management in the Colorado River Basin. Chuck was chairman of the San Juan River Recovery Implementation Program’s Biology Committee for three years (2004-2006).

Projected Duration Of Project:

The Adult Monitoring study began in 1991 (see Introduction for details). It has continued, annually, with a consistent sampling regime every year since that time. This has allowed for the compilation of one of the longest-running and most statistically powerful fisheries databases available to the SJRIP. The Adult Monitoring study was modified with just very slight changes (e.g., a reduction in sampling frequency from every RM to two out of every three RM’s) when it was incorporated as an integral part of the long-term San Juan River Monitoring Plan and Protocols (Propst et al. 2000). This suite of long-term monitoring studies were initiated in 1999 and are scheduled to run through the termination of the San Juan River Recovery Implementation Program.

Literature Cited:

Propst, D. L., S. P. Platania, D. W. Ryden, and R. L. Bliesner. 2000. San Juan River Monitoring Plan and Protocols. San Juan River Recovery Implementation Program; U. S. Fish and Wildlife Service, Albuquerque, NM. 20 pp. + appendices.

San Juan River Basin Recovery Implementation Program. 2008. Draft Long-range plan (dated 18 April 2008). San Juan River Basin Recovery Implementation Program, U. S. Fish and Wildlife Service, Albuquerque, New Mexico.

See Attached Budget For Line Item Breakdowns	\$ 13,668.00
Utah Division of Wildlife Resources - Moab, UT	
See Attached Budget For Line Item Breakdowns	\$ 4,794.00
	\$ 18,462.00

FY-2009 WORKPLAN TOTAL **\$122,034.00**

Estimated Outyear Funding (based on an annual 5% increase as agreed upon by the SJRIP Biology Committee at their 21 May 2002 meeting):

Fiscal Year 2010		\$128,135	Fiscal
Year 2011	\$134,540		
Fiscal Year 2012		\$141,270	

Under the heading "Funding for participation by other agencies." Costs for participation of the U.S. Fish and Wildlife Service, New Mexico Fish & Wildlife Conservation Office in Albuquerque, NM in FY-2009.

Personnel/Labor Costs (Federal Salary + Benefits)

Principal Biologist (GS-12-3) – 12 days @ \$390/day (1 person x 12 days x 1 trip)	\$ 4,680
Biological Technician (GS-7-2) – 18 days @ \$213/day (1 person x 6 days x 1 trip; Farmington to Four Corners) (1 person x 12 days x 1 trip; Four Corners to Clay Hills)	\$ 3,834
Administrative Officer (GS-9-6) – 1 day @ \$262/day	\$ 262
Sub Total	\$ 8,776

Travel and Per Diem (Based on Published FY-2008 Federal Per Diem Rates)

Hotel Costs – 7 nights (1 night x 2 rooms @ \$86/night; Cortez, CO) (5 nights x 1 room @ \$70/night; Farmington, NM)	\$ 522
Per Diem	
Camping Rate - 20 days @ \$29/day (2 people x 10 days x 1 trip)	\$ 580
Hotel Rate – 6 days @ \$39.00/day	\$ 234
Sub Total	\$ 1,336

Equipment

Vehicle Maintenance & Gasoline (@ \$0.60/mile) (660 miles round trip from Albuquerque, NM to Blanding, UT + 100 miles shuttling)	\$ 456
Generator fuel (45 gallons X \$3.00/gallon)	\$ 135
Equipment Maintenance, Repair, & Replacement (e.g., life jackets, hip boots, generator repair, rubber gloves, dip nets, aluminum welding, raft repair, etc.)	\$ 500
Sub Total	\$ 1,091

USFWS-NMFWCO (Albuquerque) Total **\$ 11,203**

USFWS Region 2 Regional Office Administrative Overhead (22.00%) **\$ 2,465**

USFWS Region 2 Total **\$ 13,668**

Under the heading "Funding for participation by other agencies." Costs for participation of the Utah Division of Wildlife Resources office in Moab, UT in FY-2009.

Personnel/Labor Costs (State Salary + Benefits)

Principal Biologist – 12 days @ \$265/day		<u>\$ 3,180</u>
	Subtotal	\$ 3,180

Travel and Per Diem (Based on Published FY-2007 State Per Diem Rates)

Hotel Costs – 1 night (1 night @ \$70/night)		\$ 70
Per Diem (Hotel Rate) - 1 day @ \$43/day (Camp Rate) - 10 days @ \$20/day		\$ 43
		<u>\$ 200</u>
	Subtotal	\$ 313

Equipment

Vehicle Maintenance & Gasoline (@ \$0.49/mile) (412 miles round trip from Moab, UT to Cortez, CO to Clay Hills, UT)		\$ 202
Equipment Repair, & Replacement (e.g., life jackets, hip boots, waders)		<u>\$ 300</u>
	Subtotal	\$ 502

UDWR- Moab Total		\$ 3,995
-------------------------	--	----------

UDWR- Administrative Overhead (20%)		<u>\$ 799</u>
--	--	---------------

UDWR TOTAL		\$ 4,794
-------------------	--	----------

YOY/Small Bodied Fish Monitoring Fiscal Year 2009 Project Proposal

Principal Investigators: David L. Propst and Yvette Paroz
Conservation Services Division
New Mexico Department of Game and Fish
One Wildlife Way, P.O. Box 25112
Santa Fe, NM 87504
(505 476-8103)
dpropst@state.nm.us
yparoz@state.nm.us

Background:

As set forth in Section 5.7 of the San Juan River Basin Recovery Implementation Program (SJRIP) Long-Range Plan, a long-term monitoring program “to identify changes in the endangered and other native species populations, status, distributions and habitat conditions” was to be developed by the SJRIP Biology Committee. The ichthyofaunal monitoring portion of the San Juan River Monitoring Plan and Protocols (Propst, et al., 2000) was divided into four primary areas, larval fish drift sampling, larval fish seining, young-of-year/small bodied fishes, and sub-adult and adult/large-bodied fishes. The portion of the San Juan River to be monitored extends from the confluence of the Animas and San Juan rivers (Farmington) to Lake Powell (Clay Hills Crossing). The purposes of small-bodied fish monitoring are to document occurrence and mesohabitat of young-of year Colorado pikeminnow, razorback sucker, and roundtail chub; characterize the fish assemblages of primary channel shoreline and near-shoreline mesohabitats, secondary channels, and backwaters; and document and assess changes in the abundance of common native and nonnative small-bodied fishes (including age 0 flannelmouth sucker, bluehead sucker, common carp, and channel catfish). The following work proposal for 2009 is to conduct the young-of-year/small-bodied fishes monitoring effort per protocols set forth in the San Juan River Monitoring Plan and Protocols (SJRMP). Beginning in 2003, specimens collected from each mesohabitat were preserved separately, data were recorded in database by mesohabitat, and annual reporting included summary of species occurrences by mesohabitat. During 2004, 2005, and 2006 autumn monitoring, sampling of primary channel near-shore riffle and run mesohabitats was accomplished, on a trial basis, by using a backpack electrofisher to stun fishes and capture them in a bag seine. This sampling was done every six miles in Geomorphic reaches 6 through 3.

In addition to accomplishing work (field, laboratory, data analyses, and report writing) specific to the young-of-year/small-bodied fish monitoring effort, NMGF personnel participate in telemetry studies, native-nonnative interactions studies (upper and lower San Juan), Colorado pikeminnow augmentation evaluation, and larval fish sampling of the San Juan River Basin Recovery Implementation Program. This work and budgeting for NMGF participation in these activities is included with Scopes of Work for each activity and submitted by Principal Investigator(s) for each.

Study Area:

The study area for YOY/small bodied fish monitoring extends from river mile RM 180.0 (Animas River confluence) in Farmington, New Mexico, downstream to RM 2.9 (Clay Hills Crossing), just above Lake Powell in Utah.

Collections:

Specimens collected will be inspected to determine if any rare fishes (Colorado pikeminnow, roundtail chub, and razorback sucker) are present in a sample or collection. All identifiable rare fish and all other readily identified native fishes will be measured and released. Nonnative fishes are routinely retained. Specimens from each sampled mesohabitat will be counted separately. Retained specimens will be preserved in 10% formalin and returned to the

New Mexico Department of Game and Fish Laboratory for identification, enumeration, and measurement (total length). After processing, all specimens will be accessioned to the UNM Museum of Southwestern Biology, Fish Section.

Objectives:

The objectives of this portion of the San Juan River monitoring effort are to document primary channel shoreline and near-shoreline mesohabitat, secondary channel, and backwater use by age-0 Colorado pikeminnow, razorback sucker, and roundtail chub; obtain data that will aid in the evaluation of the responses (e.g., reproduction, recruitment, and growth) of native and nonnative fishes to different flow regimes and other management actions (e.g., impediment modification); track trends in species populations (e.g., abundance and relative condition); and characterize patterns of mesohabitat use by common native and nonnative small-bodied fishes (including age 0 flannelmouth sucker, bluehead sucker, common carp, and channel catfish). The data will also be available to all researchers and may be used in conjunction with data obtained in other studies to evaluate future management activities.

Methods:

The study reach (Farmington to Clay Hills Crossing) includes geomorphic reaches 6 through 1, with Reach 1 being the most downstream. Primary channel sampling will occur every third mile within the study reach. To the extent possible, all secondary channels will be sampled. Secondary channels are defined as channels having less than 25% of the volume of flow at the time of sampling and are at least 300 m in length. Inflow at the top of a channel is not necessary for it to be classified as a secondary channel. Young-of-year/small-bodies fish monitoring will occur in conjunction with the large-bodied fish monitoring effort. Fieldwork will be accomplished in autumn (late-September through mid-October) and involves one foray through each of three macro-reaches (Farmington-Shiprock, Shiprock-Four Corners, and Four Corners-Cray Hills Crossing).

In addition to structured primary channel sampling, all backwaters and embayments ($>25 \text{ m}^2$) associated with the primary channel within each third-mile will be sampled. Large backwaters (ca. 250 m^2) outside designated sample-miles will be sampled, if possible.

Sample sites within secondary channels will be a sufficient distance from inflow to and outflow from the secondary channel to minimize primary channel faunal and physiochemical influences. Secondary channel sample sites will be at least 100 and not more than 200 m in length. Each mesohabitats (e.g., pool, riffle, riffle-eddy, and shoal) within the site will be sampled in rough proportion to its availability within the site; typically, at least five mesohabitat types will be sampled in each secondary channel with inflow. Mesohabitat of secondary channels lacking inflow is typically only pool. Each mesohabitat will be sampled separately with $3.2 \times 1.6 \text{ m}$ (4 mm mesh) drag seines. Each secondary channel sampling effort will be a minimum of 5 seine hauls (= 5 mesohabitats). The number of seine hauls, area (m^2) of seined portion of each mesohabitat, and types of mesohabitats sampled will be recorded on standard field forms. Specimens collected in each mesohabitat will be inspected to determine if any rare fishes (Colorado pikeminnow, roundtail chub, and razorback sucker) are present in the seine. If a rare fish is captured, it will be identified, total length ($\pm 1.0 \text{ mm}$) and mass ($\pm 1.0 \text{ g}$) determined, and released. Any rare fish $>150 \text{ mm TL}$ will be scanned to determine presence of a PIT tag. If none is present, the specimen will be implanted with a PIT tag having a unique alphanumeric code. All pertinent data (i.e., total and standard lengths, mass, PIT tag code, mesohabitat, water depth, substrate, and cover) on rare fish captured will be recorded. All readily identified native fishes will be identified and counted; all specimens $>150 \text{ mm TL}$ (flannelmouth and bluehead suckers) will be weighted, measured, and released; all specimens $<150 \text{ mm TL}$ will be measure and released. All nonnative fishes will be retained and relevant information recorded by mesohabitat of capture. All retained specimens will be preserved separately by mesohabitat in 10% formalin and returned to the New Mexico Department of Game and Fish Laboratory for identification, enumeration, and measurement (total length). Field collection number, habitat number, and river mile will be recorded on a water-proof label and placed in each specimen container. Location of site (UTM) will be determined with a GPS unit. Identification of all retained rare fishes will be confirmed by personnel of the Museum of Southwestern Biology. Preserved specimens will be accessioned to the University of New Mexico Museum of Southwestern Biology.

Within each third-mile, shoreline habitats of the primary channel will be sampled. At each designated mile, all mesohabitats (e.g., riffle, debris pool, and shoal) along 200 m (near center of mile) of shoreline will be sampled. All mesohabitats present will be sampled in rough proportion to their availability within the site. Regardless of number of mesohabitats present at a primary channel site, at least 5 seine hauls will be made with a drag seine (3.2 x 1.6 m, 4mm mesh). Shoreline (river right or left) sampled will be dependent upon accessibility of the shoreline. Where more than one shoreline is accessible (and can be seined efficiently), that with greater habitat diversity/complexity will be sampled. Alternatively, primary channel island shorelines may be sampled. Procedures for specimen processing, retention, and accounting will be same as for secondary channels.

Backwaters and embayments (>25 m²) not located within structured primary channel sampling sites, but within each designated mile, also will be sampled. During periods of low flow, secondary channel mouths frequently function as backwaters or embayments. In this monitoring effort, secondary channel mouths without surface inflow from upstream will be treated as backwater/embayment habitat. The maximum number of backwaters or embayments sampled will be one per mile. Three seine hauls typically will be made in each backwater or embayment sampled. One seine will be across mouth of backwater (or embayment), one in middle, and one near head of backwater (or embayment). Specimens collected in each seine haul will be preserved separately. All specimens collected, except rare fishes, will be retained and returned to the laboratory for identification and enumeration. All rare fish will be measured and released; those >150 mm will be PIT tagged. Data collection and recording of relevant information (including GPS determined location) will be the same as for secondary and primary channels.

Sampling effort for all seine collections will be number fish/unit area. Electrofishing sampling effort will be number fish/unit area and number fish/electrofishing minute.

Ambient temperature and water quality data (water temperature, dissolved oxygen, conductivity, and salinity) will be measured in each sampled secondary channel, at primary channel sites, and in backwaters/embayments. Secondary channel water quality data will be obtained a sufficient distance from the inflow to the secondary channel to minimize primary channel influences. All water quality data for each sample will be recorded on standard field forms.

Products:

Data collected during the 2009 monitoring effort will be summarized by geomorphic reach. Minimally, the annual report will report density per species (number/m²) per geomorphic reach (primary and secondary channels and backwaters) and rare fishes and the mesohabitats in which each was found. Uni- and multivariate statistics will be used to characterize relationships among species densities and environmental variables, longitudinal patterns of species abundance, and species somatic conditions. River discharge data (Shiprock gage) will be used to assess the effect of discharge volume on species density estimates. All data obtained during 2009 monitoring activities will be electronically recorded in format determined by the SJRIP Biology Committee. The annual report (including electronic database) will be submitted to the SJRIP Biology Committed by 31 March 2010.

Literature Cited:

Propst, D.L., S. P. Platania, D.W. Ryden, and R. Bliesner. 2000. San Juan River Monitoring Plan and Protocols. San Juan Basin Recovery Implementation Program. U.S. Fish and Wildlife Service, Albuquerque, NM.

FISCAL YEAR 2009 BUDGET—SMALL-BODIED FISHES MONITORING¹

FIELD

PERSONNEL

PROJECT LEADER (1)

Tasks - Annual monitoring primary channel, secondary channel, and backwater habitats, San Juan River, Farmington to Clay Hills.

196 hrs	196 hrs
\$39.90/hr (base salary) + \$13.17 (benefits)	\$53.067/hr
TOTAL PROJECT LEADER SALARY	\$10,401.13

PROJECT BIOLOGISTS (2)

Tasks—Annual monitoring primary channel, secondary channel, and backwater habitats, San Juan River, Farmington to Clay Hills.

152 hrs ea ²	304 hrs
\$28.35.00/hr (base salary) + \$9.36 (benefits)	\$37.71/hr
TOTAL PROJECT BIOLOGISTS SALARY	\$11,463.84

PER DIEM

17 days/project biologist	34 days
\$85.00/day (standard NM in-state rate)	\$85.00/day
TOTAL PER DIEM	\$2,890.00

TRAVEL

4 x 4 vehicles (2) 400 mi (round-trip Farmington) ea.	800 miles
75 mi/day x 5 days ea.	750 miles
600 mi (round-trip Clay Hills) ea.	1200 miles
\$0.32/mile (standard NM rate)	\$0.32/mile
TOTAL VEHICLE	\$880.00

FIELD EQUIPMENT & SUPPLIES

Seines (6) @ \$50.00 ea	\$300.00
Whirlpacks (500) @ \$50.00/500	\$ 50.00
Formalin (30 gal) @ \$25/5gal	\$150.00
TOTAL EQUIPMENT & SUPPLIES	\$500.00

TOTAL FIELD**\$26,134.97****SPECIMEN MANAGEMENT****PERSONNEL****PROJECT BIOLOGISTS (2)**

Tasks—processing (sorting, identification, and data-entry) ca. 400 primary channel seining samples, 150 secondary channel seining samples, and 20 backwater seining samples.

Since 2000, an annual average of 31,000 specimens (retained and released) have been processed.

320 hrs ea.	640 hrs
\$28.35.00/hr (base salary) + \$9.36 (benefits)	\$37.71/hr
TOTAL SPECIMEN MGMT SALARY	\$24134.40

LABORATORY SUPPLIES

Ethyl alcohol (50 gal) @ \$450.00/50 gal	\$450.00
Specimen containers (misc. vials & jars)	\$500.00
TOTAL LABORATORY SUPPLIES	\$950.00

TOTAL SPECIMEN MANAGEMENT**\$25,084.40**

DATA SYNTHESIS & REPORT PREPARATION**PERSONNEL****PROJECT LEADER (1)**

Tasks—data analysis, data synthesis, report drafting (primary channel, secondary channel, backwater, and summary sections), report review, and report revision.

120 hrs	120 hrs
\$39.90/hr (base salary) + \$13.17 (benefits)	\$53.067/hr
TOTAL PROJECT LEADER SALARY	\$6,367.68

PROJECT BIOLOGISTS (2)

Tasks—data management, data QA/QC, data analysis, data synthesis, table and graph preparation, report drafting (primary channel, secondary channel, and backwaters sections), and report revision.

200 hrs ea.	400 hrs
\$28.35.00/hr (base salary) + \$9.36 (benefits)	\$37.71/hr
TOTAL PROJECT BIOLOGISTS SALARY	\$15,084.00

SECRETARY/CLERK

Tasks—time record keeping & reporting, billing, supply orders, and budget management.

40 hrs.	40 hrs
\$21.00/hr (salary) + \$6.93 (benefits)	\$27.93/hr
TOTAL SECRETARY/CLERK SALARY	\$1117.20

TOTAL DATA SYNTHESIS & RPT PREPARATION **\$22,568.88**

REVIEWS AND MEETINGS**PERSONNEL****PROJECT BIOLOGISTS (1)**

Tasks—attendance at 3 Biology Committee meeting annually (28 hrs. ea) and annual report review (excluding NMGF; 24 hrs).

108 hrs	108 hrs
\$28.35.00/hr (base salary) + \$9.36 (benefits)	\$37.71/hr
TOTAL PROJECT BIOLOGIST SALARY	\$4,072.68

SECRETARY/CLERK

Tasks—travel arrangements, etc.

20 hrs	20 hrs
\$21.00/hr (salary) + \$6.93 (benefits)	\$27.93/hr
TOTAL SECRETARY/CLERK SALARY	\$558.60

PER DIEM**PROJECT BIOLOGIST (1) (includes 3 Biology & 1 Coordination Committee meetings)**

15 days @ \$85.00/day (standard NM in-state rate)	\$1,275.00
6 days @ \$115.00/day (standard NM out-of-state rate)	\$690.00
TOTAL PER DIEM	\$1,965.00

TRAVEL

VEHICLE

5 Biology & Coordination Committee meetings (Farmington) @ 400 miles ea. 2000 miles @ \$0.32/mile (standard NM rate)	\$640.00
2 Biology & Coordination Committee meetings (Durango) @ 500 miles ea. 1000 miles @ \$0.32/mile (standard NM rate)	\$320.00
TOTAL VEHICLE	\$960.00

TOTAL REVIEWS & MEETINGS **\$ 7,556.28**

TOTAL **\$ 81,344.53**
INDIRECT COSTS (10%) **\$ 8,134.45**

GRAND TOTAL **\$ 89,478.98**

¹Budget does not include in-kind contributions of about \$40,000 per year in salary and benefits, equipment, and some supplies. In-kind includes field time, data analyses and report preparation, and project administration.

²16 additional hours per biologist to cover overtime associated with field work

Outyear Funding (based on 5% annual cost of living increases):

Fiscal Year 2000	\$57,200
Fiscal Year 2001	51,700
Fiscal Year 2002	51,700
Fiscal Year 2003	49,775
Fiscal Year 2004	63,545
Fiscal Year 2005	72,645
Fiscal Year 2006	72,885
Fiscal Year 2007	\$81,246
Fiscal Year 2008	\$91,882
Fiscal Year 2009	\$89,479

SAN JUAN RIVER
LARVAL RAZORBACK SUCKER AND LARVAL COLORADO PIKEMINNOW SURVEYS
FISCAL YEAR 2009 PROJECT PROPOSALS

Summary Regarding Proposed FY 2009 Budget Increase for Larval Fish Surveys

Starting with FY 2007 funds, the responsibility for the San Juan River larval razorback sucker and larval Colorado pikeminnow studies was shifted from the University of New Mexico (UNM) to the New Mexico Department of Game and Fish (NMGF). An RFP for that work was let by NMGF and awarded to American Southwest Ichthyological Researchers (ASIR). A key provision of that RFP was the seamless transition of the project from UNM to NMGF. By the time the new contract was in place, funds for the 2007 effort were already allocated and the budget for FY 2008 study was well along in the SJRBRIP review process. As such, no attempts were made to adjust the FY 2007 and FY 2008 budgets to reflect new contracting costs. The FY 2009 Scope of Work and associated budget is the first opportunity under the new contract that a request for modification could be submitted.

What follows is an abbreviated summary of information regarding development of the requested budget increase for these two projects. During 2007, in an effort to have future budgets reflect actual costs, we worked with NMGF to initiate stricter accounting practices for these projects than had been available while the contract was at UNM. The results of that process and a little over one-year of being in charge of the contract is a new budget and higher cost than had been previously provided to the SJRBRIP. We intend to refine the costs of these projects on an annual basis and provide an explanation of any changes in future budgets.

Key parameters in determining the new budgets:

- Budgets of the two projects (razorback sucker and Colorado pikeminnow) are combined
- Costs are determined per trip (six trips per year)
- Study Area is from Cudei (RM 141.53) to Clay Hills Crossing (RM 2.9)
- Study Area is broken into two sections (upper and lower) for sampling purposes
- Discrete sampling crews work each study section (= two sampling crews)
- Discrete sampling crews = two complete sets of all necessary field equipment
- Sampling (launch of rafts) occurs simultaneously in the two reaches
- There is no cost differentiation (field or laboratory) between individual monthly trips – (we are attempting to acquire that information in 2008)
- Effort is concentrated on larval fish
- Use current reporting protocols (follow & provide formal response to reviewer comments)
- Required meeting attendance for researchers is twice per year
- Budget information is based on W. Howard Brandenburg and Michael A. Farrington as Principal Investigators (would be higher if we did not have them in charge of projects)
- Indirect Costs not changed from original but are not reflective of actual costs

SAN JUAN RIVER LARVAL RAZORBACK SUCKER SURVEY
FISCAL YEAR 2009 PROJECT PROPOSAL

Principal Investigators: W. Howard Brandenburg and Michael A. Farrington
 American Southwest Ichthyological Researchers, L.L.C. (ASIR)
 800 Encino Place NE
 Albuquerque, New Mexico 87102-2606
 505.247.9337 (voice) 505.247.2522 (facsimile)
 howard_brandenburg@asirllc.com michael_farrington@asirllc.com

and

Co-Principal Investigator: David L. Propst
New Mexico Department of Game and Fish
Conservation Services Division
One Wildlife Way
Santa Fe, New Mexico 87505
505.476.8103
dpropst@state.nm.us

Project History:

The apparent absence of razorback sucker in the San Juan River drainage necessitated experimental stocking of adults (n=672) of this species in 1994 between Hogback, New Mexico, and Bluff, Utah. In their 1995 report of activities, Ryden and Pfeifer (1996) suggested that the majority of the 1994 experimentally stocked razorback sucker would achieve sexual maturity in 1996 and spawning by those individuals might begin a few years afterwards.

At the November 1996 San Juan River Basin Biology Committee integration meeting, it was suggested that the Colorado pikeminnow, *Ptychocheilus lucius*, larval fish drift study (= Passive Drift Netting Study; RM 128.0 and RM 53.3; July-August) be expanded in an attempt to document spawning of the stocked razorback sucker (presumed to be during April-May). In addition to temporal differences in spawning between Colorado pikeminnow and catostomids (suckers), researchers were attempting to document reproduction by hatchery reared razorback sucker whose spawning potential was unknown. Sampling for larval razorback sucker was to be conducted to determine if the stocked population of adult razorback sucker would spawn in this system. Conversely, data from the passive drift-netting study continued to document Colorado pikeminnow reproduction in the San Juan River and, because of this certainty, larval fish sampling efforts for this fish would (initially) be different than those for razorback sucker.

Numerous Upper Colorado River Basin researchers reported light-traps as one of the best means of collecting larval razorback sucker. Most of their (Upper Colorado River basin researchers) efforts were concentrated in floodplain habitats during high spring flows. Light-trap sampling was employed during the first year (calendar year 1997) of the San Juan River larval razorback sucker survey. The lack of inundated floodplain habitats in the San Juan River, in comparison to the Upper Colorado River Basin, meant that the light-traps would have to be set in low velocity riverine habitats. The only previous San Juan River fish investigations that had employed light-traps were in 1994 and 1995 (conducted by the National Park Service) near the San Juan River-Lake Powell confluence. That sampling effort produced an extremely large number of larval fish (ca. 25,000) from a modest number of samples (n=20), of which over 99% were red shiner. Similar sampling in 1995 yielded 25,455 specimens in 47 light-traps samples and as in 1994, red shiner numerically dominated the catch. Both sampling efforts were conducted during July-August but neither Colorado pikeminnow nor razorback sucker was present in the 1994-1995 light-trap samples.

During the 1997 razorback sucker larval fish survey, light-traps were set nightly in low-velocity habitats between Aneth and Mexican Hat, Utah, from late March through mid-June. The traps were distributed at dusk and retrieved about four hours later. Fish taken in those samples were preserved in the field. Sampling success during the 1997 razorback sucker larval fish study was poor. While there were over 200 light-trap sets, those sampling efforts produced only 297 fish. Of those, about 200 (66%) were larval suckers (either flannelmouth sucker or bluehead sucker). Larval razorback sucker were not present in the 1997 sampling survey. While there were probably several factors to account for the poor light trap catch rate, a principal factor was the limited access to suitable habitats. Light-traps are most effective when set in habitats with little or no water velocity. During our driving survey of riverine habitats in the region (March 1997), we identified numerous locations that appeared to be suitable sites for light trap sampling. However, high spring flow in the San Juan River eliminated virtually all previously identified low velocity habitats. Further driving reconnaissance failed to yield additional locations to set light traps. We determined that being limited to specific collecting sites was not the most efficient means of collecting large numbers of individuals; a prerequisite for this study.

In 1998 a new study design was developed to allow for the sampling of a greater portion of the San Juan River and the collection of a significantly larger number of larval fish through out several river reaches. An inflatable raft was used to traverse the San Juan River and allowed us the opportunity to sample habitats that were either not formerly accessible or observable under the constraints of the previous sampling protocol. Six sampling forays were conducted at approximately bi-weekly intervals from 17 April to 6 June 1998 between the Four Corners (RM 127.5) and Mexican Hat, Utah (RM 53.3). Both active (seining) and passive (light-traps) sampling techniques were used to collect larval fish. The primary sampling method was a fine mesh larval seine. If appropriate aquatic mesohabitats could be located, light-traps would be set adjacent to nightly campsites of the sampling crew.

The 1998 sampling protocol resulted in 183 collections containing over 13,000 specimens between river miles 128.0 and 53.3 with the majority of these individuals (n=9,960) being larval catostomids. This 43-fold increase in number of specimens, as compared with 1997, provided substantially better resolution of spawning periodicity of the catostomid community. In addition, the 1998 samples produced enough individuals for us to determine, with a high degree of confidence, if razorback sucker reproduction occurred in the San Juan River during that period. None of the aforementioned information was obtainable from 1997 light-trap samples. In 1998, two larval razorback sucker were collected providing verification of spawning by the hatchery reared stocked population.

The use of active sampling to determine the reproductive success of razorback sucker has proven to be effective. To date, the results of this investigation have provided ten consecutive years of unequivocal documentation of reproduction in the San Juan River by razorback sucker that have been stocked as part of the San Juan River Basin Recovery Implementation Program (Table 1). The data collected during the larval razorback sucker survey provide not only valuable data concerning the distribution (spatial and temporal), duration and magnitude of razorback sucker reproductive efforts but also equally informative data on the reproductive efforts of other native catostomids in the San Juan River.

Table 1. Collection information razorback sucker (*Xyrtex*) collected during the larval razorback sucker survey, 1998 - 2007.

<i>Year</i>	<i>Sampling method</i>	<i>Study Area (River Miles)</i>	<i>River Miles sampled</i>	<i>Percent change</i>	<i>Specimens collected</i>	<i>Xyrtex n=</i>
1998	Larval seine Light traps	127.5 – 53.3	74.2	na	13,608	2
1999	Larval seine Light traps	127.5 – 2.9	124.6	+ 40.4%	20,348	7
2000	Larval seine Light traps	127.5 – 2.9	124.6	na	11,473	129
2001	Larval seine Light traps	141.5 – 2.9	138.6	+ 10.1%	95,628	50
2002	Larval seine Light traps	141.5 – 2.9	138.6	na	56,034	813
2003	Larval seine Light traps	141.5 – 2.9	138.6	na	41,181	472
2004	Larval seine	141.5 – 2.9	138.6	na	14,642	41
2005	Larval seine	141.5 – 2.9	138.6	na	19,139	19

2006	Larval seine	141.5 – 2.9	138.6	na	25,127	202
2007	Larval seine	141.5 – 2.9	138.6	na	22,093	200

This work is being conducted as required by the 31 March 2000 San Juan River Basin Recovery Implementation Program Monitoring Plan and Protocol. The objectives of this specific monitoring effort are identified and listed below. Where applicable, these objectives are related to the specific tasks listed in the Long Range Plan set forth by the San Juan River Basin Recovery Implementation Program (SJRBRIP).

Project Modifications:

There have been numerous modifications to the original (1997) sampling methodology of the San Juan River larval razorback sucker survey as well as changes in reporting priorities, protocol, and format. The extent of the study area and aspects of the longitudinal sampling have been modified to improve spatial comparisons. The study area for this project has been expanded (in 2001) by 46.5% (64.4 river miles) and now includes from the middle of Reach 5 (Cudei, New Mexico) to the downstream-most end of Reach 1 (Clay Hills Crossing, Utah; total of 138.6 miles of critical habitat sampled). Beginning in 2003, the entire 139 river miles of the study area was sampled in single uninterrupted trips (10-12 field days per trip) rather than in two temporally discrete sections as done in previous years (1998 – 2002). Since greater numbers of larval razorback sucker were collected (as well as detailed information regarding the native fish community), the SJRBRIP Biology Committee voted to elevated the razorback larval survey from an “experimental” project to a monitoring program. This change allowed for comparisons of catch per unit effort (CPUE) with the program designated river reaches and facilitated integration of the larval survey data with that of the other monitoring activities (i.e., small bodied fish, adult monitoring, habitat, etc).

Conducting the larval razorback sucker survey under this new protocol not only provided discreet reach information but also provided greater temporal resolution in respect to the longitudinal distribution of razorback sucker larvae and potential environmental cues required by razorback sucker for spawning. Disadvantages to this top to bottom approach were that the duration of the monthly sampling trips (10-12 field days) made them more subject to abiotic fluctuations (floods, flow spikes). Annually, at least one trip (an average) had to be cut short due to large flood events or low water events in the lower canyon. The abbreviated trips were subsequently resumed once conditions improved (usually 1-2 weeks later). Large flood events not only disrupted the temporal resolution of the single-continuous pass effort but it also reduces sampling efficiency as many low velocity habitat are flooded by rising water levels thereby transporting larval and early juvenile fish in the drift. Additional costs were incurred because of the need to return to the field to complete the sampling effort for that month.

Another problem of that resulted from cancelled or disrupted trips during the single-continuous pass effort was that designated camp sites in the lower canyon (scheduled and coordinated by BLM) had to be canceled and often new camp sites could not be provided. Logistically, a single raft could not carry enough supplies for a crew of two/three for 11-12, so food and water and fuels for cooking had to be transported to and stored at Mexican Hat for re-supply. Concerns were also made regarding the safety and fatigue to crews due to the long days (10-12 hours) and labor-intensive trip schedules. To mitigate these effects as well as gain even greater temporal resolution of the longitudinal distribution of razorback sucker larvae, the protocol was changed to allow sampling of the upper (RM 142 – 53) and lower (RM 53 – 3) sections of the San Juan River simultaneously. This effort began in 2007 and utilized two fully equipped and discrete crews (comprised of 2 people per crew) with one crew sampling the upper and the other sampling the lower portion of the study area simultaneously (Table 2). In 2008, addition participation of our staff with other SJRBRIP projects made the new simultaneous sampling effort a necessity so that our staff could meet obligations to assist the other researcher with their work.

Table 2. Summary project and project modification of the larval razorback sucker survey from 1997 to 2008.

<i>Year</i>	<i>Sampling method</i>	<i>Study Area (River Miles)</i>	<i>Specimens collected</i>	<i>Field modification</i>	<i>Laboratory modification</i>
1997	Light Trap	99 – 75	297		
1998	Larval Seine Light Trap	127.5 – 53.3	13,608	study area expanded; active sampling	
1999	Larval Seine Light Trap	127.5 – 2.9	20,348	study area expanded; upper-lower reaches sampled separately; nonsynchronous	
2000	Larval Seine Light Trap	127.5 – 2.9	11,473		
2001	Larval Seine Light Trap	141.5 – 2.9	95,628	study area expanded; upper-lower reaches sampled separately; nonsynchronous	
2002	Larval Seine Light Trap	141.5 – 2.9	56,034		
2003	Larval Seine Light Trap	141.5 – 2.9	41,181	upper-lower reaches sampled monthly in one uninterrupted trip (11-12 day runs)	CPUE data used for integration in reporting
2004	Larval Seine	141.5 – 2.9	14,642		Reports merged Trend data
2005	Larval Seine	141.5 – 2.9	19,139		
2006	Larval Seine	141.5 – 2.9	25,127		
2007	Larval Seine	141.5 – 2.9	22,093	Two rafts-two crews; upper-lower reaches samples synchronous	Analyzed catch with habitat data
2008	Larval Seine	141.5 – 2.9			

Study Area:

The study will be the San Juan River between Cudei, New Mexico (RM 141.5) and the Clay Hills Crossing boat landing (RM 2.9) just above Lake Powell in Utah. As in all post 1999 sampling efforts, the study will include making collections in reaches of the San Juan River under the jurisdiction of the National Park Service.

Objectives:

- 1) Determine the spawning periodicity of razorback sucker and other native catostomids between mid-April and mid June and examine potential correlation with temperature and discharge (Task 4.1.5)

- 2) Attempt to validate presumed spawning period of San Juan River catostomids using data from the razorback sucker and Colorado pikeminnow larval fish studies.
- 3) Provide comparative analysis of the reproductive effort of catostomids.
- 4) Determine the relative annual reproductive success of razorback sucker (Task 4.1.1)
- 5) Provide annual summaries of monitoring results.
- 6) Provide detailed analysis of data collected to determine progress towards endangered species recovery in three years and thence every five years.

Methods:

Field work:

Sampling for Colorado pikeminnow larvae will be conducted in the San Juan River between RM 141.5 and RM 2.9 from early July through early September using sampling techniques that will provide sufficient numbers of fish necessary to meet study objectives. Access to the river will be gained through the use of inflatable rafts equipped with all of the necessary equipment and provisions needed for trips of up to ten days. The study area will be divided into an "upper" section (Cudei, NM, to Mexican Hat, UT) and a "lower" section (Mexican Hat, UT, to Clay Hills crossing, UT). Separate field crews will launch their rafts simultaneously in each of the two sections and proceed through their designated study area. The vehicle and raft trailer used by the field crew working in the upper section will be left at the Cudei launch site and subsequently be shuttled to Valles Trading Post in Mexican Hat, UT, where it will be placed in paid storage. The vehicle shuttle (with trailer) for the upper reach sampling effort has typically been performed gratis by personnel from the Farmington Office of the Bureau of Indian Affairs Office. Starting in 2008, this service will be performed by personnel from the N.M. Fishery resources Office stationed in Farmington. At this time, there is no charge for this service.

The sampling crew for the lower reach will launch from and store their vehicle and raft trailer at Valles Trading Post in Mexican Hat, UT, where a commercial shuttle (from Valles) will take the vehicle to Clay Hills crossing. The cost for this service is included under the travel and per diem section of our budget. Cold storage facilities are also available at Valles Trading Post and will be used on an as needed basis. Because crews sampling the lower section of the study area will be in a high use recreational area, advance reservations are required. All trips for 2009 must be scheduled by late January 2009 and submitted to the Bureau of Land Management (BLM) Office at Monticello, Utah. Designated camping permits for our lower reach sampling crews will be obtained and must be strictly adhered to in addition to other BLM- San Juan River recreation Area regulations (i.e., low impact and pack-out policies). Low flow conditions often prevalent during the study period make several sections of the river more difficult to navigate (especially in the lower reach). Our field crews are required to render assistance to boaters stuck in rapids or otherwise in distress and report all such encounters to the appropriate BLM personnel (will add one extra day to the lower reach sampling effort).

Sampling efforts for larval fish will be concentrated in low velocity habitats and employ small mesh seines (1 m x 1 m x 0.0) to collect fish. Retained specimens will be placed in Whirl-paks containing a solution of 10% buffered formalin and a tag inscribed with unique alphanumeric code that is also be recorded on the field data sheet. For seine samples, the lengths (to 0.1 m) of each seine haul and total number of hauls will be measures and recorded. Catch per unit effort for seine samples will be reported as the number of fish per 100 m².

Native species large enough to be positively identified will be measured (standard length) and returned to the river. Post-larval endangered fish species collected during this study will be photographed, a small

portion of tissue from the fin clipped and retained in 95% EtOH (in the case of potential razorback sucker hybrids) and scanned with a FS2001 PIT tag reader for the presence of a PIT tag. Specimens of sufficient size but lacking a PIT tag will be injected with a tag. All PIT tag information will be recorded in the field data sheet and subsequently forwarded to the SJRBRIP for integration in the program's PIT tag database.

For each sampling locality, river mile will be determined to the nearest tenth of a mile using the San Juan River Basin Recovery Implementation Program 2003 Standardized Map Set. Universal Transverse Mercator (UTM) coordinates and zone will be determined with a Garmin Navigation Geographic Positioning System Instrument for each sampling locality. Mesohabitat type, length, maximum and minimum depths, water clarity (determined with a Secchi disc), and substrata will be recorded for each sampling locality. Multi-parameter YSI units will be used to determine the following water quality parameters at each site sampled: pH, temperature, salinity, conductivity, specific conductance, and dissolved oxygen. A minimum of one digital photo will also be taken of each specific habitat sampled.

StowAway Tidbit temperature loggers will be set to record water temperatures hourly and deployed near Four Corners bridge, in McElmo Creek, and at Clay Hills crossing. The data from each temperature recorder will be downloaded monthly during the study period but remain in the river after completion of the annual study effort to record river water temperatures throughout the year.

Field Work, Safety:

Personnel participating in field work are required to successfully completed an International Rescue Instructors Association (IRIA) level 2 swiftwater rescue class and American Red Cross CPR/AED training. Type III personal flotation devices (PFD's) will be worn by sampling personnel at all times while working. As PFD's lose flotation capacity due to UV exposure, compression of material, and oil and grit impregnation, and since each crewmembers PFD will be used for approximately 45 days per season, the PFD's will be annually replaced. Simms Guideweight Gore-Tex waders and boots will be issued to all personnel along with 3 mm neoprene gloves (necessary in April and May). In addition to personal camping gear and rain jackets, all personnel will be required to provide and use wide brimmed hats, sunscreen, and sunglasses (provided at no cost to the program).

Both rafts used for this project will carry an extensively stocked first aid kit replete with items necessary for most minor medical situation. Additionally, the first aid kit will contain a suite of items (i.e., splints, neck braces, butterfly stitches, snakebite kits) needed address more serious medical conditions. Because formalin is used in the preservation of specimens, several vials of eyewash solution will be incorporated into each first aid kit. First aid kits will be inventoried after each sampling trip and used and/or expired items replaced. In the upper reach of the study area, personal cell phones and PDA's will be used (at no cost to the program) to contact outside parties should a medical situation arise. In the lower study area reach (canyon bound; where cell phones do not have service) an Iridium 9505-satellite phone will be provided for sampling crews to be used in case of an emergency.

Formalin used for fish preservation will be transported in heavy-duty LPDE carboys. Extensive exposure to UV light makes the carboys susceptible to decomposition and cracking and requires that they be inspected monthly and not used for more than two years. Safety rope throw bags will be similarly inspected and retired from use accordingly. Rafts will be equipped with raft recovery (Z-line) kits, well stocked supply and repair kits, extra oar and oar blade, and two spare hand pumps to help ensure that crews do not become stranded due to raft damage.

Laboratory Work:

Samples will be returned to the lab immediately after each field trip is completed and processed following a multi-step procedure. To maintain the larval fish in good condition (necessary to ensure accurate identification) the samples must be transferred from whirl-packs to glass jars and the field fluids replaced with a solution of 5% buffered formalin. Cyprinid and catostomid larvae are extremely small and

transparent especially at early developmental stages. To minimize the potential loss of fish in individual seine hauls, it is best to retain the entire contents of each seine haul. A negative result of this technique is that, in addition to larval fish, whirl-pack samples usually contain considerable debris, detritus, and silt. Another important step in processing of individual samples is to separate fish from the detritus. This necessary portion of the process is labor intensive and can be quite tedious. During this process initial sorting of fish based on age class (age 0 [larvae] and age 1+) occurs. Samples that contain a large number of larval fish, especially those proto or mesolarvae, often must be sorted twice.

After the fish are separated from the debris, personnel with San Juan River Basin larval fish identification expertise identify individual specimens to species. Bottom-lit stereomicroscopes equipped with polarized filters (that enhance the delineation of myomeres, pterygiophores, and fin rays) are used to assist with the identifications. Larval fish keys are referenced to assist in species specific determinations (e.g., Contributions to a guide to the cypriniform fish larvae of the Upper Colorado River System [Snyder 1981], Catostomid fish larvae and early juveniles of the Upper Colorado River basin, Morphological descriptions, comparisons, and computer interactive key [Snyder and Muth 2004], and Identifications of larval fishes of the Great Lakes Basin [Auer 1982]). Age-0 specimens are separated from age-1+ specimens using published literature on growth (Snyder 1981, Snyder and Muth 2004).

Age classes are enumerated, measured (minimum and maximum size [mm standard length] for each species at each site), transferred to 70% ethyl alcohol, and catalogued in the Division of Fishes of the Museum of Southwestern Biology (MSB) at the University of New Mexico (UNM). Both total length (TL) and SL of Colorado pikeminnow and razorback sucker are obtained using electronic calipers. The ontogenetic stage of Colorado pikeminnow and razorback sucker obtained in this study is determined based on the definitions provided by Snyder (1981).

Reporting and permitting:

Beginning in 2004, data from the two San Juan River larval fish surveys (razorback sucker and Colorado pikeminnow) were analyzed collectively and presented in a single report. This created for a whole picture of the reproductive activities of the entire ichthyofaunal community in the San Juan River using the same criterion used as the other monitoring programs.

The results in our annual report pertain almost exclusively to age-0 fish (i.e., as fish > age-0 are not “larval fish” and are not the focus of this effort, they are not included in analysis). The number of fish >age-0 collected during the study is presented in an Appendix. Differences in mean CPUE are determined by species between years using a one-way Analysis of Variance (ANOVA). A Poisson distribution provided the best fit to the raw data. A variety of transformations (e.g., logarithmic, reciprocal, square root) were applied on the mean CPUE data for between year comparisons. A natural log transformation yielded the best variance-stabilizing qualities and produced a relatively normal distribution. Pair-wise comparisons between years (2003 – 2007) were made for each species and significance (i.e., $p < 0.05$) was determined using the Tukey-Kramer HSD test. Finally, a nonparametric ANOVA (Kruskal–Wallis test) was used on various data sets to compare results to the parametric analyses. While both ANOVA and Kruskal-Wallis were used to analyze data, data transforms enabled use of parametric analysis in all cases. The assumption of homogeneity of variances was assessed using the more conservative variance ratio criterion of <3:1 (Box, 1954), as opposed to <4:1 (Moore, 1995), among years. All species data sets met this more rigorous criterion and in most cases the variance ratio was <2:1 among years. Additionally, the significance values between parametric and nonparametric techniques were nearly identical and so only the parametric analysis will be presented.

Hatching dates for larval Colorado pikeminnow are calculated using the formula:

$$-76.7105 + 17.4949(L) - 1.0555(L)^2 + 0.0221(L)^3, \text{ where } L = \text{length (mm TL).}$$

Spawning dates larval Colorado pikeminnow are then estimated by adding five days to the post-hatch ages to account for incubation time at 20 - 22°C (Nesler et al. 1988). Hatching dates of razorback sucker larvae

were calculated by subtracting the average length of larvae at hatching (8.0 mm TL) from the total length at capture divided by 0.3 mm (Bestgen et al. 2002), which was the average daily growth rate of wild larvae observed by Muth et al. (1998).

This study is initiated prior to spring runoff and completed near the end of summer (late September). Daily mean discharge during the study period is acquired from U.S. Geological Survey Gauge (# 09379500) near Bluff, Utah. Water temperatures (mean, maximum, and minimum) are acquired from our temperature loggers and additional data loggers (maintained by Keller-Bliesner Engineering) at the Colorado State Highway 160 bridge crossing (RM 119.2) and Mexican Hat, Utah (RM 53.3).

In addition to the annual report of the study provided to the SJRBRIP, reports summarizing fish collecting activities and specimens captured are also required annually under scientific collection permits provided by the New Mexico Department of Game and Fish, Navajo Nation, and state of Utah. The aforementioned reports include (at a minimum) site localities, GPS coordinates, and fish collected. An annual report of activities is a BLM (Monticello Field Office) requirement under our access permit to the San Juan River below San Island (Bluff UT) and designated camps in the lower reaches of the river.

Meetings:

Researchers are required to attend a minimum of two meetings annually and report on annual monitoring projects. The two meetings (February and May) require researchers present PowerPoint presentations outlining the results and that years findings. Each meeting lasts about three days (which includes travel time).

Products:

A draft report of the 2009 larval razorback sucker sampling activities (combined with 2009 larval Colorado pikeminnow sampling activities) will be prepared and distributed to the San Juan River Basin Biology Committee for review by 31 March 2010. Upon receipt of written comments, that report will be finalized and disseminated to members of the San Juan River Basin Biology Committee by 1 June 2010. Electronic copies of the 2009 collection data will be transferred to the San Juan River database manager. Fish collected from this study will be curated in the Division of Fishes, Museum of Southwestern Biology (MSB), Department of Biology, at the University of New Mexico under a MSB contract with the SJRBRIP. Original field notes will be retained in the Division of Fishes and collection information electronically stored in a permanent MSB database program. These data and any maps generated from them will be available to the San Juan River Basin Biology Committee via hard-copy reports and electronically.

SAN JUAN RIVER LARVAL COLORADO PIKEMINNOW SURVEY FISCAL YEAR 2009 PROJECT PROPOSAL

Principal Investigators: Michael A. Farrington and W. Howard Brandenburg
American Southwest Ichthyological Researchers, L.L.C. (ASIR)
800 Encino Place NE
Albuquerque, New Mexico 87102-2606
505.247.9337 (voice) 505.247.2522 (facsimile)
michael_farrington@asirllc.com howard_brandenburg@asirllc.com

and

Co-Principal Investigator: David L. Propst
 New Mexico Department of Game and Fish
 Conservation Services Division
 One Wildlife Way
 Santa Fe, New Mexico 87505
 505.476.8103
 dpropst@state.nm.us

Project History:

Beginning in spring 1995, personnel from the Division of Fishes, Museum of Southwestern Biology (MSB), at the University of New Mexico assumed responsibility for the San Juan River larval fish passive drift-netting study. This project, formerly conducted by the Utah Division of Wildlife Resources, continued through 2001 with only minor changes in sampling protocol. Between 1995 and 2001, a total of four larval Colorado pikeminnow were collected using this sampling method at two different collecting locations (Four Corners, NM and Mexican Hat, UT).

The limited number of wild adult San Juan River Colorado pikeminnow (versus stocked individuals) was reflected in the extremely low catch rate of larval Colorado pikeminnow. Numerous adult and sub-adult pikeminnow have now been stocked into the San Juan River in an effort to augment the diminished wild population with the Colorado pikeminnow augmentation plan calling for continued stocking efforts in the San Juan River over the next 10 years. The San Juan River Basin Biology Committee hopes, as was documented with stocked razorback sucker, that reproduction among stocked pikeminnow will occur and can be documented through the sampling of larval fish.

As the number of adult (reproductively mature) Colorado pikeminnow in the San Juan River increases (due to both stocking and recruitment), so does the probability of elevated levels of spawning by this species. The San Juan River Basin Biology Committee began exploring the possibility of expanding the sampling effort for larval Colorado pikeminnow in fiscal year 2003. One means of accomplishing this task was to include an additional sampling site (increasing from two to three sites) for the passive drift-netting study. Another suggestion was to perform targeted sampling for Colorado pikeminnow similar to that performed for larval razorback sucker. In the case of the latter sampling effort, discussion regarding sampling that would target larval Colorado pikeminnow centered around expanding the duration of the current larval razorback sucker survey (April-June) or development of a discrete (new) project.

These and other items were considered and evaluated during the February 2002 San Juan River Basin Biology Committee meeting. The Committee recommended the immediate expansion of the larval razorback sucker survey (April-June) to include the months of July, August, and September with seining efforts to target sampling for larval Colorado pikeminnow.

Beginning in July of 2002, using funds from FY 2002 that had been appropriated for use at the two larval drift-netting stations, Museum of Southwestern Biology (MSB) personnel began an active sampling regime that mirrored the sampling protocol successfully used in the larval razorback sucker survey. About 140 river miles of the San Juan River (Cudei, NM, to Clay Hills crossing, UT) was sampled monthly by multiple personnel using inflatable rafts to access suitable habitats within the new study area. Five wild larval Colorado pikeminnow have been collected between 2002 and 2007, at five discrete sites, within the aforementioned study area.

Between 1995 and 2007, the combined sampling methodologies (passive and active) resulted in the collection of nine Colorado pikeminnow. Back-calculated spawning dates, based on those nine individual larvae, range between 24 June and 18 July (Table 1) and are generally associated with the descending limb of the spring run-off hydrograph.

Table 1. Summary of larval and YOY Colorado pikeminnow collected in the San Juan River during larval drift-netting/larval seining (1993-2007) and back-calculated dates of spawning.

Field Number	MSB Catalog Number	N	Total Length (mm)	Collected	Date Spawned	River Mile	Sample Method
JPS95-205	26187	1	9.2	02 Aug 1995	15 Jul 1995	53.0	drift netting
JPS95-207	26191	1	9.0	03 Aug 1995	17 Jul 1995	53.0	drift netting
WHB96-037	29717	1	8.6	02 Aug 1996	18 Jul 1996	128.0	drift netting
FC01-054	50194	1	8.5	01 Aug 2001	17 Jul 2001	128.0	drift netting
MAF04-046	53090	1	14.2	22 Jul 2004	24 Jun 2004	46.3	larval seine
MAF04-059	53130	1	18.1	26 Jul 2004	25 Jun 2004	17.0	larval seine
MAF07-139	NA	1	14.9	25 Jul 2007	27 Jun 2007	107.7	larval seine
MAF07-157	NA	1	17.5	27 Jul 2007	27 Jun 2007	74.9	larval seine
WHB07-078	NA	1	15.6	25 Jul 2007	27 Jun 2007	33.7	larval seine
TOTAL		9					

Over 542,000 fish have been collected between 1995 and 2007 under the larval Colorado pikeminnow survey. Of those, about 84% (N=453,350) were collected after 2001 when the sampling protocol switched from passive to active sampling (2002).

This work is being conducted as required by the 31 March 2000 San Juan River Basin Recovery Implementation Program Monitoring Plan and Protocol. The objectives of this specific monitoring effort are identified and listed below. Where applicable, these objectives are related to the specific tasks listed in the Long Range Plan set forth by the San Juan River Basin Recovery Implementation Program (SJRBRIP).

Project Modifications:

There have been numerous modifications to the 2002 active sampling methodology of the San Juan River larval Colorado pikeminnow survey as well as changes in reporting priorities, protocol, and format. The extent of the study area and aspects of the longitudinal sampling have been modified to improve spatial comparisons. The study area for this project includes from the middle of Reach 5 (Cudei, New Mexico) to the downstream-most end of Reach 1 (Clay Hills Crossing, Utah; total of 138.6 miles of critical habitat sampled). Beginning in 2003, the entire 139 river miles of the study area was sampled in single uninterrupted trips (10-12 field days per trip) rather than in two temporally discrete sections as done in 2002.

Conducting the Colorado pikeminnow survey under this new protocol (single-continuous pass) not only provided discreet reach information but also provided greater temporal resolution in respect to the longitudinal distribution of larval fishes and potential environmental cues required for spawning. Disadvantages to single-continuous pass effort were that the duration of the monthly sampling trips (10-12

field days) made them more subject to abiotic fluctuations (floods, flow spikes). Annually, at least one trip (an average) had to be cut short due to large flood events or low water events in the lower canyon.

The abbreviated trips were subsequently resumed once conditions improved (usually 1-2 weeks later). However, numerous problems were associated with cancelled or disrupted trips during the single-continuous pass effort. Designated camp-sites in the lower canyon (scheduled and coordinated by BLM) would have to be canceled and often new camp sites could not be provided. The large flood events not only disrupted the temporal resolution of the single-continuous pass effort but it also reduces sampling efficiency as many low velocity habitat are flooded by rising water levels thereby transporting larval and early juvenile fish in the drift.

Finally, additional costs were incurred because of the need to return to the field to complete the sampling effort for that month.

Besides the threat of weather related cancellation of single-continuous pass effort, concerns arose regarding fatigue and potential compromise to safety of crews on these extended, long-day (10-12 hours), and labor-intensive trip schedules. To mitigate these effects as well as gain even greater temporal resolution of the longitudinal distribution of Colorado pikeminnow larvae the protocol was changed to allow sampling of the upper (RM 142 – 53) and lower (RM 53 – 3) sections of the San Juan River simultaneously. This effort began in 2007 and utilized two fully equipped and discrete crews (comprised of 2 people per crew) with one crew sampling the upper and the other sampling the lower portion of the study area simultaneously. In 2008, addition participation of our staff with other SJRBRIP projects made this new simultaneous sampling effort a necessity so that our staff could meet obligations to assist the other researcher with their work.

Study Area:

The study will be the San Juan River between Cudei, New Mexico (RM 141.5) and the Clay Hills Crossing boat landing (RM 2.9) just above Lake Powell in Utah and will include making collections in reaches of the San Juan River under the jurisdiction of the National Park Service.

Objectives:

- 1) Determine the relative annual reproductive success of Colorado pikeminnow (Task 5.1.2.1)
- 2) Provide annual summaries of monitoring results
- 3) Provide detailed analysis of data collected to determine progress towards endangered species recovery annually and thence every five years (Task 5.1.1.2)
- 4) Provide comparative analysis of the reproductive success of the San Juan River fishes (Task 5.1.2.4)
- 5) Attempt to validate presumed spawning period of Colorado pikeminnow

Methods:*Field work:*

Sampling for Colorado pikeminnow larvae will be conducted in the San Juan River between RM 141.5 and RM 2.9 from early July through early September using sampling techniques that will provide sufficient numbers of fish necessary to meet study objectives. Access to the river will be gained through the use of inflatable rafts equipped with all of the gear and provisions needed for trips of up to ten days. The study area will be divided into an “upper” section (Cudei, NM, to Mexican Hat, UT) and a “lower” section (Mexican Hat, UT, to Clay Hills crossing, UT). Separate field crews will launch their rafts simultaneously in each of the two sections and proceed through their designated study area. The vehicle and raft trailer used by the field crew working in the upper section will be left at the Cudei launch site and subsequently be shuttled to Valles Trading Post in Mexican Hat, UT, where it will be placed in paid storage. The vehicle shuttle (with trailer) for the upper reach sampling effort has typically been performed gratis by personnel from the Farmington Office of the Bureau of Indian Affairs Office. Starting in 2008, this service will be performed by personnel from the N.M. Fishery Resources Office stationed in Farmington, NM. At the time of submittal of this Scope of Work, there was no charge being incurred for this service.

The sampling crew for the lower reach will launch from and store their vehicle and raft trailer at Valles Trading Post in Mexican Hat, UT, where a commercial shuttle (from Valles) will take the vehicle to Clay Hills crossing. The cost for this shuttle service is included under the travel and per diem section of our budget. Cold storage facilities are also available at Valles Trading Post and will be used on an as needed basis.

Because crews sampling the lower section of the study area will be in a high use recreational area, advance reservations for camp-sites will be required. All trips for 2009 must be scheduled by late January 2009 and submitted to the Bureau of Land Management (BLM) Office at Monticello, Utah. Designated camping permits for our lower reach sampling crews will be obtained and must be strictly adhered to in addition to other BLM-San Juan River recreation Area regulations (i.e., low impact and pack-out camping policies). Low flow conditions that are often prevalent during the study period make several sections of the river more difficult to navigate (especially in the lower reach). Our field crews are required to render assistance to boaters stuck in rapids or otherwise in distress and report all such encounters to the appropriate BLM personnel (rescues or rendering assistance may add one extra day to the lower reach sampling effort between June and August).

Sampling efforts for larval fish will be concentrated in low velocity habitats and employ small mesh seines (1.2 m x 1.2 m x 0.8 mm mesh) to collect fish. Retained specimens will be placed in Whirl-paks containing a solution of 10% buffered formalin and a tag inscribed with unique alphanumeric code that is also be recorded on the field data sheet. The lengths (to 0.1 m) of each seine haul and total number of hauls per sample site will be measures and recorded. Catch per unit effort for sample site will be reported as the number of fish per 100 m².

Native species large enough to be positively identified will be measured (standard length; SL) and returned to the river. Post-larval endangered fish species collected during this study will be photographed, a small portion of tissue from the fin clipped and retained in 95% EtOH (for purposes of assessing potential razorback sucker hybrids) and scanned with a FS2001 PIT tag reader for the presence of a PIT tag. Specimens of sufficient size but lacking a PIT tag will be injected with a tag. All PIT tag information will be recorded in the field data sheet and subsequently forwarded to the SJRB RIP for integration in the program's PIT tag database.

For each sampling locality, river mile will be determined to the nearest tenth of a mile using the San Juan River Basin Recovery Implementation Program 2003 Standardized Map Set. Universal Transverse Mercator (UTM) coordinates and zone will be determined with a Garmin Navigation Geographic Positioning System Instrument for each sampling locality. Mesohabitat type, length, maximum and minimum depths,

water clarity (determined with a Secchi disc), and substrata will be recorded for each sampling locality. Multi-parameter YSI units will be used to determine the following water quality parameters at each site sampled: pH, temperature, salinity, conductivity, specific conductance, and dissolved oxygen. A minimum of one digital photo will also be taken of each specific habitat sampled.

StowAway Tidbit temperature loggers will be set to record water temperatures hourly and deployed near Four Corners bridge, in McElmo Creek, and at Clay Hills crossing. The data from each temperature recorder will be downloaded monthly during the study period but remain in the river after completion of the annual study effort to record river water temperatures throughout the year.

Field Work, Safety:

Personnel participating in field work are required to successfully complete both an International Rescue Instructors Association (IRIA) level 2 swiftwater rescue class and American Red Cross CPR/AED training. Type III personal flotation devices (PFD's) will be worn by sampling personnel at all times while working. As PFD's lose flotation capacity due to UV exposure, compression of material, and oil and grit impregnation, and since each crewmembers PFD will be exposed to intensive use for at least 45 days per season, PFD's will be replaced annually. Simms Guideweight Gore-Tex waders and boots will be issued to all personnel along with 3 mm neoprene gloves (necessary in April and May). Due to cold water conditions, waders are a necessity during early season trips (April and May). As waders are worn throughout the day during these first two trips, they must be light-weight, flexible, and durable. In addition to personal camping gear and rain jackets, all personnel will be required to provide and use wide brimmed hats, sunscreen, and sunglasses (provided at no cost to the program).

Both rafts used for this project will carry an extensively stocked first aid kit replete with items necessary for most minor medical situation. Additionally, the first aid kit will contain a suite of items (i.e., splints, neck braces, butterfly stitches, snakebite kits) needed address more serious medical conditions. Because formalin is used in the preservation of specimens, several vials of eyewash solution will be incorporated into each first aid kit. First aid kits will be inventoried after each sampling trip and used and/or expired items replaced. In the upper reach of the study area, personal cell phones and PDA's will be used (at no cost to the program) to contact outside parties should a medical situation arise. In the lower study area reach (canyon bound; where cell phones do not have service) an Iridium 9505-satellite phone will be provided for sampling crews to be used in case of an emergency.

Formalin used for fish preservation will be transported in heavy-duty LPDE carboys. Extensive exposure to UV light makes the carboys susceptible to decomposition and cracking and requires that they be inspected monthly and not used for more than two years. Safety rope throw bags will be similarly inspected and retired from use accordingly. Rafts will be equipped with raft recovery (Z-line) kits, well stocked supply and repair kits, extra oar and oar blade, and two spare hand pumps to help ensure that crews do not become stranded due to raft damage.

Laboratory Work:

Samples will be returned to the lab immediately after each field trip is completed and processed following a multi-step procedure. To maintain the larval fish in good condition (necessary to ensure accurate identification) the samples must be transferred from Whirl-packs to glass jars and field fluids replaced with a solution of 5% buffered formalin. Cyprinid and catostomid larvae are extremely small and transparent especially at early developmental stages. To minimize the potential loss of fish in individual seine hauls, it is best to retain the entire contents of each seine haul. A negative result of this technique is that, in addition to larval fish, Whirl-pack samples usually contain considerable debris, detritus, and silt. Another important step in processing of individual samples is to separate fish from the detritus. This necessary portion of the process is labor intensive and can be quite tedious. During this process initial sorting of fish based on age class (age 0 [larvae] and age 1+) occurs. Samples that contain a large number of larval fish, especially those with protolarvae or mesolarvae often must be sorted twice.

After the fish are separated from the debris, personnel with San Juan River Basin larval fish identification expertise identify individual specimens to species. Bottom-lit stereomicroscopes equipped with polarized filters (that enhance the delineation of myomeres, pterygiophores, and fin rays) are used to assist with the identifications. Larval fish keys are referenced to assist in species specific determinations (e.g., Contributions to a guide to the cypriniform fish larvae of the Upper Colorado River System [Snyder 1981], Catostomid fish larvae and early juveniles of the Upper Colorado River basin, Morphological descriptions, comparisons, and computer interactive key [Snyder and Muth 2004], and Identifications of larval fishes of the Great Lakes Basin [Auer 1982]). Age-0 specimens are separated from age-1+ specimens using published literature on growth and body length at specific ages (Snyder 1981, Snyder and Muth 2004).

Age classes are enumerated, measured (minimum and maximum size [mm standard length] for each species at each site), transferred to 70% ethyl alcohol, and catalogued in the Division of Fishes of the Museum of Southwestern Biology (MSB) at the University of New Mexico (UNM). Both total length (TL) and SL of Colorado pikeminnow and razorback sucker are obtained using electronic calipers. The ontogenetic stage of Colorado pikeminnow and razorback sucker obtained in this study is determined based on the definitions provided by Snyder (1981).

Reporting and permitting:

Beginning in 2004, data from the two San Juan River larval fish surveys (razorback sucker and Colorado pikeminnow) were analyzed collectively and presented in a single report. This new format provided a more complete picture of the reproductive activities of the entire ichthyofaunal community in the San Juan River using the same criterion used as the other monitoring programs.

The results in our annual report pertain almost exclusively to age-0 fish (i.e., as fish > age-0 are not “larval fish” and are not the focus of this effort, they are not included in analysis). The number of fish >age-0 collected during the study is presented in an Appendix. Differences in mean CPUE are determined by species between years using a one-way Analysis of Variance (ANOVA). A Poisson distribution provided the best fit to the raw data. A variety of transformations (e.g., logarithmic, reciprocal, square root) were applied on the mean CPUE data for between year comparisons. A natural log transformation yielded the best variance-stabilizing qualities and produced a relatively normal distribution. Pair-wise comparisons between years (2003 – 2007) were made for each species and significance (i.e., $p < 0.05$) was determined using the Tukey-Kramer HSD test. Finally, a nonparametric ANOVA (Kruskal-Wallis test) was used on various data sets to compare results to the parametric analyses. While both ANOVA and Kruskal-Wallis were used to analyze data, data transforms enabled use of parametric analysis in all cases. The assumption of homogeneity of variances was assessed using the more conservative variance ratio criterion of <3:1 (Box, 1954), as opposed to <4:1 (Moore, 1995), among years. All species data sets met this more rigorous criterion and in most cases the variance ratio was <2:1 among years. Additionally, the significance values between parametric and nonparametric techniques were nearly identical and so only the parametric analysis will be presented.

Hatching dates for larval Colorado pikeminnow are calculated using the formula:

$-76.7105 + 17.4949(L) - 1.0555(L)^2 + 0.0221(L)^3$, where L=length (mm TL).

Spawning dates larval Colorado pikeminnow are then estimated by adding five days to the post-hatch ages to account for incubation time at 20 - 22°C (Nesler et al. 1988). Hatching dates of razorback sucker larvae were calculated by subtracting the average length of larvae at hatching (8.0 mm TL) from the total length at capture divided by 0.3 mm (Bestgen et al. 2002), which was the average daily growth rate of wild larvae observed by Muth et al. (1998).

This study is initiated prior to spring runoff and completed near the end of summer (late September). Daily mean discharge during the study period is acquired from U.S. Geological Survey Gauge (# 09379500) near Bluff, Utah. Water temperatures (mean, maximum, and minimum) are acquired from our temperature

loggers and additional data loggers (maintained by Keller-Bliesner Engineering) at the Colorado State Highway 160 bridge crossing (RM 119.2) and Mexican Hat, Utah (RM 53.3).

In addition to the annual report of the study provided to the SJRBRIP, reports summarizing fish collecting activities and specimens captured are also required annually under scientific collection permits provided by the New Mexico Department of Game and Fish, Navajo Nation, and state of Utah. The aforementioned reports include (at a minimum) site localities, GPS coordinates, and fish collected. An annual report of activities is a BLM (Monticello Field Office) requirement under our access permit to the San Juan River below San Island (Bluff, UT) and designated camps in the lower reaches of the river.

Meetings:

Researchers are required to attend a minimum of two meetings annually and report on annual monitoring projects. The two meetings (February and May) require researchers present PowerPoint presentations outlining the results and that years findings. Each meeting lasts about three days (which includes travel time).

Products:

A draft report of the 2009 larval razorback sucker sampling activities (combined with 2009 larval Colorado pikeminnow sampling activities) will be prepared and distributed to the San Juan River Basin Biology Committee for review by 31 March 2010. Upon receipt of written comments, that report will be finalized and disseminated to members of the San Juan River Basin Biology Committee by 1 June 2010. Electronic copies of the 2009 collection data will be transferred to the San Juan River database manager. Fish collected from this study will be curated in the Division of Fishes, Museum of Southwestern Biology (MSB), Department of Biology, at the University of New Mexico under a MSB contract with the SJRBRIP. Original field notes will be retained in the Division of Fishes and collection information electronically stored in a permanent MSB database program. These data and any maps generated from them will be available to the San Juan River Basin Biology Committee via hard-copy reports and electronically.

Project Title: 2009 San Juan River larval endangered fishes survey

Proposed Budget based on Six Sampling Trips

Personnel			
FIELD WORK			
UPPER REACH (RM 141.5 - 53.3)			
<i>Cudei Diversion to Mexican Hat</i>			
Research Associate (W.H. Brandenburg or M. A. Farrington)		60 staff days	\$21,000
<i>Field data collection – 10 days per trip x 6 trips</i>			
Field Assistant (C.C. McBride or A.L. Barkalow)		60 staff days	\$12,000
<i>Field data collection – 10 days per trip x 6 trips</i>			
LOWER REACH (RM 53.3 - 2.9)			
<i>Mexican Hat to Clay Hills</i>			
Research Associate (W.H. Brandenburg or M. A. Farrington)		60 staff days	\$21,000
<i>Field data collection – 10 days per trip x 6 trips</i>			
Field Assistant (C.C. McBride or A.L. Barkalow)		60 staff days	\$12,000
<i>Field data collection – 10 days per trip x 6 trips</i>			

LAB WORK		
UPPER AND LOWER REACH SAMPLES COMBINED (i.e., not fully differentiated under this task)		
Research Associate (W.H. Brandenburg and M. A. Farrington)	120 staff days	\$42,000
<i>Laboratory effort – 10 days per trip per each reach (x 2) x 6 trips</i>		
<i>TASKS: Laboratory identification, developmental staging, specialized endangered fish processing, data entry, data query and review, database development</i>		
Research Associate (C.C. McBride and A.L. Barkalow)	120 staff days	\$24,000
<i>Laboratory effort – 10 days per trip per each reach (x 2) x 6 trips</i>		
<i>TASKS: Post-trip sample processing, juvenile identification, post-identification – processing, measures, review of counts</i>		
OFFICE WORK (REPORT DEVELOPMENT)		
UPPER AND LOWER REACH SAMPLES COMBINED (i.e., not fully differentiated under this task)		
Research Associate (W.H. Brandenburg and M. A. Farrington)	80 staff days	\$28,000
<i>Office effort – 40 days staff member (= 40 days per discrete study)</i>		
<i>TASKS: Data analysis, draft report preparation, post-review redraft and submission, development and submission of formal responses to reviewer comments, development of presentation of study for annual meetings, annual reporting related to state and tribal permitting of sampling activities</i>		
Personnel (continued)		
PROJECT OVERSIGHT AND ADMINISTRATION		
Senior Research Associate (S.P. Platania or R.K.Dudley)	12 staff days	\$6,000
<i>Oversight effort – one day per month</i>		
<i>TASKS: Project coordination, project and data review, data management, report review, scope and budget preparation, project billing and accounting</i>		
Personnel (Field, Lab, Office, Oversight) Subtotal		\$166,000
SJRBRIP MEETINGS		
Two meetings per year required of researchers; 3 days/meeting		
Research Associates (W.H. Brandenburg and M. A. Farrington)	12 staff days	\$4,200
<i>2 meetings x 2 people x 3 days = 12 staff days</i>		
Senior Research Associate (S.P. Platania or R.K.Dudley)	6 staff days	\$3,000
<i>2 meetings x 3 days = 6 staff days</i>		
Personnel (Meetings) Subtotal		\$7,200
Personnel Total		\$173,200

Materials and Supplies**FIELD RELATED****Safety training and dedicated First Aid Gear**

<i>American Red Cross CPR/AED training x 4</i>	\$75/person	\$300
<i>IRIA Level 2 swiftwater rescue class x 2</i>	\$325/person	\$650
<i>Type III Personal Floatation Devices (PFD) x 4</i>	\$115/person	\$460
<i>Fire Extinguisher annual recharge x 2</i>	\$25/unit	\$50
<i>First Aid Kit item update and replacement x 2</i>	\$50/unit	\$100
<i>Light-weight Gore-Tex waders x 2</i>	\$325/person	\$650
<i>Iridium 9505A Satellite Phone (five year depreciation)</i>	\$204/yr	\$204
<i>Satellite phone monthly service x 6</i>	\$35/mo	\$210
<i>Cell phones x 2</i>	not charged	\$0

Safety training and dedicated first aid gear Subtotal		\$1,200 \$2,624
--	--	-----------------------------------

Raft and rafting associated gear

<i>NRS Raft Supplies (average of \$500/raft/year 2007 – 2008)</i>	\$500/yr	\$1,000
<i>AIRE 156R sealed floor pocket (self-bailing) x 1</i>	\$305/yr	\$350
<i>ThorShield 2250 Tarpaulin x 2 (five year depreciation)</i>	\$125/yr	\$250
<i>Sherwin Williams Tile Clad Epoxy Paint (two gallons)</i>	\$110/yr	\$110
<i>Trailer (for raft) maintenance x 2</i>	\$250/unit/yr	\$500
<i>Raft depreciation x 1</i>	not charged	\$0
<i>Trailer depreciation x 2</i>	not charged	\$0
<i>Pit tag reader FS2001 with pass through wand x 1 (\$3,100)</i>	Program ?	

Raft and rafting associated gear Subtotal		\$2,210
--	--	----------------

Fish Sampling Gear

<i>Larval seines x 4 per year</i>	\$75/seine	\$300
<i>Preservation materials (carboys, fluid, tags, whirl paks)</i>	\$350/crew	\$700
<i>Open reel coated fiberglass measuring tape (metric) x 2</i>	\$50/tape	\$100
<i>Thermometers x 12</i>	\$12/piece	\$144

Fish sampling gear Subtotal		\$1,244
------------------------------------	--	----------------

Water Quality and Electronic Sampling Gear

<i>GPS Unit (replacement average one per year)</i>	\$125/unit	\$125
<i>Digital camera (use, memory card, depreciation) x 2</i>	\$50/unit	\$100
<i>YSI Water Quality (calibration solutions, membranes) x 2</i>	\$90/piece	\$180
<i>Water temperature data logger x 4</i>	\$125/piece	\$500

Water quality and electronic sampling gear Subtotal		\$905
--	--	--------------

OFFICE RELATED

<i>Computer, scanner, and printer use and supplies, software upgrades, electronic storage media, presentation software</i>	project cost	\$1,000
--	--------------	---------

Office materials and supplies Subtotal		\$1,000
---	--	----------------

Materials and Supplies Total		\$6,559 \$7,983
-------------------------------------	--	-----------------------------------

Travel and Per Diem**FIELD WORK**

UPPER REACH (RM 141.5 - 53.3) <i>Cudei Diversion to Mexican Hat</i>		
Travel - 4 x 4 pick up truck and raft trailer <i>575 miles round-trip per trip x 6 trips = 3,450 miles</i>	\$.58 \$1.00/mi	\$2001 \$3,450
Per Diem - 6 field days; 0 hotel day <i>6 days x 2 people x 6 trips = 72 field per diem days</i>	\$45/day	\$3,240
Truck and Trailer Shuttle from Cudei to Mexican Hat <i>Shuttle service provided gratis by USFWS – NMFRO</i>	\$0/shuttle	\$0
Truck and Trailer Storage at Valles Trading Post <i>Daily storage rate \$5/vehicle x 4 days x 6 trips = 24 days</i>	\$5/day	\$120
LOWER REACH (RM 53.3 - 2.9) Mexican Hat to Clay Hills		
Travel - 4 x 4 pick up truck and raft trailer <i>775 miles round-trip per trip x 6 trips = 4,6750 miles</i>	\$.58 \$1.00/mi	\$2697 \$4,670
Per Diem - 4 field days; 1 hotel day (combined total from below) <i>4 days x 2 people x 6 trips = 48 field per diem days</i>	\$45/day	\$2,160
<i>1 day x 2 people x 6 trips = 12 hotel per diem days</i>	\$95/day	\$1,140
Truck and Trailer Shuttle from Mexican Hat to Clay Hills <i>Valles Trading Post at \$325/shuttle x 6 trips</i>	\$325/shuttle	\$1,950
Travel and Per Diem (Field) Subtotal		\$13,308\$16,730
SJRBRIP MEETINGS		
Travel (everybody in one vehicle) <i>425 miles round-trip per trip x 2 trips = 850 miles</i>	\$.58 \$1.00/mi	\$493 \$850
Per Diem - hotel days <i>3 days x 2 trips x 3 people = 18 hotel per diem days</i>	\$95/day	\$1,710
Travel and Per Diem (Meeting) Subtotal		\$2,203\$2,560
Travel and Per Diem Total		\$15,51119,290
Personnel Total	\$	173,200
Materials and Supplies Total	\$	6,5597,983
Travel and Per Diem Total	\$	15,52119,290
Project Subtotal	\$	195,270200,473
IDC (15%)	\$	29,29130,071
GRAND TOTAL	\$	224,561230,544

Note: Budget adjusted to \$224,561 to remove overage in allowed mileage expenses and in *Safety training and dedicated First Aid Gear.*

San Juan River Specimen Curation Fiscal Year 2009 Project Proposal

Principal Investigators: Alexandra M. Snyder and Thomas F. Turner
Division of Fishes - Museum of Southwestern Biology
University of New Mexico
Albuquerque, NM 87131
(505) 277-6005
amsnyder@unm.edu
turnert@unm.edu

Background

Personnel from the Division of Fishes, Museum of Southwestern Biology (MSB), at the University of New Mexico are responsible for two inter-related programs on the San Juan River. First, the MSB fish division is the repository for specimens collected and retained by researchers with the San Juan River Recovery Implementation Program (SJR-RIP). Fish specimens taken under these programs are initially sorted by the principal investigators, held until they have submitted their yearly-progress reports, and then specimens are received by MSB personnel. The collections are accessioned, specimens transferred from formalin to alcohol, identifications verified, individuals enumerated, length ranges recorded (largest and smallest specimens in a cataloged "lot" or single collection), collection data verified, incorporated into a database (MS Access 2007) and transferred to permanent labels (DataMax ® I-Class™ thermal transfer printer and 5 mil polyester paper). This protocol is standard for all major collections and museums of natural history (i.e., Smithsonian Institution, University of Michigan Museum of Zoology, etc.). Second, it is standard practice that, prior to complete incorporation into the MSB collections and database, all specimens received from the SJR-RIP be examined by qualified personnel (in that particular field of study) in an effort to verify the original identification and collection information. This system provides a final check (safeguard mechanism) to minimize the likelihood of misidentification of San Juan River fish species with particular attention on Colorado pikeminnow and razorback sucker specimens. Any changes in species identifications are noted and reported to the SJR-RIP principal investigator along with the entire data set (i.e., listing of collection locality data, collectors, date, original field number, species, number of specimens, length ranges, and museum catalog number).

The number of fish specimens processed by the MSB Division of Fishes under the San Juan River Basin Recovery Program can fluctuate, greatly, between years. One reason for this that samples sent to MSB by non-MSB researchers are often not fully processed until almost one year following the collection of specimens. This lag period, between date of collection and complete MSB processing, is necessary as individual SRJ-RIP researchers must perform preliminary sorting and require the specimens for preparation of their reports. Other factors such as annual variability of sampling conditions and initiation of new or completion of old projects has resulted in marked changes (increase/decrease) in the number of samples and specimens. (An example of this occurred between 2001 and 2002 when drift sampling for larval Colorado pikeminnow was eliminated in favor of seine sampling.)

Discussion of this issue with the San Juan River Biology Committee resulted in the recommendation that the annual budget for the San Juan River Specimen Curation and Larval Fish Identification reflect an "average" year of sample processing. Almost all MSB-San Juan River Basin archived specimens are the result of collections made under the auspices of the San Juan River Basin Recovery Implementation Program Monitoring Plan and Protocol. The Biology Committee recognized that some years would require more effort from MSB than budgeted while other years might not require the same high level of activity. A relatively stable budget has allowed for uninterrupted processing of incoming samples. This budget has also allowed for the processing of backlogged SJR-RIP collections due to years of exceptionally high fish capture or during the early years (1992-1995) when sufficient numbers of MSB staff were not available to process and database collections. As of 2007, over 1,000,000 SJR-RIP specimens (along with associated locality and ecological data) have been incorporated with the MSB Division of Fishes archives and are available to researchers.

Study Area

This project does not involve the collection of specimens but instead the processing, curation, and data capture for all collections taken by different research components of the San Juan River Recovery Implementation Program. The annual sampling area for these researchers is the San Juan River between the outfall of Navajo Reservoir (RM 223.0) and the Clay Hills boat landing (RM 2.9) just above Lake Powell in Utah.

Objectives

1. Provide a permanent repository for San Juan River fish collections, field notes, and associated data
2. Verify species identifications, enumerate specimens, and report to principal investigators
3. Maintain a GIS reference database for current material
4. Assist principal investigators with secondary collection sorting and identifications as time and resources permit

Methods

The primary task to be completed under this project is the processing and curation of fish specimens generated by research projects executed under the auspices of the San Juan River Basin Recovery Implementation Program. Samples are transferred to the MSB Division of Fishes, by the principal investigator of a project, once that individual has completed their work and prepared the necessary reports. (This usually infers a lag-time of one year between date of collection of specimens and the transfer to the MSB Division of Fishes.) Upon receipt, field collections are removed from field jars or sample bags and the "field formalin," transferred into known (lab mixed) solutions of buffered 5% -10% formalin, if this has not already been done by SJR-RIP principle investigators. The collections are organized on shelves by field number and checked against the appropriate field data sheets. Once collections have been sufficiently fixed in formalin, the fish larvae and adults are transferred from formalin into ethanol. (Some examples of the larval fishes will be maintained in buffered 5% formalin to conserve certain morphological features.) Specimens are sorted by species (if this has not been done) into distinct lots or their identifications verified by MSB staff (curatorial and field); the number of individuals in these species lots are counted and the size range measured in millimeter standard length. Original field sheets, received from SJR-RIP principle investigators are digitally captured (scanned and tif images "cleaned") and the pdf files linked to the MSB fish collections MS Access database. Locality data are entered into a MS Access 2007 database table (Nbottom) and the species lots entered into a separate table (Ntop). The electronic catalog assigns MSB numbers to each lot, deposition information (jar, vial, tank, genetic), and produces chemical/preservative resistant jar labels using a thermal transfer printer and polyester labeling paper. Once the collections are labeled, they are filed by taxa and drainage, in the permanent archives of the MSB fish division, which is a secure room that is controlled for temperature (18°C) and light (UV).

In addition to the aforementioned responsibilities, the Division of Fishes is available and has frequently assisted principal investigators by taking on the added responsibility of processing (a limited number) of their unsorted collections (without requesting additional funding). Specimens are sorted, identified, counted, measured, catalogued, and data submitted to the principal investigator for inclusion in reports. In cases where the amount of backlogged material in the possession of the principal investigator was beyond our capabilities, supplemental funds have been sought so that additional personnel can be hired (under the supervision of the permanent staff) to process the excess material.

Products

Fishes collected under the auspices of the San Juan River Basin Recovery Implementation Program (SJR-RIP) will be curated and maintained, in perpetuity, in the Division of Fishes, Museum of Southwestern Biology (MSB), Department of Biology, at the University of New Mexico, Albuquerque NM.

A more detailed report on the 2009 San Juan River specimen curation and larval fish identification activities at the MSB Division of Fishes, University of New Mexico will be prepared in draft form and distributed by 31 March 2010 to the San Juan River Biology Committee for review. Upon receipt of written comments, this report will be finalized and disseminated to members of the San Juan River Biology Committee by 1 June 2010.

All original field notes, data sheets, and photographs will be retained in both the MSB Division of Fishes document archives and digital collections, these data to be permanently stored in a secure (backed up to an external hard drive)

MS Access 2007 database program managed by the MSB-Fishes Collections Manager and Graduate Student Research Assistant.

Electronic copies of the field and collection data will be made available to data managers with the various projects under the San Juan River Basin Recovery Implementation Program once the specimen collections and their data have been successfully incorporated into the MSB Division of Fishes.

Budget Fiscal Year 2009

Personnel:			
	Graduate student RA (Data manager and GIS)	\$	7500.00
	Graduate student health benefit	\$	679.00
	RA Fringe benefits 1%	\$	75.00
	RA Tuition 1 semester	\$	2487.00
	RA GPA fee 1 semester	\$	25.00
	Undergraduate student Curatorial Assistants (2)	\$	7920.00
	Undergraduate student fringe benefits 1%	\$	80.00
Section Subtotal		\$	18,766.00
Equipment and Supplies:			
	95% ethanol preservative	\$	3000.00
	Specimen jars, Buna-N gaskets, and polypropylene caps	\$	2000.00
	Permanent specimen labels-5 mil polyester paper	\$	2000.00
	DataMax ® Printer maintenance and calibration costs	\$	423.00
Section Subtotal		\$	7423.00
Total (Direct Costs)		\$	26,189.00
Administrative Overhead (15%)		\$	3928.00
Grand Total		\$	30,117.00

Out-year funding (based on 5% increases):

Fiscal Year 2010

\$31,623.00

San Juan River Basin Recovery Implementation Program Geomorphology and Habitat 2009 Scope of Work

Technical Approach

Element 1. Water Temperature

Background

Eight temperature recorders have been in place since summer of 1992 at the locations shown in Table 1. From 1992-1999, OMNIDATA DP-230 data pod loggers sampled water temperature every 10 minutes and stored maximum, minimum and mean temperature for each day. Optic StowAway temperature loggers from Onset Corporation were utilized from 1999-2006. In 2006, these recorders were replaced with Onset Corporation HOBO Water Temp Pro loggers. They record water temperature every 15-minutes. Table 1 also shows the periods of record at each site. The missing data were caused by equipment problems or vandalism.

Proposed Methods

Data Collection

Onset Corporation HOBO Water Temp Pro loggers with built-in thermocouple temperature sensors will be installed in the locations described in Table 1. Loggers will be installed in existing enclosures that have been used over the past 15 years. Where enclosures are deteriorated, missing or badly placed, they will be upgraded as necessary to provide protection to the equipment.

The recorders will be inspected and read twice each year, once in the spring and once in the fall. Battery condition will be monitored and loggers changed out when the battery life falls below that required to continue until the next reading point.

Following each download, data will be quality checked and bad data removed. As noted in the background, vandalism, natural causes or equipment malfunction can cause loss of data that are beyond our control. Every attempt will be made to assure quality data within the scope described, but some missing data will be inevitable. Data integrity similar to that in the existing database will be provided.

Data Storage

The records will be maintained in a Microsoft Access database. The main data table will store the 15-minute data and will be constructed as shown in Table 2. Data tables summarizing daily maximum, minimum and average temperatures will be generated for each of the eight sites by query of the main data table and stored in the database in the format shown in Table 3. Table 4 shows the information stored to describe each session, including geo-spatial data to allow importation into a geographic Information System.

Data Analysis and Reporting

After the fall logger download, data for the water year will be compiled and the daily average temperatures plotted along with the daily hydrograph of the San Juan River at Four Corners, New Mexico. A summary report will be prepared that will include presentation of the daily average temperature data with a discussion of data collection procedures, data quality and repair requirements during the season. Anomalous data, if any, will also be discussed.

Deliverables

- An annual draft report prepared and submitted by March 31, 2010
- A final report submitted by June 1, 2010.
- An updated temperature database with all data collected to date, updated through September 2009 by June 1, 2010.
- Attendance at the annual report meeting and one additional Biology Committee meeting

Table 1. Water temperature monitoring locations and period of record

Location	RM	Period of Record
<i>Active Temperature Recording Sites</i>		
Near Navajo Dam	225.0	7/9/1999 to 9/15/06
Archuleta - San Juan at USGS Gage Location	218.6	7/23/92 to 9/15/06
Farmington - San Juan at USGS Gage Location	180.1	8/5/92 to 1/16/96, 7/8/99 to 11/4/01, 10/3/02 to 9/15/06
Shiprock - San Juan at USGS Gage Location	148.0	7/8/99 to 9/16/06
Four Corners - San Juan at USGS Gage Location	119.4	10/7/94 to 3/11/96*, 7/9/99 to 10/19/06
Montezuma Creek - San Juan at Montezuma Creek Bridge	93.6	8/9/92 to 1/11/93, 2/25 to 3/14/93, 4/14 to 5/10/93, 5/28/93 to 3/11/05, (sensor stolen. Replaced 10/31/05) 10/31/05 to 9/16/06
Mexican Hat - San Juan near Bluff Gage Location	52.1	7/9/99 to 3/27/02 , 9/18/02 to 8/1/06
Farmington - Animas at USGS Gage Location	n/a	8/5/92 to 4/14/97, 5/7/97 to 8/26/97, 10/15/97 to 6/4/98, 7/8/99 to 9/15/06

Table 2. Temperature database main table format

Temp			
ID	RecDate	RecTime	DegC
4C	7/9/1999	4:04:27 PM	23.48
4C	7/9/1999	4:49:27 PM	23.74

Table 3. Daily temperature summary table format

AnimasFarminton				
ID	RecDate	Tmax	Tmin	Tavg
AF	7/8/1999	22.11	18.36	19.2225580437239
AF	7/11/1999	20.13	15.81	17.9729165037473

Table 4. Temperature station description database table

StationID					
ID	Location	Notes	Lat	Lon	Datum
4C	Four Corners	Located at the Four Corners USGS gage	37.00195	-109.0311	NAD83
AF	Animas at Farmington	Located an the Animas at Farmington USGS gage	36.72154	-108.2017	NAD83
AR	Archuleta	Located at the Archuleta USGS gage	36.80278	-107.699	NAD83
FM	Farmington	Located at the Farmington USGS gage	36.72221	-108.2251	NAD83
MC	Montezuma Creek	Located left bank at sheet piling upstream side of the Mont. Ck bridge	37.2579	-109.3096	NAD83
MH	Mexican Hat	Located right bank near the USGS mini-monitor enclosure upstream of Mex Hat bridge	37.15059	-109.8669	NAD83
ND	Navajo Dam	Base of Navajo Dam on river left immediately downstream of outlet	36.80484	-107.6148	NAD83
SR	Shiprock	Located at the Shiprock USGS gage	36.781	-108.6899	NAD83

Element 2. Habitat Mapping

Background

Habitat mapping completed during the period 1992 - 1997 has been used to develop flow/habitat relationships used in the flow recommendation process (Holden, 1999). Annual habitat mapping was included in the San Juan River Basin Recovery Implementation Program (SJRIP) long-term monitoring program in 1998 and has continued through 2006. Results from 1998 through 2002 were used to evaluate the flow recommendations, resulting in a recommendation to examine modification of operating rules to focus more heavily on the high flow portion of the recommendations (Miller, 2005).

Proposed Methods

Base Photography Preparation

USBR will acquire digital videography of the San Juan River from the San Juan River at the Animas River confluence (RM 180) downstream to below Clay Hills Crossing (RM 0). We will coordinate with USBR to acquire the videography at a flow of 500 to 1,000 cfs in late July. Digital single frames will be captured from this videography to provide full coverage of the river with about 20% overlap. River mile marks will be placed on these frames using the SJRIP standard river mile GIS coverage. These digital images will be archived to DVD for later interpretation, if needed.

Base photography maps will be prepared at a scale of approximately 1 inch = 75 feet for the two detailed study reaches discussed in Element 3. The digital frames captured for these detailed reaches will be rectified to 2005 digital orthographic quads (DOQ's) prior to printing the field maps. Field maps will be printed on 11" x 17" pages and included in sheet protectors for field mapping.

Field Mapping

The 27 aquatic habitat types and 5 associated terrestrial types listed in Table 5 will be delineated on the base photographs (1 inch = 75 ft scale) by visual inspection. Mapping will be completed in the detailed reaches at the same time as the fish surveys are completed. Each polygon delineated will be marked with its corresponding code, as shown in Table 5. The date of mapping will be recorded on the beginning map sheet for each day's mapping along with the name of the mapper.

All mappers will be experienced in mapping aquatic habitat in the San Juan River. In as much as the mapping process is interpretive, annual reviews will be conducted among the mapping crews prior to mapping to assure the best possible reproducibility in interpretation among mappers.

Following field mapping, the field sheets will be reviewed and missing codes or non-closed polygons corrected prior to processing.

Data Processing

Once the field mapping sheets are reviewed and edited, they will be scanned at a resolution of 300 dpi and then rectified to 2005 DOQ's to remove distortion. After rectification, the habitat polygons will be digitized and coded in ArcView to produce a shape file and database with habitat perimeter and area by type and river mile. The data will then be extracted and summarized by count and area per river mile for analysis. Average flow at mapping for each detailed reach will also be extracted from USGS gage data, using the gage or gages most representative of the reach.

Table 5. Eight general categories of habitat types on the San Juan River

Low Velocity Types	Run Types	Riffle Types	Back-Water Types	Shoal Types	Slack-Water Types	Vegetation Associated Habitat Types	Associated Terrestrial Types
pool	shoal/run	riffle	back water	sand shoal	slack water	overhanging vegetation	cobble bars
debris pool	run	shore riffle	back water pool	cobble shoal	pocket water	Inundated vegetation	sand bars
rootwad pool	scour run	riffle chute	embayment				islands
eddy	shore run	shoal/riffle					boulders
edge pool	undercut run	chute					rootwad piles
riffle eddy	run/riffle	rapid					

Data Analysis

No data analysis is included since river-wide mapping is not included. Detailed reach analysis is included in Element 4.

Schedule

Base photography will be acquired in late July 2009 (flow permitting) and the base maps prepared in early August 2009. Files will be archived in September 2009.

Deliverables

The following deliverables are associated with this task:

- Archived Digital image files
- Digital and print copies of the detailed reach base photography

Element 3. Geomorphology

Background

From 1998 through 2004, long-term monitoring of channel morphology continued semi-annual (pre- and post-runoff) measurement of channel cross-sections first established in 1992. As a result of the 2003 analysis of channel response, the Biology Committee and Peer Review Panel recommended going to a 5-year measurement interval for these cross-sections, with the next survey in 2009, using the protocol outlined in the Long Term Monitoring Program.

Proposed Methods

The 14 cross-sections identified in 1999 as part of the long term monitoring plan will be surveyed post-runoff every 5 years (next survey in 2009) for analysis of long-term change and compared to previous surveys to determine trends. The survey will be completed

using a marked cable and previously established bench marks across the active channel. Survey intervals shall be not greater than 10 ft. A self-leveling optical level, total station or RTK sub-centimeter accuracy GPS may be used for this survey. Flood plain that has had active flow since the last survey will be surveyed with GPS or total station equipment.

Analysis of the change in cross-section geometry and substrate in relation to hydrographic conditions will be completed to monitor response of the system to flow recommendations. Only long-term response to cumulative conditions are possible since surveys are only completed every 5 years.

Deliverables

The following will be provided for this element:

- Cross-section survey plots in AutoCad of each cross-section compared to previous years
- Report of findings at annual researcher's meeting
- Draft report, March 31, 2009
- Final report, June 1, 2009
- Attendance at the annual report meeting and one additional Biology Committee meeting

Element 4. Habitat Use of Young-of-Year Endangered Fish

Background

During integration of San Juan River Basin Recovery Implementation Program (SJRIP) monitoring data from 1999-2003, it became obvious that integration of habitat data and fish data was extremely difficult (Miller 2005) since these two data sets were taken at different levels of detail. Adult fish monitoring data were too coarse to allow correlation with habitat data while habitat mapping units were too large to see details that were often the focus of sampling by larval and juvenile fish sampling programs. While larval and small-bodied fish sampling collect habitat data, the habitat categories do not match those in the habitat mapping program. Finally, although GPS locations are provided for recently collected larval and small-bodied fish sampling programs, the accuracy is not sufficient to place them on the habitat maps with sufficient precision to correlate the two data sets.

Backwater habitat has been theorized as important to larval and young juvenile endangered fishes. Backwater habitat is low in abundance in the San Juan River and has declined substantially since 1995 (Bliesner and Lamarra, 2006). However, sampling for age 0 and age 1 Colorado pikeminnow in the last several years has indicated that they use other low velocity habitat that is not necessarily mapped by the standard mapping program (Golden, et al, 2006).

To identify the habitat utilized by young endangered fishes and to provide information to allow this habitat to be mapped more broadly in the river, the following objectives as stated in the RFP will be addressed in Element 4:

1. Sample for young-of-year Colorado pikeminnow and razorback sucker within the two complex reaches to determine habitat use of endangered fish.

2. Map habitat in each complex reach each time fish sampling occurs.
3. Use supplemental data on young Colorado pikeminnow and razorback sucker captures of any size class throughout the San Juan River from other SJRIP sampling efforts and use these data to add to the habitat use information in the complex reaches.

The mapping, combined with detailed channel topology measurements and hydraulic modeling, would also provide insight into the mechanism or process for creation and maintenance of these complex reaches and provide a better understanding of the loss or creation of backwater habitats or other low velocity habitats used by the endangered fishes.

Proposed Methods

Detailed Reach Fish Survey and Habitat Mapping

The two complex reaches will be sampled once in March 2009 prior to spring runoff and once in August 2009 during base flow for young endangered fishes. Each complex reach will be thoroughly sampled twice during each sampling period for a total of three days per reach to increase sample density in all habitat types and improve the likelihood for sampling the rare fish. The second sample for each sampling period will be conducted at each site with at least one day of "rest" to minimize impacts from sampling activities. The two samples will allow for a more thorough sampling of the complex reach at essentially one time/flow period, increasing the chance that young endangered fish will be caught. An additional complex reach may be sampled if flow and sampling conditions allow, which will increase the potential to capture young endangered fish. The following procedure will be followed:

- Habitat will be mapped for a reach at the detailed level described Element 2 above.
- Each of the previously defined habitat types (Bliesner and Lamarra, 2006) mapped will be identified for abundance, and habitats to be sampled will be randomly selected in general relation to their abundance before each of the two sampling runs, except for runs and riffles which dominate the San Juan River. Runs and riffles will be sampled but to a lesser degree than they are represented in the reach since they generally are over 90 percent of the habitat in the San Juan River. All backwaters, shoals, and other low-velocity habitats, habitats known to be preferred by young Colorado pikeminnow (Golden et al. 2006) will be sampled. A standard seining technique using a 5 m x 2 m x 3 mm double-weighted seine will be used in March as the target Colorado pikeminnow will be of young-of-the-year size. A block seining technique using a 9 m x 2 m x 6 mm double-weighted seine at the bottom of a sample and another similar seine to work down to the bottom seine will be used in the July-August monitoring trip as the fish will be larger, of the age-1 size.
- The approximate area sampled in each location will be drawn on the base map overlaying the habitat mapping as the sample is taken.
- During fish sampling, additional habitats will be mapped that correspond to areas sampled for fish that were too small to include in the original mapping. Habitat descriptions will also be refined if needed to best describe sampling sites based on the 2007 sampling results.

- Following sampling, depth, substrate, velocity and cover will be measured in the sampled area. Depths and velocities will be taken at three locations with both depth and velocity measured at the same location to represent the average conditions in the sampled area. Care will be taken to target only one habitat type during individual seine hauls. If the sampled area crosses several habitats, at least one set of depths and velocities will be taken in each habitat type. Velocity vectors will be sketched on the habitat map at each location.

Information collected at each seining location will include: river mile location, GPS location (UTM), habitat type, cover, seine type, water temperature, area sampled (length and width), depth and velocity at three or more locations, cover, and substrate type. All fish collected, except for larvae, will be identified to species and counted. Up to 50 randomly selected individuals of each species will be measured at each complex reach except for Colorado pikeminnow and razorback sucker, which will all be measured. This will provide information on the general size and age of the fishes that are collected at each reach during each sampling trip and the habitat use of these fishes. Native fishes will be returned to the habitat alive, and nonnative fishes will be retained. A separate data sheet will be used for each seine location. Multiple seine haul runs may be made in large (>100 m²) habitats.

Colorado pikeminnow and razorback sucker captured during the first pass through each reach will be VIE tagged if they are not PIT tagged. This will allow for a population estimate to be made if any are recaptured on the second pass through the reach. A PIT tag reader will be taken on sampling trips. All Colorado pikeminnow and razorback sucker over 150 mm TL captured during sampling will be scanned for PIT tags and tagged if they do not already have a PIT tag. If PIT tags are present, the number will be recorded.

Obtain Other Endangered Fish Capture Data.

It is anticipated that data on Colorado pikeminnow and razorback sucker habitat use will also be collected by other SJRIP studies within the complex reaches, as well as outside those reaches. In 2007, one trip was scheduled with the larval fish study and the small-bodied monitoring program to standardize nomenclature for habitat mapping for improved data integration. Data on larval fishes will be obtained during the annual April-June razorback sucker larval fish and July-September larval Colorado pikeminnow surveys currently being conducted under the SJRIP (Brandenburg and Farrington 2006). Additionally, the fall standardized small bodied fish sampling (Paroz et al. 2006) will also provide information on fish captures by habitat type in the complex reaches and at other locations. Adult monitoring (Ryden 2006), and nonnative removal studies (Davis 2006, Jackson 2006) will begin to obtain GPS points for larger endangered fish captures in 2007, which may provide a means to correlate pikeminnow and razorback habitat use at the time of capture to larger scale mapped habitats, although at a gross scale. With sampling efforts occurring throughout the year, these data can provide greater seasonal habitat use information. Capture data and the associated habitat information will be gathered from the respective researchers for all sections of the river and any captures that occurred within the complex reaches will be analyzed with the capture data from the field efforts from this study. Additional captures and the corresponding habitat information outside the complex reaches will be evaluated, but habitat mapping for these areas will be coarse and will not provide the same scale for detailed analysis as gathered by this study in the complex reaches. The results will be used to verify the more detailed habitat use information

collected within the detailed reaches as the level of detail of this opportunistically gathered data allows.

Analyze Fish Capture/Habitat Utilization Data.

Habitat use information for endangered fish will be developed using all capture data from this complex reach study, as well as information from other studies with reliable habitat information. Data from the second pass will be tested to determine influence of the first sampling pass on the second pass results. Any influence found will be considered in the analysis of the full data set. Individual sample habitats noted during the fisheries sampling efforts will be compared with the habitat availability developed from the habitat mapping portion of the study during each sample period. It is intended that the proportional use of individual habitat types will be evaluated relative to all habitats sampled using a chi-square goodness-of-fit test (Muhlfeld et al. 2001, Golden et al. 2006) to show habitat preference. Other statistical methods may be employed as the data warrant. The habitat use/preference information developed through this analysis will be used to determine the habitats important to the fish and those to be looked at closely in modeling.

Information on fish gathered by other studies noted in #4 above may not fit all the requirements (all habitat types sampled, randomness, available habitat mapping) of the chi-square goodness-of-fit analysis. These data, along with capture data from outside of the complex reaches, will be summarized and compared to the habitat use information from this study. This may provide information on habitats not easily sampled with seines, potentially expanding the habitat use information.

In addition, the basic fishery information from the complex reach sampling will be collated and summarized to show effort, fish captured, CPUE, endangered fish size and other pertinent information. These data will be presented for all fish species and the summary of this information included in the annual report.

In addition to analysis of habitat use by the endangered fishes, the habitat usage by each member of the fish assemblage will be analyzed (when sufficient data exist) and compared to usage by the endangered fish. By using the full assemblage, statistical power is added to the relationship analysis, particularly in the event that endangered fish captures are small in number in the two reaches.

If the number of fish captures is too small to provide statistically significant results, recommendations will be made to modify the sampling plan for any future studies to obtain adequate numbers of young Colorado pikeminnow.

Deliverables

The following will be provided for this element:

- Draft report, March 31, 2010
- Final report, June 1, 2010
- GIS shape files and database files of habitat mapping and fish capture data
- Attendance at the annual report meeting and one additional Biology Committee meeting

**SJRIP PIT TAGS
2009 Project Proposal**

Mark McKinstry UC-735
Bureau of Reclamation
125 South State Street, Room 6107
Salt Lake City, UT 84138-1147
Phone 801-524-3835
FAX 801-524-5499
mmckinstry@uc.usbr.gov

BACKGROUND:

PIT tags are used to individually mark fish for use in movement studies and for mark-recapture estimates in the San Juan River Basin. PIT tags are not specific to any particular project, but are used by several different projects. PIT tags and readers purchased for the SJRIP will be combined with the purchase made for the UCRIP to save money by purchasing larger quantities and save expenses associated with administering the contract. All PIT tags and readers will be shipped to USFWS in Grand Junction C/O Chuck McAda at:

U.S. Fish and Wildlife Service
Colorado River Fishery Project
764 Horizon Drive, Building B
Grand Junction, Colorado 81506-3946
Phone: 970-245-9319 (19)

TASKS – 2009

1. Purchase PIT tags and readers and distribute to end-users

In FY2009, \$45,000 is allocated in the workplan to purchase 15,000 PIT tags. The purchase of PIT tags and readers is done under a fully competed contract with BioMark in Boise, ID.

FY 2009 BUDGET

Funding source	Amount carried forward from FY08	Projected expenditure in FY09
FY2009 Annual funding		\$45,000
Total	\$0.0	\$45,000

Projected funding:

FY-2010 \$50,000.00

FY-2011 \$50,000.00

FY-2012 \$50,000.00

**Program Coordinator's Office
Fiscal Year 2009 Draft Proposal**

U.S. Fish and Wildlife Service
2105 Osuna NE Albuquerque, New Mexico 87113
David_Campbell@fws.gov (505) 761-4745
sharon_whitmore@fws.gov (505) 761-4753

Background

The San Juan River Recovery Implementation Program (Program) is designed to simultaneously address endangered fish species recovery and development of water resources within the Basin. The Program includes representatives from not only Federal agencies, but also the States of Colorado and New Mexico, the Jicarilla Apache Nation, the Southern Ute Indian Tribe, the Ute Mountain Ute Tribe, the Navajo Nation and the water development interests, most of which have legal mandated responsibilities to the endangered fish and/or the water resources.

The Service is responsible for directing and coordinating the overall Program. As stated in the Program Document, the Service will appoint a Program Coordinator who will be responsible for overall Program coordination, a senior level biologist and a Program Assistant to dissemination of information about Program activities. Public Law 106-392 specifically authorizes the use of base funding to fund program management.

Tasks

1. Coordinate the activities of the Biology, Hydrology and Coordination Committees.
2. Insure that approved recovery activities are implemented.
3. Disseminate information to involved state, federal, and tribal agencies.
4. Coordinate Program activities with the Upper Basin Recovery Implementation Program.
5. Coordinate outreach activities with the Upper Basin Recovery Implementation Program; disseminate information on Program activities to the public through brochures, newsletters and/or the website.
6. Forward plans and recommendations to the Coordination Committee for review and approval.
7. Annual Work Plan:
 - A. Work with the Biology and Hydrology Committees to identify and expedite individual projects that are needed to accomplish the long range plan for each of the recovery elements.
 - B. Draft an annual work plan consisting of high priority individual projects, formulated within the available funding.
 - C. Forward the work plan to the Coordination Committee for review and approval.
8. Coordinate an annual assessment of the Program's recovery progress as outlined in the Program Document.
9. Maintain a list of interested parties and provide those parties with the meeting dates, times, locations, and agendas for Program meetings.
10. Provide draft and final summaries of meetings to committee members.
11. Report to the Coordination Committee at each meeting the status of Program activities and research projects, and accomplishment of milestones; report any problems with maintaining schedules and provide recommendations for solving those problems; implement the recommendations of the Coordination Committee to resolve scheduling problems.
12. Provide support materials for annual funding efforts with the U.S. Congress and state legislatures.

San Juan River Recovery Program Program Management Budget 2009		
Personnel (salary and benefits)	USFWS Funding	Program Base Funding
Coordinator	30,426	0
Assistant Program Coordinator	54,060	54,060
Program Assistant	28,413	28,413
IT-Support	8,000	0
Program Biologist	0	68,060
USFWS Hydrologist	10,000	15,000
Personnel Subtotal	\$130,899.00	\$165,533.00
Travel		
Coordinator (70 days@ \$86)	0	6,020
Coordinator (35trips @ 400miles) * .41	5,740	0
Program Assistant (30 days@ \$86)	0	2,580
Senior Biologist Travel to Farmington (30 days@ \$86)	2,580	0
Senior Biologist (35trips @ 400miles) * .41	4,920	0
Travel to Denver	0	2,000
Denver, 5 days @ \$144	0	1,440
Travel to DC	0	2,000
DC, 7 days @ \$202	0	1,414
Hydrologist Support		5,000
Travel Subtotal	\$13,240.00	\$20,454.00
Committee Meeting Support		
General Office Supplies	2,000	2,000
Meeting space	0	0
Farmington@ \$100/day	0	1,200
Durango @ \$300/day	0	300
Mailings	0	500
Public Notices - (\$80/meeting)	0	2,500
Printing/publication	0	2,500
Gas	1,500	1,500
Misc	500	500
Support Subtotal	\$4,000.00	\$11,000.00
Consultant/ Professional Fees		
Annual Program Assessment	0	\$0
Consultant/ Professional Sub-Total		0
	USFWS Funding	Base Funding
Budget Subtotal	\$148,139	\$196,987
FY 2008 Carry over funds	0	-\$30,000
Total	\$148,139	\$176,987
Total	0	
Administrative charge (22%)	0	\$38,937
Grand Total	\$148,139	\$215,924

FY 2009 Reclamation Program Management

Mark McKinstry UC-735
Bureau of Reclamation
125 South State Street, Room 6107
Salt Lake City, UT 84138-1147
Phone 801-524-3835
FAX 801-524-5499
mmckinstry@uc.usbr.gov

Relationship to SJRIP: Supports Program goals and management by supporting approved activities

Study Goals, Objectives, and End Product: Program Management funds support Reclamation staff involved in program management. Funds are used for the administration of funding agreements, including issuing requisitions for program supplies, and the preparation and oversight of work conducted under interagency agreements, cooperative agreements, contracts, and grants. The funds are also used for formation and participation of the technical and peer-review committees, implementation of committee assignments not specifically identified in a scope of work, reporting, and coordination of water operations. Management support for Capital fund projects, including technical oversight, budgeting, preparation of bids and funding agreements is covered in a separate scope of work. Participation in Hydrology and Biology Committee meetings and business is paid for separately by Reclamation with funds unrelated to the SJRIP.

Task Description and Schedule

Task 1: Manage and administer funding for Recovery Program projects related to the Biology Committee activities. Funding Recovery Program projects requires establishment or modification of approximately 50—60 Reclamation funding agreements or contracts each year. Each financial agreement requires multiple steps and activities, including: submission of requests for Federal assistance for Recovery Program-approved projects; working with Recovery Program's office on funding issues; reviewing and approving (if warranted) project budgets; writing SOWs for RFPs, requesting obligations to cover funding agreement or contract awards; awarding agreements or contract funding to recipients; maintaining agreement and contract filing system including agreement instruments, invoices, and accruals; reviewing and tracking budgets; participating in audits; reviewing and approving invoices; performing periodic site visits to monitor project performance and progress; filing advanced procurement reports; organizing and participating on TPECs; drafting requests for proposals (RFPs); evaluating proposals and awarding contracts; performing agreement closeouts; answering agreement inquiries from auditors, assistance recipients, and the Recovery Program; recording project performance and status of deliverables; and filing recipient performance reports.

Deliverables/Due Dates: Requests from the Recovery Program for funding are processed as they are received. Other deadlines for committee activities are set by the Recovery Program participants during the development of the annual workplan. An annual report on program management activities will be delivered during the annual meeting each year (usually February/March).

Budget FY09**Task 1: Biology Committee Annual Funding Administration****A) Labor**

Position	Salary total/hr	No. persons	Total Hours	Total cost
Reclamation Acquisitions Manager	\$110.00	1	30	\$3,300.00
Biology Committee Technical Representation for Contracts and Agreements*	\$70.00	1	600	\$42,000.00
Lead contract officer	\$110.00	1	40	\$4,400.00
Agreement/Contract Specialist	\$65.00	1	600	\$39,000.00
Agreement specialist	\$50.00	2	800	\$40,000.00
Total				\$128,700.00

* Funding for Reclamation to participate in the Biology Committee is funded by Reclamation and not the SJRIP.

B) Travel

Position	Destination	Purpose	Days	Lodging per day/total	Per diem per day/total	Other*	Airfare total	Total
Reclamation Technical representative	Farmington	Contract support for CC meetings or field trips	3 trips @ 2 days/trip	\$90/\$540	\$45/\$270	\$200	\$2,000	\$3,010.00
Reclamation representative	Denver	Program mtg.	1 trip @ 2 days	\$125/\$250	\$65/\$130	\$60	\$300	\$740.00
Acquisitions Manager	Farmington	CC mtg.	1 day	\$90	\$45	\$50	\$800	\$985.00
Lead contract officer	Farmington	CC/BC mtg.	2 trips @ 2 days	\$90/\$180	\$45/\$135	\$50	\$1,600	\$1,965.00

Lead contract officer	Denver	Program mtg.	1 trip @ 2 days	\$125	\$65/\$130	\$60	\$300	\$615.00
Agreement specialist	Farmington	MC mtg.	1 trip @ 2 days	\$90/\$180	\$45/\$90	\$50	\$800	\$1120.00
Total								\$8,435.00

*Taxi \$20; Parking \$10; Rental car \$50/trip

Budget Summary

FY-2009

Labor			
	Task 1		\$128,700.00
Total labor			\$128,700.00
Total travel			\$8,435.00
Grand total			\$137,135.00¹

¹ This total budget represents a 0.44% increase over the FY2008 budget.

FY-2010 \$144,500.00***

FY-2011 \$152,000.00***

FY-2012 \$160,000.00***

***Includes ~5% adjustment