



Region 4 Inventory and Monitoring Network

Mobile Acoustical Bat Monitoring Annual Summary Report

2014

FELSENTHAL NATIONAL WILDLIFE REFUGE



The Region 4 Inventory and Monitoring Network coordinated acoustical bat monitoring on 50 National Wildlife Refuges and 2 Ecological Services field offices in Regions 2, 3, and 4 during 2014. Surveys establish baseline inventories of bat species at each station and contribute to a landscape-level understanding of bat population trends and habitat associations. Bat call data were collected using Anabat SD2 detectors along up to 30 mile mobile transects during June and July each year following the procedures outlined in the Mobile Bat Acoustical Survey Protocol (USFWS 2012*).

This report summarizes bat calls collected along transects at your station in 2014 and provides annual species detections from 2012 and 2013 for comparison. Calls were classified using Bat Call ID ([BCID] version 2.6a) software and restricted to calls with 5 or more pulses. This annual summary report package contains summary information on transect surveys, and a digital folder containing shapefiles and BCID output files. Summary tables include all classified species observations including those that did not have an associated spatial reference.

All submitted call data and survey metadata have been archived and are available on the Mobile Acoustical Bat Monitoring SharePoint site (<https://fishnet.fws.doi.net/regions/4/nwrs/IM/bats>). Bat call files, GPS data, and survey metadata sheets were reviewed for quality assurance prior to generation of this report. Some submitted data were necessarily excluded due to identified errors in collection processes. All acoustical bat data will be summarized in a comprehensive 2014 annual report following station-level reporting.

**U.S Fish and Wildlife Service. 2012. Mobile Bat Acoustical Survey Protocol, U.S. Fish and Wildlife Service, Region 4, Division of Refuges*

FELSENTHAL NATIONAL WILDLIFE REFUGE

Route Name: FslNWR

Survey date	# bat calls	Route completed?	GPS data?	Survey notes
6/11/2014	16	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Note that this survey route has been modified (shortened) to avoid an area that is commonly prone to flooding. Route length is now only 20.66 miles rather than 28.8 miles
7/7/2014	84	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Route length is now only 20.66 miles and is a permanent change from the original 28.8 miles due to areas prone to flooding.

FELSENTHAL NATIONAL WILDLIFE REFUGE

FslNWR Species Filter 13: CORA, EPFU, LABO, LACI, MYAU, MYSE, NYHU, PESU

Length of transect (miles): 20.66 (28.8 miles in 2012 and 2013 surveys)

Nightly Summary

	<u>Survey date</u>	<u>Total observed</u>	<u>Bats/mile</u>
Big brown bat			
	7/7/2014	15	0.73
Eastern red bat			
	6/11/2014	5	0.24
	7/7/2014	14	0.68
Evening bat			
	6/11/2014	6	0.29
	7/7/2014	24	1.16
Hoary bat			
	7/7/2014	1	0.05
Tricolored bat			
	6/11/2014	4	0.19
	7/7/2014	26	1.26
Unknown			
	6/11/2014	1	0.05
	7/7/2014	4	0.19

Species Summary 2012-2014

Below are the results of bat species classification generated from automated acoustic bat ID software in BCID version 2.6a. Automated acoustical bat classification has inherent limitations based in part on call quality, species filter constraints, and statistical model agreement parameters. Using a conservative approach to generate robust species classifications, calls with fewer than 5 pulses were not classified; therefore the estimates presented here may under-represent the actual number of bats detected on a survey night. Classified calls were geo-referenced using a custom python script to combine GPS locations with individual calls based on unique Date and Time parameters. Accuracy of call classification varies among species but is reported to be >80% correct. Measures of confidence in species identification for each survey night are available as a maximum-likelihood estimator p-value for each observed species in the BCID output files included with this summary report package. BCID version 2.6a does not classify the following species (Seminole bat - LASE, Northern yellow bat - LAIN, Brazilian free-tailed bat - TABR). If these species occur within your monitoring area, the calls generally will be classified to a species with the closest model agreement or classified to "unknown". Species summary tables include all classified calls including those without a spatial reference.

FELSENTHAL NATIONAL WILDLIFE REFUGE

FslNWR Species Filter 13: CORA, EPFU, LABO, LACI, MYAU, MYSE, NYHU, PESU

Length of transect (miles): 20.6 (28.8 miles for 2012 and 2013 surveys)

Annual Summary

	<u>Total observed</u>	<u># Survey nights</u>	<u># Bats/night</u>	<u># Bats/mile</u>
Big brown bat				
2012	17	2	8.5	0.30
2013	66	6	11	0.40
2014	15	2	7.5	0.36
Eastern red bat				
2012	25	2	12.5	0.43
2013	46	6	7.67	0.28
2014	19	2	9.5	0.46
Evening bat				
2012	47	2	23.5	0.82
2013	76	6	12.67	0.46
2014	30	2	15	0.73
Hoary bat				
2012	1	2	0.5	0.02
2013	11	6	1.83	0.07

FELSENTHAL NATIONAL WILDLIFE REFUGE

2014	1	2	0.5	0.02
Northern long-eared bat				
2012	1	2	0.5	0.02
Tricolored bat				
2012	65	2	32.5	1.13
2013	122	6	20.33	0.74
2014	30	2	15	0.73
Unknown				
2012	5	2	2.5	0.1
2013	10	6	1.67	0.06
2014	5	2	2.5	0.12