

NATIONAL WETLANDS INVENTORY

NOTES TO USERS

1:100,000 SCALE MAP

ST LOUIS SE

NATIONAL WETLANDS INVENTORY

1:100,000 MAP NARRATIVE

ST LOUIS SE

INTRODUCTION

The U.S. Fish and Wildlife Service, Office of Habitat Resources, is conducting an inventory of the wetlands of the United States. The National Wetlands Inventory (NWI) is establishing a wetland data base in both map and computer forms for the entire country. The NWI information will serve to identify the current status of U.S. wetlands and can be used as a reference point from which future changes in wetlands can be evaluated.

PURPOSE

The purpose of Notes to Users is to provide general information regarding the production of NWI maps and wetlands found within a relatively similar geographic area. Notes to Users are not intended to include a complete description of all wetlands found in the area nor provide complete plant species information.

MAP PREPARATION

Wetland classification for the NWI maps is in accordance with the "Classification of Wetlands and Deepwater habitats of the United States" by L. M. Cowardin, et al., 1979.

Wetland classification and delineations were produced by photo interpretation of high level aerial photography. The photography used was NHAP color infrared at a scale of 1:58,000. The photography was taken during April of 1984 and March of 1985. To correctly classify the wetlands, ground truthing, soil surveys, and input from regional U.S.F.W.S. personnel were used to relate the various photographic signatures to actual wetland identification and classification. Initial ground truthing with the photography occurred on June 2, 1986. Collateral data included U.S.G.S. topographic maps (7.5 and 15 minute series), and vegetation information.

The user of the map is cautioned that, due to the limitation of mapping primarily through aerial photointerpretation, a small percentage of wetlands may have gone unidentified. Changes in the landscape or habitat could have occurred since the time of photography. Therefore, some discrepancies between the map and current field conditions may exist. Any discrepancies that are encountered in the use of this map should be brought to the attention of Ron Erickson, Regional Wetlands Coordinator; U.S. Fish and Wildlife Service, Region 3, Federal Building, Ft. Snelling, Twin Cities, MN 55111.

GEOGRAPHY

The area being discussed in this report covers the southern portion of Illinois east of the Mississippi River and falling between longitude 90°00'-90°30'W, and latitude 38°00'-38°30'N. Important wetland areas include the broad floodplains associated with the Mississippi River. Bailey's Ecoregion Classification (1980) places this area into the Prairie Parkland Province, Oak-Hickory-Bluestem Parkland Section and the Eastern Deciduous Forest Province, the Oak-Hickory Section. The area is characterized by its gently rolling topography, with steep bluffs and ravines bordering the valley floodplains. Due to the Karst topography present in this area, numerous small depressions or sink-holes dot the landscape. The vegetation consists of both prairies and deciduous forests, resulting primarily from the local conditions of soil and slope. Soil is an important element of hydric conditions and is one of the criteria used to define wetlands. Surveys, prepared by the Soil Conservation Service, serve as an aid in properly identifying wetland habitats. The forested stands are generally restricted to the floodplains, depressions, and on moist slopes. The prairies consist largely of grasses, however, much of the area is now under till.

CLIMATE

Climatic conditions for the study area fall into the subhumid classification. Temperatures are high in the summer (can average up to 55°F) and are mild in winter. The average annual precipitation is between 35-60 inches and occurs primarily during the growing season.

WETLAND COMMUNITIES

Wetlands and deepwater habitats within the area under discussion, fall within the Palustrine, Lacustrine, and Riverine systems. Deepwater habitats are areas which are permanently flooded and are characterized by open water on the aerial photography. These habitats are present in the Riverine and Lacustrine systems, while wetland habitats are present in all systems. (See table 1).

Numerous palustrine unconsolidated bottom (UB) areas are present within the study area, with the majority being excavated (x) or impounded (h) farm ponds. All of the farm ponds are delineated as intermittently exposed (G), except in instances where the pond is 1-2 acres in size. In this case, the pond may be classified as semipermanently flooded (F). Other unconsolidated bottom areas present are natural ponds.

Palustrine water bodies are often vegetated with a rooted vascular and floating vascular aquatic bed. This mixed community of mosses, duckweed (*Lemna* spp.), and other weeds, was often not discernable on the photography. These aquatics are only classified when visible on the photography or ground truthed as such.

Palustrine emergents (EM) are abundant in this area, ranging from temporarily (A) to seasonally (C) and semipermanently (F) flooded water regimes. Species commonly encountered in the temporarily flooded habitats include reed canary grass (Phalaris sp.), sedges (Carex spp.), goldenrod (Solidago spp.), dock (Rumex spp.), and smartweed (Polygonum spp.). Species present in the seasonally and semipermanently flooded water regimes are cattail (Typha latifolia), river bullrush (Scirpus fluviatilis), dock, reed canary grass, arrowhead (Sagittaria spp.), cutgrass (Leersia spp.), and spikerush (Eleocharis spp.). Emergent habitats found under semipermanently flooded conditions are often located in old river meanders and sloughs. Cattail and arrowhead are common in these areas.

Basins located in agricultural fields are often found to support emergents when not drained or otherwise manipulated. Basins, which can be described as depressions, potholes, or meander scars, hold water early in the growing season, but are later farmed. The farmed modifier (f) is used when delineating such emergent habitats.

Palustrine scrub shrub (PSS) and palustrine forested (PFO) wetlands are characterized by woody species. The scrub shrub habitats are often found on the perimeters of emergent wetlands, with either temporary or seasonal water regimes. Willow is the dominant species of the scrub shrub habitat, along with the commonly found buttonbush (Cephalanthus occidentalis) and cottonwood saplings. The palustrine forested wetlands, which primarily occur within the floodplains of the Mississippi River, generally flood on a temporary basis. Common species occurring in these temporarily flooded areas are silver maple (Acer saccharinum), willow, elm (Ulmus spp.), cottonwood, box elder (Acer negundo), and sycamore (Platanus occidentalis). Old meander scars and depressions often support seasonally flooded forests, supporting species such as silver maple, willow, river birch (Betula nigra), cottonwood, and green ash (Fraxinus pennsylvanica). Forests along the banks of the Mississippi are found to be temporarily flooded, with seasonal forests restricted to depressions, scars, and sloughs. No semipermanently flooded forests were observed in the field.

The riverine system includes the classes: unconsolidated bottom (UB), unconsolidated shore (US), and streambed (SB). Unconsolidated bottom and shore are restricted to the riverine lower perennial (R2) subsystem. Most streams emptying into the Mississippi floodplain were identified as perennial by U.S.G.S. and are labeled accordingly (R2UBH). Small perennial and intermittent streams are often entrenched drawing down quickly after flooding. Forested areas along these channels often do not pond or retain water. Intermittent streams are classified as having semipermanent flow (R4SBF). U.S.G.S. topographic information is used to determine intermittent and perennial breaks along the riverine systems. Many rivers and streams are excavated (x) to improve drainage. In instances of streamside vegetation, the stream is delineated as a palustrine feature.

Natural or artificial open water bodies greater than 20 acres are classified as lacustrine. These large bodies have been created through excavation (x) or by impounding (h) rivers and streams. Generally, these large open water systems are classified as limnetic (L1) with an unconsolidated bottom, and are considered to be permanently flooded (H).

BIBLIOGRAPHY

Bailey, Robert G., 1980. Description of the Ecoregions of the United States. U.S. Department of Agriculture Forest Service, Ogden, UT.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe, 1977. Classification of Wetlands and Deepwater Habitats of the United States. U.S. Fish and Wildlife Service, Washington, D.C.

TABLE 1

NWI CODE	NWI DESCRIPTION	COMMON DESCRIPTION	CHARACTERISTIC VEGETATION AND PHYSIOGRAPHIC FEATURES
R4SB	Riverine, intermittent, streambed	Creek, Streambed, Canal	Unvegetated. Sand to cobble-gravel
R2UB	Riverine, lower perennial, unconsolidated bottom	River	Unvegetated. Mud to sand, cobble-gravel
R2US	Riverine, lower perennial, unconsolidated shore	River flat	Unvegetated. Sand to cobble gravel
L1UB	Lacustrine, limnetic, unconsolidated bottom	Open water lake	Unvegetated. Sand to mud
L2UB	Lacustrine, littoral, unconsolidated bottom	Shallow lake	Unvegetated. Sand to mud
L2US	Lacustrine, littoral, unconsolidated shore	Lake shore	Unvegetated. Sand to cobble-gravel
L1AB	Lacustrine, limnetic, aquatic bed	Pond weeds, water weeds	Duckweed (<u>Lemna</u> sp.)
PUB	Palustrine unconsolidated bottom	Open water, pond	Unvegetated. Sand to mud
PAB	Palustrine aquatic bed	Pond weeds, water weeds	Duckweed (<u>Lemna</u> sp.) Pondweed (<u>Potamogeton</u> sp.)
PEM	Palustrine emergents	Marsh or meadow	Cattail (<u>Typha latifolia</u>) Reed canary grass (<u>Phalaris</u> sp.)

TABLE 1

NWI CODE	NWI DESCRIPTION	COMMON DESCRIPTION	CHARACTERISTIC VEGETATION AND PHYSIOGRAPHIC FEATURES
PEM (con't)	Palustrine emergents	Marsh or meadow	Bullrushes (<u>Scirpus</u> sp.) Spike Rush (<u>Eleocharis</u> sp.) Sedges (<u>Carex</u> sp.) Arrowhead (<u>Sagittaria</u> sp.) Smartweed (<u>Polygonum</u> sp.) Cutgrass (<u>Leersia</u> sp.) Dock (<u>Rumex</u> sp.) Cocklebur (<u>Xanthium</u> sp.) Giant Ragweed (<u>Ambrosia trifida</u>) Velvetleaf (<u>Abutilon theophrasti</u>) Goldenrod (<u>Solidago</u> sp.) Milkweed (<u>Asclepias</u> sp.)
PSS	Palustrine scrub shrub	Shrub wetland	Willow (<u>Salix</u> sp.) Buttonbush (<u>Cephalanthus occidentalis</u>)
PFO	Palustrine forested	Forested wetland	Silver maple (<u>Acer saccharinum</u>) Cottonwood (<u>Populus deltoides</u>) Willow (<u>Salix</u> sp.) Box elder (<u>Acer negundo</u>) Green ash (<u>Fraxinus pennsylvanica</u>)

NWI#66

TABLE 1

NWI CODE	NWI DESCRIPTION	COMMON DESCRIPTION	CHARACTERISTIC VEGETATION AND PHYSIOGRAPHIC FEATURES
PFO (cont)	Palustrine forested	Forested wetland	Slippery elm (<u>Ulmus rubra</u>) American elm (<u>Ulmus americana</u>) Honey locust (<u>Gleditisa triacanthos</u>) Sycamore (<u>Platanus occidentalis</u>) River birch (<u>Betula nigra</u>) Red Mulberry (<u>Morus rubra</u>)

NWI#66