

WYOMING - MONTANA COAL REGION WETLAND INVENTORY
U. S. FISH AND WILDLIFE SERVICE
1:100,000 MAP NARRATIVE REPORT
-SHERIDAN SE-

INTRODUCTION

In 1974 the U. S. Fish and Wildlife Service directed its Office of Biological Services to complete an inventory of the Nation's wetlands. As part of this overall objective an effort began in September 1977 to delineate and classify photointerpretable wetlands within the Powder River Basin of Wyoming and Montana. Wetland maps at 1:100,000 scale and wetlands overlay maps at 1:24,000 scale are produced at National Wetland Inventory headquarters in St. Petersburg, Florida. Final Wyoming-Montana wetland maps are available at U. S. Fish and Wildlife Regional Office in Denver, Colorado. An integral part of all final wetland maps is the completion of narrative reports, which are available for each 1:100,000 inventoried. The following narrative report provides both basic and specific data which aids the user in understanding not only the general area of the Sheridan SE quadrangle, but also representative details of its wetland habitat.

MAP PREPARATION

Contractor for this project was Martel Laboratories, Inc., of St. Petersburg, Florida. Aerial photographic interpretation was performed by Alexander Kosinski and Charles Storrs. This report was prepared by Thomas Owens and Alexander Kosinski. Regional Wetland Coordinator is Charles Elliott, U. S. Fish and Wildlife Service, P. O. Box 25486, Denver, Colorado 80225.

Wetland delineation and classification for the Sheridan SE 1:100,000 wetland map was done on color infrared aerial photographs. Most of the map was covered by middle altitude photography taken in May, June, and July 1976 at a scale of 1:31,680. The western one-third and northeastern corner of Sheridan SE was not covered by photography and was not delineated. Limited field checking was conducted in late summer and early autumn of 1977.

Map users are cautioned that mapping with aerial photography has limitations. Through stereoscopic examination of photography wetlands are identified and classified on the basis of color, tone, texture, pattern, site, size, local ecology, and cultural patterns. Aerial photographs reflect conditions during the year and season they were taken. In addition, most wetlands on this map were delineated using high altitude aerial photography, which presents limitations. Small wetlands are too diminutive at this scale to delineate. This is also true for many stream channels whose width was less than the width of a pen line. Resolution limitations inherent in high altitude imagery cause problems in accurately recognizing ground conditions.

Any noted discrepancies regarding wetland omissions, inclusions or errors should be given to the U. S. Fish and Wildlife Regional Wetland Coordinator who is located in Denver, Colorado.

AREA DESCRIPTION

Bailey's Ecoregions

Sheridan SE falls into two provinces in Bailey's Ecoregions of the United States which classifies land into a hierarchal system based upon bioclimatic, geologic, and geomorphic criteria. Covering the quadrangle's western 20 percent is the Rocky Mountain Province, Douglas-Fir Section (3112M). The remaining 80 percent is in the Great Plains Short Grass Steppe Province, Grama-Needlegrass-Wheatgrass Section (3111L).

Hammond's Land-Surface Forms

Hammond's Land-Surface Forms systematically characterize United States' topography. Sheridan SE falls into three forms in this system. The western 20 percent is in the Rocky Mountain Division (IV), Middle Rocky Mountain Subdivision (3), High Mountains (IV-3D6), which have less than 20 percent of area gently sloping and local relief over 3,000 feet. The other two forms are in the Interior Division (III), Upper Missouri Broken Lands Subdivision (10). The central 40 percent is in Plains with Hills (III-10B3b) which have 50 to 80 percent of area gently sloping, local relief 300 to 500 feet, and 50 to 75 percent of gentle slope in lowland. The eastern 40 percent is in Open High Hills (III-10C4b) characterized by 20 to 50 percent of area gently sloping, local relief 500 to 1,000 feet, and 50 to 75 percent of gentle slope in lowland. -

Hydrologic Mapping Units

Hydrologic mapping units are part of an effort by the U. S. Geological Service to provide a series of uniform, Nationally consistent maps which accurately delineate hydrographic boundaries for Federal and State water agencies. Units are designated by eight-digit numbers, which are tied to a computer file (Catalog of Information on Water Data) which contains information on water data activities (Langford and Kapinas, 1979). All hydrologic mapping units in Sheridan SE are in the Missouri Region (10). 10090201 in the southwestern corner covers 2 percent of the map and 10090206 covers 18 percent in the northwestern corner. 10090205 covers 42 percent in the west-central portion of the map and 10090202 covers 38 percent in the east.

Topography

Sheridan SE's plains (since the Bighorn Mountains on the quadrangle's western edge was not delineated, this area will not be discussed in the remaining sections -- only the plains area will be addressed)

topography is a result of erosion upon layers of alternating shales and sandstones. These layers differ in permeability, which coupled with high relief, directly influences drainage density. When impermeable shale is exposed, water infiltration is poor and runoff is high, which favors development of many headward eroding streams and gully channels. If drainages are closely spaced, vegetation cover sparse, and rain occurs in thunderbursts, then badland topography will result. When permeable sandstone is exposed, rainfall is more completely absorbed, causing less runoff and a less intricate, more widely spaced drainage pattern develops. Instead of ridge and valley terrain typical of shale, sandstone expresses itself as a relatively smooth, flat to rolling hill topography.

Geology

All surface bedrock in Sheridan SE's plain section is sedimentary, consisting of alternating layers of sandstone, siltstone, and shale. The oldest deposits are Cambrian at the foot of the Bighorns, becoming younger farther east. The youngest, the Eocene Wasatch Formation covers the quadrangle's eastern two-thirds, with the older Fort Union and Lance Formations to the west.

The Lance Formation was deposited late in the Cretaceous Period when a sea was shrinking eastward. As this sea was displaced eastward, it moved its associated alluvial and coal deposits with it.

The Fort Union Formation was deposited during subtropical Paleocene environment and contains wide deposits of coals where ancient swamps once existed. Accumulation of eroded sediments from the Rockies on swampy ground resulted in alternating layers of carbonaceous shale and fine grained sandstone intermingled with coal seams.

The younger Wasatch Formation was deposited during the Eocene, a time when basin filling continued, but with a gradual diminution of coal swamps as the climate changed to humid temperate conditions. The Wasatch Formation consists of fine- to coarse-grained sandstones with interbedded shales, and less coal in more recent strata. Pink, yellow, and red silts are characteristic of the Eocene strata, which are often eroded into badland topography.

Where coal seams outcrops have ignited from lightning, spontaneous combustion, or man-made fires, beds of baked rock called clinker or "scoria" are found. Reddish clinker beds are sandstone and shale beds that have been subjected to melting and baking by heat and gases rising from underlying burning coal seams. Depending on availability of oxygen, clinker zones may follow the coal outcrop underground and extend for almost a mile.

Soils

Soil is an important element of wetlands; it is one criterion used to define wetlands. "The substrate (of wetlands) is predominately

undrained hydric soil" (Cowardin, 1979). The National Wetland Inventory, in cooperation with the U. S. Soil Conservation Service, is preparing a list of hydric soils to accompany the Cowardin, et al., wetland classification system.

Soils of Sheridan SE are a result of semi-arid climate with its associated vegetation and organisms acting upon Tertiary sandstones and shales. These soils are light gray-brown, often possessing a lime-carbonate (caliche) layer at shallow depths. The zone of altered bedrock is three feet below the surface or less, with slopes having thinner soil cover and bedrock outcrops. Soils developed on steeply sloping uplands and ridges are easily eroded, resulting in dissected, broken ground. Loamy soils derived from clinker beds have developed on steep slopes, are easily eroded, and have low agricultural potential. Loams developed from sandstones and shales on steep slopes are also easily eroded and have low agricultural productivity. Loams developed from sandstones and shales on steep slopes are also easily eroded and have low agricultural productivity. Soils on gently sloping uplands have moderate to high erodibility and moderate agricultural productivity. Near large streams soils that have developed on alluvium are alkaline, have wind erodibility, and are subject to flooding.

Climate

Sheridan SE is a temperate and semi-arid climate and experiences wide variation in temperature and precipitation between winter and summer. The growing season averages about 120 days per year. Average annual precipitation is 13 inches. Two-thirds of annual precipitation occurs between March and August of an average year. One-third occurs as snow, which annually averages 50 inches. Snowfall is generally well distributed through winter and spring, but December usually receives heaviest accumulations. Summer-time hailstorms occur, which are capable of causing serious crop damages. Prevailing winds are from the northwest, which have their highest velocities in spring.

Mean Monthly Temperature January	21.50F.
Mean Monthly Temperature July	73.00F.
Lowest Minimum Temperature January	-40.00F.
Highest Maximum Temperature July	105.00F.

WETLANDS

Community Descriptions

Sheridan SE has wetland types reflective of semi-arid climate with ridge and gully terrain that is interrupted by expanses of rolling terrain. This quadrangle is characterized by many artificial impoundments and "intermittent wetlands," almost all of which are in the Palustrine system.

Palustrine System

Artificial Impoundments

Many gullies, intermittent streams, and natural springs are dammed. These water storage structures create many wetlands on this map. These wetlands are classified as Palustrine Open Water (POW), Palustrine Emergent (PEM), Palustrine Flat (PFL) or combinations of these classes. Size varies from less than one-half acre catch basins to reservoirs of dozens of acres.

Palustrine Open Water impoundments are semi-permanently flooded and provide a reliable source of water throughout the growing season most years. These wetlands have limited aquatic vegetation.

Palustrine Flat impoundments are usually unvegetated and contain silty mineral substrates. However, some flats contain vegetation in low densities or in widely scattered clumps. Species included are western wheatgrass (Agropyron smithii), spikerush (Eleocharis spp.), and bulrush (Scirpus spp.).

Palustrine Emergent impoundments possess higher densities of hydrophytic vegetation than do flats. Water regime influences species composition. Spikerush, either in pure stands or mixed with western wheatgrass is present at sites with temporary to seasonal flooding. Bulrush is associated with seasonal to semi-permanent flooding.

Local factors, such as fluctuations in the quantity of spring meltwater runoff, variability of flow from natural springs and seeps, and the unpredictability of thunderstorm recharge, play an important role in the diversity of water regimes in all three classes of impoundment wetlands.

Intermittent Wetlands

Intermittent wetlands are naturally occurring wetland features that show up on flat to rolling topography as saucer-shaped shallow depressions. These depressions often occur in groups, which may contain dozens of depressions scattered over square miles.

Terminology used by ASCS refer to these depressions as "clay overflows." Some depressions appear to have developed from gradual accumulation of clay colluvium. This clay is deposited by runoff from eroding shale scarps into basins.

Other shallow depressions seem to occur near the transition zone between two differing rock types; specifically sandstone and shale. It appears that the permeable uppermost sandstone layer has been weathered thinly enough in spots to permit breakdown of underlying shale. This deterioration of shale produces clay, which compacts differentially when exposed to moisture, creating a shallow basin as it settles. The basin collects more rainfall as it deepens, further hastening the weather process. At the same time, it serves as a collection basin for fine silts and surface clays washed in by runoff from surrounding uplands.

In both cases, clay is the common denominator in the genesis of these shallow depressions. Intermittent wetlands are classified as either Palustrine Emergent, Palustrine Flat, or combinations of them.

Palustrine Flats have substrates of silt-clay. Both non-vegetated and pioneer sparsely vegetated (less than 10 percent area coverage) varieties are present on this quadrangle. Some non-vegetated flats are disturbed by cattle trampling. Vegetation includes western wheatgrass and spikerush. Flats are often saline. Water regimes range from intermittently through seasonally flooded.

Palustrine Emergent depressions fall into three categories: (1) homogeneous purestands of vegetation, (2) vegetation occurring in distinct concentric rings that differ in species composition or density, (3) vegetation existing in a random patchwork of amorphous shapes, also differentiated by species composition or density. Western wheatgrass is usually dominant, since most depressions are intermittently flooded. Where surface flooding is temporary to seasonal, spikerush and bulrush are in association with western wheatgrass. Saline conditions are common.

Riverine System

Riverine Lower Perennial Open Water (R2OW) and Riverine Lower Perennial Beach-Bar (R2BB) are delineated along the Powder River. Lower perennial streams have slow moving water, substrates of sand and silt, low dissolved oxygen levels, and well developed floodplains. Open water is delineated in the main channel where there is a permanent flow of water. Beach-bar is delineated on the sides of the main channel where old channels exist. Beach-bars are flooded during high water and are exposed during low water. The substrate is sand, silt or gravel and only pioneer vegetation is found on them. This vegetation is usually washed away during the next high water.

Wetland Values

The function of impoundments is to store water for cattle and sheep. Grazing is the main agricultural land use in this area because of low precipitation and shallow soils. Where farmland does exist, the principle crops of wheat, oats, barley, and hay are grown with dry land farming methods. Intermittent wetlands are grazed by livestock if they are vegetated.

All streams in Sheridan SE are rated as low production trout waters; often incapable of sustaining a fishery (Wyoming Stream Fishery Classification, 1976).

Kearney Creek is rated as a "high priority fishery resource" by the Fish and Wildlife Service. This means it probably has an endangered fish species, has intensively used habitat by a "highly valued" population, and has low potential for restoration to an equivalent fishery following water development. French, Muddy, Billy, and Poison Creeks are "substantial"

fishery resources." They are occasionally used by "highly valued" fish populations and have moderate potential for restoration (Stream Evaluation Map, Wyoming, 1978).

Water storage impoundments constructed for domestic livestock use are used by local wildlife such as birds (meadowlark, horned lark), rodents (mice, voles), small herbivorous mammals (rabbits, hares), and fur bearing predators (coyote, fox), may include these impoundments in their home territories and use them as their source of drinking water. In periods of drought, a single isolated impoundment may be valuable as the only source of water available for miles around.

Standing water is present in intermittent wetlands during periods of shorebird migration and is used by birds as resting and feeding areas. Waterfowl use larger open water impoundments or flooded intermittent lakes as migration habitat and breeding areas.

Floodplains of the Powder River, Crazy Woman, Rock and Clear Creeks are critical habitat for whitetail deer.

Wetland Loss and Vulnerability

Increased development in coal strip mining and its associated facilities has begun in Sheridan SE. Low sulfur coal exists in vast deposits in east-central Wyoming and much of this coal occurs in Sheridan SE. Coal mining operations currently in the planning stage are expected to disturb thousands of acres by the end of this century.

In Sheridan SE, boundaries of known strippable coal incorporate a large number of intermittent wetlands and impoundments in the Palustrine system. Wetlands will be lost if strip mining operations include them within their sites.

Large scale surface mining will alter existing rock and soil patterns, which will change geochemical characteristics. Geochemistry is the vital factor controlling distribution of soil types, vegetation, water quality, and to some extent, air quality.

Many coal seams in Sheridan SE are important aquifers. Disturbance of these aquifers during coal operations will reduce water quality. Increased erosion, sedimentation, and discharge of toxic wastes in ground water that is pumped from the mine will be detrimental to stream water quality and consequently ground water quality. Regional impact of mining on ground water table has not been determined.

Petroleum and natural gas exploration will be developed faster in the future. However, where currently operating wells are situated near wetlands, the only disturbance appears in the immediate vicinity of the well. Intermittent lakes near wells show no evidence of disturbance.

REFERENCES

- Bailey, R. G. 1978. Ecoregions of the United States. U. S. Forest Service, Intermountain Region. Ogden, Utah.
- Breckenridge, Roy M., Gary B. Glass, Forrest K. Root, and William G. Wendell. 1974. Campbell County, Wyoming, Geologic Map Atlas and Summary of Land, Water, and Mineral Resources. County Resource Series No. 3. The Geological Survey of Wyoming, in cooperation with the Mineral Division, Wyoming Department of Economic Planning and Development, and U. S. Bureau of Mines.
- Britton, Nathaniel Lord, and Hon. Addison Brown. 1970. An Illustrated Flora of the Northern United States and Canada. In Three Volumes. Dover Publications, Inc. New York.
- Burt, William H., and Richard P. Grossenheider. 1976. A Field Guide to the Mammals. Houghton Mifflin Company. Boston.
- Cowardin, Lewis M., Virginia Carter, Francis C. Golet, and Edward T. LaRoe. 1979. Classification of Wetlands and Deep-Water Habitats of the United States. Office of Biological Services, Fish and Wildlife Service, U. S. Department of the Interior. Washington, D.C.
- Dott, Robert H. Jr., and Roger L. Batten. 1971. Evolution of the Earth. McGraw-Hill Book Company. New York.
- Fenneman, Nevin M. 1931. Physiography of Western United States. McGraw-Hill Book Company, Inc. New York.
- Hunt, Charles B. 1974. Natural Regions of the United States and Canada. W. H. Freeman and Company. San Francisco.
- Hunt, Charles B. 1967. The Evolution of North America. W. H. Freeman and Company. San Francisco.
- Langford and Kapinas. 1979. The National Water Data Network: A Case History. Water Resources Council. Vol.15, No. 6. pp. 1687-1691.
- Thornbury, William D. 1969. Principles of Geomorphology. John Wiley and Sons, Inc. New York.
- Thornbury, William D. 1965. Regional Geomorphology of the United States. John Wiley and Sons, Inc. New York.
- U. S. Fish and Wildlife Service. 1978. Stream Evaluation Map, Wyoming. Billings, Montana. 1 sheet.
- U. S. Geological Survey. 1976. Hydrologic Unit Map - 1974, State of Wyoming. Department of the Interior, United States Geological Survey, in cooperation with the U. S. Water Resources Council.

Wyoming Game and Fish Department, Game Division, n.d. Critical Habitat Overlay (1:500,000 scale map), Wyoming Game and Fish Department. Cheyenne, Wyoming.

Wyoming Game and Fish Department, Fish Division. 1976. Stream Fishery Classification (1:1,000,000 scale map), Wyoming Game and Fish Department. Cheyenne, Wyoming.

Wyoming Game and Fish Department, Game Division, n.d. Whitetail Deer -- Powder River (year-long habitat map 1:500,000 scale). 3 sheets. Wyoming Game and Fish Department. Cheyenne, Wyoming.

Wyoming Game and Fish Department, Game Division. 1978. Whitetail Deer -- Thunder Basin (year-long habitat map 1:500,000 scale). 4 sheets. Wyoming Game and Fish Department. Cheyenne, Wyoming.