

COLORADO WETLAND INVENTORY
U. S. FISH AND WILDLIFE SERVICE
1:100,000 MAP NARRATIVE REPORT
-LA JUNTA NE-

INTRODUCTION

In 1974, the U. S. Fish and Wildlife Service directed its Office of Biological Services to complete an inventory of the Nation's wetlands. As part of this overall objective, an effort began in September 1978, to delineate and classify photo-interpretable wetlands within the eastern five-sevenths of Colorado.

Wetland maps at 1:100,000 scale and wetland overlay maps at 1:24,000 scale are produced at National Wetland Inventory headquarters in St. Petersburg, Florida. Final Colorado wetland maps are available at the U. S. Fish and Wildlife Service's Regional Office located in Denver, Colorado. An integral part of all final wetland maps is the completion of narrative reports for each 1:100,000 quadrangle inventoried. The following narrative report provides both basic and specific data which aids the user in understanding not only the general area of the La Junta NE quadrangle but also representative details of its wetland habitat.

MAP PREPARATION

Contractor for this wetland inventory was Colorado Division of Wildlife, 317 West Prospect, P. O. Box 2287, Fort Collins, Colorado 80526. Richard Hopper was the contract officer. Photo-interpretation was done by the subcontractor, Colorado State Forest Service, Foothills Campus, Colorado State University, Fort Collins, Colorado 80523. Photo-interpreters were Thomas Owens, Charles Storrs and Alexander Kosinski. Preparation of this narrative report was completed by Thomas Owens. Regional Wetland Coordinator was Charles Elliott, U. S. Fish and Wildlife Service, Denver Federal Center, P. O. Box 25486, Denver, Colorado 80225.

Wetland delineation and classification for La Junta NE 1:100,000 quadrangle was done on 1:80,000 black and white aerial photographs taken in June and July 1975 and June 1976. Photography covered 100 percent of the quadrangle. Wetland classification was done in accordance with Cowardin, et al., 1979. Specific mapping conventions developed at National Wetland Inventory headquarters were used to assist in photo-interpretation. Field checking was done March 19, 1979.

Map users are cautioned that mapping with aerial photography has limitations. Wetlands are identified and classified through stereoscopic examination of photography on the basis of tone, texture, pattern, site,

size, local ecology, and cultural patterns. Aerial photographs reflect conditions during the year and season they are taken. In addition, the 1:80,000 black and white photography used on this project was photographed for purposes other than wetland mapping. The small scale precludes delineating very small wetlands (less than 1/4-acre) and narrow linear wetlands (less than 15 feet wide). Black and white emulsion makes distinguishing between classes of vegetation (and nonvegetation) difficult. Some imagery was not photographed during the best season for wetland delineation and classification. If photographed too early or too late in the season, moist (dark) tones are not evident. The photography was four to five years old when it was interpreted and land use changes have occurred.

Any discrepancies noticed regarding wetland omissions, inclusions, or errors should be given to the U. S. Fish and Wildlife Service Regional Wetland Coordinator who is located in Denver, Colorado, and whose address is on the previous page.

Special Mapping Problems

This quadrangle has many small depressions scattered across the plains. There are two types of depressions: wetlands emergent species and non-wetlands without emergents. Non-wetland depressions without emergents are often dark in tone and conspicuous on the imagery, which is due to slightly more moisture and lush vegetation than surrounding areas, but have no lighter toned ring. Depressions with wetland species (usually) have a lighter toned ring in them which shows the extent of inundation after rain. Depressions receive water from local summer thunderstorms, are dry most of the time, and are not part of drainage systems. These wetlands are locally important, but are often difficult to see on the imagery, unless flooded after a thunderstorm.

AREA DESCRIPTION

Bailey's Ecoregions

La Junta NE 1:100,000 quadrangle falls into one province in Bailey's Description of the Ecoregions of the United States, 1978, which classifies land into a hierarchal system based upon bioclimatic, geologic, and geomorphic criteria. The province is Great Plains-Shortgrass, Grama-Buffalo Grass Section (3113L) and is characterized by bunched short grasses, with scattered trees.

Hammond's Land-Surface Forms

La Junta NE falls into two Hammond Land-Surface Forms which systematically characterizes United States' topography. Both forms are in the Interior Division (III). One form is in the Rocky Mountain Piedmont Subdivision (13) and is Tablelands of Moderate Relief (III-13B3c) which has 50 to 80

percent of area gently sloping, local relief 300 to 500 feet, 50 to 75 percent of gentle slope on upland, and covers 55 percent of the quadrangle. The second form is in the High Plains Subdivision (14) and is Smooth Plains (III-14A2c) characterized by more than 80 percent of area gently sloping, local relief 100 to 300 feet, 50 to 75 percent of gentle slope on upland, and covers 45 percent of the quadrangle.

Hydrologic Mapping Units

Four hydrologic mapping units are found in La Junta NE (Hydrologic Unit Map of Colorado, 1974). Hydrologic units are part of an effort by the United States Geological Survey to provide a series of uniform, Nationally consistent maps which accurately delineate hydrographic boundaries for Federal and State water resource agencies. Units are designed by eight-digit numbers tied to a computer file (Catalog of Information on Water Data) which contains information on water data activities (Langford and Kapinas, 1979). All units are in the Arkansas-Red-White Region (11). 11020009 covers 35 percent in the northwestern corner and 11020013 covers 27 percent in the center of the quadrangle. 11030001 covers 6 percent in the northeastern corner and 11040005 covers 32 percent in the southeastern corner.

Geography

La Junta NE is a semi-arid quadrangle covered by plains which have level to rolling topography. Two Buttes, which rise 200 feet over the plains, are the quadrangle's outstanding topographic features. Elevations range from 3,400 to 4,700 feet and vegetation is short grass prairie with cholla cactus and cottonwoods and willows along streams.

Two Buttes Reservoir is the quadrangle's only reservoir.

There are no towns of note on the quadrangle. La Junta NE's major economic activity is agriculture. Intensive irrigated agriculture occurs on the quadrangle's eastern half where center pivot sprinklers are supplied with well water. Dryland farming and ranching are practiced away from water sources.

A strip on the eastern edge, three miles wide, is in Kansas.

Geology

A large portion of La Junta NE in the north and west is covered by Cretaceous (70 to 135 million years ago) shale, limestone, and sandstone that were deposited when this area was under a sea. The rest of the quadrangle is covered by the Ogallala aquifer, an important water

bearing formation of sandstone deposited in the Tertiary Period (3 to 70 million years ago) (Chronic and Chronic, 1972; Tweto, 1979).

Soils

Soil is an important element of wetlands; it is one criterion used to define wetlands. "The substrate of wetlands is predominately undrained hydric soil" (Cowardin, et al.). The National Wetland Inventory, in cooperation with the U. S. Soil Conservation Service, is preparing a list of hydric soils to accompany the Cowardin, et al., wetland classification system.

Two major wetland soil types are found in La Junta NE: soils associated with drainages and soils associated with flood-irrigated meadows.

Where streams are intermittent, wetland soils are sandy and gravelly, unstable, excessively drained, subject to occasional flooding, and have low water tables. Native vegetation includes cottonwoods and annual forbs (Cipra, et al., 1978).

Flood-irrigated soils are found where there is a ready supply of water. These soils may not have been wetland soils originally, but since they have been irrigated, the water table is less than three feet from the surface. These soils are deep, level, range in texture from sandy loam to clay loam, and are often saline. Vegetation includes salt grass, alkali sacaton, tall grasses, sedges, rushes and cattails in low pockets (Cipra, et al.).

Climate

La Junta's climate is semi-arid and continental, with cool dry winters and warm, relatively dry, summers. Two Buttes receives 12.5 inches of precipitation annually, 15 inches of this coming as snow. January's average maximum temperature is 47.4°F., average minimum is 17.3°F. July's average maximum temperature is 92.7°F., average minimum is 61.2°F. (Benci and McKee, 1977). The growing season is 165 days (SCS data, 1978).

WETLANDS

Community Description

Lacustrine System

One lake type is found within La Junta NE which is reservoir (L10WKZ) (information in this section comes from field notes taken March 29, 1979).

Two Buttes Reservoir is the only reservoir found in La Junta NE. Water levels fluctuate as much as 30 feet and the areal extent of the open water changes significantly during the year because of filling in spring with snow melt and drawing down during summer for agricultural irrigation. The exposed shoreline is composed of rocks, gravel, sand and mud. Reservoirs normally retain some water throughout the year.

Riverine System

One stream type is delineated; intermittent stream (R4SBW). This stream type is the most common stream type on the eastern plains where there is no permanent water source to supply moisture. Intermittent streams have a sandy substrate that is very well drained which is often scoured by flash floods. These streams flow after snow melt and after local summer thunderstorms.

Palustrine System

An important palustrine type is flood-irrigated meadow (PEMKC). Flood-irrigated meadows are found along streams and below springs. Ditches are built along the meadows; upper edges to allow water to flow from upper sources and spill out over the meadows. Standing water can be found for short periods early in the growing season; soil remains moist for extended periods through the growing season. Flood-irrigated meadows have not been cultivated and vegetation is native. Meadows are hayed and grazed by cattle. Vegetation includes Juncus arcticus, Eleocharis acicularis (species identification was according to Harrington, 1955; Fasset, 1957; Weber, 1976; Nelson, 1977), along with grasses and forbs. Many flood-irrigated fields are cultivated; these were not delineated.

Scrub/shrub and scrub/shrub-emergent areas are found along streams in this quadrangle (PSSW, PSSY, PSS/EMW, PSS/EMY). The water regime is dependent upon the amount of water flowing in the stream. In the western portion, where water is relatively plentiful, the water regime tends towards the seasonal regime. In the east, where moisture is scarce, water regimes are drier. Shrub species are Salix spp.; understory emergents include Juncus arcticus and Carex spp.

Forested wetlands (PFOW, PFOY, PFO/EMW, PFO/EMY, PFO/SSW, PFO/SSY) are found throughout the quadrangle along streams. The water regime situation is the same as that of scrub/shrub areas. Tree species include Populus sargentii and Salix spp. Understory species are the same as those mentioned with scrub/shrub-emergent areas.

In the plains numerous areas (PFLW, PEMW) are delineated which are small dugouts or impoundments constructed to supply water to livestock. These flats are dependent upon local precipitation for their water supply and do not receive enough moisture to remain wet year-round. They often

have saline soils. Vegetation consists of sparse stands of Distichilis stricta, Sporobolus airoides, forbs, and grasses.

Palustrine open water (POWY) areas are uncommon in this semi-arid region. These areas are impoundments that have open water most of the year.

Intermittent depressions are also delineated on the plains (PEMJ) that are the result of: wind deflation or blowouts, solution-subsidence (leaching water removes limestone and subsequent deflation), or differential compaction of Tertiary sediments of the plains (Thornbury, 1965). Delineated depressions are generally a few feet in depth and from 100 to several hundred feet across. There is not sufficient moisture in spring after snow melt to supply them with moisture; they are filled with water after local summer thunderstorms and hold water for a few days. They are dry most of the time, but do receive enough moisture to support stands of Eleocharis spp., as well as upland vegetation.

Many palustrine wetlands are cultivated, but remain wetlands. These are called Palustrine Farmed (PF).

Wetland Values

An important wetland value is flood-irrigated hay meadow production of hay for cattle (information in values section is from Hopper, 1980). Flood-irrigated hay meadows produce hay at a much higher rate than dryland meadows do. Wildlife also benefit from wet hay meadows: ducks nest in them.

Two Buttes Reservoir and Turks Pond (a 10-acre impoundment four miles south of Two Buttes Reservoir) support winter populations of 50,000 lesser Canada geese and 10,000 mallards. In some years Turks Pond can literally be filled with geese, which creates a problem with fowl cholera. Both of these areas are important waterfowl hunting areas; Two Buttes for public hunting and Turks Pond for commercial hunting.

Two Buttes Reservoir is rated as a major warm water fishing reservoir (Stream and Lake Evaluation Map, Colorado, 1979).

Forested wetlands along streams are important habitat for mule deer use these areas, and a variety of wildlife.

Semi-permanently flooded cattail stands provide critical habitat for pheasants.

Intermittent depressions, streams, and windmills are important water sources for wildlife, such as antelope, scaled quail, and lesser prairie chicken, and livestock in this semi-arid region.

Wetland Loss and Vulnerability

Wetland loss is not a major problem in La Junta NE. This quadrangle is sparsely populated and there were few wetlands originally.

REFERENCES

- Bailey, Robert. 1978. Description of the Ecoregions of the United States. USDA-Forest Service. Intermountain Region. Ogden, Utah. 77 pp.
- Benci and McKee. 1977. Colorado Monthly Temperature and Precipitation Summary for Period 1951-1970. Climatology Report 77-1. Department of Atmospheric Science. Colorado State University. Fort Collins, Colorado.
- Chronic, John and Halka. 1972. Prairie, Peak, and Plateau. Colorado Geological Survey, Bulletin #32. 126 pp.
- Cipra, J. E., Moreland, D. C., Williams, R. D., Kornblau, M. L., Heil, R. D. 1978. Soil Resources of Colorado. C.S.U. Experiment Station and USDA Soil Conservation Service. Special Series #6. 154 pp.
- Cowardin, Carter, Golet, Laroe. 1979. Classification of Wetland and Deep-Water Habitats of the United States. USDI-Fish and Wildlife Service, National Wetland Inventory. St. Petersburg, Florida. 100 pp.
- Fasset, Norman. 1957. A Manual of Aquatic Plants. University of Wisconsin Press. Madison, Wisconsin. 405 pp.
- Fish and Wildlife Service Staff. 1979. Stream and Lake Evaluation Map, State of Colorado. USDI-Fish and Wildlife Service. Salt Lake City, Utah.
- Geological Survey Staff. 1974. Hydrologic Unit Map of Colorado. USDI-Geological Survey. Reston, Virginia.
- Harrington, H. D. 1955. Manual of Plants of Colorado. Swallow Press. Chicago. 666 pp.
- Hopper, Richard. 1980. Personal Communication.
- Langford and Kapinas. 1979. The National Water Data Network: A Case History. Water Resources Research. Vol. 15, No. 6. pp. 1687-1691.
- National Wetland Inventory Staff. 1975. Ecoregions and Land-Surface Form Map for La Junta 1:250,000 Quadrangle. USDI-Fish and Wildlife Service National Wetland Inventory. St. Petersburg, Florida.
- Nelson, Ruth. 1977. Handbook of Rocky Mountain Plants. Skyland Publishers. Estes Park, Colorado. 331 pp.
- Soil Conservation Service Staff. 1978. Temperature Extremes and Freeze Data. USDA-Soil Conservation Service. Fort Collins, Colorado.

Thornbury, William D. 1965. Regional Geomorphology of the United States.
John Wiley and Sons. New York.

Tweto, Ogden. 1979. Geologic Map of Colorado. USDI-Geological Survey.
Denver, Colorado.

Weber, William. 1976. Rocky Mountain Flora. Colorado Associated University
Press. Boulder, Colorado. 479 pp.