

COLORADO WETLAND INVENTORY
U. S. FISH AND WILDLIFE SERVICE
1:100,000 MAP NARRATIVE REPORT
-LAMAR NE-

INTRODUCTION

In 1974, the U. S. Fish and Wildlife Service directed its Office of Biological Services to complete an inventory of the Nation's wetlands. As part of this overall objective, an effort began in September 1978, to delineate and classify photo-interpretable wetlands within the eastern five-sevenths of Colorado.

Wetland maps at 1:100,000 scale and wetland overlay maps at 1:24,000 scale are produced at National Wetland Inventory headquarters in St. Petersburg, Florida. Final Colorado wetland maps are available at the U. S. Fish and Wildlife Service's Regional Office located in Denver, Colorado. An integral part of all final wetland maps is the completion of narrative reports for each 1:100,000 quadrangle inventoried. The following narrative report provides both basic and specific data which aids the user in understanding not only the general area of the Lamar NE quadrangle but also representative details of its wetland habitat.

MAP PREPARATION

Contractor for this wetland inventory was Colorado Division of Wildlife, 317 West Prospect, P. O. Box 2287, Fort Collins, Colorado 80526. Richard Hopper was the contract officer. Photo-interpretation was done by the subcontractor, Colorado State Forest Service, Foothills Campus, Colorado State University, Fort Collins, Colorado 80523. Photo-interpreters were Thomas Owens and Alexander Kosinski. Preparation of this narrative report was completed by Thomas Owens. Regional Wetland Coordinator was Charles Elliott, U. S. Fish and Wildlife Service, Denver Federal Center, P. O. Box 25486, Denver, Colorado 80225.

Wetland delineation and classification for Lamar NE 1:100,000 quadrangle was done on 1:80,000 black and white aerial photographs taken in June and July 1975 and June and July 1976. Photography covered 100 percent of the quadrangle. Wetland classification was done in accordance with Cowardin, et al., 1979. Specific mapping conventions developed at National Wetland Inventory headquarters were used to assist in photo-interpretation. Field checking was done May 23, 1979.

Map users are cautioned that mapping with aerial photography has limitations. Wetlands are identified and classified through stereoscopic examination of photography on the basis of tone, texture, pattern, site, size, local ecology, and cultural patterns. Aerial photographs

reflect conditions during the year and season they are taken. In addition, the 1:80,000 black and white photography used on this project was photographed for purposes other than wetland mapping. The small scale precludes delineating very small wetlands (less than 1/4-acre) and narrow linear wetlands (less than 15 feet wide). Black and white emulsion makes distinguishing between classes of vegetation (and nonvegetation) difficult. Some imagery was not photographed during the best season for wetland delineation and classification. If photographed too early or too late in the season, moist (dark) tones are not evident. The photography was four to five years old when it was interpreted and land use changes have occurred.

Any discrepancies noticed regarding wetland omissions, inclusions, or errors should be given to the U. S. Fish and Wildlife Service Regional Wetland Coordinator who is located in Denver, Colorado, and whose address is on the previous page.

Special Mapping Problems

This quadrangle has many small depressions scattered across the plains. There are two types of depressions: wetlands with emergent species and non-wetlands without emergents. Non-wetland depressions without emergents are often dark in tone and conspicuous on the imagery, which is due to slightly more moisture and lush vegetation than surrounding areas, but have no lighter toned ring. Depressions with wetland species (usually) have a lighter toned ring in them which shows the extent of inundation after rain. Depressions receive water from local summer thunderstorms, are dry most of the time, and are not part of drainage systems. These wetlands are locally important, but are often difficult to see on the imagery, unless flooded after a thunderstorm.

AREA DESCRIPTION

Bailey's Ecoregions

Lamar NE 1:100,000 quadrangle falls into two provinces in Bailey's Description of the Ecoregions of the United States, 1978, which classifies land into a hierarchal system based upon bioclimatic, geologic, and geomorphic criteria. The first province is Great Plains-Shortgrass, Grama-Buffalo Grass Section (3113L) and is characterized by bunched short grasses, with scattered trees. This province covers 97 percent of the quadrangle. The second province is Tall Grass Prairie, Blue-stem-Grama Prairie Section (2533L) characterized by tall grasses and park land. This province covers 3 percent of the quadrangle in the northeastern corner.

Hammond's Land-Surface Forms

Lamar NE falls into three Hammond Land-Surface Forms which systematically characterizes United States topography (Ecoregion and Land-Surface Form Map, 1975). All forms are in the Interior Division (III). The first form is in the Rocky Mountain Piedmont Subdivision (13). In the north-western corner, covering 17 percent of the quadrangle, are Tablelands of Moderate Relief (III-13B3c) which have 50-80 percent of area gently sloping, local relief 300-500 feet, and 50-75 percent of gentle slope on upland. The other two forms are in the High Plains Subdivision (14). In the south are Smooth Plains (III-14A2c) characterized by more than 80 percent of area gently sloping, local relief 100-300 feet, and 50-75 percent of gentle slope on upland. This form covers 70 percent of the quadrangle. Covering 13 percent in the north are Irregular Plains (III-14B2c) which have 50-80 percent of area gently sloping, local relief 100-300 feet, and 50-75 percent of gentle slope on upland.

Hydrologic Mapping Units

Seven hydrologic mapping units are found in Lamar NE (Hydrologic Unit Map of Colorado, 1974). Hydrologic units are part of an effort by the United States Geological Survey to provide a series of uniform, Nationally consistent maps which accurately delineate hydrographic boundaries for Federal and State water resource agencies. Units are designated by eight-digit numbers tied to a computer file (Catalog of Information on Water Data) which contains information on water data activities (Langford and Kapinas, 1979). Three units are in the Missouri Region (10). 10260702 covers 2 percent in the north, 10260001 covers 15 percent in the north-eastern corner, and 10260004 covers 14 percent in the east. Four units are in the Arkansas-Red-White Region (11). 11030002 covers 14 percent in the southeastern corner and 11020011 covers 35 percent in the west. 11020012 covers 15 percent in the southwest and 11020009 covers 5 percent in the southwestern corner.

Geography

Lamar NE is a semi-arid quadrangle covered by plains which have level to rolling topography. Elevations are from 3,800-4,700 feet and vegetation is short grass prairie with cottonwoods and willows along streams. There are no permanently flowing streams on this quadrangle.

Cheyenne Wells, the quadrangle's largest town, is supported by agriculture, which is Lamar NE's major economic activity. Intensive irrigated agriculture occurs on the quadrangle's eastern half where center pivot sprinklers are supplied with well water. Dryland farming and ranching are practiced away from water sources.

A strip on the eastern edge, 3 miles wide, is in Kansas.

Geology

Several Cretaceous (70-135 million years ago) formations consisting of shale and limestone are in Lamar NE. These formations were deposited when eastern Colorado was under a sea. Large areas in the eastern half of Lamar NE are covered by the Ogallala aquifer, an important water bearing formation of sandstone deposited in the Tertiary Period (3-70 million years ago). Over half the quadrangle is covered by Quaternary (present to 3 million years ago) eolian deposits of dune sand, silt, and loess. Quaternary deposits of gravel and alluvium are found in scattered areas. Modern alluvium is scattered along flood plains on intermittent streams.

Soils

Soil is an important element of wetlands; it is one criterion used to define wetlands. "The substrate of wetlands is predominately undrained hydric soil" (Cowardin, et al.). The National Wetland Inventory, in cooperation with the U. S. Soil Conservation Service, is preparing a list of hydric soils to accompany the Cowardin, et al., wetland classification system.

Two major wetland soil types are found in Lamar NE: soils associated with drainages and soils associated with flood-irrigated meadows.

Where streams are intermittent, wetland soils are sandy and gravelly, unstable, excessively drained, subject to occasional flooding, and have low water tables. Native vegetation includes cottonwoods and annual forbs (Cipra, et al., 1978).

Flood-irrigated soils are found where there is a ready supply of water. These soils may not have been wetland soils originally, but since they have been irrigated the water table is less than 3 feet from the surface. These soils are deep, level, range in texture from sandy loam to clay loam, and are often saline. Vegetation includes salt grass, alkali sacaton, tall grasses, sedges, rushes and cattails in low pockets (Cipra, et al.).

Climate

Lamar NE's climate is semi-arid and continental, with cool dry winters and warm, relatively dry, summers. Cheyenne Wells receives 15.5 inches of precipitation annually, 21 inches of this coming as snow. January's average maximum temperature is 45.0°F., average minimum is 14.9°F. July's average maximum temperature is 91.6°F., average minimum is 59.5°F. (Benci and McKee, 1977). The growing season is 155 days (SCS data, 1978).

WETLANDS

Community Description

Lacustrine System

No lakes are found in Lamar NE (information in this section comes from field notes taken May 27, 1979).

Riverine System

One stream type is delineated; intermittent stream (R4SBW). This stream type is the most common stream type on the eastern plains where there is no permanent water source to supply moisture. Intermittent streams have a sandy substrate that is very well drained which is often scoured by flash floods. These streams flow after snow melt and after local summer thunderstorms.

Palustrine System

An important palustrine type is flood-irrigated meadow (PEMKC). Flood-irrigated meadows are found along streams and below springs. Ditches were built along the meadows' upper edges to allow water to flow from upper sources and spill out over the meadows. Standing water can be found for short periods early in the growing season; soil remains moist for extended periods through the growing season. Flood-irrigated meadows have not been cultivated and vegetation is native. Meadows are hayed or grazed by cattle. Vegetation includes Juncus arcticus, Eleocharis acicularis (species identification was according to Harrington, 1955; Fasset, 1957; Weber, 1976; Nelson, 1977), along with grasses and forbs. Many flood-irrigated fields are cultivated; these were not delineated.

Scrub/shrub and scrub/shrub-emergent areas are found along streams in this quadrangle (PSSW, PSSY, PSS/EMW, PSS/EMY). The water regime is dependent upon the amount of water flowing in the stream. In the western portion, where water is relatively plentiful, the water regime tends toward the seasonal regime. In the east, where moisture is scarce, water regimes are drier. Shrub species are Salix spp.; understory emergents include Juncus arcticus and Carex spp.

Forested wetlands (PFOW, PFOY, PFO/EMW, PFO/EMY, PFO/SSW, PFO/SSY) are found throughout the quadrangle along streams. The water regime situation is the same as that of scrub/shrub areas. Tree species include Populus sargentii and Salix spp. Understory species are the same as those mentioned with scrub/shrub-emergent areas.

On the plains numerous areas (PFLW, PEMW) are delineated which are small dugouts or impoundments constructed to supply water to livestock. These flats are dependent upon local precipitation for their water supply, and

do not receive enough moisture to remain wet year-round. They often have saline soils. Vegetation consists of sparse stands of Distichilis stricta, Sporobolus airoides, forbs and grasses.

Intermittent depressions are also delineated on the plains (PEMJ) that are the result of: wind deflation or blowouts, solution-subsidence (leaching water removes limestone and subsequent deflation), or differential compaction of Tertiary sediments of the plains (Thornbury, 1965). Delineated depressions are generally a few feet in depth, and from 100 to several hundred feet across. There is not sufficient moisture in spring after snow melt to supply them with moisture; they are filled with water after local summer thunderstorms and hold water for a few days. They are dry most of the time, but do receive enough moisture to support stands of Eleocharis spp., as well as upland vegetation.

Many palustrine wetlands are cultivated, but remain wetlands. These are called Palustrine Farmed (PF).

Wetland Values

An important wetland value is flood-irrigated hay meadow production of hay for cattle (information in values section is from Hopper, 1980). Flood-irrigated hay meadows produce hay at a much higher rate than dryland meadows do. Wildlife also benefit from wet hay meadows: ducks nest in them.

Forested wetlands along streams are important habitat for mule deer and a variety of wildlife.

Semi-permanently flooded cattail stands provide critical habitat for pheasants.

Intermittent depressions, streams, and windmills are important water sources for wildlife, such as antelope and scaled quail, and livestock in this semi-arid region.

Wetland Loss and Vulnerability

Wetland loss is not a major problem in Lamar NE. This quadrangle is sparsely populated and there were few wetlands originally.

REFERENCES

- Bailey, Robert. 1978. Description of the Ecoregions of the United States. USDA-Forest Service. Intermountain Region. Ogden, Utah. 77 pp.
- Benci and McKee. 1977. Colorado Monthly Temperature and Precipitation Summary for Period 1951-1970. Climatology Report 77-1. Department of Atmospheric Science. Colorado State University. Fort Collins, Colorado.
- Chronic, John and Halka. 1972. Prairie, Peak, and Plateau. Colorado Geological Survey, Bulletin #32. 126 pp.
- Cipra, J. E.; Moreland, D. C.; Williams, R. D.; Kornblau, M. L.; Heil, R. D. 1978. Soil Resources of Colorado. C.S.U. Experiment Station and USDA-Soil Conservation Service. Special Series #6. 154 pp.
- Cowardin, Carter, Golet, Laroe. 1979. Classification of Wetland and Deep-Water Habitats of the United States. USDI-Fish and Wildlife Service, National Wetland Inventory. St. Petersburg, Florida. 100 pp.
- Fasset, Norman. 1957. A Manual of Aquatic Plants. University of Wisconsin Press. Madison, Wisconsin. 405 pp.
- Geological Survey Staff. 1974. Hydrologic Unit Map of Colorado. USDI-Geological Survey. Reston, Virginia.
- Harrington, H. D. 1955. Manual of Plants of Colorado. Swallow Press. Chicago. 666 pp.
- Hopper, Richard. 1980. Personal Communication.
- Langford and Kapinas. 1979. The National Water Data Network: A Case History. Water Resources Research. Vol. 15, No. 6. pp. 1687-1691.
- National Wetland Inventory Staff. 1975. Ecoregions and Land-Surface Form Map for Lamar 1:250,000 Quadrangle. USDI-Fish and Wildlife Service National Wetland Inventory. St. Petersburg, Florida.
- Nelson, Ruth. 1977. Handbook of Rocky Mountain Plants. Skyland Publishers. Estes Park, Colorado. 331 pp.
- Soil Conservation Service Staff. 1978. Temperature Extremes and Freeze Data. USDA-Soil Conservation Service. Fort Collins, Colorado.
- Thornbury, William D. 1965. Regional Geomorphology of the United States. John Wiley and Sons. New York.
- Tweto, Ogden. 1979. Geologic Map of Colorado. USDI-Geological Survey. Denver, Colorado.

Weber, William. 1976. Rocky Mountain Flora. Colorado Associated University Press. Boulder, Colorado. 479 pp.