

WETLAND USER NOTES -- Nogales SE QUADRANGLE

1. Map Preparation

Wetland classification for the National Wetlands Inventory (NWI) wetland map overlay to the Nogales SE 1:100,000 scale map are in accordance with L. M. Cowardin et al (1979). Wetland delineations and classifications were produced through the interpretation of black and white aerial photographs at a scale of 1:120,000 taken during June 1973. The photographs were viewed stereoscopically at a 6X magnification. Delineations were enlarged using a zoom-transferscope to overlays of 1:24,000 and 1:62,500. Overlays were then transferred to 1:100,000 scale base maps. Limited field checks were performed during February 1979.

The Project Officer for production of the wetland map was Warren Hagenbuck, Regional Wetlands Coordinator, (U.S. Fish and Wildlife Service, Region 2, P.O. Box 1306, Albuquerque, New Mexico 87103, (505) 766-2914). Aerial photo interpretations were completed by the School of Renewable Natural Resources, University of Arizona, Tucson 85721, Project Director: Dr. Jon Rodiek (602) 626-2313.

The user of the map is cautioned that due to mapping, primarily through photo interpretation, a small percentage of wetlands may have gone unidentified. Landscape changes could have taken place since the time of photo acquisition; therefore, discrepancies or land use changes should be provided to the Regional Wetlands Coordinator, Region 2, U.S. Fish and Wildlife Service.

2. Collateral Data

A. General Location

The area is within Santa Cruz and Cochise Counties, Arizona.

Northern boundary	31° 30' N Latitude
Southern boundary	Mexican - U. S. Border
Eastern boundary	110° W Longitude
Western boundary	111° W Longitude

B. Land Ownership

There are four (4) major land holdings within the Nogales SE quadrangle. These are:

Coronado National Forest
Fort Huachuca Park
Coronado International Mineral
BLM Lands

C. Soils

There are six (6) major soil associations found on the Nogales SE according to J. E. Jay et al (1975). These are:

Mesic Subhumid Soils -- mean annual soil temperatures of 47-59° F (8-15° C) and more than 16 inches (41 cm) mean annual precipitation.

- MH 1 Casto-Martinez-Canelo Association
- MH 2 Lithic Haplustolls-Lithic Argiustolls-Rock Outcrop Association

Thermic semi-arid soils -- mean annual soil temperatures of 59° to 72° F (15-22° C) and 10-16 in (25-41 cm) mean annual precipitation.

- TS 4 White House-Bernardino-Hathaway Association
- TS 14 Nickel-Latene-Cave Association
- TS 5 Caralampi-Hathaway Association
- TS 6 Lithic Torriorthents-Lithic Haplustolls-Rock Outcrop Association

Detailed soil information may be found in USDA Soil Conservation Service (1977) and M. L. Richardson and M L Miller (1974).

D. Biotic Communities

The descriptions of the biotic communities are summarized from C. H. Lowe (1977) and Brown, Lowe and Pase (1977). The community types are listed here in descending order of their relative dominance in the Nogales SE quadrangle.

The Semidesert Grassland, accounting for 8.0% of Arizona's total land area, is a transitional type of grass dominated landscape positioned between desert below and evergreen woodland or chapparal above. Its lower limit is about 3,500 feet in elevation and its best development is between 4,000 and 5,000 feet. In Arizona it occurs largely in the southeastern quarter of the state. A small area also occurs in the northwestern quarter near Kingman. The grasses are often bunch-growth perennials in which the bases of clumps are separated by bare ground. Where the soil is deep black gramma, blue gramma, slender gramma and others may be the dominant cover over large areas. A wider variety of shrubs, trees and cacti life forms occur on shallow-soiled rocky and gravelly hills and slopes. Mesquite trees have invaded large areas of former grassland. Mean annual precipitation ranges between 10 and 15 inches.

The Madrean Evergreen Woodland type covers 3% of Arizona's total land area. It is dominated by species of evergreen oak including silverleaf oak, Arizona oak, and Emory oak which occur primarily in the southeastern quarter of the state. They occur in hills and mountain slopes between 4,000 and 6,500 feet, reaching their greatest development in the foothills of the larger mountain ranges. The shrubs which center their

User Notes Nogales SE Arizona Wetland Map

distribution in these oakwoodland types, range upward into the pine forests or downward into the desert grasslands. Characteristic species include mountain yucca, agave, and woodland sumac. Other dry-tropic shrubs and succulents occur in varying degrees in these interior southwestern woodlands. A few cacti and grasses such as bullgrass, little blue stem and woody bunchgrass are found with blue gramma being the most common. Mean annual precipitation ranges between 12 and 22 inches.

The Plains and Great Basin Grassland spread over 15.5% of Arizona's land area, and forms a nearly continuous cover in an elevational zone between 5,000 and 7,000 feet. This type generally occurs in the eastern portion of the state. The grasses include gramma grass, needlegrass, dropseed and muhly. Plains grassland is a semi-arid habitat occasionally extending upward into the lower portion of the transition zone (ponderosa pine forest) in Northern Arizona and often is in mixtures with juniper-pinyon woodland and sagebrush. Mean annual precipitation ranges between 11 and 18 inches.

The Petran Subalpine Conifer Forest (Spruce-Fir) type accounts for .5% of Arizona's total land area. It occurs from 7,500 - 9,500 feet in elevation depending upon slope exposure. It reaches its greatest development in Arizona on the higher ranges rising from the Colorado Plateau in the central, north and east portions of the state. It is there that one of its best developments occurs in the extensive elevated mass known as the White Mountains. The dominant tree species include Douglas fir, White fir, Alpine fir, Englemann spruce, and Blue spruce. Broadleaf deciduous trees include quaking aspen, water birch and Rocky Mountain maple. Mean annual precipitation ranges between 25 to 40 inches.

The Chihuahuan Desertscrub represents 1.5% of Arizona's land area and is a shrub desert. Representative areas are found in the southeastern corner of the state, primarily in Cochise County. Tarbush, creosotebush, sandpaperbush, or Chihuahuan white-thorn comprise the four major association-types of this desert occurring in Arizona. In many areas tarbush, creosotebush and white-thorn are intermixed and associated with such species as desert sumac, ocotillo, and mesquite. The Chihuahuan Desert has a relatively poor representation of cactus life-forms due to its relatively high elevation (3,500'). Desert and grassland species form landscape mosaics controlled by marked changes in soil conditions over short distances. Mean annual precipitation is approximately 10 inches.

3. SUPPLEMENTAL DATA (N.W.I.)

A. Hydrologic Units

Major hydrologic units found within the Nogales SE quadrangle are defined by the U.S. Geologic Survey (1974). For example, the coding designates region (15) Lower Colorado, subregion (01), accounting unit (00), and cataloging unit (05).

15050202
15050301

There are two (2) groundwater areas designated within the quadrangle by the U.S. Geological Survey (1979). These are:

Upper San Pedro Basin
Upper Santa Cruz Basin

B. Geography

Landforms

Physical subdivisions and land surface forms are classified according to E. H. Hammond (1965). There are two (2) major types:

(V-4) B6a -- Intermontane Basin and Range Area; Plains with high mountains.

(V-4) B5a -- Intermontane; Basin and Range Area; Plains with low mountains.

Bailey's Ecoregions

Bailey's Ecoregion Classification, R. G. Bailey (1975), identifies the area as within the following province(s):

3140L Dry Domain; Semiarid Steppe Division; Mexican Highland Shrub Steppe Province.

3222L Dry Domain; Arid Desert Division; Creosote Bush-Bur Sage Section.

4. Wetland Communities

The wetland and deepwater habitats displayed on the Nogales SE map are dominated by those types occurring along the many short segmented live streams in the area. Notable representatives include Sonoita Creek, San Pedro River, Babocomari River, Parker Canyon Santa Cruz River, Redrock Canyon, and several in the Huachuca Mountains. Lake Patagonia, Parker Canyon Lake and Canelo Hills Cienega are fine representatives of water impoundments in this region.

- L₁OW Lacustrine, Limnetic, Open Water/Unknown Bottom (Parker Canyon Lake and Lake Patagonia)
- POW Palustrine, Open Water/Unknown Bottom (Stock tank and natural types)
- PEM Palustrine, Emergent (nonpersistent)
- PFO Palustrine, Forested
- R₂OW Riverine, Lower Perennial, Open Water/Unknown Bottom
- R₂BB Riverine, Lower Perennial, Beach/Bar
- R₄SB Riverine, Intermittent, Streambed

5. Field Checks

A. Plants Observed

<u>Common Name</u>	<u>Scientific Name</u>
Mesquite	Prosopis juliflora
Fremont cottonwood	Populus fremontii
Baccharis	Baccharis sp.
Arizona sycamore	Platanus wrightii
Emergents (Graminoids)	

<u>Common Name</u>	<u>Scientific Name</u>
White-tailed deer	Odocoileus virginianus

6. Wetland Loss and Vulnerability

Riverine types found in Sonoita Creek, Parker Canyon, Canelo Hills, the Santa Cruz River and Red Rock Canyon are most notable. Live stream segments here are a major attraction to human recreation activities. Some controlled management of these areas left unprotected is needed.